BMJ Open KAP-COVID_{GLOBAL}: a multinational survey of the levels and determinants of public knowledge, attitudes and practices towards COVID-19

Ahmed Taher Masoud ¹, Mohamed Sayed Zaazouee,² Sarah Makram Elsayed,³ Khaled Mohamed Ragab,⁴ Esraa M Kamal,¹ Yusra T Alnasser,⁵ Ahmed Assar,⁶ Anas Z Nourelden,⁷ Loai J Istaitiah,¹ Mohamed M Abd-Elgawad,¹ Ahmed T Abdelsattar,¹ Ahmed A Sofy,¹ Doaa G Hegazy,⁸ Victor Z Femía,⁹ Adriana R Mendonça,⁹ Fatma M Sayed,¹ Ahmed Elmoursi,¹⁰ Alaa Alareidi,¹¹ Ahmed K Abd-Elkawab,¹ Mohamed Abdelmonem,¹ Omar M Mohammed,¹ EzzEldeen A Derballa,¹ Kareem A El-Fas ¹², Mohamed M. Abdel-Daim ^{13,14}, Abdelrahman I. Abushouk ¹⁵ for the KAP-COVID_{GLOBAL} Investigators

To cite: Masoud AT, Zaazouee MS, Elsayed SM, *et al.* KAP-COVID_{GLOBAL}: a multinational survey of the levels and determinants of public knowledge, attitudes and practices towards COVID-19. *BMJ Open* 2021;11:e043971. doi:10.1136/bmjopen-2020-043971

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-043971>).

MMA-D and AIA contributed equally.

Received 19 August 2020
Revised 18 January 2021
Accepted 03 February 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Abdelrahman I. Abushouk;
Abdelrahman.abushouk@med.asu.edu.eg

ABSTRACT

Objective The adherence to public health recommendations to control COVID-19 spread is influenced by public knowledge, attitudes and practices (KAP). We performed this cross-sectional study to assess the levels and determinants of public KAP towards COVID-19 in a large, multinational sample.

Design Cross-sectional study (survey).

Setting The questionnaire was distributed to potential respondents via online platforms.

Participants 71 890 individuals from 22 countries.

Methods We formulated a four-section questionnaire in English, followed by validation and translation into seven languages. The questionnaire was distributed (May to June 2020) and each participant received a score for each KAP section.

Results Overall, the participants had fair knowledge (mean score: 19.24 ± 3.59) and attitudes (3.72 ± 2.31) and good practices (12.12 ± 1.83) regarding COVID-19. About 92% reported moderate to high compliance with national lockdown. However, significant gaps were observed: only 68.2% knew that infected individuals may be asymptomatic; 45.4% believed that antibiotics are an effective treatment; and 55.4% stated that a vaccine has been developed (at the time of data collection). 71.9% believed or were uncertain that COVID-19 is a global conspiracy; 36.8% and 51% were afraid of contacting doctors and Chinese people, respectively. Further, 66.4% reported the pandemic had moderate to high negative effects on their mental health. Female gender, higher education and urban residents had significantly ($p \leq 0.001$) higher knowledge and practice scores. Further, we observed significant correlations between all KAP scores.

Conclusions Although the public have fair/good knowledge and practices regarding COVID-19, significant gaps should be addressed. Future awareness efforts should target less advantaged groups and future studies should develop new strategies to tackle COVID-19 negative mental health effects.

Strengths and limitations of this study

- Besides our large sample size (of both individuals and countries), such comparative study can help international organisations focus their efforts on countries and population groups with less developed public knowledge, attitudes and practices (KAP) against COVID-19.
- In addition, we analysed the association between KAP and demographic factors, history of COVID-19, as well as the correlation between different scores and scales of mental health effects and compliance to lockdown. However, this study is not without limitations.
- First, as a cross-sectional study, the temporal relevance of our findings may change with time or implementation of large-scale prevention measures.
- Second, the elderly population (most vulnerable to COVID-19) only represents 3.9% of our sample. This is probably related to the online distribution of the questionnaire, which is likely to draw younger populations. The requirement of access to electronic devices and the internet may have limited the reach based on wealth and literacy.
- Third, as a self-reported questionnaire, respondents may have opted towards socially desired choices rather than their actual KAP. Fourth, due to variations in the population size of the included countries, our fixed minimum sample size may have been less representative of more populous nations as India, Brazil and the USA.

INTRODUCTION

Starting in China in December 2019, SARS-CoV-2 (the causative agent of COVID-19) has spread to almost every country worldwide.¹ As of 16 February 2021, over 110

million confirmed cases have been reported globally with more than 2.4 million deaths.² The disease is transmitted by respiratory droplets. After an incubation period of 2–14 days, patients may develop fever, cough, dyspnoea, fatigue and sore throat or are commonly asymptomatic.^{3,4} The main cause of death is fatal pneumonia and respiratory distress. Adults with chronic diseases and those over 65 years of age are the most vulnerable.⁵ Although various drugs are under trial, the management remains mainly supportive. Therefore, prevention measures as mass vaccination, social distancing, face masks and public awareness campaigns are key players in controlling the pandemic.⁶

However, there is lack of data on the awareness and practices of different populations and their influence on COVID-19 burden. Multiple cross-sectional studies have been conducted in many countries, for example, an early questionnaire was developed by Zhong *et al*⁷ in China, and it was later applied in other countries as Italy,⁸ India,⁹ Malaysia,¹⁰ Pakistan¹¹ and Colombia.¹² Another survey in the USA revealed that a large portion of the public lacked critical knowledge about COVID-19 and were not changing their daily routine and hygiene practices as per the recommendations of health authorities.¹³ However, when coupled with extensive governmental awareness efforts, the public awareness on COVID-19 can be significantly improved as revealed by recent studies from Saudi Arabia¹⁴ and Nigeria,¹⁵ which would reduce infection rates and alleviate the medical and economic burdens of the disease.

The success of prevention efforts is tied to public adherence and the latter has been linked to public knowledge, attitudes and practices (KAP).^{16,17} A large-scale, horizontal evaluation of KAP towards COVID-19 across different countries is lacking. Plus, this evaluation was not performed in most low-to-middle income countries. In the present multinational survey, we aimed to assess the levels of public KAP in different countries towards COVID-19 and to determine the factors that could influence public practices in this regard. Our findings may have implications for public awareness efforts worldwide.

MATERIALS AND METHODS

Study design and participants

We conducted a multinational, cross-sectional study to assess public KAP towards COVID-19 in 22 countries using an online self-administered questionnaire during the period of 10 May to 25 June 2020. The study was conducted and reported in consistence with the Strengthening the Reporting of Observational Studies in Epidemiology checklist (online supplemental appendix 1). Any citizen of the included countries above the age of 18 who agreed to fill the questionnaire was eligible to participate. There were no demographic restrictions on participation.

Sampling

We used a convenience sampling method for data collection. The sample size was calculated for each country using

the equation: $n = z^2 P(1-P)/d^2$.¹⁸ Under a 95% CI, 50% response distribution and 0.05 margin of error, a sample of 384 participants was considered as a minimum sample to represent large populations. However, due to the limitations of convenience sampling and online surveying, we empowered our sample by including a design effect (DE) factor in the equation. According to previous studies, the minimal acceptable DE for convenience-sampled studies is 2.^{19,20} Therefore, an adjusted minimum sample of 768 (384×2) participants was considered for each country.

Questionnaire development

The questionnaire was developed using the frequently asked questions on the WHO and Centers for Disease Control and Prevention websites in addition to the previously published national surveys of COVID-19/other pandemics awareness.^{7,21–23} Experts from the departments of Community Medicine & Public Health and Internal Medicine (division of infectious diseases) at Fayoum University (Fayoum, Egypt) formulated the questionnaire. The questionnaire was revised by the departments' heads for face validity, relevance, comprehensiveness and clarity of each section, and some details were improved.

The final four-section questionnaire included:

- Sociodemographic data: that collected participants' age, gender, country, residence (urban/rural), educational level, whether they or a family member/friend had been diagnosed (by a medical doctor) with COVID-19.
- Knowledge about COVID-19: consisted of 28 questions about COVID-19 mode of transmission, vulnerable groups for infection, symptoms, treatment, prevention measures and mortality rate. The answer to each question was Yes/No/I don't know choices, except for the question about the mortality rate. Cronbach's alpha values for the knowledge assessment section were 0.76, 0.55, 0.70, 0.60, 0.75, 0.70, 0.60 and 0.64 for English, Arabic, French, Indonesian, Nepali, Pakistani, Sinhala and Portuguese languages, respectively.
- Attitudes towards COVID-19: consisted of eight questions assessing optimism about the current situation; responsible public health attitudes; stigma against symptomatic individuals, healthcare professionals and Chinese people; and whether the participant believes in conspiracy theories about the disease. The possible answers to each question were Agree/Uncertain/Disagree. Cronbach's alpha values for the attitudes assessment section were 0.60, 0.60, 0.77, 0.60, 0.66, 0.64, 0.72 and 0.60 for English, Arabic, French, Indonesian, Nepali, Pakistani, Sinhala and Portuguese languages, respectively.
- In addition, participants were asked to rate their fear of infection and the negative impact of the pandemic on their mental health on a scale from 1 to 10.
- Practices regarding COVID-19: included 14 questions describing different practices regarding coughing and sneezing, hand washing, wearing masks and contact with people. The available answers to each

question were Yes/Sometimes/No. In addition, the participants were asked to rate their overall compliance with the lockdown or the measures applied by their country on a scale from 1 to 10. The Cronbach's alpha values for the practices assessment questionnaire were 0.77, 0.67, 0.66, 0.66, 0.67, 0.55, 0.68 and 0.55 for English, Arabic, French, Indonesian, Nepali, Pakistani, Sinhala and Portuguese languages, respectively. The full version of the questionnaire can be found in [tables 1–4](#).

The questionnaire was developed in English and was then translated into the native languages of the included countries (Arabic, French, Indonesian, Nepali, Portuguese, Pakistani and Sinhala). For each language, two bilinguals initially performed forward translation, then another bilingual performed a backward translation; the translated versions were compared and checked until a final draft was agreed on. We checked the internal consistency of the questions in each section by calculation of Cronbach's alpha using the data of the first 150 responses from each language; these 150 responses were not included in the final analysis.

Data collection and handling

We recruited collaborators between 20 April and 1 May 2020 in a snowball fashion. The recruited collaborators were given an orientation session about the nature of the study and the data collection strategy. We assigned a central investigator from each country to monitor the data collection process to ensure the adequate contribution of all collaborators (≥ 100 participants) and to avoid over-representation of some cities over others within each country. Each collaborator was granted access to view their responses only, while the central investigator had access to all responses of the country. All collaborators are listed in online supplemental appendix 2.

On 10 May, we started data collection using Google Forms, distributed on social media platforms (repeated posting on Facebook, Twitter, WhatsApp and LinkedIn), online websites, blogs and contact with non-governmental organisations and academic institutions in the included countries. Each participant was allowed to answer the survey only once and no duplicates were included. After the data collection, we used Microsoft Excel for data cleaning. The results of each country were translated automatically to English and were combined in one data-sheet for analysis.

The correct responses to knowledge questions were given a score of 1, while incorrect/I don't know answers were given a score of 0 (hence knowledge maximum score was of 28). The knowledge score of each participant was classified based on the modified Bloom's cut-off points into poor ($<60\%$: <16.8), fair (60% – 79% : 16.8 – 22.1) and good ($\geq 80\%$: 22.2 – 28). In terms of attitudes, the proper attitude was given a score of +1, the improper attitude was given a score of –1 and uncertain was given a score of 0 (hence a maximum positive attitudes score of 8). Regarding practice questions, the

Table 1 Demographic characteristics and COVID-19-confirmed infection rates among survey respondents

Demographics	Count (%) (n=71 890)
Age mean $\pm$ SD (years)	27.64 $\pm$ 9.78
≤30 years	53 048 (73.8)
31–50 years	16 073 (22.3)
>50 years	2 769 (3.9)
Gender	
Male	28 449 (40)
Female	42 601 (60)
Country	
Algeria	4 900 (6.81)
Brazil	839 (1.16)
Egypt	6 830 (9.49)
Ghana	1 847 (2.56)
India	1 464 (2.04)
Indonesia	4 444 (6.18)
Iraq	2 092 (2.92)
Ireland	1 026 (1.43)
Jordan	5 882 (8.20)
Lebanon	3 380 (4.70)
Libya	4 300 (6.00)
Morocco	1 755 (2.44)
Nigeria	3 449 (4.80)
Nepal	2 657 (3.70)
Palestine	5 993 (8.33)
Pakistan	1 723 (2.40)
Saudi Arabia	1 616 (2.24)
South Africa	1 979 (2.75)
Sri Lanka	1 793 (2.50)
Sudan	4 381 (6.09)
Syria	6 576 (9.14)
UK	2 160 (3.00)
USA	804 (1.12)
Education	
High school	7 577 (10.54)
Undergraduate	44 436 (61.80)
Graduated	16 269 (22.65)
Prefer not to say	3 608 (5.01)
Residency	
Urban	57 653 (80.2)
Rural	14 237 (19.8)
Have you had a confirmed infection with COVID-19?*	
Yes	1 326 (1.84)
No	70 559 (98.16)
Do you know a friend or a family member who had a confirmed COVID-19 infection?*	
Yes	9 935 (13.82)
No	61 952 (86.18)

The presented data are count (valid %) unless otherwise specified.

*Confirmed infection was explained to participants as having diagnosis by a licensed healthcare professional.

Table 2 Answers to knowledge questions about COVID-19 among survey respondents

	Yes	No/I don't know
COVID-19 is a serious disease.	58 121 (80.8)	13 769 (19.2)
Regarding the mode of transmission of the virus		
Eating wild animals (eg, bats)	37 829 (52.6)	34 061 (47.4)
Droplet transmission	65 126 (90.6)	6764 (9.4)
Contact with infected surfaces then putting your hand on your face, mouth or nose	68 819 (95.7)	3071 (4.3)
The most vulnerable group to infection is		
Children (0–18 years)	32 973 (45.9)	38 917 (54.1)
Adults (18–50 years)	29 448 (41.0)	42 442 (59.0)
Elderly (more than 50 years)	64 736 (90.0)	7154 (10.0)
Adults with chronic diseases	67 282 (93.6)	4608 (6.4)
The clinical symptoms of COVID-19 include		
Fever	68 565 (95.4)	3325 (4.6)
Fatigue	57 076 (79.4)	14 814 (20.6)
Dry cough	66 882 (93.0)	5008 (7.0)
Myalgia	50 672 (70.5)	21 218 (29.5)
Stuffy nose	21 127 (29.4)	50 763 (70.6)
Runny nose	23 449 (32.6)	48 441 (67.4)
Sneezing	14 094 (19.6)	57 796 (80.4)
Shortness of breath	68 589 (95.4)	3301 (4.6)
Diarrhoea	35 293 (49.1)	36 597 (50.9)
Asymptomatic	49 057 (68.2)	22 833 (31.8)
Regarding treatment of COVID-19		
There is effective cure for it.	53 148 (73.9)	18 742 (26.1)
The treatment is symptomatic only.	51 140 (71.1)	20 750 (28.9)
Antibiotics are an effective treatment.	32 628 (45.4)	39 262 (54.6)
There are various drugs under trial.	58 231 (81.0)	13 659 (19.0)
A vaccine has been developed.	39 798 (55.4)	32 092 (44.6)
Regarding prevention of COVID-19, which of the following is effective?		
Wearing medical masks	67 365 (93.7)	4524 (6.3)
Avoiding crowded places	70 509 (98.1)	1380 (1.9)
Isolation of infected individuals	70 551 (98.1)	1338 (1.9)
Healthy diet and avoiding high-fat-containing diet	18 159 (25.3)	53 730 (74.7)
To what extent does COVID-19 cause death?	Less than 15%: 42 522 (59.1)	Wrong answers: 29 368 (40.9)

Data are presented as count (%).

correct practice was given a score of 1, (sometimes) was given a score of 0.5 and incorrect practice was given a score of 0 (hence a maximum practices score of 14). The participants' responses to scale questions (from 1 to 10) were classified as low (1–3), moderate (4–7) or high (8–10).

Statistical analysis

We used SPSS (V.24, IBM) for data analysis. Quantitative outcomes (eg, scores) were presented as mean±SDs. Associations were analysed using the independent samples

t-test and one-way analysis of variance (ANOVA) with post hoc Hochberg test, while the correlation between different scores was assessed using Pearson correlation tests. We used Tableau software (Seattle, Washington) for geographical map presentation.

Patient and public involvement

It was not appropriate or possible to involve patients or the public in the design, or conduct, or reporting, or dissemination plans of our research.

Table 3 Answers to attitude questions about COVID-19 among survey respondents

	Positive attitude	Uncertain	Negative attitude
Do you believe that COVID-19 is a global conspiracy?	20 191 (28.1)	32 559 (45.3)	19 140 (26.6)
Do you believe that COVID-19 pandemic will finally end?	34 053 (47.4)	30 851 (42.9)	6986 (9.7)
Do you believe that your country will be able to control COVID-19 situation soon?	32 526 (45.2)	28 603 (39.8)	10 760 (15.0)
Since the outbreak, I seek more medical information about COVID-19 to keep updates.	59 215 (82.4)	9090 (12.6)	3585 (5.0)
Since the outbreak, I follow the recommendations to deal with the pandemic.	66 609 (92.7)	4350 (6.1)	931 (1.3)
Since the outbreak, I am afraid to contact anyone with ordinary influenza symptoms.	55 788 (77.6)	10 236 (14.2)	5866 (8.2)
Since the outbreak, I am afraid of contacting any doctors except for the utmost necessity.	45 475 (63.2)	13 292 (18.5)	13 123 (18.3)
Since the outbreak, are you afraid of eating in Chinese restaurants or contact Chinese people?	35 243 (49.0)	15 356 (21.4)	21 291 (29.6)

Data are presented as count (%).

RESULTS

Demographic characteristics and COVID-19 infection rates

The present sample comprised 71 890 respondents from 22 countries around the globe. The mean age of all participants was 27.64 ± 9.78 years and 42 601 (59.3%) were females. The majority lived in African and Asian countries, enrolled in/graduated from college education and were living in urban settings. Among those surveyed, 1326 (1.84%) reported having

been diagnosed with COVID-19 while 9935 (13.82%) reported knowing a friend or a family member who had been diagnosed with COVID-19 (table 1).

Public knowledge regarding COVID-19

The mean knowledge score among all respondents was $19.24/28 \pm 3.59$ (fair). Of them, 14 221 (19.8%), 45 087 (62.7%) and 12 582 (17.5%) had poor, fair and good knowledge levels, respectively. The majority of

Table 4 Answers to practice questions about COVID-19 among survey respondents

	Proper practice	Sometimes	Wrong practice
When coughing or sneezing, do you			
Cover your mouth and nose with a tissue?	62 946 (87.6)	6749 (9.4)	2195 (3.1)
Throw away the used tissue into the bin?	66 291 (92.2)	2731 (3.8)	2868 (4.0)
Turn your face away from the surrounding people?	67 664 (94.1)	2303 (3.2)	1923 (2.7)
As for your hands, you wash them			
Before touching your eyes and nose.	52 267 (72.7)	14 516 (20.2)	5107 (7.1)
After covering the nose while sneezing.	56 198 (78.2)	9994 (13.9)	5698 (7.9)
After coming from outside.	67 791 (94.3)	2815 (3.9)	1284 (1.8)
Using soap and water.	68 337 (95.1)	2299 (3.2)	1254 (1.7)
Using concentrated alcohol.	39 729 (55.3)	16 862 (23.5)	15 299 (21.3)
Regarding wearing a face mask, you			
Wear a face mask in crowded places.	58 939 (82.0)	5082 (7.1)	7869 (10.9)
Wear a face mask outside in general (not crowded).	37 364 (52.0)	12 371 (17.2)	22 155 (30.8)
Never use a face mask.	46 829 (65.1)	12 813 (17.8)	12 247 (17.0)
Regarding the preventive measures from infection, you			
Avoid contact with an infected person.	69 776 (97.1)	1037 (1.4)	1077 (1.5)
Avoid touching and shaking hands.	61 981 (86.2)	7044 (9.8)	2865 (4.0)
Avoid going to crowded places.	63 201 (87.9)	6684 (9.3)	2005 (2.8)

Data are presented as count (%).

respondents agreed that COVID-19 is a serious disease (80.8%); correctly identified droplet transmission (90.6%) and contact with surfaces covered with infected droplets (95.7%) as the mode of transmission; correctly identified elderly subjects (90%) and adults with chronic disease (93.6%) as the vulnerable groups to COVID-19 infection; and the majority could identify the correct prevention measures against COVID-19 infection. However, we detected some critical knowledge gaps, for example, only 68.2% knew that infected individuals may be asymptomatic. Regarding treatment, 73.9% stated that there is an effective cure for COVID-19, 45.4% stated that antibiotics are an effective treatment and 55.4% stated that a vaccine has been developed (at the time of data collection), while only 59.1% identified the correct mortality rate for COVID-19 (table 2).

Data analysis showed that demographic factors influenced knowledge scores, being significantly higher (≤ 0.001) in females, urban residents, those with higher education or who knew a family member or a friend who had a confirmed diagnosis with COVID-19 disease. Interestingly, those who reported a confirmed COVID-19 diagnosis before had a lower knowledge level. The one-way ANOVA test showed that the mean knowledge levels differed across the surveyed countries ($p < 0.001$), with the highest mean scores from Brazil, Egypt, Jordan, Sudan and Syria and the lowest mean scores from Indonesia, Nigeria, Pakistan and India (online supplemental appendix 3).

Public attitudes towards COVID-19

The mean attitudes score towards COVID-19 among the surveyed respondents was $3.72/8 \pm 2.31$. Some positive attitudes were observed, for example, the majority of respondents (>80%) stated that since the outbreak, they seek updated medical information and recommendations about COVID-19. However, 71.9% believed or were uncertain that COVID-19 is a global conspiracy; >50% were uncertain or not optimistic that the pandemic will finally end or that their government will be able to control COVID-19 situation; 36.8% were uncertain or afraid of contacting doctors except for utmost necessity; and 51% were afraid or uncertain about contacting Chinese people and eating in Chinese restaurants (table 3). When the respondents were asked to rate their fear of getting COVID-19, 20021 (27.8%), 33752 (46.9%) and 18117 (25.2%) reported low, moderate and high levels of fear, respectively. Further, 47712 (66.4%) reported that the pandemic had moderate to high negative effects on their mental health.

Similar to knowledge levels, the overall attitude score was significantly higher in females ($p = 0.002$) or those who knew a family member or a friend with a confirmed COVID-19 diagnosis ($p = 0.003$). However, those with previous COVID-19 diagnosis had less positive attitude scores ($p < 0.001$) compared with those without COVID-19 diagnosis history. Further, the overall attitudes score, fear of getting COVID-19 and the negative mental health

Figure 1 Geographic representation of (A) fear of getting COVID-19, (B) negative mental health effects of COVID-19 pandemic, and (C) compliance with governmental lockdown/traffic ban across the 22 countries. The colour gradient (from light to dark) in every map represents country scores (from low to high).

impact varied by country (online supplemental appendix 3, figure 1A,B).

Public practices regarding COVID-19

The mean practices score ($12.12/14 \pm 1.83$) and answers to individual questions showed good practices towards COVID-19. The majority of respondents indicated that they usually follow proper practices regarding hand washing, coughing and sneezing, wearing face masks and social distancing. Few gaps were, however, noted. Although 82% indicated that they usually wear face masks in crowded places, only 52% responded that they usually wear face masks outside in general and 17% replied that they never wear face masks (table 4). When the respondents were asked about their overall compliance to their national lockdown/traffic ban, 5856 (8.1%), 19166 (26.7%) and 46868 (65.2%) reported low, moderate and high compliance levels, respectively.

Likewise, females, those with higher education, residing in urban areas or knowing an individual who had a COVID-19 diagnosis had better practice scores ($p \leq 0.001$). However, those who experienced COVID-19 diagnosis reported significantly lower practice scores than those who did not. The one-way ANOVA test revealed that the overall practices score and compliance to national lockdown/traffic ban varied by country (online supplemental appendix 3; figure 1C).

Correlation between KAP towards COVID-19

We recorded significant positive correlations ($p < 0.001$) between KAP scores in our sample, although the magnitude of these correlations in our sample was weak. For example, knowledge scores were positively correlated to attitudes ($r = 0.05$) and practice ($r = 0.12$) scores, while

attitude scores were positively correlated ($r=0.276$) to practice scores.

Interestingly, knowledge was inversely associated with fear of getting COVID-19 ($r=-0.04$) and negative mental health effects of the pandemic ($r=-0.02$) and was directly associated with compliance to lockdown ($r=0.11$). Likewise, better attitudes were associated with lockdown compliance ($r=0.08$) and practice scores were directly correlated to fear of getting COVID-19 ($r=0.167$) (online supplemental appendix 3).

DISCUSSION

The current cross-sectional study assessed the levels and determinants of KAP towards COVID-19 in 22 countries around the globe. Our results show that the public in those countries had fair knowledge and good attitudes towards COVID-19. We, however, uncovered many gaps in the public understanding and behaviours towards COVID-19. For example, one-third of our participants did not know that infected individuals can be asymptomatic, which increases their risk of exposure to the disease. Further, about half of the participants thought that antibiotics may be an effective treatment and about 74% thought that a curative treatment exists, which may give them a false sense of security. Another alarming finding is that almost half of our participants held negative/uncertain attitudes regarding contacting Chinese people and more than one-third had similar attitudes towards doctors.

Our analysis showed that 82% of respondents usually wear face masks in crowded places, but only 52% wear masks outdoors in general. This finding is relevant for public awareness programmes. Several studies and predictive models showed that wearing face masks can reduce COVID-19 spread.^{24 25} In compliance with the building evidence, major public health authorities around the world unanimously recommend wearing face masks outdoors in general, not just in crowded places.^{26 27} However, the compliance rates to these recommendations vary between and within countries. Our study highlights the importance of public awareness about the value of masks in preventing infections and slowing the spread of COVID-19.

In the current study, we found a significant positive correlation between knowledge and attitudes, which coincides with several former studies on COVID-19.^{7 28 29} However, the magnitude of correlation in our study was weak, similar to a former Indonesian study.³⁰ This is probably because although knowledge is essential in shaping attitudes, this is not absolute and several other factors may be involved. A stronger correlation was found between attitudes and practices, indicating that promoting knowledge alone is insufficient and effective interventions to improve practices should target promoting both adequate knowledge and positive attitudes. Interestingly, our analysis also showed lower knowledge scores in those who reported having a confirmed diagnosis with COVID-19. This can be explained in the light of our finding

that knowledge scores were directly correlated to practice scores and compliance with lockdown/traffic ban orders. This suggests that good knowledge translates into safe practices, which can reduce one's risk of COVID-19 infection.

We found an inverse correlation between knowledge level and fear of getting COVID-19. This implies that improving the knowledge about COVID-19 can alleviate public anxiety and panic. During the severe acute respiratory syndrome (SARS) epidemic (2002–2004), misinformation led to excessive public panic and resistance to comply with public health guidelines.^{31 32} We could also infer that people's knowledge would not be improved just by communicating daily increases in COVID-19 cases. In the same vein, about two-thirds of our participants reported moderate to high negative mental health effects for the pandemic. These effects had significant inverse correlations with knowledge and attitude scores; however, the magnitude of the correlations was weak. Several studies showed multiple risk factors for anxiety and mental health problems related to COVID-19, including social media use, worry about economy and personal finances, working in COVID-19 hot spots and being pregnant.^{33 34} Therefore, poor knowledge and attitudes may contribute—among a multitude of factors—towards the growing incidence of mental health issues, being reported worldwide.

The association between KAP scores and demographic characteristics in the current study was consistently significant. For example, females had better KAP scores towards COVID-19 than males. This finding echoes previous studies by Al-Hanawi *et al*³⁵ and Azlan *et al*.¹⁰ In addition, those living in rural areas had lower knowledge and practice scores than their urban counterparts. This may be attributed to relying on digital sources of information with easier access in urban settings or the higher levels of education in urban areas, which were also associated with higher KAP scores in the present study.

Most of the included countries in the current analysis are low-to-middle income countries. These countries had varied KAP levels and also were significantly different when assessed on three rating questions (fear of COVID-19, negative mental health effects and compliance to lockdown). Other studies have assessed KAP levels in countries that have been included in this analysis (eg, USA, UK, Egypt, Saudi Arabia, Pakistan and Indonesia) and countries outside our scope (eg, Malaysia, Turkey and Italy). To put our study in context, we performed a comprehensive review of published public KAP studies in the literature about COVID-19 (online supplemental appendix 4). The majority of these studies showed good public knowledge and practices across different countries, especially those conducted in the later 3 months (probably due to the growing public awareness about COVID-19).

Practical and research recommendations

Although we did not explore sources of knowledge about COVID-19 in this study, previous works highlighted television and social media as the primary sources of knowledge. Using these platforms should be optimised to deliver evidence-based information to the most vulnerable groups, for example, less educated and those living in rural areas. Political leaders and stakeholders should take action to eliminate fear and discrimination against healthcare professionals and Asian community members.³⁶ Research-wise, future studies should evaluate other populations, not surveyed in the present study; considering the relatively low Cronbach's alpha values (<0.6) in few language translations in our study, these studies should perform validation through pilot testing and revision. In addition, they should test the value of innovative strategies in mitigating mental health effects of public health disasters like COVID-19.

CONCLUSION

The current multinational cross-sectional study showed fair public knowledge on COVID-19; however, it uncovered several gaps in the public understanding and practices about the diseases. Moreover, it highlighted the negative mental health effects of COVID-19 pandemic. Some demographic groups were less advantaged than others including the less educated and those living in rural areas. Future awareness efforts should target those groups and develop innovative strategies to mitigate negative mental health effects, as well as discriminatory behaviours against Asians and healthcare professionals.

Author affiliations

¹Faculty of Medicine, Fayoum University, Fayoum, Egypt

²Faculty of Medicine, Al-Azhar University, Assiut, Egypt

³Faculty of Medicine, October 6 University, Giza, Egypt

⁴Faculty of Medicine, Minia University, Minia, Egypt

⁵Faculty of Medicine, Damascus University, Damascus, Syrian Arab Republic

⁶Faculty of Medicine, Menoufia University, Shebin El-Kom, Egypt

⁷Faculty of Medicine, Al-Azhar University, Cairo, Egypt

⁸Global Clinical Scholars Research Training, Harvard Medical School, Boston, Massachusetts, USA

⁹Universidade do Vale do Sapucaí, Pouso Alegre, Minas Gerais, Brazil

¹⁰College of Medicine, University of Kentucky, Lexington, Kentucky, USA

¹¹School of Environmental and Biological Sciences, Rutgers University, New Brunswick, New Jersey, USA

¹²College of Clinical Pharmacy, King Faisal University, Hofuf, Saudi Arabia

¹³Faculty of Veterinary Medicine, Suez Canal University, Ismailia, Egypt

¹⁴College of Science, King Saud University, Riyadh, Saudi Arabia

¹⁵Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA

Twitter Abdelrahman I. Abushouk @Shoukmed

Collaborators Mariam Salah Moris; Ayman Essa Nabhan; Mohammed Jehad Al-kfarna; Hala Aladwan; Amira Yasmine Benmelouka; Manar Mohammed Hosny; Sara Rajab Araara; Alaa Ahmed Elshanbary; Mariam Ahmed Maray; Ahmed Sultan; Merna Ahmed Riad; Radwa Mohamed Awadalla; MennatulRahman Mohamed Daa-ElEnsaf; Ahmed Saeed Ahmed; Ahmed Fares Ghannam; Mohamed Mahmoud Abdelkarem; Mohamed Marey Yahya; Salama Ahmed Ali; Sara Gamal Fayad; Mohamed Essam; Noha Ahmed Ammar; Maryam Abd-Elmalak Shafik; Osama Mohamed Rokaby; Israa Mohamed Elshahawy; Aya Mosad Elhelesy; Ahmed Bostamy Elshnory; Esraa Ghanem; Mostafa Mahmoud Meshref; Manar Ahmed

Kamal; Katrina Taha Al-Bank; Salam Muhammad Sharif; Houssein Deeb; Tarek Al Soufi; Yara Issa; Danny Salem Knaizeh; Bana Zuhair Alafandi; Hasan Hassan Raslan; Farah AL Bakkar; Mohamed AlHashemi; Sami Jomaa; Sabah Refaieh; Raghad Dannan; Laith Alsabek; Mohammad Yaser Haidar; Hala Jamal Redwan; Nataly Mazen Salhab; Mahmoud Aref Aldrini; Qusai N Zrekat; Sojoud Saleem Alabed; Omar A. Safarini; Ammar Ahmad Thabaleh; Ithar Moufak Barghouthi; Roaa Waleed Abu-Ereban; Ayat Abed-Albaset Mahamid; Bushra Majd Barghouthe; Sahar kamel Balasi; Mo'men Helmi Suleiman; Faris Jamal Abu-Za'nounneh; Obada Ahmad Al-Jayyousi; Mohammad Hasan Ismail; Batool Emad Al-Masri; Israa Aayed Al-Odat; Mahmoud Omar Alshneikat; Tasneem suhail Abu-Alkhair; Hani Adnan Bashir; Mustafa Ismat Aburumman; Mais Hutham Sabri; Hiba Ramadan; Hayat Ghaseb Abu-Alkhoun; Malak Eyad Abu-Qaddoura; Rahmani Meriem; Mohamed Elkhaili Bouaich; Benslimane Sahar; Khennoussi Amel; Ahelam Zerga; Yassamine ouerdane; Hamel Asma; Nedjima Mouhoubi; Nour Salem; Affaf Sahih; Manal Benatia; Wiame Benhabiles; Boutheyna Drid; Sara Menzer; Krazdi Asma; Abdulkarim Aldoukali Babaa; Mabruka Mohamed Algallal; Nawal Aldokali Muhammed; Mohaned Mohammed Ziltni; Mohammed Salem Mansour; Islam Ammar; Mohammed Salem Mansour; Reem Khaled Wishah; Asma Abubakr Saleh; Sahar Azaz; Amal Sharif Eljali; Ahmad Bouhuwaish; Alarabi Alsalem Ali Alarabi; Ahmed Ateia Alzedam; Hamdan Hilan; Mazen Bashir Ahmed; Hiba Mahgoub Eltayeb; Arwa F. Hassan; Anfal Mahmoud Alkhalifa; Walaa Elnaiem; Suad Elsadig Yousif; Lina Sameer Ibrahim; Tareq Fouad Neme; Elaf Mohamed Elhassan; Mona Muhe-Eldeen Eshag; Tayseer Hatim Mohammed; Sjda Ameen Merghany; Victor Carlos Nuvunga; Priscilla Sarfo Adu; Attah Al-Hassan Dawuni; Eliham Salifu; Isaac Mawunyega Kwaku; Faisal Tikuma Abdallah; Rachel Laadi Sulia; Avantika Pandey; Neetish Patel; Motilal Nehru; Shweta Patel; Kamni raj Bavoria; Avantika Pandey; Neetish Rani Patel; Sreenath Sreekantan; Mohammed J Al-Awady; Renas Husain Isa; Mohanad Jawad Kadhim; Raad A. Alharmoosh; Maram Al-Enzi; Fatemah Alalawi; Osman Kamal Osman Elmahi; Muhammad Mahmoud; Dhouha Daassi; Mohamed M. Khodairy; Hasan Mirza; Pratha Rajesh Taiwade; Sanju gautam; Febtrias Mandeabuti Prasetyo; Dewi Anggraini; Dini Setyowati; Ninuk Hariyani; Risa Haryati Tambunan; Bernike Yuriska Metabuti Prasetyo; Theresia Pakaedith Lodang Hurint; Dewi Ayu Ratna Sari; Bernike Yuriska Metabuti Prasetyo; Mulia Daniel Sihotang; Oktavia Manuama; Kuber Bajgain; Suresh Panthee; Buddha Bahadur Basnet; Bimala Panthee; Nanda Kumar Tharu; Rakesh Kumar Lama; Ramesh Kumar Yadav; Sandeep Khattri; Amrita Acharya; Nashib Pandey; Hanane Amer Chamma; Hadi Mohammad Fateh Shammaa; Tarek Abdulkarim Baroud; Marc Samir Machaalani; Bachir Toufik Zrayka; Amir Rabi Al Ayoubi; Lemir Majed Lemir Ahmad El Ayoubi; Ilham Hassan Said-Salman; Jad Samer Al Masri; Anthony Dany Daher; Fouad Mario Assaf; Diala Samer Al Masri; Miguel Michel Farraj; Manal Ali El Ahmar; Louna Karam; Apareka Gamage Dinusha Madhubhashini Perera; Uchini Shermilie Bandaranayake; Miyuru Chandradasa; Jayaweera Arachchige Asela Sampath Jayaweera; Ayesha Lakmali Weerasingha; Ayesha Lakmali Weerasingha; Veranga Kavithri Wickramasinghe; Jayakodi Arachchige Isuru Sohan; Oloyede Oyedibu Oyebayo; Adewuyi A. Tunde; Anwar Jamal Abdulnasir; Jamal Raihan Abdulnasir; Salisu Danjuma Gezawa; Shuaibu Omeiza Salawudeen; Kazeem Bidemi Okesina; Nuraddeen Wada; Yakubu Egigogo Raji; Anthonia Omotola Ishabiyi; Patrick Hosea Olayiwola; Muhammed Olatunbosun Ogunlola; Abdultaofeek Abayomi; Bouya ilyass; Othmane Lamoihi; Achraf Hafdi; Hiba Lazrek; Lahmouz Nouhaila; Irfan Ullah; Asma Nawaz; Khayam ul haq; Abdul Rafay; Kainat Khan; Latif ullah khattak; Noreen Aslam; Ei Cho Lin; Taghreed Saud Almansouri; Maheswaran Warren Archunan; Hasan Hazim Alsaratei; Asif Mahmood; Doaa Hamed Sobeih.

Contributors ATM, MSZ, SME: idea conception, study design. KMR, EMK, YTA, AAS, AZN, LJI, MMA, ATA, AAS, DGH: questionnaire formulation, validation and translation. VZF, ARM, FMS, AE: data curation, analysis and interpretation. AAI, AKA, MA, OMM, EAD, KAE: manuscript drafting. MAD, AIA: study design, analysis planning and supervision. All authors contributed to data collection and all involved investigators reviewed the manuscript and approved it for publication.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Map disclaimer The depiction of boundaries on this map does not imply the expression of any opinion whatsoever on the part of BMJ (or any member of its group) concerning the legal status of any country, territory, jurisdiction or area or of its authorities. This map is provided without any warranty of any kind, either express or implied.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study protocol was approved (R223) by the Institutional Review Board of the Faculty of Medicine, Fayoum University (Fayoum, Egypt).

Consent was obtained at the start of the online questionnaire after explaining the goal and methods of the study. No personal data were collected.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Data are available from the corresponding author upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Ahmed Taher Masoud <http://orcid.org/0000-0003-4304-6781>

Kareem A El-Fas <http://orcid.org/0000-0001-8203-8499>

Mohamed M. Abdel-Daim <http://orcid.org/0000-0002-4341-2713>

Abdelrahman I. Abushouk <http://orcid.org/0000-0003-1399-6487>

REFERENCES

- Zhu N, Zhang D, Wang W, *et al.* A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med Overseas Ed* 2020;382:727–33.
- Worldometers. Covid-19 coronavirus pandemic, 2021. Available: <https://www.worldometers.info/coronavirus/> [Accessed 16 Feb 2021].
- Guan W-J, Ni Z-Y, Hu Y, *et al.* Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;382:1708–20.
- CDC. Symptoms of coronavirus disease 2019 (COVID-19), 2020. Available: <https://www.cdc.gov/coronavirus/2019-ncov/about/symptoms.html> [Accessed 10 Aug 2020].
- CDC. People Who Are at Higher Risk for Severe Illness - Coronavirus, 2020. Available: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/groups-at-higher-risk.html> [Accessed 12 Aug 2020].
- Wiersinga WJ, Rhodes A, Cheng AC, *et al.* Pathophysiology, transmission, diagnosis, and treatment of coronavirus disease 2019 (COVID-19): a review. *JAMA* 2020;324:782.
- Zhong B-L, Luo W, Li H-M, *et al.* Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. *Int J Biol Sci* 2020;16:1745–52.
- Gallè F, Sabella EA, Da Molin G, *et al.* Understanding knowledge and behaviors related to CoVid-19 epidemic in Italian undergraduate students: the EPICO study. *Int J Environ Res Public Health* 2020;17:3481.
- Dkhar SA, Quansar R, Saleem SM, *et al.* Knowledge, attitude, and practices related to COVID-19 pandemic among social media users in J&K, India. *Indian J Public Health* 2020;64:S205–10.
- Azlan AA, Hamzah MR, Sern TJ, *et al.* Public knowledge, attitudes and practices towards COVID-19: A cross-sectional study in Malaysia. *PLoS One* 2020;15:e0233668.
- Mubeen S, Kamal S, Kamal S. Knowledge and awareness regarding spread and prevention of COVID-19 among the young adults of karachi. *J Pak Med Assoc* 2020;70:1–74.
- Bates BR, Villegas Botero A, Grijalva MJ. Knowledge, attitudes, and practices towards COVID-19 among Colombians during the outbreak: an online cross-sectional survey. *J Commun Healthc* 2020;13:262–70.
- Wolf MS, Serper M, Opsasnick L, *et al.* Awareness, attitudes, and actions related to COVID-19 among adults with chronic conditions at the onset of the U.S. outbreak: a cross-sectional survey. *Ann Intern Med* 2020;173:100–9.
- Almofada SK, Alherbisch RJ, Almuhray NA, *et al.* Knowledge, attitudes, and practices toward COVID-19 in a Saudi Arabian population: a cross-sectional study. *Cureus* 2020;12:e8905.
- Reuben RC, Danladi MMA, Saleh DA, *et al.* Knowledge, attitudes and practices towards COVID-19: an epidemiological survey in north-central Nigeria. *J Community Health* 2020;6:1–14.
- Chirwa GC. “Who knows more, and why?” Explaining socioeconomic-related inequality in knowledge about HIV in Malawi. *Scientific African* 2020;7:e00213.
- Chirwa GC, Sithole L, Jamu E. Socio-Economic inequality in comprehensive knowledge about HIV in Malawi. *Malawi Med J* 2019;31:104–11.
- Daniel W. *Biostatistics: a foundation for analysis in the health sciences*. 7th ed. New York: John Wiley & Sons, 1999.
- Wejnert C, Heckathorn DD. Web-Based network sampling: efficiency and efficacy of respondent-driven sampling for online research. *Sociological Methods & Research* 2008;37:105–34.
- Wejnert C, Pham H, Krishna N, *et al.* Estimating design effect and calculating sample size for respondent-driven sampling studies of injection drug users in the United States. *AIDS Behav* 2012;16:797–806.
- Albano L, Matuozzo A, Marinelli P, *et al.* Knowledge, attitudes and behaviour of hospital health-care workers regarding influenza A/H1N1: a cross sectional survey. *BMC Infect Dis* 2014;14:208.
- Hasan F, Khan MO, Ali M. Swine flu: knowledge, attitude, and practices survey of medical and dental students of karachi. *Cureus* 2018;10:e2048.
- Wong LP, Sam I-C. Knowledge and attitudes in regard to pandemic influenza A(H1N1) in a multiethnic community of Malaysia. *Int J Behav Med* 2011;18:112–21.
- Greenhalgh T, Schmid MB, Czypionka T. Face masks for the public during the covid-19 crisis. *BMJ* 2020;2:m1435.
- Eikenberry SE, Mancuso M, Iboi E, *et al.* To mask or not to mask: modeling the potential for face mask use by the general public to curtail the COVID-19 pandemic. *Infect Dis Model* 2020;5:293–308.
- Brooks JT, Butler JC, Redfield RR. Universal masking to prevent SARS-CoV-2 Transmission-The time is now. *JAMA* 2020. doi:10.1001/jama.2020.13107. [Epub ahead of print: 14 Jul 2020].
- WHO. Coronavirus disease (COVID-19) advice for the public: when and how to use masks, 2020. Available: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public/when-and-how-to-use-masks> [Accessed 10 Aug 2020].
- Salman M, Mustafa ZU, Asif N, *et al.* Knowledge, attitude and preventive practices related to COVID-19: a cross-sectional study in two Pakistani university populations. *Drugs Ther Perspect* 2020;1:1–7.
- Kumar J, Katto MS, Siddiqui AA, *et al.* Knowledge, attitude, and practices of healthcare workers regarding the use of face mask to limit the spread of the new coronavirus disease (COVID-19). *Cureus* 2020;12.
- Sari DK, Amelia R, Dharmajaya R. Positive correlation between general public knowledge and attitudes regarding COVID-19 outbreak 1 month after first cases reported in Indonesia. *J Community Health* 2020:1–8.
- Person B, Sy F, Holton K, *et al.* Fear and stigma: the epidemic within the SARS outbreak. *Emerg Infect Dis* 2004;10:358.
- Tao N. An analysis on reasons of SARS-induced psychological panic among students. *Journal of Anhui Institute of Education* 2003;21:78–9.
- Li X, Lu P, Hu L, *et al.* Factors associated with mental health results among workers with income losses exposed to COVID-19 in China. *Int J Environ Res Public Health* 2020;17:5627.
- Holingue C, Badillo-Goicoechea E, Riehm KE, *et al.* Mental distress during the COVID-19 pandemic among US adults without a pre-existing mental health condition: findings from American trend panel survey. *Prev Med* 2020;139:106231.
- Al-Hanawi MK, Angawi K, Alshareef N, *et al.* Knowledge, attitude and practice toward COVID-19 among the public in the Kingdom of Saudi Arabia: a cross-sectional study. *Front Public Health* 2020;8:217.
- Devakumar D, Shannon G, Bhopal SS, *et al.* Racism and discrimination in COVID-19 responses. *Lancet* 2020;395:1194.