BMJ Open Methodology for task-shifting evidence-based psychological treatments to non-licensed/lay health workers: protocol for a systematic review

Kathryn E Kanzler ^{1,2,3}, Lisa Smith Kilpela,^{1,2,4} Jaqueline Pugh,⁵ Luz M Garcini,^{1,5} Christine S Gaspard,⁶ James Aikens,⁷ Erick Reynero,¹ Joel Tsevat,^{1,5,8} Eliot Santana Lopez,^{2,3} Yajaira Johnson-Esparza,³ Amelie G Ramirez,⁹ Erin P Finley^{1,2,5,10}

To cite: Kanzler KE, Kilpela LS, Pugh J, *et al.* Methodology for task-shifting evidence-based psychological treatments to non-licensed/lay health workers: protocol for a systematic review. *BMJ Open* 2021;**11**:e044012. doi:10.1136/bmjopen-2020-044012

► Prepublication history and additional material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-044012>).

Received 19 August 2020
Revised 23 December 2020
Accepted 15 January 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Dr Kathryn E Kanzler;
kanzler@uthscsa.edu

ABSTRACT

Introduction ‘Task-shifting’ or ‘task-sharing’ is an effective strategy for delivering behavioural healthcare in lower resource communities. However, little is known regarding the actual steps (methods) in carrying out a task-shifting project. This paper presents a protocol for a systematic review that will identify steps in adapting an evidence-based psychological treatment for delivery by lay/non-licensed personnel.

Methods and analysis A systematic review of peer-reviewed, published studies involving a non-licensed, non-specialist (eg, community health worker, promotor/a, peer and lay person) delivering an evidence-based psychological treatment for adults will be conducted. Study design of selected articles must include a statistical comparison (eg, randomised controlled trials, quasiexperimental trials, pre-post designs and pragmatic trials). Study selection will follow the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. Databases including PubMed, the Cochrane Library, Cochrane Central Register of Controlled Trials, SCOPUS, Cumulative Index to Nursing and Allied Health Literature, APA PsycInfo and Google Scholar will be searched from 2000 to 2020. Risk of bias will be assessed using the Cochrane Collaboration’s Risk of Bias (RoB 2) tool, and publication bias will be evaluated with the Cochrane GRADE approach. A narrative synthesis will be conducted for all included studies, and a summary table following Proctor’s framework for operationalising implementation strategies will be included. This protocol was developed following the 2015 guidelines of Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols.

Ethics and dissemination This review will analyse data from published studies only; thus, it will not require institutional board review. Findings will be presented at conferences, to the broader community via the Community Health Worker Translational Advisory Board and social media, and the final systematic review will be published in a peer-reviewed journal.

Strengths and limitations of this study

- This protocol describes a planned systematic review to identify best practices for task-shifting evidence-based psychological treatments to non-licensed/lay health workers.
- We will use established operationalised terms to identify and describe implementation strategies.
- Studies will be identified via a thorough search strategy using independent data extraction techniques, with risk of bias mitigation strategies.
- This protocol adheres to the 17-item checklist, Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols.
- The review will only include studies in English and focuses on non-licensed/lay health workers (eg, not nurses or teachers) and identifies only studies examining delivery of psychological treatments (ie, not education or other programming).

INTRODUCTION

Mental health disorders are common worldwide,¹ and although there are evidence-based psychological treatments (EBPTs) that improve health outcomes, most of the people who need treatment do not receive it.² Mental health treatment is often provided by licensed mental health professionals in specialty settings.³ These specialty providers may not be available due to workforce shortages, cost of care and/or access difficulty, especially for lower income and economically challenged populations,⁴ who disproportionately suffer from physical and mental health concerns.^{5 6}

Dissemination and implementation of EBPTs (eg, cognitive-behavioural therapy) to communities in need requires a multidimensional approach with innovative delivery methods.⁷ ‘Task-shifting’ or ‘task-sharing’

is one notable strategy that has emerged over the past two decades, largely in low-income and middle-income countries, as a method for delivering healthcare in lower resource communities. As described by the WHO, with task-shifting, 'specific tasks are moved, where appropriate, from highly qualified health workers to health workers with shorter training and fewer qualifications in order to make more efficient use of the available human resources for health'.⁸

Task-shifting has great promise in improving access to EBPTs. A recent review of 27 trials in low-income and middle-income countries found that task-shifted EBPTs delivered by lay persons in primary care and community settings produced a pooled effect size of 0.49.⁹ Findings from this review indicate that EBPTs can be task-shifted and maintain effectiveness while delivered in non-traditional settings that improve scalability. Lay health workers, such as community health workers or promotor/as, are often trusted members of their communities and perform many important roles, such as delivering physical healthcare^{10 11} and mental healthcare^{12–14} and providing pandemic-related support.^{15 16} Lay health workers help increase access to healthcare in lower resource areas around the world. Therefore, task-shifting EBPTs to lay personnel or paraprofessionals can help reduce disparities in access to evidence-based mental healthcare, thus improving health equity.^{8 17}

Task-shifted psychological treatments

Many different EBPTs have been effectively task-shifted with cultural and contextual adaptations. For example, Patel and colleagues¹⁸ developed a task-shifted treatment programme for moderately severe to severe depression based on Behavioral Activation¹⁹, an established EBPT. Their rigorous randomised controlled trial (RCT) in India showed that the Healthy Activity Program (HAP), delivered by non-specialist lay counselors in primary care, significantly improved patients' levels of depression (moderate effect size) and led to remission in almost two-thirds of patients treated.¹⁸ Likewise, the 'Friendship Bench' programme by Chibanda and colleagues²⁰ was a task-shifting study conducted in Zimbabwe to address depression and common mental disorders. Their treatment programme was based on Problem Solving Therapy,²¹ an established EBPT²², and was delivered by lay health workers in a population with a high prevalence of people living with HIV. Results of their initial non-controlled clinical trial and a later RCT²³ showed that this approach was efficacious in reducing psychological morbidity; a large cluster RCT is now underway.²⁴ In both the HAP and the Friendship Bench studies, lay health workers were trained and supervised by licenced professional specialists (ie, clinical psychologists and/or psychiatrists).

While task-shifting is a recognised method for disseminating EBPTs, the best practice procedural steps for how to task-shift are unclear. A recognised problem in implementation research is that strategies are 'often

inconsistently labeled and poorly described ... lack operational definitions ... and are part of "packaged" approaches whose specific elements are poorly understood' (p. 254).²⁵ Efforts to scale up EBPTs are critical, yet there is no straightforward roadmap for how to implement task-shifting in new settings. This gap in the literature leaves interested stakeholders without clear guidance in deploying this promising implementation strategy. Therefore, we seek to operationalise such strategies used in task-shifting projects. Our research question is: what are the best practices in task-shifting an EBPT for delivery by lay/non-licenced personnel, including methods of (A) adapting the treatment for the new delivery context; (B) training lay personnel; (C) implementing the new treatment protocol and maintaining fidelity; and (D) sustaining the task-shifted programme over time?

Objectives

This protocol outlines our specific methods and planned analyses for a systematic review. This paper adheres to the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P).²⁶ The systematic review seeks to identify specific task-shifting strategies using established definitions as described by Proctor and colleagues.^{25 27} In Proctor's framework for operationalising implementation strategies, there are seven dimensions to consider: the actor, the action, action target, temporality, dose, the implementation outcome affected and the justification. Our review will focus on identifying each of these strategies employed in task-shifting studies, operationally defined in [table 1](#). The most important outcome of this review is to identify best practices for conducting task-shifting implementation projects.

METHODS AND ANALYSIS

Eligibility criteria

Types of studies

Study inclusion criteria

(1) Studies must involve a non-licenced, non-specialist (eg, community health worker, promotor/a, peer and lay person) who is delivering the intervention. (2) Studies must address a 'behavioural health' problem, broadly defined as any psychological/mental health issue (eg., depression, eating disorders and substance use) and/or physical health concern (eg, chronic disease management, health behaviour change, lifestyle changes and adherence) using behavioural/psychological strategies. (3) The treatment components that have been shifted must be derived from an EBPT that has been found efficacious in at least one prior peer-reviewed RCT (eg, cognitive-behavioural therapy, motivational interviewing, behavioural activation and interpersonal psychotherapy). (4) The studies must include a statistical comparison of some kind. The comparator condition must be any of the following: baseline functioning of participants (as in pre-post design), or in an RCT, the control group must be attentional control, a waitlist control, a non-treated

Table 1 Dimensions and definitions of task-shifting strategies for the proposed systematic review based on Proctor et al.'s framework for operationalising implementation strategies

Dimension	Operational definition for the planned systematic review
Actors	Those persons delivering the implementation strategy.
Actions	The methodology used to (A) adapt the treatment for the new delivery context, (B) train the lay personnel to protocol adherence, (C) implement the new treatment protocol with fidelity and (D) sustain the new programme.
Action target	The focus of the task-shifting strategy, including the type of personnel delivering the intervention and the recipients.
Temporality	The order/sequence of the action strategies.
Dose	The amount or intensity of the actions and how much those doses differ from standard/non-shifted EBPTs.
Implementation outcome affected	Identification of which outcome—acceptability, adoption, appropriateness, feasibility, fidelity, implementation cost, penetration and/or sustainability—is being targeted by the actions identified.
Justification	Theoretical, empirical and/or pragmatic rationale for the strategies used to implement their intervention.

EBPTs, evidence-based psychological treatments.

group, a treatment-as-usual group, a group receiving a different form of treatment or a group receiving treatment delivered by an expert provider (e.g. licenced psychologist). Eligible study designs also include RCTs, quasiexperimental trials, pre-post designs, pragmatic trials (eg, using stepped wedge or cluster RCT designs). (5) Eligible studies must report evidence of effectiveness of the task-shifting strategy (ie, clinical outcomes) by using a study design that statistically analyses outcomes using a comparator/control.

Study exclusion criteria

(1) Studies that deliver care solely using a licenced or specialist/non-lay person (eg, nurse and educator); (2) studies focused solely on task-shifting a primarily medical task (eg, HIV treatment and prenatal care); (3) studies reporting psychological/behavioural treatments that have not been previously proven effective as outlined above or not involving treatment (eg, screening only); (4) patient education studies with no behavioural intervention (eg, nutritional information only); (5) studies not involving a comparison; (6) descriptive studies, case reports or exclusively qualitative studies; (7) studies not published in a peer-reviewed journal (eg, dissertations, poster or paper presentations and newsletter articles); (8) books and book chapters; and (9) study protocol publications.

Types of participants

Participant inclusion criteria

(1) Study participants must have received a psychological/behavioural-based (ie, non-pharmacological) intervention for a 'behavioural health' problem, broadly defined as any psychological/mental health problem (eg, depression, eating disorder, parent-child behavioural issues and substance use) and/or physical health concern (eg, chronic disease management, lifestyle changes and adherence); (2) study participants must have received interventions delivered by non-licenced, non-specialists;

and (3) study participants must be adults aged 18 years and older.

Participant exclusion criteria

(1) Patients treated using pharmacological, surgical or medical procedures as the primary intervention tested in the study; and (2) participants treated exclusively by licenced health professionals (eg, physicians and nurses).

Setting and timeframe

Inclusion criteria for setting and timeframe

(1) Task-shifting research studies conducted in high-income, low-income and middle-income countries; and (2) studies conducted in any setting (eg, healthcare or community settings) or region (eg, urban and rural).

Exclusion criteria for setting and timeframe

Studies conducted prior to 2000 (ie, the approximate time when task-shifting was first reported).

Report characteristics

Information sources

The search strategy will be adapted for each of the following sources/databases: PubMed, the Cochrane Library, Cochrane Central Register of Controlled Trials, SCOPUS, Cumulative Index to Nursing and Allied Health Literature, APA PsycInfo and Google Scholar. The search will cover the time frame from January 2000 to July 2020. Peer-reviewed published literature will be sought, and posters, dissertations and presentations; descriptive or protocol articles; books and book chapters; and studies not published in English will be excluded. Unpublished studies will not be sought. The search will be rerun prior to the final analysis, and any further studies identified will be retrieved for inclusion.

Search strategy

The search strategy will be developed and overseen by a medical librarian in consultation with the primary

Table 2 Key word search terms

Key Words	Search terms
Task-shifting	task-shifting; task-sharing; "care sharing"
Lay workers	community health worker; church; community based facilitator; community based organization; health manpower; lay counsellor; lay counselor; lay health worker; non-licensed; nonprofessional; non-specialist; nonspecialists; patient care teams; patient navigator; peer; peer-coach; peer-counsellor; peer-counselor; peer-facilitator; promotor; promotora; promotoras; promotores; self care; self management; shared care; traditional healer; unlicensed
EBPTs	psychological; psychological treatment; psychological intervention; empirically-supported psychological treatment; evidence-based psychological treatment; evidence-based behavioral treatment; evidence-based behavioural treatment; mental health; cognitive behavioral therapy; cognitive behavioural therapy; behavioral therapy; behavioural therapy; interpersonal therapy; acceptance and commitment therapy; psychotherapy; motivational interviewing; interpersonal therapy

EBPTs, evidence-based psychological treatments.

researchers throughout the review (see online supplemental file 1 for PubMed search strategy example). Medical Subject Headings and free-text terms relating to lay health workers and the implementation of task-shifting/sharing will be included (see table 2).

Study screening and selection

Using a coding guide and form, two reviewers will independently search titles and abstracts to remove publications not meeting inclusion criteria. Full texts will be retrieved. Multiple reports of the same study will be linked together (collated) per Cochrane guidelines (see Handbook sec 4.6.2)²⁸ so that the unit of interest is each study, not each article. For example, if a single study was split into separate publications such as a protocol paper, report of the actual study, a qualitative analysis on acceptability and a follow-up, it will be counted as one study, and each of these articles will be searched for relevant data. We will examine prior reviews on the same topic and employ hand-searching of the reference lists for articles identified in the inclusion stage. We will iteratively refine our search strategy to refine the coding guide and form. We will add indexing terms as needed during the preliminary development of the inclusion article guide.

Data extraction

A data extraction chart has been developed by the team to aid in extracting the specific task-shifting steps from the included articles (see online supplemental file 2 for example of study extraction table). Data extracted will follow established definitions as described by Proctor and colleagues.^{25 27} Additional data we anticipate extracting include year of study publication, design, setting, participant demographics, geographic location of study, type of personnel delivering the intervention, demographics of personnel delivering intervention and reported effect sizes.

Two coauthors will independently conduct data extraction, and an additional author will review the data extracted for completion and accuracy. When necessary, consensus will be reached through discussions with an independent fourth author. Missing data will be sought

out by contacting study investigators for unreported data and/or additional details. Data will be recorded in an Excel spreadsheet.

Assessing risk of bias

Reviewers will consider the quality of studies and risk of bias using the Cochrane Collaboration's Risk of Bias (RoB 2) tool,²⁹ incorporating considerations for evaluating psychotherapy outcome research.³⁰ Although the RoB 2 is focused on RCTs, it is applicable to other types of study designs (eg, quasiexperimental and pre-post).²⁹ All domains will be assessed, including bias related to the randomisation process, deviations from intended interventions, missing outcome data, measurement of the outcome, selection of reported result and overall bias. Two authors will review and independently rate risk of bias in each domain as 'low risk of bias', 'some concerns', or 'high risk of bias'. When necessary, disagreements will be resolved by reaching consensus with a third reviewer.

Using the same procedures, we will evaluate publication bias by using the relevant section of the Cochrane GRADE tool³¹ and also mitigate publication bias by searching as extensively as possible using diverse databases, reviewing reference lists and any related systematic reviews. While many systematic reviews grade evidence of a particular treatment, the purpose of this review is to identify specific methods; therefore, we are restricting our search to published literature only.

Data synthesis and analysis

Data synthesis will primarily involve descriptive statistics with tables and graphs to visually communicate findings. Descriptive statistics will be employed to categorise and tally the different types of methodologies used in task-shifting studies, based on standardised language for operationalising implementation strategies,²⁵ to include actors, actions, action targets, implementation outcome affected, temporality, dose and justification. Frequency counts and measures of central tendency will be included. We will report effect sizes for each study. Although we are not grading evidence (ie, incorporating all GRADE

criteria,³² we have developed strict inclusion criteria for rigour and quality as outlined above.

We will consider different subgroups of studies in our review, such as design, population, personnel delivering intervention, location of studies and setting. Although this review is descriptive (no inferential statistical analyses are planned), different tables for each subgroup will be developed. Various subgroups are important to consider separately because differences are anticipated in methodologies depending on each subgroup, for example: design (randomised vs non-randomised trial), population (those with physical health vs mental health concerns), personnel delivering the intervention (community health worker vs other lay personnel), location of studies (low-income and middle-income countries vs high-income countries) and setting (community vs clinical or clinically affiliated).

Patient

Patients/public were not involved in choosing the methods or plans for dissemination of this protocol. However, we will seek feedback on dissemination plans for the forthcoming systematic review from our Community Health Worker Translational Advisory Board.

Ethics and dissemination

This review will analyse data from published studies only; thus, it will not require institutional board review. Any important protocol amendments will be documented in the methods section of the planned systematic review manuscript. Findings will be presented at conferences, to the broader community via the Community Health Worker Translational Advisory Board and social media, and the final systematic review will be published in a peer-reviewed journal.

Author affiliations

¹Center for Research to Advance Community Health (ReACH), University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

²Department of Psychiatry & Behavioral Sciences, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

³Department of Family & Community Medicine, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

⁴Barshop Institute for Longevity and Aging Studies, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

⁵Department of Medicine, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

⁶Briscoe Library, Long School of Medicine, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

⁷Department of Family Medicine, University of Michigan, Ann Arbor, Michigan, USA

⁸Departments of Population Health and Internal Medicine, Dell Medical School, University of Texas, Austin, Texas, USA

⁹Department of Population Health Sciences, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

¹⁰Center for the Study of Healthcare Innovation, Implementation, and Policy (CSHIP), VA Greater Los Angeles Healthcare System, Los Angeles, California, USA

Twitter Kathryn E Kanzler @DrKatieK

Contributors KK: literature search, study design, writing, critical revision and referencing; LSK and JP: study design, writing and critical revision; LMG: study design and writing; CSG: search strategy and writing; ER: literature search and writing; ESL, YJ-E and JA: writing and critical revision; AGR and JT: study design

and critical revision; EPF: study design, literature review, writing and critical revision.

Funding This work was supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health under Award Number K23DK123398 (PI: KK) and by National Institute on Aging of the National Institutes of Health grant number K76AG060003-01A1 (PI: LSK).

Disclaimer The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Kathryn E Kanzler <http://orcid.org/0000-0001-7253-7067>

REFERENCES

- 1 Kessler RC, Aguilar-Gaxiola S, Alonso J, *et al*. The global burden of mental disorders: an update from the who world mental health (WMH) surveys. *Epidemiol Psychiatr Soc* 2009;18:23–33.
- 2 Wang PS, Aguilar-Gaxiola S, Alonso J, *et al*. Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the who world mental health surveys. *Lancet* 2007;370:841–50.
- 3 Kazdin AE, Blase SL. Rebooting psychotherapy research and practice to reduce the burden of mental illness. *Perspect Psychol Sci* 2011;6:21–37.
- 4 Ogbeide SA, Landoll RR, Nielsen MK, *et al*. To go or not go: patient preference in seeking specialty mental health versus behavioral consultation within the primary care behavioral health consultation model. *Fam Syst Health* 2018;36:513–7.
- 5 Braveman PA, Cubbin C, Egerter S, *et al*. Socioeconomic disparities in health in the United States: what the patterns tell us. *Am J Public Health* 2010;100 Suppl 1:S186–96.
- 6 Pollack CE, Cubbin C, Sania A, *et al*. Do wealth disparities contribute to health disparities within racial/ethnic groups? *J Epidemiol Community Health* 2013;67:439–45.
- 7 Kazdin AE, Rabbitt SM. Novel models for delivering mental health services and reducing the burdens of mental illness. *Clin Psychol Sci* 2013;1:170–91.
- 8 World Health Organization. *Task shifting: rational redistribution of tasks among health workforce teams: global recommendations and guidelines*. Geneva: World Health Organization, 2007.
- 9 Singla DR, Kohrt BA, Murray LK, *et al*. Psychological treatments for the world: lessons from low- and middle-income countries. *Annu Rev Clin Psychol* 2017;13:149–81.
- 10 Norris SL, Chowdhury FM, Van Le K, *et al*. Effectiveness of community health workers in the care of persons with diabetes. *Diabet Med* 2006;23:544–56.
- 11 Scott K, Beckham SW, Gross M, *et al*. What do we know about community-based health worker programs? A systematic review of existing reviews on community health workers. *Hum Resour Health* 2018;16:39.
- 12 Barnett ML, Gonzalez A, Miranda J, *et al*. Mobilizing community health workers to address mental health disparities for underserved populations: a systematic review. *Adm Policy Ment Health* 2018;45:195–211.
- 13 Barnett ML, Lau AS, Miranda J. Lay health worker involvement in evidence-based treatment delivery: a conceptual model to address disparities in care. *Annu Rev Clin Psychol* 2018;14:185–208.

- 14 Hoeft TJ, Fortney JC, Patel V, *et al.* Task-Sharing approaches to improve mental health care in rural and other low-resource settings: a systematic review. *J Rural Health* 2018;34:48–62.
- 15 Bhaumik S, Moola S, Tyagi J, *et al.* Community health workers for pandemic response: a rapid evidence synthesis. *BMJ Glob Health* 2020;5:e002769.
- 16 Boyce MR, Katz R. Community health workers and pandemic preparedness: current and prospective roles. *Front Public Health* 2019;7:62.
- 17 Kazdin AE. *Innovations in psychosocial interventions and their delivery: leveraging cutting-edge science to improve the world's mental health*. Oxford: Oxford University Press, 2018.
- 18 Patel V, Weobong B, Weiss HA, *et al.* The healthy activity program (HAP), a lay counsellor-delivered brief psychological treatment for severe depression, in primary care in India: a randomised controlled trial. *Lancet* 2017;389:176–85.
- 19 Dimidjian S, Hollon SD, Dobson KS, *et al.* Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the acute treatment of adults with major depression. *J Consult Clin Psychol* 2006;74:658–70.
- 20 Chibanda D, Mesu P, Kajawu L, *et al.* Problem-Solving therapy for depression and common mental disorders in Zimbabwe: piloting a task-shifting primary mental health care intervention in a population with a high prevalence of people living with HIV. *BMC Public Health* 2011;11:828.
- 21 Nezu AM, Nezu CM, Perri MG. *Problem-solving therapy for depression: theory, research, and clinical guidelines*. Hoboken: John Wiley & Sons, 1989.
- 22 Bell AC, D'Zurilla TJ. Problem-solving therapy for depression: a meta-analysis. *Clin Psychol Rev* 2009;29:348–53.
- 23 Chibanda D, Weiss HA, Verhey R, *et al.* Effect of a primary care-based psychological intervention on symptoms of common mental disorders in Zimbabwe: a randomized clinical trial. *JAMA* 2016;316:2618–26.
- 24 Chibanda D, Bowers T, Verhey R, *et al.* The Friendship bench programme: a cluster randomised controlled trial of a brief psychological intervention for common mental disorders delivered by lay health workers in Zimbabwe. *Int J Ment Health Syst* 2015;9:21.
- 25 Proctor EK, Powell BJ, McMillen JC. Implementation strategies: recommendations for specifying and reporting. *Implement Sci* 2013;8:139.
- 26 Moher D, Shamseer L, Clarke M, *et al.* Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
- 27 Kirchner JE, Waltz TJ, Powell BJ. Implementation strategies. dissemination and implementation research in health: translating science to practice 2017;2:245–66.
- 28 Cochrane. *Cochrane Handbook for systematic reviews of interventions version 6.0*. London: Cochrane, 2019.
- 29 Sterne JAC, Savović J, Page MJ, *et al.* Rob 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366:l4898.
- 30 Munder T, Barth J. Cochrane's risk of bias tool in the context of psychotherapy outcome research. *Psychother Res* 2018;28:347–55.
- 31 Ryan R, Hill S. How to GRADE the quality of the evidence. Cochrane Consumers and Communication Group. Version 3.0 December 2016. Available: <http://cccrgr.cochrane.org/author-resources>
- 32 Schünemann H, Brożek J, Guyatt G, eds. *Handbook for grading the quality of evidence and the strength of recommendations using the GRADE approach*, 2013. <https://gdt.gradepr.org/app/handbook/handbook.html>