BMJ Open National, clinical cohort study of late effects among survivors of acute lymphoblastic leukaemia: the ALL-STAR study protocol

Liv Andrés-Jensen ¹, Mette Tiedemann Skipper ², Kristian Mielke Christensen,¹ Pia Hedegaard Johnsen,³ Katrine Aagaard Myhr,⁴ Martin Kaj Fridh,¹ Kathrine Grell,^{1,5} A. M. L. Pedersen,⁶ Sune Leisgaard Mørck Rubak,² Martin Ballegaard,^{7,8} Arne Hørlyck,⁹ Rikke Beck Jensen,¹⁰ Trine-Lise Lambine,¹¹ Kim Gjerum Nielsen,^{1,8} Ruta Tuckuviene,¹² Peder Skov Wehner,¹³ Birgitte Klug Albertsen,^{2,14} Kjeld Schmiegelow,^{1,8} Thomas Leth Frandsen¹

To cite: Andrés-Jensen L, Skipper MT, Mielke Christensen K, *et al.* National, clinical cohort study of late effects among survivors of acute lymphoblastic leukaemia: the ALL-STAR study protocol. *BMJ Open* 2021;11:e045543. doi:10.1136/bmjopen-2020-045543

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-045543>).

Received 06 October 2020
Revised 15 December 2020
Accepted 07 January 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Liv Andrés-Jensen;
liv.andres-jensen.02@regionh.dk

ABSTRACT

Introduction More than 90% of patients diagnosed with childhood acute lymphoblastic leukaemia (ALL) today will survive. However, half of the survivors are expected to experience therapy-related chronic or late occurring adverse effects, reducing quality of life. Insight into underlying risk trajectories is warranted. The aim of this study is to establish a Nordic, national childhood ALL survivor cohort, to be investigated for the total somatic and psychosocial treatment-related burden as well as associated risk factors, allowing subsequent linkage to nation-wide public health registers.

Methods and analysis This population-based observational cohort study includes clinical follow-up of a retrospective childhood ALL survivor cohort (n=475), treated according to a common Nordic ALL protocol during 2008–2018 in Denmark. The study includes matched controls. Primary endpoints are the cumulative incidence and cumulative burden of 197 health conditions, assessed through self-report and proxy-report questionnaires, medical chart validation, and clinical examinations. Secondary endpoints include organ-specific outcome, including cardiovascular and pulmonary function, physical performance, neuropathy, metabolic disturbances, hepatic and pancreatic function, bone health, oral and dental health, kidney function, puberty and fertility, fatigue, and psychosocial outcome. Therapy exposure, acute toxicities, and host genome variants are explored as risk factors.

Ethics and dissemination The study is approved by the Regional Ethics Committee for the Capital Region in Denmark (H-18035090/H-20006359) and by the Danish Data Protection Agency (VD-2018–519). Results will be published in peer-reviewed journals and are expected to guide interventions that will ameliorate the burden of therapy without compromising the chance of cure.

INTRODUCTION

Five-year overall survival rates for childhood acute lymphoblastic leukaemia (ALL) has

Strengths and limitations of this study

- The first population-based, uniformly treated, Nordic childhood and young adult acute lymphoblastic leukaemia (ALL) survivor cohort to be systematically evaluated for treatment-related organ-wide morbidity and psychosocial impact.
- Additional strengths include use of matched controls, systematic subjective and objective assessments of outcome and alignment of adverse effects definitions with established classification systems.
- Estimation of the cumulative burden includes both persistent and recurrent events, thereby describing the total disease burden more comprehensively than traditional measures, such as cumulative incidence.
- Initial cross-sectional analyses are limited by a follow-up of 3.5–14 years, acknowledging that the full burden of ALL and leukaemic therapy may not yet have emerged.
- Self-report and systematic clinical data will supplement data from the Danish nationwide public demographic, socioeconomic and health registries.

climbed from less than 10% to above 90% during the last seven decades as a result of chemotherapy intensification, refined risk stratification and improved supportive care.^{1–3} However, cure comes at a cost.⁴ All patients experience acute, but transient toxicities during therapy (eg, infections, mucositis and peripheral neuropathy), and half of the patients are burdened by at least one severe, potentially life-threatening organ toxicity, such as acute pancreatitis, thromboembolism, or severe neurotoxicity.^{5–8} Acute organ damage resulting from leukaemia and/or therapy may persist (eg, insulin-dependent

diabetes)⁹ or emerge (eg, osteonecrosis)¹⁰ several years into survival as late effects. The 30-year-old survivor will have experienced an average of 5.4 health conditions, and half of survivors face at least one chronic health condition 20 years from ALL diagnosis, both measures significantly exceeding those among siblings and community controls.^{11 12} This disproportion in morbidity rate seems to amplify with increasing age.¹² Certain therapeutic exposures have been reduced over time (eg, reduction in cardiotoxic anthracycline exposure and limited use of cranial irradiation in first-line therapy protocols),^{13 14} which has decreased the risk of early mortality and lowered the prevalence of treatment-related cancers, severe cognitive deficits, hypothalamic-pituitary dysfunction and immunological disease.^{12 14} However, the number of health conditions per individual survivor is still double of that in controls, now dominated by endocrine system disorders, and musculoskeletal, neurological and cardiovascular conditions,¹² maintaining risk of impaired physical, cognitive and emotional functions, socioeconomic achievements and health-related quality of life.^{11 15–18} Therapy exposure and genetic susceptibility are considered the two major risk factors predicting late morbidity.¹⁹ Risk of specific late effects are associated with cumulative chemotherapy exposures,^{20–29} but no safe lower doses have been established. Specific genetic variants have been linked with survivor obesity, reduced bone mineral density (BMD), cardiotoxicity, skeletal muscle dysfunction and neurocognitive dysfunction,^{30–33} however, clinical recommendations await harmonisation of outcomes, larger sample sizes and replication in independent cohorts.³⁰

To date, many late effects studies are observational and focus on one or few organ systems, cross-sectional in design, or emerge from single centres with resulting risk of selection bias and limited sample power. A few very large (>10 000 patients), multi-institutional childhood cancer survivor cohort studies exist, such as the US Childhood Cancer Survivor Study,³⁴ the Nordic Adult Life after Childhood Cancer in Scandinavia study³⁵ and the British Childhood Cancer Survivor Study.³⁶ Most studies emerging from these cohorts rely on self-reported and register-based data, while few collect systematic clinical data, none of which include Nordic survivors.³⁷

During the 10-year period of 2008–2018, patients diagnosed with Philadelphia chromosome-negative (Ph⁻) ALL at age 1–45 years in the Nordic and Baltic countries, have been treated according to the Nordic Society of Paediatric Haematology and Oncology (NOPHO) ALL2008 protocol described elsewhere.^{3 5 38} Two thirds of this cohort were below 18 years of age at diagnosis. The aim of the Acute Lymphoblastic Leukemia Survivor Toxicity And Rehabilitation (ALL-STAR) study is to establish a national, Danish NOPHO ALL2008 survivor cohort, to be assessed for the total somatic and psychosocial treatment-related burden, using both self-report and systematic clinical data that will supplement data from the

Danish nationwide public demographic, socioeconomic and health registries.³⁹

METHODS

Study aims

Primary aim

To quantify the burden of health conditions occurring during the first decade of survivorship among survivors of childhood and young adult ALL.

Secondary aims

To describe, in detail, organ-specific health among survivors as compared with matched controls and normative data; and to investigate host genome variants, therapy exposure and acute toxicities as risk factors for organ-specific late effects and for the overall cumulative burden. Furthermore, to establish a platform for life-long follow-up in this survivor cohort, facilitating longitudinal studies and linkage to nationwide public, socioeconomic and health registries.

Study design

The ALL-STAR study is a national, Danish, population-based, observational cohort study, including prospective clinical follow-up in a retrospective childhood and young adult ALL survivor cohort. This present protocol primarily focuses on the initial, cross-sectional phase of the project.

Participants

Study population

Survivors treated in Denmark according to the NOPHO ALL2008 protocol and minimum 1 year from therapy cessation (3.5 years from ALL diagnosis) are eligible. This corresponds to all patients (n=475) aged 1–45 years at diagnosis with Ph⁻ B-cell precursor or T-cell ALL, diagnosed between July 2008 and October 2018 (the complete NOPHO ALL2008 protocol period). A control group consisting of age- and sex-matched individuals (1:1) is recruited for the study. First-degree relatives to survivors and individuals with previous or ongoing cancer, previous or ongoing chemotherapy, and/or previous or ongoing radiation therapy are excluded as controls. Study recruitment for the initial cross-sectional studies opened in February 2019 and will terminate in September 2022, allowing for the entire survivor cohort to meet the inclusion criteria.

Recruitment

Eligible survivors are identified through the NOPHO ALL2008 registry. Contact information is obtained through medical charts and provided by treating physicians to the research team. Written information is sent to parents if survivor's age is <15 (addressed to survivor and parents), to parents and the survivor if aged 15–17.9, and to survivors if aged ≥18 years. Contact is obtained through subsequent phone calls. All survivors and/or families are invited to an on-site meeting for further information

before consent. Controls are recruited via participating survivors (or parents) who appoint and invite friends or relatives to participate. Potential controls (or parents) contact the research team, with subsequent written and oral information provided as described for survivors, before possible consent.

Endpoints

Primary endpoints and hypothesis

The two primary endpoints are cumulative incidence and cumulative burden of 197 health conditions presented in [table 1](#).

We hypothesise that the cumulative incidence and cumulative burden are significantly higher among survivors of childhood and young adult ALL already during the first decade of survival, when compared with age- and sex-matched controls, and normative data.

The measure of cumulative burden considers occurrence of multiple health conditions and recurrent events, thereby encompassing the total disease burden more comprehensively than cumulative incidence, which includes only first occurrence of an event.^{40 41} The 197 conditions investigated in the ALL-STAR study are defined and graded according to the St Jude Children's Research Hospital modification of the National Cancer Institute's Common Terminology Criteria for Adverse Events (CTCAE; V.4.03),⁴² developed to optimise characterisation of long-term and late-onset health conditions among childhood cancer survivors.⁴³ Ten additional health conditions (generalised muscle weakness, reduced cardiopulmonary fitness, sarcopenic obesity and seven oral/dental conditions) have been added. Definitions and grading criteria for all conditions are provided in online supplemental table 1.

Secondary endpoints

Secondary endpoints include: (1) organ-specific outcome, including metabolic syndrome (MetS) and body composition, hepatic function, pancreatic function, bone health, cardiac function, pulmonary function, peripheral and autonomic neuropathy, physical performance, renal function, pubertal timing, fertility, oral and dental health, and (2) psychosocial outcome, including emotional distress, fatigue and health-related quality of life.

An overview of the ALL-STAR study is provided in [figure 1](#).

Data collection

Data are collected through questionnaires and clinical evaluations. An overview of ALL-STAR source data used for evaluation of the 197 health conditions is provided in online supplemental table 1.

Questionnaire data

An electronic questionnaire is sent to participants prior to clinical examinations. The questionnaire is adjusted according to self-report (age ≥ 15 years), proxy report (age < 18), age, sex, and status as survivor or control. Items include demographics, use of healthcare services,

medicine and nutritional supplements, medical history (including age at diagnosis of a confirmed condition), pubertal status and fertility, familial dispositions, health behaviour, mental health, quality of life and socioeconomic status. An overview of questionnaire versions and content is provided in online supplemental table 2.

Clinical data

All clinical examinations are performed at either Copenhagen University Hospital or Aarhus University Hospital, as requested by the participant. Examinations are performed during 10 hours in a single day. The examination programme and description of clinical investigators are provided in online supplemental table 3.

Evaluation of metabolic syndrome and body composition

Endpoints include MetS, insulin sensitivity, lipid profile, abdominal circumference, body mass index (BMI) for adults and BMI standard deviation (SD) scores for children based on national reference material,^{44 45} lean body mass, and fat mass. Adult MetS is defined according to the National Cholesterol Education Program Adult Treatment Panel criteria,⁴⁶ and includes central obesity, dyslipidaemia, hypertension and elevated fasting glucose. Paediatric MetS is defined using the same measures.⁴⁷ Questionnaire items include metabolic disease and family disposition, prescription medicine, nutritional supplements and lifestyle (diet, level of physical activity, and sedentary behaviour). Clinical investigations include anthropometrics (weight, height and abdominal circumference) and blood pressure (described in cardiovascular section). Fasting blood samples include glucose, glycated haemoglobin, insulin, proinsulin c-peptide, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, very low-density lipoprotein cholesterol, triglycerides, glucagon and leptin. Homeostatic Model Assessment for Insulin resistance score is calculated as $\text{fasting plasma glucose (mmol/L)} \times \text{fasting plasma insulin (}\mu\text{U/L)} / 22.5$.⁴⁸ Body composition (fat and lean body mass, and android/gynoid fat distribution) is assessed with dual-energy X-ray absorptiometry (DXA) (Lunar Prodigy, GE Healthcare, Wisconsin, USA; or Hologic Horizon A, Hologic, Marlborough, Massachusetts, USA), and is adjusted for sex and pubertal stage.

Evaluation of hepatic function and gallbladder

Endpoints include signs of hepatocellular damage, impaired liver synthesis, iron overload, hepatic fibrosis, portal hypertension and gall bladder disease (cholestasis, cholangitis and cholelithiasis). Questionnaire items include symptoms of hepatic disease (fatigue, abdominal pain, nausea, jaundice, discoloured urine and itchy skin), hepatic diagnoses, alcohol consumption and use of prescription medicine. Fasting blood samples include alanine transaminase, aspartate transaminase, alkaline phosphatase, gamma-glutamyl transferase, bilirubin, lactate dehydrogenase, prothrombin time, international normalised ratio, albumin, ferritin, transferrin saturation,

Table 1 Health conditions investigated in the ALL-STAR study

Auditory-hearing	Gastrointestinal	Infections	Neurological	Pulmonary
Cholesteatoma	Bowel perforation	Bronchial/lung infection C/R	Autonomic dysfunction	Asthma
Hearing loss	Coeliac disease	Endocarditis	Cavernoma	COPD
Tinnitus	Constipation	Gastrointestinal infection	Cerebellar dysfunction	Epistaxis, C/R
Vertigo	Dysphagia	Genitourinary infection	Cerebral necrosis	Obstructive sleep apnoea
	Enterocolitis	Hepatitis B, chronic	Cerebrovascular accident	Obstructive ventilatory defect
			Cerebrovascular disease	Pleural space disorders
Cardiovascular	Oesophageal varices	Hepatitis C, chronic		
Aortic root aneurysm	Oesophagitis	HIV infection	Cranial nerve disorder	Pneumonitis
Arteriovenous malformation	Faecal incontinence	Lymphatic infection	Dysarthria	Pulmonary diffusion defect
Atrioventricular heart block	Gastritis/duodenitis	Meningoencephalitis	Generalised muscle weakness	Pulmonary embolism
Bradycardia, sinus	GORD	Osteomyelitis	Headaches, C/R	Respiratory tract haemorrhage
				Restrictive ventilatory defect
Cardiopulm. fitness, reduced	Gastrointestinal fistulas	Otitis media, C/R	Hydrocephalus	Tracheal aspiration
Conduction abnormalities	Gastrointestinal haemorrhage	Pelvic inflammatory disease	Hydrosyngomyelia	Tracheal stenosis
Congestive heart failure	Gastrointestinal necrosis	Pharyngitis/tonsillitis, C/R	Intracranial haemorrhage	
Coronary artery disease	Gastrointestinal obstruction	Sinusitis, C/R	Movement disorders	
Cor pulmonale	Gastrointestinal strictures	Soft tissue infection	Multiple sclerosis	Psychiatric
Dysrhythmia	Gastrointestinal ulcer		Narcolepsy	Anxiety
Heart valve disorder	Gastroparesis syndrome	Musculoskeletal	Nerve root disorder	Depression
High total cholesterol	Malabsorption syndrome	Amputation	Neurogenic bladder	Other psychiatric disorders
Hypertension	Pancreatic insufficiency	Arthralgia	Neurogenic bowel	
Hypertriglyceridaemia	Pancreatitis	Arthritis	Neuromuscular disorders	Renal/urinary
LV systolic dysfunction	Proctitis	Bone mineral density deficit	Paralytic disorder	Acute kidney injury
Pericarditis		Hernia	Peripheral motor neuropathy	Chronic haematuria
Prolonged (QTc) interval	Hepatobiliary	Intervertebral disc disorder	Peripheral sensory neuropathy	Chronic kidney disease
Pulmonary hypertension	Cholecystitis/cholelithiasis	Kyphosis	Pseudomeningocele	Incontinence
Raynaud phenomenon	Fibrosis/cirrhosis	Limb length discrepancy	Pseudotumour cerebri	Obstructive uropathy
RV systolic dysfunction	Hepatic failure	Osteonecrosis	Seizures	Urinary bladder dysfunction
Tachycardia, sinus	Hepatopathy	Palatal defects, acquired		Urinary tract calculi
Thromboembolic event	Portal hypertension	Scoliosis	Ocular/visual	Vesicoureteral reflux, acquired
			Cataract	
Vascular disease	Veno-occlusive disease	Skeletal spine disorder	Diplopia	Reproductive/genital
		Slipped capital femoral epiphysis	Dry eye syndrome	Abnormal sperm concentration
Endocrine	Haematological			
Abnormal glucose metabolism	Anaemia	Oral/dental	Eyelid function disorder	Cervical dysplasia
Adrenal insufficiency	Coagulopathy	Dental caries	Glaucoma	Dysfunctional uterine bleeding
				Endometriosis
Adult GH deficiency	Iron overload	Dental erosion	Ocular disease, non-infectious	Erectile dysfunction
Childhood GH deficiency	Neutropaenia	Gingivitis	Ocular surface disease	Genitourinary adhesions
Diabetes insipidus	Polycythaemia	Periodontitis	Photophobia	Hypogonadism, central
GH excess	Thrombocytopenia	Sialadenitis, acute	Phthisis bulbi	Leydig cell insufficiency
Hyperparathyroidism	Thrombocytosis	Sialadenitis, chronic	Retinal detachment	Polycystic ovarian syndrome
Hyperprolactinaemia		Salivary gland dysfunction	Retinopathy	Precocious puberty
Hyperthyroidism	Immunological	TMJ disorder	Strabismus	Primary ovarian failure
Hypoparathyroidism	Autoimmune disorders	Dental maldevelopment	Visual acuity, reduced (OD)	Prostatic hypertrophy, benign
Hypothyroidism	Graft-versus-host disease		Visual acuity, reduced (OS)	
				Vaginal fistula
Overweight/obesity	Immunodeficiency	Neoplasms	Visual field deficit	Vaginal stenosis
Sarcopenic obesity		Benign neoplasms		
SIADH		Malignant neoplasms		
Underweight				

ALL-STAR, Acute Lymphoblastic Leukemia Survivor Toxicity And Rehabilitation; COPD, chronic obstructive pulmonary disease; CR, chronic/recurrent; GH, growth hormone; GORD, gastro-oesophageal reflux disease; LV, left ventricle; OD, right eye; OS, left eye; QTc, corrected QT interval; RV, right ventricle; SIADH, syndrome of inappropriate antidiuretic hormone secretion; TMJ, temporomandibular joint.

Figure 1 ALL-STAR study overview. *Estimated survivors from 475 patients and overall survival rate of 90%. **Estimated number of participants based on minimum recruitment rate of 60%. ALL-STAR, Acute Lymphoblastic Leukemia Survivor Toxicity And Rehabilitation; NOPHO, Nordic Society of Paediatric Haematology and Oncology.

alfa-1 antitrypsin, platelets and immunoglobulins. Serum is stored for evaluation of viral antibodies (hepatitis A, B and C, cytomegalovirus and Epstein-Barr virus) in case of elevated transaminases. Hepatic ultrasound is performed using a convex, abdominal probe (C1-5, GE Logiq E9 or E10, GE Healthcare, Chicago, USA). The liver, gall-bladder, gall ducts, liver artery and spleen are evaluated for appearance and size. Portal vein flow is measured using Doppler ultrasound. Ultrasound 2D shear wave elastography is performed on the right liver lobe for evaluation of parenchymal stiffness and fibrosis.⁴⁹

Evaluation of pancreatic function

Endpoints include pancreatic disease, endocrine and exocrine function, and morphological appearance. Questionnaire items include pancreatic symptoms and disease, diet and alcohol consumption. Fasting blood samples include pancreatic amylase, lipase, glucose, glycosylated haemoglobin, insulin and proinsulin c-peptide. Faecal elastase-1 is evaluated in survivors only and associated with occurrence of asparaginase-associated pancreatitis during ALL therapy. Pancreatic ultrasound is performed using a convex, abdominal probe (C1-5, GE Logiq E9 or E10, GE Healthcare, Chicago, USA) for anterior-posterior measurement of caput and corpus, and for evaluation of morphological changes (tumour, inflammation, oedema, pseudocysts and ductal ectasia).

Evaluation of bone mineral density

Endpoints include bone mineral density (BMD), bone fractures and biomarkers of bone metabolism. Questionnaire items include bone symptoms and joint symptoms, history of fractures and low BMD, age of menarche, hormone supplements, calcium and vitamin D intake, and

family history of bone disease. Biomarkers include ionised calcium, phosphate, alkaline phosphatase, parathyroid hormone and 25-OH vitamin D. Bone mineral content, bone area, and BMD is determined by DXA of total body, and lumbar spine (L1-L4) and dual-femur scans. Adult bone mineral values are analysed as age-specific and sex-specific SD and expressed as z-scores and T-scores. Paediatric bone mineral values are analysed as age-specific and sex-specific SD and expressed as z-scores. All paediatric bone mineral values are corrected for height.

Evaluation of cardiovascular function

Endpoints include left and right ventricular systolic and diastolic dysfunction, ventricular volumes and left ventricular mass, myocardial fibrosis, myocardial iron deposits, atrial thrombosis, conduction abnormalities and hypertension. Questionnaire items include cardiovascular symptoms, diagnoses, prescription medicine and familial disposition to cardiovascular disease. Automated oscillometric blood pressure is measured in all, and manual auscultatory systolic blood pressure is measured in children (<18 years). Cardiac biomarkers include N-terminal pro b-type natriuretic peptide and troponin-T. Cardiac electrical activity is assessed with a 12-lead ECG. Imaging includes speckle tracking echocardiography and cardiac MRI. Echocardiography is performed using a Vivid E95 ultrasound scanner (GE Vingmed Ultrasound AS, Horten, Norway) to obtain measurements of left ventricular output and volumes, as well as global longitudinal systolic strain. An intravenous contrast agent (SonoVue)⁵⁰ is used to improve imaging. Cardiac MRI is performed using a whole body 1.5T scanner for assessment of left ventricular systolic function, volumes, and

mass. Multiparametric characterisation, including T1, T2, T2* and mapping is performed for visualisation of possible myocardial oedema and fibrosis, and to evaluate myocardial iron content.^{51 52} Intravenous gadolinium-containing contrast (0.15 mmol/kg Gadovist) is used for assessment of extracellular volume fraction in survivors (not controls).

Evaluation of pulmonary function

Endpoints include function of conductive and acinar airways, resistance of airways, alveolar volumes, alveolar membrane diffusion, lung clearance index, capillary volume, total diffusion capacity, spirometric volumes, and flows and reversibility. Questionnaire items include respiratory symptoms, pulmonary diagnoses, smoking and use of prescription medicine. Prebronchodilator and post-bronchodilator spirometry is performed using a Jaeger MasterScreen Bodybox and Jaeger Vyntus Spiro (CareFusion, Hochberg; Vyair Medical, Bayern) for measurement of forced expiratory volume during the first second, forced vital capacity (FVC), and forced expiratory flow between 25% and 75% of FVC. Impulse oscillometry, including reversibility test, is performed at University Hospital of Aarhus using a Jaeger MasterScreen Bodybox and Jaeger Vyntus IOS system (CareFusion, Hochberg; Vyair Medical, Bayern) to evaluate lung volumes, resistance and elasticity of the lung. Nitrogen multiple breath washout is performed using an Exhalyzer D, N2 and SF6 option (Eco Medics AG, Dürnten, Switzerland) to assess Lung Clearance Index, including indexes of acinar and conductive airways ventilation heterogeneity. Single breath diffusion capacity for carbon monoxide and nitrogen oxide, corrected for haemoglobin, is performed using a Jaeger Vyntus Body (CareFusion, Hochberg) for real-time single breath diffusion to evaluate diffusion membrane capacity and pulmonary capillary volume. All pulmonary measures are compared with normative data^{53–58} and with data from the matched controls.

Evaluation of peripheral and autonomic neuropathy

Endpoints include sensory nerve, motor nerve and autonomic nerve dysfunction. Peripheral neuropathy is assessed according to recommendations from the American Academy of Neurology, the American Association of Electrodiagnostic Medicine and the American Academy of Physical Medicine and Rehabilitation,⁵⁹ and using the Total Neuropathy Score, which includes self-report of sensory, motor and autonomic symptoms, pin sensibility, vibration sensibility (quantitative threshold using Vibrometer type IV (Somedic Electronics, Solna, Sweden) and Biothesiometer model PVD-LP (Bio-Medical Instruments, Ohio, USA), muscle strength (Medical Research Council grade 5 score), deep tendon reflexes, and amplitude of sural and peroneal nerves.⁶⁰ Abnormal values corresponding to quantitative vibration threshold are calculated as percentages relative to the upper limit of normal values based on data from matched ALL-STAR controls and previously published normative data.^{61–63} Electroneurography is performed using standard surface recording

techniques (Neuroline 715, Ambu). A G3 Keypoint platform (Dantec, Natus, USA) is used for amplification, filtration and storing of signals (motor: 2 Hz to 10 kHz; sensory: 20 Hz to 10kHz). Electroneurography includes unilateral evaluation of sensory nerve action potential from the sural nerve, compound motor action potential from the peroneal nerve, the distal motor latency in the peroneal nerve and both sensory and motor nerve conduction velocities. Recorded values are compared with ALL-STAR control values and with unpublished national (Danish age-specific and gender-specific reference values), and reported as SD from the expected mean (z-scores). Autonomic neuropathy is evaluated with the validated 31-item composite autonomic symptom questionnaire, COMPASS 31,^{64 65} which assesses orthostatic intolerance, vasomotor, secretomotor, gastrointestinal, bladder and pupillomotor function. Objective evaluation of autonomic dysfunction is performed by analysis of heart rate variability (HRV) obtained from a 21 min Holter recording (13 min reclined resting period followed by shift to standing position with continued ECG recording for 8 min). The RR-interval data set derived from the ECG segments is analysed for heart rate and HRV. Analysis of HRV is performed in both the time and frequency domain and changes elicited by the standing position is calculated.

Evaluation of physical performance

Endpoints include cardiorespiratory fitness, level of physical activity, physical function and muscle strength. Questionnaire items include levels of sedentary behaviour and of physical activity, which is objectified with a 7-day continuous accelerometer measurement (ActiGraph model GT3X+, ActiGraph, Pensacola, Florida, USA). Cardiorespiratory fitness is evaluated using an electronically braked cycle ergometer (Lode Corival Pediatric or Monark Ergonomic 839 E) following a modified Godfrey protocol.^{66 67} Ventilation and gas exchange data are determined breath-by-breath (INNOCOR ergo-spirometry-system, INNO00010, Innovision, DK-5260 Odense, Denmark or Jaeger Master Screen Vyntus CPX and JLAB software package). Peak oxygen uptake (VO₂ peak) is defined as the highest mean over 60 s. Results are reported as SD from the expected mean (z-scores), derived from the age-matched and sex-matched ALL-STAR controls. Muscle strength is evaluated as isometric muscle strength (knee extensor, upper body (Gym 2000) with a strain gauge (US2A100 kg, Hottinger, Germany) and custom-made amplifier), and hand grip strength (Saehan hand dynamometer, Glanford Electronics, Scunthorpe). Dynamic muscle strength is evaluated by the counter movement jump test (FP4, HUR Labs Oy, Tampere, Finland) and physical functional tests (sit-to-stand 30 s and 60 s⁶⁸ and timed up and go).⁶⁹

Evaluation of kidney function

Endpoints include kidney function and kidney size. Questionnaire items include urinary tract symptoms and disease, and use of prescription medicine. Fasting blood samples, include creatinine, urea, cystatin C, sodium, potassium,

phosphate, albumin, ionised calcium and parathyroid hormone. Estimated glomerular filtration rate is evaluated using creatinine in adults,⁷⁰ and using creatinine, urea and cystatin in children.⁷¹ Proteinuria is determined by the albumin-creatinine ratio in a morning urine specimen. A spot urine dip stick test is performed to evaluate haematuria and urine pH. Kidney size is evaluated by ultrasonography using a convex abdominal probe (C1-5, GE Logiq E9 or E10, GE Healthcare, Chicago, USA) as longitudinal anterior-posterior length, width and cross-sectional anterior-posterior length, and abnormal appearance (eg, hydronephrosis, cysts or tumours) is registered. Kidney size is related to body weight, height and lean mass.

Evaluation of pubertal status and gonadal function

Pubertal stage is evaluated in all participants aged >8 years at time of evaluation. Endpoints include age at menarche/spermarche, pubertal stage and levels of reproductive hormones. Questionnaire items include age at menarche and use of contraception for females, age at spermarche for males and hormonal substitution. Pubertal stage is evaluated according to Tanner⁷² in combination with testicular volume for males, estimated by comparative palpation using a Prader orchidometer.⁷³ Results are compared with Danish puberty normograms.^{74 75} Measurements of serum follicle stimulating hormone, luteinising hormone, estradiol, testosterone, insulin-like peptide 3, inhibin B, anti-Müllerian hormone and sex hormone binding globulin are performed, and levels are compared with sex-matched and age-matched national reference material as well as with ALL-STAR controls.

Evaluation of fertility

Fertility is evaluated in all participants aged ≥18 years. Endpoints include impaired fertility (defined as previously diagnosed impaired fertility or infertility and/or use of assisted reproductive technology), number and health of offspring. Questionnaire data include items regarding health behaviour, previous relationship/marital status, attempts to conceive, menopause, results from fertility investigations, natural conceptions, use of assisted reproductive technology, pregnancies, miscarriages, voluntary terminations, stillbirths, live births and health of offspring. Reproductive hormones and testicular volume are assessed as described in the previous section.

Evaluation of oral health and salivary secretion

Oral health assessments are performed at the Copenhagen University Hospital. Endpoints include oral symptoms, impaired oral and dental health (mucosal changes, caries, erosions, gingivitis, periodontitis, malocclusions, temporomandibular dysfunction and salivary gland dysfunction) and changes in salivary microbiome and proteome. Questionnaire items include oral symptoms, current medication, dietary supplements, tobacco and alcohol consumption and oral hygiene habits. Xerostomia is assessed by means of the bother index,⁷⁶ xerostomia inventory-8⁷⁷ and visual analogue scale (0–10).⁷⁸ Additional evaluation of oral health includes a

clinical examination with palpation of the temporomandibular joint, salivary glands and regional lymph nodes; assessments of the morphological and functional occlusion, dental status (decayed missing filled teeth scores, dental erosion scores, dental malformations and delayed eruptions), periodontal status (gingival inflammation and dental plaque indices and assessment of periodontal probing depth) and mucosal status (mucosal changes and signs of dryness using clinical oral dryness score).⁷⁹ Unstimulated and chewing-stimulated whole saliva flow rates are measured according to the drooling method.⁸⁰ DNA is extracted from the saliva samples and the microbiome characterised by Human Oral Microbe Identification (using next-generation sequencing) with bacterial species-level identification based on 16S rDNA comparisons. Salivary proteome analyses are performed by means of 2D electrophoresis (2DE) and matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry. Western blot analysis and ELISA are used for validation of the 2DE results.

Evaluation of other somatic health conditions

Iron overload is evaluated with plasma levels of ferritin and transferrin saturation and with cardiac MRI T2* mapping. Hyperthyroidism and hypothyroidism is investigated with questionnaire data and measurement of thyroid stimulating hormone and free thyroxine. Hyperparathyroidism is evaluated with parathyroid hormone and total and ionised calcium. Additional health conditions specified in [table 1](#), but not described above, are evaluated using questionnaire data and medical chart validation as reported in online supplemental table 1.

Evaluation of mental health and quality of life

Endpoints include psychiatric disease, social relations, emotional distress (anxiety, depression and fatigue), executive function and health-related quality of life. Questionnaire items include psychiatric diagnoses and psychopharmaceuticals. Additional psychosocial data are collected via instruments from Paediatric Quality of Life Inventory, Patient Reported Outcomes Measurement Information System, and the Behavior Rating Inventory of executive Function system. Specific instruments and versions are provided in online supplemental table 4.

Use of previously collected data for risk factor analysis

ALL patient characteristics (date of diagnosis, tumour burden, immunophenotype, cytogenetics, minimal residual disease levels, final risk stratification, participation in protocol randomisations and occurrence of 19 well-defined acute toxicities) are obtained from the NOPHO ALL2008 registry.³⁸ Medical chart data are used for validation of registered acute toxicities and for validation of self-reported health conditions (including dates). Genotyping for genome wide association studies has been performed previously on germline DNA from samples obtained after clinical remission using the exome-enriched Illumina Infinium Omni2.5Exome-8 BeadChip arrays and is available for approximately 70% of the survivor population. Whole blood samples are collected

from participants (including controls) not previously genotyped and samples are analysed using the same platform.

Data management

Study data were collected and managed using Research Electronic Data Capture (REDCap) tools hosted at Copenhagen University Hospital.^{81 82} REDCap is a secure, web-based software platform designed to support data capture for research studies, providing: (1) an intuitive interface for validated data capture; (2) audit trails for tracking data manipulation and export procedures; (3) automated export procedures for seamless data downloads to common statistical packages; and (4) procedures for data integration and interoperability with external sources.

Statistical considerations

Cumulative incidence of the 197 health-conditions is calculated (individually and grouped according to organ system and severity grade) for survivors and controls and presented with 95% confidence intervals (CI). Gray's test is used for comparison between the two groups. Cumulative burden of the 197 health conditions is calculated (individually and grouped according to organ system and severity grades) in survivors and controls by the method of mean cumulative count, estimating the mean number of recurring or multiple health events in the presence of competing risk,⁸³ and presented with 95% CIs. For this purpose, health conditions are categorised pre-analysis into three subtypes (chronic, non-recurrent; single, recurrent; and chronic, recurrent) based on clinical definitions of recurrence and chronicity, as described by others.⁴⁰ Based on a projected sample size of 250 survivors and 250 controls (survival rate of 90% in the Danish NOPHO ALL2008 patient cohort and a conservative recruitment rate of 60%), a 16% prevalence of chronic disease among Danish children,⁸⁴ and significance level $\alpha=0.05$, we will have a power of 87% to detect a HR of 2.0 in a Cox regression model, which is the lowest HR previously reported in a study comparing leukaemia survivors with sibling controls.⁸⁵ Secondary outcomes (ie, description of organ-specific functions) are compared between survivors and controls using relevant regression models (eg, linear regression models, logistic regression models or Cox regression models). For survivors, associations of primary and secondary outcomes with therapy exposure and occurrence of acute toxicities are explored using multiple regression models. Association with single nucleotide polymorphisms (SNPs) is explored with genome-wide association analysis and using a candidate gene approach where relevant SNPs have been identified. Differences in demographics between survivor participants and non-participants are explored using Fisher's exact test for binary data and Student's/Welch's t-test or Wilcoxon rank-sum test.

Patient and public involvement

In preparation of this study, we performed a qualitative interview study among adolescent and young adult

NOPHO ALL2008 survivors to identify the challenges they perceive as most important.¹⁸ Results from this study have been incorporated into the ALL-STAR examination programme, securing focus on the issues most important to the survivors.

Longitudinal follow-up

The cohort is invited for follow-up examinations corresponding to 5-year intervals from diagnosis, thereby aligning survival time at data capture. Results from the initial cross-sectional studies will inform which examinations to prioritise and additional endpoints can be added to address new hypotheses. The ALL-STAR database will facilitate repeated invitations and longitudinal data capture. Participants will provide new informed consent before repeated examinations.

ETHICS AND DISSEMINATION

The study has been approved by the Regional Ethics Committee for the Capital Region in Denmark (no. H-18035090 and H-20006359), the Danish Data Protection Agency (no. VD-2018-519) and is conducted in accordance with the Helsinki Declaration. Written informed consent is obtained from all participants prior to enrolment. All participants can choose to participate in only parts of the study, can choose not to be informed of results, and can withdraw consent at any time. Study results will be published in international peer-reviewed scientific journals and presented at relevant conferences.

DISCUSSION

The ALL-STAR cohort is the first Nordic, population-based childhood and young adult ALL survivor cohort treated according to the same ALL protocol, to be uniformly evaluated for treatment-related morbidity across several organ systems, using both self-report data and objective, clinical investigations.

The evolution of long-term, adverse consequences of antileukaemic therapy is complex and dependent not only on therapy exposure and genetic susceptibility, but also on age at diagnosis, pre-existing morbidity, individual resources, resources among family and peers, and on surrounding healthcare structures and societal structures. The comparison of a Nordic survivor cohort with survivor cohorts from other Western countries will provide insight into the consequences of differing systems, as well as differences in treatment regimens used. Scandinavian countries, and particularly Denmark, have access to extensive register data including health information and socioeconomic information, linked to the individual person and surrounding network through personal identification numbers.³⁹ Coupling of systematic survivor data obtained in the ALL-STAR study to national register data will enable contextualising association studies novel to survivorship research.

Strengths of the ALL-STAR study include a population-based design, a homogenous and well characterised cohort, the combination of both subjective and objective assessments of outcome, and inclusion of matched controls. The ALL-STAR cohort has already been characterised by prospective registration of tumour burden, immunophenotype, cytogenetics, treatment response and occurrence of 19 severe toxicities, performed by treating physicians from time of diagnosis until end of therapy.³⁸ These data enable high precision evaluation of risk factors and confounders associated with our main outcome of interest. Similarly, high quality survivor data is secured through self-report data, medical chart validation and systematic clinical examinations across most organ systems and using both well-established and novel, advanced techniques. This combination of data enables inclusion of survivor perspective while minimising risk of recall bias and inaccurate information. Furthermore, the systematic screening of organ functions facilitates capture of low-grade conditions, which are likely to be underestimated if using retrospective or self-report data only.⁴⁰ Since the investigated health conditions are not pathognomic, the inclusion of matched controls, often missing in survivorship studies, will support conclusions made regarding the risk patterns, which we seek to identify.

Despite vast amounts of late effects research being performed, no international consensus on how to characterise therapy-related late effects currently exists. Lack of precise definitions and uniform grading result in substantial variability across studies, hindering meaningful pooling and comparisons of results.⁸⁶ Cancer and therapy-related acute toxicities are traditionally defined and graded according to the CTCAE system and systematic modifications proposed by St Jude researchers has improved its utility for also addressing the long-term effects among childhood cancer survivors.⁴³ We have aligned our outcome definitions accordingly and by introducing 10 additional health conditions of relevance for cancer survivorship research, we wish to further increase the relevance of this classification system.

The magnitude of late effects is often described with cumulative incidence; however, we encourage the use of the cumulative burden, which includes recurrent events as opposed to only first events, thereby reflecting the burden more comprehensively. This is illustrated by a study investigating therapy-related cardiovascular conditions among survivors of childhood and young adult Hodgkin's lymphoma. By the age of 50 years, cumulative incidence of all-grade cardiovascular health conditions did not differ between survivors and controls, while the cumulative burden of the same conditions was almost double in survivors compared with controls.⁴⁰ Since none of the measures, however, reflect survivor perspective, we have included the perceived impact of the individual health conditions, psychosocial function and health-related quality of life as reported by the participants. Coupling of objective findings with subjective survivor experiences is essential for interpretation of results and

for guiding healthcare workers and researchers when choosing which survivorship issues to prioritise.

There are several limitations to our study. A survival time of 3.5–14 years at evaluation means that the full spectrum of therapy-related conditions will not be captured. Establishment of the ALL-STAR cohort within the first decade of survival, however, forms the platform for longitudinal follow-up, which is within the overall scope of the project. Interpretation of initial data will be challenged by a heterogeneous distribution in both age and survival time, but this will be adjusted for in the statistical analysis. Several of the health conditions evaluated in the ALL-STAR study rely on questionnaire data and subsequent medical chart validation. Low-grade conditions, which are not screened for are, therefore, likely to be missed. Finally, the ALL-STAR study outcomes are not exhaustive in describing the treatment-related burden. Important outcomes, such as semen quality, ovarian reserve and neurocognitive function were omitted to ensure feasibility of the initial ALL-STAR initiative. Future studies of these important outcomes in the ALL-STAR cohort will benefit from the infrastructure already created.

In conclusion, the ALL-STAR study is establishing a Nordic childhood and young adult ALL survivor cohort to be systematically assessed for the cumulative incidence and burden of treatment-related morbidity. Results will supplement existing evidence regarding risk of treatment-related morbidity and the underlying possible biological, physical and psychosocial mechanisms, while providing novel insight into differences in survivorship related to specific therapy regimens, health cultures and healthcare systems. Findings may guide future toxicity risk grouping and preventive interventions during treatment and follow-up, ultimately ameliorating the burden of therapy without compromising chance of cure.

Author affiliations

¹Department of Pediatrics and Adolescent Medicine, Copenhagen University Hospital, Copenhagen, Denmark

²Department of Pediatrics and Adolescent Medicine, Aarhus University Hospital, Aarhus, Denmark

³Department of Cardiology, Aarhus University Hospital, Aarhus, Denmark

⁴Department of Cardiology, Copenhagen University Hospital, Copenhagen, Denmark

⁵Section of Biostatistics, University of Copenhagen, Copenhagen, Denmark

⁶Section of Oral Medicine/Oral Biology and Immunopathology, Department of Odontology, University of Copenhagen, Copenhagen, Denmark

⁷Department of Neurology, Zealand University Hospital Roskilde, Roskilde, Denmark

⁸Institute of Clinical Medicine, University of Copenhagen Faculty of Health and Medical Sciences, Copenhagen, Denmark

⁹Department of Radiology, Aarhus University Hospital, Aarhus, Denmark

¹⁰Department of Growth and Reproduction, Copenhagen University Hospital, Copenhagen, Denmark

¹¹Department of Radiology, Copenhagen University Hospital, Copenhagen, Denmark

¹²Department of Pediatrics, Aalborg University Hospital, Aalborg, Denmark

¹³Department of Pediatric Hematology and Oncology, Odense University Hospital, Odense, Denmark

¹⁴Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

Acknowledgements The authors wish to acknowledge valuable contributions made by Dr Margrethe Bastholm Bille; Associate professor Niels Holtum Birkebæk; Dr Alfred Peter Born; Dr Vibeke Brix; Sisse Højlund Carlsen; Pernille Fleggaard; Dr Christian Alcaraz Frederiksen; Dr Peter Orm Hansen; Professor Won Yong Kim; Dr

Karen Markussen Linnet; Professor Bente Lomholt Langdahl; Associate professor Hanne Baekgaard Larsen; Dr Sonja Izquierdo Riis Meyer; Professor Rasmus Mogelvang; Kamilla Tofing-Olesen; Dr Redi Pecini; Professor Steen Hvittfeldt Poulsen; Dr Ida Maria Schmidt. Figure 1 is created with BioRender.com.

Contributors LAJ, KS, and TLF devised the study concept, design and protocol. LAJ drafted the manuscript. PHJ and KAM contributed to the cardiac protocol. MKF contributed to the protocol regarding physical performance. AMLP devised and wrote the protocol regarding oral health. SLMR and KGN contributed to the pulmonary protocol. MB contributed to the protocol regarding neuropathy. AH and TLL devised the ultrasound protocols. RBJ, MTS and KMC contributed to the protocol regarding metabolic syndrome and bone density, and MTS to the pancreatic protocol. KG monitored the plan for statistical analyses. LAJ, MTS, KMC, RT, PSW, BKA, KS and TLF contributed to the logistical aspects of the study. All authors contributed to writing and reviewing the manuscript, and made the decision to submit the manuscript for publication.

Funding This study has received funding from the Danish Children's Cancer Foundation (2017–2082; 2019–5966), The Danish Cancer Society (R192-A11.590), Cancer Survivorship and General Late effects (CASTLE, F-2 23 724–03), The Danish Cancer Research Foundation (FID20823), NordForsk (ID 91 172), Engineer Otto Christensen's Foundation (101459), University of Aarhus (grant number not applicable), Holm's Memorial Foundation (20 006–1975), AP Moeller Foundation (18-L-0225 L-0225), Farmer of Oelufgaard Memorial Foundation (grant number not applicable) and Skagen Teddy Bear Museum (grant number not applicable). The funding bodies have not had a role in the design, manuscript writing or in the decision to submit the manuscript for publication, nor will they have in the collection of data, analysis or interpretation.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer-reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Liv Andrés-Jensen <http://orcid.org/0000-0003-3437-8338>

Mette Tiedemann Skipper <http://orcid.org/0000-0002-6281-4814>

REFERENCES

- Hunger SP, Mullighan CG. Acute lymphoblastic leukemia in children. *N Engl J Med* 2015;373:1541–52.
- Pui C-H, Yang JJ, Hunger SP, et al. Childhood acute lymphoblastic leukemia: progress through collaboration. *J Clin Oncol* 2015;33:2938–48.
- Toft N, Birgens H, Abrahamsson J, et al. Results of NOPHO ALL2008 treatment for patients aged 1–45 years with acute lymphoblastic leukemia. *Leukemia* 2018;32:606–15.
- Schmiegelow K, Frandsen TL. The cost of cure. *Lancet Haematol* 2018;5:e504–5.
- Toft N, Birgens H, Abrahamsson J, et al. Toxicity profile and treatment delays in NOPHO ALL2008-comparing adults and children with Philadelphia chromosome-negative acute lymphoblastic leukemia. *Eur J Haematol* 2016;96:160–9.
- Wolthers BO, Frandsen TL, Baruchel A, et al. Asparaginase-associated pancreatitis in childhood acute lymphoblastic leukaemia: an observational Ponte di Legno toxicity Working Group study. *Lancet Oncol* 2017;18:1238–48.
- Rank CU, Toft N, Tuckuviene R, et al. Thromboembolism in acute lymphoblastic leukemia: results of NOPHO ALL2008 protocol treatment in patients aged 1 to 45 years. *Blood* 2018;131:2475–84.
- Millan NC, Pastrana A, Gutter MR, et al. Acute and sub-acute neurological toxicity in children treated for acute lymphoblastic leukemia. *Leuk Res* 2018;65:86–93.
- Wolthers BO, Mogensen PR, Frandsen TL, et al. Insulin-Dependent diabetes: a chronic complication to acute pancreatitis in childhood acute lymphoblastic leukemia. *Pediatr Blood Cancer* 2019;66:e27437.
- Mogensen SS, Harila-Saari A, Mäkitie O, et al. Comparing osteonecrosis clinical phenotype, timing, and risk factors in children and young adults treated for acute lymphoblastic leukemia. *Pediatr Blood Cancer* 2018;65:e27300.
- Mody R, Li S, Dover DC, et al. Twenty-Five-Year follow-up among survivors of childhood acute lymphoblastic leukemia: a report from the childhood cancer Survivor study. *Blood* 2008;111:5515–23.
- Mulrooney DA, Hyun G, Ness KK, et al. The changing burden of long-term health outcomes in survivors of childhood acute lymphoblastic leukaemia: a retrospective analysis of the ST Jude lifetime cohort study. *Lancet Haematol* 2019;6:e306–16.
- Vora A, Andreano A, Pui C-H, et al. Influence of cranial radiotherapy on outcome in children with acute lymphoblastic leukemia treated with contemporary therapy. *J Clin Oncol* 2016;34:919–26.
- Armstrong GT, Chen Y, Yasui Y, et al. Reduction in late mortality among 5-year survivors of childhood cancer. *N Engl J Med* 2016;374:833–42.
- Jacola LM, Edelstein K, Liu W, et al. Cognitive, behaviour, and academic functioning in adolescent and young adult survivors of childhood acute lymphoblastic leukaemia: a report from the childhood cancer Survivor study. *Lancet Psychiatry* 2016;3:965–72.
- Vetsch J, Wakefield CE, Robertson EG, et al. Health-related quality of life of survivors of childhood acute lymphoblastic leukemia: a systematic review. *Qual Life Res* 2018;27:1431–43.
- Anestini AS, Lippé S, Robaey P, et al. Psychological risk in long-term survivors of childhood acute lymphoblastic leukemia and its association with functional health status: a PETALE cohort study. *Pediatr Blood Cancer* 2018;65:e27356.
- Andrés-Jensen L, Larsen HB, Johansen C, et al. Everyday life challenges among adolescent and young adult survivors of childhood acute lymphoblastic leukemia: an in-depth qualitative study. *Psychooncology* 2020;29:1630–7.
- Robison LL, Hudson MM. Survivors of childhood and adolescent cancer: life-long risks and responsibilities. *Nat Rev Cancer* 2014;14:61–70.
- Teepen JC, van Leeuwen FE, Tissing WJ, et al. Long-Term risk of subsequent malignant neoplasms after treatment of childhood cancer in the DCOG later study cohort: role of chemotherapy. *J Clin Oncol* 2017;35:2288–98.
- Turcotte LM, Liu Q, Yasui Y. Chemotherapy and risk of subsequent malignant neoplasms in the childhood cancer survivor study cohort. In: *Journal of Clinical Oncology*. Vol 37. American Society of Clinical Oncology 2019:3310–9.
- Green DM, Liu W, Kutteh WH, et al. Cumulative alkylating agent exposure and semen parameters in adult survivors of childhood cancer: a report from the ST Jude lifetime cohort study. *Lancet Oncol* 2014;15:1215–23.
- Sklar CA, Mertens AC, Mitby P, et al. Premature menopause in survivors of childhood cancer: a report from the childhood cancer Survivor study. *J Natl Cancer Inst* 2006;98:890–6.
- Nysom K, Holm K, Lipsitz SR, et al. Relationship between cumulative anthracycline dose and late cardiotoxicity in childhood acute lymphoblastic leukemia. *J Clin Oncol* 1998;16:545–50.
- Krischer JP, Epstein S, Cuthbertson DD, et al. Clinical cardiotoxicity following anthracycline treatment for childhood cancer: the pediatric Oncology Group experience. *J Clin Oncol* 1997;15:1544–52.
- Lipshultz SE, Lipsitz SR, Sallan SE, et al. Chronic progressive cardiac dysfunction years after doxorubicin therapy for childhood acute lymphoblastic leukemia. *J Clin Oncol* 2005;23:2629–36.
- Kadan-Lottick NS, Dinu I, Wasilewski-Masker K, et al. Osteonecrosis in adult survivors of childhood cancer: a report from the childhood cancer Survivor study. *J Clin Oncol* 2008;26:3038–45.
- Cheung YT, Krull KR. Neurocognitive outcomes in long-term survivors of childhood acute lymphoblastic leukemia treated on contemporary treatment protocols: a systematic review. *Neurosci Biobehav Rev* 2015;53:108–20.
- Tay CG, Lee VWM, Ong LC, et al. Vincristine-Induced peripheral neuropathy in survivors of childhood acute lymphoblastic leukaemia. *Pediatr Blood Cancer* 2017;64:e26471.

- 30 Clemens E, van der Kooi ALF, Broer L, *et al.* The influence of genetic variation on late toxicities in childhood cancer survivors: a review. *Crit Rev Oncol Hematol* 2018;126:154–67.
- 31 Kraljinovic M, Elbared J, Drouin S, *et al.* Polymorphisms of ABCC5 and NOS3 genes influence doxorubicin cardiotoxicity in survivors of childhood acute lymphoblastic leukemia. *Pharmacogenomics J* 2016;16:530–5.
- 32 Nadeau G, Ouimet-Grennan E, Aaron M, *et al.* Identification of genetic variants associated with skeletal muscle function deficit in childhood acute lymphoblastic leukemia survivors. *Pharmacogenomics Pers Med* 2019;12:33–45.
- 33 Petrykey K, Lippé S, Robaey P, *et al.* Influence of genetic factors on long-term treatment related neurocognitive complications, and on anxiety and depression in survivors of childhood acute lymphoblastic leukemia: the Petale study. *PLoS One* 2019;14:e0217314.
- 34 Robison LL, Mertens AC, Boice JD, *et al.* Study design and cohort characteristics of the childhood cancer Survivor study: a multi-institutional collaborative project. *Med Pediatr Oncol* 2002;38:229–39.
- 35 Asdahl PH, Winther JF, Bonnesen TG, *et al.* The adult life after childhood cancer in Scandinavia (ALiCCS) study: design and characteristics. *Pediatr Blood Cancer* 2015;62:2204–10.
- 36 Hawkins MM, Lancashire ER, Winter DL, *et al.* The British childhood cancer Survivor study: objectives, methods, population structure, response rates and initial descriptive information. *Pediatr Blood Cancer* 2008;50:1018–25.
- 37 Winther JF, Kenborg L, Byrne J. Childhood cancer Survivor cohorts in Europe. *Acta Oncol* 2015;54:655–68.
- 38 Frandsen TL, Heyman M, Abrahamsson J, *et al.* Complying with the European Clinical Trials directive while surviving the administrative pressure – an alternative approach to toxicity registration in a cancer trial. *Eur J Cancer* 2014;50:251–9.
- 39 Frank L. Epidemiology. when an entire country is a cohort. *Science* 2000;287:2398–9.
- 40 Bhakta N, Liu Q, Yeo F, *et al.* Cumulative burden of cardiovascular morbidity in paediatric, adolescent, and young adult survivors of Hodgkin's lymphoma: an analysis from the ST Jude lifetime cohort study. *Lancet Oncol* 2016;17:1325–34.
- 41 Bhakta N, Liu Q, Ness KK, *et al.* The cumulative burden of surviving childhood cancer: an initial report from the ST Jude lifetime cohort study (SJLIFE). *Lancet* 2017;390:2569–82.
- 42 U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute. Common terminology criteria for adverse events (CTCAE) | protocol development | CTEP, 2010. Available: https://ctep.cancer.gov/protocolDevelopment/electronic_applications/ctc.htm#ctc_40 [Accessed 12 Aug 2020].
- 43 Hudson MM, Ehrhardt MJ, Bhakta N, *et al.* Approach for classification and severity grading of long-term and late-onset health events among childhood cancer survivors in the St. Jude lifetime cohort. *Cancer Epidemiol Biomarkers Prev* 2017;26:666–74.
- 44 Cole TJ, Bellizzi MC, Flegal KM, *et al.* Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000;320:1240–3.
- 45 Tinggaard J, Aksglaede L, Sørensen K, *et al.* The 2014 Danish references from birth to 20 years for height, weight and body mass index. *Acta Paediatr* 2014;103:214–24.
- 46 Grundy SM, Cleeman JI, Daniels SR, *et al.* Diagnosis and management of the metabolic syndrome: an American heart Association/National heart, lung, and blood Institute scientific statement. *Circulation* 2005;112:2735–52.
- 47 Cook S, Weitzman M, Auinger P, *et al.* Prevalence of a metabolic syndrome phenotype in adolescents: findings from the third National health and nutrition examination survey, 1988–1994. *Arch Pediatr Adolesc Med* 2003;157:821–7.
- 48 van der Aa MP, Knibbe CAJ, Boer Ade, *et al.* Definition of insulin resistance affects prevalence rate in pediatric patients: a systematic review and call for consensus. *J Pediatr Endocrinol Metab* 2017;30:123–31.
- 49 Masuzaki R, Kanda T, Sasaki R, *et al.* Noninvasive assessment of liver fibrosis: current and future clinical and molecular perspectives. *Int J Mol Sci* 2020;21:4906.
- 50 Lang RM, Badano LP, Mor-Avi V, *et al.* Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of echocardiography and the European association of cardiovascular imaging. *J Am Soc Echocardiogr* 2015;28:1–39.
- 51 Kramer CM, Barkhausen J, Bucciarelli-Ducci C, *et al.* Standardized cardiovascular magnetic resonance imaging (CMR) protocols: 2020 update. *J Cardiovasc Magn Reson* 2020;22:17.
- 52 Schulz-Menger J, Bluemke DA, Bremerich J, *et al.* Standardized image interpretation and post-processing in cardiovascular magnetic resonance – 2020 update : Society for Cardiovascular Magnetic Resonance (SCMR): Board of Trustees Task Force on Standardized Post-Processing. *J Cardiovasc Magn Reson* 2020;22:19.
- 53 Quanjer PH, Borsboom GJ, Brunekreef B, *et al.* Spirometric reference values for white European children and adolescents: Polgar revisited. *Pediatr Pulmonol* 1995;19:135–42.
- 54 Quanjer PH, Stanojevic S, Cole TJ, *et al.* Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012;40:1324–43.
- 55 Frei J, Jutla J, Kramer G, *et al.* Impulse oscillometry: reference values in children 100 to 150 cm in height and 3 to 10 years of age. *Chest* 2005;128:1266–73.
- 56 Anagnostopoulou P, Latzin P, Jensen R, *et al.* Normative data for multiple breath washout outcomes in school-aged Caucasian children. *Eur Respir J* 2020;55:1901302.
- 57 Stanojevic S, Graham BL, Cooper BG, *et al.* Official ERS technical standards: global lung function initiative reference values for the carbon monoxide transfer factor for Caucasians. *Eur Respir J* 2017;50:1700010.
- 58 Zavorsky GS, Hsia CCW, Hughes JMB, *et al.* Standardisation and application of the single-breath determination of nitric oxide uptake in the lung. *Eur Respir J* 2017;49:1600962.
- 59 England JD, Gronseth GS, Franklin G, *et al.* Distal symmetric polyneuropathy: a definition for clinical research: report of the American Academy of Neurology, the American association of Electrodiagnostic medicine, and the American association of physical medicine and rehabilitation. *Neurology* 2005;64:199–207.
- 60 Cornblath DR, Chaudhry V, Carter K, *et al.* Total neuropathy score: validation and reliability study. *Neurology* 1999;53:1660–4.
- 61 Bloom S, Till S, Sönksen P, *et al.* Use of a biothesiometer to measure individual vibration thresholds and their variation in 519 non-diabetic subjects. *Br Med J* 1984;288:1793–5.
- 62 Maffei L, Premrou V, Roldan P, *et al.* Vibration perception threshold in the screening of sensorimotor distal symmetric polyneuropathy: the need of more accurate age-specific reference values. *J Diabetes Sci Technol* 2014;8:621–2.
- 63 Hilz MJ, Axelrod FB, Hermann K, *et al.* Normative values of vibratory perception in 530 children, juveniles and adults aged 3–79 years. *J Neurol Sci* 1998;159:219–25.
- 64 Sletten DM, Suarez GA, Low PA, *et al.* Compass 31: a refined and abbreviated composite autonomic symptom score. *Mayo Clinic Proceedings* 2012;87:1196–201.
- 65 Treister R, O'Neil K, Downs HM, *et al.* Validation of the composite autonomic symptom scale 31 (COMPASS-31) in patients with and without small fiber polyneuropathy. *Eur J Neurol* 2015;22:1124–30.
- 66 Thorsteinsson T, Helms AS, Adamsen L, *et al.* Study protocol: rehabilitation including social and physical activity and education in children and teenagers with cancer (respect). *BMC Cancer* 2013;13:1.
- 67 Shepherd RJ, Allen C, Benade AJ, *et al.* The maximum oxygen intake. An international reference standard of cardiorespiratory fitness. *Bull World Health Organ* 1968;38:757–64.
- 68 Bohannon RW. Sit-to-stand test for measuring performance of lower extremity muscles. *Percept Mot Skills* 1995;80:163–6.
- 69 Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991;39:142–8.
- 70 Levey AS, Stevens LA, Schmid CH, *et al.* A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;150:604–12.
- 71 Schwartz GJ, Muñoz A, Schneider MF, *et al.* New equations to estimate GFR in children with CKD. *J Am Soc Nephrol* 2009;20:629–37.
- 72 Tanner JM. *Growth at adolescence*. Oxford: Blackwell Scientific Publications, 1962.
- 73 Prader A. Testicular size: assessment and clinical importance. *Triangle* 1966;7:240–3.
- 74 Lawaetz JG, Hagen CP, Mieritz MG, *et al.* Evaluation of 451 Danish boys with delayed puberty: diagnostic use of a new puberty nomogram and effects of oral testosterone therapy. *J Clin Endocrinol Metab* 2015;100:1376–85.
- 75 Lindhardt Johansen M, Hagen CP, Mieritz MG, *et al.* Pubertal progression and reproductive hormones in healthy girls with transient Thelarche. *J Clin Endocrinol Metab* 2017;102:1001–8.
- 76 Challacombe SJ, Osailan SM, Proctor GBCarpenter G, ed. *Dry mouth: a clinical guide on causes, effects and treatments*. Groeningen: Springer Berlin Heidelberg, 2015.
- 77 Thomson WM, van der Putten G-J, de Baat C, *et al.* Shortening the xerostomia inventory. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;112:322–7.

- 78 Pedersen AM, Reibel J, Nauntofte B. Primary Sjögren's syndrome (pSS): subjective symptoms and salivary findings. *J Oral Pathol Med* 2007;28:303–11.
- 79 Osailan SM, Pramanik R, Shirlaw P, *et al.* Clinical assessment of oral dryness: development of a scoring system related to salivary flow and mucosal wetness 2012;114:597–603.
- 80 Pedersen A, Reibel J, Nordgarden H, *et al.* Primary Sjögren's syndrome: salivary gland function and clinical oral findings. *Oral Dis* 2008;5:128–38.
- 81 Harris PA, Taylor R, Thielke R, *et al.* Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.
- 82 Harris PA, Taylor R, Minor BL, *et al.* The REDCap Consortium: building an international community of software platform partners. *J Biomed Inform* 2019;95:103208.
- 83 Dong H, Robison LL, Leisenring WM, *et al.* Estimating the burden of recurrent events in the presence of competing risks: the method of mean cumulative count. *Am J Epidemiol* 2015;181:532–40.
- 84 Sundhedsstyrelsen (Danish National Health Authority). Sociale Forholds Betydning for Håndteringen AF Børn Med Kronisk Sygdom, 2012. Available: <http://www.sst.dk> [Accessed 29 Sep 2020].
- 85 Oeffinger KC, Mertens AC, Sklar CA, *et al.* Chronic health conditions in adult survivors of childhood cancer. *N Engl J Med* 2006;355:1572–82.
- 86 Streefkerk N, Fioole LCE, Beijer JGM, *et al.* Large variation in assessment and outcome definitions to describe the burden of long-term morbidity in childhood cancer survivors: a systematic review. *Pediatr Blood Cancer* 2020;67:e28611.