BMJ Open Training general practitioners in melanoma diagnosis: a scoping review of the literature

Evelyne Harkemanne ^{1,2} Marie Baeck ^{1,2} Isabelle Tromme^{1,3}

To cite: Harkemanne E, Baeck M, Tromme I. Training general practitioners in melanoma diagnosis: a scoping review of the literature. *BMJ Open* 2021;**11**:e043926. doi:10.1136/bmjopen-2020-043926

► Prepublication history and additional materials for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-043926>).

Received 18 August 2020
Revised 03 March 2021
Accepted 09 March 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Service de dermatologie, Cliniques universitaires Saint-Luc, Bruxelles, Belgique

²Pôle de pneumologie et dermatologie, Institut de Recherche Expérimentale et Clinique, UCLouvain, Bruxelles, Belgique

³Clinique du mélanome, Institut Roi Albert II, Cliniques universitaires Saint-Luc, Bruxelles, Belgique

Correspondence to

Dr Evelyne Harkemanne; evelyne.harkemanne@uclouvain.be

ABSTRACT

Background General practitioners (GPs) play a key role in early melanoma detection. To help GPs deal with suspicious skin lesions, melanoma diagnostic training programmes have been developed. However, it is unclear whether these programmes guarantee the acquisition of skills that will be applied by GPs in their daily clinical practice and maintained over time.

Objectives This scoping review aimed to examine and compare educational programmes designed to train GPs in melanoma diagnosis using clinical (naked eye) examination alone or dermoscopy±clinical examination, and sought to inform on the long-term sustainability of the GPs' acquired skills.

Eligibility criteria Studies eligible for inclusion evaluated educational programmes for teaching diagnosis of melanoma to GPs. MEDLINE, EMBASE and Cochrane databases were searched for relevant articles from 1995 to May 2020.

Results Forty-five relevant articles were found assessing 31 educational programmes. Most programmes that improved the diagnostic accuracy and long-term performances of the GPs, that is, increase in confidence, decrease in dermatologist referral for benign skin lesions and improvement in the benign/malignant ratio of excised skin lesions, trained the GPs in clinical diagnosis, followed by dermoscopy. To maintain long-term performances, these programmes provided refresher training material.

Conclusion This review shows that studies generally report positive outcomes from the training of GPs in melanoma diagnosis. However, refresher training material seemed necessary to maintain the acquired skills. The optimal form and ideal frequency for these updates have yet to be defined.

INTRODUCTION

Early melanoma detection is essential to reduce the morbidity and mortality of patients with melanoma.¹ Given the increased incidence of this aggressive skin cancer, primary care physicians (PCPs) play a key role in early melanoma diagnosis.^{2–4} PCPs include a number of healthcare professionals who provide first and continuing medical care to a patient. In this review, we focus on general practitioners (GPs) who take care of patients in community settings and are, in most

Strengths and limitations of this study

- Systematic review was conducted following the guidelines of the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews checklist.
- It thoroughly evaluates educational programmes on melanoma diagnosis for general practitioners.
- Specifically, the review examines the long-term effect of educational programmes and the value of providing regular refresher training sessions after the training.
- This review led inevitably to some publication bias as only English-language peer-reviewed articles were included.

countries, the first point of contact for any patient with a health issue.

To improve the diagnostic accuracy of melanoma by GPs, specific educational training programmes have been developed. At first, training courses focused on melanoma diagnosis by clinical (naked eye) examination alone. A systematic review,⁵ published in 2011, reported on 20 studies that evaluated 13 educational interventions in clinical melanoma diagnosis for PCPs. All the evaluated interventions improved diagnostic accuracy and melanoma management. Later, educational programmes that included dermoscopy training were created and then evaluated for primary care. To date, dermoscopy has been the most widely used non-invasive in vivo technique in clinical practice to assess skin tumours.⁶ It involves the use of a handheld device that allows the observation of skin structures invisible to the naked eye. However, the sensitivity and specificity of the technique are operator-dependent (trained vs untrained physicians).⁷ Ninety-two percent sensitivity and 95% specificity can be achieved for melanoma diagnosis by a trained dermatologist combining visual inspection and in vivo dermoscopy.⁸ In primary care, dermoscopy has also been shown to be an effective

tool for the triage of suspicious pigmented skin lesions when performed by properly trained PCPs.^{9 10} Yet, the minimum training required to reach competence is still unknown.¹¹

Previously published reviews^{5 11–14} on training programmes in melanoma diagnosis for GPs focused on the content, teaching method, outcome measures and study-by-study efficacy of the evaluated educational interventions. However, they did not assess whether the GPs' acquired skills were measured in the short or long term. Yet, given the increasing burden of melanoma on general practice, it is crucial to know whether these programmes are capable of teaching GPs easily applicable and sustainable skills in melanoma diagnosis and management. This scoping review aimed to explore educational programmes training GPs in melanoma diagnosis using clinical (naked eye) examination alone and diagnosis using dermoscopy±clinical examination. Educational programmes were examined regarding training content, teaching method, training duration, availability of refresher training material and outcome measures. This review also specifically sought to inform on the long-term sustainability of the skills acquired during these training programmes.

MATERIAL AND METHODS

To carry out this literature review, a scoping review seemed the most appropriate research method. Indeed, the studies we identified, which provided evidence on the efficacy of educational programmes in melanoma diagnosis for GPs, showed a wide range of study designs and heterogeneous outcome measures. This observation made it impossible to formally assess the quality of these studies and to perform a meta-analysis, leading to a narrative synthesis of our research results. To conduct this scoping review, the framework developed by Arksey and O'Malley,¹⁵ subsequently refined by Levac *et al*,¹⁶ and the guidelines of the PRISMA-ScR¹⁷ (Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews) checklist were followed.

Eligibility criteria

Studies eligible for inclusion in this review (box 1) evaluated educational programmes teaching either clinical diagnosis of melanoma or diagnosis using dermoscopy±clinical examination and designed primarily for PCPs including GPs. The population of interest included qualified GPs and GP trainees. Specialists and GPs working in hospital settings and/or specialised clinics were excluded. Studies that included training programmes for PCPs other than GPs were not eligible. Studies where no participant training in melanoma diagnosis was proposed and studies evaluating exclusively non-melanoma skin cancer detection were also excluded. Studies evaluating teledermoscopy and computer-aided diagnosis of melanoma were not assessed as they do not require specific education in melanoma recognition by the participants. Only studies assessing the type of educational programme

Box 1 Inclusion and exclusion criteria for article selection

Inclusion criteria:

Articles

- ▶ Articles written in English.
- ▶ Study articles and descriptive articles of educational programmes.

Population

- ▶ Qualified GPs and GP trainees working in community settings.

Intervention

- ▶ Studies evaluating educational programmes in clinical (naked eye) diagnosis and/or diagnosis of melanoma using dermoscopy.

Outcome(s)

- ▶ Studies assessing the type of educational programme and its short-term and/or long-term efficacy on the skills acquired by the GPs.

Exclusion criteria:

Articles

- ▶ Articles not subject to peer review and written in languages other than English.

Population

- ▶ Studies involving specialists, medical students, non-GPs, GPs working in hospital settings and/or specialised skin cancer clinics.

Intervention

- ▶ Studies evaluating exclusively non-melanoma skin cancer.
- ▶ Teledermoscopy studies.
- ▶ Studies on computer-aided diagnosis of melanoma.

Outcome(s)

- ▶ No method of measuring outcomes was ruled out.

Abbreviations: GPs, general practitioners.

and its short-term and/or long-term efficacy on the skills acquired by the GPs were included. Finally, only peer-reviewed and English-language articles were included.

Data sources and study selection

MEDLINE, EMBASE and Cochrane databases were searched for relevant articles published from 1995 to May 2020. Studies were selected for inclusion independently by three authors (EH, MB and IT), with IT providing the final decision in the event of disagreement. The studies were not assessed for bias, as the risk of bias assessment was reported as not applicable to scoping reviews in the 2018 PRISMA-ScR guidelines.¹⁷

To extensively cover the literature on the subject, four categories of terms were identified: (1) GPs, family doctor, general medicine, family practice and PCPs; (2) education and continuing medical education; (3) melanoma, malignant melanoma, cutaneous melanoma and skin neoplasms; and (4) diagnosis and early detection of cancer. In MEDLINE, the following Medical Subject Headings were used: *general practitioners* OR *family practice* OR *primary care physicians* OR *general practice* AND *melanoma* AND *diagnosis*. No limits were defined. In EMBASE, Emtree terms were exploded: *general practitioner*, *family doctor*, *primary care*, *family physician*, *primary care physician*, *melanoma*, *diagnosis* and *education*. In the Cochrane database, the following terms were searched: *melanoma* AND *diagnosis* AND *general practitioners* OR *family medicine* AND *dermoscopy* (see online supplemental table 1 for search

Table 1 Definition of study categories

Criteria	Categories	Definition
Training content	Epidemiology	Background information on rates of melanoma cancer, risk factors, localisation and evolution of melanomas
	Clinical diagnosis	Naked eye melanoma recognition
	Clinical algorithm	Use of a pre-existing algorithm as a learning tool to aid for clinical diagnosis
	Dermoscopic diagnosis	Recognition of melanoma using dermoscopy
	Dermoscopic algorithm	Use of an algorithm as a learning tool to aid for dermoscopic diagnosis
	Management	Determination of a plan of action for a skin lesion, that is, reassurance, follow-up or lesion excision
Teaching method	Live	Presentation by a speaker to a group of participants
	Scientific literature	Use of educational books, posters, letters, CD-ROMs or videos
	E-learning	Interactive online tutorials including audio and visual information
	Self-assessment	Learning by the participant himself using educational material
Refresher training material	Teledermatology feedback	Feedback from a dermatologist on the image and clinical history of a suspicious lesion at a distance, using remote internet-based technologies
Outcome measures	Competences	Acquired skills, which are evaluated in a training setting on clinical and/or dermoscopic photographs of skin lesions
	Diagnostic accuracy	Ability of the participants to discriminate between melanoma and benign lesions
	Knowledge	Report of conceptual understanding
	Appropriate management	Determination of the right plan of action for a skin lesion
	Performances	Changes in real-life practice measured in a clinical setting, that is, changes in the benign/malignant ratio of excised lesions, the number of total-body skin examinations performed, confidence of the general practitioners, changes in referral rates to a dermatologist and decrease in the incidence of advanced melanomas
Evaluation	Short-term	Measurement of outcomes immediately or up to 3 months after the training
	Long-term	Measurement of outcomes at ≥ 6 months after the training

strategies). In addition, the reference lists of included studies were screened as a source of further relevant articles.

Data extraction

Two authors (EH and MB) reviewed all included articles and independently collected data. Extracted data included authors, year of publication, origin of the article, study design, number of participating GPs, type of educational programme, type of outcome measures and short-term and/or long-term evaluation of these outcomes. The type of educational programme included training content, teaching method, training duration and refresher training material (if provided). To facilitate comparison with data found in previous reviews, all these data were reported into categories adapted from those presented by Fee *et al.*¹⁴

Table 1 gives the definition of the different categories. The training content was subdivided into six components: epidemiology, clinical diagnosis, clinical algorithm, dermoscopic diagnosis, dermoscopic algorithm and management. The teaching method was considered as live, in the form of scientific literature, e-learning or self-assessment. The refresher training material specified

the material available for participants to refresh their skills after the training. The outcome measures were expressed either in terms of competence or in terms of performance, according to the assessment approach of continuing medical education programmes proposed by Moore.¹⁸ Finally, since the limits between short-term and long-term evaluation of a medical educational programme are not standardised, arbitrary limits have been chosen based on the observations made during this literature review.

Patient and public involvement

No patient and public involvement was required for this review.

RESULTS

In total, 325 articles were identified from the electronic database searches, as shown in the PRISMA flowchart (figure 1).¹⁹ At the end of the study selection process, 45 relevant articles, which assessed 31 educational interventions, were included in the review analysis.

Study designs

Thirty-six interventional studies with a range of study designs were found: 11 randomised controlled trials

Figure 1 Preferred Reporting Items for Systematic reviews and Meta-Analyses flowchart of the study selection process.¹⁷ GP, general practitioner.

(RCTs),^{9 10 20–29} 19 diagnostic accuracy studies,^{30–48} 3 cohort studies^{49–51} and 3 case-control studies.^{52–54} Five of the 31 training programmes were assessed twice.^{25 26 28 32 49}

Four systematic reviews were identified: one on the training of PCPs in clinical melanoma diagnosis,⁵ two on the training of PCPs in dermoscopy for melanoma diagnosis^{12 13} and one on the use of dermoscopy in primary care.¹¹ A scoping review on the training of PCPs in dermoscopy¹⁴ was also included. The final three articles were descriptive articles of the educational programmes and study protocols.^{55–57}

Educational programmes

The educational programmes in melanoma diagnosis for GPs varied in terms of content, teaching method and outcome measures. The characteristics of these training programmes are summarised in table 2.

Training content

Of the 31 educational programmes, 15 involved the training of GPs in clinical diagnosis, 5 involved dermoscopic diagnosis alone and 11 involved the training of GPs in both of these melanoma diagnostic methods. Twelve (80%) of the clinical diagnostic training programmes also involved learning of epidemiology and

11 (73%) involved learning of management guidelines for suspicious lesions. Only seven (47%) programmes teaching clinical diagnosis used an algorithm to teach melanoma recognition, with the ABCD(E) rule⁵⁸ (Asymmetry, uneven Borders, uneven Colours, Diameter >6 mm and Evolution) being most commonly taught. Of the dermoscopic training programmes, 12 (80%) included learning of at least one dermoscopic algorithm (Menzies method,^{24 35 37 40} three-point checklist,^{9 40 41} seven-point checklist,^{10 35} triage amalgamated dermoscopic algorithm (TADA),^{44 47} the ABCD rule,^{35 38} BLINCK⁴⁰ and pattern analysis^{35 54}). The Menzies method,⁵⁹ the ABCD rule,⁶⁰ pattern analysis and seven-point checklist⁶¹ were designed originally for trained physicians and were later tested as effective when used by non-experts.⁶² Other algorithms, such as the three-point checklist⁶³ and the TADA,⁴⁴ were initially created for use by PCPs. In addition, two educational programmes included training on other diagnostic tools, such as sequential digital dermoscopy imaging³⁷ and polaroid instant camera photography.²⁶

Teaching method, training duration and refresher training material

Live training courses and the use of educational books, posters or videos (literature) were the two preferred

Table 2 Characteristics of educational programmes in melanoma diagnosis for GPs

Article	Study design	Study participants	Training content				Teaching method	Training duration	Refresher raining material
Author, year, location ^{ref.}			Epidemiology diagnosis	Clinical diagnosis algorithm	Dermoscopic diagnosis	Dermoscopic algorithm	Management		
Marra <i>et al.</i> , 2020, The Netherlands ⁵³	Case-control study	185 (83; 102)*	X	X	X (optional)	X	X	E-learning	2 hours
Sawyers <i>et al.</i> , 2020, Canada ⁴⁸	DA study	33 GPs			X	TADA step I		Live	3.5 hours
Augustsson <i>et al.</i> , 2019, Sweden ⁵⁴	Case-control study	43 GPs (27;16)*			X	Pattern analysis		Live	1 day
Seiverling, 2019, USA ⁴⁷	DA study	59 GPs	X	X	X	TADA		Live	75 min
Beecher <i>et al.</i> , 2018, Ireland ⁴⁶	DA study	23 GP trainees	X	X			X	Live, literature	1 hour
Secker <i>et al.</i> , 2017, The Netherlands ⁴⁵	DA study	293 PCPs including ? GPs	X	X	X		X	Live, literature and e-learning	1 day
Rogers <i>et al.</i> , 2016, USA ⁴⁴	DA study	16 GPs			X	TADA		Live	1 day
Badertscher <i>et al.</i> , 2011, 2015, Switzerland ^{29,56}	RCT	78 GPs (39;39)*	X	X	Lumio			Live	1 day
Gulati <i>et al.</i> , 2015, UK ⁵¹	Cohort study	967 GPs	X	X			X	E-learning	PD
Koelink <i>et al.</i> , 2014, The Netherlands ¹⁰	RCT	53 GPs	X	X	X	Seven-point checklist		Live	4-hour clinical diagnosis; 6-hour dermoscopy
Grange <i>et al.</i> , 2013, France ⁵⁰	Cohort study	398 GPs	X	X				Live, Literature	2.5 hours
Markova <i>et al.</i> , 2013, USA ²⁸	RCT	46 GPs (20;26)*						E-learning	CD-ROM+regular information sheets
Mikkilineni <i>et al.</i> , 2001 and 2002, USA ^{42,43}	DA study	7 GPs	X	X			X	Live Literature	2 hours
Eide <i>et al.</i> , 2013 ⁴¹ (Shaikh <i>et al.</i> , 2012 ⁵⁵) INFORMED	DA study	54 PCPs including 9 GPs (20%)	X	X	X	Three-point checklist	X	E-learning, self-assessment	2-hour web-based learning
Bourne <i>et al.</i> , 2012, Australia ⁴⁰	DA study	3 GPs	X	X	X	BLINCK Three-point checklist Menzies method		Live, literature	UD
Shariff <i>et al.</i> , 2010, UK ³⁹	DA study	94 GPs	X					Literature	PD
Grimaldi <i>et al.</i> , 2009, Italy ³⁸	DA study	13 GPs	X	X	X	ABCD rule		Live, e-learning, self-assessment	PD
Menzies <i>et al.</i> , 2009, Australia ³⁷	DA study	63 GPs	X			Menzies method		Live, literature, e-learning	2-hour workshop
Peuvrel <i>et al.</i> , 2009, France ³⁶	DA study	210 GPs	X	X		ABCD(E)	X	Live, literature (CD-ROM)	2 hours

Continued

Table 2 Continued

Article	Study design	Study participants	Training content				Teaching method	Training duration	Refresher raining material
Author, year, location ^{ref.}			Epidemiology diagnosis	Clinical diagnosis	Clinical algorithm	Dermoscopic diagnosis	Dermoscopic algorithm	Management	
Youl <i>et al.</i> 2007, Australia ⁴⁹	Cohort study	16 GPs							Self-assessment, paper-based training sessions
Raasch <i>et al.</i> 2000, Australia ²³	RCT	46 GPs (23;23)*	X	X	ABCD(E)	X		X	PD
Argenziano <i>et al.</i> 2006, Italy and Spain ⁵	RCT	73 GPs (36;37)*		X	ABCD	X	Three-point checklist	Live	1 day
Doliantis <i>et al.</i> 2005, Australia ³⁵	DA study	35 GPs				X	Menzies method Seven-point checklist ABCD rule Pattern analysis	Literature, e-learning, self-assessment	PD
Carl <i>et al.</i> 2005, Italy ²⁴	DA study	41 GPs	X	X	ABCD(E)		X	Live	4 hours
de Gannes <i>et al.</i> 2004, Canada ²⁷	RCT	27 GPs (10;17)*	X	X				Literature (video format)	12 min
English <i>et al.</i> 2003, Australia ²⁶	RCT	468 GPs (245;228)*		X+polaroid instant camera			X	Literature	>8hours
Del Mar and Green, 1995, Australia ²⁰	RCT	93 GPs (48;45)*							1 hour
Brochez <i>et al.</i> 2001, Belgium ³³	DA study	146 GPs	X	X			X	Live, literature	2 hours
Harris <i>et al.</i> 1999 and 2001, USA ^{30, 32}	DA study	232 GPs 17 GPs	X	X	ABCD Seven-point Glasgow checklist		X	Literature, e-learning	1 hour
Westerhoff <i>et al.</i> Australia 2000 ²⁴	RCT	74 GPs				X	Menzies method	Live, literature	1-hour live
Bedlow <i>et al.</i> 2000, UK ³¹	DA study	17 GPs		X				Live, literature	UD
Gerbert <i>et al.</i> 1998 and 2002, USA ^{22, 25}	RCT	52 GPs (26;26)*	X	X			X	Live, literature, e-learning	>3hours
Dolan <i>et al.</i> 1997, USA ²¹	RCT	82 PCPs, including 16 GPs (46;36)*	X	X				Live	2 hours
Girgis <i>et al.</i> 1995, Australia ⁵²	Case-control study	41 GPs (24;17)*	X	X			X	Live, literature	>6hours

*Number of participants in the intervention group and control group.

ABCD, Asymmetry, uneven Borders, uneven Colours and Diameter >6 mm; ABCD(E), Asymmetry, uneven Borders, uneven Colours, Diameter >6 mm and Evolution; DA, diagnostic accuracy; GPs, general practitioners; INFORMED, Internet curriculum for melanoma early detection; PCPs, primary care physicians; PD, participant-dependent; RCT, randomised controlled trial; TADA, triage amalgamated dermoscopic algorithm; UD, undisclosed.

teaching methods of clinical diagnostic training programmes. Five training programmes also used an e-learning approach.^{25 28 30 51 53} The most common teaching method used in dermoscopic training programmes was live training. This approach was combined with literature and/or e-learning in six programmes. Three programmes also used self-assessment. Overall, the teaching method did not appear to have influenced the programme outcomes. Duration of training varied from 75 min to 1 day, was not specified in two studies^{31 40} and was participant-dependent in six studies using self-assessment methods. Six dermoscopic diagnostic training programs^{29 37 38 41 49 54} and three programmes in clinical diagnosis^{27 36 50} provided regular refresher training material such as unlimited e-learning access or self-assessment training sessions.

Training outcomes

Table 3 summarises the outcome measures of the studies. In the selected studies, the GPs' competences were generally measured in the short term and their performances measured in the long term after the training.

Eight clinical diagnostic training programmes and seven dermoscopic training programmes only assessed the short-term efficacy of their programme (table 3A). For these studies, the competences most often evaluated were diagnostic accuracy and appropriate management measured in a training setting. The most evaluated short-term performance, measured in a clinical setting, was the GPs' confidence in diagnosing melanoma. With the exception of two studies, all showed a positive impact of their intervention.^{21 23} Four clinical diagnostic training programmes and three dermoscopic training programmes (one teaching dermoscopy alone⁵⁴) only assessed long-term performances (table 3B). The most evaluated long-term performances, measured in daily clinical practice, were the GPs' diagnostic accuracy and the benign/malignant ratio of excised lesions. Three studies^{9 36 49} reported improvement in the GPs' performances in melanoma diagnosis. The other studies reported no improvement.

Finally, three clinical diagnostic and seven dermoscopic training programmes assessed the short-term and long-term outcome of their training (table 3C). Except for one,²³ all these training programmes demonstrated improvement in the GPs' competences, measured in a training setting in the short term. In the long term, eight training programs^{10 37 38 41 49 50 53 54} reported significant improvement in the GPs' performances for the diagnosis of melanoma and benign skin lesions. This led to either a decrease in the referral rates to dermatologists^{37 41 53} or a decrease in the ratio of benign/malignant excised skin lesions.^{37 49} Among the major studies, Koelink *et al*¹⁰ found that their dermoscopic training programme improved the GPs' long-term performances with up to 1.25 times greater diagnostic accuracy for skin lesions including melanomas. In a French department, Grange *et al*⁵⁰ observed an impressive reduction in the incidence of advanced melanomas (Breslow thickness ≥ 3 mm) during the 3-year period after

their GP training campaign in clinical melanoma diagnosis. A very recent study by Marra *et al*⁵³ assessing 1662 referrals reported better quality of referrals by GPs trained in melanoma diagnosis than by untrained GPs, potentially leading to less unnecessary referrals. However, two educational programs^{28 29} were unable to maintain the GPs' acquired performances in the long term.

DISCUSSION

This scoping review aimed to explore educational programmes training GPs in melanoma diagnosis using clinical (naked eye) examination alone and diagnosis using dermoscopy±clinical examination. Educational programmes were examined regarding training content, teaching method, training duration, availability of refresher training material and outcome measures. This review also specifically sought to inform on the long-term sustainability of the skills acquired during these training programmes.

Types of educational programmes with positive long-term outcomes

Most reported educational programmes that improved long-term diagnostic accuracy and changed GPs' melanoma practice patterns trained their participating GPs in dermoscopy combined with clinical diagnosis. This teaching method is supported by a recent Cochrane review⁸ in which dermoscopy alone was found to be less accurate than clinical examination followed by dermoscopy for the diagnosis of melanoma. The Cochrane review results also suggested that dermoscopic algorithms were the most useful method to train non-experts in dermoscopy. In our review, we found that five of the programmes with long-term positive impact used dermoscopic algorithms to teach melanoma diagnosis.^{10 37 38 41 54}

Unfortunately, the substantial number of training hours necessary to become competent in dermoscopy is the main reported factor limiting its use in general practice.^{64 65} At this time, there is no evidence on the optimal length of training, even though it has been demonstrated that diagnostic accuracy of dermoscopy depends on the degree of training of the practitioner.⁶⁶ One study suggested that 1 day of live training in dermoscopy was sufficient to build the confidence of GPs with special interest in melanoma diagnosis.⁶⁷ We found two RCTs that demonstrated sustained improvement in GPs' diagnostic accuracy, both of which proposed live training in dermoscopy over 1 day or 10 hours.^{9 10} Because the duration of training in dermoscopy is a limiting factor for most GPs, it is important to keep in mind that training in clinical melanoma diagnosis has also been shown to improve the GPs' performances while requiring less training time (in this review, a mean duration of 2.5 hours was observed for clinical diagnostic training programmes).^{36 50}

However, we found that educational programmes teaching dermoscopy have been more likely to assure positive long-term outcomes than programmes teaching clinical examination alone. One of the reasons could be that the latter used measures of performance such as GPs'

Table 3 Outcome measures of the melanoma diagnostic educational programmes for general practitioners

(A) Studies with short-term outcomes												
Studies	Measured competence in training setting			Measured performance in clinical setting			Timing after the training					
(First author and year of publication)	DA	Knowledge	AM	B/M ratio	TBSE	DA	Confidence	Decrease in VTM incidence	RR	Immediately	At 1 month	At 3 months
Dermoscopic diagnostic training programmes												
Sawyers (2020) ⁴⁸	+									X		
Seiverling (2019) ⁴⁷	+									X		
Secker (2017) ⁴⁵	+		+									X
Rogers (2016) ⁴⁴	+									X		
Bourne (2012) ⁴⁰	+									X		
Dolanitis (2005) ³⁵	+									X		
Westerhoff (2000) ²⁴	+		+							X		
Clinical diagnostic training programmes												
Beecher (2018) ⁴⁶	+		+							X		X
Carli (2005) ³⁴	+		+							X		
Mikkilineni (2001, 2002) ^{42 43}			+		+			+		X		
Brochez (2001) ³³	+									X		
Harris (2001) ³²	+		+					+		X		
Harris (1999) ³⁰	+		+					+		X		
Raasch (2000) ²³	-							+			3-month period	
Bedlow (2000) ³¹	+		+							X		
Gerbert (1998, 2002) ^{22 25}	+		+					+		X		
Dolan (1997) ²¹	-		+							X	X	
Girgis (1995) ⁵²	-		+					+		X		X
(B) Studies with long-term outcomes												
Studies	Measured competence in training setting			Measured performance in clinical setting			Timing after the training					
	DA	Knowledge	AM	B/M ratio	TBSE	DA	Confidence	Decrease in VTM incidence	RR			
Dermoscopic diagnostic training programmes												
Shariff (2010) ³⁹						-				At 11 months		
Youl (2007) ⁴⁹				+		+				6-month period		
Argenziano (2006) ⁹						+			+	16-month period		
Clinical diagnostic training programmes												
Gulati (2015) ⁵¹						-		+		8-month period		

Continued

Table 3 Continued

Peuvrel (2009) ³⁶	+			+															15-month period
de Gannes (2004) ²⁷	-	-																	At 6 months
English (2003) ²⁶																			21-month period
Del Mar (1995) ²⁰																			24-month period
(C) Studies with short-term and long-term outcomes																			
Studies	Measured competence in training setting				Timing after the training				Measured performance in clinical setting				Timing after the training						
	DA	Knowledge	AM		Immediately after	At 1-3 months		B/M ratio	TBSE	DA	Confidence	Decrease in VTM incidence	RR						
Dermoscopic diagnostic training programmes																			
Augustsson (2019) ⁵⁴	+				X														At 6 months (here competence measure)
Badertscher (2015) ²⁹	+				X					-	+								At 12 months
Koelink (2014) ¹⁰	+		+		X					+	+								At 8 months
Eide (2013) ⁴¹	+	+	+	+	X					+	+								At 6 months
Youl (2007) ⁴⁹								+											6-month period
Raasch (2000) ²³	-				3-month period														
Grimaldi (2009) ³⁸	+				X					+									6-month period
Menzies (2009) ³⁷	+	+	+	+	X			+											6-month period
Clinical diagnostic training programmes																			
Marra (2020) ⁵³	+									+	+								At 10 months
Grange (2013) ⁵⁰	+	+			X							+							3-year period
Markova (2013) ²⁸										-									At 12 months
Mikkilineni (2001, 2002) ^{42 43}		+	+		X														

To note that Menzies *et al*³⁷ and Carli *et al*³⁴ provided only descriptive statistics of their study results. For the other studies, a p value <0.05 was considered significant, except for Peuvrel *et al*³⁶ (p<0.001).

+ = significant improvement; - = no significant improvement.
AM, appropriate management; B/M, benign/malignant; DA, diagnostic accuracy; RR, decrease in dermatologist referral rates for benign lesions and increase in referral rates for malignant lesions; TBSE, total-body skin examination; VTM, very thick melanomas.

confidence level and number of total-body skin examinations performed before and after training, which did not reflect GPs' diagnostic ability in clinical practice. On the one hand, measuring the confidence of GPs in their own ability to diagnose melanoma is more useful in assessing the quality of a training programme than evaluating skills acquired by participants. From a pedagogical point of view, participants feel more confident when they know how to use the teaching content in daily practice, but this does not define their true diagnostic competence.¹⁸ On the other hand, the number of total-body skin examinations performed may be useful in measuring GPs' awareness of skin tumours, but not for evaluation of GPs' diagnostic skills.

Long-term improvement of the GPs' performances in clinical settings

The GPs' long-term performances measured in clinical settings were assessed for 15 educational programmes: 6 in clinical diagnosis and 9 in dermoscopic diagnosis. Ten showed a positive impact on the GPs' performances measured over periods ranging from 6 to 19 months. The most frequent observations were a decrease in referral rates to dermatologists for benign skin lesions and an improvement in the benign/malignant ratio of excised skin lesions. The internet curriculum for melanoma early detection (INFORMED) group⁴¹ reported an increase in melanoma diagnosis during a screening campaign by GPs trained with their programme in 2016.⁶⁸ Furthermore, a decrease in the incidence of advanced melanomas was shown in a French department over a 3-year period after their training programme in clinical melanoma diagnosis.⁵⁰ Unfortunately, two educational programs^{28 29} failed to maintain the GPs' acquired performances at 1 year after the end of the training. The reasons might be that Markova *et al*²⁸ chose to assess the number of total-body skin examinations performed but did not evaluate the GPs' diagnostic accuracy and that Badertscher *et al*²⁹ trained their GPs to use Lumio, a polarised magnifying glass with 2× magnification instead of a standard dermoscopy device (10× magnification).

To retain acquired diagnostic skills over the long term, results of a recent RCT suggested the need for 'refresher sessions at regular intervals'.^{69 70} In our review, nine (60%) educational programmes evaluated in the long term provided refresher training material. Seven of these programmes were successful. In 2014, Grange *et al*⁵⁰ produced a CD-ROM containing their teaching material and sent regular information about melanoma to the GPs. The INFORMED group⁴¹ provided GPs with a free unlimited access to their web-based course. Menzies *et al*³⁷ gave participating GPs an educational textbook and an unlimited e-learning access. Grimaldi *et al*³⁸ and Youl *et al*⁴⁹ also ensured self-assessment refresher training sessions. Marra *et al*⁵³ found a 10-month sustainability of the diagnostic accuracy of their trained GPs and assumed that daily use of the obtained knowledge during the study period achieved this effect. Only Koelink *et al*¹⁰ who evaluated the longest post-training period (19 months) and who demonstrated

sustainability of diagnostic skills did not specify whether update training modalities were provided.

However, the ideal frequency and form of updates have never been studied. An RCT, taking place in the English National Defibrillator Programme, determined that update session intervals after a medical education session should not exceed 7 months to limit the loss of acquired skills and maintain the participants' confidence.⁷¹ In the UK, a survey among GPs with special interest in dermatology stated that self-assessment learning was the most popular for refresher sessions.⁶⁷ Nevertheless, they also showed that 36% of GPs who use dermoscopy in their clinical practice reported to have never updated their training skills. We found that the most appreciated form of self-assessment updates was the unlimited access to an e-learning course. In the future, this enthusiasm for online training could lead to the development of smartphone applications to train GPs in melanoma diagnosis. Some newly developed applications have currently been evaluated among medical students⁷² and dermatology residents.⁷³ Initial results already looked very promising.

Finally, the variability of refresher training material provided in the educational programmes and the heterogeneity of outcome measures did not allow more robust conclusions to be drawn on the most beneficial training modality for sustainable improvements in GPs' diagnostic skills.

LIMITATIONS

This scoping review has some limitations and led inevitably to certain publication biases. We used keywords for the selection of articles, and only peer-reviewed articles were included. By limiting our research to English-language articles, some studies may also have been missed. It is also very likely that melanoma diagnostic training programmes exist in unpublished forms, for example, in university continuing medical education programmes. Moreover, we focused only on studies assessing melanoma diagnostic training methods among GPs. Therefore, we may have failed to mention some educational programmes for primary care in this review. Furthermore, educational programmes which were not evaluated in studies were overlooked. Finally, this review covers educational programmes over a 25-year period. As technology has evolved considerably over this time, some teaching methods and refresher training materials have been overshadowed by interactive online tutorials (e-learning)—all the more so with the health crisis caused by COVID-19 during which distance learning methods have developed very rapidly.

CONCLUSION

In conclusion, educational programmes trained GPs in melanoma diagnosis using clinical examination alone or dermoscopy±clinical diagnosis. Most reported programmes that improved the long-term diagnostic accuracy and changed routine performances of the

GPs (ie, decrease in dermatologist referral of benign skin lesions and improvement in the benign/malignant ratio of excised skin lesions) trained their participating GPs in both diagnostic methods. The preferred teaching methods were live and e-learning, but the teaching method did not seem to influence the GPs' acquired performances. It is important to note that the educational programmes that achieved long-term sustainability of GPs' performances in daily clinical practice provided refresher training material. However, no conclusions on the most beneficial training modality to sustainably improve GPs' diagnostic skills could be drawn given the heterogeneity of outcome measures and study designs. Therefore, the optimal form and ideal frequency of these updates have yet to be defined.

Acknowledgements The authors would like to thank Graziella Messina, Research Assistant at the Health Sciences Library of the Catholic University of Louvain, UCLouvain in Brussels, for her help in developing the research strategy for this review.

Contributors EH developed the research protocol, completed the literature search and screened the articles for inclusion, extracted the data, synthesised the findings, interpreted the results and drafted the manuscript. MB screened the articles for inclusion, extracted data and critically revised the manuscript. IT developed the protocol, independently reviewed the articles for inclusion, extracted data, interpreted the results and critically revised the manuscript. All authors approved the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Evelyne Harkemanne <http://orcid.org/0000-0001-9936-9868>

Marie Baeck <http://orcid.org/0000-0003-0499-7939>

REFERENCES

- Petrie T, Samatham R, Witkowski AM. Melanoma early detection: big data, bigger picture. *J Invest Dermatol* 2019;139:25–30.
- van Rijsingen MCJ, van Bon B, van der Wilt GJ, et al. The current and future role of general practitioners in skin cancer care: an assessment of 268 general practitioners. *Br J Dermatol* 2014;170:1366–8.
- Buckley D, McMonagle C. Melanoma in primary care. The role of the general practitioner. *Ir J Med Sci* 2014;183:363–8.
- Grange F, Barbe C, Mas L, et al. The role of general practitioners in diagnosis of cutaneous melanoma: a population-based study in France. *Br J Dermatol* 2012;167:1351–9.
- Goulart JM, Quigley EA, Dusza S, et al. Skin cancer education for primary care physicians: a systematic review of published evaluated interventions. *J Gen Intern Med* 2011;26:1027–35.
- Bakos RM, Blumetti TP, Roldán-Marín R, et al. Noninvasive imaging tools in the diagnosis and treatment of skin cancers. *Am J Clin Dermatol* 2018;19:3–14.
- Piccolo D, Ferrari A, Peris K, et al. Dermoscopic diagnosis by a trained clinician vs. a clinician with minimal dermoscopy training vs. computer-aided diagnosis of 341 pigmented skin lesions: a comparative study. *Br J Dermatol* 2002;147:481–6.
- Dinnes J, Deeks JJ, Chuchu N, et al. Dermoscopy, with and without visual inspection, for diagnosing melanoma in adults. *Cochrane Database Syst Rev* 2018;96.
- Argenziano G, Puig S, Zalaudek I, et al. Dermoscopy improves accuracy of primary care physicians to triage lesions suggestive of skin cancer. *J Clin Oncol* 2006;24:1877–82.
- Koelink CJL, Vermeulen KM, Kollen BJ, et al. Diagnostic accuracy and cost-effectiveness of dermoscopy in primary care: a cluster randomized clinical trial. *J Eur Acad Dermatol Venerol* 2014;28:1442–9.
- Jones OT, Jurascheck LC, van Melle MA, et al. Dermoscopy for melanoma detection and triage in primary care: a systematic review. *BMJ Open* 2019;9:e027529.
- Herschorn A. Dermoscopy for melanoma detection in family practice. *Can Fam Physician* 2012;58:740–5.
- Chia A, Trevena L. A systematic review of training to improve melanoma diagnostic skills in general practitioners. *J Cancer Educ* 2016;31:730–5.
- Fee JA, McGrady FP, Rosendahl C. Training primary care physicians in dermoscopy for skin cancer detection: a scoping review. *J Cancer Educ* 2019.
- Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol Theory Pract* 2005.
- Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci* 2010;5:69.
- Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018;169:467–73.
- Moore DE. How physicians learn and how to design learning experiences for them: an approach based on an interpretive review of evidence. In: *Continuing education in the health professions: improving healthcare through lifelong learning*. 2008.
- Moher D, Liberati A, Tetzlaff JAD. PRISMA 2009 flow diagram. *Prism* 2009.
- Del Mar CB, Green AC. Aid to diagnosis of melanoma in primary medical care. *BMJ* 1995;310:492–5.
- Dolan NC, Ng JS, Martin GJ, et al. Effectiveness of a skin cancer control educational intervention for internal medicine housestaff and attending physicians. *J Gen Intern Med* 1997;12:531–6.
- Gerbert B, Bronstone A, Wolff M, et al. Improving primary care residents' proficiency in the diagnosis of skin cancer. *J Gen Intern Med* 1998;13:91–7.
- Raasch BA, Hays R, Buettner PG. An educational intervention to improve diagnosis and management of suspicious skin lesions. *J Contin Educ Health Prof* 2000;20:39–51.
- Westerhoff K, McCarthy WH, Menzies SW. Increase in the sensitivity for melanoma diagnosis by primary care physicians using skin surface microscopy. *Br J Dermatol* 2000;143:1016–20.
- Gerbert B, Bronstone A, Maurer T. The effectiveness of an Internet-based tutorial in improving primary care physicians' skin cancer triage skills. *J Cancer Educ* 2002.
- English DR, Burton RC, del Mar CB, et al. Evaluation of aid to diagnosis of pigmented skin lesions in general practice: controlled trial randomised by practice. *BMJ* 2003;327:375.
- de Gannes GC, Ip JL, Martinka M, et al. Early detection of skin cancer by family physicians: a pilot project. *J Cutan Med Surg* 2004;8:103–9.
- Markova A, Weinstock MA, Risica P, et al. Effect of a web-based curriculum on primary care practice: basic skin cancer triage trial. *Fam Med* 2013;45:558–68.
- Badertscher N, Tandjung R, Senn O. A multifaceted intervention: no increase in general practitioners' competence to diagnose skin cancer (minSKIN) - randomized controlled trial. *J Eur Acad Dermatology Venerol* 2015.
- Harris JM, Salasche SJ, Harris RB. Using the Internet to teach melanoma management guidelines to primary care physicians. *J Eval Clin Pract* 1999;5:199–211.
- Bedlow AJ, Cliff S, Melia J, et al. Impact of skin cancer education on general practitioners' diagnostic skills. *Clin Exp Dermatol* 2000;25:115–8.

- 32 Harris JM, Salasche SJ, Harris RB. Can Internet-based continuing medical education improve physicians' skin cancer knowledge and skills? *J Gen Intern Med* 2001;16:50–6.
- 33 Brochez L, Verhaeghe E, Bleyen L, et al. Diagnostic ability of general practitioners and dermatologists in discriminating pigmented skin lesions. *J Am Acad Dermatol* 2001;44:979–86.
- 34 Carli P, De Giorgi V, Crocetti E, et al. Diagnostic and referral accuracy of family doctors in melanoma screening: effect of a short formal training. *Eur J Cancer Prev* 2005;14:51–5.
- 35 Dolianitis C, Kelly J, Wolfe R, et al. Comparative performance of 4 dermoscopic algorithms by nonexperts for the diagnosis of melanocytic lesions. *Arch Dermatol* 2005;141:1008–14.
- 36 Peuvrel L, Quereux G, Jumbou O, et al. Impact of a campaign to train general practitioners in screening for melanoma. *Eur J Cancer Prev* 2009;18:225–9.
- 37 Menzies SW, Emery J, Staples M, et al. Impact of dermoscopy and short-term sequential digital dermoscopy imaging for the management of pigmented lesions in primary care: a sequential intervention trial. *Br J Dermatol* 2009;161:1270–7.
- 38 Grimaldi L, Silvestri A, Brandi C, et al. Digital epiluminescence dermoscopy for pigmented cutaneous lesions, primary care physicians, and telediagnosis: a useful tool? *J Plast Reconstr Aesthet Surg* 2009;62:1054–8.
- 39 Shariff Z, Roshan A, Williams AM, et al. 2-Week wait referrals in suspected skin cancer: does an instructional module for general practitioners improve diagnostic accuracy? *Surgeon* 2010;8:247–51.
- 40 Bourne P, Rosendahl C, Keir J, et al. BLINCK-A diagnostic algorithm for skin cancer diagnosis combining clinical features with dermoscopy findings. *Dermatol Pract Concept* 2012;2:202a12.
- 41 Eide MJ, Asgari MM, Fletcher SW, et al. Effects on skills and practice from a web-based skin cancer course for primary care providers. *J Am Board Fam Med* 2013;26:648–57.
- 42 Mikkilineni R, Weinstock MA, Goldstein MG, et al. The impact of the basic skin cancer triage curriculum on provider's skin cancer control practices. *J Gen Intern Med* 2001;16:302–7.
- 43 Mikkilineni R, Weinstock MA, Goldstein MG. The impact of the basic skin cancer triage curriculum on providers' skills, confidence, and knowledge in skin cancer control. *Prev Med* 2002.
- 44 Rogers T, Marino ML, Dusz SW, et al. A clinical aid for detecting skin cancer: the triage amalgamated dermoscopic algorithm (TadA). *J Am Board Fam Med* 2016;29:694–701.
- 45 Secker LJ, Buis PAJ, Bergman W, et al. Effect of a dermoscopy training course on the accuracy of primary care physicians in diagnosing pigmented lesions. *Acta Derm Venereol* 2017;97:263–5.
- 46 Beecher SM, Keogh C, Healy C. Dedicated general practitioner education sessions can improve diagnostic capabilities and may have a positive effect on referral patterns for common skin lesions. *Ir J Med Sci* 2018;187:959–63.
- 47 Seiverling EV, Ahrens HT, Greene A, et al. Teaching benign skin lesions as a strategy to improve the triage amalgamated dermoscopic algorithm (TadA). *J Am Board Fam Med* 2019;32:96–102.
- 48 Sawyers EA, Wigle DT, Marghoob AA, et al. Dermoscopy training effect on diagnostic accuracy of skin lesions in Canadian family medicine physicians using the triage Amalgamated Dermoscopic algorithm. *Dermatol Pract Concept* 2020;10:e2020035.
- 49 Youl PH, Raasch BA, Janda M, et al. The effect of an educational programme to improve the skills of general practitioners in diagnosing melanocytic/pigmented lesions. *Clin Exp Dermatol* 2007;32:365–70.
- 50 Grange F, Woronoff AS, Bera R, et al. Efficacy of a general practitioner training campaign for early detection of melanoma in France. *Br J Dermatol* 2014;170:123–9.
- 51 Gulati A, Harwood CA, Rolph J, et al. Is an online skin cancer toolkit an effective way to educate primary care physicians about skin cancer diagnosis and referral? *J Eur Acad Dermatol Venereol* 2015;29:2152–9.
- 52 Girgis A, Sanson-Fisher RW, Howe C, et al. A skin cancer training programme: evaluation of a postgraduate training for family doctors. *Med Educ* 1995;29:364–71.
- 53 Marra E, van Rijsingen MCJ, Alkemade JAC, et al. The effect of a dermato-oncological training programme on the diagnostic skills and quality of referrals for suspicious skin lesions by general practitioners. *Br J Dermatol* 2021;184:538–44.
- 54 Augustsson A, Paoli J. Effects of a 1-day training course in dermoscopy among general practitioners. *Dermatol Pract Concept* 2019;9:195–9.
- 55 Shaikh WR, Geller A, Alexander G, et al. Developing an interactive web-based learning program on skin cancer: the learning experiences of clinical educators. *J Cancer Educ* 2012;27:709–16.
- 56 Badertscher N, Rosemann T, Tandjung R, et al. minSKIN does a multifaceted intervention improve the competence in the diagnosis of skin cancer by general practitioners? Study protocol for a randomised controlled trial. *Trials* 2011;12:165.
- 57 Weinstock MA, Goldstein MG, Dubé CE, et al. Basic skin cancer triage for teaching melanoma detection. *J Am Acad Dermatol* 1996;34:1063–6.
- 58 Friedman RJ, Rigel DS, Kopf AW. Early detection of malignant melanoma: the role of physician examination and self-examination of the skin. *CA Cancer J Clin* 1985;35:130–51.
- 59 Menzies SW. A method for the diagnosis of primary cutaneous melanoma using surface microscopy. *Dermatol Clin* 2001;19:299–305.
- 60 Stolz W, Riemann A, Cagnetta AB. Abcd rule of dermoscopy: a new practical method for early recognition of malignant melanoma. *Eur J Dermatology* 1994.
- 61 Argenziano G, Fabbrocini G, Carli P, et al. Epiluminescence dermoscopy for the diagnosis of doubtful melanocytic skin lesions. *Arch Dermatol* 1998;134.
- 62 Pagnanelli G, Soyer HP, Argenziano G, et al. Diagnosis of pigmented skin lesions by dermoscopy: web-based training improves diagnostic performance of non-experts. *Br J Dermatol* 2003;148:698–702.
- 63 Soyer HP, Argenziano G, Zalaudek I, et al. Three-point checklist of dermoscopy. A new screening method for early detection of melanoma. *Dermatology* 2004;208:27–31.
- 64 Chappuis P, Duru G, Marchal O, et al. Dermoscopy, a useful tool for general practitioners in melanoma screening: a nationwide survey. *Br J Dermatol* 2016;175:744–50.
- 65 Oliveria SA, Heneghan MK, Cushman LF, et al. Skin cancer screening by dermatologists, family practitioners, and internists: barriers and facilitating factors. *Arch Dermatol* 2011;147:39–44.
- 66 Kittler H, Pehamberger H, Wolff K, et al. Diagnostic accuracy of dermoscopy. *Lancet Oncol* 2002;3:159–65.
- 67 Jones OT, Jurascheck LC, Utukuri M, et al. Dermoscopy use in UK primary care: a survey of GPs with a special interest in dermatology. *J Eur Acad Dermatol Venereol* 2019;33:1706–12.
- 68 Weinstock MA, Ferris LK, Saul MI, et al. Downstream consequences of melanoma screening in a community practice setting: first results. *Cancer* 2016;122:3152–6.
- 69 Robinson JK, Jain N, Marghoob AA, et al. A randomized trial on the efficacy of mastery learning for primary care provider melanoma opportunistic screening skills and practice. *J Gen Intern Med* 2018;33:855–62.
- 70 Robinson JK, MacLean M, Reavy R. Dermoscopy of concerning pigmented lesions and primary care providers' referrals at intervals after randomized trial of mastery learning. *J Gen Intern Med* 2018.
- 71 Woollard M, Whitfield R, Newcombe RG, et al. Optimal refresher training intervals for AED and CPR skills: a randomised controlled trial. *Resuscitation* 2006;71:237–47.
- 72 Lacy FA, Coman GC, Holliday AC, et al. Assessment of smartphone application for teaching intuitive visual diagnosis of melanoma. *JAMA Dermatol* 2018;154:730–1.
- 73 Nelson KC, Gershenwald JE, Savory SA, et al. Telementoring and smartphone-based answering systems to optimize dermatology resident dermoscopy education. *J Am Acad Dermatol* 2019;81:e27–8.