BMJ Open Areca nut consumption with and without tobacco among the adult population: a nationally representative study from India

Prashant Kumar Singh ^{1,2}, Amit Yadav,³ Lucky Singh,⁴ Sumit Mazumdar ⁵, Dharendra N Sinha,⁶ Kurt Straif,^{7,8} Shalini Singh^{2,9}

To cite: Singh PK, Yadav A, Singh L, *et al.* Areca nut consumption with and without tobacco among the adult population: a nationally representative study from India. *BMJ Open* 2021;11:e043987. doi:10.1136/bmjopen-2020-043987

► Prepublication history and supplemental material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-043987>).

Received 20 August 2020

Accepted 19 April 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Prashant Kumar Singh; prashants.geo@gmail.com

ABSTRACT

Objective Areca nut is one of the most widely consumed substances globally, after nicotine, ethanol and caffeine and classified as carcinogenic to humans. This study examines the disparity and determinants of areca nut consumption with and without tobacco in India.

Design Nationally representative cross-sectional study. **Participants** We used the nationally representative Global Adult Tobacco Survey 2016–2017. The analytical sample size was 74 037 individual's aged 15 years and above with a response rate of 92.9%.

Measures Current consumption of areca nut without tobacco and with tobacco.

Method We examined determinants of areca nut consumption (without tobacco and with tobacco) using multinomial logistic regression, accounting for the survey design.

Results About 23.9% (95% CI 23.1 to 24.8) of the adult population consume areca nut, that is, approximately 223.79 million people in India; majority of users (14.2%–95% CI 13.5 to 14.9) consumed areca nut with tobacco. When compared with women, men were more likely to consume areca nut (with tobacco relative risk (RR)=2.02; 95% CI 1.85 to 2.21 and without tobacco RR=1.13; 95% CI 1.07 to 1.20). Age, marital status, education, occupation, caste, religion and region were significantly associated with areca nut consumption. However, the direction and magnitude of association differ with respect to the areca nut consumption with and without tobacco.

Conclusion The ongoing tobacco control efforts would not address the majority of areca nut users until greater attention to areca nut consumption with and without tobacco is reflected in health policies in India.

INTRODUCTION

Areca nut is one of the most widely consumed substances globally, after nicotine, ethanol and caffeine.^{1,2} Owing to its addictive properties, areca nut is estimated to be consumed by hundreds of millions of people across various countries.³ However, addiction to areca nut is primarily prevalent in many Asia-Pacific countries and by emigrants from these countries in other parts of world.³ It is not only known

Strengths and limitations of this study

- Using a nationally representative survey with a high response rate, this study disentangled the current prevalence of areca nut consumption with and without tobacco in India, which has significant policy implications.
- The study provided detailed information on socioeconomic determinants of areca nut consumption, with and without tobacco, and separately for men and women, which may further guide future policy.
- The survey covers only people 15 years and older, whereas areca nut consumption often starts at younger age.
- The survey cannot provide insights into trends of areca nut consumption over time.

by several, sometimes local names, but also consumed in several forms, for example, pan masala, gutkha, mawa, dohra, kharra, betel and so on with or without tobacco.^{4,5} Some forms of consumption may also include other constituents, such as betel leaf, slaked lime and various spices.

The International Agency for Research on Cancer classified areca nut consumption with or without tobacco as carcinogenic to humans.⁶ A meta-analysis based on 50 studies worldwide reported increased relative risks for cancer of the oral cavity and oropharynx for the Indian subcontinent and areca nut consumption with tobacco (Relative Risk 7.03; 95% CI 4.68 to 10.56) and areca nut consumption without tobacco (Relative Risk 3.22; 95% CI 2.11 to 4.92).⁷ A global systematic review based on 62 studies concluded that consumption of areca nut affects almost all organs of the human body, including the brain, heart, lungs, gastrointestinal tract and reproductive organs; and causes or aggravates pre-existing conditions such as neuronal injury, myocardial infarction, cardiac

arrhythmias, hepatotoxicity, asthma, central obesity, type II diabetes, hyperlipidaemia, metabolic syndrome.⁸ It has harmful effects on the fetus when used during pregnancy.⁸ Previous studies observed that areca nut dependency among users⁹ and its withdrawal effects¹⁰ were similar to those observed among nicotine users.¹⁰ It is also a gateway product in children who start using different kinds of areca nut products at an early age.¹¹

Despite growing scientific evidence of high addictiveness and several ill effects (8–11) associated with areca nut consumption, research on areca nut has not received much attention.³ The large global and national movement that addresses tobacco control under the ambit of the WHO Framework Convention on Tobacco Control (FCTC) has focused primarily on smoking and has been less effective in controlling smokeless tobacco (SLT).¹² The regulatory framework for areca nut control has also remained limited to prescribing health warnings on areca nut products by the Food Safety and Standard Authority of India. Furthermore, use of tobacco and nicotine as an ingredient in any food item is also prohibited under FSSAI regulations, thereby restricting mixing of tobacco in areca nut products and vice versa.¹ Although tobacco control policies are applicable to areca nut products which contain tobacco a considerable number of people now consume areca nut without tobacco, which poses greater public health challenges in controlling and regulating the substance.¹³

A comprehensive search of the literature revealed that studies on areca nut use in India lack representativeness and published studies were restricted to a specific geographical area or population groups. None of the published studies have examined diverse habits of areca nut consumption, its disparity and determinants using a nationally representative survey. Also, a recent global review calls for more research to better understand the epidemiology of areca nut consumption across different populations and geographies.³

India, with a population of over 1.30 billion, exhibits one of the highest socioeconomic and demographic heterogeneities ever experienced anywhere in the world at the regional level.¹⁴ There is considerable evidence of marked regional inequities in tobacco use,¹⁵ health and healthcare¹⁶ and mortality outcomes¹⁷ in India. These differences are primarily the outcomes of differences in community-level development, population composition, state health expenditure, poverty levels, status of women and availability, accessibility and affordability of maternal and child healthcare services and their utilisation.^{18–20}

While India's share to overall areca nut production and consumption remains at the top in the world, no attempts have been made to explore the patterns and determinants of the consumption of areca nut based on large-scale representative surveys. This study aims to examine the disparity and determinants of areca nut consumption, with and without tobacco using the nationally representative Global Adult Tobacco Survey (GATS) conducted in 2016–2017.

METHODS AND MATERIALS

Study design and participants

We used the nationally representative cross-sectional GATS 2016–2017, conducted in all 29 states and three union territories (UTs) of India.²¹ The study included whole GATS sample of 74 037 adults aged 15 and above. A multistage sampling design separately for rural and urban areas was adopted to draw a representative sample considering the 2011 census population figures. The person-level response rate was 96.0% (95.6% in urban areas and 96.3% in rural areas). The overall response rate, calculated as the product of response rates at the household and person level, was 92.9%.

The sampling was done independently in each state/UT; and within the state/UT, it was done independently for urban and rural areas. In urban areas, a three-stage sampling process was adopted. At the first level, the list of all the wards from all cities and towns of the state/UT constituted the urban sampling frame, from which a required sample of wards (primary sampling units—PSUs) was selected using probability proportional to size (PPS) sampling. At the second level, a list of all census enumeration blocks (CEBs) in each selected ward constituted the sampling frame from which one CEB was selected by PPS from each ward. At the third level, a list of all residential households in each selected CEB constituted the sampling frame, from which a sample of required number of households was selected.

In rural areas, a two-stage sampling process was adopted. At the first stage of sampling, all villages in the state/UT formed the sampling frame. All small villages having less than five households were removed from the sampling frame. Villages with 5 to 49 households as per Census of India, 2011 were linked with the neighbouring larger villages. The required number of PSUs (villages) within each stratum was selected according to PPS sampling. At the second stage, a list of all residential households in each selected village constituted the sampling frame, from which a sample of the required number of households was selected.

A household listing operation was carried out in each sample area. All large villages with 300 or more households were segmented into three or more segments (depending on village size) of almost equal proportions, each being about 100–200 households. From all the segments in each large village, two segments were selected by using PPS sampling. Thirty households (plus three more, accounting for nonresponse) were selected from the list of households by systematic random sampling. The 33 selected households in a PSU were divided into two groups: (1) households for interview of a male member and (2) households for interview of a female member; this was in proportion to the total sample size of male and female interviews in a state. From the total number of male/female members aged 15 or above in a household, one member was randomly selected for the interview.

Further details related to survey methodology, sampling design, household and individual selection,

data collection, management and monitoring procedures have been described elsewhere.²¹

Dependent variables

The outcome variable was current consumption of areca nut use, assessed based on the following questions covered in the GATS:

- i. Do you consume *pan masala* without tobacco? (response options: yes, no and refused)
- ii. Do you consume betel quid without tobacco? (response options: yes, no and refused)
- iii. Do you consume areca nut of any type, plain, powdered or flavoured? (response options: yes, no and refused)
- iv. Betel quid with tobacco? (response: on average, how many times a day do you use).
- v. *Gutka*, areca nut–tobacco lime mixture, or *mawa*? (response: on average, how many times a day do you use)
- vi. *Pan masala* with tobacco? (response: on average, how many times a day do you use).

Based on the above-mentioned questions asked in GATS, we constructed three sets of variables: (1) areca nut consumption only without tobacco, (2) areca nut consumption only with tobacco and (3) areca nut consumption with and without tobacco, dual use. Definition of specific products can be found with the GATS two national report.²²

Independent variables

A range of socioeconomic (education, occupation, caste, religious affiliation and wealth quintile), demographic (age, sex, marital status), awareness-related and contextual level variables included in this study, which were found to be associated with areca nut consumption in previous studies.^{23–27} These variables include age (categorised as 15–18, 19–23, 24–30, 31–40, 41–50, 51–60 and 60+) and sex as male and female. Individual's education was measured as: (1) no formal education, (2) below primary, (3) primary completed, (4) below secondary, (5) secondary completed, (6) completed higher secondary, (7) completed college/university and (8) completed postgraduate level. Individual's occupation on the other side was assessed based on self-reported information as (1) student, (2) government sector, (3) nongovernment sector, (4) casual/daily labourer, (5) self-employed, (6) homemaker, (7) retired and (8) unemployed.

A wealth index was calculated based on availability of electricity, flush toilet, radio, television, fixed telephone or cell phone, refrigerator, washing machine, moped/scooter/motorcycle and car using principal component analysis (PCA) methodology.²⁸ There are various ways to assign weighting values to the indicator variables. Ad hoc weights, such as assigning '1' for a bicycle, '3' for a motorcycle and '5' for a car or truck, work to a certain extent, but they are arbitrary and are difficult to assign when the wealth ordering is not readily apparent. For this reason, Filmer and Pritchett recommended using PCA to assign the indicator weights, the procedure that is used for the

wealth index.²⁹ This procedure first standardises the indicator variables (calculating z scores); then the factor coefficient scores (factor loadings) are calculated and, finally, for each household, the indicator values are multiplied by the loadings and summed to produce the household's index value. In this process, only the first of the factors produced is used to represent the wealth index. The resulting sum is itself a standardised score with a mean of 0 and an SD of 1.²⁸ Individuals were divided into five wealth quintiles based on their household score ranges from 1 being poorest to 5 being wealthiest, with each category representing 20% of the score.²⁸

A composite knowledge variable, which measures the poor health impact of SLT use, was constructed based on the following information asked in the survey: SLT causes serious illness (yes/no), SLT causes oral cancer (yes/no), SLT causes dental diseases (yes/no), SLT causes harm to fetus during pregnancy (yes/no) and do you think SLT leads to addiction (yes/no). The new knowledge variable was categorised as: (1) 'no, to all five awareness', (2) 'no, to at least one awareness' and (3) 'yes, to all five awareness'.

Caste (social group) as categorised based on individual's self-reporting as Scheduled Castes (SCs), Scheduled Tribes (STs), Other Backward Classes (OBCs) and others. This broad categorisation of caste is based on their socioeconomic disadvantage in education, health, nutrition and employment by federal government. For instance, a study has shown that as compared with other caste, children (age 2–5 years) and adolescents (age 6–18 years) belonging to STs had the greatest risk of mortality (OR=1.94, 95% CI 1.47 to 2.57), followed by those from SCs (OR=1.35, 95% CI 1.05 to 1.74) and OBCs (OR=1.33, 95% CI 1.05 to 1.67).¹⁷ Other studies have also shown lower enrolment and completion of education among SCs and STs due to various factors.^{30–31} Religion captures self-reported follower/believer of Hinduism, Islam, Christianity and others (which mainly include Sikhs, Jains, Buddhists and nonbelievers). The study also considered place of residence as rural and urban as well as all 29 states and three UTs in the analysis.

Analytical strategy

At first, prevalence of areca nut consumption with and without tobacco at national and subnational levels along with rural–urban and male–female differences was analysed. χ^2 tests were performed to examine whether variations in areca nut consumption across independent variables were statistically significant. To examine the associated between areca nut consumption with various socioeconomic and demographic characteristics, multinomial logistic regression was used. In the multinomial logit regression, it is assumed that log odds of outcome/dependent variable either follow linear form or non-binary form; each outcome/dependent variable is modelled relative to the baseline group or outcome.³² In this study, we have considered (1) 'non-areca nut user (baseline group)', (2) 'areca nut consumption only with tobacco',

(3) 'areca nut consumption only without tobacco' and (iv) 'areca nut consumption with and without tobacco, dual use'. The study reported the relative risk ratio along with 95% CIs.³³ We calculated the population burden based on GATS weighted sample population figures, which were provided in the GATS India report.²² The analysis was adjusted for sampling weights and multistage sampling design using *svy* command in STATA. Analysis was carried out in STATA, V.15³⁴

Patient and public involvement

No patients were involved in the development of the research question, the outcome measures or the design of the study.

RESULTS

Descriptive statistics

Of the 74037 respondents, 40265 (48.9%) were women and 33772 (51.1%) were men and 47549 (65.5%) individuals resided in rural areas. One out of four respondents had no formal education and nearly 78% were aware about the adverse health effects of SLT consumption (online supplemental table 1).

We found that overall, betel quid without tobacco (8.7%; 95% CI 6.7 to 10.2) was consumed largely, followed by areca nut of any type (8%; 95% CI 5.9 to 10.3) at the national level (table 1). Among men, the prevalence of *gutka*, areca nut–tobacco lime mixture or *mawa* was consumed the most (17.8%; 95% CI 15.1 to 20.2), whereas, among women, betel quid without tobacco was largely consumed (9.0%; 95% CI 6.1 to 11.9). In urban areas, both betel quid without tobacco and areca nut of any type were largely consumed, while in the rural areas, it was mainly betel quid without tobacco. Regional pattern suggests that betel quid with tobacco was predominately consumed in many north-eastern states, while betel quid without tobacco was mainly used in south (online supplemental table 2).

Regional disparity in areca nut consumption

We found 23.9 (95% CI 23.1 to 24.8) adults were consuming areca nut at national level and 14.2% (95% CI 13.6 to 14.9) were consuming areca nut without tobacco (table 2). Figure 1 shows considerable variations in areca nut consumption across states and UTs of India. In many states, areca nut consumption in any form was over 40% among men (like, Uttar Pradesh, Assam Meghalaya, Mizoram and Manipur) and women (like Karnataka and all north-eastern states except Nagaland). Areca nut consumption without tobacco was largely being consumed across north-eastern states, apart from other bigger states like Karnataka (28.8%; 95% CI 25.6 to 32.1), Tamil Nadu (25.5%; 95% CI 21.9 to 29.5) and Maharashtra (20%; 95% CI 17.0 to 22.5). Nearly 223.4million people out of the total 932 488 000 population aged 15 and above consume areca nut in India (table 3). The distribution of areca nut users both in terms of population and

Table 1 Prevalence (in %) and number of users of different types of areca nut in India, GATS 2016–2017

	Total		Men		Women		P value of difference between men and women (χ^2 test)	Urban		Rural		P value of difference between urban and rural areas (χ^2 test)
	% (95% CI)	Users (in 000)	% (95% CI)	Users (in 000)	% (95% CI)	Users (in 000)		% (95% CI)	Users (in 000)			
Tobacco products												
Pan masala without tobacco	4.8 (3.2 to 5.6)	44 759	6.2 (4.1 to 8.3)	57 814	3.2 (1.9 to 5.1)	29 840	<0.001	5.2 (3.1 to 7.7)	48 489	4.5 (2.2 to 6.8)	41 962	>0.005
Pan masala with tobacco	2.8 (1.6 to 3.8)	26 110	4.5 (2.8 to 5.7)	41 962	1.1 (0.6 to 2.1)	10 257	<0.001	2.3 (1.1 to 3.8)	21 447	3.1 (1.2 to 5.2)	28 907	>0.005
Betel quid without tobacco	8.7 (6.7 to 10.2)	81 126	8.4 (5.9 to 10.8)	78 329	9.0 (6.1 to 11.9)	83 924	<0.001	9.1 (6.8 to 12.6)	84 856	8.4 (5.7 to 10.8)	78 329	>0.005
Betel quid with tobacco	5.8 (3.8 to 7.2)	54 084	7.1 (5.2 to 9.3)	66 207	5.5 (3.2 to 7.8)	51 287	<0.001	4.3 (2.8 to 6.3)	40 097	6.6 (4.2 to 8.3)	61 544	<0.005
Areca nut of any type	8.0 (5.9 to 10.3)	74 599	8.3 (5.8 to 11.2)	77 397	7.7 (5.1 to 9.2)	71 802	<0.001	9.1 (6.5 to 13.1)	84 856	7.5 (4.8 to 9.8)	69 937	>0.005
Gutka, areca nut-tobacco lime mixture or mawa	6.8 (5.7 to 8.6)	63 409	17.8 (15.1 to 20.2)	100 709	2.7 (1.2 to 4.1)	25 177	<0.001	6.3 (3.1 to 8.7)	58 747	7.1 (4.5 to 9.8)	66 207	<0.005
GATS: Global Adult Tobacco Survey.												

GATS, Global Adult Tobacco Survey.

Table 2 Prevalence (in %) of areca nut use in different forms across states and UTs of India, GATS 2016–2017

States/UTs	Areca nut use without tobacco only	Areca nut use with tobacco only	Dual use	Any form
North region				
Jammu and Kashmir	0.5 (0.2 to 1.2)	0.9 (0.5 to 1.5)	0 (0.0 to 0.1)	1.4 (0.8 to 2.1)
Himachal Pradesh	1.2 (0.8 to 1.9)	0.3 (0.1 to 0.8)	0 (0.0 to 0.2)	1.5 (1.1 to 2.3)
Punjab	1 (0.6 to 1.6)	2 (0.9 to 4.2)	0.1 (0.0 to 0.6)	3.1 (1.8 to 5.2)
Chandigarh	1.7 (1.1 to 2.7)	1.6 (0.9 to 3.0)	0.1 (0.0 to 0.3)	3.4 (2.3 to 5.1)
Uttarakhand	17.5 (14.0 to 21.6)	3.8 (2.9 to 4.8)	1.3 (0.7 to 2.1)	22.6 (18.5 to 27.1)
Haryana	2.6 (1.7 to 4.0)	2.4 (1.0 to 5.3)	0.2 (0.1 to 0.6)	5.2 (3.3 to 8.1)
Delhi	15.7 (13.0 to 18.7)	2 (1.3 to 2.9)	1 (0.6 to 1.7)	18.7 (15.6 to 22.0)
Central region				
Rajasthan	7.5 (6.3 to 9.1)	6.5 (5.3 to 7.9)	1.4 (0.7 to 2.5)	15.4 (13.4 to 17.6)
Uttar Pradesh	18.3 (16.4 to 20.3)	12.4 (11.0 to 14.0)	3.5 (2.7 to 4.4)	34.2 (31.9 to 36.5)
Chhattisgarh	8.4 (6.4 to 10.8)	7.4 (5.9 to 9.3)	1.2 (0.7 to 1.8)	17 (14.2 to 20.1)
Madhya Pradesh	8.6 (7.2 to 10.2)	13.4 (11.3 to 15.9)	2.1 (1.4 to 3.0)	24.1 (21.2 to 27.2)
East region				
West Bengal	13.4 (10.7 to 16.7)	4.1 (3.0 to 5.5)	3.7 (2.8 to 5.0)	21.2 (18.2 to 24.6)
Jharkhand	8.3 (6.6 to 10.3)	8.2 (6.4 to 10.5)	1.1 (0.6 to 2.0)	17.6 (14.3 to 21.4)
Odisha	11.3 (8.6 to 14.6)	11.5 (9.6 to 13.8)	4.8 (3.4 to 6.6)	27.6 (23.1 to 32.6)
Bihar	7 (5.4 to 9.1)	5 (4.0 to 6.3)	0.5 (0.3 to 0.9)	12.5 (10.5 to 14.8)
Northeast region				
Sikkim	11.5 (8.9 to 14.9)	1.5 (0.9 to 2.6)	0.7 (0.4 to 1.2)	13.7 (10.8 to 17.4)
Arunachal Pradesh	17.4 (14.4 to 20.8)	18.6 (13.0 to 25.8)	3.7 (2.6 to 5.2)	39.7 (32.6 to 47.0)
Nagaland	9.6 (7.6 to 12.2)	15.1 (12.7 to 17.8)	5.8 (4.3 to 7.8)	30.5 (27.3 to 34.0)
Manipur	23.7 (20.2 to 27.5)	21 (18.3 to 24.0)	6 (4.6 to 7.8)	50.7 (47.0 to 54.4)
Mizoram	52.9 (48.3 to 57.5)	0.8 (0.4 to 1.7)	4.4 (3.2 to 5.9)	58.1 (53.0 to 62.9)
Tripura	14.3 (11.6 to 17.4)	17 (14.0 to 20.6)	14 (11.0 to 17.7)	45.3 (42.0 to 48.7)
Meghalaya	63.2 (57.5 to 68.5)	1.6 (0.9 to 2.8)	6 (4.3 to 8.2)	70.8 (65.4 to 75.5)
Assam	45.6 (42.8 to 48.4)	11.4 (9.9 to 13.2)	10.6 (9.2 to 12.2)	67.6 (64.7 to 70.5)
West region				
Gujarat	8.3 (6.5 to 10.5)	11.1 (8.8 to 13.9)	1.1 (0.8 to 1.7)	20.5 (17.8 to 23.5)
Maharashtra	19.6 (17.0 to 22.5)	6.9 (5.5 to 8.8)	2.9 (2.0 to 4.2)	29.4 (25.3 to 34.0)
Goa	17.3 (14.6 to 20.4)	1.3 (0.7 to 2.2)	1.1 (0.7 to 1.9)	19.7 (16.7 to 23.1)
South region				
Andhra Pradesh	6.7 (5.2 to 8.7)	1.3 (0.7 to 2.4)	1.2 (0.6 to 2.4)	9.2 (7.1 to 11.7)
Telangana	8.7 (6.9 to 11.0)	2.8 (1.8 to 4.3)	2 (1.2 to 3.3)	13.5 (11.3 to 16.2)
Karnataka	28.8 (25.6 to 32.1)	7.7 (6.2 to 9.4)	4.3 (3.4 to 5.3)	40.8 (36.3 to 45.2)
Kerala	3.1 (2.2 to 4.3)	2.6 (1.9 to 3.6)	0.4 (0.2 to 1.0)	6.1 (4.7 to 7.9)
Tamil Nadu	25.5 (21.9 to 29.5)	3.7 (2.5 to 5.6)	1.2 (0.8 to 1.7)	30.4 (27.1 to 34.0)
Puducherry	17.7 (14.7 to 21.1)	1.9 (1.0 to 3.4)	1.4 (0.9 to 2.1)	21 (17.6 to 24.9)
India	14.2 (13.6 to 14.9)	7.3 (6.9 to 7.7)	2.4 (2.2 to 2.7)	23.9 (23.1 to 24.8)

GATS, Global Adult Tobacco Survey; UTs, union territories.

proportion across states was as follows: Uttar Pradesh with 49.9 million users contributes to nearly 22% of all areca nut users, followed by Maharashtra with 26.7 million users (12%), Karnataka with 19.8 million (9%) and Tamil Nadu

with 17.7 million users (8%). Together, these four states share nearly 51% of all areca nut users in the country. Not much difference exists between urban and rural areas in areca nut usage patterns (online supplemental table

Figure 1 Geographical variation in areca nut use among adult men and women in India. GATS, Global Adult Tobacco Survey.

3). In 18 states/UTs, however, areca nut consumption was higher in urban areas than rural counterparts. In 13 states/UTs, the opposite pattern was evident.

Demographic and socioeconomic differences in areca nut consumption

Consumption of areca nut in any form was higher among men as compared with women both at national level as well as in a majority of states (online supplemental table 4). All forms of areca nut consumption were higher in the age group 31–50 years (table 4) as compared with other age categories. 28.8% men (95% CI 27.7 to 30.0) and 27.1% widowed/separated/divorced (95% CI 25.3 to 29.1) were consuming areca nut. Individuals who had completed below the primary level of schooling consumed higher proportion of areca nut. Areca nut consumption was highest among daily wage labourers (30.2%; 95% CI 28.7 to 31.7). We found that a high percentage of STs (25.6%; 95% CI 23.0 to 27.5) and Muslims (30.8%; 95% CI 28.4 to 33.2) were consuming areca nut.

Determinants of areca nut consumption: regression analysis

Regression results suggest that as compared with 15–18 age group, the likelihood of areca nut consumption with tobacco and dual use was higher in higher age groups (table 5); except that areca nut consumption without tobacco was lower among the age group 51 and above, as compared with the 15–18 age groups. Probability of areca nut consumption was higher among men as compared with women for all three forms. The likelihood of areca nut consumption without tobacco was higher across all the educational categories as compared with those who had no formal education. However, the probability of

areca nut consumption with tobacco and in dual-form was declining with increase in the education level of respondents. The likelihood of areca nut consumption with tobacco and dual-use was significantly higher among SCs than Other castes. Probability of all the three forms of areca nut consumption was higher among Muslims as compared with Hindus.

DISCUSSION

The findings of the study revealed that nearly one out of every four adults in India consumes areca nut, that is, almost 223.79 million users, making areca nut consumption a bigger public health challenge than use of SLT (199 million users) in dealing with substance use and addiction in the country. The large number of users of areca nut, a known carcinogen, presents a huge public health challenge for the country. Moreover, nearly 10% consume areca nut with tobacco. Thus, considering the wide range of adverse health impacts, effective implementation on banning of tobacco as an ingredient with areca nut products under regulation 2.3.4 of the Food Safety and Standards Regulation, 2011 and ban on manufacture and sale of areca nut products, as implemented in some of the states, is urgently needed.¹

We found considerable regional and socioeconomic differences in the consumption of areca nut. In four states, Meghalaya, Assam, Mizoram and Manipur, over half of the population consume areca nut. Furthermore, Karnataka, Uttar Pradesh, Tamil Nadu, Maharashtra and Odisha constitute nearly 55% of the country's areca nut users. As far as other determinants are concerned,

Table 3 Population and share of areca nut use by states and UT of India, GATS 2016–2017

States/UTs	Population	Share (in %)
Chandigarh	33 040	0.0
Sikkim	68 448	0.0
Himachal Pradesh	88 112	0.0
Jammu and Kashmir	121 264	0.1
Puducherry	211 680	0.1
Goa	237 779	0.1
Arunachal Pradesh	415 800	0.2
Nagaland	459 940	0.2
Mizoram	488 040	0.2
Punjab	699 081	0.3
Haryana	1 048 632	0.5
Manipur	1 131 624	0.5
Tripura	1 316 418	0.6
Meghalaya	1 493 184	0.7
Kerala	1 650 843	0.7
Uttarakhand	1 756 575	0.8
Delhi	2 761 914	1.2
Chhattisgarh	3 262 714	1.5
Andhra Pradesh	3 654 056	1.6
Telangana	3 809 088	1.7
Jharkhand	4 261 840	1.9
Rajasthan	7 900 200	3.5
Odisha	8 984 904	4.0
Bihar	9 095 000	4.1
Gujarat	9 813 760	4.4
Madhya Pradesh	13 145 827	5.9
West Bengal	15 475 728	6.9
Assam	15 833 272	7.1
Tamil Nadu	17 753 296	7.9
Karnataka	19 834 738	8.9
Maharashtra	26 753 412	12.0
Uttar Pradesh	49 932 289	22.3
India	223 797 120	100.0

GATS, Global Adult Tobacco Survey; UTs, union territories.

the findings confirmed that age, gender, marital status, education, occupation, castes and religion are significantly associated with areca nut consumption. However, the direction of association differs with respect to areca nut consumption with and without tobacco. Cheaper and abundant availability, due to large-scale domestic production of areca nut, could be one of the key reasons for such high prevalence in the country.

We found protective effect of secondary and above level education in the case of areca nut consumption with and without tobacco. A study from Pakistan also observed

that the consumption of areca nut users increased by grade among school children aged 4–16 years.²⁷ Areca nut consumption was higher among men than among women, a finding that is consistent with other studies conducted in Tamil Nadu and Assam in India^{26 35} and countries like Thailand and Taiwan.¹³ It may be because areca nut consumption results in staining of teeth, which may not be liked by young and adult women. The age-wise pattern suggests that areca nut consumption without tobacco began to decline from age 51 onwards. But in the case of areca nut consumption with tobacco and in both forms, it increased with age.

Similar to other studies from India and other neighbouring countries,^{24 36} we also observed higher consumption of areca nut with tobacco among daily wage/casual labourers. This study further adds that areca nut consumption without tobacco too was largely consumed by daily wage/casual labourers, followed by nongovernment sector. Evidences suggest that many misconceptions including consuming areca nut improve concentration, pleasure, helps in anxiety and muscle relaxation and suppresses appetite increases the likelihood of consumption among those who are engaged in casual labour and have long working hours.^{23 37 38} We found higher consumption of areca nut among STs and SCs than other caste groups. Furthermore, Muslims were more likely to consume all three forms of areca nut as compared with Hindus. Previous studies documented higher consumption of tobacco including SLT, among SCs/STs and Muslims.^{25 39}

Urban–rural differences by state suggest that in 18 states, areca nut consumption was higher in urban areas than in rural areas. Regression results also revealed higher consumption of areca nut without tobacco in urban areas than rural counterparts. This is likely due to higher awareness about harms related to tobacco use in urban areas than rural counterparts. Studies from India and Pakistan documented that *pan masala* and *gutka* are very popular even in urban areas due to aggressive advertising, targeting middle class and adolescents, which improved sale many tobacco and related products including areca nut.⁴⁰

Our study had some limitations. Information related to areca nut consumption in different forms in the GATS was based on respondents self-reporting. Thus, the study cannot rule out social desirability bias—a tendency among some people to respond to questions in a way which they deem to be more acceptable than would be their ‘correct’ answer.⁴¹ The nomenclature of various areca nut products in geographically diverse country like India could be a source of concern, which is difficult to capture in the large-scale surveys. Considering the cross-sectional design of the survey, we did not examine the cause and effect relationship between socioeconomic characteristics and areca nut consumption. Similarly, the available data did not allow us to estimate trends of areca nut usage over time, but future analyses of repeated GATS may inform on important trends. Another limitation is that the study

Table 4 Areca nut use pattern by demographic and socioeconomic characteristics, GATS 2016–2017

Background variables	Areca nut use without tobacco only	Areca nut use with tobacco only	Dual use	Any forms
	% 95% CI	% 95% CI	% 95% CI	% 95% CI
Age	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
15–18	15.7 (13.9 to 17.6)	1.9 (1.4 to 2.5)	0.7 (0.5 to 1.2)	18.3 (16.5 to 20.3)
19–23	14.8 (13.3 to 16.4)	5.4 (4.5 to 6.4)	1.3 (0.9 to 1.9)	21.5 (19.8 to 23.3)
24–30	14.1 (13.1 to 15.2)	8.2 (7.3 to 9.1)	2.4 (2.0 to 2.9)	24.7 (23.3 to 26.1)
31–40	14.6 (13.6 to 15.6)	9.3 (8.6 to 10.1)	2.7 (2.4 to 3.2)	26.7 (25.4 to 28.0)
41–50	15.2 (14.1 to 16.3)	8.1 (7.3 to 9.0)	3.1 (2.6 to 3.7)	26.4 (25.0 to 27.9)
51–60	13.5 (12.2 to 14.8)	8 (7.0 to 9.0)	3.2 (2.6 to 3.9)	24.6 (23.1 to 26.3)
60+	11.1 (9.9 to 12.3)	7.8 (6.6 to 9.0)	3.2 (2.6 to 4.0)	22 (20.2 to 23.9)
Sex	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
Female	13.2 (12.5 to 14.0)	3.4 (3.0 to 3.8)	2.3 (2.0 to 2.6)	18.9 (18.0 to 19.9)
Male	15.2 (14.4 to 16.1)	11 (10.4 to 11.8)	2.6 (2.3 to 2.9)	28.8 (27.7 to 30.0)
Marital Status	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
Married	13.7 (13.1 to 14.4)	8.1 (7.6 to 8.6)	2.7 (2.4 to 3.0)	24.5 (23.6 to 25.4)
Unmarried	15.8 (14.6 to 17.1)	4.6 (3.9 to 5.3)	1 (0.8 to 1.4)	21.4 (20.0 to 22.9)
Widowed/Separated/ Divorced	14.3 (12.8 to 15.9)	8.3 (7.1 to 9.7)	4.5 (3.7 to 5.5)	27.1 (25.3 to 29.1)
Education	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
No formal education	11.2 (10.3 to 12.1)	8.7 (8.0 to 9.6)	3.7 (3.2 to 4.2)	23.6 (22.4 to 25.0)
<Primary completed	14.7 (13.4 to 16.1)	9.7 (8.6 to 10.9)	2.9 (2.4 to 3.6)	27.4 (25.6 to 29.2)
Primary completed	15.4 (14.1 to 16.9)	10 (8.9 to 11.2)	2.5 (2.0 to 3.0)	27.9 (26.2 to 29.7)
<Secondary completed	16 (14.7 to 17.3)	8.5 (7.6 to 9.5)	2.4 (1.9 to 2.9)	26.8 (25.3 to 28.3)
Secondary completed	15.1 (13.8 to 16.5)	5.1 (4.4 to 5.8)	1.7 (1.3 to 2.4)	21.9 (20.5 to 23.5)
Higher Secondary completed	15.6 (14.0 to 17.3)	4.4 (3.6 to 5.3)	1.4 (0.9 to 2.1)	21.4 (19.7 to 23.3)
College/University completed	15.5 (13.9 to 17.3)	3 (2.3 to 3.8)	1.1 (0.7 to 1.7)	19.6 (17.8 to 21.5)
Post-graduate completed	12.8 (10.7 to 15.2)	2.8 (1.6 to 4.6)	0.9 (0.4 to 2.4)	16.5 (14.0 to 19.3)
Occupation	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
Student	14.8 (13.2 to 16.6)	0.8 (0.5 to 1.2)	0.3 (0.2 to 0.6)	15.8 (14.2 to 17.6)
Government Employee	17.4 (14.9 to 20.3)	6.7 (5.1 to 8.8)	1.9 (1.1 to 3.4)	26.1 (23.2 to 29.2)
Non-government Employee	16.9 (15.3 to 18.6)	10.4 (8.7 to 12.3)	2.7 (2.0 to 3.8)	30 (27.7 to 32.5)
Daily Wage/Casual Labourer	15.3 (14.2 to 16.5)	11.4 (10.4 to 12.4)	3.5 (3.0 to 4.1)	30.2 (28.7 to 31.7)
Self-employed	14.9 (13.8 to 16.2)	11.7 (10.7 to 12.7)	3.1 (2.7 to 3.7)	29.7 (28.2 to 31.4)
Homemaker	12.4 (11.6 to 13.3)	3.5 (3.1 to 4.1)	1.9 (1.6 to 2.3)	17.9 (16.9 to 18.9)
Retired	9.1 (7.2 to 11.6)	6.3 (4.1 to 9.5)	2.6 (1.2 to 5.7)	18 (14.7 to 22.0)
Unemployed able to work	14.6 (11.9 to 17.8)	6.5 (4.6 to 9.2)	2.8 (1.7 to 4.6)	23.9 (20.3 to 28.0)
Unemployed unable to work	10.7 (8.5 to 13.2)	7 (5.2 to 9.4)	3.4 (2.1 to 5.4)	21.1 (18.2 to 24.4)
Knowledge of adverse health impact of SLT use	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
No	11.5 (8.3 to 15.7)	7.7 (5.6 to 10.4)	3.7 (2.2 to 6.1)	22.9 (18.6 to 27.8)
Partial	15.2 (14.1 to 16.3)	9 (8.2 to 9.9)	3.4 (2.9 to 3.9)	27.5 (26.1 to 29.0)
Full	14 (13.4 to 14.8)	6.8 (6.4 to 7.3)	2.2 (2.0 to 2.4)	23.1 (22.2 to 24.0)

Continued

Table 4 Continued

Background variables	Areca nut use without tobacco only	Areca nut use with tobacco only	Dual use	Any forms
	% 95% CI	% 95% CI	% 95% CI	% 95% CI
Caste	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
Others	14.2 (13.2 to 15.2)	6.1 (5.5 to 6.9)	2.6 (2.2 to 3.0)	22.9 (21.6 to 24.3)
Scheduled Castes	12.3 (11.1 to 13.5)	8.4 (7.5 to 9.3)	2.6 (2.2 to 3.0)	23.2 (21.8 to 24.6)
Scheduled Tribes	14.3 (12.7 to 16.0)	8.5 (7.2 to 9.5)	2.8 (2.1 to 3.3)	25.6 (23.0 to 27.5)
Oother Backward Castes	15.1 (14.2 to 16.1)	7.3 (6.8 to 7.9)	2.3 (2.0 to 2.6)	24.7 (23.5 to 25.9)
Religion	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
Hindu	13.6 (12.9 to 14.3)	7.3 (6.9 to 7.8)	2.3 (2.1 to 2.6)	23.2 (22.3 to 24.1)
Muslim	19 (17.2 to 20.9)	8.5 (7.3 to 9.9)	3.3 (2.8 to 4.0)	30.8 (28.4 to 33.2)
Christian	16.2 (13.6 to 19.1)	3.5 (2.8 to 4.4)	2.3 (1.4 to 3.7)	21.9 (18.9 to 25.3)
Others	8.4 (6.5 to 10.7)	4.1 (2.7 to 6.0)	1.7 (1.0 to 3.1)	14.1 (11.4 to 17.4)
Wealth Quintile	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
Poorest	12.1 (11.1 to 13.1)	9.6 (8.8 to 10.4)	3.1 (2.7 to 3.6)	24.7 (23.4 to 26.2)
Poorer	13.7 (12.8 to 14.8)	8.6 (7.8 to 9.4)	2.9 (2.5 to 3.4)	25.2 (23.9 to 26.6)
Middle	14.6 (13.5 to 15.8)	8.2 (7.3 to 9.2)	2.4 (1.9 to 2.9)	25.2 (23.7 to 26.8)
Richer	16.1 (14.8 to 17.5)	5.4 (4.7 to 6.1)	2 (1.5 to 2.6)	23.4 (21.9 to 25.0)
Richest	15.7 (14.3 to 17.3)	2.7 (2.2 to 3.3)	1.1 (0.8 to 1.6)	19.5 (18.0 to 21.1)
Place of Residence	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
Urban	15.7 (14.6 to 16.9)	6.1 (5.4 to 6.8)	2 (1.7 to 2.4)	23.8 (22.3 to 25.4)
Rural	13.5 (12.7 to 14.2)	7.9 (7.4 to 8.5)	2.7 (2.4 to 3.0)	24.1 (23.1 to 25.0)
Region	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)	(χ^2 p value<0.001)
North	5.6 (4.9 to 6.4)	2 (1.4 to 2.8)	0.4 (0.3 to 0.6)	8 (7.0 to 9.1)
Central	13.6 (12.5 to 14.8)	11.1 (10.2 to 12.1)	2.6 (2.2 to 3.2)	27.3 (25.9 to 28.8)
East	10.1 (8.9 to 11.6)	6.1 (5.4 to 6.9)	2.4 (2.0 to 3.0)	18.7 (17.1 to 20.4)
Northeast	39.9 (37.8 to 42.0)	11.9 (10.8 to 13.2)	9.6 (8.6 to 10.7)	61.4 (59.3 to 63.5)
West	15.7 (13.8 to 17.8)	8.3 (7.0 to 9.8)	2.3 (1.6 to 3.2)	26.3 (23.3 to 29.5)
South	17.3 (15.8 to 18.8)	3.9 (3.3 to 4.6)	1.9 (1.6 to 2.3)	23.1 (21.4 to 24.9)

χ^2 p value<0.001.

GATS, Global Adult Tobacco Survey.

is based on 15 years and older population, whereas the areca nut habits often start at younger age. The future Global Youth Tobacco Survey should have areca nut-related questions similar to GATS, so that detailed usage pattern of areca nut could be examined among younger population of the country.

CONCLUSION

It is now well established that areca nut consumption in any form is highly addictive, a well-known risk factor for oral, pharynx and oesophageal cancers and is associated with many adverse health effects. This study adds to the existing knowledge that areca nut consumption in India was much higher than the overall SLT. Moreover, a significant proportion of areca nut was consumed along with tobacco, which elevates the adverse health impacts and

comorbidities further. Thus, it calls for urgent policy intervention to prevent both new generations from taking up areca nut consumption habit and helping current users to quit. Such policy efforts to control areca nut consumption should be guided by the huge differences in its consumption across states, gender and socioeconomic groups in India. Unlike tobacco, for which the WHO FCTC provides evidence-based policies, no global policy exists for the regulation and control of areca nut consumption and its cessation. Also, there is a need for further research and population-based interventions to find treatment for areca nut dependence. In addition, research is needed to examine the intention to quit among areca nut users, separately for all three categories—those who consume areca nut with tobacco, without tobacco and those who consume in both the forms—to develop an appropriate

Table 5 Multinomial regression analysis showing determinants of areca nut use, India GATS 2016–2017

Background variables	Areca nut use without tobacco only			Areca nut use with tobacco only			Dual use		
	RRR	95% CI	P value	RRR	95% CI	P value	RRR	95% CI	P value
Age									
15–18 (ref.)	1.00			1.00			1.00		
19–23	0.91	0.81 to 1.02	0.106	1.73	1.33 to 2.25	<0.001	1.53	1.03 to 2.26	0.034
24–30	0.92	0.81 to 1.04	0.202	2.37	1.83 to 3.08	<0.001	2.26	1.54 to 3.33	<0.001
31–40	0.92	0.81 to 1.04	0.191	2.71	2.08 to 3.52	<0.001	2.72	1.84 to 4.03	<0.001
41–50	0.91	0.80 to 1.04	0.159	2.42	1.85 to 3.16	<0.001	2.59	1.74 to 3.86	<0.001
51–60	0.77	0.67 to 0.89	<0.001	2.14	1.62 to 2.81	<0.001	2.53	1.68 to 3.80	<0.001
60+	0.65	0.56 to 0.75	<0.001	2.04	1.54 to 2.70	<0.001	2.36	1.55 to 3.58	<0.001
Sex									
Female (ref.)	1.00			1.00			1.00		
Male	1.13	1.07 to 1.20	<0.001	2.02	1.85 to 2.21	<0.001	1.81	1.72 to 1.92	0.001
Marital status									
Married (ref.)	1.00			1.00			1.00		
Unmarried	0.94	0.86 to 1.02	0.160	1.06	0.93 to 1.20	0.382	1.12	0.92 to 1.36	0.273
Widowed/separated/divorced	1.37	1.24 to 1.50	<0.001	1.62	1.42 to 1.83	<0.001	1.59	1.35 to 1.88	<0.001
Education									
No formal education (ref.)	1.00			1.00			1.00		
<Primary completed	1.23	1.14 to 1.34	<0.001	1.01	0.91 to 1.12	0.867	0.93	0.80 to 1.08	0.318
Primary completed	1.20	1.11 to 1.30	<0.001	1.06	0.96 to 1.17	0.264	0.82	0.70 to 0.96	0.016
<Secondary completed	1.37	1.27 to 1.48	<0.001	0.99	0.89 to 1.09	0.774	0.90	0.77 to 1.05	0.177
Secondary completed	1.23	1.13 to 1.34	<0.001	0.79	0.70 to 0.89	<0.001	0.75	0.62 to 0.91	0.003
Higher secondary completed	1.23	1.11 to 1.35	<0.001	0.74	0.64 to 0.86	<0.001	0.74	0.59 to 0.93	0.011
College/university completed	1.18	1.06 to 1.31	0.003	0.53	0.44 to 0.63	<0.001	0.53	0.40 to 0.71	<0.001
Post-graduate completed	1.18	1.02 to 1.36	0.023	0.35	0.25 to 0.48	<0.001	0.43	0.27 to 0.68	<0.001
Occupation									
Student (ref.)	1.00			1.00			1.00		
Government employee	1.15	0.99 to 1.34	0.06	3.43	2.49 to 4.73	<0.001	2.56	1.63 to 4.0	<0.001
Non-government employee	1.59	1.39 to 1.81	<0.001	4.87	3.64 to 6.53	<0.001	4.17	2.75 to 6.30	<0.001
Daily wage/casual labourer	1.65	1.46 to 1.87	<0.001	4.51	3.40 to 5.99	<0.001	3.95	2.66 to 5.87	<0.001
Self-employed	1.30	1.15 to 1.47	<0.001	4.34	3.27 to 5.75	<0.001	2.87	1.94 to 4.27	<0.001
Homemaker	1.26	1.11 to 1.42	<0.001	2.61	1.95 to 3.49	<0.001	2.11	1.42 to 3.13	<0.001
Retired	0.95	0.77 to 1.17	0.604	2.28	1.56 to 3.33	<0.001	2.21	1.29 to 3.77	0.004
Unemployed able to work	0.91	0.76 to 1.08	0.287	2.91	2.09 to 4.05	<0.001	2.04	1.28 to 3.25	0.003
Unemployed unable to work	1.24	1.01 to 1.52	0.039	2.56	1.79 to 3.67	<0.001	2.49	1.51 to 4.09	<0.001

Continued

Table 5 Continued

Background variables	Areca nut use without tobacco only			Areca nut use with tobacco only			Dual use		
	RRR	95% CI	P value	RRR	95% CI	P value	RRR	95% CI	P value
Knowledge of adverse health impact of SLT use									
No (ref.)	1.00			1.00			1.00		
Partial	1.37	1.13 to 1.66	0.001	1.36	1.06 to 1.76	0.017	1.60	1.12 to 2.28	0.009
Full	1.22	1.01 to 1.48	0.056	1.04	0.81 to 1.34	0.759	1.23	0.87 to 1.74	0.249
Caste									
Others (ref.)	1.00			1.00			1.00		
Scheduled castes	1.05	0.97 to 1.13	0.197	1.17	1.05 to 1.29	0.004	1.25	1.07 to 1.46	0.005
Scheduled tribes	1.11	1.02 to 1.20	0.016	0.96	0.85 to 1.08	0.459	0.91	0.77 to 1.08	0.295
Other backward castes	0.95	0.89 to 1.01	0.087	0.96	0.88 to 1.05	0.346	0.80	0.70 to 0.92	0.001
Religion									
Hindu (ref.)	1.00			1.00			1.00		
Muslim	1.35	1.26 to 1.45	<0.001	1.22	1.11 to 1.35	<0.001	1.41	1.22 to 1.63	<0.001
Christian	0.83	0.77 to 0.91	<0.001	0.59	0.52 to 0.68	<0.001	0.58	0.49 to 0.69	<0.001
Others	0.61	0.55 to 0.68	<0.001	0.50	0.42 to 0.60	<0.001	0.36	0.27 to 0.47	<0.001
Wealth quintile									
Poorest (ref.)	1.00			1.00			1.00		
Poorer	0.94	0.88 to 1.00	0.062	0.96	0.88 to 1.04	0.321	1.11	0.98 to 1.25	0.097
Middle	1.02	0.95 to 1.11	0.573	0.97	0.88 to 1.08	0.628	1.00	0.85 to 1.17	0.964
Richer	1.03	0.96 to 1.12	0.403	0.79	0.70 to 0.89	<0.001	0.79	0.66 to 0.95	0.01
Richest	1.04	0.94 to 1.15	0.305	0.54	0.46 to 0.63	<0.001	0.83	0.67 to 1.03	0.096
Place of residence									
Urban (ref.)	1.00			1.00			1.00		
Rural	0.94	0.90 to 0.99	0.024	0.94	0.87 to 1.01	0.092	1.07	0.95 to 1.20	0.262
Region									
North (ref.)	1.00			1.00			1.00		
Central	2.28	2.07 to 2.51	<0.001	6.08	5.26 to 7.04	<0.001	6.45	4.78 to 8.71	<0.001
East	2.01	1.82 to 2.22	<0.001	3.38	2.90 to 3.95	<0.001	6.25	4.64 to 8.44	<0.001
Northeast	11.85	10.84 to 12.95	<0.001	14.81	12.77 to 17.18	<0.001	51.51	38.9 to 68.23	<0.001
West	3.4	3.10 to 3.73	<0.001	4.84	4.14 to 5.65	<0.001	5.85	4.26 to 8.03	<0.001
South	3.67	3.37 to 4.01	<0.001	2.29	1.95 to 2.68	<0.001	6.14	4.57 to 8.25	<0.001

GATS, Global Adult Tobacco Survey; Ref, Reference; RRR, relative risk ratio; SLT, smokeless tobacco.

intervention model for cessation. This information may be collected within the GATS survey by adding a few additional questions on areca nut for future analysis. Given that areca nut consumption follows a complex pattern by socioeconomic status (SES) and regional trajectories, separately for with and without tobacco, future research is needed to explore the various intersections between SES and areca nut consumption in different regions of India to gain better clarity.

Author affiliations

¹Division of Preventive Oncology & Population Health, ICMR - National Institute of Cancer Prevention and Research, Noida, India

²WHO FCTC Global Knowledge Hub on Smokeless Tobacco, ICMR - National Institute of Cancer Prevention and Research, Noida, India

³The International Union Against Tuberculosis and Lung Disease (The Union) South East Asia Office, New Delhi, India

⁴ICMR - National Institute of Medical Statistics, New Delhi, India

⁵Centre for Health Economics, University of York, Heslington, York, UK

⁶School of Preventive Oncology, Patna, India

⁷Boston College USA, Boston, Massachusetts, USA

⁸ISGlobal, Barcelona, Spain

⁹ICMR - National Institute of Cancer Prevention and Research, Noida, India

Twitter Prashant Kumar Singh @prashantks_pks and Amit Yadav @ayadav24

Contributors PKS conceived the study. PKS and LS performed the statistical analysis. PKS and LS analysed and interpreted the data. PKS and AY drafted the manuscript. SM, DNS, KS and SS provided comments and contributed to the development of the final draft of the manuscript. All authors have supervised and approved the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Map disclaimer The depiction of boundaries on the map(s) in this article does not imply the expression of any opinion whatsoever on the part of BMJ (or any member of its group) concerning the legal status of any country, territory, jurisdiction or area or of its authorities. The map(s) are provided without any warranty of any kind, either express or implied.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The second round of GATS obtained ethical clearance from the Ethics Committee of Tata Institute of Social Sciences. No ethics clearance was required for this study, as we performed a secondary analysis using publicly available data.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. Data are available upon reasonable request. Data used by the study are available by emailing prashants.geo@gmail.com.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Prashant Kumar Singh <http://orcid.org/0000-0003-1742-5534>

Sumit Mazumdar <http://orcid.org/0000-0002-9278-4396>

REFERENCES

- Gupta PC, Ray CS, Papke RL, et al. Perspectives on areca nut with some global implications: symposium report. *Translational Research in Oral Oncology* 2018;3:2057178X1881406.
- Warnakulasuriya S, Peters TJ. Introduction: biology, medical and socio-economic aspects of areca nut use. *Addict Biol* 2002;7:75–6.
- Mehrtash H, Duncan K, Parascandola M, et al. Defining a global research and policy agenda for betel quid and areca nut. *Lancet Oncol* 2017;18:e767–75.
- National Institute of Cancer Prevention and Research (ICMR). Commonly used smokeless tobacco products around the globe, 2020. Available: <https://untobaccocontrol.org/kh/smokeless-tobacco/paan-betel-quid-tobacco/>
- Patidar KA, Parwani R, Wanjar SP, et al. Various terminologies associated with areca nut and tobacco chewing: a review. *J Oral Maxillofac Pathol* 2015;19:69.
- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Betel-quid and areca-nut chewing and some areca-nut derived nitrosamines. *IARC Monogr Eval Carcinog Risks Hum* 2004;85:1–334.
- Guha N, Warnakulasuriya S, Vlaanderen J, et al. Betel quid chewing and the risk of oral and oropharyngeal cancers: a meta-analysis with implications for cancer control. *Int J Cancer* 2014;135:1433–43.
- Garg A, Chaturvedi P, Gupta PC. A review of the systemic adverse effects of areca nut or betel nut. *Indian J Med Paediatr Oncol* 2014;35:3.
- Mirza SS, Shafique K, Vart P, et al. Areca nut chewing and dependency syndrome: is the dependence comparable to smoking? A cross sectional study. *Subst Abuse Treat Prev Policy* 2011;6:23.
- Papke RL, Horenstein NA, Stokes C. Nicotinic activity of arecoline, the psychoactive element of "Betel Nuts", suggests a basis for habitual use and anti-inflammatory activity. *PLoS One* 2015;10:e0140907.
- Chandra PS, Mulla U. Areca nut: the hidden Indian 'gateway' to future tobacco use and oral cancers among youth. *Indian J Med Sci* 2007;61:319–21.
- Mehrotra R, Yadav A, Sinha DN, et al. Smokeless tobacco control in 180 countries across the globe: call to action for full implementation of WHO FCTC measures. *Lancet Oncol* 2019;20:e208–17.
- Gupta PC, Warnakulasuriya S. Global epidemiology of areca nut usage. *Addict Biol* 2002;7:77–83.
- James KS. India's Demographic Change: Opportunities and Challenges. *Science* 2011;333:576–80.
- Singh PK, Yadav A, Lal P. Dual burden of smoked and smokeless tobacco use in India, 2009–2017: a repeated cross-sectional analysis based on global adult tobacco survey. *Nicotine Tob Res* 2020.
- Singh PK. Trends in child immunization across geographical regions in India: focus on urban-rural and gender differentials. *PLoS One* 2013;8:e73102.
- Subramanian SV, Nandy S, Irving M, et al. The mortality divide in India: the differential contributions of gender, caste, and standard of living across the life course. *Am J Public Health* 2006;96:818–25.
- Kumar C, Singh PK, Rai RK. Under-five mortality in high focus states in India: a district level geospatial analysis. *PLoS One* 2012;7:e37515.
- Farahani M, Subramanian SV, Canning D. Effects of state-level public spending on health on the mortality probability in India. *Health Econ* 2010;19:1361–76.
- Kravalda Øystein. Child mortality in India: the community-level effect of education. *Popul Stud* 2004;58:177–92.
- TISS, MoHFW. *Global adult tobacco survey GATS 2 India 2016-17*. New Delhi, India, 2017.
- Tata Institute of Social Sciences and Ministry of Health and Family Welfare Government of India. Global adult tobacco survey GATS 2 India 2016-2017 report, 2017. Available: https://untobaccocontrol.org/kh/smokeless-tobacco/wp-content/uploads/sites/6/2018/06/GATS_2.pdf
- Ansari ZA, Bano SN, Zulkifli M. Prevalence of tobacco use among power loom workers - a cross-sectional study. *Indian J Community Med* 2010;35:34.
- Bandyopadhyay A, Irfan M. Educational and wealth inequalities in smokeless tobacco use: an analysis of rural-urban areas of Bangladesh and India. *Subst Abuse* 2019;13:117822181882507.
- Bhan N, Karan A, Srivastava S, et al. Have socioeconomic inequalities in tobacco use in India increased over time? trends from the national sample surveys (2000-2012). *Nicotine Tob Res* 2016;18:1711–8.

- 26 Rajan G, Ramesh S, Sankaralingam S. Areca nut use in rural Tamil Nadu: a growing threat. *Indian J Med Sci* 2007;61:332–7.
- 27 Shah SMA, Merchant AT, Luby SP, *et al*. Addicted schoolchildren: prevalence and characteristics of areca nut chewers among primary school children in Karachi, Pakistan. *J Paediatr Child Health* 2002;38:507–10.
- 28 Rutstein SO, Johnson K. The DHS wealth index. *DHS Comp. Reports* 2004;6. No.<http://dhsprogram.com/pubs/pdf/CR6/CR6.pdf>
- 29 Filmer D, Pritchett LH. Estimating wealth effects without expenditure data--or tears: an application to educational enrollments in states of India. *Demography* 2001;38:115–32.
- 30 Borooah VK. Caste, religion, and health outcomes in India, 2004–14. *Econ Polit Wkly* 2018;53:65.
- 31 Desai S, Dubey A. Caste in 21st century India: competing narratives. *Econ Polit Wkly* 2012;46:40.
- 32 Scott Long J. *Regression models for categorical and limited dependent variables*. 7. USA: Indiana University, 1997.
- 33 Haan P, Uhlenborff A. Estimation of multinomial logit models with unobserved heterogeneity using maximum simulated likelihood. *Stata J* 2006;6:229–45.
- 34 Stata S. *Release 15. statistical software*. Coll Station TX: StataCorp LP, 2017.
- 35 Mahanta B, Phukan N, Mohapatra PK. Betel nut chewing habit and its determinants among adolescent boys and girls in an industrial town of Assam, India. *International Journal of Health Sciences and Research*.
- 36 Hashibe M, Jacob BJ, Thomas G, *et al*. Socioeconomic status, lifestyle factors and oral premalignant lesions. *Oral Oncol* 2003;39:664–71.
- 37 Gunaseelan R, Sankaralingam S, Ramesh S, *et al*. Areca nut use among rural residents of Sriperambudur Taluk: a qualitative study. *Indian J Dent Res* 2007;18:11.
- 38 Thankappan KR, Thresia CU. Tobacco use & social status in Kerala. *Indian J Med Res* 2007;126:300.
- 39 Rani Met *al*. Tobacco use in India: prevalence and predictors of smoking and chewing in a national cross sectional household survey. *Tob Control* 2003;12:4e–4.
- 40 Tobacco Free Initiative, World Health Organization. Tobacco and youth in the South East Asian region. *Indian J Cancer* 2002;39:5–34.
- 41 Fisher RJ. Social desirability bias and the validity of indirect Questioning. *J Consum Res* 1993;20:303–15.