

BMJ Open Addressing maternal and child health equity through a community health worker home visiting intervention to reduce low birth weight: retrospective quasi-experimental study of the Arizona Health Start Programme

Samantha Sabo ,¹ Patrick Wightman,² Kelly McCue ,¹ Matthew Butler,³ Vern Pilling,⁴ Dulce J Jimenez ,¹ Martín Celaya,⁵ Sara Rumann⁵

To cite: Sabo S, Wightman P, McCue K, *et al.* Addressing maternal and child health equity through a community health worker home visiting intervention to reduce low birth weight: retrospective quasi-experimental study of the Arizona Health Start Programme. *BMJ Open* 2021;**11**:e045014. doi:10.1136/bmjopen-2020-045014

► Prepublication history and supplemental material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-045014>).

Received 19 September 2020
Revised 09 April 2021
Accepted 12 April 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Samantha Sabo;
samantha.sabo@nau.edu

ABSTRACT

Objective To test if participation in the Health Start Programme, an Arizona statewide Community Health Worker (CHW) maternal and child health (MCH) home visiting programme, reduced rates of low birth weight (LBW), very LBW (VLBW), extremely LBW (ELBW) and preterm birth (PTB).

Design Quasi-experimental retrospective study using propensity score matching of Health Start Programme enrolment data to state birth certificate records for years 2006–2016.

Setting Arizona is uniquely racially and ethnically diverse with comparatively higher proportions of Latino and American Indian residents and a smaller proportion of African Americans.

Participants 7212 Health Start Programme mothers matched to non-participants based on demographic, socioeconomic and geographic characteristics, health conditions and previous birth experiences.

Intervention A statewide CHW MCH home visiting programme.

Primary and secondary outcome measures LBW, VLBW, ELBW and PTB.

Results Using Health Start Programme's administrative data and birth certificate data from 2006 to 2016, we identified 7212 Health Start Programme participants and 53948 matches. Programme participation is associated with decreases in adverse birth outcomes for most subgroups. Health Start participation is associated with statistically significant lower rates of LBW among American Indian women (38%; average treatment-on-the-treated effect (ATT): 2.30; 95% CI –4.07 to –0.53) and mothers with a pre-existing health risk (25%; ATT: –3.06; 95% CI –5.82 to –0.30). Among Latina mothers, Health Start Programme participation is associated with statistically significant lower rates of VLBW (36%; ATT: 0.35; 95% CI –0.69 to –0.01) and ELBW (62%; ATT: 0.31; 95% CI (–0.52 to –0.10)). Finally, Health Start Programme participation is associated with a statistically significant lower rate of PTB for teen mothers (30%; ATT: 2.81; 95% CI –4.71 to –0.91). Other results were not statistically significant.

Strengths and limitations of this study

- The large sample sizes of the intervention and comparison groups enabled calculation of more reliable estimates of the Health Start Programme's effect with respect to relatively rare and infrequent outcomes for the programme as a whole and for select demographic subgroups.
- The primary limitation is the identifying assumption (common to all propensity-score matching analyses) that participation in the Health Start Programme is fully explained by observable characteristics.
- The analysis may have limited external validity for populations that differ along socioeconomic status, race and ethnicity.
- The study design and limited information on programme implementation mean the present study is not able to identify which specific programmatic features are associated with outcomes we find.

Conclusion A state health department-operated MCH home visiting intervention that employs CHWs as the primary interventionist may contribute to the reduction of LBW, VLBW, ELBW and PTB and could improve birth outcomes statewide, especially among women and children at increased risk for MCH inequity.

INTRODUCTION

Growing maternal and child health (MCH) inequities are largely associated with multi-level social and structural determinants of health, many of which are beyond the proximal control of any individual or community.^{1,2} Here, we focus on low birth weight (LBW), very LBW (VLBW) and extremely LBW (ELBW), generally accepted to be a result of preterm birth (PTB) or fetal growth restriction^{3,4} and associated with several interlocking

Percent Low Birthweight (< 2500 g) by County of Maternal Residence, 2010–2016




Figure 1 Low birth weight (LBW) in Arizona, 2010–2016. Map demonstrates the per cent of LBW, defined as <2500 g, by county of maternal residence in Arizona. Map courtesy of and permission by Arizona Health Start Programme, Arizona Department of Health Services. This map is not under copyright.

socioecological risk factors, including poverty, discrimination and access and quality of care among others.⁵ Rates of LBW, VLBW and ELBW in the USA consistently track with socioeconomic and ethnoracial health inequities.⁶ Women and children of colour, specifically African American, Latina and American Indian communities, experience disproportionately higher rates of LBW and VLBW^{7 8} (figures 1–3).

At the individual level, LBW, VLBW, ELBW and PTB are influenced by several behaviours associated with living in poverty and experiencing personally and institutionally mediated racism and discrimination, including smoking,⁹ poor nutrition,¹⁰ acute stress,^{11 12} prenatal depression^{13 14} and late or no prenatal care. Such factors may be exacerbated by interpersonal relationships, short interpregnancy intervals,¹⁵ relationship stress,¹¹ interpersonal violence¹² and lack of social support.¹² It is widely accepted that societal-level influences, including exposure to discrimination,¹² living in disadvantaged neighbourhoods,¹⁶ experiencing low socioeconomic status (SES)¹² and lacking access to health insurance,¹² all contribute to the cumulative physiological damage on the body caused by environmental stressors.¹⁷ From a life course perspective, LBW, VLBW, ELBW and PTB have several implications for health equity, including cost of care,¹⁸ decreased long-term educational attainment and earnings, and the predisposition for adult-onset chronic diseases.^{19–21} Furthermore, documented differences in birth weight and SES in the US compound health equity

through observed correlations in economic standing across generations.²²

For more than half a century, community health workers (CHWs) have served as trusted members of communities experiencing health inequity, defined as the unfair and preventable differences in health and well-being.²³ As a workforce, CHWs are trusted intermediaries between people and systems and are consistently effective within clinical-based and community-based contexts and with high priority health issues such as managing chronic diseases, improving birth outcomes and maintaining child wellness.^{23 24} Among national expansion of state and tribal health departments, Medicaid services, health plans and provider groups that promote CHW integration,²⁵ CHWs have emerged as a healthcare workforce able to enhance the patient experience of care, improve population health, reduce cost of care and improve the experience of providing care among clinicians and staff.^{26 27}

While early childhood home visitation has a goal of improving birth outcomes, studies using random assignment have seen significant challenges in doing so, even with registered nurses as the home visitors.²⁸ We conducted a quasi-experimental evaluation of a long-standing state health department-operated programme administered within county, clinic and community-based organisations in 14 rural and urban counties across Arizona to assess whether Health Start Programme participation is associated with improved birth and birth-related outcomes.²⁹ Our study aims to contribute to the gap in

Very Low Birthweight (< 1500 g) per 1,000 Live Births by County of Maternal Residence, 2010–2016




Figure 2 Very low birth weight (VLBW) in Arizona, 2010–2016. Map demonstrates the incidence of VLBW, defined as <1500 g, per 1000 live births by county of maternal residence in Arizona. Map courtesy of and permission by Arizona Health Start Programme, Arizona Department of Health Services. This map is not under copyright.

Percent Preterm Births (< 37 weeks) by County of Maternal Residence, 2010–2016




Figure 3 Preterm births (PTBs) in Arizona, 2010–2016. Map demonstrates the per cent of PTBs, defined as <37 weeks of gestation, by county of maternal residence in Arizona. Map courtesy of and permission by Arizona Health Start Programme, Arizona Department of Health Services. This map is not under copyright.

literature for rigorous evaluations of CHW-led home visiting programmes targeting MCH equity.

Established in 1994, the Arizona Health Start Programme is a primary prevention home visitation intervention aimed at improving MCH outcomes among medically and socially high risk, racially and ethnically diverse, rural and urban mothers with children aged 0–2 years.^{30 31} MCH home visiting services are delivered by trained CHWs who, through CHW core competency and specialised MCH training, connect clients to prenatal care and increase continuity of care during and after pregnancy.²⁹ Broadly, CHWs identify, screen and enrol eligible women; provide perinatal and postpartum education and social support; provide referral and advocacy services; and emphasise timely immunisations and developmental assessments for children.^{29 31 32} During pregnancy, CHWs provide education (eg, perinatal nutrition, physical symptoms and changes, labour and delivery, and breastfeeding); assist with access and enrolment to continuous perinatal care; screen, educate and follow-up for maternal behavioural health disorders, alcohol, tobacco and drug cessation; and intimate partner violence. The Health Start Programme's CHWs are trained to motivate and support their clients through behaviour change activities that promote personal agency and self-efficacy, and achieve Health Start Programmatic goals, including reducing LBW outcomes.

This paper is part of a broader research agenda assessing the impact of the Arizona Health Start Programme on newborn health, maternal healthcare utilisation and early child health. The specific objective of the analysis presented here is to assess the effectiveness of the Health Start Programme with respect to newborn health, particularly whether participation is associated with lower rates of LBW, VLBW, ELBW and PTB compared with non-participating women. Our methodological strategy is explicitly motivated and guided by Home Visiting Evidence of Effectiveness (HomVEE) standards, established by the US Department of Health and Human Services.³³

This evaluation builds on a previous assessment of the Health Start Programme, which found that programme participation was associated with a reduction in the likelihood of LBW.³¹ Accordingly, our primary hypothesis is that mothers and children that participated in this MCH home visitation programme during the study period of 2006–2016 will experience lower rates of LBW, VLBW, ELBW and PTB.

METHODS

Study design and data

This study uses a quasi-experimental design to estimate the impact of Health Start Programme participation on LBW, VLBW, ELBW and PTB. All data on births and mothers are derived from Arizona's Vital Records Birth Database, maintained by the Arizona Department of Health Services for the years 2006–2016. This information

was linked to Health Start Programme enrolment data to identify participants. Both the Health Start Programme enrolment information and birth certificate records are administrative data sources, established and maintained primarily for public health monitoring. Detailed Health Start Programme evaluation protocol, including the process used to merge the birth certificate records and Health Start Programme enrolment data, Health Start intervention components, and CHW training is published elsewhere.²⁹

Outcomes

The primary outcomes of our study are LBW, birth weight less than 2500g despite gestational age; VLBW, birth weight less than 1500g; ELBW, birth weight less than 1000g; and PTB, a gestational age of less than 37 completed weeks of pregnancy.

Patient and public involvement

Patients and the public were not involved in the design or conducting and reporting of the study.

Study population

Women can enrol in Health Start Programme during pregnancy and/or if they have a child of 2 years of age or younger and can remain in the programme as long as they maintain this criterion. Non-pregnant women enrolling with children who are 2 years of age or younger still receive home visits and preconception and inter-conception health education, which is associated with improved birth outcomes.^{28 34} We limited the treatment population to women who were enrolled prior to giving birth to the children whose birth certificates comprise our data source. This resulted in 7212 Health Start Programme-attributed births and 53948 matches from a total of 966809 non-Health Start Programme's births statewide (table 1). Approximately 13% of the Health Start Programme population were enrolled in the programme between 9 months and 15 months prior to the birth of the target child, due to previous programme eligibility.

Intervention group

A detailed description of the protocol used to link Health Start Programme enrolment records to birth certificate records can be found elsewhere.²⁹ Briefly, Health Start Programme administrative data were curated to create a longitudinal panel of all Health Start Programme enrollees, which was queried against the Arizona Department of Health Services Vital Records Birth Database using a combination of first name, last name and date of birth. The final linked dataset included an internally generated random unique identifier, measures of linkage quality, the mother's Health Start Programme enrolment date and the child's birth date.

Propensity score matching and synthetic control groups

To limit bias in our estimate of the effectiveness of the Health Start Programme, we followed HomVEE guidelines for observational/retrospective evaluations.

Table 1 Matching results (baseline equivalence) for statewide and rural border county HSP participants

	Statewide					Rural border counties				
	Non-HSP	HSP	Matches	P value	SD	Non-HSP	HSP	Matches	P value	SD
N	966 809	7212	53948			55223	2393	7045		
Maternal age (years)										
Age <20	9.9	17.5	16.9	0.321	0.017	12.4	21.4	21.3	0.944	0.013
Age 20–24	25.3	34.4	34.8	0.564		29.8	37.5	37.5	0.976	
Age 25–30	34.0	28.4	28.6	0.825		33.7	24.8	24.4	0.763	
Age>30	30.8	19.7	19.7	1.000		24.1	16.3	16.7	0.697	
Race/ethnicity										
White	42.4	24.0	23.7	0.653	0.027	28.2	14.7	15.4	0.492	0.054
American Indian	6.0	11.8	12.0	0.700		1.0	0.6	0.6	0.852	
Latina	41.8	59.1	59.7	0.436		65.9	80.9	81.1	0.825	
Other race/ethnicity	9.8	5.1	4.6	0.142		4.9	3.8	2.9	0.077	
Maternal nativity										
Mother born in USA	73.6	68.6	69.2	0.461	0.014	68.4	68.4	69.8	0.288	0.045
Mother born in Mexico	18.7	27.9	27.5	0.602		27.8	30.1	29.1	0.448	
Mother born outside USA	7.6	3.5	3.3	0.550		3.8	1.5	1.1	0.201	
Mother's education										
Less than high school	21.7	32.7	32.9	0.811	0.052	22.2	29.1	29.2	0.936	0.037
High school/GED	28.8	35.7	35.7	0.972		34.9	38.2	39.2	0.458	
Some post-secondary	25.2	23.8	23.5	0.710		26.0	25.4	24.3	0.385	
4-year degree or more	22.5	7.3	7.0	0.420		16.1	7.1	6.9	0.821	
Education missing	1.7	0.3	0.7	0.003		0.7	0.1	0.2	0.489	
Insurance/payer										
Private/commercial insurance	41.1	13.7	13.7	0.904	0.014	36.5	13.5	12.7	0.391	0.025
Medicaid	53.8	82.5	82.8	0.613		56.2	83.1	84.0	0.413	
Other insurance	5.1	3.8	3.5	0.424		7.3	3.4	3.3	0.936	
Married	54.5	37.8	37.4	0.680	0.007	56.3	39.1	37.7	0.342	0.027
Cohabiting	75.6	62.4	62.8	0.570	0.014	71.6	58.5	58.8	0.814	0.009
First birth	36.9	41.6	40.5	0.160	0.023	36.3	51.8	51.0	0.563	0.017
Pre-existing health risk	8.3	11.3	10.8	0.276	0.018	7.4	9.8	8.5	0.120	0.045

Matches: comparison group identified via propensity score. All models control for median income at the zip code level, county of residence and year of birth. Participant subgroups matching models may include additional interactions between controls in order to achieve baseline equivalence. Full tables available upon request.

GED, general education development test; HSP, Health Start Programme; SD, standardised difference.

Specifically, we generated a synthetic control group via propensity score matching, using information provided in birth certificate records. Propensity score matching is used to estimate counterfactual outcomes for a given treatment, that is, what would have happened in the absence of treatment. Matching methods are quasi-experimental, mimicking treatment–control group study designs by identifying a comparison group that is statistically similar to the treatment group. Treatment effectiveness is measured by the difference in outcomes between the two groups, sometimes referred to as the average treatment-on-the-treated effect (ATT).

The ‘propensity score’ is the parameter used to identify the control group and is typically the estimated probability (propensity) that a given individual is part of the treatment group, based on the individual’s characteristics. For this analysis, the propensity scores used to identify each mother’s nearest statistical neighbour were estimated via logistic regression. We allowed for multiple nearest-neighbour matches (ie, ties) to the same Health Start Programme (treatment) observation, based on the estimated propensity score. Thus, the population rates among the matched control groups, for both the covariates used in the matching model

and the outcome measures, are the weighted averages across all matches.

Control group candidates were first limited to mothers residing within the same geographic regions and who gave birth within the same calendar years as the Health Start Programme participants, to implicitly account for prevailing economic and policy conditions based on time and location, as well as any programme parameters that may have shifted as Health Start Programme developed over the study period.³⁵ Next, we selected variables for the matching model itself, which was complicated because Arizona revised the birth certificate form in 2014. As a result, some variables were constructed from component fields (on either side of the revision) in order to generate consistent measures. The final roster of controls were chosen based on their association with both treatment status (ie, Health Start Programme participation) and the outcomes of interest.³⁶ Because Health Start Programme eligibility focuses largely on social and medical risks, we prioritised inclusion of related birth certificate fields. Additionally, in order to achieve an evidence-based rating of 'Moderate', the highest rating for which matched comparison group study designs are eligible,³³ HomVEE requires (at the time of this writing) at least two direct measures of SES. We used maternal education (less than high school, high school, some college, 4 years or more of post-secondary education) and the mother's primary insurance payer (private/commercial, Medicaid and all others) as a proxy for SES.

To mitigate issues with propensity score estimates, we coarsened maternal age (eg, <20 years, 21–25 years, 26–30 years and >30 years) and our indirect SES measure, the median household income of the mother's zip code (decile indicators). All other variables were categorical and include the mother's race/ethnicity, country of birth, marital and cohabiting status, whether or not this was her first birth, the presence of pre-existing health conditions (non-gestational diabetes and/or hypertension, and/or a previous PTB), the county of residence and the year of the delivery. Our variable selection is guided by the HomVEE standards, which require that the covariates used to balance the treatment and control groups be associated with both treatment status and the outcomes of interest,³⁶ and is limited to information available in the data sources.

A critical criterion associated with the HomVEE moderate rating is baseline equivalence, requiring any differences between the treatment and matched-control groups be not statistically significant at the (α) 5% level. Consequently, we verified that our models meet this standard for the Health Start Programme population as a whole, and for each subgroup for whom we separately estimated Health Start Programme's effectiveness. Interaction terms were added to the core set of variables described here to achieve balance and meet the HomVEE standard, as demonstrated in tables 1–3, and more fully in the supplemental tables. In meeting this criterion, the present study can successfully replicate a blocked

experimental design, based on observable characteristics. In addition, the standardised differences (SDs) for all analyses do not exceed 0.2, a typical cut-off for a 'small' effect size.

Descriptive statistics and baseline equivalence

We accessed records for 966809 total births in Arizona from 2006 to 2016 for this study. Among these births, 7212 were to Health Start Programme participants. The first two columns of table 1 show the descriptive statistics for these two populations. Compared with all statewide births, Health Start Programme participants were more likely to be younger than 24 years of age, born in Mexico, have Medicaid insurance, and have pre-existing health risks, and less likely to identify as white or other race/ethnicity, have an education greater than high school degree or GED, and be married or cohabiting. Among Health Start Programme participants, a majority (59%) identify as Latina, over 30% were from a rural border county, 12% identify as American Indian, 17% were teen mothers and 11% had pre-existing health risks. Just 2.6% identify as black and 1.8% as Asian/Pacific Islander, together with those for whom race/ethnicity information is missing, comprise the 'other' category for this variable.

These statistics illustrate both the diversity of the Arizona population and the heterogeneity of Health Start Programme participants. While the propensity score explicitly controls for these (observed) differences in identifying the matches in the non-participant population, we take the additional step of implicitly controlling for unobserved potential differences by evaluating the effectiveness of Health Start Programme separately for selected demographic subgroups. Conditioning on these characteristics further limits the potential for selection bias by narrowing the comparisons.

From the 966809 total births, 53948 (unweighted) were identified as matched controls. Baseline equivalence between Health Start Programme participants and their matches is detailed in tables 1–3 and online supplemental table 1 for all variable comparisons by subgroup. The impact of Health Start Programme is measured by the difference in rates for each outcome measure between the treatment and matched control groups, the ATT.^{37 38} All ATTs were estimated in Stata V.14 using the `teffects` command suite, which, following Abadie and Imbens (2006) takes into account the fact that the propensity scores used to identify the control group are estimated (and not observed) when calculating the SEs and CIs around the difference in rates (ATT).³⁷

RESULTS

Tables 4 and 5 compare the ATT of Health Start Programme participation on LBW, VLBW, ELBW and PTB, across five subgroups. Below, we detail the statistically significant results. All other results were not statistically significant.

Table 2 Matching results (baseline equivalence) for Latina and American Indian HSP participants

	Latina				American Indian					
	Non-HSP	HSP	Matches	P value	SD	Non-HSP	HSP	Matches	P value	SD
N	404 188	4259	32 502			58 358	852	2259		
Maternal age (years)										
Age <20	14.2	18.5	17.8	0.431	0.020	16.0	18.8	17.1	0.377	0.043
Age 20–24	29.5	33.3	33.1	0.890		32.0	35.8	36.4	0.801	
Age 25–30	31.4	27.3	27.6	0.752		29.9	28.6	29.1	0.831	
Age >30	25.0	21.0	21.5	0.578		22.1	16.8	17.4	0.748	
Race/ethnicity										
White	0.0	0.0	0.0		<0.001	0.0	0.0	0.0		<0.001
American Indian	0.0	0.0	0.0			100.0	100.0	100.0	1.000	
Latina	100.0	100.0	100.0	1.000		0.0	0.0	0.0		
Other race/ethnicity	0.0	0.0	0.0			0.0	0.0	0.0		
Maternal nativity										
Mother born in USA	52.1	51.8	52.6	0.474	0.024	99.2	99.9	99.2	0.037	0.101
Mother born in Mexico	44.3	46.9	46.3	0.602		0.2	0.0	0.5	0.034	
Mother born outside USA	3.6	1.3	1.1	0.372		0.5	0.1	0.3	0.465	
Mother's education										
Less than high school	36.6	38.8	39.9	0.304	0.027	27.6	31.0	28.5	0.261	0.075
High school/GED	33.0	35.1	34.5	0.570		38.3	40.3	42.6	0.326	
Some post-secondary	19.6	19.5	18.9	0.441		26.9	26.2	26.8	0.784	
4-year degree or more	10.1	6.2	6.2	0.893		6.3	2.1	1.5	0.365	
Education missing	0.5	0.3	0.3	0.710		0.8	0.5	0.6	0.808	
Insurance/payer										
Private/commercial insurance	23.7	10.1	9.7	0.491	0.016	12.3	5.2	5.3	0.913	0.032
Medicaid	70.3	84.9	85.4	0.483		85.0	93.4	93.7	0.844	
Other insurance	6.0	5.0	4.9	0.842		2.7	1.4	1.1	0.510	
Married	43.0	39.4	39.7	0.740	0.007	21.5	14.3	12.3	0.226	0.059
Cohabiting	71.0	63.6	63.8	0.822	0.005	53.7	50.5	50.8	0.885	0.098
First birth	33.7	40.7	39.8	0.414	0.018	31.7	34.5	34.9	0.879	0.007
Pre-existing health risk	8.0	10.7	10.1	0.357	0.020	14.6	16.4	15.7	0.693	0.019

Matches: comparison group identified via propensity-score. All models control for median income at the zip code level, county of residence and year of birth. Participant subgroups matching models may include additional interactions between controls in order to achieve baseline equivalence. Full tables available upon request. GED, general education development test; HSP, Health Start Programme; SD, standardised difference.

**Table 3** Matching results (baseline equivalence) for teen mothers and mothers with pre-existing health risks

	Teen mothers (Age <20 years)					Mothers with pre-existing health risks*				
	Non-HSP	HSP	Matches	P value	SD	Non-HSP	HSP	Matches	P value	SD
N	95 750	1264	6810			79 912	817	2101		
Maternal age (years)										
Age <20	100.0	100.0	100.0	1.000	<0.001	3.8	5.9	5.1	0.516	0.045
Age 20–24	0.0	0.0	0.0			16.6	25.7	26.1	0.866	
Age 25–30	0.0	0.0	0.0			33.1	34.9	33.8	0.639	
Age>30	0.0	0.0	0.0			46.5	33.5	35.0	0.532	
Race/ethnicity										
White	23.9	22.2	22.9	0.669	0.065	37.3	20.9	20.2	0.714	0.023
American Indian	9.7	12.7	12.1	0.673		10.7	17.1	17.5	0.845	
Latina	59.7	62.2	63.0	0.681		40.7	55.9	56.5	0.803	
Other race/ethnicity	6.6	2.9	2.0	0.123		11.4	6.0	5.8	0.833	
Maternal nativity										
Mother born in USA	79.7	83.2	84.8	0.286	0.043	74.4	66.8	66.2	0.793	0.014
Mother born in Mexico	18.3	16.1	14.6	0.295		17.3	27.7	28.3	0.783	
Mother born outside USA	2.0	0.6	0.6	0.861		8.3	5.5	5.5	1.000	
Mother's education										
Less than high school	55.3	57.6	57.3	0.876	0.043	20.1	29.4	31.4	0.381	0.044
High school/GED	36.4	34.7	35.2	0.802		27.2	34.5	33.5	0.676	
Some post-secondary	7.6	7.3	6.9	0.698		30.2	30.0	29.1	0.705	
4-year degree or more	0.0	0.0	0.1	0.267		21.4	5.8	5.5	0.830	
Education missing	0.6	0.3	0.3	0.989		0.8	0.2	0.3	0.873	
Insurance/payer										
Private/commercial insurance	11.6	6.0	5.4	0.493	0.027	41.5	13.6	13.6	1.000	0.073
Medicaid	84.4	90.6	91.1	0.629		54.8	84.1	82.9	0.506	
Other insurance	4.0	3.4	3.5	0.913		3.7	2.3	3.5	0.143	
Married	12.0	11.7	11.3	0.755	0.012	56.8	43.2	45.2	0.426	0.039
Cohabiting	48.6	41.9	42.3	0.809	0.020	76.6	68.1	70.3	0.335	0.103
First birth	80.6	80.8	82.6	0.237	0.047	28.4	30.7	29.9	0.706	0.019
Pre-existing health risk	3.1	3.8	3.6	0.834	0.008	100.0	100.0	100.0	1.000	

Matches: comparison group identified via propensity-score. All models control for median income at the zip code level, county of residence and year of birth. Participant subgroups matching models may include additional interactions between controls in order to achieve baseline equivalence. Full tables available upon request.

*Pre-existing health risks defined as the presence of pre-existing (non-gestational) diabetes and hypertension. HSP, Health Start Programme; SD, standardised difference.

LBW: we found low rates of LBW for all subgroups except among Latina women, with statistically significant lower rates of LBW among Health Start Programme participants who identified as American Indian and participants with a known pre-existing health risk (table 4). Among American Indian mothers, the LBW rate was 2.30% points lower for Health Start Programme mothers compared with their matched controls. Put differently, the LBW rate was 38% lower for American Indian mothers, a difference of 20 fewer LBW cases, compared with the control group. The LBW rate for Health Start Programme mothers with pre-existing health risks is 25% lower (3.06% points), a

difference of 25 fewer cases of LBW, compared with their matched controls. In Arizona, mothers who identify as American Indian and mothers with pre-existing health risks have higher rates of LBW compared with Arizona statewide rates, 5.96% and 10.41%, respectively, versus 5.52% statewide (table 6).

VLBW and ELBW: in Arizona, all subgroups have higher rates of VLBW and ELBW compared with the statewide rates (table 6). We found a reduction in VLBW and ELBW among all participants, except teens, with significant Health Start Programme's effects among participating mothers who identified as Latina. Among Latina mothers,

Table 4 ATTs of HSP on birth weight

HSP population	VLBW									
	HSP %	Matched %	ATT	95% CI*	P value*	HSP %	Matched %	ATT	95% CI*	P value*
Statewide	5.96	5.92	0.04	-0.65 to 0.73	0.918	0.67	0.83	-0.16	-0.41 to 0.08	0.368
Rural border counties	5.47	6.03	-0.56	-1.80 to 0.69	0.381	0.63	0.84	-0.22	-0.67 to 0.23	0.342
Latina	5.85	5.46	0.39	-0.49 to 1.26	0.385	0.63	0.98	-0.35	-0.69 to -0.01	0.044
American Indian	3.76	6.05	-2.30	-4.07 to -0.53	0.011	0.35	0.64	-0.29	-0.87 to 0.29	0.323
Teen mothers (age: <20 years)	7.04	7.76	-0.72	-2.61 to 1.17	0.455	1.19	1.23	-0.04	-0.88 to 0.79	0.918
Pre-existing health risk†	9.42	12.49	-3.06	-5.82 to -0.30	0.030	1.10	1.63	-0.53	-1.56 to 0.51	0.318

*ATT, CI and p value based on estimated propensity score.

†Pre-existing health risks defined as the presence of pre-existing (non-gestational) diabetes and hypertension, and/or previous PTB.

ATTs, average treatment-on-the-treated effects; HSP, health start program; LBW, low birth weight; PTB, preterm birth; VLBW, very low birth weight.

Table 5 ATTs of HSP participation on birth weight and PTB

HSP population	ELBW						PTB								
	HSP %	Matched %	ATT	95% CI*	P value*	HSP %	Matched %	ATT	95% CI*	P value*	HSP %	Matched %	ATT	95% CI*	P value*
Statewide	0.22	0.45	-0.23	-0.39 to -0.06	0.007	7.38	7.48	-0.28	-0.87 to 0.66	0.793	7.38	7.48	-0.28	-0.87 to 0.66	0.793
Rural border counties	0.21	0.50	-0.29	-0.60 to 0.02	0.068	5.81	6.82	-1.01	-2.31 to 0.28	0.126	5.81	6.82	-1.01	-2.31 to 0.28	0.126
Latina	0.19	0.50	-0.31	-0.52 to -0.10	0.005	7.16	7.66	-0.50	-1.48 to 0.48	0.314	7.16	7.66	-0.50	-1.48 to 0.48	0.314
American Indian	0.23	0.26	-0.03	-0.42 to 0.37	0.894	8.10	7.89	0.20	-1.99 to 2.40	0.855	8.10	7.89	0.20	-1.99 to 2.40	0.855
Teen mothers (age: <20 years)	0.47	0.30	0.18	-0.27 to 0.62	0.440	6.65	9.45	-2.81	-4.71 to -0.91	0.004	6.65	9.45	-2.81	-4.71 to -0.91	0.004
Pre-existing health risk†	0.12	0.30	-0.17	-0.52 to 0.17	0.222	16.03	15.69	0.34	-2.96 to 3.65	0.314	16.03	15.69	0.34	-2.96 to 3.65	0.314

*ATT and p value based on estimated propensity score.

†Pre-existing health risks defined as the presence of pre-existing (non-gestational) diabetes and hypertension.

ATTs, average treatment-on-the-treated effects; ELBW, extremely low birth weight; HSP, Health Start Programme; PTB, preterm birth.

**Table 6** Birth outcomes, Arizona population level rates (2006–2016)

	Statewide	Latina	Border	American Indian	Low education	Prior risk	Age <20 years
Low birth weight	5.52%	5.45%	5.38%	5.96%	6.24%	10.41%	6.83%
Very low birth weight	0.83%	0.85%	0.85%	0.92%	0.93%	1.82%	0.99%
Extremely low birth weight	0.37%	0.40%	0.38%	0.39%	0.41%	0.72%	0.45%
Preterm birth	7.97%	8.12%	6.81%	8.88%	9.05%	16.47%	8.66%

the VLBW and ELBW rates were 0.35% and 0.31% points lower, respectively, for Health Start Programme mothers compared with their matched controls (tables 4 and 5). Health Start Programme participation is associated with a statistically significant lower VLBW rate (36%) and ELBW rate (62%) for Latina mothers, which translates to 15 and 13 fewer women having VLBW and ELBW birth outcomes, respectively, compared with Latina women not enrolled in Health Start Programme.

PTB: we found reductions in the rate of PTB for all groups except American Indian women and women with a pre-existing health risk ($p>0.05$), with statistically significant lower rates of PTB among teen mothers (age: <20 years). In Arizona, almost all subgroups have higher rates of PTB compared with the state rate of 7.97% (table 6). Compared with the matched control group, the PTB rate for Health Start Programme teen mothers is 30% (2.81% points) lower, equaling 35 fewer cases of PTB.

DISCUSSION

Consistent with our hypothesis, women enrolled in the Health Start Programme prior to giving birth over the study period of 2006–2016 had improved birth outcomes, with the most statistically significant effects among American Indian women, women with pre-existing health risks, Latina women and teen mothers. While these effects appear small, the outcomes are rare occurrences but with large consequences. To illustrate, the -0.2% -point effect on the ELBW rate for all Health Start Programme participants is small in absolute terms, but the effect represents substantial economic savings. Gilbert *et al*³⁹ estimates that the early healthcare costs associated with a surviving ELBW infant is approximately \$202 700, compared with \$1100 for a healthy infant. The -0.2% -point effect translates to approximately 16 fewer ELBW cases, translating to an estimated cost savings of \$3.2 million. This is only an example of the monetary costs associated with these infrequent but expensive outcomes, which, when they occur in already disadvantaged populations such as Health Start Programme participants, are likely to be magnified and multiplied over time.

These findings provide important evidence supporting the efficacy of CHW-led home visiting interventions generally, and specifically CHW's ability to address MCH equity in ethnographically and geographically diverse, and socioeconomically disadvantaged populations of mothers and children.⁴⁰ To the best of our knowledge, this is the largest study of a CHW-led MCH home visiting

programme on incidence of LBW, VLBW, ELBW and PTB in the USA to date.

Our evaluation is consistent with an earlier evaluation of the Health Start Programme, which found that Health Start Programme participation was associated with a reduction in the likelihood of LBWs.³¹ Our study relies on a substantially larger sample size, longer observational period and a more sophisticated matching methodology.²⁹ Moreover, the subgroup analyses are a significant new contribution. Although several rigorous studies of prenatal home visitation programmes exist, most use a combination of licensed health professionals, such as nurses and social workers to achieve outcomes.^{41–43} Unique to the Health Start Programme is the utilisation of CHWs as the sole and primary MCH interventionist and home visitor.⁴⁴ The evaluation of the Arizona Health Start Programme is one of very few empirical studies in which CHWs are the primary interventionists that operate outside of a clinical setting and not as a member of a primary care or prenatal care coordination team.^{40–42 45 46} In the USA, and consistent with our findings, CHWs have contributed to the initiation of any, early, and adequate prenatal care, and decreased incidence of PTB and LBW among health disparate populations.^{31 45–52} This study further contributes to CHW effectiveness research in home visitation.

Limitations

Among the smaller limitations, if Health Start mothers are incorrectly linked to state birth certificate data, attenuation bias is possible, resulting in an underestimate (in absolute value) of the magnitude of the resulting coefficients. Linkage quality measures (eg, per cent match on names, etc) suggest this is not a serious concern. The presence of or interaction of Health Start Programme with other maternal health or home visiting programmes may be another source of bias in our estimated treatment effects. Reviewing available evidence, less than 1% of Health Start Programme mothers reported participating in similar CHW or home visiting programmes in the state; therefore, to the extent that the matched mothers similarly participate in another home visiting programme, the potential bias likely is toward finding no effect.

More generally, given the uniqueness of Arizona's population and the consequent demographic heterogeneity among Health Start Programme participants, the present analysis may have limited external validity with respect to other states and populations. A larger limitation (common to matching methods in general) is the

identifying assumption that Health Start Programme participation is fully explained by observable characteristics. Health Start Programme considers social risk factors (eg, domestic violence, lack of family/social support, inconsistent or uncertain employment) as criteria for enrolment. We are unable to observe and account for such factors in this analysis because this information is not available for non-participants (and is rarely collected in large populations in general). At the same time, to the extent that (1) these risks are associated with adverse MCH outcomes, (2) are likely over-represented in the Health Start Programme population and (3) unaccounted for in our control groups, the results presented here likely underestimate a more accurate Health Start Programme's effect.

A related limitation concerns the timing of data collection (at delivery, from the birth certificate) relative to enrolment (prior to delivery). An important identifying assumption of propensity score matching is that the variables used to match are associated with but not affected by treatment status. In the case of controls such as mother's age, race/ethnicity, country of origin and (to a reasonable degree of certainty) pre-existing health conditions, this assumption is easily confirmed. For other controls such as education and insurance, marital and/or cohabitation status, and possible county of residence, it is less certain to be true. For example, a mother may enrol in Health Start Programme without a high school degree and then graduate or complete a GED prior to giving birth. For 86% of Health Start Programme mothers, the window in which they would be able to make this or other changes is 9 months or less, and for 63% it is 6 months or less. Insurance status may be the most likely factor to change. To the extent that 'positive' changes (ie, enrolling in Medicaid) are associated with positive outcomes, such transitions are likely to lead the Health Start Programme impact to be underestimated, because these mothers will be matched to non-participant mothers 'better off' than they were at baseline.

A final limitation associated with the methodology used here is that we are not able to disentangle which programmatic elements of Health Start Programme (eg, different education or home visit formats) and participant engagement (eg, timing and length of enrolment, and degree of engagement) are associated with the effects we report. While this type of mediation pathway analysis is outside the scope of the present study, it is important to note that we are able to show that Health Start Programme is associated with improved outcomes, however, not why or how.

While these limitations are important to keep in mind, the present study has considerable strengths. The birth certificate data enabled us to account for many important factors associated with birth outcomes at the population level. This includes measures of SES, education and insurance status, which are valuable proxies for many social risks. In addition, the volume of enrolment and birth certificate data and the extended time period over which it was collected made possible the calculation of relatively

precise estimates of the effects of the Health Start Programme on low-frequency birth outcomes. Together these results provide valuable high-level evidence of the effectiveness of Health Start Programme.

Implications for public health policy and research

Our limitations represent important next steps in our research agenda, which, in addition to examining health-care utilisation and child development outcomes, will look more closely at the Health Start Programme population itself. Programme administrators collect extensive risk and participation information on Health Start Programme participants. We also plan to use Medicaid claims data (over 80% of Health Start Programme participants are insured by Medicaid) to generate the critical health utilisation and social-risk factors information associated with poor birth outcomes. By comparing programme effects between different participants and implementations, we will be able to address some of these relationships and advance our understanding of the mediating pathways that connect Health Start Programme participation to birth (and other) outcomes.

Several public health entities, including the WHO, the Centers for Disease Control and Prevention, the Institute of Medicine²⁵ and the Community Preventive Services Task Force,⁵³ conclude CHW-centred interventions are effective across disease areas, for example, prevention, screening and management of chronic illnesses such as diabetes, hypertension, cancer and cardiovascular disease,⁵⁴ and across contexts, including community and clinical settings. At the federal level, in 2014, the centres for Medicaid and Medicare-issued guidance to allow states to reimburse for preventive services offered by non-licensed professionals such as CHWs. Such recognition signals opportunity for expansion of CHW home visiting within MCH systems of care. Our study provides much needed evidence to guide policymakers and practitioners on integration and financing of CHW prenatal home visitation. The Arizona Health Start Programme and its 25-year commitment to strengthening CHW MCH home visiting is a healthcare innovation that can improve birth weight and PTB outcomes among ethnoracially, socioeconomically and geographically diverse mothers and infants of Arizona.

CONCLUSIONS

A state health department-operated MCH home visiting intervention that employs CHWs as the primary interventionists may reduce the risk of LBW, VLBW, ELBW and PTB among certain subgroups of women at increased risks for MCH related inequities (ie, American Indian, Latina, women with pre-existing health conditions and teens). Considering the multi-level social and structural determinants of adverse birth outcomes and its impact throughout the life course, results of this evaluation contribute to MCH home visiting research and policy. Specifically, this study advances CHW effectiveness

research as sole and primary home visitation interventionists within MCH.

Author affiliations

¹Center for Health Equity Research, College of Health and Human Services, Northern Arizona University, Flagstaff, Arizona, USA

²Center for Population Science and Discovery, University of Arizona Health Sciences, Tucson, Arizona, USA

³Department of Economics, Brigham Young University, Provo, Utah, USA

⁴Center for Biomedical Informatics and Biostatistics, University of Arizona, Tucson, Arizona, USA

⁵Arizona Department of Health Services, Bureau of Women's and Children's Health, Phoenix, Arizona, USA

Acknowledgements We would like to acknowledge the several offices at the Arizona Department of Health Services (ADHS) that have supported this partnership, including the Bureau of Women and Children's Health, the Health Start Program, the Bureau of Public Health Statistics, the Immunization Program, and Vital Statistics. We are deeply appreciative of their commitment to this study designed to improve the health of women and children of Arizona. We also acknowledge the members of the ADHS Human Subjects Review Board for continued guidance in data sharing and data protection and security. We acknowledge the hundreds of the Health Start CHWs who have worked tirelessly to improve maternal and child health equity among thousands of mothers, infants, and children. We would also like to thank Drs. Lorraine McKelvey and Kimberly McCombs-Thornton for their valuable insights and feedback.

Contributors VP had full access to the identified data in the study and takes responsibility for its integrity. All analyses were conducted on a de-identified (limited) dataset by PW, who is responsible for accuracy of the data analysis. Study concept and design: SS, PW, MB, SR. Acquisition of data: SS, PW, MB, VP, SR. Analysis and interpretation of data: PW, MB, SS, KM, SR, Celaya. Drafting of the manuscript: SS, KM, PW. Critical revision of the manuscript for important intellectual content: SS, KM, PW, MC, SR. Obtained funding: SS. Administrative, technical, or material support: KM, DJJ. Study supervision: SS, PW, SR.

Funding This work is supported through an inter-agency contract from the Arizona Department of Health Services. Funding for the study is provided by the Arizona State Lottery through the Arizona Department of Health Services for the time period of July 1, 2017–June 30, 2022. Health Resources Services Administration (HRSA) Maternal, Infant, Early Childhood Home Visiting (MIECHV) provided 17 months of additional federal funding through the Arizona Department of Health Services during the study period. Role of Sponsor: The funding source, Arizona Department of Health Services, worked closely with the authors to design and conduct the study; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Map disclaimer The depiction of boundaries on the map(s) in this article does not imply the expression of any opinion whatsoever on the part of BMJ (or any member of its group) concerning the legal status of any country, territory, jurisdiction or area or of its authorities. The map(s) are provided without any warranty of any kind, either express or implied.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Approved by the University of Arizona Research Institutional Review Board (protocol number: 1701128802) on 25 January 2017.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement At the conclusion of the study, de-identified data may be available upon request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Samantha Sabo <http://orcid.org/0000-0003-4509-9182>

Kelly McCue <http://orcid.org/0000-0002-8383-0338>

Dulce J Jimenez <http://orcid.org/0000-0003-2962-7467>

REFERENCES

- 1 National Academies of Sciences, Engineering, and Medicine. *The root causes of health inequity: Committee on community-based solutions to promote health equity in the United States*. Washington, DC: National Academies of Sciences, Engineering, and Medicine, 2017.
- 2 Krieger N, Van Wye G, Huynh M, et al. Structural racism, historical Redlining, and risk of preterm birth in New York City, 2013–2017. *Am J Public Health* 2020;110:1046–53.
- 3 Paneth NS. The problem of low birth weight. *Future Child* 1995;5:19–34.
- 4 World Health Organization (WHO). *Global nutrition targets 2025: low birth weight policy brief*. Geneva: World Health Organization, 2014.
- 5 Rimer BK. *Theory at a glance a guide for health promotion practice*. 2nd ed. Bethesda, MD: U.S. Dept. of Health and Human Services, National Institutes of Health, National Cancer Institute, 2005.
- 6 Martinson ML, Reichman NE. Socioeconomic inequalities in low birth weight in the United States, the United Kingdom, Canada, and Australia. *Am J Public Health* 2016;106:748–54.
- 7 Martin JA, Hamilton BE, Osterman MJK, et al. Births: final data for 2017. *National Vital Statistics Reports* 2018;67:1–50.
- 8 Health Resources & Services Administration (HRSA). *Child health USA 2014*. Rockville: Maternal and Child Health Bureau, 2015.
- 9 Pineles B. *Smoking in pregnancy: from effects to solutions [Dissertation]*. Los Angeles: University of Southern California, 2013.
- 10 Thompson JMD, Wall C, Becroft DMO, et al. Maternal dietary patterns in pregnancy and the association with small-for-gestational-age infants. *Br J Nutr* 2010;103:1665–73.
- 11 Almeida J, Bécares L, Erbetta K, et al. Racial/Ethnic inequities in low birth weight and preterm birth: the role of multiple forms of stress. *Matern Child Health J* 2018;22:1154–63.
- 12 Brewin D, Nannini A. Using a life course model to examine racial disparities in low birth weight during adolescence and young adulthood. *J Midwifery Womens Health* 2014;59:417–27.
- 13 Gress-Smith JL, Luecken LJ, Lemery-Chalfant K, et al. Postpartum depression prevalence and impact on infant health, weight, and sleep in low-income and ethnic minority women and infants. *Matern Child Health J* 2012;16:887–93.
- 14 Tomita A, Labys CA, Burns JK. Depressive symptoms prior to pregnancy and infant low birth weight in South Africa. *Matern Child Health J* 2015;19:2179–86.
- 15 Dolan SM. Interpregnancy interval and congenital anomalies. *Am J Obstet Gynecol* 2014;210:498–9.
- 16 Ncube CN, Enquobahrie DA, Albert SM, et al. Association of neighborhood context with offspring risk of preterm birth and low birthweight: a systematic review and meta-analysis of population-based studies. *Soc Sci Med* 2016;153:156–64.
- 17 McEwen BS. Stress, MBS. Stress, adaptation, and disease. allostasis and allostatic load. *Ann N Y Acad Sci* 1998;840:33–44.
- 18 Russell RB, Green NS, Steiner CA, et al. Cost of hospitalization for preterm and low birth weight infants in the United States. *Pediatrics* 2007;120:e1–9.
- 19 Black SE, Devereux PJ, Salvanes KG. From the cradle to the labor market? the effect of birth weight on adult outcomes. *Q J Econ* 2007;122:409–39.
- 20 Gueorguieva R, Morse SB, Roth J. Length of prenatal participation in WIC and risk of delivering a small for gestational age infant: Florida, 1996–2004. *Matern Child Health J* 2009;13:479–88.
- 21 Kogan MD. Social causes of low birth weight. *J R Soc Med* 1995;88:611–5.
- 22 Currie J. Healthy, wealthy, and wise: socioeconomic status, poor health in childhood, and human capital development. *J Econ Lit* 2009;47:87–122.
- 23 Balcazar H, Rosenthal EL, Brownstein JN, et al. Community health workers can be a public health force for change in the United

- States: three actions for a new paradigm. *Am J Public Health* 2011;101:2199–203.
- 24 Kangovi S, Grande D, Trinh-Shevrin C. From rhetoric to reality--community health workers in post-reform U.S. health care. *N Engl J Med* 2015;372:2277–9.
 - 25 Pittman M, Sunderland A, Broderick A. *Bringing community health workers into the mainstream of U.S. health care*. 5. Washington, DC: Institute of Medicine, 2015.
 - 26 Sikka R, Morath JM, Leape L. The quadruple aim: care, health, cost and meaning in work. *BMJ Qual Saf* 2015;24:608–10.
 - 27 Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of the provider. *Ann Fam Med* 2014;12:573–6.
 - 28 Lee H, Crowne SS, Estarziou M. *The effects of home visiting on prenatal health, birth outcomes, and health care use in the first year of life: final implementation and impact findings from the mother and infant home visiting program Evaluation-Strong start*. Washington, DC: Office of Planning, Research, and Evaluation, Administration for Children and Families, US. Department of Health and Human Services, 2019.
 - 29 Sabo S, Butler M, McCue K, et al. Evaluation protocol to assess maternal and child health outcomes using administrative data: a community health worker home visiting programme. *BMJ Open* 2019;9:e031780.
 - 30 Meister JS, Warrick LH, de Zapién JG, et al. Using lay health workers: case study of a community-based prenatal intervention. *J Community Health* 1992;17:37–51.
 - 31 Hussaini SK, Holley P, Ritenour D. Reducing low birth weight infancy: assessing the effectiveness of the health start program in Arizona. *Matern Child Health J* 2011;15:225–33.
 - 32 Rumann S. *Health start core training manual: a self-paced manual for CHW staff*. Arizona: Arizona Department of Health Services, 2014.
 - 33 Home Visiting Evidence of Effectiveness (HomVEE). Review process: producing study ratings. U.S. Department of Health & Human Services, 2020. Available: <https://homvee.acf.hhs.gov/review-process/Producing%20Study%20Ratings>
 - 34 Rosener SE, Barr WB, Frayne DJ, et al. Interconception care for mothers during well-child visits with family physicians: an implicit network study. *Ann Fam Med* 2016;14:350–5.
 - 35 Dehejia RH, Wahba S. Propensity score-matching methods for nonexperimental causal studies. *Rev Econ Stat* 2002;84:151–61.
 - 36 Caliendo M, Kopeinig S. Some practical guidance for the implementation of propensity score matching. *J Econ Surv* 2008;22:31–72.
 - 37 Abadie A, Imbens GW. Large sample properties of matching estimators for average treatment effects. *Econometrica* 2006;74:235–67.
 - 38 Abadie A, Imbens GW. Matching on the estimated propensity score. *Econometrica* 2016;84:781–807.
 - 39 Gilbert WM, Nesbitt TS, Danielsen B. The cost of prematurity: quantification by gestational age and birth weight. *Obstet Gynecol* 2003;102:488–92.
 - 40 Pan Z, Veazie P, Sandler M, et al. Perinatal health outcomes following a community health Worker-Supported Home-Visiting program in Rochester, New York, 2015–2018. *Am J Public Health* 2020;110:1031–3.
 - 41 Roman L, Raffo JE, Zhu Q, et al. A statewide Medicaid enhanced prenatal care program: impact on birth outcomes. *JAMA Pediatr* 2014;168:220–7.
 - 42 Meghea CI, Raffo JE, Zhu Q, et al. Medicaid home visitation and maternal and infant healthcare utilization. *Am J Prev Med* 2013;45:441–7.
 - 43 Roman LA, Raffo JE, Meghea CI. Maternal perceptions of help from home visits by nurse-community health worker teams. *Am J Public Health* 2012;102:643–5.
 - 44 Bradley PJ, Martin J. The impact of home visits on enrollment patterns in pregnancy-related services among low-income women. *Public Health Nurs* 1994;11:392–8.
 - 45 Williams CM, Cprek S, Asaolu I, et al. Kentucky health access nurturing development services home visiting program improves maternal and child health. *Matern Child Health J* 2017;21:1166–74.
 - 46 Redding S, Conrey E, Porter K, et al. Pathways community care coordination in low birth weight prevention. *Matern Child Health J* 2015;19:643–50.
 - 47 Poland ML, Giblin PT, Waller JB, et al. Effects of a home visiting program on prenatal care and birthweight: a case comparison study. *J Community Health* 1992;17:221–9.
 - 48 Coughlin RL, Kushman EK, Copeland GE, et al. Pregnancy and birth outcome improvements for American Indians in the healthy start project of the Inter-Tribal Council of Michigan, 1998–2008. *Matern Child Health J* 2013;17:1005–15.
 - 49 Brown KK, Johnson C, Spainhower M, et al. Is timing of enrollment associated with birth outcomes? findings from a healthy start program in Kansas. *Matern Child Health J* 2017;21:25–31.
 - 50 DeAngelis KR, Doré KF, Dean D, et al. Strengthening the healthy start workforce: a mixed-methods study to understand the roles of community health workers in healthy start and inform the development of a standardized training program. *Matern Child Health J* 2017;21:65–74.
 - 51 Bouye KH. *The resource mothers program: how community health workers can reduce low-birth weight among African-American clients in WIC programs*. Columbus: The Ohio State University, 2005.
 - 52 Lee E, Mitchell-Herzfeld SD, Lowenfels AA, et al. Reducing low birth weight through home visitation: a randomized controlled trial. *Am J Prev Med* 2009;36:154–60.
 - 53 Community Preventive Services Taskforce. Community guide: community health workers, 2017. Available: <https://www.thecommunityguide.org/content/community-health-workers>
 - 54 Kim K, Choi JS, Choi E, et al. Effects of community-based health worker interventions to improve chronic disease management and care among vulnerable populations: a systematic review. *Am J Public Health* 2016;106:e3–28.

Statewide Results: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value difference‡ SD
	Non-HSP	ORs	95% CI	HSP	MCG	
N	966,809			7,212	53,674	
<u>Birth outcomes</u>						
Low birth weight	5.5%			6.0%	5.9%	0.918
Very low birth weight	0.8%			0.7%	0.8%	0.263
Extremely low birth weight	0.4%			0.2%	0.4%	0.018
Preterm birth	8.0%			7.4%	7.5%	0.814
<u>Covariates/Matching controls</u>						
<i>Mother's age</i>						
Age<20	9.9%	1.35***	(1.24, 1.47)	17.5%	17.2%	0.553 0.011
Age 20-24	25.3%	1.19***	(1.12, 1.27)	34.4%	34.7%	0.649
Age 25-30†	34.0%			28.4%	28.5%	0.868
Age>30	30.8%	0.91**	(0.85, 0.98)	19.7%	19.6%	0.867
<i>Mother's Race/Ethnicity & Nativity</i>						
White	42.4%	0.93	(0.82, 1.06)	24.0%	24.0%	1.000 0.017
American Indian	6.0%	1.43***	(1.23, 1.66)	11.8%	12.0%	0.662
Hispanic	41.8%	1.29***	(1.13, 1.46)	59.1%	59.2%	0.892
Other race/ethnicity†	9.8%			5.1%	4.8%	0.338
Born in US†	73.6%			68.6%	69.5%	0.280 0.022
Born in Mexico	18.7%	1.68***	(1.57, 1.80)	27.9%	27.4%	0.479
Born outside US	7.6%	1.18*	(1.00, 1.39)	3.5%	3.2%	0.307
Hispanic & born outside US	1.5%	0.76*	(0.55, 1.03)	0.8%	0.7%	0.848
<i>Mother's education</i>						
Less than high school†	21.7%			32.7%	32.7%	0.985 0.027
High school/GED	28.8%	0.84***	(0.79, 0.90)	35.7%	35.8%	0.862
Some post-secondary education	25.2%	0.81***	(0.76, 0.87)	23.8%	23.6%	0.845
4-year degree or more	22.5%	0.59***	(0.53, 0.66)	7.3%	7.4%	0.949
Education missing	1.7%	0.41***	(0.27, 0.62)	0.3%	0.2%	0.104
<i>Method of payment</i>						
Private/commercial insurance	41.1%	0.92	(0.80, 1.07)	13.7%	13.8%	0.866 0.014
AHCCCS (AZ Medicaid)	53.8%	2.34***	(2.05, 2.68)	82.5%	82.7%	0.792
Other insurance†	5.1%			3.8%	3.5%	0.398
<i>Marital cohabitation status</i>						
Married	54.5%	1.06	(0.92, 1.23)	37.8%	37.2%	0.513
Married & cohabiting	52.0%	0.93	(0.80, 1.09)	34.7%	34.5%	0.820
Mother & father cohabiting	75.6%	0.99	(0.93, 1.05)	62.4%	62.3%	0.973
<i>Health factors</i>						
First birth	36.9%	1.27***	(1.20, 1.34)	41.6%	40.8%	0.327
Pre-existing health risk	8.3%	1.11**	(1.02, 1.19)	11.3%	10.8%	0.326

Statewide Results: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
<i>Mother's county of residence</i>							
Maricopa†	62.5%			26.4%	26.8%	0.572	0.031
Cochise, Santa Cruz	2.4%	15.75***	(14.54, 17.07)	16.8%	17.1%	0.625	
Gila	0.7%	6.56***	(5.55, 7.76)	2.4%	2.3%	0.784	
Mohave	2.2%	4.50***	(3.95, 5.12)	4.2%	4.1%	0.676	
Pima	13.4%	1.60***	(1.46, 1.76)	9.0%	9.1%	0.839	
Coconino, Navajo	3.7%	6.73***	(6.09, 7.43)	11.9%	11.2%	0.160	
Yavapai	2.1%	5.57***	(4.94, 6.27)	5.2%	5.2%	0.970	
Yuma	3.3%	7.99***	(7.38, 8.65)	16.4%	16.7%	0.622	
Apache, Graham, Greenlee, Pinal, La Paz	7.4%	1.85***	(1.66, 2.07)	5.9%	5.6%	0.453	
County missing	2.2%	2.06***	(1.46, 2.89)	1.8%	2.0%	0.626	
<i>Median income of mother's zip code</i>							
Bottom income decile	2.9%	0.95	(0.74, 1.23)	2.4%	2.0%	0.113	0.043
Income decile 2	6.7%	1.93***	(1.56, 2.39)	15.3%	15.6%	0.565	
Income decile 3	10.4%	2.33***	(1.89, 2.88)	15.5%	15.8%	0.614	
Income decile 4	6.5%	2.65***	(2.14, 3.29)	10.4%	10.8%	0.372	
Income decile 5	8.4%	2.28***	(1.85, 2.82)	14.6%	14.7%	0.888	
Income decile 6	11.7%	1.73***	(1.40, 2.14)	8.6%	8.5%	0.905	
Income decile 7	11.9%	2.22***	(1.81, 2.74)	13.7%	12.8%	0.111	
Income decile 8	10.7%	3.15***	(2.55, 3.88)	12.7%	12.9%	0.746	
Income decile 9	12.1%	1.26*	(0.99, 1.60)	2.8%	2.8%	0.960	
Top income decile†	10.4%			1.4%	1.3%	0.472	
Zip code missing	8.3%	1.43**	(1.07, 1.92)	2.6%	2.7%	0.639	
<i>Birth certificate year</i>							
2006-07†	20.8%			0.6%	0.6%	0.915	0.022
2008	10.0%	3.45***	(2.38, 5.00)	1.0%	1.0%	0.867	
2009	9.3%	9.93***	(7.18, 13.74)	2.7%	2.8%	0.878	
2010	8.7%	30.03***	(22.14, 40.74)	7.4%	7.4%	1.000	
2011	8.5%	57.58***	(42.67, 77.70)	13.5%	13.9%	0.513	
2012	8.6%	63.19***	(46.84, 85.24)	14.9%	14.9%	0.925	
2013	8.5%	70.92***	(52.59, 95.64)	15.9%	15.8%	0.820	
2014	8.7%	68.23***	(50.60, 92.01)	15.8%	16.1%	0.649	
2015	8.5%	67.63***	(50.13, 91.24)	14.9%	14.3%	0.278	
2016	8.5%	59.81***	(44.30, 80.77)	13.1%	13.3%	0.749	
<i>Other balancing factors</i>							
County, median income missing	0.02	1.36	(0.86, 2.14)	0.01	0.02	0.268	
American Indian*college	0.00	0.83	(0.51, 1.34)	0.00	0.00	0.479	

*** p<.01; ** p<.05; * p<.01

‡ p-value for difference in outcome rates between HSP and matches accounts for estimated propensity score.

† Reference category/categories.

HSP: Health Start Participants; MCG: Matched Control Group; SD: Standardized Differences (variables w/more than two categories).

Latina Mothers: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)		Match Results		p-value difference‡ SD	
	Non-HSP	ORs	95% CI	HSP	MCG	
N	404,188			4,259	32,502	
<u>Birth outcomes</u>						
Low birth weight	5.5%			5.8%	5.5%	0.385
Very low birth weight	0.9%			0.6%	1.0%	0.044
Extremely low birth weight	0.4%			0.2%	0.5%	0.005
Preterm birth	8.1%			7.2%	7.7%	0.314
<u>Covariates/Matching controls</u>						
<i>Mother's age</i>						
Age<20	14.2%	1.23***	(1.10, 1.38)	18.5%	17.8%	0.431
Age 20-24	29.5%	1.13***	(1.04, 1.23)	33.3%	33.1%	0.890
Age 25-30†	31.4%			27.3%	27.6%	0.752
Age>30	25.0%	0.99	(0.90, 1.08)	21.0%	21.5%	0.578
<i>Mother's Nativity</i>						
Mother born in US†	52.1%			51.8%	52.6%	0.474
Mother born in Mexico	44.3%	1.66***	(1.55, 1.78)	46.9%	46.3%	0.602
Mother born outside US	3.6%	0.88	(0.67, 1.15)	1.3%	1.1%	0.372
<i>Mother's education</i>						
Less than high school†	36.6%			38.8%	39.9%	0.304
High school/GED	33.0%	0.83***	(0.77, 0.89)	35.1%	34.5%	0.570
Some post-secondary education	19.6%	0.79***	(0.72, 0.87)	19.5%	18.9%	0.441
4-year degree or more	10.1%	0.67***	(0.57, 0.77)	6.2%	6.2%	0.893
Education missing	0.5%			0.3%	0.3%	0.710
<i>Method of payment</i>						
Private/commercial insurance	23.7%	0.91	(0.75, 1.09)	10.1%	9.7%	0.491
AHCCCS (AZ Medicaid)	70.3%	2.19***	(1.86, 2.58)	84.9%	85.4%	0.483
Other insurance†	6.0%			5.0%	4.9%	0.842
<i>Marital cohabitation status</i>						
Married	43.0%	1.16	(0.96, 1.40)	39.4%	39.7%	0.740
Mother & father cohabiting	71.0%	1.05	(0.97, 1.14)	63.6%	63.8%	0.822
Married & cohabiting	41.1%	0.90	(0.74, 1.11)	36.2%	36.6%	0.719
<i>Health factors</i>						
First birth	33.7%	1.39***	(1.29, 1.50)	40.7%	39.8%	0.414
Pre-existing health risk	8.0%	1.09	(0.98, 1.21)	10.7%	10.1%	0.357

Latina Mothers: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
<i>Mother's county of residence</i>							
Cochise	1.7%	3.18***	(2.80, 3.61)	11.9%	10.8%	0.109	0.041
Maricopa	63.9%	0.16***	(0.15, 0.18)	24.7%	25.2%	0.565	
Pima	15.2%	0.26***	(0.23, 0.29)	8.8%	8.4%	0.536	
Santa Cruz	1.5%	2.95***	(2.56, 3.39)	9.8%	9.9%	0.884	
Yuma	5.8%	1.43***	(1.29, 1.58)	23.7%	24.6%	0.349	
Other county†	8.7%			18.7%	18.7%	0.978	
County missing	3.1%	0.44***	(0.28, 0.70)	2.4%	2.4%	1.000	
<i>Median income of mother's zip code</i>							
Bottom income decile†	3.1%			2.2%	2.2%	0.941	0.045
Income decile 2	10.4%	1.34***	(1.07, 1.67)	21.3%	20.6%	0.440	
Income decile 3	15.8%	1.52***	(1.22, 1.90)	18.3%	18.6%	0.675	
Income decile 4	8.8%	1.55***	(1.23, 1.95)	10.6%	11.1%	0.464	
Income decile 5	9.8%	1.49***	(1.19, 1.86)	15.4%	15.9%	0.474	
Income decile 6	12.5%	0.96	(0.76, 1.22)	8.1%	7.9%	0.842	
Income decile 7	11.4%	1.15	(0.92, 1.45)	11.6%	10.9%	0.337	
Income decile 8	9.6%	1.50***	(1.18, 1.90)	7.6%	7.7%	0.807	
Income decile 9	7.5%	0.53***	(0.38, 0.74)	1.5%	1.6%	0.538	
Top income decile	3.7%	0.82	(0.56, 1.21)	0.9%	0.7%	0.217	
Zip code missing	7.5%	0.47***	(0.31, 0.71)	2.7%	2.7%	1.000	
<i>Birth certificate year</i>							
2006-07†	22.2%			0.8%	0.7%	0.799	0.026
2008	10.4%	3.27***	(2.09, 5.11)	1.1%	1.0%	0.675	
2009	9.4%	8.90***	(6.02, 13.16)	2.8%	2.7%	0.739	
2010	8.4%	29.66***	(20.62, 42.66)	8.0%	7.6%	0.519	
2011	8.0%	51.27***	(35.86, 73.30)	13.5%	13.5%	0.949	
2012	8.1%	54.72***	(38.29, 78.20)	14.9%	14.9%	0.951	
2013	8.1%	57.82***	(40.46, 82.64)	15.1%	15.1%	0.976	
2014	8.5%	56.42***	(39.49, 80.62)	15.2%	15.9%	0.420	
2015	8.5%	56.24***	(39.34, 80.40)	14.9%	14.7%	0.807	
2016	8.5%	52.89***	(36.96, 75.67)	13.8%	14.0%	0.778	
County, median income missing	2.9%	2.19**	(1.18, 4.09)	1.9%	2.1%	0.489	

*** p<.01; ** p<.05; * p<.01

‡ p-value for difference in outcome rates between HSP and matches accounts for estimated propensity score.

† Reference category/categories.

HSP: Health Start Participants; MCG: Matched Control Group; SD: Standardized Differences (variables w/more than two categories).

Rural Border County Residents: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
N	55,223			2,393	7,045		
<u>Birth outcomes</u>							
Low birth weight	5.4%			5.5%	6.0%	0.381	
Very low birth weight	0.8%			0.6%	0.8%	0.342	
Extremely low birth weight	0.4%			0.2%	0.5%	0.068	
Preterm birth	6.8%			5.8%	6.8%	0.126	
<u>Covariates/Matching controls</u>							
<i>Mother's age</i>							
Age<20	12.4%	1.36***	(1.16, 1.58)	21.4%	21.3%	0.944	0.013
Age 20-24	29.8%	1.23***	(1.10, 1.38)	37.5%	37.5%	0.976	
Age 25-30†	33.7%			24.8%	24.4%	0.763	
Age>30	24.1%	1.05	(0.92, 1.20)	16.3%	16.7%	0.697	
<i>Mother's Race/Ethnicity & Nativity</i>							
White	28.2%	0.63***	(0.49, 0.81)	14.7%	15.4%	0.492	0.054
American Indian	1.0%	0.55**	(0.31, 0.98)	0.6%	0.6%	0.852	
Hispanic	65.9%	0.96	(0.75, 1.24)	80.9%	81.1%	0.825	
Other race/ethnicity†	4.9%			3.8%	2.9%	0.077	
Mother born in US†	68.4%			68.4%	69.8%	0.288	0.045
Mother born in Mexico	27.8%	1.19***	(1.07, 1.32)	30.1%	29.1%	0.448	
Mother born outside US	3.8%	0.65**	(0.43, 0.99)	1.5%	1.1%	0.201	
Hispanic & born outside US	0.9%	0.84	(0.38, 1.84)	0.4%	0.3%	0.438	
<i>Mother's education</i>							
Less than high school†	22.2%			29.1%	29.2%	0.936	0.037
High school/GED	34.9%	0.98	(0.88, 1.09)	38.2%	39.2%	0.458	
Some post-secondary education	26.0%	0.97	(0.85, 1.09)	25.4%	24.3%	0.385	
4-year degree or more	16.1%	0.71***	(0.59, 0.86)	7.1%	6.9%	0.821	
Education missing†	0.7%			0.1%	0.2%	0.489	
<i>Method of payment</i>							
Private/commercial insurance	36.5%	1.20	(0.93, 1.55)	13.5%	12.7%	0.391	0.025
AHCCS (AZ Medicaid)	56.2%	3.52***	(2.80, 4.43)	83.1%	84.0%	0.413	
Other insurance†	7.3%			3.4%	3.3%	0.936	
<i>Marital cohabitation status</i>							
Married	56.3%	0.98	(0.77, 1.25)	39.1%	37.7%	0.342	
Mother & father cohabiting	71.6%	1.00	(0.89, 1.12)	58.5%	58.8%	0.814	
Married & cohabiting	53.4%	1.00	(0.76, 1.30)	35.7%	35.0%	0.586	
<i>Health factors</i>							
First birth	36.3%	1.84***	(1.66, 2.03)	51.8%	51.0%	0.563	
Pre-existing health risk	7.4%	1.01	(0.87, 1.17)	9.8%	8.5%	0.120	

Rural Border County Residents: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value difference‡ SD	
	Non-HSP	ORs	95% CI	HSP	MCG		
<i>Mother's county of residence</i>							
Cochise	30.4%	1.51***	(1.35, 1.68)	32.8%	32.4%	0.781	0.008
Santa Cruz, Yuma†	69.6%			67.2%	67.6%	0.781	
<i>Median income of mother's zip code</i>							
Bottom income decile	0.2%	1.07	(0.57, 2.00)	0.6%	0.4%	0.316	0.058
Income decile 2	20.2%	1.41**	(1.01, 1.96)	29.1%	29.7%	0.657	
Income decile 3	10.3%	1.19	(0.84, 1.67)	15.6%	15.7%	0.937	
Income decile 4	7.9%	0.90	(0.63, 1.28)	9.5%	10.4%	0.288	
Income decile 5	23.4%	1.08	(0.77, 1.51)	16.5%	16.5%	0.938	
Income decile 6	4.6%	1.25	(0.85, 1.86)	3.2%	2.8%	0.448	
Income decile 7	25.4%	1.09	(0.78, 1.52)	17.7%	17.4%	0.790	
Income decile 8	4.0%	1.18	(0.82, 1.68)	5.6%	5.2%	0.524	
Top income quintile†	3.3%			1.8%	1.5%	0.367	
Zip code missing	0.7%	0.82	(0.37, 1.80)	0.3%	0.3%	0.796	
<i>Birth certificate year</i>							
2006-08†	30.3%			1.0%	1.1%	0.776	0.039
2009	9.8%	8.24***	(5.14, 13.19)	2.7%	2.5%	0.648	
2010	9.4%	31.26***	(20.48, 47.73)	9.3%	9.5%	0.804	
2011	8.9%	53.78***	(35.48, 81.52)	14.6%	15.2%	0.598	
2012	8.8%	61.63***	(40.58, 93.59)	16.0%	16.5%	0.639	
2013	8.2%	64.13***	(42.29, 97.24)	15.8%	15.6%	0.905	
2014	8.3%	55.64***	(36.60, 84.59)	14.0%	13.0%	0.352	
2015	8.2%	54.71***	(35.92, 83.33)	13.4%	13.8%	0.704	
2016	8.2%	53.77***	(35.31, 81.88)	13.2%	12.8%	0.637	

*** p<.01; ** p<.05; * p<.01

‡ p-value for difference in outcome rates between HSP and matches accounts for estimated propensity score.

† Reference category/categories.

HSP: Health Start Participants; MCG: Matched Control Group; SD: Standardized Differences (variables w/more than two categories).

American Indians: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
N	58,358			852	2,259		
<u>Birth outcomes</u>							
Low birth weight	6.0%			3.8%	6.1%	0.011	
Very low birth weight	0.9%			0.4%	0.6%	0.323	
Extremely low birth weight	0.4%			0.2%	0.3%	0.894	
Preterm birth	8.9%			8.1%	7.9%	0.855	
<u>Covariates/Matching controls</u>							
<i>Mother's age</i>							
Age<20	16.0%	1.37**	(1.07, 1.76)	18.8%	17.1%	0.377	0.043
Age 20-24	32.0%	1.17*	(0.98, 1.40)	35.8%	36.4%	0.801	
Age 25-30†	29.9%			28.6%	29.1%	0.831	
Age>30	22.1%	0.94	(0.76, 1.16)	16.8%	17.4%	0.748	
<i>Mother's education</i>							
Less than high school†	27.6%			31.0%	28.5%	0.261	0.075
High school/GED	38.3%	0.92	(0.77, 1.09)	40.3%	42.6%	0.326	
Some post-secondary education	26.9%	0.75***	(0.62, 0.92)	26.2%	26.8%	0.784	
4-year degree or more	6.3%	0.37***	(0.22, 0.62)	2.1%	1.5%	0.365	
Education missing†	0.8%			0.5%	0.6%	0.808	
<i>Method of payment</i>							
Private/commercial insurance	12.3%	0.49**	(0.25, 0.95)	5.2%	5.3%	0.913	0.032
AHCCCS (AZ Medicaid)	85.0%	1.34	(0.75, 2.42)	93.4%	93.7%	0.844	
Other insurance†	2.7%			1.4%	1.1%	0.510	
<i>Marital cohabitation status</i>							
Married	21.5%	0.73	(0.42, 1.26)	14.3%	12.3%	0.226	
Mother & father cohabiting	53.7%	1.02	(0.87, 1.19)	50.5%	50.8%	0.885	
Married & cohabiting	19.5%	1.15	(0.64, 2.08)	12.7%	11.6%	0.505	
<i>Health factors</i>							
First birth	31.7%	0.97	(0.81, 1.15)	34.5%	34.9%	0.879	
Pre-existing health risk	14.6%	0.92	(0.75, 1.12)	16.4%	15.7%	0.693	

American Indians: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value difference‡ SD	
	Non-HSP	ORs	95% CI	HSP	MCG		
<i>Mother's county of residence</i>							
Maricopa†	27.2%			56.5%	58.2%	0.463	0.092
Apache	15.4%	0.20***	(0.15, 0.28)	5.3%	3.6%	0.101	
Coconino	12.2%	0.65***	(0.54, 0.79)	23.5%	23.5%	1.000	
Navajo	17.4%	0.22***	(0.16, 0.29)	7.0%	6.7%	0.774	
Other county	24.8%	0.12***	(0.09, 0.16)	6.6%	7.2%	0.632	
County missing	3.0%	0.21***	(0.11, 0.42)	1.2%	0.8%	0.465	
<i>Median income of mother's zip code</i>							
Bottom income decile	20.8%	0.91	(0.58, 1.42)	5.4%	4.0%	0.170	0.135
Income decile 2	15.8%	1.58**	(1.08, 2.33)	11.3%	11.3%	1.000	
Income decile 3	12.1%	1.69***	(1.17, 2.43)	13.8%	15.3%	0.410	
Income decile 4	7.4%	1.32	(0.87, 1.99)	6.2%	5.9%	0.761	
Income decile 5	6.7%	1.79***	(1.20, 2.67)	7.5%	9.2%	0.220	
Income decile 6	9.5%	1.46*	(0.99, 2.14)	9.4%	10.3%	0.516	
Income decile 7	6.1%	2.37***	(1.63, 3.46)	10.9%	8.3%	0.071	
Income decile 8	7.8%	5.07***	(3.59, 7.15)	28.2%	28.4%	0.914	
Income decile 9†	4.7%			4.3%	4.0%	0.732	
Top income decile†	1.4%			0.6%	0.8%	0.593	
Zip code missing	7.7%	1.52	(0.85, 2.71)	2.3%	2.6%	0.755	
<i>Birth certificate year</i>							
2006†	10.3%			0.2%	0.2%	1.000	0.114
2007	10.6%	0.99	(0.14, 7.01)	0.2%	0.1%	0.564	
2008	10.3%	6.22**	(1.39, 27.84)	1.4%	2.1%	0.269	
2009	10.0%	14.60***	(3.47, 61.36)	3.3%	4.2%	0.308	
2010	9.3%	32.45***	(7.91, 133.08)	6.8%	7.6%	0.513	
2011	9.3%	78.07***	(19.29, 315.98)	15.3%	15.7%	0.789	
2012	8.9%	75.95***	(18.75, 307.64)	14.9%	14.7%	0.892	
2013	8.7%	96.64***	(23.88, 391.04)	16.7%	15.6%	0.554	
2014	7.8%	110.30***	(27.27, 446.12)	17.6%	17.5%	0.949	
2015	7.3%	98.90***	(24.40, 400.83)	14.7%	15.3%	0.734	
2016	7.5%	56.31***	(13.79, 229.86)	8.9%	6.9%	0.127	

*** p<.01; ** p<.05; * p<.01

‡ p-value for difference in outcome rates between HSP and matches accounts for estimated propensity score.

† Reference category/categories.

HSP: Health Start Participants; MCG: Matched Control Group; SD: Standardized Differences (variables w/more than two categories).

Education Less than high school: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value difference‡ SD
	Non-HSP	ORs	95% CI	HSP	MCG	
N	209,365			2,361	18,683	
<u>Birth outcomes</u>						
Low birth weight	6.2%			6.9%	7.2%	0.666
Very low birth weight	0.9%			0.8%	0.7%	0.582
Extremely low birth weight	0.4%			0.2%	0.2%	0.937
Preterm birth	9.0%			8.2%	9.0%	0.248
<u>Covariates/Matching controls</u>						
<i>Mother's age</i>						
Age<20	25.3%	1.50***	(1.29, 1.74)	30.8%	30.1%	0.569 0.023
Age 20-24	28.3%	1.12*	(0.99, 1.27)	27.1%	26.9%	0.819
Age 25-30†	25.4%			21.1%	21.4%	0.803
Age>30	21.0%	1.09	(0.95, 1.24)	20.9%	21.7%	0.522
<i>Mother's Race/Ethnicity & Nativity</i>						
White	16.3%	0.57***	(0.44, 0.75)	14.6%	14.4%	0.869 0.010
American Indian	7.7%	1.37**	(1.03, 1.81)	11.2%	11.1%	0.890
Hispanic	70.7%	1.00	(0.79, 1.26)	70.1%	70.2%	0.924
Other race/ethnicity†	5.3%			4.2%	4.4%	0.774
Mother born in US†	53.5%			58.6%	58.8%	0.883 0.024
Mother born in Mexico	41.5%	1.60***	(1.43, 1.79)	38.5%	37.9%	0.675
Mother born outside US	5.0%	1.06	(0.80, 1.40)	2.9%	3.3%	0.451
<i>Method of payment</i>						
Private/commercial insurance	8.1%	0.83	(0.60, 1.15)	3.3%	3.2%	0.935 0.039
AHCCCS (AZ Medicaid)	87.4%	2.00***	(1.57, 2.53)	93.3%	94.0%	0.311
Other insurance†	4.5%			3.5%	2.8%	0.182
<i>Marital cohabitation status</i>						
Married	29.7%	0.88	(0.64, 1.19)	28.8%	27.4%	0.300
Mother & father cohabiting	62.7%	0.95	(0.86, 1.06)	55.7%	54.7%	0.501
Married & cohabiting	28.4%	1.30	(0.93, 1.80)	26.9%	25.8%	0.428
<i>Health factors</i>						
First birth	30.7%	1.08	(0.96, 1.21)	35.2%	35.1%	0.951
Pre-existing health risk	7.7%	0.99	(0.86, 1.14)	10.2%	10.7%	0.536

Education Less than high school: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value difference‡ SD
	Non-HSP	ORs	95% CI	HSP	MCG	
<i>Mother's county of residence</i>						
Maricopa†	65.2%			31.2%	32.2%	0.453
Cochise	1.2%	20.38***	(17.24, 24.08)	10.0%	10.2%	0.735
Mohave	2.5%	4.26***	(3.34, 5.44)	3.5%	3.6%	0.937
Apache, Coconino, Navajo	4.3%	4.52***	(3.69, 5.53)	13.6%	11.7%	0.054
Pima	11.7%	1.86***	(1.59, 2.17)	9.6%	9.2%	0.690
Santa Cruz	0.8%	19.83***	(16.22, 24.24)	6.7%	6.6%	0.861
Yavapai	2.1%	9.74***	(8.13, 11.67)	8.0%	8.2%	0.831
Yuma	3.9%	6.07***	(5.24, 7.02)	12.8%	13.7%	0.367
Other county	6.5%	1.08	(0.84, 1.37)	3.3%	3.3%	0.935
County missing	1.8%	1.91***	(1.23, 2.95)	1.4%	1.2%	0.699
<i>Median income of mother's zip code</i>						
Bottom income decile†	5.1%			3.1%	2.5%	0.253 0.074
Income decile 2	11.6%	1.61***	(1.24, 2.09)	17.2%	17.3%	0.908
Income decile 3	17.5%	2.19***	(1.70, 2.84)	17.8%	18.6%	0.474
Income decile 4	10.8%	2.61***	(1.98, 3.43)	11.0%	12.0%	0.254
Income decile 5	10.3%	2.12***	(1.62, 2.77)	14.5%	14.7%	0.869
Income decile 6	12.4%	1.83***	(1.39, 2.42)	9.2%	8.3%	0.304
Income decile 7	10.1%	2.11***	(1.61, 2.77)	11.1%	10.0%	0.217
Income decile 8	9.1%	3.71***	(2.84, 4.85)	12.1%	12.3%	0.790
Income decile 9	5.2%	1.39	(0.93, 2.08)	1.6%	1.9%	0.373
Top income decile	1.6%	1.02	(0.52, 2.00)	0.4%	0.3%	0.637
Zip code missing	6.4%	1.66**	(1.09, 2.52)	2.1%	1.9%	0.677
<i>Birth certificate year</i>						
2006†	13.2%			0.5%	0.5%	0.834 0.060
2007	13.0%	1.13	(0.52, 2.48)	0.6%	0.4%	0.393
2008	11.7%	2.88***	(1.47, 5.62)	1.3%	1.3%	0.897
2009	10.0%	7.89***	(4.28, 14.55)	3.0%	3.0%	0.932
2010	8.5%	25.70***	(14.32, 46.11)	8.0%	8.0%	0.957
2011	7.5%	52.69***	(29.57, 93.87)	14.1%	15.2%	0.323
2012	7.4%	56.48***	(31.70, 100.64)	14.7%	14.1%	0.534
2013	6.8%	61.57***	(34.54, 109.74)	14.6%	14.2%	0.709
2014	7.7%	57.23***	(32.13, 101.95)	15.1%	16.1%	0.336
2015	7.3%	62.01***	(34.80, 110.51)	14.9%	13.7%	0.244
2016	6.9%	57.20***	(32.05, 102.08)	13.1%	13.6%	0.669
<i>Other balancing factors</i>						
White*Apache, Coconino, Navajo	0.6%	5.41***	(3.95, 7.39)	4.4%	3.9%	0.344

*** p<.01; ** p<.05; * p<.01

‡ p-value for difference in outcome rates between HSP and matches accounts for estimated propensity score.

† Reference category/categories.

HSP: Health Start Participants; MCG: Matched Control Group; SD: Standardized Differences (variables w/more than two categories).

Pre-Existing Health Risk: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value difference‡ SD	
	Non-HSP	ORs	95% CI	HSP	MCG		
N	79,912			817	2,101		
<u>Birth outcomes</u>							
Low birth weight	10.4%			9.4%	12.5%		0.030
Very low birth weight	1.8%			1.1%	1.6%		0.318
Extremely low birth weight	0.7%			0.1%	0.3%		0.330
Preterm birth	16.5%			16.0%	15.7%		0.839
<u>Covariates/Matching controls</u>							
<i>Mother's age</i>							
Age<20	3.8%	0.93	(0.66, 1.32)	5.9%	5.1%		0.516
Age 20-24	16.6%	1.11	(0.92, 1.35)	25.7%	26.1%		0.866
Age 25-30†	33.1%			34.9%	33.8%		0.639
Age>30	46.5%	0.79***	(0.66, 0.93)	33.5%	35.0%		0.532
<i>Mother's Race/Ethnicity & Nativity</i>							
White	37.3%	1.16	(0.80, 1.66)	20.9%	20.2%		0.714
American Indian	10.7%	2.06***	(1.37, 3.09)	17.1%	17.5%		0.845
Hispanic	40.7%	1.50**	(1.04, 2.16)	55.9%	56.5%		0.803
Other race/ethnicity†	11.4%			6.0%	5.8%		0.833
Mother born in US†	74.4%			66.8%	66.2%		0.793
Mother born in Mexico	17.3%	1.70***	(1.39, 2.09)	27.7%	28.3%		0.783
Mother born outside US	8.3%	1.88***	(1.24, 2.84)	5.5%	5.5%		1.000
Hispanic & born outside US	1.4%	0.84	(0.39, 1.81)	1.2%	1.2%		1.000
<i>Mother's education</i>							
Less than high school†	20.1%			29.4%	31.4%		0.381
High school/GED	27.2%	0.97	(0.81, 1.16)	34.5%	33.5%		0.676
Some post-secondary education	30.2%	0.94	(0.77, 1.15)	30.0%	29.1%		0.705
4-year degree or more	21.4%	0.50***	(0.35, 0.71)	5.8%	5.5%		0.830
Education missing†	0.8%			0.2%	0.3%		0.873
<i>Method of payment</i>							
Private/commercial insurance	41.5%	0.94	(0.56, 1.56)	13.6%	13.6%		1.000
AHCCS (AZ Medicaid)	54.8%	2.56***	(1.58, 4.14)	84.1%	82.9%		0.506
Other insurance†	3.7%			2.3%	3.5%		0.143
<i>Marital cohabitation status</i>							
Married	56.8%	1.04	(0.68, 1.59)	43.2%	45.2%		0.426
Married & cohabiting	54.1%	0.97	(0.61, 1.53)	40.1%	42.6%		0.315
Mother & father cohabiting	76.6%	1.05	(0.87, 1.27)	68.1%	70.3%		0.335
<i>Health factors</i>							
First birth	28.4%	1.29***	(1.08, 1.53)	30.7%	29.9%		0.706

Pre-Existing Health Risk: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
<i>Mother's county of residence</i>							
Maricopa†	61.5%			33.5%	34.4%	0.715	0.111
Cochise, Santa Cruz	1.7%	11.81***	(9.16, 15.23)	12.1%	10.9%	0.438	
Gila	0.7%	3.62***	(2.08, 6.29)	1.8%	1.2%	0.314	
Mohave	1.3%	4.30***	(2.80, 6.61)	3.2%	3.3%	0.889	
Pima	14.4%	1.38**	(1.08, 1.76)	11.0%	12.1%	0.487	
Coconino, Navajo	4.9%	3.20***	(2.39, 4.29)	11.3%	10.6%	0.692	
Yavapai	1.6%	3.04***	(1.91, 4.84)	2.6%	1.8%	0.312	
Yuma	3.4%	5.27***	(4.19, 6.63)	16.5%	17.7%	0.512	
Apache, Graham, Greenlee, Pinal, La Paz	8.5%	1.36**	(1.00, 1.84)	6.9%	6.1%	0.547	
County missing	1.8%	1.32	(0.42, 4.22)	1.1%	1.7%	0.294	
<i>Median income of mother's zip code</i>							
Bottom income decile†	4.5%			4.3%	4.0%	0.804	0.084
Income decile 2	7.9%	1.60**	(1.09, 2.36)	15.2%	15.9%	0.682	
Income decile 3	10.9%	2.00***	(1.36, 2.94)	18.7%	20.6%	0.351	
Income decile 4	6.4%	1.76***	(1.15, 2.69)	9.8%	9.7%	0.934	
Income decile 5	8.6%	1.58**	(1.05, 2.38)	12.0%	11.3%	0.644	
Income decile 6	12.2%	1.19	(0.78, 1.83)	7.7%	7.1%	0.637	
Income decile 7	11.2%	1.66**	(1.10, 2.50)	12.0%	12.5%	0.763	
Income decile 8	10.7%	2.81***	(1.89, 4.17)	15.2%	14.6%	0.728	
Income decile 9	12.4%	0.91	(0.54, 1.55)	3.2%	3.1%	0.887	
Top income decile	9.9%	0.48*	(0.22, 1.07)	1.0%	0.5%	0.247	
Zip code missing	5.3%	0.32	(0.08, 1.32)	1.0%	0.9%	0.795	
<i>Birth certificate year</i>							
2006-07†	10.9%			0.2%	0.2%	1.000	0.121
2008	6.2%	2.80	(0.47, 16.78)	0.4%	0.1%	0.317	
2009	5.8%	8.35***	(1.77, 39.38)	1.0%	0.7%	0.592	
2010	6.0%	21.33***	(5.01, 90.84)	2.7%	2.4%	0.755	
2011	6.2%	46.48***	(11.31, 191.13)	6.5%	5.1%	0.245	
2012	6.8%	52.98***	(12.97, 216.43)	8.8%	7.2%	0.237	
2013	8.0%	57.56***	(14.14, 234.22)	11.3%	13.5%	0.176	
2014	15.5%	61.20***	(15.17, 246.90)	23.5%	25.0%	0.489	
2015	17.2%	62.52***	(15.51, 252.07)	25.8%	25.8%	1.000	
2016	17.6%	46.31***	(11.46, 187.06)	19.8%	19.8%	1.000	
<i>Other balancing factors</i>							
County, median income missing	0.01	3.85	(0.53, 28.16)	0.01	0.01	0.781	

*** p<.01; ** p<.05; * p<.01

‡ p-value for difference in outcome rates between HSP and matches accounts for estimated propensity score.

† Reference category/categories.

HSP: Health Start Participants; MCG: Matched Control Group; SD: Standardized Differences (variables w/more than two categories).

Age Less than 20: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
N	95,750			1,264	6,810		
<u>Birth outcomes</u>							
Low birth weight	6.8%			7.0%	7.8%	0.455	
Very low birth weight	1.0%			1.2%	1.2%	0.918	
Extremely low birth weight	0.5%			0.5%	0.3%	0.440	
Preterm birth	8.7%			6.6%	9.5%	0.004	
<u>Covariates/Matching controls</u>							
<i>Mother's Race/Ethnicity & Nativity</i>							
White	23.9%	1.51**	(1.06, 2.17)	22.2%	22.9%	0.669	0.065
American Indian	9.7%	1.69**	(1.13, 2.51)	12.7%	12.1%	0.673	
Hispanic	59.7%	1.62***	(1.15, 2.30)	62.2%	63.0%	0.681	
Other race/ethnicity†	6.6%			2.9%	2.0%	0.123	
Mother born in US†	79.7%			83.2%	84.8%	0.286	0.043
Mother born in Mexico	18.3%	1.42***	(1.19, 1.68)	16.1%	14.6%	0.295	
Mother born outside US	2.0%			0.6%	0.6%	0.861	
<i>Mother's education</i>							
Less than high school†	55.3%			57.6%	57.3%	0.876	0.043
High school/GED	36.4%	0.83***	(0.74, 0.95)	34.7%	35.2%	0.802	
Some post-secondary education	7.6%	0.81*	(0.64, 1.01)	7.3%	6.9%	0.698	
4-year degree or more	0.0%			0.0%	0.1%	0.267	
Education missing	0.6%			0.3%	0.3%	0.989	
<i>Method of payment</i>							
Private/commercial insurance	11.6%	1.19	(0.80, 1.79)	6.0%	5.4%	0.493	0.027
AHCCCS (AZ Medicaid)	84.4%	2.11***	(1.51, 2.94)	90.6%	91.1%	0.629	
Other insurance	4.0%			3.4%	3.5%	0.913	
<i>Marital cohabitation status</i>							
Married	12.0%	1.09	(0.66, 1.79)	11.7%	11.3%	0.755	
Married & cohabiting	11.2%	0.94	(0.55, 1.60)	10.3%	10.0%	0.792	
Mother & father cohabiting	48.6%	0.99	(0.87, 1.13)	41.9%	42.3%	0.809	
<i>Health factors</i>							
First birth	80.6%	0.93	(0.80, 1.08)	80.8%	82.6%	0.237	
Pre-existing health risk	3.1%	0.88	(0.64, 1.21)	3.8%	3.6%	0.834	

Age Less than 20: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
<i>Mother's county of residence</i>							
Maricopa†	58.5%			22.2%	23.1%	0.601	0.089
Cochise, Santa Cruz	2.7%	25.32***	(21.01, 30.52)	24.5%	23.9%	0.710	
Gila	1.1%	9.17***	(6.53, 12.89)	3.8%	3.2%	0.386	
Mohave	2.7%	4.77***	(3.47, 6.55)	4.2%	4.0%	0.841	
Pima	13.4%	1.13	(0.86, 1.48)	5.5%	4.9%	0.530	
Coconino, Navajo	4.7%	7.21***	(5.68, 9.16)	13.0%	11.6%	0.303	
Yavapai	2.3%	5.28***	(3.86, 7.23)	4.4%	5.3%	0.266	
Yuma	4.5%	8.57***	(7.02, 10.45)	16.0%	17.6%	0.287	
Apache, Graham, Greenlee, Pinal, La Paz	8.3%	1.47**	(1.09, 1.98)	4.4%	4.0%	0.622	
County missing	1.8%	5.68***	(3.16, 10.20)	2.1%	2.4%	0.589	
<i>Median income of mother's zip code</i>							
Bottom income decile†	4.7%			2.0%	1.9%	0.885	0.085
Income decile 2	10.2%	2.59***	(1.68, 3.98)	20.2%	20.6%	0.767	
Income decile 3	14.9%	2.71***	(1.75, 4.20)	13.1%	14.1%	0.487	
Income decile 4	9.3%	3.53***	(2.25, 5.54)	10.9%	11.3%	0.752	
Income decile 5	10.6%	2.75***	(1.77, 4.29)	14.2%	15.1%	0.500	
Income decile 6	12.5%	1.79**	(1.12, 2.86)	6.3%	6.3%	1.000	
Income decile 7	10.8%	3.39***	(2.19, 5.26)	14.2%	12.0%	0.099	
Income decile 8	9.8%	5.30***	(3.42, 8.22)	13.1%	13.1%	0.953	
Income decile 9	7.1%	2.90***	(1.71, 4.92)	2.9%	3.2%	0.729	
Top income decile	2.8%	1.81	(0.88, 3.70)	0.9%	0.9%	0.834	
Zip code missing	7.2%	1.82*	(0.92, 3.57)	2.1%	1.6%	0.303	
<i>Birth certificate year</i>							
2006-07†	25.9%			0.9%	0.9%	1.000	0.049
2008	12.1%	3.33***	(1.62, 6.88)	1.5%	1.3%	0.610	
2009	10.9%	9.39***	(4.99, 17.66)	4.0%	3.9%	0.918	
2010	9.4%	25.82***	(14.20, 46.93)	8.9%	8.3%	0.571	
2011	8.3%	58.17***	(32.42, 104.36)	17.2%	18.0%	0.602	
2012	8.0%	62.60***	(34.90, 112.30)	18.4%	19.0%	0.721	
2013	7.3%	55.81***	(30.95, 100.64)	14.1%	13.4%	0.644	
2014	6.7%	58.33***	(32.34, 105.23)	13.3%	13.3%	1.000	
2015	5.9%	60.64***	(33.55, 109.59)	12.4%	11.9%	0.715	
2016	5.5%	45.33***	(24.89, 82.55)	9.2%	9.9%	0.542	
<i>Other balancing factors</i>							
Married & pre-existing health risk	0.3%	0.83	(0.31, 2.24)	0.4%	0.3%	0.739	
County, median income missing	0.01	0.64	(0.24, 1.68)	0.01	0.01	0.575	

*** p<.01; ** p<.05; * p<.01

‡ p-value for difference in outcome rates between HSP and matches accounts for estimated propensity score.

† Reference category/categories.

HSP: Health Start Participants; MCG: Matched Control Group; SD: Standardized Differences (variables w/more than two categories).

First Birth: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
N	356,315			3,003	18,391		
<u>Birth outcomes</u>							
Low birth weight	6.5%			6.7%	6.5%	0.722	
Very low birth weight	1.0%			0.8%	0.9%	0.583	
Extremely low birth weight	0.5%			0.3%	0.5%	0.222	
Preterm birth	7.9%			6.8%	7.4%	0.314	
<u>Covariates/Matching controls</u>							
<i>Mother's age</i>							
Age<20	21.7%	1.24***	(1.09, 1.42)	34.0%	34.2%	0.892	0.032
Age 20-24	32.2%	1.12**	(1.00, 1.25)	40.8%	41.9%	0.387	
Age 25-30†	28.3%			18.1%	17.3%	0.437	
Age>30	17.8%	0.88	(0.74, 1.03)	7.1%	6.6%	0.442	
<i>Mother's Race/Ethnicity & Nativity</i>							
White	46.3%	0.94	(0.78, 1.13)	27.2%	27.9%	0.564	0.042
American Indian	5.2%	1.19	(0.94, 1.50)	9.8%	10.0%	0.762	
Hispanic	38.2%	1.29***	(1.07, 1.56)	57.7%	57.6%	0.979	
Other race/ethnicity†	10.3%			5.3%	4.4%	0.119	
Mother born in US†	78.2%			77.6%	79.1%	0.150	0.038
Mother born in Mexico	13.6%	1.52***	(1.33, 1.72)	19.3%	17.9%	0.164	
Mother born outside US	8.3%	1.04	(0.81, 1.35)	3.1%	3.0%	0.763	
Hispanic & born outside US	1.3%	0.71	(0.41, 1.23)	0.6%	0.6%	1.000	
<i>Mother's education</i>							
Less than high school†	18.0%			27.7%	27.7%	0.957	0.032
High school/GED	29.0%	0.91*	(0.83, 1.01)	37.9%	37.4%	0.709	
Some post-secondary education	25.2%	0.92	(0.82, 1.03)	25.7%	25.5%	0.906	
4-year degree or more	26.3%	0.64***	(0.54, 0.76)	8.5%	8.9%	0.582	
Education missing†	1.4%			0.3%	0.4%	0.312	
<i>Method of payment</i>							
Private/commercial insurance	45.7%	1.18	(0.93, 1.49)	16.1%	16.0%	0.972	0.014
AHCCS (AZ Medicaid)	49.3%	2.93***	(2.36, 3.64)	80.4%	80.2%	0.820	
Other insurance†	5.0%			3.5%	3.8%	0.583	
<i>Marital cohabitation status</i>							
Married	47.4%	1.05	(0.83, 1.32)	28.0%	27.8%	0.840	
Married & cohabiting	45.0%	0.88	(0.68, 1.13)	25.3%	25.7%	0.722	
Mother & father cohabiting	71.0%	1.00	(0.92, 1.10)	54.9%	55.6%	0.551	
<i>Health factors</i>							
Pre-existing health risk	6.4%	1.13*	(0.98, 1.29)	8.4%	7.7%	0.367	

First Birth: Birth Outcomes

	Propensity Score Model (Logit, Y=Health Start)			Match Results		p-value	
	Non-HSP	ORs	95% CI	HSP	MCG	difference‡	SD
<i>Mother's county of residence</i>							
Maricopa†	62.7%			20.7%	21.1%	0.680	0.056
Cochise, Santa Cruz	2.4%	22.42***	(19.78, 25.41)	20.0%	20.3%	0.748	
Gila	0.6%	10.12***	(7.87, 13.01)	2.7%	2.8%	0.752	
Mohave	2.1%	5.05***	(4.12, 6.20)	4.2%	3.9%	0.647	
Pima	14.3%	1.60***	(1.38, 1.87)	7.8%	8.1%	0.668	
Coconino, Navajo	3.5%	8.85***	(7.57, 10.34)	12.1%	11.2%	0.278	
Yavapai	2.2%	5.71***	(4.69, 6.95)	4.6%	4.0%	0.252	
Yuma	3.3%	12.82***	(11.33, 14.51)	21.3%	21.8%	0.638	
Apache, Graham, Greenlee, Pinal, La Paz	6.5%	1.89***	(1.56, 2.28)	4.6%	4.2%	0.488	
County missing	2.3%	2.69***	(1.59, 4.54)	2.1%	2.5%	0.301	
<i>Median income of mother's zip code</i>							
Bottom income decile†	2.5%			1.5%	1.2%	0.260	0.046
Income decile 2	6.1%	2.66***	(1.93, 3.65)	15.3%	15.4%	0.914	
Income decile 3	9.7%	3.09***	(2.24, 4.25)	13.8%	14.7%	0.319	
Income decile 4	6.0%	3.93***	(2.83, 5.45)	11.1%	11.4%	0.713	
Income decile 5	8.3%	3.11***	(2.25, 4.29)	15.4%	15.0%	0.719	
Income decile 6	11.6%	2.21***	(1.58, 3.08)	7.3%	7.0%	0.616	
Income decile 7	12.5%	3.31***	(2.40, 4.55)	16.0%	15.4%	0.547	
Income decile 8	11.2%	4.03***	(2.92, 5.55)	11.8%	11.7%	0.936	
Income decile 9	13.0%	2.12***	(1.47, 3.07)	3.2%	3.4%	0.719	
Top income decile	11.1%	1.54**	(1.01, 2.35)	1.6%	1.6%	1.000	
Zip code missing	8.1%	2.51***	(1.64, 3.84)	3.1%	3.3%	0.715	
<i>Birth certificate year</i>							
2006-07†	21.1%			0.8%	0.7%	0.881	0.004
2008	10.0%	3.24***	(1.91, 5.49)	1.2%	1.0%	0.621	
2009	9.4%	9.27***	(5.88, 14.63)	3.2%	2.9%	0.452	
2010	8.8%	27.27***	(17.77, 41.84)	8.4%	8.6%	0.782	
2011	8.6%	49.91***	(32.76, 76.05)	14.6%	15.1%	0.586	
2012	8.7%	55.53***	(36.47, 84.55)	16.7%	16.9%	0.809	
2013	8.4%	56.72***	(37.21, 86.45)	15.1%	14.1%	0.306	
2014	8.5%	55.20***	(36.22, 84.12)	15.1%	15.4%	0.693	
2015	8.3%	54.05***	(35.43, 82.44)	14.3%	14.0%	0.795	
2016	8.2%	41.16***	(26.89, 62.99)	10.8%	11.2%	0.650	
<i>Other balancing factors</i>							
Age<20 & born in Mexico	3.7%	0.88	(0.71, 1.08)	5.4%	4.9%	0.382	
County, median income missing	2.0%	1.24	(0.63, 2.43)	1.6%	2.1%	0.151	

*** p<.01; ** p<.05; * p<.01

‡ p-value for difference in outcome rates between HSP and matches accounts for estimated propensity score.

† Reference category/categories.

HSP: Health Start Participants; MCG: Matched Control Group; SD: Standardized Differences (variables w/more than two categories).