BMJ Open Cohort profile: an observational longitudinal data collection of health aspects in a cohort of female cancer survivors with a history of pelvic radiotherapy – a population-based cohort in the western region of Sweden

Linda Åkeflo ¹, Gail Dunberger,² Eva Elmerstig,³ Viktor Skokic,¹ Gunnar Steineck,¹ Karin Bergmark¹

To cite: Åkeflo L, Dunberger G, Elmerstig E, *et al.* Cohort profile: an observational longitudinal data collection of health aspects in a cohort of female cancer survivors with a history of pelvic radiotherapy—a population-based cohort in the western region of Sweden. *BMJ Open* 2021;**11**:e049479. doi:10.1136/bmjopen-2021-049479

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-049479>).

Received 25 January 2021
Accepted 06 July 2021

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Mrs Linda Åkeflo;
linda.akeflo@vgregion.se

ABSTRACT

Purpose The study 'Health among women after pelvic radiotherapy' was conducted in response to the need for more advanced and longitudinal data concerning long-term radiotherapy-induced late effects and chronic states among female cancer survivors. The objective of this paper is to detail the cohort profile and the study procedure in order to provide a sound basis for future analyses of the study cohort.

Participants Since 2011, and still currently ongoing, participants have been recruited from a population-based study cohort including all female patients with cancer, over 18 years of age, treated with pelvic radiotherapy with curative intent at Sahlgrenska University Hospital in Gothenburg, in the western region of Sweden, which covers 1.7 million of the Swedish population. The dataset presented here consists of baseline data provided by 605 female cancer survivors and 3-month follow-up data from 260 individuals with gynaecological, rectal or anal cancer, collected over a 6-year period.

Findings to date Data have been collected from 2011 onwards. To date, three studies have been published using the dataset reporting long-term radiation-induced intestinal syndromes and late adverse effects affecting sexuality, the urinary tract, the lymphatic system and physical activity. These projects include the evaluation of interventions developed by and provided in a nurse-led clinic.

Future plans This large prospective cohort offers the possibility to study health outcomes in female pelvic cancer survivors undergoing a rehabilitation intervention in a nurse-led clinic, and to study associations between demographics, clinical aspects and long-term late effects. Analysis focusing on the effect of the interventions on sexual health aspects, preinterventions and postinterventions, is currently ongoing. The cohort will be expanded to comprise the entire data collection from 2011 to 2020, including baseline data and data from 3-month and 1-year follow-ups after interventions. The data will be used to study conditions and treatment-induced late effects preintervention and postintervention.

Strengths and limitations of this study

- The major strength of the study is the large representative population-based cohort (n=605).
- Uses a large longitudinal dataset consisting of female cancer survivors' entries from 6 months to several years post pelvic radiotherapy treatment.
- Highlighting treatment-induced cancer survivorship diseases and chronic states improves the possibilities for developing further effective treatments.
- One limitation of the study is the reliance on self-reported data, which has the potential for response bias.
- The interventions that were provided varied over the course of the study, which is considered a potential limitation.

INTRODUCTION

An increasing number of individuals now live longer after a cancer diagnosis than was previously common, and more and more patients are living with treatment-induced late effects, treatment-induced cancer survivorship diseases and survivorship chronic states.^{1–3} Pelvic radiotherapy affects the intestinal and urinary tracts, the lymphatic system and sexual health.^{4–10} In a video at <https://www.jostrust.org.uk/video/letstalk-pelvic-radiation-disease-after-cervical-cancer>, some of more than 20 million cancer survivors in Europe describe how radiation-induced impaired intestinal health affects them. In Sweden, approximately 3500 women are diagnosed with pelvic cancer (gynaecological, anal or rectal cancer) every year.¹¹ In 2011, we started a nurse-led clinic focusing on treatment-induced late effects among pelvic cancer survivors with the intent of developing

self-care strategies and treatments. We have previously reported on the benefits of the clinic regarding the improved quality of life and psychosocial well-being among female pelvic cancer survivors.¹²

We aim to describe a population-based cohort consisting of female cancer survivors treated with pelvic radiotherapy using a dataset from a longitudinal study that has been ongoing since 2011. A further aim is to describe the data collection procedure, the interventions provided in a nurse-led clinic and the characteristics of the study cohort. In addition, a few basal empirical results will be presented to illustrate the study cohort. The objective of this paper is to detail the cohort profile and the study procedure in order to provide a sound basis for future analyses in the study cohort.

COHORT DESCRIPTION

Setting

In 2009, the Swedish government proposed a new regulation: 'A National Cancer Strategy' (SOU2009:1).¹³ In collaboration with the Swedish county councils and regions, six Regional Cancer Centres (RCC) were established. Based on the national strategy proposals, the RCC in Western Sweden financed a nurse-led clinic at Sahlgrenska University Hospital. The nurse-led clinic was founded in 2011 by a senior consultant physician who has a PhD and is specialised in gynaecological oncology, and an oncology nurse with a PhD in oncology. The team currently consists of three clinical oncology nurses who are specially trained and clinically specialised in understanding and addressing issues concerning pelvic cancer survivorship late effects. The cancer survivors receive education about radiotherapy-induced late effects, basic anatomy and physiology and together with the nurse, decide on supportive care actions regarding medication, nutrition, and coping with sexual, psychological and social challenges. In cases with more specific needs, intercurrent diseases, or a suspected recurrence of cancer, the patient meets the senior medical consultant who has the primary medical responsibility for the clinic. The senior medical consultant meets regularly with the team and discusses plans for patients' medication with the staff member who will issue the prescriptions. The healthcare programme developed in the clinic is based on programmes developed by others and on our previous studies in this area.^{4 14–20}

Patient and public involvement

Patients and/or the public are not involved in the design, conducting, reporting or dissemination plans in this research. Results from future studies from the cohort will be submitted for publication in peer-reviewed journals and presented at relevant conferences.

Participants

Study participants are recruited from two different cohorts: (1) a population-based study cohort group

including all female patients with cancer treated with curative intent from 2007 onwards who are continuously identified from medical records at Sahlgrenska University Hospital in Sweden and (2) a referred patient group including all female patients with cancer referred to the rehabilitation clinic. Inclusion criteria are female cancer survivors who completed pelvic radiotherapy at least 6 months prior to inclusion. Exclusion criteria are cancer recurrence, inability to comprehend or answer the questionnaire, and poor proficiency in the Swedish language. Data are collected from female cancer survivors with a history of pelvic radiotherapy in Western Sweden, an area that includes 20% of the Swedish population. The tradition of collecting registry data in Sweden offers good opportunities for studying cancer survivors without selection-induced problems. The dataset contains information about the preintervention and postintervention needs of women who received pelvic radiotherapy at Sahlgrenska University Hospital between 2007 and 2016. Female cancer survivors referred to the nurse-led clinic are also invited to participate in the study.

Invited women

An introductory letter is sent to eligible study participants. Shortly after, a research secretary phones them and gives oral information about the study, asks if they are willing to participate and, if so, asks them to give permission to be sent a written informed consent and the questionnaire. When the baseline questionnaire is returned, the participant receives an invitation to attend the nurse-led clinic where individual healthcare interventions are conducted. The code number from the questionnaire (followed throughout the data collection) is entered into the patient-database FileMaker Pro17 Advanced that was specifically designed to suit the study. The study participant receives a follow-up questionnaire three and twelve months after the completed intervention. The research secretary keeps track of each patient and, in case questionnaires are delayed or missing, sends two reminders at time points determined in advance (see figure 1).

Referred patients

Patients referred from oncology healthcare providers, general practitioners and through private referrals, who meet the inclusion criteria are also invited to participate in the study. The data collection procedure follows the same principles as for the invited women in the population-based cohort.

Study-specific questionnaires

As shown in table 1, the baseline questionnaire consists of 175 questions, divided into eight main sections and one concluding chapter. Information is obtained about the frequency, intensity, duration, and quality of each symptom, and the degree of distress it causes the patient. An example of one question is: 'how many times per week (approximately) have you had bowel movements during the past 6 months?' with possible answers: 'about

Figure 1 Flowchart of pelvic cancer rehabilitation ongoing data collection procedure and the procedure for patient inclusion, contact with participants and the follow-up points for questionnaires.

every other day', 'less than once a day', 'once a day', 'up to two times a day', '2–3 times per day', '3–4 times per day' or 'five times or more per day'. To facilitate statistical analyses, the answers are coded into values, ranked and placed into groups. Variables such as age and number of children are categorised into groups.

Several sections allow free comments

The questionnaire also addresses the extent to which the symptoms affect social functioning. In the concluding chapter, the participants are invited to visit the nurse-led clinic. Some questions require the patients to rank their most distressing symptoms. In the follow-up questionnaires, the study participants' health status and symptoms are measured, and the interventions conducted at the nurse-led clinic are evaluated.

Diagnostic and alleviative means

The questionnaire serves as a basis for the interventions conducted at the first visit to the clinic. The patient's current health status reveals the areas that need to be focused on. Current symptoms and self-care strategies are discussed. New strategies are then suggested and decided on together with the patient. The duration of the contact and the number of follow-ups varies from 1 month to 1 year depending on the symptoms and the effect of treatments and interventions.

Intestinal health

The assessment is based on five syndromes: urgency to defecate, faecal leakage, excessive gas production, excessive mucus discharge and blood discharge.²¹ Symptoms, such as leakage of mucus and blood, abdominal bloating, signs of bacterial overgrowth and bile salt malabsorption, are also assessed. Diagnostic means, including blood tests such as electrolytes, blood counts, vitamin B12 and serum magnesium, are taken when indicated. An algorithm

developed by the research group serves as a guide in the clinical setting.¹² A mobile application has also been developed that was specially designed for radiation-induced intestinal syndromes and which can show graphs of the variation in intestinal health over time. The application serves as a complement to existing methods of patients' self-assessment and self-management, and helps to support the nurse's and patient's conversation and decision-making.^{22 23} The objective of the interventions is to restore or improve intestinal health and includes medical treatment, pelvic floor muscle training and techniques using cognitive training to control intestinal function. Details of these interventions can be accessed elsewhere.^{12 24}

Sexual health

The PLISSIT-model (Permission, Limited Information, Specific Suggestions, and Intensive Therapy) developed by Annon²⁵ is used to address sexual health concerns. The first three levels in the model are helpful for most patients. When specific suggestions are not sufficient, the patient is recommended to take part in intensive therapy, which is sometimes given in the clinical setting and other times with a sexologist or psychotherapist after a referral is sent. Female sexual dysfunction is a complex condition, including diverse aspects, definitions and classifications.²⁶ To assess and treat sexual dysfunction, previous sexual practices and experiences are discussed, with particular attention given to patients' integrity, special needs and preferences. Structured information about common physical late effects involving vaginal changes and sexual problems is provided as well as information concerning psychological issues influencing sexual health; all information given is in accordance with evidence-based knowledge and practice in the field.^{1 4 17 27–29} Patients receive information about and instructions on the use of vaginal dilator therapy and topical oestrogen, and they are also given guidance and suggestions related to lack of sexual desire. When requested, specific sexual devices, films and literature are suggested to encourage and help the patient to regain sexual function, which for some patients is helpful in managing issues of body image perception, self-esteem and intimacy.

Sexual abuse

Patients with a history of sexual abuse have an increased risk of developing sexual problems after cancer treatment, and a large proportion of women with cervical cancer and those experiencing dyspareunia postcancer treatment have been sexually abused.³⁰ In cases where this is reported, experiences of sexual abuse are carefully and sensitively discussed with the patient and, in some cases, this may lead to referral to a psychologist or therapist.

Urinary tract health

Symptoms of urgency, nocturia, urinary retention, urinary tract infection and pelvic pain are addressed. The diagnostic methods used include evaluations of urinary

Table 1 The study-specific baseline questionnaire, divided into eight sections and one concluding chapter

Section	Question areas
Sociodemographic	Gender Age Marital status Number of children Level of education Employment status
Quality of life and well-being	Quality of life Depression Worry Anxiety
Body perception and self-image	Femininity Self-esteem Fertility Childbirth Vaginal and perianal injury related to childbirth or physical trauma
Intestinal and defecation habits	Loose stools Faecal incontinence Urgency to defecate Excessive gas Abdominal pain Having a stoma Medical treatment related to intestinal symptoms
Micturition habits and urinary tract symptoms	Urinary frequency Urgency to urinate Nocturia Having a urinary catheter Medical treatment due to symptoms
Sexual health	Menopause Use of systemic hormone replacement therapy Use of topical oestrogen Impaired lubrication Vaginal shortness Vaginal inelasticity Dyspareunia
Sexual abuse	Experience of sexual abuse Experience of sexual harassment Age when experienced sexual abuse or harassment Extent to which the experience affects sexual life
Lymphoedema	Heaviness in legs, genitals and abdomen If diagnosed with lymphoedema, current lymphoedema treatment
Concluding chapter	Self-reported needs Invite to visit the nurse-led clinic for assessment and counselling

frequency and of symptoms, which are divided into irritative symptoms, obstructive symptoms and bleeding. Patients can keep a 72-hour voiding diary, including urine volume and fluid intake, which is a helpful resource for this discussion. Guidance and recommendations are given about topical oestrogen, pelvic floor muscle training, medical treatments, self-care modifications and behavioural interventions, all in accordance with previous evidence-based knowledge.^{31 32} In cases of severe urinary tract symptoms, referrals are sent to a urotherapist, physiotherapist, or urologist.

Lymphoedema

Lower limb lymphoedema is a common non-curable chronic complication with multifactorial pathophysiology.³³ Early detection and treatment of lymphoedema are essential in order to prevent complications.

Self-reported or objectively assessed swelling or heaviness in the lower limbs leads to referral for lymphatic therapy. The primary therapy is the use of individually tailored compression garments and second-line therapy consists of manual lymphatic drainage with intermittent use of pumps and other self-care strategies.^{34 35}

Statistical analyses

Initially, data from the questionnaires were entered into EpiDataSoftware V.3.1 (EpiData Association). Since only one answer to each question is allowed, in the event of two answers the procedure was to alternate entering the first provided answer and the second provided answer. In order to facilitate the analysis of responses to the open-ended questions, the answers to these questions were transcribed in Microsoft Word (2016). R V.3.5.2. was

used for statistical analysis of the data. The results will be reported in means, medians and percentages.

FINDINGS TO DATE

Three studies based on the cohort have been published.^{12 36 37} Two papers focused on gastrointestinal side effects and one on sexual health aspects. It was found that the majority of the women reported a change in bowel habits and, in almost half of them, the effect was considerable.¹² Women with faecal leakage were less likely to practise physical activity than survivors without leakage, and survivors who practised weekly physical activity experienced better quality of life and were less frequently in a depressed mood than women not physical active.³⁶ Furthermore, a statistically significant increase in psychological distress and sexual health impairment was found among women with a history of sexual abuse compared with women without such experience.³⁷

Characteristics of study participants

From January 2011 to June 2017, we identified a total of 791 patients in the population-based cohort who met the inclusion criteria. Of the total sample, 684 (86%) individuals gave oral consent to participate in the study and of these, 464 (68%) completed the baseline questionnaire.

During the same period, 184 referred patients met the inclusion criteria and were invited to participate in the study. Of those invited to participate, 141 (76.6%) completed the baseline questionnaire and of these, 131 (92%) took part in interventions.

The responses and completeness of the questionnaires from both study groups are shown in the schematic diagram (see figure 2). In table 2, we present the baseline characteristics and demographics in the dataset, which consists of 605 participants: 464 (76.7%) from the invited cohort and 161 (23.3%) from the referred cohort.

The majority of the study participants had a history of gynaecological cancer and had been treated with radiotherapy in combination with surgery. In the population-based study cohort, the mean age was higher and two times as many were retired compared with the referred group. In the referred group, two times as many were on sick leave. In total, 379 (63%) of the 605 study participants agreed to visit the clinic (see table 3). Sixty-seven (14.4%) of the women in the population-based study group who declined to visit the clinic reported that they had radiotherapy-induced late effects.

In this paper, we describe the population-based cohort consisting of female cancer survivors treated with pelvic radiotherapy in the western region of Sweden. The data

Figure 2 Schematic diagram of data collected from two different study cohorts from January 2011 to June 2017: a population-based cohort and patients referred from oncology healthcare providers, general practitioners and through private referrals who met the inclusion criteria. The diagram includes study response rate, completeness of questionnaires and reasons for loss of participants.

Table 2 Characteristics of the study participants

Variable	Total	Invited	Referred
Participants, n (%)	605	464 (76.7)	141 (23.3)
Cancer type, n (%)			
Endometrial cancer	216 (35.7)	181 (39.0)	35 (24.8)
Cervical cancer	132 (21.8)	80 (17.2)	52 (36.9)
Ovarian cancer	2 (0.3)	1 (0.2)	1 (0.7)
Vaginal cancer	5 (0.8)	3 (0.6)	2 (1.4)
Vulvar cancer	21 (3.5)	19 (4.1)	2 (1.4)
Anal cancer	80 (13.2)	58 (12.5)	22 (15.6)
Rectal cancer	145 (24.0)	122 (26.3)	23 (16.3)
Other	4 (0.7)		4 (2.8)
Age in years			
Mean	64.5	66.5	57.6
SD	12.6	11.5	13.6
Missing, (%)	11	7 (1.5)	4 (2.8)
Years since radiotherapy, grouped			
0	35 (5.8)	6 (1.3)	29 (20.6)
1	219 (36.1)	166 (35.8)	53 (37.6)
2	98 (16.1)	86 (18.5)	12 (8.5)
3	139 (23.0)	131 (28.2)	8 (5.7)
≥4	105 (17.3)	69 (14.9)	36 (25.5)
Missing	9 (1.5)	6 (1.3)	3 (2.1)
Mean	2.6	2.2	3.9
SD	3.4	1.2	6.7
Cancer treatment, n (%)			
External radiotherapy, only	145 (24.0)	101 (21.8)	44 (31.2)
External radiotherapy and brachytherapy	20 (3.3)	16 (3.4)	4 (2.8)
External radiotherapy, brachytherapy and surgery	180 (29.7)	157 (33.8)	23 (16.3)
External radiotherapy and surgery	260 (43.0)	190 (40.9)	70 (49.6)
Marital status, n (%)			
Married or living with a partner	402 (66.4)	309 (66.6)	93 (66.0)
Widow	70 (11.6)	59 (12.7)	11 (7.8)
Has a partner but lives alone	30 (5.0)	15 (3.2)	15 (10.6)
Single	102 (17.0)	80 (17.2)	22 (15.6)
Missing	1 (0.2)	1 (0.2)	
Education level, n (%)			
Elementary school	173 (29.1)	150 (32.3)	23 (16.3)
Secondary school	227 (38.2)	169 (36.4)	58 (41.1)
College/university	194 (32.0)	135 (29.1)	59 (41.8)
Missing	11 (1.8)	10 (2.2)	1 (0.7)
Employment status, n (%)			
Student	4 (0.7)	3 (0.6)	1 (0.7)
Unemployed job seeker	12 (2.0)	10 (2.2)	2 (1.4)
Employed	162 (27.0)	116 (25.0)	46 (32.6)
Housewife	4 (0.7)	2 (0.2)	2 (1.4)
On sick leave	54 (9.0)	17 (3.7)	37 (26.2)

Continued

Table 2 Continued

Variable	Total	Invited	Referred
Disability pension	35 (5.7)	26 (5.6)	9 (6.4)
Retired	328 (54.7)	284 (61.2)	44 (31.2)
Missing	6 (1.0)	6 (1.3)	
Resident, n (%)			
In a big city	182 (30.6)	131 (28.2)	51 (36.2)
In a small or medium-sized city	309 (50.2)	244 (52.6)	61 (43.3)
In the countryside	116 (18.9)	87 (18.8)	29 (20.6)
Missing	2 (0.3)	2 (0.4)	
Smoking, n (%)			
Does not smoke	448 (74.0)	339 (73.1)	109 (77.3)
Smokes	67 (11.0)	52 (11.2)	17 (12.1)
Missing	88 (14.5)	73 (15.7)	15 (10.6)

N (number) and proportion (%) of women are presented.

collection procedure, the interventions provided and the characteristics of the study cohort are also outlined. In addition, a few basal empirical results are reported to illustrate the study population. The major strength of the study is the large population-based cohort since it creates the possibility of studying cancer survivors without selection-induced problems and makes it representative of the reference population consisting of an increasing number of female cancer survivors treated with pelvic radiotherapy. The longitudinal study design enables future investigations of long-term treatment-induced late effects, diseases and chronic states. Moreover, it will be possible to evaluate the long-term outcomes of interventions and the treatments provided.

The data collection has generated a large dataset consisting of patient entries from 6 months to several years post pelvic radiotherapy. Over a period of six and a half years, almost 1000 female pelvic cancer survivors have been invited to participate in the study; the dataset consists of 605 cancer survivors. In the population-based cohort group, 68% completed a baseline questionnaire. Treatment and interventions concerning physical, psychosocial and sexual issues were offered to patients in both the population-based cohort and the referred cohort, and these were evaluated. As shown in table 3, 53.4% of the study participants in the population-based cohort agreed to visit the clinic, which may indicate

the proportion of cancer survivors with unmet needs. Improved self-care strategies, increased clinical knowledge and developments in technology currently provide healthcare professionals with possible methods to help cancer survivors manage and treat the late effects of pelvic radiotherapy treatment. We believe that highlighting treatment-induced cancer survivorship diseases and chronic states may increase the likelihood of further effective treatments being developed.

Cancer survivorship issues have advanced from being neglected to gradually being given increasingly greater attention in healthcare practice as well as in research. This is reflected by the development of national guidelines and the organisation of national and international scientific conferences and meetings in this subject area. Efforts are being made to both understand how treatment-induced late effects manifest in pelvic cancer survivors and to find strategies to deal with these late effects. Twenty years have passed since researchers within our team⁴ first observed that patients with gynaecological cancer suffered from vaginal changes affecting their sexual health. More recently, we have identified five syndromes impairing pelvic cancer survivors' intestinal health. Results published in 2017²¹ simplify the search for ways to prevent, manage and help minimise the occurrence and intensity of survivorship diseases. One promising ongoing study being conducted by Schofield *et al* in Australia is

Table 3 Number (N) and proportion (%) of study participants who agreed to visit the clinic

	Total N (%)	Invited N (%)	Referred N (%)
No, I have no late effects and do not need to visit the clinic	108 (17.8)	107 (23.1)	1 (0.7)
No, I have late effects, but I do not want to visit the clinic	71 (11.7)	67 (14.4)	4 (2.8)
Yes, I want to visit the clinic	379 (62.6)	248 (53.4)	131 (92.9)
Missing	47 (7.8)	42 (9.1)	5 (3.5)

N (number) and proportion (%) of women are presented.

evaluating a care programme similar to ours.^{38 39} Furthermore, in 2015 Andreyev *et al*¹⁵ published a guide for the management of intestinal problems, including an algorithm that, in our opinion, is a useful tool for clinicians. However, the pathophysiological changes, which are described in numerous previous studies, are not yet fully understood.^{19–21} Inflammatory and fibrotic processes in the gut wall are probably of importance in explaining the various symptoms. The processes may relate to both the intestinal tract and other organs located in the pelvis. Hofsjö *et al* in our research group⁴⁰ recently studied vaginal changes and found morphological explanations for changes in the vaginal wall. Biopsies from the vaginal connective tissue affected by radiotherapy showed dense collagen and entangled elastin fibres, a finding that may explain common symptoms such as reduced vaginal elasticity during intercourse, reduced lubrication, and dyspareunia.

In Sweden, promising steps are being taken towards developing a programme for national coordination of cancer rehabilitation practice. Cerna *et al*,²³ in a study concerning self-management from an educational perspective, observed that patients and nurses can together create tailor-made solutions. The nurses focus on encouraging the patients to self-reflect and on maintaining the patient's motivation to continue to engage in self-care. The ongoing establishment of oncology nurse navigators⁴¹ is also an initiative that has the potential to increase the supportive care given to cancer survivors. In the 1980s, research in the psychological field⁴² showed that patients have a high risk of anxiety related to oncology treatment. Recent studies report that patients have a lower risk of future anxiety and depression when their needs are addressed during treatment.^{43 44} In our opinion, when planning future follow-up in clinical cancer care, these findings need to be taken into account, irrespective of the patient's cancer diagnosis. We suggest that, in the future, healthcare should provide advanced specialist expertise in the management of severe treatment-induced late effects.

Sexual concerns are generally not addressed or discussed as much as patients would like,⁴⁵ and patients generally wait for healthcare professionals to raise the subject.⁴⁶ In our clinical setting, sexual health conversations are integrated into the clinical work, routines for assessment and treatment have been created, and it becomes clearer when a patient should be referred to specialists and sexologists. The PLISSIT-model²⁵ has been used for addressing sexual health concerns in both our own clinical setting and in that of others. Clinical experience shows that sexual function might improve and even return to prediagnosis level through frequent clinical follow-up, which is consistent with results from previous studies.⁴⁷ Parts of WHO's definition of sexual and reproductive health state that: 'sexual health requires a positive and respectful approach to sexuality and sexual relationships',⁴⁸ so healthcare professionals need to be able to talk openly to patients about sexuality. It has

been suggested that healthcare professionals should actively engage in training to improve their communication skills in order to overcome common communication barriers.^{49 50} Our clinical experience has shown that several patients have felt relieved when being able to speak openly about sexual issues as well as other private health concerns. The baseline questionnaire seems to serve as a therapeutic tool that can be followed up in counselling.

One could argue that the observational study design is a limitation in this study and that a randomised clinical trial might provide data that are more reliable. However, for ethical reasons, we found it necessary to offer all patients in the population the best available intervention. Hence, we considered that the observational study design was best suited to this purpose, a design previously shown to produce reasonably useful results.⁵¹

One important limitation of the study is the reliance on self-reported data, which has the potential for response bias. However, we considered self-reported data to provide a wider range of responses than data collected using other data collection instruments.⁵² To avoid information-related problems, we took advance preventive action by using questionnaires based on the clinometric method that we have introduced, developed, used and described in previous research projects.^{4–6 53} We employed epidemiological methods, introduced into the cancer survivorship field by a hierarchical step-model, to manage bias and confounding.⁵⁴ This method was considered appropriate to measure causal relationship, and to examine different symptoms and characteristics in treatment-induced side effects. In short, the method comprises a quantitative prephase of semi-structured interviews with persons suitable for the study. Thereafter, a face-to-face-validation is conducted to ensure satisfactory internal consistency.

The wide range in length of time since completion of treatment may be considered a weakness in the study; however, this will probably also facilitate measurement of symptom progression in future analyses. Another potential limitation is that the interventions provided varied over the course of the study due to the increase in both ours and others' understanding of the complexity and pathophysiological mechanisms of symptoms and late effects. The minor changes in the interventions during the study need to be considered in future analysis. The results may also be affected by non-participation and loss to follow-up. We can only speculate about the reasons for these, such as patients not having time, not feeling motivated or being unwilling to recall their previous cancer experience. Possible reasons for patients declining an offer to visit the clinic despite having troublesome symptoms could be due to the symptoms being too severe to enable travel to the clinic. Age-related problems and long-distance transportation can be other reasons. The data collected from the referred patients will allow analysis of the prevalence of symptoms and unmet needs observed by other healthcare providers. Since the data

were collected in Sweden, we do not know to what extent our analysis will be applicable to other populations.

Through frequent lectures for patients and health-care professionals, and the implementation of treatment strategies directly to patients, we apply the knowledge we have acquired and use it in the clinic and with the cancer survivors themselves. It is also worth mentioning that our rehabilitation clinic serves as a model for similar clinics in other regions of Sweden that have been established as part of the national strategy financed by the Swedish government. The current nurse-led clinic may serve as the beginning of a future tertiary centre to develop interventions and treatments for cancer survivors.

The extent to which the interventions provided in the individualised nurse-led rehabilitation might improve health in female cancer survivors treated with pelvic radiotherapy is currently unclear. The dataset from the 3-month follow-up questionnaire was prepared and used in recently published studies, while preparation of the dataset from the 1-year follow-up questionnaire is ongoing. To the best of our knowledge, our study cohort is one of only a few published population-based cohorts of female pelvic cancer survivors with treatment-induced late effects receiving individualised interventions with a focus on physical and sexual health after radiotherapy. The interventions developed, outlined and provided will hopefully contribute to further development of evidence-based management strategies in pelvic cancer rehabilitation and will be reported in future papers.

Author affiliations

¹Division of Clinical Cancer Epidemiology, Department of Oncology, Institute of Clinical Science, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

²Department of Health Care Sciences, Ersta Sköndal Bräcke University College, Stockholm, Sweden

³Centre for Sexology and Sexuality studies, Malmö Universitet, Malmö, Sweden

Contributors Guarantor of integrity of the entire study: GS. Study concepts and design: KB, GS, GD. Data analysis: LA, GD, KB, EE, GS. Statistical analysis: VS. Manuscript preparation: LA, KB, GD, EE. Manuscript editing: LA. All authors have read and approved the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval All procedures in the study, which involved human participants, were in accordance with the ethical standards of the Regional Ethics Review Board in Gothenburg (D 686-10) and with the 1964 Declaration of Helsinki and its later amendments. The research was approved by the Regional Ethics Review Board in Gothenburg (D 686-10), Gothenburg University. Written informed consent was obtained from all individual participants included in the study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given,

and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Linda Åkeffo <http://orcid.org/0000-0002-6795-1092>

REFERENCES

- Allemani C, Matsuda T, Di Carlo V, *et al*. Global surveillance of trends in cancer survival 2000–14 (CONCORD-3): analysis of individual records for 37 513 025 patients diagnosed with one of 18 cancers from 322 population-based registries in 71 countries. *Lancet* 2018;391:1023–75.
- Miller KD, Siegel RL, Lin CC. Cancer treatment and survivorship statistics 2016;66:271–89.
- Steineck G, Schmidt H, Alevronta E, *et al*. Toward restored bowel health in rectal cancer survivors. *Semin Radiat Oncol* 2016;26:236–50.
- Bergmark K, Åvall-Lundqvist E, Dickman PW, *et al*. Vaginal changes and sexuality in women with a history of cervical cancer. *N Engl J Med* 1999;340:1383–9.
- Dunberger G, Lind H, Steineck G, *et al*. Self-Reported symptoms of faecal incontinence among long-term gynaecological cancer survivors and population-based controls. *Eur J Cancer* 2010;46:606–15.
- Lind H, Waldenström A-C, Dunberger G, *et al*. Late symptoms in long-term gynaecological cancer survivors after radiation therapy: a population-based cohort study. *Br J Cancer* 2011;105:737–45.
- Bober SL, Kingsberg SA, Faubion SS. Sexual function after cancer: paying the price of survivorship. *Climacteric* 2019;22:558–64.
- Sekse RJT, Dunberger G, Olesen ML, *et al*. Lived experiences and quality of life after gynaecological cancer-An integrative review. *J Clin Nurs* 2019;28:1393–421.
- Morris KA, Haboubi NY. Pelvic radiation therapy: between delight and disaster. *World J Gastrointest Surg* 2015;7:279–88.
- Yoshihara M, Shimono R, Tsuru S, *et al*. Risk factors for late-onset lower limb lymphedema after gynecological cancer treatment: a multi-institutional retrospective study. *Eur J Surg Oncol* 2020;46:1334–8.
- Socialstyrelsen. Statistical database, 2017. Available: <https://www.socialstyrelsen.se/en/statistics-and-data/statistics/>
- Dunberger G, Bergmark K. Nurse-Led care for the management of side effects of pelvic radiotherapy: what does it achieve? *Curr Opin Support Palliat Care* 2012;6:60–8.
- 2009:11 S (2009) A National Strategy for the Future - Summary. Available: <https://www.regeringen.se/49b6a7/contentassets/e343b40615eb46b395e5c65ca38d1337/summary-sou-200911>
- Andreyev J. Gastrointestinal symptoms after pelvic radiotherapy: a new understanding to improve management of symptomatic patients. *Lancet Oncol* 2007;8:1007–17.
- Andreyev HJN, Muls AC, Norton C, *et al*. Guidance: the practical management of the gastrointestinal symptoms of pelvic radiation disease. *Frontline Gastroenterol* 2015;6:53–72.
- Schover LR, van der Kaaij M, van Dorst E, *et al*. Sexual dysfunction and infertility as late effects of cancer treatment. *European Journal of Cancer Supplements* 2014;12:41–53.
- Schover LR, Fife M, Gershenson DM. Sexual dysfunction and treatment for early stage cervical cancer. *Cancer* 1989;63:204–12.
- Bergmark K, Åvall-Lundqvist E, Dickman PW, *et al*. Patient-rating of distressful symptoms after treatment for early cervical cancer. *Acta Obstet Gynecol Scand* 2002;81:443–50.
- Dunberger G, Lind H, Steineck G, *et al*. Loose stools lead to fecal incontinence among gynecological cancer survivors. *Acta Oncol* 2011;50:233–42.
- Andreyev HJN, Wotherspoon A, Denham JW, *et al*. "Pelvic radiation disease": new understanding and new solutions for a new disease in the era of cancer survivorship. *Scand J Gastroenterol* 2011;46:389–97.
- Steineck G, Skokic V, Sjöberg F, *et al*. Identifying radiation-induced survivorship syndromes affecting bowel health in a cohort of gynecological cancer survivors. *PLoS One* 2017;12:e0171461.
- Lindroth T, Islind AS, Steineck G. From narratives to numbers: data work and patient-generated health data in consultations. *Studies in health technology and informatics* 2018;247.
- Cerna K, Ivarsson J, Weilenmann A. Supporting self-management of radiation-induced bowel and bladder dysfunction in pelvic-cancer rehabilitation: an ethnographic study. *Journal of clinical nursing* 2019.
- Andreyev HJN, Davidson SE, Gillespie C, *et al*. Practice guidance on the management of acute and chronic gastrointestinal problems arising as a result of treatment for cancer. *Gut* 2012;61:179–92.

- 25 Annon JS. The PLISSIT model: a proposed conceptual scheme for the behavioral treatment of sexual problems. *J Sex Educ Ther* 1976;2:1–15.
- 26 Basson R, Wierman ME, van Lankveld J, *et al.* Summary of the recommendations on sexual dysfunctions in women. *J Sex Med* 2010;7:314–26.
- 27 Carter J, Goldfrank D, Schover LR. Simple strategies for vaginal health promotion in cancer survivors. *J Sex Med* 2011;8:549–59.
- 28 DuHamel K, Schuler T, Nelson C, *et al.* The sexual health of female rectal and anal cancer survivors: results of a pilot randomized psycho-educational intervention trial. *J Cancer Surviv* 2016;10:553–63.
- 29 Faithfull S, White I. Delivering sensitive health care information: challenging the taboo of women's sexual health after pelvic radiotherapy. *Patient Educ Couns* 2008;71:228–33.
- 30 Bergmark K, Åvall-Lundqvist E, Dickman PW, *et al.* Synergy between sexual abuse and cervical cancer in causing sexual dysfunction. *J Sex Marital Ther* 2005;31:361–83.
- 31 Aoki Y, Brown HW, Brubaker L, *et al.* Urinary incontinence in women. *Nat Rev Dis Primers* 2017;3:17042.
- 32 Leddy LS. Management of lower urinary tract symptoms after pelvic radiation in females. *Curr Urol Rep* 2018;19:106.
- 33 Lawenda BD, Mondry TE, Johnstone PAS. Lymphedema: a primer on the identification and management of a chronic condition in oncologic treatment. *CA: A Cancer Journal for Clinicians* 2009;59:8–24.
- 34 Dunberger G, Lindquist H, Waldenström A-C, *et al.* Lower limb lymphedema in gynecological cancer survivors--effect on daily life functioning. *Support Care Cancer* 2013;21:3063–70.
- 35 Bakar Y, Tuğral A. Lower extremity lymphedema management after gynecologic cancer surgery: a review of current management strategies. *Ann Vasc Surg* 2017;44:442–50.
- 36 Lindgren A, Dunberger G, Steineck G, *et al.* Identifying female pelvic cancer survivors with low levels of physical activity after radiotherapy: women with fecal and urinary leakage need additional support. *Support Care Cancer* 2020;28:2669–81.
- 37 Åkeflo L, Elmerstig E, Dunberger G, *et al.* Sexual health and wellbeing after pelvic radiotherapy among women with and without a reported history of sexual abuse: important issues in cancer survivorship care. *Support Care Cancer* 2021. doi:10.1007/s00520-021-06263-0. [Epub ahead of print: 18 May 2021].
- 38 Schofield P, Juraskova I, Bergin R, *et al.* A nurse- and peer-led support program to assist women in gynaecological oncology receiving curative radiotherapy, the Pentagon study (peer and nurse support trial to assist women in gynaecological oncology): study protocol for a randomised controlled trial. *Trials* 2013;14:39.
- 39 Jefford M, Aranda S, Gough K, *et al.* Evaluating a nurse-led survivorship care package (SurvivorCare) for bowel cancer survivors: study protocol for a randomized controlled trial. *Trials* 2013;14:260.
- 40 Hofsjo A, Bohm-Starke N, Blomgren B. Radiotherapy-Induced vaginal fibrosis in cervical cancer survivors. *Acta oncologica* (Stockholm, Sweden) 2017;1–9.
- 41 Westman B, Ullgren H, Olofsson A, *et al.* Patient-Reported perceptions of care after the introduction of a new advanced cancer nursing role in Sweden. *Eur J Oncol Nurs* 2019;41:41–8.
- 42 Andersen BL, Karlsson JA, Anderson B, *et al.* Anxiety and cancer treatment: response to stressful radiotherapy. *Health Psychol* 1984;3:535–51.
- 43 Oberoi D, White VM, Seymour JF, *et al.* The influence of unmet supportive care needs on anxiety and depression during cancer treatment and beyond: a longitudinal study of survivors of haematological cancers. *Support Care Cancer* 2017;25:3447–56.
- 44 Stubblefield MD. The underutilization of rehabilitation to treat physical impairments in breast cancer survivors. *Pm R* 2017;9:S317–23.
- 45 Penalba V, Deshields TL, Klinkenberg D, DJSCiC K. Gaps in communication between cancer patients and healthcare providers: symptom distress and patients' intentions to disclose. *Support Care Cancer* 2019;27:2039–47.
- 46 Rasmusson E-M, Plantin L, Elmerstig E. 'Did they think I would understand all that on my own?' A questionnaire study about sexuality with Swedish cancer patients. *Eur J Cancer Care* 2013;22:361–9.
- 47 Bakker RM, Mens JWM, de Groot HE, *et al.* A nurse-led sexual rehabilitation intervention after radiotherapy for gynecological cancer. *Support Care Cancer* 2017;25:729–37.
- 48 WHO. Sexual health and sexuality. *Definition* 2006 http://www.who.int/reproductivehealth/topics/sexual_health/sh_definitions/en/
- 49 Chapman CH, Heath G, Fairchild P, *et al.* Gynecologic radiation oncology patients report unmet needs regarding sexual health communication with providers. *J Cancer Res Clin Oncol* 2019;145:495–502.
- 50 Hordern AJ, Street AF. Communicating about patient sexuality and intimacy after cancer: mismatched expectations and unmet needs. *Med J Aust* 2007;186:224–7.
- 51 Concato J, Shah N, Horwitz RJ. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med Overseas Ed* 2000;342:1887–92.
- 52 Althubaiti A. Information bias in health research: definition, pitfalls, and adjustment methods. *J Multidiscip Healthc* 2016;9:211–7.
- 53 Steineck G, Bergmark K, Henningssohn L, *et al.* Symptom documentation in cancer survivors as a basis for therapy modifications. *Acta Oncol* 2002;41:244–52.
- 54 Steineck G, Hunt H, Adolfsson J. A hierarchical step-model for causation of bias-evaluating cancer treatment with epidemiological methods. *Acta Oncol* 2006;45:421–9.