BMJ Open Multimorbidity in Latin America and the Caribbean: a systematic review and meta-analysis

Alessandra M Huaquía-Díaz,¹ Tarick S Chalán-Dávila,¹ Rodrigo M Carrillo-Larco,^{2,3} Antonio Bernabe-Ortiz ^{1,3,4}

To cite: Huaquía-Díaz AM, Chalán-Dávila TS, Carrillo-Larco RM, *et al.* Multimorbidity in Latin America and the Caribbean: a systematic review and meta-analysis. *BMJ Open* 2021;**11**:e050409. doi:10.1136/bmjopen-2021-050409

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-050409>).

Received 20 February 2021

Accepted 12 July 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

¹Universidad Científica del Sur, Lima, Peru

²Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK

³CRONICAS Centre of Excellence in Chronic Diseases, Universidad Peruana Cayetano Heredia, Lima, Peru

⁴School of Public Health and Administration, Universidad Peruana Cayetano Heredia, Lima, Peru

Correspondence to

Dr Antonio Bernabe-Ortiz; antonio.bernabe@upch.pe

ABSTRACT

Objective To estimate the pooled prevalence of multimorbidity (≥ 2 non-communicable diseases in the same individual) among adults of the general population of Latin American and the Caribbean (LAC).

Design Systematic review and meta-analysis.

Data sources MEDLINE, Embase, Global Health, Scopus and LILACS up to 1 July 2020.

Eligibility criteria for selecting studies The outcome was the prevalence of multimorbidity. Reports were selected whether they enrolled adult individuals (age ≥ 18 years) from the general population.

Data extraction and synthesis Reviewers extracted relevant data and assessed risk of bias independently. A random-effects meta-analysis was conducted to report pooled prevalence estimates of multimorbidity; pooled estimates by pre-specified subgroups (eg, national studies) were also pursued.

Results From 5830 results, we selected 28 reports, mostly from Brazil and 16 were based on a nationally representative sample. From the 28 selected reports, 26 were further included in the meta-analysis revealing a pooled multimorbidity prevalence of 43% (95% CI: 35% to 51%; I^2 : 99.9%). When only reports with a nationally representative sample were combined, the pooled prevalence was 37% (95% CI: 27% to 47%; I^2 : 99.9%). When the ascertainment of multimorbidity was based on self-reports alone, the pooled prevalence was 40% (95% CI: 31% to 48%; I^2 : 99.9%); this raised to 52% (95% CI: 33% to 70%; I^2 : 99.9%) for reports including self-reported and objective diagnosis.

Conclusions Our results complement and advance those from global efforts by incorporating much more reports from LAC. We revealed a larger presence of multimorbidity in LAC than previously reported.

PROSPERO registration number CRD42020196177.

INTRODUCTION

The Academy of Medical Sciences defines multimorbidity as ‘the existence of two or more medical chronic conditions in a single individual’.¹ Subjects with multimorbidity tend to increase healthcare utilisation and costs of primary and secondary care services²; also, multimorbidity has a subsequent impact on quality of life.³

Strengths and limitations of this study

- Most of the analysed data came from Brazil preventing region representativity.
- High heterogeneity was present in the analysis, mainly due to different conditions included to define multimorbidity.
- Participant's age seems to explain high heterogeneity as age range in studies is very wide.
- A bias due to self-reporting of conditions may underestimate the real burden of multimorbidity in Latin American and the Caribbean region.

In high-income countries, multimorbidity rates are heterogeneous but seem depend on individual's age. Thus, in a cross-sectional study using the data set of medical practices in Scotland, the prevalence of multimorbidity was 23% using a list of 40 conditions, and was present mainly in older people.⁴ Moreover, multimorbidity seems to be increasing in low-income and middle-income countries (LMIC), where data are yet scarce.⁵ The increase of life expectancy in Latin American and the Caribbean (LAC) has been associated with greater incidence of non-communicable conditions,⁶ with the consequent emergence of multimorbidity.

Multimorbidity prevalence has been explored and summarised in some systematic reviews around the world^{7 8} and results ranged from 5% to 98%; nevertheless, their results were mainly informed by data from high-income countries. In a relatively recent systematic review,⁹ 31 LMIC were included with a prevalence of multimorbidity of 30%, compared with an estimate of 38% in high-income countries. However, only nine studies from the LAC region were included. Moreover, pooled estimates by region were not elucidated, preventing to have appropriate indicators of the burden of multimorbidity in this region. A more recent systematic review has reported a pooled prevalence

of multimorbidity in LMIC between 3% and 90%, with almost 80% of the studies being from Brazil, China, South Africa, India, Mexico and Iran.¹⁰

The lack of evidence about multimorbidity may have important consequences for research, public health and clinical management in LAC region. For example, multimorbidity was not appropriately defined up to 2018; in addition, whether estimates depend on sex or setting characteristics (ie, rural vs urban areas) should be also studied. Moreover, the need of surveillance systems to assess multimorbidity may be elucidated as these estimates have not been estimated in LAC region. Thus, from the public health perspective may not be easy to take appropriate decisions or implement adequate strategies to tackle the problem of multimorbidity.

As a result, we aimed at providing robust evidence about multimorbidity prevalence estimates in LAC region through a systematic review and meta-analysis of population-based surveys. These results evidence may help to guide interventions and policies so that they can focus on the most pressing frequent multimorbidity phenotypes in LAC.

METHODS

Protocol

This systematic review was registered in PROSPERO. We aimed to identify the population-based prevalence of multimorbidity in LAC, and to study whether this prevalence varies by multimorbidity definitions, sex and urban–rural settings.

Eligibility criteria

Reports were selected whether they enrolled adult individuals (age ≥ 18 years) from the general population. We focused on LAC populations; therefore, we excluded studies with LAC individuals in countries outside the LAC region, and studies with only foreign subjects in LAC nations. Population-based studies were defined as those following a random sampling approach, and such sample was taken from the general population. On the contrary, studies addressing specific populations (eg, pregnant women), those with individuals with specific conditions (eg, people with hypertension) or subjects with specific risk factors (eg, obese or alcohol disorders) were excluded.

The outcome of interest was the prevalence of multimorbidity, defined as the existence of ≥ 2 chronic conditions in the same person.¹ Other different definitions of multimorbidity were considered in this review (eg, ≥ 3 , ≥ 4 or ≥ 5 conditions) as the current definition (≥ 2 conditions) is relatively recent. In addition, the presence of chronic conditions could have been measured, self-reported or a combination of these approaches.

Information sources

The search was conducted on 10 January 2020 and then updated on 1 July 2020. We used Ovid search engine,

comprising MEDLINE, Embase and Global Health databases; and in parallel, we also searched Scopus and LILACS. In all of these, searching was carried out without time or language restriction. The search strategy and terms used is detailed in online supplemental tables 1–3.

Study selection

Results from each search engine were downloaded and saved in EndNote where duplicates were removed. After that, information was transferred to Rayyan, an open access online tool for systematic reviews.¹¹ Titles and abstracts were reviewed by two researchers in an independent way, and disagreements were solved by a third party. After this screening phase, selected reports were downloaded and independently studied in detail by two researchers, and similarly, discrepancies were solved by a third party. Finally, selected studies were examined again to check for data duplication, that is, different reports that used the same data (eg, multiple reports based on the same underlying data). In this case, the paper with more information or the one with the largest sample size was included in the review and meta-analysis.

Data collection

An extraction template form was built by the authors and tested with a random sample of selected studies. After starting data collection, the form was not further modified. This form included study characteristics: study design, country, if it was a nationally representative sample, sample size, year of data collection, age range, age mean, proportion of women and if it was urban, rural or both. The extraction form also collated the definition of multimorbidity used, self-reported or a combination of self-reported and measured, the number and a list of chronic conditions studied, and the prevalence of multimorbidity (overall, by sex, and by rural or urban settings).

Risk of bias of individual studies

Risk of bias of selected studies was evaluated using the Newcastle-Ottawa Quality Assessment Scale adapted for cross-sectional studies as in a previous report.¹² This tool is focused on selection process (representativeness, sample size and non-respondents), and the assessment of the outcome (independent blind assessment, self-report or not description). The items of this scale were implemented in an Excel spreadsheet and assessed independently by two reviewers; discrepancies were solved by a third party.

Summary measures

Our systematic review followed the Preferred Reporting Items for Systematic Review and Meta-Analysis toolkit (see checklist in online supplemental table 4). We presented a qualitative and quantitative summary. The qualitative summary described the characteristics of the study (as listed above), whereas the quantitative summary explored pooled prevalence estimates.

Statistical analyses were performed using Stata V.16 for Windows (StataCorp). The ‘metaprop’ command attains

a pooled estimate as a weighted average, by fitting a logistic-normal random-effect model without covariates, but random intercepts.¹³ After that, the pooled estimate was calculated using the Freeman-Tukey arcsine transformation as suggested in literature.¹⁴

Because the selected studies were different in nature, scope (eg, national surveys vs community/subnational studies, or urban vs rural settings) and sample size, we conducted random-effects meta-analysis for comparing estimates in specific subgroups (eg, national studies only). In addition, stratified meta-analysis (eg, by sex, and by urban/rural settings) was pursued. Sensitivity analyses were also conducted that focused on age by including studies with individuals aged 50+ and 60+ years, and also with studies whose data was collected from 2010 and onwards, as well as 2015 and onwards. Besides, as many of the studies were conducted in Brazil, a comparison of pooled estimated between Brazil and other countries together was also carried out.

Finally, meta-regression was also conducted as a high level of heterogeneity was expected. Meta-regression command in Stata investigates if between-study heterogeneity can be explained by one or more of the variables included in the review.

Patient and public involvement

Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of our research.

RESULTS

Selection process

The search strategy yielded 5830 titles and abstracts after removing duplicates (see figure 1); of these 66 were studied in detail and 28 reports met the inclusion criteria.

Figure 1 Flowchart of studies included in the systematic review.

For quantitative analyses, 26 reports were included (2 were excluded as the definition of multimorbidity was ≥ 5 chronic conditions), representing information for 12 LAC countries.

General characteristics of selected studies

As shown in table 1, Brazil contributes with a total of 20 data points^{15–34}; Mexico with 4 data points^{23 35–37}; Peru with 3 data points^{36 38 39}; Colombia with 2 data points^{23 40}; and Argentina,⁴¹ Cuba,³⁶ Dominican Republic,³⁶ El Salvador,²³ Jamaica,²³ Panama,²³ Puerto Rico³⁶ and Venezuela³⁶ with 1 data point each. On the other hand, only 16 studies analysed a nationally representative sample (table 1).

Data from 37 country-level points were used to calculate pooled estimates using a definition of multimorbidity of ≥ 2 chronic conditions. Only one country (Brazil) used an additional definition of multimorbidity (ie, ≥ 3 chronic conditions) in eight different reports. Fourteen reports had information to calculate estimates for urban settings,^{15 17–19 22 25–27 29 33 35 38 39 42} whereas seven had information from rural areas.^{18 25 27 35 38 39 42}

Finally, the number of conditions evaluated to define multimorbidity ranged from 5 to 29, with a mean of 12.3 (SD: 5.7) conditions (online supplemental table 5). Hypertension and type 2 diabetes were the only conditions included in all the definitions of multimorbidity.

Synthesis of results

The meta-analysis included 26 studies with information with a pooled estimate of multimorbidity defined as ≥ 2 chronic conditions of 43% (95% CI: 35% to 51%; I^2 : 99.9%; see figure 2); whereas the pooled estimate for multimorbidity using ≥ 3 chronic conditions was 40% (95% CI: 22% to 57%; I^2 : 99.9%).

As many of the studies were from Brazil, the pooled estimate of multimorbidity (≥ 2 chronic conditions) for this country was 50% (95% CI: 37% to 63%; I^2 : 99.9%), whereas the pooled estimate for other countries together was 35% (95% CI: 26% to 43%; I^2 : 99.7%). Nationally representative samples had a pooled estimate of multimorbidity of 37% (95% CI: 27% to 47%; I^2 : 99.9%), while this estimate was 48% (95% CI: 35% to 61%; I^2 : 99.9%) for non-nationally representative samples. Similarly, when multimorbidity was assessed as self-reported, pooled prevalence was 40% (95% CI: 31% to 48%; I^2 : 99.9%), but the prevalence was 52% (95% CI: 33% to 70%; I^2 : 99.9%) for those which have objectively measured and self-reported chronic conditions.

On the other hand, when analysis was conducted using studies which data was collected from 2010 and onwards, pooled prevalence of multimorbidity was 48% (95% CI: 34% to 61%; I^2 : 99.9%), whereas this estimate was 44% (95% CI: 24% to 65%, I^2 : 99.8%) for studies from 2015 and onwards. This approach was used because time can have an impact on estimations due to the health transition in LMIC. Similarly, when assessing only studies including subjects of 50 years and over, the pooled prevalence of multimorbidity was 62% (95% CI: 51% to 73%;

Table 1 Characteristics of the studies included in the systematic review

First author	Country	Study design	Data collected in	Sample size	Definition of multimorbidity	Age range (years)	Age mean (years)	% women	% urban
Andrade ¹⁷	Brazil	Cross-sectional	1994–1995	1464	≥2 (self-reported)	18+	–	57.4	100
Boing ¹⁹	Brazil	Cross-sectional	2009–2010	1720	≥2 (self-reported)	20–59	–	55.6	100
De Souza Santos Machado ²²	Brazil	Cross-sectional	2005	377	≥2 (self-reported)	40–65	–	100	100
Aguiar ¹⁵	Brazil	Cross-sectional	2011	622	≥2 (self-reported)	50+	64.1	100	100
Nunes ²⁹	Brazil	Cross-sectional	2008	1593	≥2 and ≥3 (self-reported)	60+	–	62.8	100
Agrawal ³⁵	Mexico	Cross-sectional	2007–2010	2725	≥2 (self-reported)	18+	63.1	61.8	73.7
Nunes ²⁶	Brazil	Cross-sectional	2012	2927	≥2 and ≥3 (self-reported)	20+	45.7	58.9	100
Valadares ³³	Brazil	Cross-sectional	2012–2013	736	≥2 (self-reported)	45–60	52.5	100	100
Bustos-Vazquez ³⁷	Mexico	Cross-sectional	2012	7967	≥2 (self-reported)	60+	69.3	53.4	–
Cavalcanti ⁴²	Brazil	Cross-sectional	2010–2011	676	≥2 (self-reported)	60+	70.0	54.6	69.4
Nunes ²⁷	Brazil	Cross-sectional	2013	60 202	≥2 and ≥3 (self-reported)	18+	43.7	55.1	86.5
Nunes ²⁸	Brazil	Cross-sectional	2008	1593	≥2 and ≥3 (self-reported)	60+	–	62.8	–
Olivares ⁴¹	Argentina	Cross-sectional	2014–2015	1044	≥2 (self-reported)	18+	43.0	65	–
Taype-Rondan ³⁹	Peru	Longitudinal	2013–2014	2433	≥2 (self-reported and measured)	35+	57.2	51.3	51.3
Amaral ¹⁶	Brazil	Cross-sectional	2010	264	≥2 (self-reported)	60+	–	61	100
Araujo ¹⁸	Brazil	Cross-sectional	2015	4001	≥2 and ≥3 (self-reported)	18+	–	52.7	86.8
Camargo-Casas ⁴⁰	Colombia	Cross-sectional	2012	2000	≥2 (self-reported)	60+	71.1	63.4	–
Costa ²⁰	Brazil	Cross-sectional	2014	1451	≥2 and ≥3 (self-reported and measured)	60+	–	63	–
Nunes ²⁵	Brazil	Longitudinal	2015–2016	9412	≥2 and ≥3 (self-reported)	50+	62.9	54	84.7

Continued

Table 1 Continued

First author	Country	Study design	Data collected in	Sample size	Definition of multimorbidity	Age range (years)	Age mean (years)	% women	% urban
Bao ³⁶	Cuba	Longitudinal	2003–2007	2944	≥2	65+	73.9	64	–
	Dominican Republic			2011	(self-reported and measured)		74.5	65	–
	Puerto Rico			2009			75.2	67	–
	Venezuela			1965			71.5	63	–
	Peru			1933			74.2	60	–
	Mexico			2003			73.7	64	–
Macinko ²³	Brazil	Cross-sectional	2013–2014	1486	≥2	18+	–	51.1	–
	Colombia			1485	(self-reported)		–	51.1	–
	El Salvador			1460			–	52.8	–
	Jamaica			1480			–	52.6	–
	Mexico			1492			–	51.4	–
	Panama			1475			–	51.6	–
Miranda ³⁸	Peru	Longitudinal	2010–2012	2890	≥2 (self-reported and measured)	35+	55.2	51	50.9
Petarlj ³¹	Brazil	Cross-sectional	2016–2017	790	≥2 and ≥3 (self-reported)	18–59	–	47.7	–
Tavares ³²	Brazil	Cross-sectional	2012	1691	≥2 (self-reported)	60+	72.5	63.7	100
Wang ³⁴	Brazil	Cross-sectional	2005–2007	2713	≥2 (self-reported)	18–64	–	52.4	–
Montes ²⁴	Brazil	Cross-sectional	2014	1336	≥5 (self-reported)	60+	–	63.1	–
Padilha Pereira ³⁰	Brazil	Cross-sectional	2014	1426	≥5 (self-reported)	60+	–	63	–
Da Silva Almeida ²¹	Brazil	Cross-sectional	2016	850	≥2 (self-reported and measured)	18+	–	61.2	–

Multimorbidity (2+): Pooled prevalence

Figure 2 Forest plot of the pooled prevalence of multimorbidity defined as ≥ 2 chronic conditions.

I^2 : 99.9%), and there was no variation when the pooled prevalence was estimated for subject of 60 years and over: 62% (95% CI: 49% to 75%; I^2 : 99.9%).

In stratified analyses, pooled prevalence of multimorbidity was 38.9% (95% CI: 28.6% to 49.1%; I^2 : 99.7%) for men and 50.5% (95% CI: 38.3% to 62.7%; I^2 : 99.8%) for women. These estimates were 38.1% (95% CI: 26.1% to 50.1%; I^2 : 87.3%) and 24.7% (95% CI: 12.5% to 36.8%; I^2 : 50.8%) for urban and rural dwellers, respectively (see online supplemental figures 1–4).

In meta-regression analysis, the number of chronic conditions defining multimorbidity was strongly associated with heterogeneity (β : 0.02 per additional condition, $p < 0.001$). Similarly, mean age was also strongly associated (β : 0.01 per additional year, $p < 0.001$) and the proportion of women involved in the study (β : 0.75, $p = 0.008$). In addition, setting (urban vs rural) was almost associated with heterogeneity (β : 0.59 compared with rural settings, $p = 0.06$).

Risk of bias

All reviewed studies had low risk of bias (online supplemental table 6). Nevertheless, sample size was not justified in three studies,^{23 36 38} and the outcome was self-reported in most of the studies except in five of them.^{21 36 38 39 41}

DISCUSSION

Summary of evidence

This systematic review provides a comprehensive analysis of the burden of multimorbidity in the LAC region.

Accordingly, the pooled prevalence of multimorbidity was 43%. Nevertheless, there was a high heterogeneity among studies included, and results varied by the characteristics of the study as well as some of population characteristics. Thus, the pooled prevalence was lower in nationally representative compared with subnational samples, whereas estimates were higher when only studies with measured and self-reported chronic conditions were included. In addition, age seems to be an important predictor as the prevalence of multimorbidity among those aged 50 years and over was high compared with the pooled estimate of multimorbidity. Finally, the pooled prevalence of multimorbidity was higher among women compared with men, highlighting the link between sex and multimorbidity and among urban compared with those rural dwellers highlighting perhaps better access to diagnostic care in urban sites.

Limitations of the review

There are some limitations in this review that should be highlighted. First, high heterogeneity is present in almost all the results. This finding may be attributable to age group inclusion criteria since a great proportion of studies enrolled individuals aged ≥ 50 years and the pooled estimate of multimorbidity was high in this group. In addition, the proportion of women enrolled in each individual study may be relevant as pointed out by the meta-regression analyses. Second, the number of chronic conditions as well as the list of them used to define multimorbidity is very dissimilar. Defining specific clusters of multimorbidity is needed to guarantee comparability between studies, but this is not usually reported. Therefore, it is relevant to standardise the definition of multimorbidity and the conditions included in such definition to estimate which clusters of multimorbidity are more frequent and relevant for LAC region. In addition to that, before recent definition of multimorbidity, some reports used other different definitions (≥ 3 or ≥ 5 chronic conditions), which could affect pooled results. Fortunately, analysis was possible to include only those with ≥ 2 chronic conditions. Third, most of the studies included in the review were from Brazil, preventing inferability to the whole region, but also highlighting the need of population-based studies on multimorbidity in other countries of the region. Of note, the pooled prevalence of multimorbidity was higher in Brazil compared with other countries, perhaps because Brazilian researchers have addressed a common definition of multimorbidity, using a list of 12 conditions. Finally, a bias due to self-reporting can affect our results. Therefore, whether multimorbidity is defined by self-report or by more objective measurements may have an impact on prevalence estimates. Awareness of some chronic conditions are usually low and varying; for example, hypertension and diabetes awareness is around 64% and 78%, respectively, in urban areas of Latin America,⁴³ but tend to be lower in rural settings.⁴⁴ Thus, our results may be underestimating the real burden of multimorbidity in the LAC region.

Public health relevance

Global trends suggest that multimorbidity is a public health challenge; thus, understanding the epidemiology of multimorbidity in LAC region may be relevant as this issue has received little attention from researchers, but especially, policymakers. Moreover, much of the response of the health system has been developed based on one specific condition or specific body system, instead of an integral approach whereby multiple conditions are addressed synergistically. Therefore, disperse and heterogeneous information have been available for this systematic review.

Our results highlight the need of implementing a surveillance system focused on multimorbidity. This can be done by including some specific conditions in routine health surveys (Demographic Health Surveys or similar) and other population-based research studies. In addition, these surveys may include the most common clusters of multimorbidity, and those with higher morbidity and mortality,⁴⁵ however, such relevant clusters need to be appropriately defined.

Our results also imply that health systems need to be adapted to face the challenge of multimorbidity which increases healthcare use and costs related to primary and secondary prevention.² This adaptation process includes the appropriate training of human resources as well as improving of health services infrastructure and care delivery. It is also needed to develop guidelines for multimorbidity care as that of National Institute for Health and Care Excellence that includes clinical assessment and adequate management,⁴⁶ but also highlight related issues as polypharmacy and life expectancy. Therefore, a holistic approach may be needed to tackle this global health problem.

CONCLUSIONS

Our systematic review shows that 4 of 10 participants have multimorbidity at the population level in LAC. There is, however, a marked variability, depending on participant's age and the number of chronic conditions assessed, highlighting the need of better designed and standardised studies to inform the landscape of multimorbidity in LAC.

Contributors AMH-D, TSC-D and AB-O conceived the idea of the systematic review and manuscript. AMH-D and TSC-D conducted the review with support of RMC-L and AB-O. RMC-L and AB-O conducted the statistical analysis. AMH-D and TSC-D drafted the first version of the manuscript, with relevant comments and suggestions of RMC-L and AB-O. All the authors approved the version submitted for publication.

Funding RMC-L is supported by a Wellcome Trust International Training Fellowship (214185/Z/18/Z).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The systematic review protocol was reviewed and approved by the Ethical Committee at the Universidad Científica del Sur, Lima, Peru.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement As this is a systematic review and meta-analysis, all data relevant to the study are included in the article or uploaded as supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Antonio Bernabe-Ortiz <http://orcid.org/0000-0002-6834-1376>

REFERENCES

- 1 The Academy of Medical Sciences. *Multimorbidity: a priority for health research*. London, UK: Academy of Medical Sciences, 2018.
- 2 Soley-Bori M, Ashworth M, Bisquera A, *et al*. Impact of multimorbidity on healthcare costs and utilisation: a systematic review of the UK literature. *Br J Gen Pract* 2021;71:e39–46.
- 3 Makovski TT, Schmitz S, Zeegers MP, *et al*. Multimorbidity and quality of life: systematic literature review and meta-analysis. *Ageing Res Rev* 2019;53:100903.
- 4 Barnett K, Mercer SW, Norbury M, *et al*. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380:37–43.
- 5 Banerjee A, Hurst J, Fottrell E, *et al*. Multimorbidity: not just for the West. *Glob Heart* 2020;15:45.
- 6 Bilal U, Hessel P, Perez-Ferrer C, *et al*. Life expectancy and mortality in 363 cities of Latin America. *Nat Med* 2021;27:463–470.
- 7 Violan C, Foguet-Boreu Q, Flores-Mateo G, *et al*. Prevalence, determinants and patterns of multimorbidity in primary care: a systematic review of observational studies. *PLoS One* 2014;9:e102149.
- 8 Pati S, Swain S, Hussain MA, *et al*. Prevalence and outcomes of multimorbidity in South Asia: a systematic review. *BMJ Open* 2015;5:e007235.
- 9 Nguyen H, Manolova G, Daskalopoulou C, *et al*. Prevalence of multimorbidity in community settings: a systematic review and meta-analysis of observational studies. *J Comorb* 2019;9:2235042X1987093.
- 10 Abebe F, Schneider M, Asrat B, *et al*. Multimorbidity of chronic non-communicable diseases in low- and middle-income countries: a scoping review. *J Comorb* 2020;10:2235042X2096191.
- 11 Ouzzani M, Hammady H, Fedorowicz Z, *et al*. Rayyan a web and mobile APP for systematic reviews. *Syst Rev* 2016;5:210.
- 12 Modesti PA, Reboli G, Cappuccio FP, *et al*. Panethnic differences in blood pressure in Europe: a systematic review and meta-analysis. *PLoS One* 2016;11:e0147601.
- 13 Nyaga VN, Arbyn M, Aerts M. Metaprop: a Stata command to perform meta-analysis of binomial data. *Arch Public Health* 2014;72:39.
- 14 Freeman MF, Tukey JW. Transformations related to the angular and the square root. *Ann Math Statist*. 1950;21:607–11.
- 15 Aguiar LB, Baccaro LF, de Souza Santos Machado V, *et al*. Disability and multimorbidity in women older than 50 years: a population-based household survey. *Menopause* 2015;22:660–6.
- 16 Amaral TLM, Amaral CdeA, Lima NSde, Cda A, NSd L, *et al*. Multimorbidity, depression and quality of life among elderly people assisted in the family health strategy in Senador Guimard, ACRE, Brazil. *Cien Saude Colet* 2018;23:3077–84.
- 17 Andrade LH, Benseñor IM, Viana MC, *et al*. Clustering of psychiatric and somatic illnesses in the general population: multimorbidity and socioeconomic correlates. *Braz J Med Biol Res* 2010;43:483–91.
- 18 Araujo MEA, Silva MT, Galvao TF, *et al*. Prevalence and patterns of multimorbidity in Amazon region of Brazil and associated determinants: a cross-sectional study. *BMJ Open* 2018;8:e023398.
- 19 Boing AF, Melo GR, Boing AC, *et al*. Associação entre depressão E doenças crônicas: um estudo populacional. *Rev Saúde Pública* 2012;46:617–23.

- 20 CdS C, Flores TR, Wendt A. Inequalities in multimorbidity among elderly: a population-based study in a City in southern Brazil. *Cad Saude Pública* 2018;34:e00040718.
- 21 da Silva Almeida IL, dos Santos SR, Morbeck de Queiroz B. Lifestyle, morbidity and multimorbidity in adult Quilombolas. *ABCS health sci* 2020;45:1325.
- 22 de Souza Santos Machado V, Valadares ALR, da Costa-Paiva LS, et al. Multimorbidity and associated factors in Brazilian women aged 40 to 65 years: a population-based study. *Menopause* 2020;19:569–75.
- 23 Macinko J, Andrade FCD, Nunes BP, et al. Primary care and multimorbidity in six Latin American and Caribbean countries. *Rev Panam Salud Publica* 2019;43:e8–e.
- 24 Montes MC, Bortolotto CC, Tomasi E, et al. Strength and multimorbidity among community-dwelling elderly from southern Brazil. *Nutrition* 2020;71:110636.
- 25 Nunes BP, Batista SRR, FBd A. Multimorbidity: the Brazilian longitudinal study of aging (ELSI-Brazil). *Rev Saude Publica* 2018;2.
- 26 Nunes BP, Camargo-Figuera FA, Guttier M, et al. Multimorbidity in adults from a southern Brazilian City: occurrence and patterns. *Int J Public Health* 2016;61:1013–20.
- 27 Nunes BP, Chiavegatto Filho ADP, Pati S, et al. Contextual and individual inequalities of multimorbidity in Brazilian adults: a cross-sectional national-based study. *BMJ Open* 2017;7:e015885.
- 28 Nunes BP, Soares MU, Wachs LS, et al. Hospitalization in older adults: association with multimorbidity, primary health care and private health plan. *Rev Saude Publica* 2017;51:43.
- 29 Nunes BP, Thumé E, Facchini LA. Multimorbidity in older adults: magnitude and challenges for the Brazilian health system. *BMC Public Health* 2015;15:1172.
- 30 Pereira BP, Bortolotto CC, Tomasi E, Padilha Pereira B, Cardozo Bortolotto C, et al. Food consumption and multimorbidity among non-institutionalized elderly people in Pelotas, 2014: a cross-sectional study. *Epidemiol Serv Saude* 2020;29:e2019050.
- 31 Petarli GB, Cattafesta M, Sant'Anna MM, et al. Multimorbidity and complex multimorbidity in Brazilian rural workers. *PLoS One* 2019;14:e0225416.
- 32 Tavares DMS, Pelizaro PB, Pegorari MS, et al. Prevalence of self-reported morbidities and associated factors among community-dwelling elderly in Uberaba, Minas Gerais, Brazil. *Cien Saude Colet* 2019;24:3305–13.
- 33 Valadares ALR, Lui-Filho JF, Costa-Paiva L, et al. Middle-aged female sexual dysfunction and multimorbidity: a population-based study. *Menopause* 2016;23:304–10.
- 34 Wang Y-P, Nunes BP, Coêlho BM, et al. Multilevel analysis of the patterns of Physical-Mental multimorbidity in general population of São Paulo metropolitan area, Brazil. *Sci Rep* 2019;9:2390.
- 35 Agrawal S, Agrawal PK. Association between body mass index and prevalence of multimorbidity in Low- and middle-income countries: a cross-sectional study. *Int J Med Public Health* 2016;6:73–83.
- 36 Bao J, Chua K-C, Prina M, et al. Multimorbidity and care dependence in older adults: a longitudinal analysis of findings from the 10/66 study. *BMC Public Health* 2019;19:585.
- 37 Bustos-Vázquez E, Fernández-Niño JA, Astudillo-García CI. Self-Rated health, multimorbidity and depression in Mexican older adults: proposal and evaluation of a simple conceptual model. *Biomedica* 2017;37:92–103.
- 38 Miranda JJ, Bernabe-Ortiz A, Gilman RH, et al. Multimorbidity at sea level and high-altitude urban and rural settings: the CRONICAS cohort study. *J Comorb* 2019;9:2235042X1987529–10.
- 39 Taype-Rondan A, Abbs ES, Lazo-Porras M, et al. Association between chronic conditions and health-related quality of life: differences by level of urbanization in Peru. *Qual Life Res* 2017;26:3439–47.
- 40 Camargo-Casas S, Suarez-Monsalve S, Zepeda MUP, et al. [Multimorbidity, depressive symptoms, and self-reported health in older adults: a secondary analysis of the sabe bogota study]. *Rev Invest Clin* 2018;70:192–7.
- 41 Olivares DEV, Chambi FRV, Chañi EMM, et al. Risk factors for chronic diseases and multimorbidity in a primary care context of central Argentina: a web-based interactive and cross-sectional study. *Int J Environ Res Public Health* 2017;14. doi:10.3390/ijerph14030251. [Epub ahead of print: 02 03 2017].
- 42 Cavalcanti G, Doring M, Portella MR, et al. Multimorbidity associated with polypharmacy and negative self-perception of health. *Rev bras geriatr gerontol* 2017;20:634–42.
- 43 Silva H, Hernandez-Hernandez R, Vinuesa R, et al. Cardiovascular risk awareness, treatment, and control in urban Latin America. *Am J Ther* 2010;17:159–66.
- 44 Lerner AG, Bernabe-Ortiz A, Gilman RH, et al. The "rule of halves" does not apply in Peru: awareness, treatment, and control of hypertension and diabetes in rural, urban, and rural-to-urban migrants. *Crit Pathw Cardiol* 2013;12:53–8.
- 45 Whitty CJM, Watt FM. Map clusters of diseases to tackle multimorbidity. *Nature* 2020;579:494–6.
- 46 National Institute for Health and Care Excellence (NICE). Multimorbidity: clinical assessment and management: NICE guideline [NG56]. NICE, 2016. Available: <https://www.nice.org.uk/guidance/ng56> [Accessed 30 Jan 2021].