BMJ Open Assessing the impact of sports and recreation facility density within school neighbourhoods on Canadian adolescents' substance use behaviours: quasi-experimental evidence from the COMPASS study, 2015–2018

Amanda Doggett ¹, Katelyn M Godin,² Olena Schell,² Suzy L Wong,² Ying Jiang,² Scott T Leatherdale¹

To cite: Doggett A, Godin KM, Schell O, *et al.* Assessing the impact of sports and recreation facility density within school neighbourhoods on Canadian adolescents' substance use behaviours: quasi-experimental evidence from the COMPASS study, 2015–2018. *BMJ Open* 2021;**11**:e046171. doi:10.1136/bmjopen-2020-046171

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-046171>).

Received 21 October 2020
Accepted 22 July 2021

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Public Health Sciences, University of Waterloo, Waterloo, Ontario, Canada
²Public Health Agency of Canada, Ottawa, Ontario, Canada

Correspondence to

Amanda Doggett;
adoggett@uwaterloo.ca

ABSTRACT

Objectives There has been relatively little exploration to date of potential protective effects within school neighbourhoods, such as those conferred by facilities that seek to promote health with respect to substance use and related harms. This study examined how the density of sports and recreation facilities in the school neighbourhood is associated with the likelihood of binge drinking, e-cigarette use, cigarette smoking and cannabis use among Canadian secondary school students.

Design Longitudinal data from the COMPASS study on Canadian youth health behaviours from 2015/2016 to 2017/2018 was linked with school neighbourhood data capturing the number of sports and recreation facilities within a 1500 m radius of schools.

Setting Secondary schools and school neighbourhoods in Alberta, British Columbia, Ontario and Quebec who participated in the COMPASS study.

Participants 16 471 youth who participated in the COMPASS study over three school years (2015/2016–2017/2018).

Primary and secondary outcome measures Binge drinking, e-cigarette use, cigarette use, cannabis use.

Results Logistic regression models using generalised estimating equations identified that greater density of sports and recreation facilities within the school neighbourhood was significantly associated with lower likelihood of binge drinking and e-cigarette use but was not associated with cigarette smoking or cannabis use.

Conclusions This research can help to support evidence-informed school community-based efforts to prevent substance-related harms among youth.

INTRODUCTION

Youth substance use is subject to considerable public health attention within Canada. Though many youth who use substances may do so without experiencing significant or long-lasting harms, research demonstrates that initiation of substance use at an earlier

Strengths and limitations of this study

- This study links multiple high-quality data sources, including youth behaviour data from the COMPASS study, neighbourhood information from the Statistics Canada census, and built environment features from the Desktop Mapping Technology Inc enhanced points of interest database.
- The COMPASS study data are drawn from a convenience sample and are therefore not generalisable.
- The COMPASS data are self-reported and as such may be subject to biases.
- The COMPASS study uses an active information passive-consent protocol, which increases sample size and reduces potential self-selection bias.

age during adolescence has been associated with future problematic substance use behaviours, as well as increased propensity for certain chronic diseases later in life.^{1–4} Substance use during adolescence, particularly high-frequency use, may also have negative impacts on mental well-being,^{5–8} as well as educational outcomes and aspirations.^{8–11} Nationally representative data from the 2018/2019 school year show that 23% of Canadian students in grades 7–12 report engaging in binge drinking, while 18% report cannabis consumption within the past year.¹² These data also suggest that the prevalence of current cigarette smoking among Canadian youth is low (3%); however, 20% of students reported current e-cigarette use, reflecting an all-time high.¹² Though there are a variety of factors that can impact propensity for substance use during adolescence, youth substance use is largely moderated by peer influence and social norms, particularly in

the context of binge drinking.^{13–15} The relatively high rates of youth substance use, coupled with the potential health impacts, illustrate the need to identify population-level opportunities to prevent substance-related harms among Canadian youth.

The school and community environments have an important influence on youth health behaviours, particularly during adolescence when individuals increasingly spend time outside of the home. The school can play an important part in moderating youth substance use through school culture, policies and prevention programmes.^{16–19} This influence extends beyond these aspects of the school itself and to the broader school neighbourhood. For example, previous research indicates that the density of tobacco retailers surrounding a school is positively associated with the likelihood of smoking initiation among students.²⁰ Similarly, e-cigarette retailer density is associated with greater likelihood of e-cigarette use in the USA,²¹ although similar Canadian research found no association.²² Research has also suggested that alcohol retailer density surrounding schools can influence students' perceived availability of substances, which in turn is related to likelihood of future substance use.²³

Current literature focuses on risk factors within the school neighbourhood related to youth substance use (eg, tobacco and alcohol retailer density), with little exploration of potential protective effects, such as those conferred by facilities that seek to promote health. One of the most robust and internationally celebrated primary prevention approaches for youth substance use is the Icelandic Model for Preventing Adolescent Substance Use developed by the Icelandic Centre for Social Research & Analysis (ICSRA), which supports youth health through focusing on leveraging protective factors within the school and community environments.²⁴ ICSRA identified that one key protective factor in Icelandic communities was increased access to physical activity, sports and leisure activities. The researchers leveraged the influence of this protective factor through ongoing, multifaceted school-based and community-based efforts, including free access to sports and leisure facilities for youth and collaborating with schools to encourage student participation in extracurricular activities.²⁵ ICSRA researchers assert that the positive impact of these community-level interventions stems from youths' increased opportunity to develop life skills and form healthy, supportive relationships with adults, which builds social capital within the community.²⁶ Iceland has witnessed substantial reductions in youth substance use over the past few decades as a result of this model, which has since been adopted in other jurisdictions internationally.²⁷

In the Canadian context, many of the population level, built environment-focused public health interventions to date have focused on increasing individuals' physical activity. Examples of these interventions include efforts to improve neighbourhood walkability, neighbourhood attractiveness and access to green spaces or recreation facilities.²⁸ There is a wealth of evidence suggesting

that physical activity can be a protective factor for youth substance use and support positive mental well-being.^{29 30} However, many studies have also identified sport participation as a risk factor for binge drinking among adolescents^{31–34} and a potential risk factor for cannabis use, although results are mixed.^{32 35 36} Research surrounding physical activity/sports and cigarette/e-cigarette use has found inverse associations.^{37–40}

The wide range of contexts for sports and recreation activities (eg, inside vs outside of the school recreation, varsity vs intramural sports) can differentially impact youth substance use.^{41 42} However, in general, engagement in these activities is seen as a positive for youth development.⁴³ Notably, the concept of sports participation being beneficial for youth extends beyond physiological impact, as social and psychological factors also play a substantial role in how these activities can be health promoting.⁴⁴ Moreover, although much of the literature focuses on potential protective effects of physical activity and sports, there is substantial rationale to support that leisure more broadly is health promoting. Leisure plays a large role in how individuals find life purpose and meaningfully engage with life,^{45 46} and leisure activities have been shown to promote youth health and protect against behaviours such as substance use.⁴⁷ As such, availability of community resources that offer spaces for youth to engage in recreation and leisure may promote health in a variety of ways, including being protective against substance use behaviours.

Although many studies have examined the relationships between individual-level physical activity, sports participation and youth substance use, there has been a lack of research surrounding broader environmental factors within the school and community that can impact substance use behaviours, such as the sports and leisure facilities that were part of ICSRA's approach in Iceland. This study aims to fill a gap in the youth substance use and school neighbourhood literature, as most of the Canadian literature examining school neighbourhood characteristics focuses on risk factors rather than protective factors. By using a 3-year linked longitudinal sample of youth in Canada at the individual level, and corresponding built environment data at the school neighbourhood level, this study examined the effect that density of sports and recreation facilities (eg, sports clubs, instructional exercise facilities, skating rinks, swimming pools, etc) in the school community may have on youth substance use. Based on existing research, including evidence from evaluations of the Icelandic Model, we hypothesise that greater density of sports and recreation facilities in the school neighbourhood may be associated with decreased propensity for substance use among youth.

METHODS

COMPASS is a prospective cohort study that collects data longitudinally from secondary school students and the school that they attend in four Canadian provinces: Ontario,

Alberta, British Columbia and Quebec.⁴⁸ The student-level data collected are self-reported via an anonymous paper-based questionnaire administered during class time. The questionnaire includes items on multiple domains, including physical activity, diet and substance use, as well as potential correlates of these behaviours and demographic information. COMPASS uses an active information passive-consent protocol.⁴⁸ A full description of COMPASS methods can be found in print⁴⁸ or online (www.compass.uwaterloo.ca).

Sample

This study uses linked longitudinal COMPASS data collected over three consecutive school years (2015/2016, 2016/2017, 2017/2018), hereout referred to as year 1 (Y1), year 2 (Y2) and year 3 (Y3). To be eligible for inclusion in this study, participants needed to complete the COMPASS student questionnaire at baseline (Y1) as grade 9, 10, or secondary I or II students (secondary I and II pertains to Quebec, where the school system structure differs from other provinces; this is equivalent to grades 7 or 8), and again in at least one follow-up year (ie, Y2 and/or Y3). A total of 16471 participants met these criteria. The primary reason for non-participation in the COMPASS study in any year is students being absent or on a spare period during the time of data collection. Student questionnaires were linked across years without the need for identifying information by generating a consistent but unique code from a series of questions on the front page of the questionnaire. The full procedure for linking participants' data over time is described elsewhere.⁴⁹ Secondary data on the types of facilities surrounding the schools were derived from Desktop Mapping Technology Inc (DMTI)'s enhanced points of interest (EPOI) database, which categorises types of area and facilities using the Standard Industrial Classification (SIC) codes system.⁵⁰ EPOI data from 2017 were linked with COMPASS data by school postal code in order to obtain the number of facilities within the neighbourhood of participating schools.⁴⁸ Socioeconomic status and urbanicity of the school neighbourhood were obtained from Statistics Canada Census 2016 data and similarly linked to participating COMPASS schools.

Dependent Variables

Four substance use behaviours, binge drinking, e-cigarette use, cigarette smoking and cannabis use, were examined in this study, and each of the substance use variables was dichotomised into a binary outcome. Consistent with existing research measures,^{12 34 51} binge drinking was examined through the question 'In the last 12 months, how often did you have 5 drinks of alcohol or more on one occasion?' where the response included 'I have never done this', 'I did not have 5 or more drinks on one occasion in the last 12 months', 'Less than once a month', 'Once a month', '2–3 times a month', 'Once a week', '2–5 times a week' and 'Daily or almost daily'. Binge drinking was examined in this study as opposed

to alcohol consumption in general because it represents the more high-risk substance use behaviour. E-cigarette use was examined through the question 'On how many of the last 30 days did you use an e-cigarette?' where the response option included 'None', '1 day', '2–3 days', '4–5 days', '6–10 days', '11–20 days', '21–29 days', '30 days (every day)'. Cigarette smoking was examined through the question 'On how many of the last 30 days did you smoke one or more cigarettes?' where the response options included 'None', '1 day', '2–3 days', '4–5 days', '6–10 days', '11–20 days', '21–29 days', '30 days (every day)'. Cannabis use was examined through the question 'In the last 12 months, how often did you use marijuana or cannabis? (a joint, pot, weed, hash)' where the response options included 'I have never used marijuana', 'I have used marijuana but not in the last 12 months', 'Less than once a month', 'Once a month', '2 or 3 times a week', '4–6 times a week' and 'very day'. For comparability, all four substance use measures were dichotomised into current and non-current use, where current use was defined as engaging in the respective behaviour at least once in the past month.

Independent variables

Individual level demographic variables and control variables

The following control variables (with their response options in brackets) were included in all analyses to address potential confounding effects: grade (9, 10, 11, 12, secondary I and II), sex (male, female), ethno-racial identity (select all that apply: white, black, Asian, Aboriginal, Latin American/Hispanic and other). Personal spending money (\$0, \$1–\$5, \$6–\$10, \$11–\$20, \$21–\$40, \$41–\$100, more than \$100, I do not know how much money I get each week) was also included as an important control which may influence the propensity for substance use behaviours.^{52–54} Participants were also asked to separately indicate their moderate and vigorous physical activity per week in hours (0, 1, 2, 3, 4) and minutes (0, 15, 30, 45). Examples of what constitutes each activity were given in the questionnaire alongside the question. The measure of physical activity is an average of moderate to vigorous physical activity (MVPA) per day and was included in all analyses as a covariate to allow for separation of individual physical activity levels from any potential effects of sports and recreation facility density.

School neighbourhood level variables

Neighbourhood sports and Recreation facilities

The number of neighbourhood sports and recreation facilities within a 1500 m radius of secondary schools that participated in the COMPASS study were obtained from DMTI's EPOI. School-based research has previously used a radius between 1000 m and 2000 m to constitute the school neighbourhood^{55–58}—a radius of 1500 m was chosen for this study. The categories of interest for this project based on the SIC⁵⁹ were: Physical Fitness Facilities (SIC:7991), Membership Sports and Recreation Clubs (SIC:7997) and Amusement and

Recreation Services, not elsewhere classified (SIC:7999). Notably, the latter category was selected for inclusion because it encompasses several sports and recreation-focused items of interest not included in the former two, including swimming, gymnastics, and yoga instruction. All three SIC categories were collapsed into one overall indicator of the number of neighbourhood sports and recreation facilities. A full list of the included facilities in each SIC category can be found by referring to the SIC Major Group 79: Amusement and Recreation Services.⁵⁹ Consistent with previous research,^{20 22 23} the term ‘density’ of school neighbourhood sports and recreation facilities is used hereout in reference to the number of sports and recreation facilities within the aforementioned 1500 m radius of a school.

Alcohol and tobacco Retailers

The models created for each of binge drinking and cigarette smoking controlled for the number of alcohol and tobacco retailers in the school neighbourhood, since existing research has demonstrated that a greater number of these retail facilities can be associated with higher rates of substance use among youth.^{20 23} The DMTI EPOI for these facilities were: Liquor Stores (SIC:5921) and Tobacco Stores and Stands (SIC:5993) within a 1500 m radius of participating COMPASS schools. The number of alcohol retailers was included in the binge drinking model, and the number of tobacco retailers was included in the cigarette smoking model. Data related to cannabis and e-cigarettes retailers were not included in the corresponding models for those substance use behaviours, as data are not yet available on these locations in DMTI’s EPOI database.

Urbanicity and median income

The models adjusted for measures of urbanicity and income as potential neighbourhood-level confounders of youth substance use.^{60 61} Urbanicity data were obtained from Statistics Canada census data (2016), using GeoSearch lookup on city name.⁶² Areas were classified into ‘large urban’, ‘medium urban’, ‘small urban’ and ‘rural’ categories based on national standards.⁶³ Median household income data for the areas where schools were located was obtained from the 2016 Statistics Canada Census Profile.⁶⁴

Analysis

Descriptive statistics were used to characterise the student and school-level variables examined. Four logistic regression models were developed: one for each substance use behaviour. All models used the same 3-year linked longitudinal sample of COMPASS youth and adjusted for the same individual-level covariates. Models were categorised into ‘A’ and ‘B’ series. The A series of models focused only on the substance use examined in the outcome, while the B series of models included adjustments for the other substance use behaviours (eg, in the binge drinking model, the other three substance use behaviours

(cannabis use, cigarette smoking and e-cigarette use) were included as predictors).

Generalised estimating equations (GEE)—a semiparametric method that models population average effects—were used for all logistic regression analyses via PROC GENMOD specifying exchangeable working correlation.⁶⁵ All models accounted for the longitudinal nature of the data (within-subject correlation), as well as the clustering of students within schools (between-subject correlation). GEE is the preferred method when the aim of the research is to estimate the overall effect of a particular variable, rather than individual-specific effects.⁶⁶ The GEE approach ignores missing observations, but all available pairs are used in the estimation of the working correlation matrix.⁶⁷ The longitudinal effect of the main variable of interest (density of school neighbourhood sports and recreation facilities) was tested through an interaction term between that variable and time for all models. All analyses were conducted using SAS V.9.4 (SAS Institute).

Patient and public involvement

Youth participating in the COMPASS study were not involved in the study design. Some of the COMPASS student questionnaire measures are based on previously validated questions among youth, and as such in some cases youth may have been involved in the design. COMPASS disseminates results to stakeholders in a variety of ways including school and provincial level reports prepared yearly. More information about the COMPASS study can be found at <https://uwaterloo.ca/compass-system/>.

RESULTS

Table 1 presents school neighbourhood-level characteristics of the sample. The mean density of sports and recreation facilities within a 1500 m radius of schools was 3.01, although this varied considerably between schools and ranged from 0 to 29. At baseline, 46.17% (n=7586) of the sample were in grade 9, 46.22% (n=7008) were in grade

Table 1 Characteristics of school neighbourhoods (2016/2017) in the sample population

	Per cent of schools (N) N=78
Urbanicity	
Rural	8.97 (7)
Small urban	35.90 (28)
Medium urban	14.10 (11)
Large urban	43.59 (34)
Mean (SD)	
Median income	69 014.26 (18 726.60)
Sports and recreation facilities	3.01 (4.64)
Alcohol retailers	1.61 (2.45)
Tobacco retailers	0.48 (1.37)

10, and 11.17% (1835) were in secondary I or II. Of the baseline sample, 52.85% (n=8655) reported their sex as female, and 47.15% (n=7721) reported their sex as male. 73.72% (12090) of the sample reported their ethno-racial identity as white, while 26.28% (4311) reported an ethno-racial identity other than white. **Figure 1** presents the prevalence of each substance use behaviour across each study sample year. Item non-response for the outcome variables was low (In Y1, 49, 282, 119 and 246 students were missing responses for binge drinking, e-cigarette use, cigarette use and cannabis use, respectively).

Results of the logistic regression models are presented in **table 2**. For all models, time (ie, participants increasing in age/grade) was significantly associated with likelihood of engaging in current substance use, with the exception of model 3B, where the likelihood of smoking decreased over time. The main variable of interest (ie, density of sports and recreation facilities) was significantly associated with lower likelihood of binge drinking and e-cigarette use but was not associated with either cigarette smoking or cannabis use. For binge drinking and e-cigarette use, a greater number of sports and recreation facilities within a school neighbourhood decreased likelihood of respective substance use by 3% for each additional facility. The results pertaining to sports and recreation facilities remained consistent between the A and B series models. In the B series models, physical activity levels were significantly associated with an increased likelihood of binge drinking and e-cigarette use but were not associated with cigarette smoking or cannabis use. Interaction terms between time and each of the school

neighbourhood sports and recreation facilities measures were used to test for the presence of longitudinal effects⁶⁸ but are not presented as effect sizes were not large enough to meaningfully interpret. Models which did not include substance use retailers were also tested, but effect size and significance for sports and recreation facilities remained stable compared with the models presented.

DISCUSSION

To our knowledge, this is the first study to examine the potential impact of density of school neighbourhood sports and recreation facilities on youth substance use in Canada. Our findings indicate that the density of these facilities within the school neighbourhood may have a protective effect against youth binge drinking and e-cigarette use but is not associated with youths' cigarette smoking or cannabis use.

Patterns of substance use behaviours

E-cigarette use was the most common substance use behaviour reported by participants, followed by binge drinking and cannabis use, and then cigarette smoking. The prevalence of these different substance use behaviours aligns with expectations based on previous Canadian and US research.^{12 22 69} Likewise, consistent with previous research,^{70 71} this study observed increased likelihood of substance use over time (except for smoking, which decreased over time) in regression models as well as descriptive analyses, which showed increases in substance use prevalence rates between baseline Y1 and follow-up

Figure 1 Per cent of participants who reported each substance use behaviour in the sample population (Y1:2015/2016, Y2:2016/2017, Y3:2017/2018).

Table 2 Results of binary logistic regression models of four youth substance use behaviours (2015/2016–2017/2018)

OR (95% CI) N=16 471						
Variable†	Binge drinking		E-cigarette use		Cigarette smoking	
	Model 1A	Model 1B	Model 2A	Model 2B	Model 3A	Model 3B
Time	1.65 (1.60 to 1.70)***	1.26 (1.21 to 1.31)***	2.25 (2.18 to 2.31)***	2.11 (2.04 to 2.19)***	1.35 (1.29 to 1.41)***	0.79 (0.75 to 0.84)***
School Variables						
Sports and recreation facilities	0.97 (0.96 to 0.99)***	0.97 (0.96 to 0.99)***	0.97 (0.97 to 0.98)***	0.97 (0.97 to 0.98)***	0.98 (0.97 to 1.00)	0.99 (0.97 to 1.00)
Rural	1	1	1	1	1	1
Small urban	0.65 (0.56 to 0.75)***	0.74 (0.64 to 0.87)***	0.65 (0.56 to 0.75)***	0.71 (0.61 to 0.83)***	0.67 (0.56 to 0.80)***	0.83 (0.68 to 1.02)
Medium urban	0.59 (0.50 to 0.70)***	0.65 (0.55 to 0.78)***	0.63 (0.54 to 0.74)***	0.70 (0.59 to 0.82)***	0.58 (0.47 to 0.71)***	0.66 (0.53 to 0.83)***
Large urban	0.42 (0.36 to 0.48)***	0.52 (0.45 to 0.61)***	0.50 (0.43 to 0.57)***	0.66 (0.57 to 0.77)***	0.30 (0.25 to 0.36)***	0.37 (0.30 to 0.46)***
Alcohol retailers	1.02 (0.99 to 1.05)	1.04 (1.01 to 1.07)**				
Tobacco retailers					1.05 (1.00 to 1.10)	
Individual Variables						
Physical activity level (hours MVPA/day)	1.11 (1.09 to 1.13)***	1.11 (1.09 to 1.14)***	1.08 (1.06 to 1.10)***	1.06 (1.04 to 1.08)***	1.08 (1.05 to 1.11)***	1.03 (1.00 to 1.07)
Binge drinking				3.85 (3.59 to 4.13)***		2.77 (2.51 to 3.07)***
E-cigarette use		3.89 (3.62 to 4.17)***				
Cigarette smoking		2.60 (2.35 to 2.87)***		3.66 (3.30 to 4.07)***		4.17 (3.74 to 4.64)***
Cannabis use			3.32 (3.05 to 3.60)***	3.71 (3.42 to 4.03)***		5.99 (5.41 to 6.64)***

*p<0.05, **p<0.01, ***p<0.001.

†In addition to all variables shown in the table, all models also controlled for time-invariant covariates (measured at baseline): neighbourhood income, sex, ethno-racial identity, grade at baseline, as well as spending money as a time-varying covariate. ORs are rounded to two decimals.

MVPA, moderate to vigorous physical activity.

by Y3. The increase in e-cigarette use between Y1 and follow-up Y3 is particularly drastic compared with the other substances and is consistent with previous research having identified spikes in youth e-cigarette use in recent years.^{12 72}

Density of school neighbourhood sports and recreation facilities

Greater density of school neighbourhood sports and recreation facilities was significantly associated with a decreased likelihood of binge drinking and e-cigarette use among this sample of adolescents. The confidence that these findings do directly reflect the effects of density of these facilities, as opposed to some proxy of larger effects (eg, neighbourhood urbanicity, individual physical activity, etc) is strengthened due to the inclusion of several potential school-level and individual-level confounders in the models.

Previous Canadian research identifies that a key aspect of community sports and recreation facilities that promotes health is their ability to foster social connection and support.²⁸ Sports and recreation facilities can provide youth with opportunities for social connections through a variety of mechanisms, including sports and club participation, as well as volunteer and work opportunities. This type of social engagement in the community is a part of the foundation for the Iceland Prevention Model.²⁴ Research has demonstrated that greater accessibility to sports and leisure for youth in Iceland was protective against substance use behaviours through greater community interaction and strengthened social capital.^{25 73}

However, the density of school neighbourhood sports and recreation facilities did not produce a protective effect against all four substance use behaviours examined. This disparity may be partially explained by differences in the social contexts in which substances are typically used by adolescents (eg, using substances alone vs with peers, motivations for using substances, etc).¹⁵ Among youth, it is known that substance use behaviours are strongly influenced by social norms.^{74 75} As discussed below, the unique social contexts of different substance use behaviours may have a bearing on the likelihood for sports and recreation activities to impact substance use behaviours.

Binge drinking and e-cigarette use

Research indicates that, among young people, binge drinking most often occurs in social contexts (eg, parties).^{14 76} It may be reasonable to suggest that accessing school neighbourhood sports and recreation facilities can reduce some social motivators for binge drinking; some of the students' need for social engagement could be met through involvement with these facilities, and they may therefore be less inclined to binge drink for social reasons. Moreover, boredom is a known predictor of binge drinking,^{77 78} and a neighbourhood with a higher density of sports and recreation

facilities offers more opportunities for extracurricular activities in which youth can participate, and therefore, may decrease the likelihood that youth will engage in binge drinking out of boredom. While the overall effect of sports and recreation facility density was protective against binge drinking in this study, aforementioned research has highlighted that team sports specifically can be a risk factor for binge drinking, so there are likely unique social contexts depending on the type of sports in which youth participate.^{31 36} The complex nature of the associations between sports, recreation and binge drinking indicate that prevention efforts likely require a combination of population level and targeted approaches.

Similar to binge drinking, youth primarily report social reasons for engaging in e-cigarette use, including the need to be perceived as 'cool' by their peers.^{79 80} The large presence of online marketing of novel e-cigarette products that target youth has also likely contributed to these shifts in perception and normalisation of e-cigarette use.^{81 82} This study demonstrated protective effects of school neighbourhood sports and recreation facilities density against youth e-cigarette use, as well as binge drinking. These findings may indicate that greater access to opportunities for social connection may displace some social motivation to engage in these substance use behaviours.

Tobacco and cannabis use

In contrast, there is evidence that cigarette smoking and cannabis use may be less socially motivated behaviours, compared with binge drinking and e-cigarette use. While smoking has inherent ties to social factors, such as peer group and family opinions on smoking,^{83 84} opinions towards smoking have shifted and continue to shift in Canada to be increasingly unfavourable.⁸⁵ This can be observed even over the short time period of the present study, as smoking was the only substance use behaviour that showed decreased likelihood over time. Research has found that Canadian youth are altering their perceptions away from smoking being a 'cool' behaviour and instead considering it a negative and less socially acceptable behaviour, in part due to better knowledge of associated adverse health outcomes.⁸⁵ These changing social norms among Canadian youth may have contributed to steadily decreasing smoking rates in recent years⁸⁶ and may explain why the density of sports and recreation facilities showed no protective effects for this substance use behaviour.

The social context of cannabis use among Canadian youth appears to be more complex; while many youth report that a main motivator for cannabis use is to 'fit-in', there are many reported motivators that are not necessarily social in nature, such as improved physical or mental health, sleep or concentration.⁸⁷ Particularly relevant to the present study, Canadian youth have reported that being busy with extracurricular activities is not necessary an impediment to cannabis

use, and instead sometimes can be used as a ‘cover’ to hide use from parents.⁸⁷ Therefore, while existing research suggests that youth have some social motivators for youth cannabis use, it remains distinct from other substance use behaviours. This may partly explain the findings presented here that school neighbourhood sports and recreation facilities had no effect on cannabis use while showing significant protective effects for other substances. Notably, the present study, as well as the referenced literature, was conducted prior to non-medical cannabis legalisation for adults. Although cannabis remains an illegal substance for youth, it is unclear what effects legalisation may have on youths’ perceptions and norms around cannabis.

Implications

The present findings suggest a protective effect of school neighbourhood sports and recreation facilities, even after controlling for potential neighbourhood risk factors, including the presence of alcohol and tobacco retailers. This lends support for initiatives similar to the Iceland Prevention Model’s comprehensive approach to addressing youth substance use, which focused on leveraging community-level resources and engagement.²⁷ The relatively small effect sizes in this study are to be expected and reflect that neighbourhood sports and recreation facilities represent only a single aspect of the built environment. The built environment components of the school neighbourhood consist of many overarching elements that may influence health behaviours, including the density of sports and recreation facilities examined in the present study, as well as other types of facilities, walkability, green space, transit and many others. The number and various types of factors that influence youth health and well-being underscore the importance of multipronged, comprehensive efforts to prevent substance-related harms and that there is no ‘silver bullet’ for addressing this complex public health problem. The present study, supported by the existing literature, illustrates the importance of considering all aspects of the school community for youth health. Understanding the impact of the school neighbourhood sports and recreation facilities also illustrates potential opportunities to leverage partnerships and services between schools and community organisations towards the shared goal of supporting youth health.

Limitations

There are several limitations to note with respect to the present study. First, this study only measured associations between the built environment and individual health behaviours, and as such, it was not possible to fully understand the mechanism behind these associations. Future research should consider measuring the social context of substance use alongside frequency in order to further understand the potential connection between school neighbourhood sports and recreation facilities, social connection and substance use. Second,

this study relied on existing DMTI database categorisations as representations of the built environment and therefore were imperfect representations of sports and recreation centres in which youth might engage. As such, our study could only measure density, rather than accessibility, of the school neighbourhood sports and recreation facilities; unmeasured characteristics of the individual facilities (eg, cost, activities offered, age restrictions, etc) would affect ability for youth to use these facilities. In particular, cost barriers to participation may disproportionality impact school communities in lower-income areas, and therefore, future research should consider collecting this data and examining the accessibility of these facilities. Moreover, family-related indicators such as parental income or education may be associated with youth substance use but were not collected in this study, and as such, may confound results. Lastly, the COMPASS study uses convenience sampling and as such, the results of this study may not be generalisable outside the study population. This study has many strengths; to our knowledge, this is the first study to examine the impact of school neighbourhood sports and recreation facilities on youth substance use in Canada and is one of few studies in the literature that focus on potential protective factors of the built environment. Furthermore, the longitudinal nature of this study, combined with the large sample size, allowed for robust examination of how aspects of the school neighbourhood impact likelihood of substance use. Lastly, the use of a passive consent protocol was an important feature for a study examining substance use behaviours since it helps to limit response and self-selection biases.^{88 89}

CONCLUSIONS

This study identified a negative association between the density of school neighbourhood sports and recreation facilities and likelihood of youth binge drinking and e-cigarette use, which suggests that these facilities may have the potential to displace some substance use behaviours among youth. These findings lend support to population-level interventions aiming to prevent substance-related harms among youth through built environment features, such as sports and recreation facilities in school neighbourhoods. Potential directions for future research include examining the influence of sports and recreation facility density within school neighbourhoods on youth substance use in other study samples, as well as exploring other features of the built environment that may have protective effects on youth substance use behaviours.

Contributors AD, KMG, OS, SLW and YJ contributed to project conceptualisation methodology plan. AD conducted the formal analysis and the original draft preparation. AD, KMG, OS, SLW, YJ and STL reviewed and edited the manuscript for critical content. SL acquired funding, managed resources and provided supervision for the host study and corresponding data in this manuscript. All authors read and agreed to the published version of the manuscript.

Funding The COMPASS study has been supported by a bridge grant from the CIHR Institute of Nutrition, Metabolism and Diabetes (INMD) through the “Obesity – Interventions to Prevent or Treat” priority funding awards (OOP-110788; awarded to SL), an operating grant from the CIHR Institute of Population and Public Health (IPPH) (MOP-114875; awarded to SL), a CIHR project grant (PJT-148562; awarded to SL), a CIHR bridge grant (PJT-149092; awarded to KP/SL), a CIHR project grant (PJT-159693; awarded to KP), by a research funding arrangement with Health Canada (#1617-HQ-000012; contract awarded to SL), a CIHR-Canadian Centre on Substance Abuse (CCSA) team grant (OF7 B1-PCPEGT 410-10-9633; awarded to SL), and a SickKids Foundation New Investigator Grant, in partnership with CIHR Institute of Human Development, Child and Youth Health (IHDCYH) (Grant No. NI21-1193; awarded to KAP) funds a mixed methods study examining the impact of the COVID-19 pandemic on youth mental health, leveraging COMPASS study data. The COMPASS-Quebec project additionally benefits from funding from the Ministère de la Santé et des Services sociaux of the province of Québec, and the Direction régionale de santé publique du CIUSSS de la Capitale-Nationale. The creation of this manuscript was funded by the Research Affiliate Program from the Public Health Agency of Canada, Applied Research Branch.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The COMPASS Study received ethics approval from the University of Waterloo Office of Research Ethics as well as participating school boards (ORE #: 30118).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. COMPASS data are available for researchers upon successful completion and approval of the COMPASS data usage application (<https://uwaterloo.ca/compass-system/information-researchers>). Technical reports detailing COMPASS study methods are available online (<https://uwaterloo.ca/compass-system/publications#technical>).

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Amanda Doggett <http://orcid.org/0000-0001-8538-432X>

REFERENCES

- Degenhardt L, Coffey C, Carlin JB, *et al.* Outcomes of occasional cannabis use in adolescence: 10-year follow-up study in Victoria, Australia. *Br J Psychiatry* 2010;196:290–5 https://www.cambridge.org/core/product/identifier/S0007125000252203/type/journal_article
- Nelson SE, Van Ryzin MJ, Dishion TJ, Alcohlo DTJ. Alcohol, marijuana, and tobacco use trajectories from age 12 to 24 years: demographic correlates and young adult substance use problems. *Dev Psychopathol* 2015;27:253–77.
- Government of Canada. Preventing problematic substance use in youth [Internet], 2018. Available: <https://www.canada.ca/content/dam/phac-aspc/documents/corporate/publications/chief-public-health-officer-reports-state-public-health-canada/2018-preventing-problematic-substance-use-youth/2018-preventing-problematic-substance-use-youth.pdf>
- Schulte MT, Hser Y-I. Substance use and associated health conditions throughout the lifespan. *Public Health Rev* 2014;35:3 <http://publichealthreviews.biomedcentral.com/articles/10.1007/BF03391702>
- Conway KP, Green VR, Kasza KA, *et al.* Co-Occurrence of tobacco product use, substance use, and mental health problems among youth: findings from wave 1 (2013–2014) of the population assessment of tobacco and health (path) study. *Addict Behav* 2018;76:208–17.
- Marmorstein NR, Iacono WG, Malone SM. Longitudinal associations between depression and substance dependence from adolescence through early adulthood. *Drug Alcohol Depend* 2010;107:154–60.
- Mackie CJ, O’Leary-Barrett M, Al-Khudhairy N, *et al.* Adolescent bullying, cannabis use and emerging psychotic experiences: a longitudinal general population study. *Psychol Med* 2013;43:1033–44 http://www.journals.cambridge.org/abstract_S003329171200205X
- Meier MH, Hill ML, Small PJ, *et al.* Associations of adolescent cannabis use with academic performance and mental health: a longitudinal study of upper middle class youth. *Drug Alcohol Depend* 2015;156:207–12.
- Stiby AI, Hickman M, Munafò MR, *et al.* Adolescent cannabis and tobacco use and educational outcomes at age 16: birth cohort study. *Addiction* 2015;110:658–68.
- Latvala A, Rose RJ, Pulkkinen L, *et al.* Drinking, smoking, and educational achievement: cross-lagged associations from adolescence to adulthood. *Drug Alcohol Depend* 2014;137:106–13.
- Patte KA, Qian W, Leatherdale ST. Marijuana and alcohol use as predictors of academic achievement: a longitudinal analysis among youth in the COMPASS study. *J Sch Health* 2017;87:310–8.
- Government of Canada. Summary of results for the Canadian Student Tobacco, Alcohol and Drugs Survey 2018–19 (CSTAD) [Internet], 2019. Available: <https://www.canada.ca/en/health-canada/services/canadian-student-tobacco-alcohol-drugs-survey/2018-2019-summary.html>
- Ajilore O, Amialchuk A, Egan K. Alcohol consumption by youth: Peers, parents, or prices? *Econ Hum Biol* 2016;23:76–83.
- MacArthur GJ, Jacob N, Pound P, *et al.* Among friends: a qualitative exploration of the role of peers in young people’s alcohol use using Bourdieu’s concepts of habitus, field and capital. *Sociol Health Illn* 2017;39:30–46.
- Jenkins EK, Slemon A, Haines-Saah RJ. Developing harm reduction in the context of youth substance use: insights from a multi-site qualitative analysis of young people’s harm minimization strategies. *Harm Reduct J* 2017;14:1–11.
- Bonell C, Parry W, Wells H, *et al.* The effects of the school environment on student health: a systematic review of multi-level studies. *Health Place* 2013;21:180–91.
- Costello MJE, Leatherdale ST, Ahmed R, *et al.* Co-morbid substance use behaviors among youth: any impact of school environment? *Glob Health Promot* 2012;19:50–9.
- Evans-Whipp T, Beyers JM, Lloyd S, *et al.* A review of school drug policies and their impact on youth substance use. *Health Promot Int* 2004;19:227–34.
- Bisset S, Markham WA, Aveyard P. School culture as an influencing factor on youth substance use. *J Epidemiol Community Health* 2007;61:485–90.
- Chan WC, Leatherdale ST. Tobacco retailer density surrounding schools and youth smoking behaviour: a multi-level analysis. *Tob Induc Dis* 2011;9:9 <http://www.tobaccoinduceddiseases.org/Tobacco-retailer-density-surrounding-schools-and-youth-smoking-behaviour-a-multi,66050,0,2.html>
- Giovenco DP, Casseus M, Duncan DT, *et al.* Association between electronic cigarette marketing near schools and e-cigarette use among youth. *J Adolesc Health* 2016;59:627–34.
- Cole AG, Aleyan S, Leatherdale ST. Exploring the association between e-cigarette retailer proximity and density to schools and youth e-cigarette use. *Prev Med Rep* 2019;15:100912 <https://www.clinicalkey.es/#!/content/playContent/1-s2.0-S2211335519300877?scrollTo=%23h0000443>
- Milam AJ, Johnson SL, Furr-Holden CDM, *et al.* Alcohol outlets and substance use among high Schoolers. *J Community Psychol* 2016;44:819–32.
- Kristjánsson AL, Mann MJ, Sigfusson J, *et al.* Development and guiding principles of the Icelandic model for preventing adolescent substance use. *Health Promot Pract* 2020;21:62–9.
- Kristjánsson AL, James JE, Allegrante JP, *et al.* Adolescent substance use, parental monitoring, and leisure-time activities: 12-year outcomes of primary prevention in Iceland. *Prev Med* 2010;51:168–71 <https://www.sciencedirect.com/science/article/pii/S0091743510001763>
- Sigfúsdóttir ID, Thorlindsson T, Kristjánsson AL, *et al.* Substance use prevention for adolescents: the Icelandic model. *Health Promot Int* 2009;24:16–25 <https://academic.oup.com/heapro/article-abstract/24/1/16/677129>
- Sigfúsdóttir ID, Kristjánsson AL, Gudmundsdóttir ML. Prevención del consumo de sustancias mediante La promoción de la salud en La escuela Y en La comunidad: Enfoque transdisciplinario de Islandia. *Glob Health Promot* 2011;18:23–6 <http://www.sagepub.co.uk/journalsPermissions.nav>
- Public Health Agency of Canada. Designing healthy living [Internet], 2017. Available: https://www.canada.ca/content/dam/phac-aspc/documents/services/publications/chief-public-health-officer-reports-state-public-health-canada/2017-designing-healthy-living/PHAC_CPHO-2017_Report_E.pdf
- Spruit A, Assink M, van Vugt E, *et al.* The effects of physical activity interventions on psychosocial outcomes in adolescents: a meta-analytic review. *Clin Psychol Rev* 2016;45:56–71.

- 30 Simonton AJ, Young CC, Johnson KE. Physical activity interventions to decrease substance use in youth: a review of the literature. *Subst Use Misuse* 2018;53:2052–68 <https://www.tandfonline.com/doi/full/10.1080/10826084.2018.1452338>
- 31 Butler A, Romano I, Patte K, et al. Psychological correlates and binge drinking behaviours among Canadian youth: a cross-sectional analysis of the mental health pilot data from the COMPASS study. *BMJ Open* 2019;9:e028558.
- 32 Lau EY, Riaz NA, Qian W, et al. Protective or risky? the longitudinal association of team sports participation and health-related behaviours in Canadian adolescent girls. *Can J Public Health* 2019;110:616–25.
- 33 Kwan M, Bobko S, Faulkner G, et al. Sport participation and alcohol and illicit drug use in adolescents and young adults: a systematic review of longitudinal studies. *Addict Behav* 2014;39:497–506 http://ac.els-cdn.com/S0306460313003766/1-s2.0-S0306460313003766-main.pdf?_tid=58075f18-52c5-11e7-a80d-00000aacb35d&acdnat=1497639507_3ac94fa3fc7ad2d875aa9669fee39d69
- 34 Modecki KL, Barber BL, Eccles JS. Binge drinking trajectories across adolescence: for early maturing youth, extra-curricular activities are protective. *J Adolesc Health* 2014;54:61–6.
- 35 Boyes R, O'Sullivan DE, Linden B, et al. Gender-specific associations between involvement in team sport culture and Canadian adolescents' substance-use behavior. *SSM Popul Health* 2017;3:663–73.
- 36 Wichstrøm T, Wichstrøm L. Does sports participation during adolescence prevent later alcohol, tobacco and cannabis use? *Addiction* 2009;104:138–49.
- 37 Pokhrel P, Schmid S, Pagano I. Physical activity and use of cigarettes and e-cigarettes among young adults. *Am J Prev Med* 2020;58:580–3.
- 38 Milicic S, Piérard E, DeCicca P, et al. Examining the association between physical activity, sedentary behavior and sport participation with e-cigarette use and smoking status in a large sample of Canadian youth. *Nicotine Tob Res* 2019;21:285–92.
- 39 Larson NI, Story M, Perry CL, et al. Are diet and physical activity patterns related to cigarette smoking in adolescents? findings from project eat. *Prev Chronic Dis* 2007;4:A51.
- 40 Ali MM, Amialchuk A, Heller LR. The influence of physical activity on cigarette smoking among adolescents: evidence from add health. *Nicotine Tob Res* 2015;17:539–45.
- 41 Williams GC, Burns KE, Battista K, et al. High school sport participation and substance use: a cross-sectional analysis of students from the COMPASS study. *Addict Behav Rep* 2020;12:100298.
- 42 Zhou J, Heim D. Sports and spirits: a systematic qualitative review of emergent theories for student-athlete drinking. *Alcohol Alcohol* 2014;49:604–17.
- 43 Guèvremont A, Findlay L, Kohen D. Organized extracurricular activities: are in-school and out-of-school activities associated with different outcomes for Canadian youth? *J Sch Health* 2014;84:317–25.
- 44 Elsborg P, Nielsen G, Klinker CD, et al. Sports-based recreation as a means to address social inequity in health: why, when, where, who, what, and how. *BMC Public Health* 2019;19:1–9.
- 45 Iwasaki Y, Messina ES, Hopper T. The role of leisure in meaning-making and engagement with life. *J Posit Psychol* 2018;13:29–35.
- 46 Hopper TD, Iwasaki Y. Engagement of 'At-Risk' Youth Through Meaningful Leisure. *J Park Recreat Admi* 2017;35:20–33.
- 47 Santini ZI, Meilstrup C, Hinrichsen C, et al. Associations between multiple leisure activities, mental health and substance use among adolescents in Denmark: a nationwide cross-sectional study. *Front Behav Neurosci* 2020;14:1–10.
- 48 Leatherdale ST, Brown KS, Carson V, et al. The COMPASS study: a longitudinal hierarchical research platform for evaluating natural experiments related to changes in school-level programs, policies and built environment resources. *BMC Public Health* 2014;14:331.
- 49 Battista K, Qian W, Bredin C. Student data linkage over multiple years [Internet]. Vol. 6. COMPASS Technical Report Series, 2019. Available: <https://uwaterloo.ca/compass-system/student-data-linkage-over-multiple-years>
- 50 DMTI. DMTI Spatial: Real-time Location Intelligence [Internet]. 2019. Available: <https://www.dmtispatial.com/>
- 51 McCabe SE, Boyd CJ, Evans-Polce RJ, et al. School-Level prevalence and predictors of e-cigarette use in 8th, 10th, and 12th grade U.S. youth: results from a national survey (2015–2016). *Journal of Adolescent Health* 2020;67:531–41.
- 52 Leatherdale ST, Ahmed R. Alcohol, marijuana, and tobacco use among Canadian youth: do we need more multi-substance prevention programming? *J Prim Prev* 2010;31:99–108.
- 53 Zuckermann AME, Gohari MR, de Groh M, et al. Factors associated with cannabis use change in youth: evidence from the COMPASS study. *Addict Behav* 2019;90:158–63.
- 54 Perikleous EP, Steiropoulos P, Paraskakis E, et al. E-cigarette use among adolescents: an overview of the literature and future perspectives. *Front Public Health* 2018;6:1–9.
- 55 Stone M, Larsen K, Faulkner GEJ, et al. Predictors of driving among families living within 2km from school: exploring the role of the built environment. *Transp Policy* 2014;33:8–16.
- 56 Lane NE, Leatherdale ST, Dubin JA, et al. Student and school characteristics associated with use of nicotine replacement therapy: a multilevel analysis among Canadian youth. *Addict Behav* 2012;37:811–6.
- 57 Hobin EP, Leatherdale ST, Manske S, et al. A multilevel examination of gender differences in the association between features of the school environment and physical activity among a sample of grades 9 to 12 students in Ontario, Canada. *BMC Public Health* 2012;12:74.
- 58 Larsen K, To T, Mann R. Does the school neighbourhood play a role in health behaviour? A study on the prevalence of smoking in youths and teenagers. *American Thoracic Society International Conference [Internet]*, 2016.
- 59 United States Department of Labor. SIC division structure | occupational safety and health administration [Internet], 2018. Available: https://www.osha.gov/pls/imis/sic_manual.html
- 60 Results NS, OND U. *The future secondary school students*. 1, 2012.
- 61 Martino SC, Ellickson PL, McCaffrey DF. Developmental trajectories of substance use from early to late adolescence: a comparison of rural and urban youth. *J Stud Alcohol Drugs* 2008;69:430–40.
- 62 Statistics Canada. GeoSearch [Internet], 2017. Available: <https://www12.statcan.gc.ca/census-recensement/2016/geo/geosearch-georecherche/index-eng.cfm>
- 63 Canada S. Population Centre and Rural Area Classification 2016 [Internet]. Government of Canada, Statistics Canada, 2016. Available: <https://www.statcan.gc.ca/eng/subjects/standard/pcrac/2016/introduction/>
- 64 StatsCan. *Census Program [Internet]*. Central Statistics Office Census, 2016: 72–4. <https://www150.statcan.gc.ca/n1/en/type/data?MM=1>
- 65 Wilson JR. *Modeling Binary Correlated Responses using SAS, SPSS and R [Internet]*. Cham: Springer International Publishing, 2015. <http://www.springer.com/series/13402>
- 66 Fitzmaurice GM, Laird NM, Ware JH. *Applied longitudinal analysis [Internet]*. Wiley, 2011. Available: <https://www.wiley.com/en-ca/Applied+Longitudinal+Analysis%2C+2nd+Edition-p-9780470380277>
- 67 SAS. The GENMOD procedure - missing values [Internet], 2019. Available: https://documentation.sas.com/?cdcid=pgmsascdc&cdcVersion=9.4_3.3&docsetId=statug&docsetTarget=statug_genmod_details80.htm&locale=en
- 68 Hoffman L. *Longitudinal analysis*. Taylor & Francis, 2015.
- 69 Miech R, Patrick ME, O'Malley PM, et al. What are kids vaping? results from a national survey of US adolescents. *Tob Control* 2017;26:386–91.
- 70 Leatherdale ST, Burkhalter R. The substance use profile of Canadian youth: exploring the prevalence of alcohol, drug and tobacco use by gender and grade. *Addict Behav* 2012;37:318–22.
- 71 Gardner M, Steinberg L. Peer influence on risk taking, risk preference, and risky decision making in adolescence and adulthood: an experimental study. *Dev Psychol* 2005;41:625–35.
- 72 Aleyan S, Cole A, Qian W, et al. Risky business: a longitudinal study examining cigarette smoking initiation among susceptible and non-susceptible e-cigarette users in Canada. *BMJ Open* 2018;8:21080.
- 73 Schüttoff U, Pawlowski T, Downward P, et al. Sports participation and social capital formation during Adolescence*. *Soc Sci Q* 2018;99:683–98.
- 74 Mahalik JR, Lombardi CM, Sims J, et al. Gender, male-typicality, and social norms predicting adolescent alcohol intoxication and marijuana use. *Soc Sci Med* 2015;143:71–80.
- 75 Bryden A, Roberts B, Petticrew M, et al. A systematic review of the influence of community level social factors on alcohol use. *Health Place* 2013;21:70–85.
- 76 Ander B, Abrahamsson A, Bergnehr D. 'It is ok to be drunk, but not too drunk': party socialising, drinking ideals, and learning trajectories in Swedish adolescent discourse on alcohol use. *J Youth Stud* 2017;20:841–54.
- 77 Biolcati R, Mancini G, Trombini E. Proneness to Boredom and risk behaviors during adolescents' free time. *Psychol Rep* 2018;121:303–23.
- 78 Biolcati R, Passini S, Mancini G. "I cannot stand the boredom." Binge drinking expectancies in adolescence. *Addict Behav Rep* 2016;3:70–6.

- 79 Camenga DR, Fiellin LE, Pendergrass T, *et al.* Adolescents' perceptions of flavored tobacco products, including e-cigarettes: a qualitative study to inform FDA tobacco education efforts through videogames. *Addict Behav* 2018;82:189–94.
- 80 Romijnders KAGJ, van Osch L, de Vries H, *et al.* Perceptions and reasons regarding e-cigarette use among users and non-users: a narrative literature review. *Int J Environ Res Public Health* 2018;15:1190.
- 81 Laestadius LI, Wahl MM, Pokhrel P, *et al.* From apple to Werewolf: a content analysis of marketing for e-liquids on Instagram. *Addict Behav* 2019;91:119–27.
- 82 Hilton S, Weishaar H, Sweeting H, *et al.* E-cigarettes, a safer alternative for teenagers? A UK focus group study of teenagers' views. *BMJ Open* 2016;6:13271.
- 83 Nilsson M, Emmelin M. "Immortal but frightened"-smoking adolescents' perceptions on smoking uptake and prevention. *BMC Public Health* 2010;10 <http://www.biomedcentral.com/1471-2458/10/776>
- 84 Johnston V, Westphal DW, Earnshaw C, *et al.* Starting to smoke: a qualitative study of the experiences of Australian Indigenous youth. *BMC Public Health* 2012;12:963 <http://www.biomedcentral.com/1471-2458/12/963>
- 85 Woodgate RL, Busolo DS. A qualitative study on Canadian youth's perspectives of peers who smoke: an opportunity for health promotion. *BMC Public Health* 2015;15.
- 86 Janz T. Current smoking trends [Internet]. Statistics Canada, 2012. Available: <http://www.statcan.gc.ca/pub/82-624-x/2012001/article/11676-eng.htm%5Cnhttp://www.statcan.gc.ca/pub/82-624-x/2012001/article/11676-eng.pdf>
- 87 Mckiernan A, Fleming K. *Canadian youth perceptions on cannabis. Canadian centre on substance use and addiction.* 67, 2017.
- 88 White VM, Hill DJ, Effendi Y. How does active parental consent influence the findings of drug-use surveys in schools? *Eval Rev* 2004;28:246–60 <http://journals.sagepub.com/doi/10.1177/0193841X03259549>
- 89 Rojas NL, Sheritt L, Harris S, *et al.* The role of parental consent in adolescent substance use research. *J Adolesc Health* 2008;42:192–7 http://ac.els-cdn.com/S1054139X07003229/1-s2.0-S1054139X07003229-main.pdf?_tid=3c346012-6722-11e7-9677-00000aacb35e&acdnat=1499878427_8f0da40aa70aa39468f160a3314cf50d