




# BMJ Open Evaluating the implementation of interdisciplinary patient-centred care intervention for people with multimorbidity in primary care: a qualitative study

Patrice Ngangue <sup>1</sup>, Judith Belle Brown,<sup>2</sup> Catherine Forgues,<sup>1</sup> Mohamed Ali Ag Ahmed,<sup>1</sup> Tu Ngoc Nguyen <sup>3</sup>, Maxime Sasseville <sup>4</sup>, Christine Loignon <sup>1</sup>, Frances Gallagher,<sup>5</sup> Moira Stewart,<sup>2</sup> Martin Fortin<sup>1</sup>

**To cite:** Ngangue P, Brown JB, Forgues C, *et al.* Evaluating the implementation of interdisciplinary patient-centred care intervention for people with multimorbidity in primary care: a qualitative study. *BMJ Open* 2021;**11**:e046914. doi:10.1136/bmjopen-2020-046914

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-046914>).

Received 14 November 2020  
Accepted 02 September 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Patrice Ngangue;  
Patrice.Ngangue@Usherbrooke.ca and  
Dr Martin Fortin;  
martin.fortin@usherbrooke.ca

## ABSTRACT

**Objective** A patient-centred care interdisciplinary pragmatic intervention to support self-management for patients with multimorbidity was implemented in one region of Quebec, Canada. This embedded study aimed to evaluate the process of implementation.

**Design** A descriptive qualitative study was conducted in 2016–2017 using semistructured individual interviews. The Consolidated Framework for Implementation Research (CFIR) was used to guide the data coding, analysis and reporting of the findings.

**Setting** The study took place in seven Family Medicine Groups in one region (Saguenay-Lac-Saint-Jean) of Quebec, Canada.

**Participants** Ten managers (including two family physicians) and 19 healthcare professionals (HCPs), nurses, kinesiologists, nutritionists and a respiratory therapist, were interviewed.

**Results** Many key elements within the five CFIR domains were identified as impacting the implementation of the intervention: (1) intervention characteristics—evidence strength and quality, design quality and packaging, relative advantage and complexity; (2) outer setting—patients' needs and resources, external policies and incentives; (3) inner setting—structural characteristics, networks and communication, culture, compatibility, readiness for implementation and leadership engagement; (4) characteristics of the managers and HCPs—knowledge and belief about the intervention; (5) process—planning, opinion leaders, formally appointed internal implementation leaders, reflecting and evaluating.

**Conclusion** This study revealed the organisational and contextual aspects of the implementation based on different and complementary perspectives. With the growing demand for interdisciplinary teams in primary care, we believe that our insights will be helpful for practices, researchers, and policymakers interested in the implementation of disease prevention and management programmes for people with multiple chronic conditions in primary care.

**Trial registration number** NCT02789800.

## Strengths and limitations of this study

- This study represents one of the few contributions in Quebec to explore, in depth, the factors influencing the implementation of interdisciplinary disease prevention and management interventions for people with multiple chronic conditions in primary care.
- Using an existing framework helps to understand, describe and identify factors that predict the likelihood of implementation success.
- This evaluation of the implementation is limited to managers and healthcare professionals' perspectives and experience.
- Given the qualitative nature of the study, data interpretation could be subjective, and thus, caution should be applied in interpretation.

## BACKGROUND

Chronic diseases (CDs) such as cardiovascular and chronic respiratory diseases, cancer and diabetes are responsible for 70% of deaths worldwide, making them the leading global causes of death.<sup>1,2</sup> Furthermore, many individuals live with more than one CD, now described as multimorbidity, and defined as the coexistence of two or more CDs.<sup>3–5</sup> Given that most people accessing primary care services have more than one CD,<sup>2</sup> multimorbidity is a challenge for both the healthcare system and primary healthcare professionals (HCPs) in primary care.<sup>3–7</sup>

As reported in a recent systematic review,<sup>8</sup> a growing number of studies have investigated the effectiveness of healthcare services and patient-oriented interventions in people with multimorbidity in primary care and community settings. These interventions focused on individualised care plans, self-management support, and to a lesser extent, goal setting and



peer support. Results, in general, were mixed and inconclusive, with most interventions having limited effect on clinical outcomes and patient-reported health outcomes such as quality of life and health service utilisation, or mixed effects on hospital admission rates and medication use and adherence. Thus, the authors highlighted the importance of considering overall participant experience, the context in which interventions occur, and the need to integrate multimorbidity interventions into existing healthcare systems to support implementation and sustainability.<sup>8</sup> These recommendations are consistent with the chronic care model (CCM) focused on primary care.<sup>9–11</sup> In Canada, several innovations focusing on multimorbidity based on the CCM, self-management programmes and primary care renewal have been implemented, and very few of them have been assessed.<sup>12</sup> For example, in 2015, Quebec's health and social services system has been reformed with a centralisation process that abolished Regional Health Authorities.<sup>13</sup> The government of Quebec explicitly stated that this reorganisation was put in place to 'facilitate and simplify access of services to the population, to improve the quality and security of services, and to increase the efficacy and efficiency of the health system'.<sup>14</sup>

The Patient-Centred Innovations for Persons with Multimorbidity research programme was developed to evaluate two complex interventions implemented in two Canadian provinces (Quebec and Ontario).<sup>15</sup> This study is a part of the Quebec intervention.

The Quebec intervention was a 4-month, pragmatic, interdisciplinary intervention for preventing and managing CDs to support self-management of patients with multimorbidity in primary care. The intervention was conducted from April 2016 to July 2017. The intervention consisted of six components: (1) relocating HCPs (nurses, nutritionists, kinesiologists or respiratory therapists) into Family Medicine Groups (FMGs); (2) training HCPs on the patient-centred care (PCC) approach for patients with multimorbidity, interprofessional collaboration, motivational interviewing and self-management support; (3) forming, with key resource persons within each FMG (nurses) and an FMG coordinator, a community of practice aiming to support the integration of the intervention, to ensure the quality of the care, harmonise ongoing changes to practice and consolidate achievements; (4) assessing patients' eligibility for the programme by family physicians or nurses; (5) providing each eligible patient with 1-hour initial assessment by a primary care nurse to create an intervention plan focused on their needs, values, preferences and according to their objectives; and (6) directing patients to the most appropriate HCPs (nutritionists, kinesiologists or the respiratory therapist) who ensure to integrate them into clinical decision-making and outcome assessments according to their intervention plan. The template for the Intervention Description and Replication checklist<sup>16</sup> is available in online supplemental appendix S1.

To that end, the implementation process assessment can offer insights into the 'black box' of interventions (an

approach that can elaborate on the mechanisms through which changes in the outcomes operate) and identify how the potential effects of interventions are moderated and mediated.<sup>17</sup> Thus, this process evaluation aimed to identify barriers and facilitators in implementing an interdisciplinary PCC intervention for patients with multimorbidity in primary care.

## METHODS

### Theoretical framework

The Consolidated Framework for Implementation Research (CFIR) was used to examine the implementation of interdisciplinary intervention for people with multimorbidity in primary care in one region of Quebec, Canada. The CFIR is a compilation of 39 constructs related to implementation and divided into five domains: (1) characteristics of the intervention; (2) outer setting; (3) inner setting; (4) characteristics of the individuals involved and (5) the process of implementation. According to Damschroder *et al.*,<sup>18</sup> researchers may select the constructs from the CFIR that are most relevant for their study setting. The CFIR was chosen based on its comprehensiveness and ability to manage both breadth and depth of data to capture the intervention implementation's complexity. In addition, it includes many implementation aspects and is thus considered a helpful framework for illuminating barriers and facilitators influencing the implementation.<sup>18</sup>

### Study design and research sites

This qualitative descriptive study was part of a larger concurrent triangulation mixed-methods study.<sup>15</sup> A qualitative design allows answering questions about experience, meaning and perspective, most often from the participant's standpoint.<sup>19</sup> Furthermore, qualitative descriptive studies' goal is to summarise specific events experienced by individuals or groups of individuals.<sup>20</sup> Therefore, it is essential to use guidelines such as the Consolidated criteria for Reporting Qualitative research (COREQ) checklist to avoid inadequate reporting that can lead to inappropriate application of qualitative research in decision-making, healthcare, health policy and future research.<sup>21</sup> The COREQ was used to guide reporting (see online supplemental appendix S2).

### Participants and sampling strategy

Seven of the 11 FMGs from Saguenay-Lac-Saint-Jean, a region in Quebec, Canada, participated in both evaluation aspects (quantitative and qualitative). FMGs are primary care clinics where family physicians work with other HCPs to provide comprehensive primary care.<sup>22</sup>

A purposive sample of HCPs (family physicians, nurses, nutritionists, kinesiologists and respiratory therapist) and managers was recruited from the FMGs. A recruitment invitation was sent by email to all HCPs and managers who participated in the programme or its implementation, followed by phone call reminders.

The sample size was determined based on achieving theoretical saturation with a minimum of 12 participants.<sup>23</sup>

### Data collection

Two semistructured interview guides based on the literature review and the CFIR framework tailored to each participant group (managers and HCPs) were developed and pilot tested. The interview guides consisted of open-ended questions related to the participants' perceptions, expectations, role in the intervention implementation, and impact on their work and their organisation's functioning (see online supplemental appendix S3).

The individual semistructured interviews were conducted from October 2016 to September 2017 (6 months after the beginning of the intervention) by a research coordinator (TB), a PhD student (MS), a research assistant (BBD) trained in conducting qualitative interviews and two senior researchers (MCC, MF). The interviews lasted between 23 and 74 min (average of 47 min) and were conducted face-to-face at the various sites. The interviewers also took field notes during the interviews. All interviews were audio-recorded. The goal of the interview was explained to all participants.

### Data analysis

A hybrid process of inductive and deductive thematic content analysis in a stepwise manner, as Braun and Clarke described,<sup>24</sup> was conducted as an analysis approach. Six steps were followed through the process. First, a research assistant with qualitative expertise (CF) read all transcripts and identifies possible themes (step 1). Second, two research team members (PN, CF) developed a coding scheme based on an independent review of three transcripts. Initial codes were created as themes based on the five domains in the CFIR framework.<sup>6</sup> New codes were created if some parts of the transcripts did not directly fit into any CFIR constructs. Discussions with the research team reached an agreement on a final coding scheme. The same research team members used this to code all transcripts using NVivo V.11.0 to assist with data management (step 2). Together, PN and CF discussed and identified recurring and converging themes across participants. The codes that did not fit into the CFIR framework were evaluated and reconsidered. The refined themes were then discussed and agreed on with other research team members (MF, JBB) (steps 3 and 4). Finally, key quotes that illustrated each theme were extrapolated from the data (steps 5 and 6). Finally, credibility was established to ensure the trustworthiness of this qualitative research.

Credibility criteria involve establishing that qualitative research results are believable to the study participants.<sup>25</sup> This was done by (a) data triangulation of sources among the study participants showing, (b) iterative review of transcripts and (c) showing the findings of this study to research team members from different disciplines involved in the study. As a result, they were able to recognise the implementation issues raised by this study.

### Patient and public involvement

No patient involved.

### Findings

A total of 29 interviews were conducted. Intervention stakeholder participants included managers (n=10, including two family physicians) and 19 HCPs, including family physicians, nurses, kinesiologists, nutritionists and respiratory therapists (table 1). Information about participants invited and the number who declined or did not respond is presented in the COREQ checklist (see online supplemental appendix S2).

We have chosen to present our findings by CFIR domains. However, the complexity of the intervention and implementation processes made it difficult to separate key findings by constructs within each domain. As such, our findings are organised into seven themes that reflected participants' experiences with the intervention by the CFIR framework (see online supplemental appendix S4).

### Characteristics of the intervention

The intervention characteristics were more often perceived by the participants as facilitators rather than as barriers to implementation.

### Familiarity with intervention principles

Participants reported that they were already successfully applying certain principles being put forward in the intervention. 'Yes, [the intervention] has positive points. But you know ..., we just used methods that we already know, that we already knew to be efficient, and then reintegrate into a more structured form' (Participant HCP1).

Participants' prior experiences or practices with interdisciplinary collaboration, motivational approaches and research have facilitated the intervention implementation as described by these participants. 'We were already working in interdisciplinarity with a nutritionist and a social worker; we had done this before' (Participant Manager 1). Another participant explained: 'This

**Table 1** Characteristics of participants

	Nurses n=12	Nutritionists n=4	Kinesiologists n=2	Respiratory therapists n=1	Managers n=10
Sex, women	9	4	1	1	8
Age, range (years)	20–52	20–69	20–30	40–49	30–69
Experience, range (years)	12–24	4–18	3–6	13	1–13

approach [motivational] was used in all other settings where I worked [...]. So, we have already used it for several years' (Participant HCP2).

#### Appreciation of intervention components and relative advantage

The majority of the participants believe the intervention was well designed and innovative. For example, HCPs highly appreciated the intervention's approaches, such as the PCC and motivational approaches. 'It is a new way to interact with patients because it is focused on them, on what they want to do with their chronic conditions' (Participant HCP3).

In addition, the training offered at the beginning of the project also facilitated the implementation of the intervention by allowing the HCPs to learn the new concepts before they applied them in practice. 'Introduce us to the approach, describe it to us so that we can embrace it and then apply it' (Participant HCP4).

The participants described several advantages of the intervention. First, relocating nutritionists and kinesiologists into the FMGs improved interprofessional collaboration and patient follow-up. 'Before (this intervention), clients were referred to the hospital, it was more challenging to have communication and to follow up on what happened, being on-site we have access to the whole background of the patient, then we are more able to target and then intervene correctly' (Participant HCP5).

Another perceived advantage was that the intervention enabled HCPs to focus on their roles and use their skills to manage patients. 'I have a good opinion of it because it puts the overall competence of the nurse in the foreground, both at the evaluative level, the teaching level and at the collaborative level. It positions nursing ... in its pivotal role' (Participant Manager 2).

Some HCPs felt that the intervention's preventive rather than curative nature changed their practices, thereby reducing health and emergency services. 'I find it interesting because we see people very early, we can identify people who will develop problems for which they do not yet need medication, problems not already burdensome for the system' (Participant HCP6).

Finally, according to some managers, the financial support received to deploy the intervention facilitated its implementation. 'Receiving funds to help us get started is always welcome rather than reorganizing activities using our resources' (Participant Manager 3).

The only barrier reported by the participants was the complexity of the intervention. According to Damschroder *et al.*,<sup>18</sup> the complexity describes the perceived difficulty of implementation, reflected by duration, scope, radicalness, disruptiveness, centrality, intricacy and the number of steps required for the implementation.<sup>6</sup>

Some of the participants found the intervention complex. They viewed some concepts as difficult. They found some concepts difficult to explain to patients and not applicable to everyone. From the HCP's perspective, leaflets explaining the intervention could have been beneficial for the patients. 'We did not have any document to

give to patients, to explain what [the intervention] meant. It was difficult for us to explain it' (Participant HCP7). They also found it challenging to use the motivational approach. 'Motivational interviewing is very difficult; I feel that I lacked information a little bit' (Participant HCP8).

#### Outer setting

The components of the outer setting were perceived as present and positive in the implementation.

#### Health system reorganisation

For most managers, the health reform context had influenced the ordinary course of the intervention implementation, particularly in coordination and monitoring.

The health system reform was undoubtedly a difficulty because this great project happened simultaneously as the whole reorganization, which led to the creation of the CIUSSS. (Participant Manager 3)

Each team lost the link with its manager. Even for me, dealing with new teams and the realities from one sector to another was not easy. This change had an impact on the implementation of the approach. (Participant Manager 4)

Regarding the impact of the health system reorganisation, a manager mentioned it as 'a difficult deployment' (Participant Manager 5).

#### Inner setting

##### Internal organisation

Participants specifically discussed coordination and monitoring of patient appointments by administrative staff, the sharing of patient records and staff turnover and vacations as key issues.

Having administrative staff coordinate and monitor patient appointments was unanimously emphasised by the HCPs interviewed. They viewed this support as important for the proper conduct of the intervention. For example, a participant working in two different FMGs, one with the support of a secretary and the other without this support, stated that 'It makes things a lot easier, it is not comparable' (Participant HCP9).

The ability to navigate through the patient's electronic medical record and share information among the various HCPs supported the implementation of the intervention. 'It is very facilitating; we see the notes of the doctors, the nutritionist, the nurses. Moreover, we know why we see him [...]' (Participant HCP10).

Participants appreciated the community of practice's monthly telephone meetings to discuss the evolution of the implementation and share their knowledge and experiences. 'During these meetings, we discussed specific themes. We updated our knowledge and shared literature. We also discussed cases that we encountered, shared the experiences and difficulties of the implementation' (Participant Manager 6).

HCPs' turnover interfered in the implementation because some newly relocated HCPs were not systematically trained in the approach. Their training delayed their integration into the interdisciplinary team. 'The new staff training resulted in delays in the intervention' (Participant Manager 4).

Concerning patients' follow-up, the communication between HCPs or family physicians could vary depending on their environment's reality. In some settings, face-to-face communication was possible during the implementation of the intervention. For other settings, it is the use of indirect communication, such as email, that has facilitated interdisciplinary collaboration. 'Since the implementation of the intervention, we take 5 minutes to discuss, either at the end or before dinner [the day] ... that allows us to ensure better follow-up' (Nurse 11). For other settings, the use of email was prevalent. 'When I have to speak with doctors, it is often by e-mail. We do not see them that much. We do not have an interdisciplinary meeting with them' (Participant HCP10).

However, certain aspects of the internal organisation of FMGs, such as the availability of premises and health professionals, negatively influenced the implementation of the intervention.

The lack of infrastructure and scarcity of resources in some FMGs negatively impacted interprofessional collaboration and interdisciplinarity.

A manager summed it up nicely: 'One constraint was the lack of physical space in some FMGs. Therefore, some healthcare professionals could not meet together to discuss patient action plans' (Participant Manager 4).

Some health professionals working part-time did not feel they had enough time to discuss patient medical records with their colleagues. This situation probably influenced negatively the implementation of interprofessional collaboration, which was an essential component of the intervention.

I am here for half a day. We do not necessarily have time to go through the patient's file. I do not have time to read the notes of all my colleagues before I see my patients. (Participant HCP11)

#### Compatibility of the intervention with participants' vision and values

The principles of the intervention were in line with FMGs' vision and values, which facilitated the implementation. 'I believe in it, and that is where we have to go more and more. It is aligned with the vision of our organization, but also the concept of chronic disease management' (Participant Manager 4).

The majority of health professionals explained how the intervention's approaches, such as interprofessional collaboration, motivational approach, self-management and the PCC approach, correspond to their values and vision as HCPs.

I love it because I have always believed in interdisciplinarity, not multidisciplinary. I have always believed

in supporting self-management support and in motivational interviewing. (Participant HCP12)

Some HCPs described a coordinating nurse in each FMG responsible for supervising the entire team and ensuring that the intervention was done appropriately. 'There is also a coordinator who is always near us and who makes it work' (Kinesiologist 14).

However, some managers felt that the intervention principles were not compatible with the family physicians' philosophy and practice and perhaps hindered the physicians' involvement. 'There was a lack of collaboration. Family physicians do not adhere to the [intervention] philosophy. This approach consists of identifying client needs. They still tend to say: 'I am the doctor, I know what you need. I will tell you what you need, then apply it' (Participant Manager 6).

#### Characteristics of the individuals involved

##### Leadership engagement

The leadership conveyed by the managers played an essential role in implementing the intervention by ensuring the permanent flow of information, the mobilisation of teams and the supervision of the training of newly relocated HCPs. 'My contribution is to talk about it regularly to the teams, to mobilize them and then to make sure that I have the people in place to support. Finally, to ensure that they are adequately trained' (Participant Manager 7). Indeed, some participants felt the managers' involvement and commitment have certainly facilitated the implementation of the intervention. 'The managers are involved in the project. They believe in it, and we also believe in it at the organizational level' (Participant HCP13).

Furthermore, the presentation and promotion of the intervention with HCPs working in the FMGs, conducted jointly by researchers and managers, were highly appreciated. This activity added more credibility, enhanced the participants' consideration of adopting the intervention's approach and probably facilitated/further supported the implementation process. 'It is also good that it was clinicians, managers and researchers who presented this approach, so it brought credibility to the approach' (Participant Manager 4).

On the other hand, the low participation of some family physicians negatively influenced the implementation of the intervention. Participants' reasons were doctors' age, time to devote to interdisciplinary activities and the doctors' vision of nurses' role. 'The collaboration with the doctors was not a success. I was afraid to refer clients' (Participant HCP14).

#### Process of implementation

##### Communication

Communication, both face-to-face and through telephone meetings, has helped maintain interaction between HCPs during implementation.

Direct communication was the primary means to present the intervention to the stakeholders (FMG managers, family physicians and other HCPs). First,



meetings were organised by the investigators (research team and leaders of the CIUSSS) to present the intervention and services offered to each FMG manager. Then, when an FMG decided to implement the intervention, each FMG manager (coordinator, manager or nurse) presented the intervention to family doctors and health professionals.

My contribution is to talk to the teams as much as possible, to mobilize them. Ensure that there are people in place to support it, that people feel comfortable doing it, have the proper training, and that there are also interdisciplinary relationships between nurses and other health professionals. Make sure that they understand what the intervention consists of and how to intervene with this clientele. (Participant Manager 7)

Furthermore, some managers explained how telephone meetings during the implementation of the intervention facilitated communication with HCPs. 'At one point, we tried to use technology to make conference calls within the whole region. It was a facilitator or a success factor on some level because some teams did not speak to each other' (Participant Manager 4).

## DISCUSSION

This study aimed to evaluate the implementation of interdisciplinary PCC intervention for people with multimorbidity in primary care. Many key elements were identified as facilitating the implementation of the intervention (familiarity with intervention principles, the quality and the relative advantage of the intervention, the leadership engagement). However, some obstacles were identified such as the complexity of the intervention, the health system reorganisation, the internal organisation of FMGs, the compatibility of the intervention principles with some family physicians' philosophy and practice.

Prior work indicates an increased likelihood that stakeholders who subscribe to the principles of evidence-based interventions for the prevention and management of CDs (CCM, PCC) in primary care will be motivated to implement the programme.<sup>26</sup> Furthermore, there is evidence that innovations in CDs, which has a relative advantage over standard care, have led to greater implementation.<sup>13</sup> For example, the Canadian Heart Health Kit, a risk management and patient education resource for the prevention of cardiovascular disease and promotion of cardiovascular health for which the participants perceived the relative advantage, resulted in better implementation.<sup>27</sup> The organisational culture, the leadership of managers and leaders' presence are also recognised as successful ingredients for implementing an intervention based on the CCM in primary care.<sup>10</sup>

On the other hand, the literature also shows that hierarchical working relationships without leadership for change negatively influence healthcare changes intervention and innovation.<sup>28</sup> Thus, transforming care

practices in a primary care organisation requires a culture of support for change and learning.<sup>29 30</sup> Furthermore, leaders can be agents of change by ensuring resources and providing adequate support to staff.<sup>31</sup>

The intervention was aligned with the participants and their organisation vision. The implementation is successful if the intervention is integrated into the organisation's vision and a recognised need to adopt new care approaches to prevent and manage CDs in primary care.<sup>10 32</sup>

Although this evaluation identified many facilitators, the implementation of the intervention faced several challenges, which are essential to consider. For instance, characteristics of individuals such as individuals' beliefs, knowledge, self-efficacy and personal attributes may affect implementation.<sup>18</sup> In this study, the family physicians' lack of commitment and leadership appeared to deter implementation. The lack of involvement of family physicians in implementing complex interventions, including prevention, has been identified previously in the literature.<sup>33-35</sup> One reason for this barrier might be the inconsistency between the ideals of PCC and the reality of the healthcare system, which is oriented towards the biomedical model.<sup>30 33</sup> However, the role of family physicians remains fundamental in interdisciplinary interventions based on the CCM due to their privileged relationship with chronically ill patients.<sup>35</sup> Family physicians need to maintain a climate of trust and collaboration with other HCPs. Therefore, physicians must be involved from the outset, beginning with the development phase of the intervention to ensure their involvement.<sup>36</sup> The high turnover of HCPs and organisational changes also negatively influenced the intervention implementation. Staff turnover is a common barrier encountered when implementing CD prevention and management interventions in primary care.<sup>37</sup> The difficulty in integrating newly relocated professionals into teams' routines also influenced the implementation, particularly for interprofessional interventions. Organisational changes caused by integrating new staff and interprofessional collaboration with consequences such as lack of time, allocation of roles, distribution of work, work overload and the shortage of health professionals are regularly reported in complex interventions for CDs in primary care.<sup>38</sup> The use of information and communication technologies such as electronic health records or telehealth is often proposed as a solution to overcome these challenges.<sup>39</sup>

## Study strengths and limitations

The strength of the study is the inclusion of different HCPs and managers to reveal several views. It is a contribution to improve the prevention and management of interdisciplinary interventions for people with multiple chronic conditions in primary care. Using an existing framework within implementations, science is considered an important strength to better understand, describe and identify factors that predict implementation success. The CFIR framework

was useful in organising, analysing and categorising the data. Furthermore, applying the CFIR as an analysis lens allows comparisons with other contexts and settings. With the growing demand for interdisciplinarity in primary care, the findings of this study will be useful for practices, researchers, and policymakers interested in implementing disease prevention and management programmes for people with multiple chronic conditions in primary care at local, regional and international levels.

A limitation of this work is the lack of patients' perspective. However, we made this choice because we believe that since the patients did not actively participate in the implementation phase, they could not have shed light to understand the implementation process.

Participation in this study was voluntary and limited to managers and HCPs involved in implementing the intervention in Saguenay-Lac-Saint-Jean, Quebec. Therefore, the research team and social desirability may have influenced the perspectives of these participants.

## CONCLUSIONS

Implementing an interdisciplinary PCC intervention to prevent and manage CDs in primary care organisations is a complex process. Using the CFIR as a guide, this evaluation identified multiple elements that assisted in implementing the intervention. In addition, the results revealed organisational and contextual aspects of the implementation based on different and complementary perspectives. With the growing demand for interdisciplinarity in primary care, we believe that our insights will be useful for practices, researchers, and policymakers interested in implementing disease prevention and management programmes for people with multiple chronic conditions in primary care.

### Author affiliations

<sup>1</sup>Department of Family Medicine and Emergency Medicine, Université de Sherbrooke, Sherbrooke, Quebec, Canada

<sup>2</sup>Department of Family Medicine, Western University, London, Ontario, Canada

<sup>3</sup>Westmead Applied Research Centre, Faculty of Medicine and Health, University of Sydney, Sydney, New South Wales, Australia

<sup>4</sup>Department of Health Sciences, Université du Québec à Chicoutimi, Chicoutimi, Quebec, Canada

<sup>5</sup>School of Nursing, University of Sherbrooke, Sherbrooke, Quebec, Canada

**Acknowledgements** The authors wish to thank Maud-Christine Chouinard (MCC), Tarek Bouhali (TB), Bayero Boubakar Diallo (BDD) and Martin Desmeules (MD) for their contribution to data collection and analyses. MS was funded by the Dr Brian W Gilbert Canada Research Chair in Primary HealthCare Research (2003–2017). MF holds a Research Chair on Chronic Diseases in Primary Care (Fondation de ma vie, Hôpital de Chicoutimi and Université de Sherbrooke).

**Contributors** Conceptualisation—PN, JBB and MF. Methodology—PN, CF, JBB and MF. Software—PN and CF. Validation—MF and JBB. Formal analysis—PN and CF. Investigation—MaS and MF. Resources—MF and MoS. Data curation—MF and JBB. Writing (original draft preparation)—PN. Writing (review and editing)—PN, JBB, CF, MAAA, TNN, MaS, CL, FG, MoS and MF. Visualization—PN, JBB, CF, MAAA, TNN, MaS, CL, FG, MoS and MF. Supervision—JBB and MF. Project

administration—MF and MoS. Funding acquisition—MF and MoS. All authors have read and agreed to the published version of the manuscript. All authors listed on the manuscript contributed to the study design, data collection, data analysis and interpretation of the findings, and drafting and reviewing the final manuscript.

**Funding** The Patient-Centred Innovations for Persons with Multimorbidity research programme is funded by a team grant from the Community-Based Primary Health Care Signature Initiative of the Canadian Institutes of Health Research (grant number: 297943). Website: [http://webapps.cihirsc.gc.ca/decisions/p/project\\_details.html?applId=283684&lang=en](http://webapps.cihirsc.gc.ca/decisions/p/project_details.html?applId=283684&lang=en).

**Disclaimer** The funders had no role in study design, data collection and analysis, decision to publish or manuscript preparation.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** This study was approved by the Ethics Committee of the Chicoutimi Health and Social Services Center (Ethical code 2013-010). Informed consent was obtained from all participants.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article or uploaded as supplemental information. The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iDs

Patrice Ngangue <http://orcid.org/0000-0002-5204-7420>

Tu Ngoc Nguyen <http://orcid.org/0000-0002-8836-8920>

Maxime Sasseville <http://orcid.org/0000-0003-1694-1414>

Christine Loignon <http://orcid.org/0000-0002-9828-3090>

## REFERENCES

- World Health Organisation (WHO). *Noncommunicable diseases - key facts*. Geneva, Switzerland: WHO, 2021.
- World Health Organisation (WHO). *Noncommunicable diseases progress monitor 2017*. 231. Geneva, Switzerland: WHO, 2017.
- Garin N, Koyanagi A, Chatterji S, *et al*. Global multimorbidity patterns: a cross-sectional, population-based, multi-country study. *J Gerontol A Biol Sci Med Sci* 2016;71:205–14.
- Nguyen H, Manolova G, Daskalopoulou C, *et al*. Prevalence of multimorbidity in community settings: a systematic review and meta-analysis of observational studies. *J Comorb* 2019;9:2235042X1987093.
- World Health Organisation (WHO). *Multimorbidity : technical series on safer primary care*. Geneva, Switzerland: WHO, 2016.
- Fortin M, Bravo G, Hudon C, *et al*. Prevalence of multimorbidity among adults seen in family practice. *Ann Fam Med* 2005;3:223–8.
- Smith SM. Managing multimorbidity in primary care. *CMAJ* 2019;191:E489–90.
- Smith SM, Wallace E, O'Dowd T, *et al*. Interventions for improving outcomes in patients with multimorbidity in primary care and community settings. *Cochrane Database Syst Rev* 2021;1:CD006560.
- Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness. *JAMA* 2002;288.
- Kadu MK, Stolee P. Facilitators and barriers of implementing the chronic care model in primary care: a systematic review. *BMC Fam Pract* 2015;16:12.



- 11 Reynolds R, Dennis S, Hasan I, *et al.* A systematic review of chronic disease management interventions in primary care. *BMC Fam Pract* 2018;19:11.
- 12 Nicholson K, Terry AL, Fortin M, *et al.* Understanding multimorbidity in primary health care. *Can Fam Physician* 2015;61:e489–990.
- 13 Wankah P, Guillette M, Dumas S, *et al.* Reorganising health and social care in Québec: a journey towards integrating care through mergers. *London J Prim Care* 2018;10:48–53.
- 14 Barette G. *Projet de loi n° 10 - Loi modifiant l'organisation et la gouvernance du réseau de la santé et des services sociaux notamment par l'abolition des agences régionales*. Quebec: Edited by Québec ANd, 2014.
- 15 Stewart M, Fortin M, Patient-Centred Innovations for Persons with Multimorbidity Team\*. Patient-centred innovations for persons with multimorbidity: funded evaluation protocol. *CMAJ Open* 2017;5:E365–72.
- 16 Hoffmann TC, Glasziou PP, Boutron I, *et al.* Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;348:g1687
- 17 Nielsen K, Randall R. Opening the black box: presenting a model for evaluating organizational-level interventions. *European Journal of Work and Organizational Psychology* 2013;22:601–17.
- 18 Damschroder LJ, Aron DC, Keith RE, *et al.* Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci* 2009;4:50.
- 19 Hammarberg K, Kirkman M, de Lacey S. Qualitative research methods: when to use them and how to judge them. *Hum Reprod* 2016;31:498–501.
- 20 Kim H, Sefcik JS, Bradway C. Characteristics of qualitative descriptive studies: a systematic review. *Res Nurs Health* 2017;40:23–42.
- 21 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349–57.
- 22 Pineault R, Borgès Da Silva R, Provost S, *et al.* Impacts of Québec primary healthcare reforms on patients' experience of care, unmet needs, and use of services. *Int J Family Med* 2016;2016:1–13.
- 23 Guest G, Bunce A, Johnson L. How many interviews are enough? An experiment with data saturation and variability. *Field Methods* 2006;18:59–82.
- 24 Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3:77–101.
- 25 Leung L. Validity, reliability, and generalizability in qualitative research. *J Family Med Prim Care* 2015;4:324–7.
- 26 Ahmed S, Ware P, Visca R, *et al.* The prevention and management of chronic disease in primary care: recommendations from a knowledge translation meeting. *BMC Res Notes* 2015;8:571.
- 27 Scott SD, Plotnikoff RC, Karunamuni N, *et al.* Factors influencing the adoption of an innovation: an examination of the uptake of the Canadian heart health kit (HHK). *Implement Sci* 2008;3:41.
- 28 Sfantou DF, Laliotis A, Patelarou AE, *et al.* Importance of leadership style towards quality of care measures in healthcare settings: a systematic review. *Healthcare* 2017;5:73.
- 29 InSchottenfeld L, Petersen D, Peikes D. *Creating patient-centered team-based primary care*. Rockville, MD: Agency for Healthcare Research and Quality, 2016.
- 30 Stewart M, Brown J, Weston W. *Patient-centered medicine, transforming the clinical method*. Salem, MA: Radcliffe Medical Press, 2003.
- 31 Santana MJ, Manalili K, Jolley RJ, *et al.* How to practice person-centred care: a conceptual framework. *Health Expect* 2018;21:429–40.
- 32 Wagner EH, Austin BT, Davis C, *et al.* Improving chronic illness care: translating evidence into action. *Health Aff* 2001;20:64–78.
- 33 Rubio-Valera M, Pons-Vigués M, Martínez-Andrés M, *et al.* Barriers and facilitators for the implementation of primary prevention and health promotion activities in primary care: a synthesis through meta-ethnography. *PLoS One* 2014;9:e89554.
- 34 Sanchez A, Grandes G, Pablo S, *et al.* Engaging primary care professionals in collaborative processes for optimising type 2 diabetes prevention practice: the PREDIAPS cluster randomised type II hybrid implementation trial. *Implement Sci* 2018;13:94.
- 35 Wallace PJ. Physician involvement in disease management as part of the CCM. *Health Care Financ Rev* 2005;27:19–31.
- 36 Brand C, Scott I, Greenberg P, *et al.* Chronic disease management: time for consultant physicians to take more leadership in system redesign. *Intern Med J* 2007;37:653–9.
- 37 Davy C, Bleasel J, Liu H, *et al.* Factors influencing the implementation of chronic care models: a systematic literature review. *BMC Fam Pract* 2015;16:102.
- 38 Supper I, Catala O, Lustman M, *et al.* Interprofessional collaboration in primary health care: a review of facilitators and barriers perceived by involved actors. *J Public Health* 2015;37:716–27.
- 39 Sanchez K. Collaborative care in real-world settings: barriers and opportunities for sustainability. *Patient Prefer Adherence* 2017;11:71–4.



## Appendix S1: TiDIER CHECKLIST

Item	
Provide the name or a phrase that describes the intervention	A four-month, pragmatic, interdisciplinary intervention for the prevention and management of chronic diseases aimed at supporting self-management of patients with multimorbidity in primary care.
Describe any rationale, theory or goal of the elements essential to the intervention	See protocol: Stewart M, Fortin M; Patient-Centred Innovations for Persons with Multimorbidity Team*. Patient-Centred Innovations for Persons with Multimorbidity: funded evaluation protocol. CMAJ Open. 2017;5(2): E365–E372. doi:10.9778/cmajo.20160097
Materials: describe any physical or informational materials used in the intervention, including those provided to participants or used in intervention delivery or in the training of the intervention providers	Added information : Each healthcare professional was given a training handbook containing information about each intervention component. No specific material was used during interventions. Instead, healthcare professionals used their material.
Procedures: Describe each of the procedures, activities and /or processes used in the intervention, including any enabling or support activities	Patients were assessed for eligibility by family physicians or registered nurses. Each eligible patient was provided with a one-hour initial assessment by a primary care nurse to create an intervention plan focused on their needs and according to their objectives and; to direct patients to other healthcare professionals (nutritionists, kinesiologists or the respiratory therapist) according to their intervention plan. Each patient's intervention had to be based on the educational and coaching content of the training (patient-centered care approach for patients with multimorbidity, self-management support, and motivational interviewing). Interdisciplinary meetings

	between family physicians, nurses and other health professionals were to be held to discuss cases and harmonize the intervention plan.”
For each category of intervention provider, describe their expertise, background, and any specific training given.	Practice Nurse; Nutritionists; Kinesiologists; Respiratory therapists. The training was given to all healthcare professionals who worked with chronic diseases patients. This involved being trained on patient-centered care approach for patients with multimorbidity, self-management support, and interprofessional collaboration and motivational interviewing. Their training lasted an average 7.8 hours.
Describe the modes of delivery (such as face-to-face or by some other mechanism, such as internet or telephone) of the intervention, and whether it was provided individually or in a group	Interventions were provided individually and face-to-face.
Describe the type(s) of location(s) where the intervention occurred, including any necessary infrastructure or relevant features	Recruitment and intervention was undertake in the patient’s FMGs.
Describe the number of times the intervention was delivered and over what period of time including the number of sessions, their schedule and their duration, intensity or dose.	Once recruited, patients had an initial assessment with the nurse during approximately one hour. Based on patients’ centered care, no specific number of sessions was planned. Therefore, patients could see healthcare professionals as much (or as little) as they wanted during a 4-month period. In average, patients had 2.6 hours of interventions throughout the 4-month period.
If the intervention was planned to be personalised, titrated or adapted, then describe what, why, when and how.	See above.
If the intervention was modified during the course of the study, describe the changes (what, why, when and how)	N/A

Planned: if intervention adherence or fidelity was assessed, describe how and by whom, and if any strategies were used to maintain or improve fidelity, describe them.	Intervention adherence was assessed after intervention completion by the research team.
Actual: if intervention adherence or fidelity was assessed, describe the extent to which the intervention was delivered as planned.	Intervention adherence will be discussed in another paper that is now in the publication process.

## Appendix S2. Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist

No	Item	Guide questions/description	
<b>Domain 1: Research team and reflexivity</b>			
Personal Characteristics			
1.	Interviewer/facilitator	Which author/s conducted the interview or focus group?	None, but one author (MF) supervised closely the work of the professionals conducting the interviews
2.	Credentials	What were the researcher's credentials? <i>E.g. PhD, MD</i>	MF : MD, MSc, CMFC(F) MCC : RN, PhD BBD : PhD TB : MSc
3.	Occupation	What was their occupation at the time of the study?	MF : Researcher MCC : Researcher BBD : Research assistant TB : Research coordinator
4.	Gender	Was the researcher male or female?	MF : Male MCC : Female

No	Item	Guide questions/description	
5.	Experience and training	What experience or training did the researcher have?	MF : Family physician, MSC, senior researcher MCC : Registered nurse, PhD and senior researcher
Relationship with participants			
6.	Relationship established	Was a relationship established prior to study commencement?	No
7.	Participant knowledge of the interviewer	What did the participants know about the researcher? e.g. <i>personal goals, reasons for doing the research</i>	The study was presented to all participants before enrollment. All participants signed a consent form. The goal of the interview was explained to all participants before the interviews (page 5).
8.	Interviewer characteristics	What characteristics were reported about the interviewer/facilitator? e.g. <i>Bias, assumptions, reasons and interests in the research topic</i>	Interviews were conducted by a research coordinator (TB), a PhD student (MS), a research assistant (BBD) trained in conducting qualitative interviews, and two senior researchers (MCC, MF). All of them worked with the research team since the beginning of the study. (page 5)
<b>Domain 2: study design</b>			
Theoretical framework			
9.	Methodological orientation and Theory	What methodological orientation was stated to underpin the study? e.g. <i>grounded theory,</i>	Descriptive qualitative approach (close to Content analysis) embedded in a randomized trial (page 4)

No	Item	Guide questions/description	
		<i>discourse analysis, ethnography, phenomenology, content analysis</i>	
Participant selection			
10.	Sampling	How were participants selected? <i>e.g. purposive, convenience, consecutive, snowball</i>	Maximum variability was targeted based on age, gender, socio-economic status, practices (page 4).
11.	Method of approach	How were participants approached? <i>e.g. face-to-face, telephone, mail, email</i>	The managers and healthcare professionals were contacted by phone at their working place (page 4).
12.	Sample size	How many participants were in the study?	29 accepted to participate (page 5). - 10 managers including two family physicians - 19 healthcare professionals
13.	Non-participation	How many people refused to participate or dropped out? Reasons?	- All (29) participants contacted accepted to participate (page 6). - No participant dropped out
Setting			
14.	Setting of data collection	Where was the data collected? <i>e.g. home, clinic, workplace</i>	Primary health care practices (page 4)

No	Item	Guide questions/description	
15.	Presence of non-participants	Was anyone else present besides the participants and researchers?	No
16.	Description of sample	What are the important characteristics of the sample? <i>e.g. demographic data, date</i>	Managers including family physicians and healthcare professionals, nurses, kinesiologists, nutritionists and a respiratory therapist actively participated in the program or its implementation (page 4).
Data collection			
17.	Interview guide	Were questions, prompts, guides provided by the authors? Was it pilot tested?	The interview guides included open questions relating the perceived factors influencing the implementation of intervention developed by the research team based on theoretical framework. Interviews were in French. It was pretested and adjusted accordingly (page 4, Appendix S3).
18.	Repeat interviews	Were repeat interviews carried out? If yes, how many?	Not with this design
19.	Audio/visual recording	Did the research use audio or visual recording to collect the data?	All interviews were audio-recorded (page 5)
20.	Field notes	Were field notes made during and/or after the interview or focus group?	The interviewers took notes during and after the interviews (page 6).
21.	Duration	What was the duration of the interviews or focus group?	Average 47 minutes (page5)

No	Item	Guide questions/description	
22.	Data saturation	Was data saturation discussed?	Yes
23.	Transcripts returned	Were transcripts returned to participants for comment and/or correction?	No
<b>Domain 3: analysis and findings</b>			
Data analysis			
24.	Number of data coders	How many data coders coded the data?	Two coders (PN and CF) with regular validation with one of the authors (MF). The two coders are also co-author (page 5).
25.	Description of the coding tree	Did authors provide a description of the coding tree?	No
26.	Derivation of themes	Were themes identified in advance or derived from the data?	The first codes were based on interview guide. Many other codes derived from the theoretical framework data following a mixed coding approach (inductive and deductive). (Page 5)
27.	Software	What software, if applicable, was used to manage the data?	NVivo 11 (page 5)



No	Item	Guide questions/description	
28.	Participant checking	Did participants provide feedback on the findings?	Not indicated (ref. Braun, V., & Clarke, V. Using thematic analysis in psychology. <i>Qualitative Research in Psychology</i> 2016, 3: 77–101.)
Reporting			
29.	Quotations presented	Were participant quotations presented to illustrate the themes / findings? Was each quotation identified? e.g. <i>participant number</i>	Yes. All quotes have been translated to English by a Professional Translator
30.	Data and findings consistent	Was there consistency between the data presented and the findings?	All findings are based on the collected and analyzed data. Major findings are consistent with the quantitative results (triangulation).
31.	Clarity of major themes	Were major themes clearly presented in the findings?	Major themes are described in the manuscript according to the theoretical framework (page 6 to 11, and appendix S4).
32.	Clarity of minor themes	Is there a description of diverse cases or discussion of minor themes?	All themes are derived from the theoretical framework and described in the manuscript (page 6 to 11, and appendix S4).

### Appendix Interview guides

Participants	Questions
Managers	<ol style="list-style-type: none"> <li>1. Can you describe the program and your role? Does the program meet your expectations? How does it meet your expectations?</li> <li>2. What is your general opinion of the program?</li> <li>3. What do you think facilitates the activities of the program?</li> <li>4. How do these factors have influenced the program effectiveness?</li> <li>5. In your opinion, what are the barriers that this program faces?</li> <li>6. About this program, are there any contextual factors that serve as facilitators or barriers (Question: space, time, distance, organizational problems)</li> </ol>
Healthcare professionals	<ol style="list-style-type: none"> <li>1. How was the patient referral made? How did the communication take place (phone, paper, report, etc.)?</li> <li>2. Are you aware of whether a patient action plan/intervention plan has been developed as part of the program by one or more professionals (nurse, nutritionist, etc.) and the patient? If so, are you aware of the plan of the elements? Did you know if the plan was changed in the process of being carried out?</li> <li>3. Tell us about the dynamic between the team members?</li> <li>4. Tell us about the interdisciplinary practices in this program?</li> <li>5. Tell us about the patient's role in the healthcare team?</li> <li>6. What is your general opinion about the program?</li> <li>7. Does it meet your expectations?</li> <li>8. How would you describe the impact of the program on your patients and their families?</li> <li>9. How do program services help patients manage their chronic diseases?</li> </ol>

Appendix S4. Key elements influencing the intervention implementation based on CFIR domains

Domains	Barriers	Facilitators
<b>A. Intervention characteristics</b>		Quotes
1. Familiarity with interventions principles		Participants reported that they were already successfully applying certain principles being part of the intervention.  Participants' prior experiences or practices in interdisciplinary collaboration, motivational approach and research
2. Appreciation of intervention components and relative advantage		Most of the participants believe the intervention was well designed and innovative. For example, healthcare professionals highly appreciated the intervention's approaches, such as the patient-centred care and motivational approaches.
4. Complexity	Some concepts were difficult to explain to patients and not applicable to everyone.  It was challenging to use the motivational approach	
<b>B. Outer setting</b>		
1. Health system reorganization	The health reform context has influenced the ordinary course of the intervention implementation, particularly in terms of coordination and monitoring.	
<b>C. Inner setting</b>		
1. Internal organization	Staff turnover/holidays and vacations  Lack of infrastructures in some FMGs negatively impacted interprofessional collaboration and interdisciplinarity.	Coordination/monitoring of patient appointments  Sharing patient records
2. Compatibility	The principles of the intervention were not compatible with the family physicians' philosophy and practice. This likely hindered interprofessional collaboration and negatively influenced physician involvement	the approaches advocated by the intervention such as interprofessional collaboration, motivational approach, self-management as well as the patient-centered care approach correspond to their values and vision as healthcare professionals.

<b>D. Characteristics of individuals</b>		
1. Leadership engagement		<p>Managers involvement and commitment have certainly facilitated the implementation of the intervention.</p> <p>The presentation and promotion of the intervention with healthcare professionals working in the FMGs, conducted jointly by a group of researchers and managers, was highly appreciated.</p>
<b>E. Process of implementation</b>		
1. Communication		<p>Communication, both face-to-face and through telephone meetings, has helped maintain interaction between healthcare professionals during implementation.</p>