BMJ Open Current use of inotropes according to initial blood pressure and peripheral perfusion in the treatment of congestive heart failure: findings from a multicentre observational study

Kazuya Nagao ¹, Takao Kato ², Hidenori Yaku³, Takeshi Morimoto ⁴, Yasutaka Inuzuka⁵, Yodo Tamaki⁶, Erika Yamamoto², Yusuke Yoshikawa², Takeshi Kitai⁷, Ryoji Taniguchi⁸, Moritake Iguchi ⁹, Masashi Kato³, Mamoru Takahashi¹⁰, Toshikazu Jinnai¹¹, Tomoyuki Ikeda¹², Takafumi Kawai¹³, Akihiro Komasa¹⁴, Ryusuke Nishikawa¹⁵, Yuichi Kawase¹⁶, Takashi Morinaga¹⁷, Kanae Su¹⁸, Mitsunori Kawato¹⁹, Yuta Seko², Tsukasa Inada¹, Moriaki Inoko²⁰, Mamoru Toyofuku¹⁸, Yutaka Furukawa²¹, Yoshihisa Nakagawa²², Kenji Ando¹⁷, Kazushige Kadota¹⁶, Satoshi Shizuta ², Koh Ono², Yukihiro Sato⁸, Koichiro Kuwahara²³, Neiko Ozasa², Takeshi Kimura ²

To cite: Nagao K, Kato T, Yaku H, *et al.* Current use of inotropes according to initial blood pressure and peripheral perfusion in the treatment of congestive heart failure: findings from a multicentre observational study. *BMJ Open* 2022;12:e053254. doi:10.1136/bmjopen-2021-053254

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-053254>).

Received 09 May 2021
Accepted 22 December 2021

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Takao Kato;
tkato75@kuhp.kyoto-u.ac.jp

ABSTRACT

Objectives Current guidelines restrict the use of inotropes for the treatment for heart failure (HF) unless the patients are hypotensive or hypoperfused because of safety concerns. This study sought to characterise the contemporary real-world use of inotropes and associated long-term outcomes according to systolic blood pressure (sBP) and perfusion status.

Design A multicentre prospective cohort study.

Setting This study was nested from the Kyoto Congestive Heart Failure registry, which included consecutive Japanese patients admitted for HF.

Participants We categorised 3995 patients into two groups: sBP ≥90 mm Hg and warm profile group, and sBP <90 mm Hg or cold profile group. In each group, patients were stratified across the use of inotropes within 24 hours of hospital presentation.

Primary and secondary outcomes The primary outcome was all-cause death throughout follow-up. Secondary outcomes included cardiovascular death throughout follow-up, all-cause death during index hospitalisation and after discharge, and HF hospitalisation.

Results A total of 793 patients (20%) presented with sBP <90 mm Hg or cold profile, whereas 3202 patients had sBP ≥90 mm Hg and warm profile; 276 patients (35%) in the sBP <90 mm Hg/cold group and 312 patients (10%) in the sBP ≥90 mm Hg/warm group received initial inotropic treatment. Adjusted excess risk of inotrope use relative to no inotrope for the primary outcome measure was significant in the sBP ≥90 mm Hg/warm group (adjusted HR, 1.36; 95% CI 1.09 to 1.72, p=0.006) but not in the sBP <90 mm Hg/cold group (adjusted HR, 1.28, 95% CI 0.96 to 1.69, p=0.09). Risk for postdischarge all-cause death and HF hospitalisation was not significantly different between the patients with inotropes and no inotropes in both groups.

Strengths and limitations of this study

- This study offers the current real-world patterns and outcomes of initial inotropic treatments in congestive heart failure.
- Data are derived from a large contemporary cohort in Japan.
- The observational nature of this study precludes the demonstration of a causal relationship between initial inotrope use and outcomes.
- Despite a careful and comprehensive attempt to correct the heterogeneity of the patient characteristics, unmeasured covariates may account for outcomes.
- The exact dose of inotropic agents and duration of their use were not available.

Conclusion Inotrope use in the absence of hypotension and hypoperfusion is still common, but associated with a worse long-term prognosis.

Trial registration number UMIN000015238.

INTRODUCTION

Congestive heart failure (HF) is a life-threatening medical condition that often requires in-hospital management. Most hospitalised patients with HF are treated with diuretics to attenuate fluid overload, while others may receive intravenous vasodilators, which decrease ventricular filling pressure and relieve symptoms.^{1–5} On the other hand, inotropes can be used for patients with a severe reduction in cardiac output to maintain vital

organ perfusion and blood pressure. However, previous studies have raised concerns regarding safety of inotrope use due to increased morbidity and mortality.^{6–8} Inotropes, especially those with adrenergic mechanisms such as dobutamine and dopamine, can increase left ventricular afterload and cause tachycardia, myocardial ischaemia and arrhythmias. Non-catecholamine inotropes, such as milrinone, may increase the risk of hypotension.⁹ Thus, current American College of Cardiology (ACC)/American Heart Association (AHA) guideline states that the use of intravenous inotropic agents in hospitalised patients without documented low blood pressure or impaired perfusion is potentially harmful (Recommendation: class III, level of evidence: B).² Current European Society of Cardiology (ESC) guideline also limits the use of inotropes to the patients with hypotension or to those with signs of impaired peripheral perfusion such as ‘cold’ HF profile (Recommendation: class III, level of evidence: A).¹ However, decisions regarding the initial use of vasoactive agents still remains largely subject to clinician discretion.^{1,2,4} Despite an important decision making basis for the initiation of inotropes, few studies have considered the clinical signs of perfusion status such as the ‘cold’ or ‘warm’ profile for the evaluation of outcomes.¹⁰ In addition, most previous studies have evaluated the short-term morbidity and mortality of inotrope use, but data on long-term outcomes associating inotrope use are scarce.¹⁰

The aim of this study is to assess the current initial use of inotropic therapy and associated outcomes according to initial blood pressure and perfusion status in contemporary real-world practice in Japan.

METHODS

Patient population

The Kyoto Congestive Heart Failure (KCHF) registry is a physician-initiated, prospective, observational, multi-centre cohort study that enrolled, without exclusion, all consecutive patients who were hospitalised for congestive HF between October 2014 and March 2016 in 19 participating hospitals in Japan. All patients enrolled in the KCHF registry had signs and symptoms of HF defined by modified Framingham criteria^{11,12} and required intravenous drugs within 24 hours of hospital presentation. For those patients with repeated HF hospitalisation during the study period, the first HF hospitalisation was selected as an index hospitalisation. The overall design of the KCHF study has been described in detail.^{13–15}

All study procedures were conducted in compliance with the ethical principles of the Declaration of Helsinki. Clinical follow-up data were collected from hospital medical records or by contact with patients, their relatives or their referring physicians in October 2017.¹⁶

Definitions

Inotrope use was defined as treatment with dopamine, dobutamine or phosphodiesterase inhibitors within

24 hours of hospital presentation. Norepinephrine was categorised as a vasopressor. Levosimendan was not available in Japan and was not included in this study. According to the systolic blood pressure (sBP) and data on the HF profile including ‘warm’ (ie, adequately perfused) and ‘cold’ (hypoperfused) on presentation, we stratified all patients into two groups: sBP ≥ 90 mm Hg/warm group, and sBP < 90 mm Hg/cold group.

Outcomes

The primary outcome was all-cause death throughout a 1-year follow-up. Secondary outcomes included all-cause death during index hospitalisation and all-cause death after discharge, cardiovascular death throughout the 1-year follow-up, and HF hospitalisation. The causes of death were classified according to the Valve Academic Research Consortium definitions and adjudicated by a clinical event committee.¹⁷

Statistical analysis

Categorical variables are expressed as numbers and percentages and are compared using the χ^2 test. Continuous variables are expressed as means with SD or medians with IQRs and are compared using an analysis of variance or the Kruskal-Wallis test depending on their distributions. The cumulative incidences of the primary and secondary outcome measures, except for all-cause death during hospitalisation, were estimated by the Kaplan-Meier method and compared using the log-rank test. We estimated HRs and 95% CIs with Cox’s proportional hazards regression models. Proportional hazard assumptions for risk-adjusting variables were assessed on the plots of log (time) vs log [–log (survival)] stratified by the variable, and verified to be acceptable. In the Cox’s proportional hazards model, we simultaneously included the clinically relevant factors listed in table 1 as risk-adjusting variables and incorporated the centres as stratification variables. The risk for all-cause death during hospitalisation was assessed by a logistic regression model in which we constructed parsimonious models with the 11 clinically most relevant risk-adjusting variables listed in table 1 because of a small number of patients with outcome. Analyses of Cox’s proportional hazards model with stratification were performed with SPSS V.19 (IBM). All other analyses were performed with JMP V.12.01 software (SAS Institute) and GraphPad Prism V.6.05 (GraphPad Software, La Jolla, California, USA). In this study, all reported p values were two tailed, and $p < 0.05$ was considered to be significant.

Patient and public involvement

Patients or the public were not involved in the design, conduction, or reporting, or in the dissemination plans of our research.

RESULTS

Clinical characteristics

Among 3995 patients in this study (median age, 80 years; 1786 (45%) women), 121 (3%) had a sBP < 90 mm Hg,

Table 1 Characteristics of the patients with inotrope use and no inotrope use according to sBP and perfusion profile

	sBP ≥90 mm Hg and warm, N=3202		sBP <90 mm Hg or cold, N=793	
	No inotrope use, n=2890	Inotrope use, n=312	No inotrope use, n=517	Inotrope use, n=276
			P value	P value
Clinical characteristics				
Age, years	81 (73–86)	81 (71–87)	0.53	78 (67–85)
Age ≥80 years*†	1570 (54)	168 (54)	0.87	122 (44)
Female*†	1354 (47)	128 (41)	0.05	91 (33)
BMI, kg/m ²	23.0±4.5	22.2±4.4	<0.001	22.0±4.3
BMI <22 kg/m ² *†	1236 (43)	154 (49)	0.026	137 (50)
Aetiology				
Ischaemic	875 (30)	122 (39)	0.001	125 (45)
ACS*†	120 (4)	23 (7)	0.009	37 (13)
Cardiomyopathy	346 (12)	72 (23)	<0.001	82 (30)
Valvular	611 (21)	64 (21)	0.8	35 (13)
Medical history				
Prior hospitalisation for HF*†	954 (33)	127 (41)	0.006	126 (46)
Atrial fibrillation or flutter*†	1285 (44)	98 (31)	<0.001	85 (31)
Hypertension*†	2166 (75)	192 (62)	<0.001	147 (53)
Diabetes mellitus*	1059 (37)	117 (38)	0.77	121 (44)
Dyslipidaemia	1129 (39)	113 (36)	0.33	104 (38)
Prior myocardial infarction*	621 (21)	89 (29)	0.005	71 (26)
Prior stroke*	486 (17)	53 (17)	0.94	43 (16)
Prior PCI or CABG	722 (25)	82 (26)	0.62	79 (29)
Current smoking*	315 (11)	32 (10)	0.73	31 (11)
Ventricular tachycardia/fibrillation	74 (3)	23 (7)	<0.001	31 (11)
Chronic kidney disease	1260 (44)	151 (48)	0.1	137 (50)
Chronic lung disease*	395 (14)	40 (13)	0.68	33 (12)
Malignancy	435 (15)	48 (15)	0.88	30 (11)
Dementia	524 (18)	72 (23)	0.03	55 (20)
Social backgrounds				
Poor medical adherence	483 (17)	52 (17)	0.98	44 (16)
Living alone*	611 (21)	61 (20)	0.51	63 (23)
With occupation	335 (12)	51 (16)	0.01	40 (14)

Continued

Table 1 Continued

	sBP ≥90 mm Hg and warm, N=3202		sBP <90 mm Hg or cold, N=793	
	No inotrope use, n=2890	Inotrope use, n=312	No inotrope use, n=517	Inotrope use, n=276
Daily life activities			P value	P value
Ambulatory*	2252 (78)	235 (77)	0.67	0.36
Use of wheelchair (outdoor only)	218 (8)	29 (9)		210 (76)
Use of wheelchair (outdoor and indoor)	277 (10)	28 (9)		19 (7)
Bedridden	118 (4)	14 (5)		25 (9)
Vital signs at presentation				18 (7)
Systolic BP, mm Hg*†	152±33	134±27	<0.001	121±35
Systolic BP <90 mm Hg	0 (0)	0 (0)	N/A	52 (19)
Diastolic BP, mm Hg	86±23	79±20	<0.001	74±24
Heart rate, bpm	95±27	94±24	0.83	93±28
Heart rate <60 bpm*	191 (7)	18 (6)	0.57	27 (10)
Rhythms at presentation				0.005
Sinus rhythm	1569 (54)	201 (64)	<0.001	172 (62)
Atrial fibrillation/flutter	1121 (39)	81 (26)	<0.001	65 (24)
Tests at admission				
LVEF, %	49±16	38±16	<0.001	35±15
HFrEF (EF <40%)*†	888 (31)	193 (62)	<0.001	195 (71)
BNP, pg/mL†	653 (365–1138)	1039 (577–1864)	<0.001	1089 (498–1940)
NT-pro-BNP, pg/mL†	5417 (2577–11500)	9049 (3780–15852)	0.03	9240 (4390–22347)
Serum creatinine, mg/dL	1.1 (0.8–1.6)	1.3 (0.9–1.8)	0.046	1.3 (1.0–2)
eGFR, mL/min/1.73 m ²	45.1 (29.4–61.3)	38 (26.3–54.8)	<0.001	35.5 (24.4–53)
eGFR <30 mL/min/1.73 m ² †	748 (26)	99 (32)	0.03	106 (38)
Blood urea nitrogen, mg/dL	23 (17–34)	29 (20–45)	<0.001	30 (21–46)
Albumin, g/dL	3.5±0.5	3.4±0.5	0.01	3.4±0.5
Albumin <3.0 g/dL*	389 (13)	37 (12)	0.43	50 (18)
Sodium, mEq/L	139±4.2	138±4.3	<0.001	137±5.4
Sodium <135 mEq/L*	311 (11)	42 (13)	0.15	73 (26)
Potassium, mEq/dL	4.2±0.6	4.4±0.8	<0.001	4.5±0.8
Haemoglobin, g/L	114±23	117±24	0.019	121±24
Anaemia*†	1987 (69)	201 (64)	0.12	161 (58)

Continued

Table 1 Continued

	sBP ≥90 mm Hg and warm, N=3202		sBP <90 mm Hg or cold, N=793	
	No inotrope use, n=2890	Inotrope use, n=312	No inotrope use, n=517	Inotrope use, n=276
			P value	P value
HF-related signs and symptoms at presentation				
NYHA Class III or IV	2485 (86)	267 (86)	0.96	0.15
Cold (peripheral hypoperfusion)	0 (0)	0 (0)	N/A	0.07
Paroxysmal nocturnal dyspnoea	1957/2787 (70)	227/299 (76)	0.039	0.63
Orthopnoea	2222/2819 (79)	243/299 (81)	0.32	0.81
Dyspnoea on exertion	2664/2793 (95)	280/299 (94)	0.18	0.34
Rale	2201/2822 (78)	246/302 (81)	0.17	0.32
Oedema	2253/2832 (80)	230/304 (76)	0.11	0.7
Elevated jugular venous pressure	2142/2738 (78)	239/295 (81)	0.27	0.93
Pleural effusion (moderate/severe)	2507/2870 (87)	272/311 (87)	0.96	0.72
Lung congestion (moderate/severe)	2655/2866 (93)	291/311 (94)	0.55	0.049
Medications at presentation				
Loop diuretic	1375 (48)	169 (54)	0.03	0.1
ACEI/ARB	1334 (46)	122 (39)	0.017	0.49
β-blocker	1087 (38)	125 (40)	0.4	0.26
MRA	463 (16)	73 (23)	<0.001	0.001
Thiazide	169 (6)	28 (9)	0.03	0.4
Tolvaptan	85 (3)	21 (7)	<0.001	<0.001
Digoxin	191 (7)	19 (6)	0.72	0.94
Warfarin	624 (22)	66 (21)	0.86	0.08
DOAC	310 (11)	21 (7)	0.03	0.59
Intravenous therapy within 24 hours of presentation				
Loop diuretic	2512 (87)	250 (80)	<0.001	<0.001
Vasodilators	1603 (55)	138 (44)	<0.001	<0.001
Carperitide	1074 (37)	93 (30)	0.01	<0.001
Nitrates	833 (29)	56 (18)	<0.001	<0.001
Vasopressor (norepinephrine)	39 (1)	17 (5)	<0.001	<0.001
Inotropes	0 (0)	312 (100)	N/A	N/A
Dopamine	0	48 (15)	N/A	N/A
Dobutamine	0	263 (84)	N/A	N/A

Continued

Figure 1 Study flow chart. HF, heart failure; KCHF, Kyoto Congestive Heart Failure; sBP, systolic blood pressure.

patients with inotrope use was significantly higher than in the patients without inotrope use (figure 2A,B). After adjustments for confounders, however, the risk of inotrope use relative to no inotrope use was significant only in the sBP ≥ 90 mm Hg/warm group, whereas adjusted excess risk of inotrope use relative to no inotrope use was not significant in the sBP < 90 mm Hg/cold group (table 2). Likewise, despite a significantly higher cumulative 1 year incidence of cardiovascular death in the patients with inotrope use relative to no inotrope use in both groups (online supplemental figure 1A,B), the adjusted excess risk of inotrope use relative to no inotrope use for cardiovascular death from admission was only significant in the sBP ≥ 90 mm Hg/warm group but not in the sBP < 90 mm Hg/cold group (table 2). Even when use of norepinephrine was added to inotropes use as the inotrope/vasopressor use, adjusted excess risk of inotrope/vasopressor use relative to no inotrope/vasopressor use for the primary outcome measure was significant in the sBP ≥ 90 mm Hg/warm group (adjusted HR, 1.43, 95% CI 1.16 to 1.77, $p < 0.001$),

but not in the sBP < 90 mm Hg/cold group (adjusted HR, 1.30, 95% CI 0.98 to 1.72, $p = 0.07$).

In-hospital prognosis

During index hospitalisation, 154 patients (5%) in the sBP ≥ 90 mm Hg/warm group and 110 patients (14%) in the sBP < 90 mm Hg/cold group died. In both groups, the incidence of all-cause death during index hospitalisation was significantly higher in the patients with inotrope use than that in those with no inotrope use (figure 3A,B). The adjusted excess risk of inotrope use relative to no inotrope use for all-cause death during index hospitalisation was significant in both groups (table 2).

Postdischarge prognosis

A total of 3048 patients in the sBP ≥ 90 mm Hg/warm group and 683 patients in the sBP < 90 mm Hg/cold group were discharged alive. In the sBP ≥ 90 mm Hg/warm group, 669 patients (21%) died and 759 patients (24%) were hospitalised for HF during the follow-up period, whereas

Figure 2 Kaplan-Meier curves for all-cause death across inotrope use according to sBP and perfusion profile. (A) patients with sBP ≥ 90 mm Hg and warm; (B) patients with sBP < 90 mm Hg or cold. sBP, systolic blood pressure.

Table 2 Clinical outcomes across the patients with inotrope use and no inotrope use according to sBP and perfusion profile

Clinical outcomes	In-hospital or cumulative 1-year incidence (%)		Inotrope use versus no inotrope use			
	Inotrope use	No inotrope use	Unadjusted HR (95% CI)	P value	Adjusted HR* (95% CI)	P value
sBP ≥90 mm Hg and warm, n=3202						
Primary outcome measure						
All-cause death after admission	94 (31.1)	524 (18.6)	1.54 (1.25 to 1.87)	<0.001	1.36 (1.09 to 1.72)	0.006
Secondary outcome measures						
Cardiovascular death after admission	70 (24.2)	310 (11.3)	1.87 (1.46 to 2.37)	<0.001	1.49 (1.13 to 1.94)	0.004
In-hospital all-cause death	33 (10.6)	121 (4.1)	2.71 (1.78 to 4.01)	<0.001	1.98 (1.28 to 3.07)	0.002
Post-discharge all-cause death	61 (22.9)	421 (15.7)	1.32 (1.03 to 1.67)	0.03	1.25 (0.96 to 1.62)	0.10
HF hospitalisation	76 (30.3)	577 (22.7)	1.35 (1.07 to 1.68)	0.01	1.26 (0.99 to 1.61)	0.07
sBP <90 mm Hg or cold, n=793						
Primary outcome measure						
All-cause death after admission	101 (36.8)	130 (25.9)	1.42 (1.12 to 1.80)	0.004	1.28 (0.96 to 1.69)	0.09
Secondary outcome measures						
Cardiovascular death after admission	75 (28.9)	91 (18.7)	1.49 (1.13 to 1.97)	0.006	1.35 (0.96 to 1.88)	0.08
In-hospital all-cause death	63 (22.8)	47 (9.1)	2.96 (1.97 to 4.48)	<0.001	2.27 (1.45 to 3.57)	<0.001
Postdischarge all-cause death	39 (19.7)	88 (19.0)	0.93 (0.67 to 1.28)	0.68	0.88 (0.60 to 1.30)	0.52
HF hospitalisation	54 (27.7)	108 (25.5)	1.15 (0.84 to 1.54)	0.38	1.10 (0.76 to 1.58)	0.62

*Adjusted for the clinically relevant variables listed in [table 1](#).

HF, heart failure; sBP, systolic blood pressure.

172 patients (22%) died and 186 patients (24%) were hospitalised for HF in the sBP <90 mm Hg/cold group. In the sBP ≥90 mm Hg/warm group, cumulative 1-year incidences of all-cause death after discharge and HF hospitalisation were significantly higher in the patients with inotrope use ([figure 3C,E](#)), whereas there was negligible difference between the patients with inotrope use and no inotrope use in the sBP <90 mm Hg/cold group ([figure 3D,F](#)). After adjustment by confounders, excess risks of inotrope use relative to no inotrope use for all-cause death after discharge and HF hospitalisation were not significant in both groups ([table 2](#)).

DISCUSSION

This study from a large contemporary Japanese cohort demonstrated that 15% of the patients hospitalised with congestive HF received inotropes within 24 hours of hospital presentation. Fifty-three per cent of those patients did not have the guideline-recommended clinical indications of inotrope use such as sBP <90 mm Hg and a cold profile. Long-term mortality risk of initial inotrope use relative to no inotrope use was significant in the sBP ≥90 mm Hg/warm group, but not in the sBP <90 mm Hg/cold group. There were no differences in the risk for post-discharge all-cause death and HF hospitalisation between the inotrope use and no inotrope use in both

sBP ≥90 mm Hg/warm group and sBP <90 mm Hg/cold group.

In addition to diuretics and vasodilators, inotropes have long been used for the treatment of decompensated HF, particularly for patients with cardiogenic shock or signs of impaired perfusion. However, previous studies demonstrated that the use of inotropes in the treatment of acute decompensated HF was associated with short-term adverse outcomes.^{6–8 18} Thus, US and European guidelines restrict the use of inotropic agents unless the patients have symptomatic hypotension or signs of peripheral hypoperfusion.¹² Despite negative implications from the previous observational studies and guideline restrictions, our study demonstrated that inotrope use is still common in the current real world of hospitalised HF patients with 15% of the study participants receiving inotropes within 24 hours of hospital presentation. Notably, more than half of the patients initially treated with inotropes did not exhibit hypotension or signs of peripheral hypoperfusion on hospital presentation. The results were similar to those from the European Society of Cardiology Heart Failure Long-Term registry in which 12% of the study patients received inotropes and/or vasopressors during the first 24 hours after admission and 45.7% of these patients exhibited signs of peripheral hypoperfusion.¹⁰ In our study, crude 1-year mortality risk of the patients initially treated with inotropes was 31.1% in the sBP ≥90 mm Hg/warm group

Figure 3 Incidence of all-cause death during index hospitalisation (A, B), all-cause death after discharge (C, D) and HF hospitalisation (E, F). A, C and E, patients with sBP ≥ 90 mm Hg and warm; B, D and F, patients with sBP <90 mm Hg or cold. HF, heart failure; sBP, systolic blood pressure.

and 36.8% in the sBP <90 mm Hg/cold group, both of which were much higher than that of the patients without inotrope use. However, inotrope use had a different effect size for long-term mortality across the two groups. Adjusted excess risk of inotrope use relative to no inotrope use for 1-year mortality was significant among the patients without hypotension or

signs of impaired perfusion, whereas it was neutral among the patients presenting with hypotension or a cold profile. The results were similar with the previous reports by Kang *et al* showing that inotrope use was an independent predictor for in-hospital and 1-month postdischarge mortality in patients with initial sBP ≥ 90 mm Hg, but not in patients with an initial

sBP <90 mm Hg.¹⁹ However, the categorisation of our patients into the sBP <90 mm Hg or cold group was mainly driven by 'cold' profile rather than sBP <90 mm Hg as more than 90% of this group had cold profile, whereas 85% had sBP ≥90 mm Hg. Given the limited options for medical management of decreased cardiac output and impaired peripheral organ perfusion, inotrope use might still be an option for patients with a cold profile. On the other hand, our results might support inappropriateness of inotrope use for haemodynamically stable patients with adequate peripheral perfusion, enforcing the current guidelines that discourage the routine use of inotropes. However, it should be noted that the present results should be interpreted carefully. Although the present results come from a large contemporary HF cohort, they provide only a snapshot of current real-world clinical practice. Its observational design per se precludes the demonstration of a causal relationship between initial inotropic treatment and outcomes. We cannot deny the possibility that the increased mortality associated with the use of inotropes may be attributed to patient characteristics rather than drug-specific characteristics.

A previous trial comparing the outcomes of dobutamine with nesiritide among patients with acute decompensated HF showed that even short-term infusion of dobutamine within 3 days resulted in a significantly higher 6-month mortality compared with nesiritide.²⁰ Similarly, a 24-hour infusion of dobutamine was associated with a significantly higher 6-month mortality compared with levosimendan. These results suggest that even short-term exposure to catecholamines could affect long-term prognosis.²¹ Irreversible catecholamine-induced damage to cardiac myocytes has been suggested as an underlying mechanism.²¹ However, we found that a high 1-year mortality risk in the inotrope stratum was mainly driven by a higher in-hospital mortality, whereas the risks of postdischarge mortality and HF hospitalisation were comparable between inotrope stratum and no inotrope stratum in both the sBP ≥90 mm Hg/cold and sBP <90 mm Hg/warm groups. These results were in line with a recent observational study reporting a lack of association between inotrope use and 6-month post-discharge mortality in patients with reduced EF.²² The results may suggest that adverse effects potentially induced by inotropes such as direct myocardial toxicity, increased oxygen consumption and arrhythmogenesis are temporal and do not last a long time. The results highlight unmet needs for the development of a more effective and safe treatment strategy targeting the initial phase of HF to overcome the imminent critical condition seen in our patients in the inotrope stratum.

Limitations

This study had several limitations. First, because of its observational nature, the decision to use vasoactive agents was made by the attending physicians, which is inherently subject to selection bias and confounding. Second, although detailed data in our dataset enabled intensive adjustment with most conceivable and previously established confounders, we were unable to exclude the influence of other unmeasured confounding factors.

In addition, for the evaluation of all-cause death during index hospitalisation, we used parsimonious models because of a small number of patients with outcome. Third, the exact dose of agents and the duration of their use were not available in our dataset. Thus, we were unable to distinguish low- and high-dose infusions of inotropes. Finally, although this study included nearly 4000 patients, the sample size was still insufficient to separately analyse the impact of individual drugs categorised as inotropes.

In conclusion, this study demonstrated the current use of inotropes in initial in-hospital treatment of HF in real-world practice. Inotropes are still commonly used against guideline recommendations. Inotrope treatment is associated with a high long-term mortality risk, particularly when used in the absence of shock or peripheral hypoperfusion. Development of standardised and evidence-based approaches to initial HF treatment might improve clinical outcomes.

Author affiliations

¹Department of Cardiology, Osaka Red Cross Hospital, Osaka, Japan

²Department of Cardiovascular Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan

³Department of Cardiology, Mitsubishi Kyoto Hospital, Kyoto, Japan

⁴Department of Clinical Epidemiology, Hyogo College of Medicine, Nishinomiya, Japan

⁵Department of Cardiovascular Medicine, Shiga General Hospital, Moriyama, Japan

⁶Department of Cardiology, Tenri Hospital, Tenri, Japan

⁷Division of Heart Failure, National Cerebral and Cardiovascular Center, Suita, Japan

⁸Department of Cardiology, Hyogo Prefectural Amagasaki General Medical Center, Amagasaki, Japan

⁹Department of Cardiology, National Hospital Organization Kyoto Medical Center, Kyoto, Japan

¹⁰Department of Cardiology, Shimabara Hospital, Kyoto, Japan

¹¹Department of Cardiology, Japanese Red Cross Otsu Hospital, Otsu, Japan

¹²Department of Cardiology, Hikone Municipal Hospital, Hikone, Japan

¹³Department of Cardiology, Kishiwada City Hospital, Kishiwada, Japan

¹⁴Department of Cardiology, Kansai Electric Power Hospital, Osaka, Japan

¹⁵Department of Cardiology, Shizuoka General Hospital, Shizuoka, Japan

¹⁶Department of Cardiology, Kurashiki Central Hospital, Kurashiki, Japan

¹⁷Department of Cardiology, Kokura Memorial Hospital, Kitakyushu, Japan

¹⁸Department of Cardiology, Japanese Red Cross Wakayama Medical Center, Wakayama, Japan

¹⁹Department of Cardiology, Nishi Kobe Medical Center, Kobe, Japan

²⁰Cardiovascular Center, Tazuke Kofukai Medical Research Institute, Kitano Hospital, Osaka, Japan

²¹Department of Cardiovascular Medicine, Kobe City Medical Center General Hospital, Kobe, Japan

²²Department of Cardiovascular Medicine, Shiga University of Medical Science, Otsu, Japan

²³Department of Cardiovascular Medicine, Shinshu University Graduate School of Medicine, Matsumoto, Japan

Acknowledgements The authors thank the staff of the KCHF study and the members of the participating centres.

Contributors KN: conceptualisation, methodology, formal analysis, writing-original draft; TKato: conceptualisation, methodology, data curation, funding acquisition, interpretation, review, writing-review and editing, final approval, agreement to be accountable; HY, EY: data curation, investigation, resources; TMorimoto: methodology, formal analysis, writing-review and editing, supervision; YI, YT, YY, TKit, RT, Mlg, MaK, MTa, TJ, Tlk, TKaw, AK, RN, YK, TMorinaga, KS, MiK, YSe, Tln, Mln, MTo, YF, YN, KA, KKa, SS, KO, YSa: investigation, resources; KKu, NO: investigation, resources, funding acquisition; TKim: project administration, methodology, supervision, writing-review and editing. TKato is the guarantor.

Funding This study is supported by a grant from the Japan Agency for Medical Research and Development (18059186) to TK, KK and NO.

Disclaimer The funder had no role in the study design, collection, analysis, or interpretation of data, writing the manuscript, or the decision to submit the paper for publication.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study was approved by Ethics Committees and reference number: Kyoto University Graduate School of Medicine (approval number: E2311), Shiga General Hospital (approval number: 20141120-01), Tenri Hospital (approval number: 640), Kobe City Medical Center General Hospital (approval number: 14094), Hyogo Prefectural Amagasaki General Medical Center (approval number: Rinri 26-32), National Hospital Organisation Kyoto Medical Center (approval number: 14-080), Mitsubishi Kyoto Hospital (approved November 12, 2014), Okamoto Memorial Hospital (approval number: 201503), Japanese Red Cross Otsu Hospital (approval number: 318), Hikone Municipal Hospital (approval number: 26-17), Japanese Red Cross Osaka Hospital (approval number: 392), Shimabara Hospital (approval number: E2311), Kishiwada City Hospital (approval number: 12), Kansai Electric Power Hospital (approval number: 26-59), Shizuoka General Hospital (approval number: Rin14-11-47), Kurashiki Central Hospital (approval number: 1719), Kokura Memorial Hospital (approval number: 14111202), Kitano Hospital (approval number: P14-11-012), Japanese Red Cross Wakayama Medical Center (approval number: 328).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Kazuya Nagao <http://orcid.org/0000-0002-5043-8778>
Takao Kato <http://orcid.org/0000-0001-8213-7999>
Takeshi Morimoto <http://orcid.org/0000-0002-6844-739X>
Moritake Iguchi <http://orcid.org/0000-0001-5029-1203>
Satoshi Shizuta <http://orcid.org/0000-0003-4587-9031>
Takeshi Kimura <http://orcid.org/0000-0002-5665-4076>

REFERENCES

- 1 Ponikowski P, Voors AA, Anker SD, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2016;37:2129–200.
- 2 Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation* 2013;128:1810–52.
- 3 O'Connor CM, Starling RC, Hernandez AF, et al. Effect of nesiritide in patients with acute decompensated heart failure. *N Engl J Med* 2011;365:32–43.
- 4 Tsutsui H, Isobe M, Ito H, et al. JCS 2017/JHFS 2017 Guideline on diagnosis and treatment of acute and chronic heart failure- Digest Version. *Circ J* 2019;83:2084–184.
- 5 Mullens W, Damman K, Harjola V-P, et al. The use of diuretics in heart failure with congestion - a position statement from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail* 2019;21:137–55.
- 6 O'Connor CM, Gattis WA, Uretsky BF, et al. Continuous intravenous dobutamine is associated with an increased risk of death in patients with advanced heart failure: insights from the Flolan international randomized survival trial (first). *Am Heart J* 1999;138:78–86.
- 7 Mebazaa A, Parissis J, Porcher R, et al. Short-term survival by treatment among patients hospitalized with acute heart failure: the global ALARM-HF registry using propensity scoring methods. *Intensive Care Med* 2011;37:290–301.
- 8 Abraham WT, Adams KF, Fonarow GC, et al. In-hospital mortality in patients with acute decompensated heart failure requiring intravenous vasoactive medications: an analysis from the acute decompensated heart failure national registry (adhere). *J Am Coll Cardiol* 2005;46:57–64.
- 9 Francis GS, Bartos JA, Adatya S. Inotropes. *J Am Coll Cardiol* 2014;63:2069–78.
- 10 Mebazaa A, Motiejunaite J, Gayat E, et al. Long-term safety of intravenous cardiovascular agents in acute heart failure: results from the European Society of cardiology heart failure long-term registry. *Eur J Heart Fail* 2018;20:332–41.
- 11 McKee PA, Castelli WP, McNamara PM, et al. The natural history of congestive heart failure: the Framingham study. *N Engl J Med* 1971;285:1441–6.
- 12 Senni M, Tribouilloy CM, Rodeheffer RJ, et al. Congestive heart failure in the community: a study of all incident cases in Olmsted County, Minnesota, in 1991. *Circulation* 1998;98:2282–9.
- 13 Yamamoto E, Kato T, Ozasa N, et al. Kyoto congestive heart failure (KCHF) study: rationale and design. *ESC Heart Fail* 2017;4:216–23.
- 14 Yaku H, Ozasa N, Morimoto T, et al. Demographics, management, and in-hospital outcome of hospitalized acute heart failure syndrome patients in contemporary real clinical practice in Japan- Observations from the prospective, multicenter Kyoto Congestive Heart Failure (KCHF) Registry. *Circ J* 2018;82:2811–9.
- 15 Yaku H, Kato T, Morimoto T, et al. Association of mineralocorticoid receptor antagonist use with all-cause mortality and hospital readmission in older adults with acute decompensated heart failure. *JAMA Netw Open* 2019;2:e195892.
- 16 Japan's ethical guidelines for epidemiologic research. 2008, 2002. Available: http://www.lifescience.mext.go.jp/files/pdf/n796_01.pdf
- 17 Kappetein AP, Head SJ, Génèreux P, et al. Updated standardized endpoint definitions for transcatheter aortic valve implantation: the valve academic research Consortium-2 consensus document. *J Am Coll Cardiol* 2012;60:1438–54.
- 18 Elkayam U, Tasissa G, Binanay C, et al. Use and impact of inotropes and vasodilator therapy in hospitalized patients with severe heart failure. *Am Heart J* 2007;153:98–104.
- 19 Kang J, Cho H-J, Lee H-Y, et al. Effects of widespread inotrope use in acute heart failure patients. *J Clin Med* 2018;7:368. doi:10.3390/jcm7100368
- 20 Silver MA, Horton DP, Ghali JK, et al. Effect of nesiritide versus dobutamine on short-term outcomes in the treatment of patients with acutely decompensated heart failure. *J Am Coll Cardiol* 2002;39:798–803.
- 21 Follath F, Cleland JGF, Just H, et al. Efficacy and safety of intravenous levosimendan compared with dobutamine in severe low-output heart failure (the LIDO study): a randomised double-blind trial. *Lancet* 2002;360:196–202.
- 22 Gomes C, Terhoch CB, Ayub-Ferreira SM, et al. Prognosis and risk stratification in patients with decompensated heart failure receiving inotropic therapy. *Open Heart* 2018;5:e000923.