

BMJ Open Preventing chronic disease in overweight and obese patients with low health literacy using eHealth and teamwork in primary healthcare (HeLP-GP): a cluster randomised controlled trial

Sharon M Parker ¹, Margo Barr ¹, Nigel Stocks ², Elizabeth Denney-Wilson,³ Nicholas Zwar,⁴ Jon Karnon,⁵ Alamgir Kabir ¹, Don Nutbeam,⁶ Jackie Roseleur ⁵, Siaw-Teng Liaw ⁷, Carmel McNamara,⁸ Oliver Frank,² An Tran,¹ Richard Osborne,⁹ Annie Y S Lau,¹⁰ Mark Harris ¹

To cite: Parker SM, Barr M, Stocks N, *et al*. Preventing chronic disease in overweight and obese patients with low health literacy using eHealth and teamwork in primary healthcare (HeLP-GP): a cluster randomised controlled trial. *BMJ Open* 2022;**12**:e060393. doi:10.1136/bmjopen-2021-060393

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-060393>).

Received 22 December 2021
Accepted 13 November 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Mark Harris;
m.f.harris@unsw.edu.au

ABSTRACT

Objectives To evaluate a multifaceted intervention on diet, physical activity and health literacy of overweight and obese patients attending primary care.

Design A pragmatic two-arm cluster randomised controlled trial.

Setting Urban general practices in lower socioeconomic areas in Sydney and Adelaide.

Participants We aimed to recruit 800 patients in each arm. Baseline assessment was completed by 215 patients (120 intervention and 95 control).

Intervention A practice nurse-led preventive health check, a mobile application and telephone coaching.

Primary and secondary outcome measures Primary outcomes were measured at baseline, 6 and 12 months, and included patient health and eHealth literacy, weight, waist circumference and blood pressure. Secondary outcomes included changes in diet and physical activity, preventive advice and referral, blood lipids, quality of life and costs. Univariate and multivariate analyses of difference-in-differences (DiD) estimates for each outcome were conducted.

Results At 6 months, the intervention group, compared with the control group, demonstrated a greater increase in Health Literacy Questionnaire domain 8 score (ability to find good health information; mean DiD 0.22; 95% CI 0.01 to 0.44). There were similar differences for domain 9 score (understanding health information well enough to know what to do) among patients below the median at baseline. Differences were reduced and non-statistically significant at 12 months. There was a small improvement in diet scores at 6 months (DiD 0.98 (0.50 to 1.47); $p=0.026$) but not at 12 months. There were no differences in eHealth literacy, physical activity scores, body mass index, weight, waist circumference or blood pressure.

Conclusions Targeted recruitment and engagement were challenging in this population. While the intervention was associated with some improvements in health literacy and diet, substantial differences in other outcomes were not

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The cluster randomised design allowed testing of the nurse-led intervention among patients without contamination.
- ⇒ Recruitment of practices and patients did not meet our planned sample size.
- ⇒ We noted variable uptake of the intervention components among patients reflecting real-world general practice.
- ⇒ The measures used to assess health literacy, diet and physical activity had some limitations.
- ⇒ The study was conducted in only two urban areas of Australia and the findings may not therefore be generalised to other communities, such as rural areas.

observed. More intensive interventions and using codesign strategies to engage the practices earlier may produce a different result. Codesign may also be valuable when targeting lower socioeconomic populations.

Trial registration number Australian New Zealand Clinical Trials Registry (ACTRN 12617001508369) (<http://www.ANZCTR.org.au/ACTRN12617001508369.aspx>).

Trial protocol The protocol for this trial has been published (open access; <https://bmjopen.bmj.com/content/8/6/e023239>).

INTRODUCTION

Obesity is a complex health issue and is influenced by biological, environmental, social and psychological factors.¹ Overweight and obesity account for 8.4% of the burden of disease being a risk factor for 11 types of cancer, 3 cardiovascular conditions, chronic kidney disease, diabetes, dementia, gall-bladder disease, fatty liver, gout, back pain and osteoarthritis.² In 2017/2018, 67% of the



Australian population were overweight (body mass index (BMI) 25–29 kg/m²; 35.6%) or obese (BMI 30+ kg/m²; 31.3%) with those who were more socially disadvantaged being more likely to be overweight or obese.³ Within Australia, rates of overweight and obesity peak for men at age 55–64 years (83.6%) and for women at 65–74 years (73.3%).⁴

Current Australian guidelines recommend that people who are overweight and obese attending general practice undergo routine measurements (BMI and waist circumference) and are engaged in discussions about lifestyle risk factors and positive messaging to improve health and well-being.⁵ Behavioural interventions in primary care have been demonstrated to achieve a 5%–7% improvement in weight, blood pressure (BP) or lipids for patients, potentially preventing or delaying the onset of type 2 diabetes and cardiovascular disease.⁶ A recent systematic review and meta-analysis supports weight loss programmes delivered by primary care practitioners as they provide effective weight loss and reduction in waist circumference.⁷ Multicomponent intensive behavioural interventions (delivered by various clinicians and provided through group, individual, technology or print-based methods) have been recommended for patients with a BMI of 30 or higher.⁸ Health coaching provided by a trained professional has become a popular tool to address weight through behaviour change strategies⁹ and high-intensity behavioural counselling (12 or more sessions per year delivered in person, by phone or electronically) is accepted to produce clinically meaningful weight loss.¹⁰

The Track Study¹¹ which combined tailored weight-related behaviour change goals for patients as a basis for self-monitoring with 18 coaching calls over 12 months found intervention patients significantly more likely to lose ≥5% of their baseline weight at 6 months and 12 months. A recent retrospective analysis of 25 000 people receiving blended care behaviour change interventions (a combination of digital care and coaching)¹² supports the use of these interventions for weight loss but highlights the need for more understanding as to which elements would be best delivered by health coaches and which can be delegated to a digital device.

Patients generally accept their general practitioners' (GPs) role in management of overweight and obesity¹³; however, lower socioeconomic groups tend to be less likely to take up weight management programmes.^{14 15} Low functional health literacy (ie, health-related reading and numeracy) is more common in socioeconomically disadvantaged populations and is associated with an increased likelihood of overweight and obesity.^{16 17} It is also a potential barrier to the uptake and effectiveness of a range of preventive interventions that mediate change in lifestyle behaviours.^{18 19} Patients with low health literacy are less likely to engage in health-promoting behaviours^{20–22} and attend or complete programmes to which they have been referred.^{23 24} Interventions with multiple components to improve health literacy for behavioural risk factors have

been shown to be more effective at improving nutritional health literacy in primary care than those with single components.⁶ Other barriers to delivering weight loss management have also been identified, including low confidence levels of clinicians in obesity management,²⁵ stigmatisation of patients²⁶ and lost opportunities by providers to initiate earlier, effective weight loss conversations.²⁷

OBJECTIVES

The HeLP-GP trial aimed to evaluate a multifaceted intervention provided to overweight and obese patients attending primary care. The primary hypothesis was that the intervention would lead to improved health literacy, eHealth literacy, physiological risk factors, lifestyle behaviours and quality of life.

METHODS

Trial design

A pragmatic, two-arm, unblinded cluster randomised controlled trial. This design was chosen to provide protection against contamination within sites (general practices) as practice staff were providing the intervention. Primary and secondary outcomes were assessed at the patient level.

Participants and setting

The trial was conducted in general practices located in metropolitan and urban fringe areas of south-western and western Sydney in New South Wales and Adelaide in South Australia. Practice eligibility included:

- ▶ Geographical location in Local Government Areas with a Socio-Economic Index for Area Index of Relative Socio-economic Disadvantage²⁸ equal to or below the eighth decile.
- ▶ Using clinical software compatible with the trial data extraction and recruitment tool, *Doctors Control Panel* (DCP),²⁹ and an active internet connection.
- ▶ Participation by at least one practice nurse (PN) and one GP from the practice.
- ▶ Participation of reception staff to distribute trial materials to eligible trial participants as they present for appointments.

Patient eligibility included:

- ▶ Aged 40–74 years.
- ▶ BMI ≥28 recorded within the previous 12 months (the cut-off point for BMI was chosen to target people at higher risk and to capture people from Asian backgrounds who have a lower equivalent BMI).
- ▶ BP and total serum cholesterol recorded within the previous 12 months.
- ▶ Speaking English and/or Arabic, Vietnamese or Chinese (languages representing common migrant groups in the catchment areas—there were very few patients who spoke other languages but not English).

- ▶ Access to a smartphone or tablet device and internet connection.
- Patients were excluded if they:
- ▶ Had a diagnosis of diabetes requiring insulin or a current prescription for insulin, a diagnosis of cardiovascular disease (angina, myocardial infarction, heart failure, heart valve disease (rheumatic or non-rheumatic)), stroke (cerebrovascular accident).
 - ▶ Had experienced weight loss of >5% in the past 3 months, were taking medication for weight loss (orlistat or phentermine) or had undergone weight loss surgery.
 - ▶ Had cognitive impairment (including serious mental illness).
 - ▶ Had a physical impairment which would prohibit engaging in moderate-level physical activity.

Practice recruitment

Between March 2018 and October 2018, general practices within the specified geographical locations were approached by partner Primary Health Networks (PHNs), which are regional organisations providing quality improvement and education to general practices. Invitations to express interest were distributed through mail, email, newsletters, GP educational events, websites, Facebook groups for health professionals, discussion groups and research networks. A face-to-face meeting was held between responding practices, a PHN representative and a member of the research team to discuss participation in detail and confirm eligibility.

Randomisation

Randomisation of practices was performed by an epidemiologist (MB) who was not involved in the data collection or intervention using the SAS³⁰ v.9.4 statistical package. Practices were characterised by size (fewer than five GPs, or five or more GPs) and by state into four strata, and intervention and control lists of random numbers (six-digit) were generated for each stratum. The resultant intervention and control strata lists were combined and sorted. Four batches were created. Allocation of intervention or control was then sequentially allocated from the lists based on the date of entry of the practice into the study by an independent researcher. Batching was undertaken to ensure similar numbers of control and intervention practices at any point in time. Practices were informed in writing as to what allocation they had received.

Recruitment of patients

From October 2018 to September 2019, patients of participating practices were flagged at the point of presentation using DCP. The software was programmed with clinical inclusion/exclusion criteria to identify potential participants as they presented. Once flagged, patient information was automatically printed and attached to trial information and consent forms by the reception staff. It was not the responsibility of GPs to gain consent, but

patients could discuss the trial with their GP or PN. As DCP was only able to determine eligibility based on the information within the practice's clinical software, eligibility was also checked by a member of the practice. Patients could return their consent forms by leaving them in a secure collection point at the practice or returning them in a reply-paid envelope to the study centre (University of New South Wales, Sydney).

The HeLP-GP intervention

The intervention was a multicomponent intervention, which has been previously described and piloted.^{31 32} It aimed to increase the knowledge of patients relating to diet and physical activity and their individual skills to address weight management behaviours. It comprised:

1. A PN-led health check designed to support Australian Guidelines for the management of overweight and obesity^{5 33} and based on the 5As (Assess, Advise, Agree, Assist and Arrange).^{34 35} Review was conducted by the PN at 6 weeks and the GP at 12 weeks.
2. A lifestyle app (*mynsnapp*) modified from *healthy.me*, a personally controlled health management platform designed to help patients and consumers to manage their health.³⁶ The components of *mynsnapp* were informed by research into behaviour change through mobile and electronic platforms that suggest that goal setting and self-monitoring, and additional methods to interact with patients, particularly text messaging, can be more effective than advice alone.^{37–40} *Mynsnapp* allowed patients to set and revise physical activity and diet-based goals and to view graphs of their progress over the previous 6 weeks. A free-text diary allowed patients to document individualised content. A range of video and written resources related to diet and physical activity, linked to the app, were available for the patient to view. Text messages reminded patients to attend the follow-up with the PN and GP and once registered, each patient received one nutrition and one physical activity message each week for 6 weeks.³²
3. Health coaching via the 'Get Healthy' telephone coaching programme (<https://www.gethealthynsw.com.au/>) provided free, confidential telephone-based health coaching to support patients to reach personalised lifestyle goals relating to healthy eating, increasing physical activity, alcohol reduction and achieving and maintaining a healthy weight. Coaching was available in multiple languages with the assistance of an interpreter service.

At the health check, patients could choose to take up *mynsnapp*, Get Healthy or both. Control practices provided 'usual care' (the clinical practice routinely offered to patients by the GP and PN of the practice).

Training and implementation of the intervention

Training was completed by all participating PNs. Training comprised three online modules covering physical assessment (weight, height, BP, waist circumference and BMI), delivery of relevant lifestyle advice and promotion



of individual goal setting. The 'teach-back' method⁴¹ (asking the patient to repeat in their own words what they have understood) was encouraged to ensure they had understood and were confident with the content of the health check. PNs assisted patients to download and set up *mynsnapp* including setting goals during the health check and were encouraged to review the patient's use of the app and the progress of health coaching at the 6-week follow-up. Written and video resources were developed for PNs and patients on the installation and use of the app. PNs referred patients to Get Healthy using a trial-specific online referral form.

Patients could claim Medicare benefits (usually without out-of-pocket payments) for GP visits as part of the intervention (Medicare is Australia's national universal health insurance scheme). Patients did not pay for the PN visits. The PN health checks were reimbursed directly to the practice by the study at a rate of \$A40 per patient for the health check and \$A20 per patient for follow-up.

Patient and public involvement

Patients and members of the public were not involved in the design of this study. Consumer volunteers with the Adelaide PHN did pilot the lifestyle app (*mynsnapp*) and provide input to its final design.

Data collection and trial outcomes

The methods are described in the protocol paper.³² Table 1 provides a summary of the data collected to assess trial outcomes, the collection method and the time points of collection. A proposed 18-month follow-up of patients was abandoned due to the need to extend the period for patient recruitment and lower than expected numbers of patients being recruited to the trial. Surveys administered over the telephone were used to collect demographic and other patient data.

Primary outcomes

We used two domains of the Health Literacy Questionnaire (HLQ) (domain 8: ability to find good health information (five items) and domain 9: understand health information well enough to know what to do (five items)).⁴² The individual domains of the HLQ can be selected to identify specific health literacy strengths and challenges or to test a hypothesis.^{43 44} Domains 8 and 9 have a 5-point response option scale (cannot do or always difficult, usually difficult, sometimes difficult, usually easy or always easy). The scores for these domains are averages for the domain (with a range between 1 and 5). The electronic Health Literacy Scale (eHEALS) was used to assess digital health literacy.⁴⁵

DCP was used to extract clinical patient data related to biomedical risk factors (BMI, systolic and diastolic BP, and waist circumference). We used the measurements recorded by the GP at the nearest time point to follow-up (baseline and 12-month follow-up interviews).

Secondary outcomes

Patient self-report was used to determine lifestyle behaviours including a diet score (portions of fruit (between 0 and a maximum of 2 per day) plus portions of vegetable intake (between 0 and a maximum of 5 per day) with a range between 0 and 7 based on the sum of fruit and vegetable scores), the number of 30-minute sessions of physical activity (moderate/vigorous) per week and changes in diet and physical activity. Questions to assess these behaviours were adapted from previous research.^{46 47} The scores for diet were between 1 and 7.

Patient self-report was used to determine advice received and referral for diet, physical activity and weight loss. Patient questions also assessed quality of life (using the EQ-5D-5L standardised to UK reference population with no imputation of missing values).^{48 49} Total cholesterol, low-density lipoprotein (LDL), high-density lipoprotein (HDL) and triglyceride (TG) values were extracted by the DCP from the GP medical record at baseline and 12-month follow-up.

Sample size calculation

The original sample size calculation of 400 in each arm was based on the primary hypothesis that the intervention would lead to improved health and eHealth literacy, diet, physical activity, weight and BP. This was based on assumption of hypothesised effect sizes described in the trial protocol.³² Sample size estimates were based on a two-sided test of significance at $\alpha=0.05$, $1-\beta=0.8$ and 20% loss to follow-up. For HLQ, the anticipated mean difference was 0.4 for both domains 8 and 9 (based on domain 8 mean 3.7 (SD=0.9) and domain 9 mean 3.9 (SD 0.8)). For BMI and systolic BP, the effect sizes were 0.2, respectively (based on means of 30 (SD 6) and 131 (SD 15), respectively).

ANALYSIS

Statistical analyses were conducted on the intention-to-treat (ITT) population for both primary and secondary outcomes. The ITT population was defined as all those recruited at baseline regardless of what intervention they received and what follow-up data were available.

Summary participant baseline characteristics and primary outcomes at baseline were compared between control and intervention groups using either X^2 test, t-test or Mann-Whitney test. Means and SDs were reported for continuous outcomes and the number and percentage were reported for dichotomous outcomes at baseline, 6-month and 12-month follow-up.

To measure the effect of the intervention on the outcomes of interest (primary or secondary), we used difference-in-differences (DiD) estimate as some of the outcomes at baseline were significantly different.⁵⁰ We used generalised estimating equation (GEE) with Gaussian family and identity link function to estimate DiD accounting for the cluster (general practice)-level correlation.⁵¹ We put an interaction term for

Table 1 Patient-level outcomes

Outcome	Instrument/contributing data	Primary or secondary outcome	Data collection method	Time point for collection		
				BL	6 months	12 months
Literacy and eHealth literacy						
Health literacy	HLLQ (domains 8 and 9)	Primary	Patient survey—administered via telephone interview	x	x	x
eHealth literacy	eHEALS	Primary	Patient survey—administered via telephone interview	x	x	x
Biomedical risk factors (patient)						
Weight/height/waist circumference/BMI	Clinical record	Primary	DCP	x	–	x
Blood pressure	Clinical record	Primary	DCP	x	–	x
Lipids (total cholesterol)	Clinical record	Secondary	DCP	x	–	x
Lifestyle risk factors (patient)						
Fruit and vegetable intake	Patient self-report—servings of fruits and vegetables per day	Secondary	Patient survey—administered via telephone interview	x	x	x
Level of physical activity	Patient self-report (moderate and vigorous physical activity per week)	Secondary	Patient survey—administered via telephone interview	x	x	x
Quality of life						
QOL	EQ-5D-5L	Secondary	Patient survey—administered via telephone interview	x	–	x
Advice and referral						
Recall of advice and goal setting for diet, physical activity, weight loss	Patient survey	Secondary	Patient survey—administered via telephone interview	x	x	–
Referral to behaviour change programmes for diet, physical activity or weight loss	Patient survey	Secondary	Patient survey—administered via telephone interview	x	x	–
Economic data						
Delivery cost of intervention	Study documentation/budget	Secondary	Study administrative records/facilitator diary	Calculated for trial costs (payments for health checks, practice staff education and practice facilitation; cost of the app and telephone coaching)		
Health service costs	Medicare Benefits Scheme data	Secondary	Output from Services Australia ⁶⁵	Data collected 01 October 2017–30 June 2020		
Prescription medication	Pharmaceutical Benefits Schedule data	Secondary	Output from Services Australia ⁶⁵	Data collected 01 October 2017–30 June 2020		
BL, baseline; BMI, body mass index; DCP, Doctors Control Panel; eHEALS, electronic Health Literacy Scale; HLLQ, Health Literacy Questionnaire.						

intervention group and a dummy variable for before/after the follow-up measurement (6-month or 12-month follow-up) in the GEE model and the coefficient of the interaction term was considered as a DiD estimate.⁵² Separate models were used for estimating DiD at 6-month and 12-month follow-up. The DiD estimates were adjusted for the potential confounders which were substantially different between control and intervention groups at baseline. To adjust for possible ceiling effects, we did stratified analysis for the health literacy scores by above or below the median score at baseline. We set 5% as a level of statistical significance. We used the R V4.0.3 programming language and environment for the statistical analysis.⁵³

Economic evaluation

The extracted cost data informed a cost consequence analysis, undertaken from the Australian healthcare system perspective. We categorised costs as follows: (1) services provided or requested by GPs (excluding consultations by specialists), (2) services provided or requested by GPs or specialists (excluding services related to surgical procedures), and (3) pharmaceutical costs. The number of times participants visited a GP was also analysed. Costs and number of GP visits were calculated for the 12 months preceding and the 12 months following the enrolment date for each participant, from which unadjusted DiD estimates were derived for each of the cost categories, as well as aggregate costs and GP visits. Bootstrapping (using 1000 resamples) was used to represent the uncertainty around the DiD estimates.

RESULTS

We used the Consolidated Standards of Reporting Trials extension for cluster trials statement to guide reporting (online supplemental file 1) and summarise the flow of participants (figure 1) through the HeLP-GP trial.⁵⁴

Baseline

We recruited 215 participants to the study (120 to the intervention group and 95 to the control group) through 22 practices (clusters). Baseline characteristics of the intervention group were similar to the control group except that the proportion of men was higher (66.3% vs 50.0%). Participants in both groups were predominantly aged between 46 and 65 years, with over one-third having been born overseas (mostly from Europe or Asia) but only one-third of those born overseas had arrived in Australia in the past 10 years and one in six of all participants spoke a language other than English. A total of 39.5% had school qualifications only and 59% were employed. The median BMI was 33.3 kg/m². The intervention outcome measures at baseline were all similar to the control group except for health literacy which was lower (mean 4.0 vs 4.3 for domain 8, and 4.1 vs 4.3 for domain 9) (table 2).

Intervention uptake

There was variable uptake of the intervention components by the 120 participants in the intervention group. Eighty-five attended the nurse health check and 73 also received either *mynsnapp*, Get Healthy or both. Thirty-eight took up both *mynsnapp* and Get Healthy coaching. Of the 62 who adopted *mynsnapp*, 60 participants set goals on 132 occasions to increase vegetables, 131 to increase fruits, 97 less takeaway, 117 smaller portions, 73 less soft drink, 129 to increase physical activity time. Of the 49 who adopted Get Healthy telephone coaching, 31 set weight-related goals.

Change between baseline and 12 months

Primary outcomes

For health literacy, at 6 months, there was a greater increase in the intervention group for the HLQ domain 8 ability to find good health information score (DiD 0.22; 95% CI 0.01 to 0.44; table 3). This difference was not sustained at 12 months. There was no difference in the HLQ domain 9 understanding health information or for eHealth literacy both at 6 and 12 months. For the group that was below the median at baseline, there was also an increase in the intervention group for the HLQ domain 8 and eHealth literacy score at 6 months, and in HLQ domain 9 score at both 6 and 12 months.

There was no statistically significant effect of the intervention on BMI or BP at 12 months (table 4). The intervention group's mean BMI decreased but mean waist circumference at 12 months increased (DiD 7.08, 95% CI 2.26 to 11.90).

Secondary outcomes

There was a greater increase in diet score in the intervention group at 6 months (DiD 0.98; 95% CI 0.50 to 1.47) due to an increase in fruit intake (DiD 0.50; 95% CI 0.20 to 0.80); however, this was not sustained at 12 months. There was no statistically significant effect of the intervention on physical activity score at 6 months (table 5).

HDL fell in both groups by 7% (control) and 8% (intervention). However, total cholesterol, LDL and TGs all fell in the intervention group (table 6). There were no statistically significant effects of the intervention on lipids (total cholesterol, LDL, HDL or TG) or quality of life (EQ-5D-5L) at 12 months. Quality of life did not change in control or the intervention group (table 6).

At 6 months, the control group self-reported a decrease in the frequency of receiving advice on physical activity, whereas the level stayed the same in intervention group (DiD 16.3%, 95% CI 1.4% to 31.1%). Similarly, the frequency of weight loss counselling or referral for physical activity fell in the control group but both increased in the intervention group (weight loss counselling DiD 27.8%, 95% CI 8.8% to 46.8%; physical activity referral DiD 13.3%, 95% CI 2.32% to 24.2%). There were no statistically significant differences between the groups in frequency of receiving information on healthy eating or being referred for healthy eating or weight loss (table 7).




Figure 1 CONSORT flow diagram. CONSORT, Consolidated Standards of Reporting Trials; EOI, Expression of Interest; GPs, general practitioners; ITT, intention-to-treat; NSW, New South Wales; PN, practice nurse; Pt, Participant; SA, South Australia.

Economic analysis

The intervention costs included fixed (development of the *mynapp* app and the online training modules) and variable (practice facilitation visits, PN health check payments and telephone coaching sessions) costs. Across the 120 patients in the intervention group, the per patient fixed and variable costs were \$787 and \$558, respectively, generating a total intervention cost per patient of \$1345.

The baseline characteristics and outcome measurements of participants in the cohort providing consent to access their cost data (n=65; 33 in the intervention group and 32 in the control group) and full cohort (n=215) were similar (see online supplemental table 1). Two participants were excluded, one due to having only 6 months of cost data available after the enrolment date, and one due to extremely high pharmaceutical costs in

**Table 2** Baseline characteristics and outcomes by intervention and control

Variables	Responses	Control	Intervention	ICC
n		95	120	
Age, mean (SD)		56.2 (9.6)	58.9 (8.8)	
Gender, n (%)	Female	32 (33.7)	60 (50.0)	
	Male	63 (66.3)	60 (50.0)	
Place of birth, n (%)	Australia	59 (62.1)	66 (55.0)	
	Overseas	36 (37.9)	54 (45.0)	
Place of birth, n (%)	Australia	59 (62.8)	66 (55.0)	
	Europe	16 (17.0)	15 (12.5)	
	Asia	11 (11.7)	13 (10.8)	
	Other	7 (7.4)	25 (20.8)	
Year of arrival in Australia	Before 2000	24 (68.6)†	40 (81.6)	
	On or after 2000*	11 (31.4)	9 (18.4)	
Primary language at home, n (%)	English	88 (92.6)	96 (80.0)	
	Other	7 (7.4)	24 (20.0)	
Hospital admissions in past 12 months, n (%)	Yes	21 (22.1)	27 (22.5)	
	No	74 (77.9)	93 (77.5)	
State, n (%)	NSW	35 (36.8)	99 (82.5)	
	SA	60 (63.2)	21 (17.5)	
Qualification, n (%)	School only	38 (40.0)	47 (39.2)	
	Professional or technical	30 (31.6)	40 (33.3)	
	University degree	18 (18.9)	26 (21.7)	
	Other	9 (9.5)	7 (5.8)	
Current working status, n (%)	Working	56 (58.9)	71 (59.7)	
	Retired	20 (21.1)	28 (23.5)	
	Other	19 (20.0)	20 (16.8)	
HLQ8 (ability to find good health information)	Mean (SD)	4.3 (0.5)	4.0 (0.8)	0.0262
	Median (IQR)	4.0 (4.0–4.8)	4.0 (4.0–4.6)	
HLQ9 (understanding health information well enough to know what to do)	Mean (SD)	4.3 (0.5)	4.1 (0.7)	0.0230
	Median (IQR)	4.0 (4.0–4.8)	4.0 (4.0–4.6)	
eHealth literacy	Mean (SD)	29.2 (6.3)	27.4 (7.3)	0.0026
	Median (IQR)	32.0 (26.0–32.0)	29.0 (23.5–32.0)	
Diet	Mean (SD)	3.1 (1.6)	3.2 (1.6)	–0.0288
	Median (IQR)	3.0 (2.0–4.0)	3.0 (2.0–4.0)	
Physical activity	Mean (SD)	2.9 (2.3)	2.7 (2.5)	0.0176
	Median (IQR)	2.0 (1.0–4.0)	2.0 (1.0–4.0)	
Body mass index (BMI)	Mean (SD)	34.9 (6.9)	34.7 (5.3)	0.0122
	Median (IQR)	33.0 (30.3–36.3)	33.3 (30.5–37.2)	
Waist	Mean (SD)	112.9 (15.2)	109.4 (13.6)	0.0263
	Median (IQR)	110.0 (104.0–121.0)	108.5 (99.0–115.5)	
Systolic blood pressure	Mean (SD)	130.7 (14.1)	130.6 (14.6)	–0.0214
	Median (IQR)	132.0 (121.0–140.0)	131.0 (120.0–139.0)	
Diastolic blood pressure	Mean (SD)	81.3 (9.1)	79.2 (11.9)	0.0098
	Median (IQR)	81.0 (75.5–87.5)	80.0 (70.0–86.0)	

Missing values: health literacy domain 8 (n=4); health literacy domain 9 (n=3); eHealth (n=3); diet (n=1); BMI (n=1); waist circumference (n=78); systolic blood pressure (n=1); diastolic blood pressure (n=1).

*There were 17.1% (n=6) and 2.0% (n=1) of people who recently (on or after 2009) moved to Australia in control and intervention groups, respectively.

†Denominator for these percentages is the number of people who were born outside Australia (n=84); there were three missing values for those who were born outside Australia (n=87).

HLQ8, Health Literacy Questionnaire domain 8; HLQ9, Health Literacy Questionnaire domain 9; ICC, intracluster correlation coefficient; NSW, New South Wales; SA, South Australia.

Table 3 Effect of intervention on health literacy score at 6 and 12 months of follow up—intention-to-treat analysis

Outcome	Time point	Control		Intervention		Effect size*	Crude DiD (95% CI)†	Adjusted DiD (95% CI)†
		n	Mean (SD)	n	Mean (SD)			
HLQ8 (ability to find good health information)	Baseline	94	4.3 (0.5)	117	4.0 (0.8)		Ref	Ref
	6-month follow-up	79	4.3 (0.6)	68	4.2 (0.7)	0.31	0.22 (0.00 to 0.44)	0.22 (0.01 to 0.44)
	12-month follow-up	72	4.4 (0.5)	54	4.3 (0.6)	0.36	0.16 (−0.08 to 0.39)	0.15 (−0.08 to 0.39)
HLQ9 (understanding health information well enough to know what to do)	Baseline	95	4.3 (0.5)	117	4.1 (0.7)		Ref	Ref
	6-month follow-up	79	4.4 (0.5)	68	4.3 (0.7)	0.16	0.11 (−0.09 to 0.32)	0.13 (−0.07 to 0.33)
	12-month follow-up	72	4.4 (0.5)	54	4.4 (0.5)	0.40	0.20 (−0.03 to 0.43)	0.20 (−0.03 to 0.44)
eHealth literacy	Baseline	93	29.2 (6.3)	119	27.4 (7.3)			
	6-month follow-up	78	28.3 (6.3)	68	28.0 (5.8)	0.25	1.60 (−0.40 to 3.59)	1.60 (−0.39 to 3.58)
	12-month follow-up	70	29.4 (5.9)	52	29.5 (6.1)	0.32	1.94 (−0.48 to 4.36)	1.82 (−0.65 to 4.29)
Below median value (baseline)								
HLQ8 (ability to find good health information)	Baseline	53	3.9 (0.2)	73	3.6 (0.7)		Ref	Ref
	6-month follow-up	43	4.1 (0.5)	38	4.2 (0.6)	0.72	0.34 (0.08 to 0.60)	0.34 (0.09 to 0.59)
	12-month follow-up	43	4.3 (0.5)	32	4.2 (0.7)	0.33	0.19 (−0.06 to 0.44)	0.19 (−0.06 to 0.43)
HLQ9 (understanding health information well enough to know what to do)	Baseline	49	3.9 (0.3)	71	3.7 (0.6)		Ref	Ref
	6-month follow-up	40	4.2 (0.5)	35	4.3 (0.7)	0.49	0.27 (0.06 to 0.48)	0.28 (0.08 to 0.48)
	12-month follow-up	40	4.3 (0.5)	29	4.5 (0.5)	0.8	0.32 (0.12 to 0.53)	0.33 (0.12 to 0.54)
eHealth literacy score	Baseline	41	23.8 (5.2)	69	22.5 (5.3)		Ref	Ref
	6-month follow-up	34	25.6 (7.1)	34	26.7 (4.8)	0.40	2.40 (−0.21 to 5.02)	2.34 (−0.39 to 5.06)
	12-month follow-up	27	26.5 (6.2)	25	29.5 (4.7)	0.42	4.12 (1.48 to 6.75)	3.77 (0.96 to 6.59)
Above median value (baseline)								
HLQ8 (ability to find good health information)	Baseline	41	4.8 (0.3)	44	4.7 (0.3)		Ref	Ref
	6-month follow-up	35	4.4 (0.6)	28	4.2 (0.7)	0.15	−0.09 (−0.45 to 0.27)	−0.44 (−2.27 to 1.39)
	12-month follow-up	28	4.5 (0.5)	20	4.4 (0.6)	0	−0.04 (−0.41 to 0.33)	−0.18 (−2.04 to 1.67)
HLQ9 (understanding health information well enough to know what to do)	Baseline	46	4.7 (0.3)	46	4.7 (0.3)		Ref	Ref
	6-month follow-up	39	4.6 (0.4)	31	4.3 (0.7)	0.53	−0.27 (−0.55 to 0.01)	−0.25 (−0.54 to 0.03)
	12-month follow-up	32	4.5 (0.4)	23	4.4 (0.6)	0.39	−0.17 (−0.41 to 0.07)	0.17 (−0.41 to 0.08)
eHealth literacy score	Baseline	52	33.5 (3.0)	50	34.1 (3.1)		Ref	Ref
	6-month follow-up	42	30.8 (4.3)	33	29.5 (6.5)	0.35	−1.90 (−4.50 to 0.70)	−1.77 (−4.36 to 0.82)
	12-month follow-up	42	31.1 (4.9)	26	30.0 (7.0)	0.28	−1.70 (−5.25 to 1.85)	−1.68 (−5.18 to 1.81)

Bold values signifies $p < 0.05$.

*Cohen's d.

†Adjusted for age, gender and state.

DiD, difference-in-differences; HLQ8, Health Literacy Questionnaire domain 8; HLQ9, Health Literacy Questionnaire domain 9.

the 12 months prior to enrolment for the treatment of age-related macular degeneration, a condition unrelated to the focus of the intervention.

Online supplemental table 1C presents the mean crude cost DiD between the 12 months prior and after recruitment to the trial. Excluding the outlier participant with high pharmaceutical costs, mean costs were higher in the intervention group in all cost categories, but there were no statistically significant differences between the intervention and control groups for the alternative cost categories (GP costs, GP and specialist costs and Pharmaceutical Benefit Scheme (PBS) costs) nor for the aggregated cost. Including the participant with outlier PBS

costs, mean costs are lower in the intervention group for comparisons including PBS cost data, but the CIs remain very wide (online supplemental table 1D).

There were no adverse events or harms reported during the trial.

DISCUSSION

In this trial of an intervention involving a PN health check, a mobile app and phone coaching in primary healthcare, we found positive effects on some outcomes (health literacy and diet at 6 months) but not on physical activity, weight or other outcomes. The primary

Table 4 Effect of intervention on anthropometry and blood pressure at 12 months of follow up—intention-to-treat analysis

Outcome	Time point	Control		Intervention		Effect size	Crude DiD (95% CI)	Adjusted DiD (95% CI)*
		n	Mean (SD)	n	Mean (SD)			
BMI, kg/m ²	Baseline	94	34.9 (6.9)	120	34.7 (5.3)		Ref	Ref
	12-month follow-up	49	32.9 (5.7)	52	34.3 (6.0)	0.27	1.45 (−0.16 to 3.06)	1.22 (−0.46 to 2.90)
Waist circumference, cm	Baseline	49	112.9 (15.2)	88	109.4 (13.6)		Ref	Ref
	12-month follow-up	20	107.0 (9.6)	49	112.4 (15.6)	0.62	8.24 (2.73 to 13.74)	7.08 (2.26 to 11.90)
Systolic blood pressure, mm Hg	Baseline	95	130.7 (14.1)	119	130.6 (14.6)		Ref	Ref
	12-month follow-up	64	133.0 (15.3)	50	130.8 (14.6)	0.17	−2.13 (−8.18 to 3.92)	−1.48 (−7.34 to 4.38)
Diastolic blood pressure, mm Hg	Baseline	95	81.3 (9.1)	119	79.2 (11.9)		Ref	Ref
	12-month follow-up	64	82.7 (8.6)	50	77.6 (9.1)	0.12	−2.84 (−5.94 to 0.25)	−3.18 (−6.50 to 0.14)

*Adjusted for age, gender and state.
BMI, body mass index; DiD, difference-in-differences.

hypothesis was that the intervention would lead to improved health literacy, health behaviours and positive changes in weight and other physiological measures. There were some differences between intervention and control groups at baseline but minimal differences in the outcomes and these were unlikely to have had a major influence on the findings. Health literacy improved in the intervention group at 6 months, although there was no further change by 12 months. Additionally, eHealth literacy improved only among those whose baseline health literacy was below the median. Although similar proportions of participants in both groups set goals for

diet and physical activity, patients in the intervention group were more likely to report an improved diet score (due to a greater increase in fruit intake) compared with the control group. There was no difference in the physical activity score between the intervention and control groups. A lack of change in physical activity outcomes may reflect a need for group rather than individual approaches to physical activity promotion for people from migrant or low socioeconomic backgrounds.⁵⁵ The intervention was tailored to patients' needs and motivation but was not codesigned or specifically tailored to differences in individual cultural and religious beliefs

Table 5 Effect of intervention on physical activity and diet score at 6 and 12 months of follow up—intention-to-treat analysis

Outcome	Time point	Control		Intervention		Effect size*	Crude DiD (95% CI)	Adjusted DiD (95% CI)
		n	Mean (SD)	n	Mean (SD)			
Total physical activity score	Baseline	95	2.9 (2.3)	120	2.7 (2.5)		Ref	Ref
	6-month follow-up	79	3.6 (2.6)	68	3.0 (2.3)	0.16	−0.45 (−1.06 to 0.15)	−0.56 (−1.19 to 0.06)
	12-month follow-up	72	3.6 (2.5)	54	3.9 (2.2)	0.21	0.47 (−0.47 to 1.42)	0.38 (−0.59 to 1.35)
Diet score	Baseline	95	3.1 (1.6)	119	3.2 (1.6)		Ref	Ref
	6-month follow-up	79	3.1 (1.7)	68	4.1 (1.5)	0.56	0.98 (0.48 to 1.48)	0.98 (0.50 to 1.47)
	12-month follow-up	72	3.8 (1.5)	54	3.9 (1.9)	0	−0.04 (−0.51 to 0.44)	0.05 (−0.41 to 0.50)
Vegetable intake	Baseline	95	1.8 (1.2)	120	1.8 (1.2)		Ref	Ref
	6-month follow-up	79	1.9 (1.3)	68	2.3 (1.3)	0.31	0.46 (0.02 to 0.90)	0.46 (0.03 to 0.89)
	12-month follow-up	72	2.4 (1.2)	54	2.3 (1.4)	0.46	−0.14 (−0.53 to 0.26)	−0.07 (−0.44 to 0.31)
Fruit intake	Baseline	95	1.3 (0.9)	119	1.4 (1.0)		Ref	Ref
	6-month follow-up	79	1.2 (0.9)	68	1.8 (0.8)	0.59	0.49 (0.20 to 0.79)	0.50 (0.20 to 0.80)
	12-month follow-up	72	1.4 (0.9)	54	1.6 (0.9)	0.11	0.03 (−0.23 to 0.30)	0.05 (−0.22 to 0.32)

Bold values signifies p<0.05.
*Cohen's d.
DiD, difference-in-differences.

Table 6 Effect of intervention on the secondary outcomes intention-to-treat analysis (who had two different measurements at baseline and 12 months)

Outcome	Time point	Control		Intervention		Crude DiD (95% CI)	Adjusted DiD (95% CI)*
		n	Mean (SD)	n	Mean (SD)		
HDL cholesterol	Baseline	90	1.4 (0.4)	109	1.3 (0.4)	Ref	Ref
	12-month follow-up	43	1.3 (0.3)	31	1.2 (0.4)	0.02 (−0.09 to 0.14)	0.04 (−0.08 to 0.16)
LDL cholesterol	Baseline	77	2.8 (0.9)	108	2.9 (0.8)	Ref	Ref
	12-month follow-up	25	2.9 (1.2)	28	2.7 (0.7)	−0.28 (−0.71 to 0.15)	−0.26 (−0.67 to 0.15)
Triglyceride	Baseline	92	1.7 (0.8)	114	1.7 (0.8)	Ref	Ref
	12-month follow-up	46	1.7 (0.8)	32	1.5 (0.8)	−0.20 (−0.50 to 0.09)	−0.22 (−0.52 to 0.09)
Total cholesterol	Baseline	93	4.9 (0.9)	115	4.9 (1.0)	Ref	Ref
	12-month follow-up	51	4.9 (1.2)	33	4.6 (0.8)	−0.32 (−0.65 to 0.01)	−0.31 (−0.64 to 0.01)
Quality of life change (mean (SD))	Baseline	95	0.88 (0.12)	120	0.87 (0.12)	Ref	Ref
	12-month follow-up	72	0.87 (0.16)	54	0.90 (0.11)	0.04 (0.00 to 0.08)	0.04 (0.00 to 0.08)

*Adjusted for age, gender and state.
DiD, difference-in-differences; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

and practices, which may mediate changes in physical activity.⁵⁶

Although there were small changes in health literacy and diet, the intervention was not associated with differences in clinical endpoints such as BMI, BP, lipids or in quality of life after adjustment for age, gender and state. This may be because we did not recruit our required sample size or because the intervention lacked sufficient intensity and duration, as has been observed in other studies.¹⁰ The lack of change in physical activity, especially at 12 months, may also have contributed, and changes in BP and lipids may have been confounded by treatment with medications since most patients' BP and lipids were

within recommended guideline levels at baseline. Further research is thus required to evaluate digital interventions which allow tailoring to patients' differing health literacy and culture and actively supported in their use over a longer period.

Only two-thirds of the patients in the intervention group received the full intervention (ie, received the health check with *mynapp* and/or Get Healthy coaching components). This was influenced by patient choice through discussion with their clinicians reflecting the real-world setting of Australian general practice. This variable engagement with the different components of the intervention may have reduced its overall effectiveness.

Table 7 Effect of intervention on the secondary outcomes (from survey data)—intention-to-treat analysis

Outcome	Time point	Control		Intervention		Crude DiD (95% CI)	Adjusted DiD (95% CI)*
		n	% (n)	n	% (n)		
Info or advice healthy eating	Baseline	95	27.4 (26)	120	44.2 (53)	Ref	Ref
	6-month follow-up	79	17.7 (14)	68	39.7 (27)	5.01 (−18.73 to 28.76)	3.30 (−21.10 to 27.69)
Info or advice physical activity	Baseline	95	30.5 (29)	120	40.8 (49)	Ref	Ref
	6-month follow-up	79	11.4 (9)	68	39.7 (27)	18.03 (3.19 to 32.86)	16.27 (1.40 to 31.14)
Info or advice weight loss	Baseline	95	34.7 (33)	120	43.3 (52)	Ref	Ref
	6-month follow-up	79	13.9 (11)	68	51.5 (35)	29.07 (10.41 to 47.74)	27.83 (8.83 to 46.84)
Referral to healthy eating	Baseline	95	11.6 (11)	120	10.0 (12)	Ref	Ref
	6-month follow-up	79	10.1 (8)	68	22.1 (15)	13.46 (−3.25 to 30.16)	14.46 (−2.35 to 31.27)
Referral to physical activity	Baseline	95	8.4 (8)	120	3.3 (4)	Ref	Ref
	6-month follow-up	79	5.1 (4)	68	13.2 (9)	13.24 (2.45 to 24.04)	13.28 (2.32 to 24.24)
Referral to weight loss	Baseline	95	7.4 (7)	120	7.5 (9)	Ref	Ref
	6-month follow-up	79	7.6 (6)	68	10.3 (7)	2.49 (−7.68 to 12.66)	2.50 (−7.75 to 12.74)

Bold values signifies p<0.05.
*Adjusted for age, gender and state.
DiD, difference-in-differences.



However, patients in the intervention group were more likely to recall being offered information or referral for physical activity or weight loss counselling than their counterparts in the control group.

In the cost analyses, low recruitment made the study insufficiently powered to draw meaningful conclusions. There was no evidence of difference in numbers of GP visits, Medicare Benefits Schedule or PBS costs between the groups over the period of the study. Despite some positive changes in some behavioural endpoints (health literacy and diet), there were no changes in clinical endpoints such as weight or other physiological measures, or in quality of life at 12 months. Trials of weight loss in primary care often show little or no change.⁵⁷ However, previous studies involving the use of apps and behavioural counselling by healthcare providers have proven successful even in low socioeconomic groups where goals were individually tailored to the patient's level of health literacy and the interventions were of moderate to high intensity.¹¹ This suggests that the intervention in the current study may have been more effective if it was more tailored to the patient's individual health literacy needs.

There were several limitations to our study. Like other studies, this study failed to achieve its planned sample size due to major challenges recruiting practices and patients despite considerable effort and an extension to the time frame of the study.⁵⁸ Post-hoc power calculations, based on our results, showed that with a sample of 100 in each arm, we would be able to detect a mean difference in diet score of 0.6–0.7 (servings per day) and a mean difference in the Health Literacy Scale scores of 0.2–0.3. Both these differences are less than in previous studies and may not be clinically meaningful.^{32 59} For all the other measures, the differences that were able to be detected were larger than expected from moderate-intensity interventions (mean physical activity score difference of 1.5, mean BMI difference of 5.5 kg/m², mean BP change of 15 mm Hg, mean cholesterol difference of 0.8).¹⁰ Our recruitment challenges suggest the need for greater efforts to increase the perceived benefits (such as improved access to quality care) and decrease barriers (especially time) associated with participation in studies such as this in the future.

There were five primary outcomes (including two HLQ domains, eHEALS, weight and BP). Furthermore, the health literacy measures were assessed at both 6 and 12 months increasing the likelihood of a type 1 error (ie, finding a significant difference). The study was conducted in only two urban areas of Australia and the findings may not therefore be generalised to other communities such as rural areas. Lastly, the measures of health literacy, diet and physical activity had some limitations, and may have not been sensitive enough to capture all changes due to the intervention.

Assessments of patient socioeconomic variables and health literacy indicate that the study fell short in recruiting its target population of people with low socioeconomic status and low health literacy. At baseline, levels of health literacy were higher than anticipated and were

in fact comparable with overweight or obese patients in the general population who were part of the national health literacy survey.⁶⁰ Our figures for 'born overseas' are higher than the Australian average but 'language spoken at home' and 'employment status' are similar to the Australian average.⁶¹ It is therefore possible that the requirements for written consent and engagement with the research study may have tended to discourage those with lower English language literacy, as has been found in some research.⁶² Furthermore, uptake by the participants in our study in the various components of the intervention varied. Previous research has identified that socioeconomic factors have impacts on intervention/trial uptake, intervention adherence and trial attrition.⁶³ Future research could consider using codesign principles to help better engage specific population groups, as well as GPs and PNs working with these groups, in the research design and development of the intervention.⁶⁴

CONCLUSION

This trial of a multifaceted intervention designed to support better preventive care for overweight and obese patients from low socioeconomic areas in the real-world environment of Australian general practice showed some short-term improvement in health literacy and diet but did not show any change in weight or other physiological variables. It was insufficiently powered for cost analysis. While there was evidence that the intervention was implemented as planned, there was variable uptake of its components, and it may therefore have been of insufficient intensity to achieve sustained change in weight and other primary outcomes. However, any preventive intervention in primary care needs to be sustainable and tailored to its capacity.

Author affiliations

¹Centre for Primary Health Care and Equity, University of New South Wales, Sydney, New South Wales, Australia

²Discipline of General Practice, Adelaide Medical School, The University of Adelaide, Adelaide, South Australia, Australia

³Sydney Nursing School, University of Sydney, Sydney, New South Wales, Australia

⁴Health Sciences & Medicine, Bond University, Gold Coast, Queensland, Australia

⁵Flinders Medical Research Institute, Flinders University, Adelaide, South Australia, Australia

⁶Public Health, The University of Sydney, Sydney, New South Wales, Australia

⁷School of Population Health, University of New South Wales, Sydney, New South Wales, Australia

⁸College of Nursing and Health Sciences, Flinders University, Adelaide, New South Wales, Australia

⁹Centre for Global Health and Equity, Swinburne University of Technology, Burwood, Victoria, Australia

¹⁰Australian Institute of Health Innovation, Faculty of Medicine and Health Sciences, Macquarie University, Sydney, New South Wales, Australia

Correction notice The article has been corrected since it was published online. There is a minor error in the DiD value in the results section of the abstract. The text has been corrected from "There was a small improvement in diet scores at 6 months (DiD 0.78 (0.10 to 1.47); $p=0.026$)" to "There was a small improvement in diet scores at 6 months (DiD 0.98 (0.50 to 1.47); $p=0.026$)."

Twitter Elizabeth Denney-Wilson @denneywilson, Siaw-Teng Liaw @EIAstee and Richard Osborne @richardosborne4

Contributors MH was the chief investigator, is guarantor, led the development of and implementation of the study, interpreted the data and drafted the manuscript. SMP developed the trial processes, coordinated the trial across sites, contributed to the development of the data collection tools, collected and analysed data and drafted the manuscript. MB cleaned the data, designed the analysis plan, conducted the analysis, interpreted the data and contributed to the draft of the manuscript. NS led the development and implementation of the study in SA, was instrumental in designing the DCP module for the trial, interpreted the data and contributed to the manuscript. ED-W contributed to the design of the study, developed the training modules, and contributed to the interpretation of the data and the manuscript. NZ contributed to the trial design, trial implementation and interpretation of the data for the manuscript. JK contributed to the trial design, designed the economic analysis and interpreted these data for the manuscript. AK cleaned the DCP data, designed the analysis plan, conducted the analysis, developed the data tables and contributed to the draft of the manuscript. DN contributed to the trial design, trial implementation and interpretation of the data for the manuscript, particularly the health literacy content. JR conducted the economic analysis and interpreted the data for inclusion in the manuscript. S-TL contributed to the trial design, trial implementation and interpretation of the data for the manuscript. CM liaised with SA practices to collect patient and practice data, collated the data and contributed to the interpretation of the data for the manuscript. OF was instrumental in designing the DCP module for the trial and troubleshooting data collection using DCP; interpreted the outcome data and contributed to the manuscript. AT liaised with NSW practices to collect patient and practice data, collated and cleaned the data, and contributed to the management and interpretation of the data. RO contributed to the trial design, particularly the tools to collect patient data and interpretation of data particularly the health literacy outcome data. He contributed to the draft manuscript. AYSL designed and developed the *mynsnapp* app and the data collected via the application and interpretation and plan for analysing these data. All authors approved the final version for publication and agree to be accountable for the integrity of the content, and responsible for any issues that arise from publication of the trial data.

Funding This work was supported by the National Health and Medical Research Council (NHMRC) of Australia (project grant APP1125681, 2017).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval This trial was approved by the University of New South Wales Human Research Ethics Committee (HC17474). The University of Adelaide Human Research Ethics Committee ratified this approval. Written consent was obtained from all participating practices to conduct the trial in the practice and access practice data; individual consent was obtained from all participating GPs and PNs. Patients provided written consent to participate in the trial and additional written consent was obtained for the researchers to access individual health service usage data (Medicare Benefits Schedule (MBS)) and pharmaceutical use (Pharmaceutical Benefit Scheme (PBS)) according to protocols governing access to these data through Services Australia.⁴² All practices received \$A1000 payment to cover the administrative costs of participation. To compensate them for their time, patients from both groups who completed the baseline and 6-month follow-up received \$A30 shopping voucher and then an additional \$A30 voucher if they completed the 12-month follow-up.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Individual participant data that underlie the results reported in this article will be available within 12 months of publication after deidentification (text, tables, figures and appendices) to investigators whose proposed use of the data has been approved by our Ethics Committee.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which

permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Sharon M Parker <http://orcid.org/0000-0001-7904-6420>
 Margo Barr <http://orcid.org/0000-0002-3007-0216>
 Nigel Stocks <http://orcid.org/0000-0002-9018-0361>
 Alamgir Kabir <http://orcid.org/0000-0002-3762-8307>
 Jackie Roseleur <http://orcid.org/0000-0003-4990-1312>
 Siaw-Teng Liaw <http://orcid.org/0000-0001-5989-3614>
 Mark Harris <http://orcid.org/0000-0002-0705-8913>

REFERENCES

- 1 Kaiser KA, Carson TL, Dhurandhar EJ, *et al*. Biobehavioural approaches to prevention and treatment: a call for implementation science in obesity research. *Obes Sci Pract* 2020;6:3–9.
- 2 Australian Institute of Health and Welfare. *Australian burden of disease study: impact and causes of illness and death in Australia 2015*. Canberra: AIHW, 2019.
- 3 Australian Institute of Health and Welfare. *Overweight and obesity: an interactive insight*. Canberra: Australian Department of Health, 2020. <https://www.aihw.gov.au/reports-data/behaviours-risk-factors/overweight-obesity/overview/2021>
- 4 Australian Bureau of Statistics. *National health survey: first results 2017–18: Australian Bureau of statistics*, 2018.
- 5 National Health and Medical Research Council (NHMRC). *Clinical practice guidelines for the management of overweight and obesity in adults, adolescents and children in Australia*. Melbourne: NHMRC, 2013.
- 6 Taggart J, Williams A, Dennis S, *et al*. A systematic review of interventions in primary care to improve health literacy for chronic disease behavioral risk factors. *BMC Fam Pract* 2012;13:49.
- 7 Madigan CD, Graham HE, Sturgiss E, *et al*. Effectiveness of weight management interventions for adults delivered in primary care: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2022;377:e069719.
- 8 , Curry SJ, Krist AH, *et al*, US Preventive Services Task Force. Behavioral weight loss interventions to prevent obesity-related morbidity and mortality in adults: US preventive services Task force recommendation statement. *JAMA* 2018;320:1163–71.
- 9 Mastellos N, Gunn LH, Felix LM, *et al*. Transtheoretical model stages of change for dietary and physical exercise modification in weight loss management for overweight and obese adults. *Cochrane Database Syst Rev* 2014;2:Cd008066.
- 10 Tronieri JS, Wadden TA, Chao AM, *et al*. Primary care interventions for obesity: review of the evidence. *Curr Obes Rep* 2019;8:128–36.
- 11 Bennett GG, Steinberg D, Askew S, *et al*. Effectiveness of an APP and provider counseling for obesity treatment in primary care. *Am J Prev Med* 2018;55:777–86.
- 12 Schirmann F, Kanehl P, Jones L. What intervention elements drive weight loss in Blended-Care behavior change interventions? A real-world data analysis with 25,706 patients. *Nutrients* 2022;14. doi:10.3390/nu14142999. [Epub ahead of print: 21 Jul 2022].
- 13 Tan D, Zwar NA, Dennis SM, *et al*. Weight management in general practice: what do patients want? *Med J Aust* 2006;185:73–5.
- 14 Graham J, Tudor K, Jebb SA, *et al*. The equity impact of brief opportunistic interventions to promote weight loss in primary care: secondary analysis of the BWEL randomised trial. *BMC Med* 2019;17:1–9.
- 15 Ahern AL, Aveyard P, Boyland EJ, *et al*. Inequalities in the uptake of weight management interventions in a pragmatic trial: an observational study in primary care. *British Journal of General Practice* 2016;66:e258–63.
- 16 Michou M, Panagiotakos DB, Costarelli V, VJCEjoph C. Low health literacy and excess body weight: a systematic review. *Cent Eur J Public Health* 2018;26:234–41.
- 17 Australian Bureau of Statistics. *Health literacy, Australia: ABS*, 2006.
- 18 Joshi C, Jayasinghe UW, Parker S, *et al*. Does health literacy affect patients' receipt of preventative primary care? A multilevel analysis. *BMC Fam Pract* 2014;15:171.
- 19 Osborn C, Paasche-Orlow MK, Bailey SC. The mechanisms linking health literacy to behavior and health status. *Am J Health Behav* 2011;35:118–28.
- 20 von Wagner C, Knight K, Steptoe A, *et al*. Functional health literacy and health-promoting behaviour in a national sample of British adults. *J Epidemiol Community Health* 2007;61:1086–90.

- 21 Wolf MS, Gazmararian JA, Baker DW. Health literacy and health risk behaviors among older adults. *Am J Prev Med* 2007;32:19–24.
- 22 Lim S, Beauchamp A, Dodson S, et al. Health literacy and fruit and vegetable intake in rural Australia. *Public Health Nutr* 2017;20:2680–4.
- 23 von Wagner C, Steptoe A, Wolf MS, et al. Health literacy and health actions: a review and a framework from health psychology. *Health Educ Behav* 2009;36:860–77.
- 24 Schillinger D, Bindman A, Wang F, et al. Functional health literacy and the quality of physician-patient communication among diabetes patients. *Patient Educ Couns* 2004;52:315–23.
- 25 Baillargeon J-P, St-Cyr-Tribble D, Xhignesse M, et al. Impact of an educational intervention combining clinical obesity preceptorship with electronic networking tools on primary care professionals: a prospective study. *BMC Med Educ* 2020;20:361.
- 26 Arora M, Barquera S, Farpour Lambert NJ, et al. Stigma and obesity: the crux of the matter. *Lancet Public Health* 2019;4:e549–50.
- 27 Caterson ID, Alfadda AA, Auerbach P, et al. Gaps to bridge: misalignment between perception, reality and actions in obesity. *Diabetes Obes Metab* 2019;21:1914–24.
- 28 Australian Bureau of Statistics. *Socioeconomic indexes for areas (SEIFA) index of relative socio-economic disadvantage (IRSd)*. Canberra (Aust: ABS, 2016.
- 29 Doctors Control Panel (DCP). Adelaide SA2021. Available: <https://www.doctorscontrolpanel.com.au/> [Accessed 23 Nov 2021].
- 30 SAS Institute Inc. Statistical analysis software Cary, NC, USA2021. Available: https://www.sas.com/en_au/software/stat.html
- 31 Faruqi N, Lloyd J, Ahmad R, et al. Feasibility of an intervention to enhance preventive care for people with low health literacy in primary health care. *Aust J Prim Health* 2015;21:321–6.
- 32 Parker SM, Stocks N, Nutbeam D, et al. Preventing chronic disease in patients with low health literacy using eHealth and teamwork in primary healthcare: protocol for a cluster randomised controlled trial. *BMJ Open* 2018;8:e023239.
- 33 National Health and Medical Research Council (NHMRC). *Clinical practice guidelines for the management of overweight and obesity in adults, adolescents and children in Australia - Systematic Review*. Melbourne: NHMRC, 2013.
- 34 Tobacco Use and Dependence Guideline Panel. *Treating tobacco use and dependence*. Rockville, MD: US Department of Health and Human Services, 2008.
- 35 Jay M, Gillespie C, Schlair S, et al. Physicians' use of the 5As in counseling obese patients: is the quality of counseling associated with patients' motivation and intention to lose weight? *BMC Health Serv Res* 2010;10:159.
- 36 Lau AYS, Sintchenko V, Crimmins J, et al. Impact of a web-based personally controlled health management system on influenza vaccination and health services utilization rates: a randomized controlled trial. *J Am Med Inform Assoc* 2012;19:719–27.
- 37 Webb TL, Joseph J, Yardley L, et al. Using the Internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *J Med Internet Res* 2010;12:e4.
- 38 DiFilippo KN, Huang W-H, Andrade JE, et al. The use of mobile apps to improve nutrition outcomes: a systematic literature review. *J Telemed Telecare* 2015;21:243–53.
- 39 Payne HE, Lister C, West JH, et al. Behavioral functionality of mobile apps in health interventions: a systematic review of the literature. *JMIR Mhealth Uhealth* 2015;3:e20.
- 40 Zhao J, Freeman B, Li M. Can mobile phone Apps influence people's health behavior change? an evidence review. *J Med Internet Res* 2016;18:e287.
- 41 Ha Dinh TT, Bonner A, Clark R, et al. The effectiveness of the teach-back method on adherence and self-management in health education for people with chronic disease: a systematic review. *JBIM Database System Rev Implement Rep* 2016;14:210–47.
- 42 Osborne RH, Batterham RW, Elsworth GR, et al. The grounded psychometric development and initial validation of the health literacy questionnaire (HLQ). *BMC Public Health* 2013;13.
- 43 Simpson RM, Knowles E, O'Cathain A. Health literacy levels of British adults: a cross-sectional survey using two domains of the health literacy questionnaire (HLQ). *BMC Public Health* 2020;20:1–3.
- 44 Yadav UN, Lloyd J, Hosseinzadeh H, et al. Levels and determinants of health literacy and patient activation among multi-morbid COPD people in rural Nepal: findings from a cross-sectional study. *PLoS One* 2020;15:e0233488.
- 45 Norman CD, Skinner HA. eHEALS: the eHealth literacy scale. *J Med Internet Res* 2006;8:e27.
- 46 Smith BJet al. Promoting physical activity in general practice: a controlled trial of written advice and information materials. *Br J Sports Med* 2000;34:262–7.
- 47 Hendrie GA, Baird D, Golley RK, et al. The CSIRO healthy diet score: an online survey to estimate compliance with the Australian dietary guidelines. *Nutrients* 2017;9:47.
- 48 Herdman M, Gudex C, Lloyd A, et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res* 2011;20:1727–36.
- 49 Devlin NJ, Shah KK, Feng Y, et al. Valuing health-related quality of life: an EQ-5D-5L value set for England. *Health Econ* 2018;27:7–22.
- 50 Lechner M. The estimation of causal effects by Difference-in-Difference methods. *Foundations and Trends(R) in Econometrics* 2011;4:165–224.
- 51 Hardin JW. Generalized estimating equations (GEE). In: Everitt BS, Howell DC, eds. *Encyclopedia of statistics in behavioral science*, 2005.
- 52 Michael L. The estimation of causal effects by difference-in-difference methods. *Foundations and Trends in Econometrics* 2010;4:165–224.
- 53 R Core Team. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing, 2020.
- 54 Campbell MK, Piaggio G, Elbourne DR, et al. Consort 2010 statement: extension to cluster randomised trials. *BMJ* 2012;345:e5661.
- 55 Cleland CL, Tully MA, Kee F, et al. The effectiveness of physical activity interventions in socio-economically disadvantaged communities: a systematic review. *Prev Med* 2012;54:371–80.
- 56 Dominick GM, Dunsiger SI, Pekmezi DW, et al. Moderating effects of health literacy on change in physical activity among Latinas in a randomized trial. *J Racial Ethn Health Disparities* 2015;2:351–7.
- 57 Booth HP, Prevost TA, Wright AJ, et al. Effectiveness of behavioural weight loss interventions delivered in a primary care setting: a systematic review and meta-analysis. *Fam Pract* 2014;31:643–53.
- 58 Perkins D, Harris MF, Tan J, et al. Engaging participants in a complex intervention trial in Australian general practice. *BMC Med Res Methodol* 2008;8:55.
- 59 Muscat DM, Song W, Cvejic E, et al. The impact of the chronic disease self-management program on health literacy: a pre-post study using a multi-dimensional health literacy instrument. *Int J Environ Res Public Health* 2020;17:58.
- 60 Australian Bureau of Statistics. *National health survey: health literacy*, 2018.
- 61 Statistics ABo. Cultural diversity: Census. Canberra: Australian government, 2021. Available: [https://www.abs.gov.au/statistics/people-and-communities/cultural-diversity-census/latest-release#:~:text=Key%20statistics,-27.6%20per%20cent&text=Top%205%20languages%20used%20at,Punjabi%20\(0.9%20per%20cent\)2022](https://www.abs.gov.au/statistics/people-and-communities/cultural-diversity-census/latest-release#:~:text=Key%20statistics,-27.6%20per%20cent&text=Top%205%20languages%20used%20at,Punjabi%20(0.9%20per%20cent)2022)
- 62 Kripalani S, Heerman WJ, Patel NJ, et al. Association of health literacy and Numeracy with interest in research participation. *J Gen Intern Med* 2019;34:544–51.
- 63 Birch JM, Jones RA, Mueller J, et al. A systematic review of inequalities in the uptake of, adherence to, and effectiveness of behavioral weight management interventions in adults. *Obes Rev* 2022;23:e13438.
- 64 Halvorsrud K, Kucharska J, Adlington K, et al. Identifying evidence of effectiveness in the co-creation of research: a systematic review and meta-analysis of the International healthcare literature. *J Public Health* 2021;43:197–208.
- 65 Services Australia. Statistical information and data 2021 Australian government Canberra;. Available: <https://www.servicesaustralia.gov.au/organisations/about-us/reports-and-statistics/statistical-information-and-data>

Supplementary 1. CONSORT checklist when reporting a cluster randomised trial: HeLP GP Trial.

Section/Topic	Item No	Standard Checklist item	Extension for cluster designs	Page No *
Title and abstract				
	1a	Identification as a randomised trial in the title	Identification as a cluster randomised trial in the title	Title page
	1b	Structured summary of trial design, methods, results, and conclusions (for specific guidance see CONSORT for abstracts) ^{1,2}	See table 2	Abstract
Introduction				
Background and objectives	2a	Scientific background and explanation of rationale	Rationale for using a cluster design	Page 3
	2b	Specific objectives or hypotheses	Whether objectives pertain to the cluster level, the individual participant level or both	Page 3
Methods				
Trial design	3a	Description of trial design (such as parallel, factorial) including allocation ratio	Definition of cluster and description of how the design features apply to the clusters	Page 3
	3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons		Page 8
Participants	4a	Eligibility criteria for participants	Eligibility criteria for clusters	Page 3/4
	4b	Settings and locations where the data were collected		Page 3/4
Interventions	5	The interventions for each group with sufficient details to allow replication, including how and when they were actually administered	Whether interventions pertain to the cluster level, the individual participant level or both	Page 6/7
Outcomes	6a	Completely defined pre-specified primary and	Whether outcome measures pertain to the cluster level, the	Table 1

		secondary outcome measures, including how and when they were assessed	individual participant level or both	
	6b	Any changes to trial outcomes after the trial commenced, with reasons		NA
Sample size	7a	How sample size was determined	Method of calculation, number of clusters(s) (and whether equal or unequal cluster sizes are assumed), cluster size, a coefficient of intracluster correlation (ICC or k), and an indication of its uncertainty	Page 9
	7b	When applicable, explanation of any interim analyses and stopping guidelines		NA
Randomisation:				
Sequence generation	8a	Method used to generate the random allocation sequence		Page 5
	8b	Type of randomisation; details of any restriction (such as blocking and block size)	Details of stratification or matching if used	Page 5
Allocation concealment mechanism	9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	Specification that allocation was based on clusters rather than individuals and whether allocation concealment (if any) was at the cluster level, the individual participant level or both	Page 5
Implementation	10	Who generated the random allocation sequence, who enrolled participants, and who assigned participants to interventions	Replace by 10a, 10b and 10c	See 10a – 10c
	10a		Who generated the random allocation sequence, who enrolled clusters, and who assigned clusters to interventions	Page 5

	10b		Mechanism by which individual participants were included in clusters for the purposes of the trial (such as complete enumeration, random sampling)	Page 5
	10c		From whom consent was sought (representatives of the cluster, or individual cluster members, or both), and whether consent was sought before or after randomisation	Page 8
Blinding				
	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how		NA
	11b	If relevant, description of the similarity of interventions		NA
Statistical methods				
	12a	Statistical methods used to compare groups for primary and secondary outcomes	How clustering was taken into account	Page 13
	12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses		Page 13
Results				
Participant flow (a diagram is strongly recommended)				
	13a	For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome	For each group, the numbers of clusters that were randomly assigned, received intended treatment, and were analysed for the primary outcome	Figure 1
	13b	For each group, losses and exclusions after randomisation, together with reasons	For each group, losses and exclusions for both clusters and individual cluster members	Figure 1
Recruitment				
	14a	Dates defining the periods of recruitment and follow-up		Page 4/5

	14b	Why the trial ended or was stopped		NA
Baseline data	15	A table showing baseline demographic and clinical characteristics for each group	Baseline characteristics for the individual and cluster levels as applicable for each group	Table 2
Numbers analysed	16	For each group, number of participants (denominator) included in each analysis and whether the analysis was by original assigned groups	For each group, number of clusters included in each analysis	Page 14
Outcomes and estimation	17a	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval)	Results at the individual or cluster level as applicable and a coefficient of intracluster correlation (ICC or k) for each primary outcome	ICC included in Table 2 Effect size included in Tables 3, 4 and 5
	17b	For binary outcomes, presentation of both absolute and relative effect sizes is recommended		Absolute differences provided
Ancillary analyses	18	Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishing pre-specified from exploratory		NA
Harms	19	All important harms or unintended effects in each group (for specific guidance see CONSORT for harms ³)		NA
Discussion				
Limitations	20	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant, multiplicity of analyses		Page 23/24
Generalisability	21	Generalisability (external validity, applicability) of the trial findings	Generalisability to clusters and/or individual participants (as relevant)	Page 24
Interpretation	22	Interpretation consistent with results, balancing benefits and harms, and		Conclusions

		considering other relevant evidence	
Other information			
Registration	23	Registration number and name of trial registry	Title page
Protocol	24	Where the full trial protocol can be accessed, if available	Title page
Funding	25	Sources of funding and other support (such as supply of drugs), role of funders	Page 25

REFERENCES

- ¹ Hopewell S, Clarke M, Moher D, Wager E, Middleton P, Altman DG, et al. CONSORT for reporting randomised trials in journal and conference abstracts. *Lancet* 2008, 371:281-283
- ² Hopewell S, Clarke M, Moher D, Wager E, Middleton P, Altman DG at al (2008) CONSORT for reporting randomized controlled trials in journal and conference abstracts: explanation and elaboration. *PLoS Med* 5(1): e20
- ³ Ioannidis JP, Evans SJ, Gotzsche PC, O'Neill RT, Altman DG, Schulz K, Moher D. Better reporting of harms in randomized trials: an extension of the CONSORT statement. *Ann Intern Med* 2004; 141(10):781-788.

Supplementary Tables S1

Table S1(a): Baseline characteristics by intervention and control for full cohort and cohort for cost analysis

Variables	Responses	Full cohort		Cohort for cost analysis	
		Control	Intervention	Control	Intervention
n	215	95	120	32	33
Age, mean (SD)		56.2 (9.6)	58.9 (8.8)	61.0 (9.8)	60.5 (8.1)
Gender, n(%)	Female	32 (33.7)	60 (50.0)	10 (31.3)	17 (51.5)
	Male	63 (66.3)	60 (50.0)	22 (68.7)	16 (48.5)
Place of Birth, n(%)	Australia	59 (62.1)	66 (55.0)	17 (53.1)	20 (60.6)
	Overseas	36 (37.9)	54 (45.0)	15 (46.9)	13 (39.4)
Primary language at home, n(%)	English	88 (92.6)	96 (80.0)	31 (96.9)	25 (75.8)
	Other	7 (7.4)	24 (20.0)	1 (3.1)	8 (24.2)
Hospital admissions in past 12 months, n(%)	Yes	21 (22.1)	27 (22.5)	8 (25.0)	7 (21.2)
	No	74 (77.9)	93 (77.5)	24 (75.0)	26 (78.8)
State	NSW	35 (36.8)	99 (82.5)	6 (18.8)	28 (84.9)
	SA	60 (63.2)	21 (17.5)	26 (81.2)	5 (15.1)

Table S1(b): Outcome measurement at baseline by control and intervention for full cohort and cohort for cost analysis

Variables	Measure ¹	Full cohort		Cohort for cost analysis	
		Control	Intervention	Control	Intervention
n	215	95	120	32	33
Health literacy domain 8	Mean (SD)	4.3 (0.5)	4.0 (0.8)	4.4 (0.5)	4.3 (0.5)
	Median (IQR)	4.0 (4.0, 4.8)	4.0 (4.0, 4.6)	4.0 (4.0, 4.8)	4.0 (4.0, 4.8)
Health literacy domain 9	Mean (SD)	4.3 (0.5)	4.1 (0.7)	4.3 (0.5)	4.3 (0.5)
	Median (IQR)	4.0 (4.0, 4.8)	4.0 (4.0, 4.6)	4.1 (4.0, 4.8)	4.0 (4.0, 5.0)
eHealth	Mean (SD)	29.2 (6.3)	27.4 (7.3)	29.2 (6.6)	28.6 (6.0)
	Median (IQR)	32.0 (26.0, 32.0)	29.0 (23.5, 32.0)	32.0 (26.0, 32.0)	30.5 (25.5, 32.0)
Diet	Mean (SD)	3.1 (1.6)	3.2 (1.6)	3.4 (1.5)	3.3 (1.5)
	Median (IQR)	3.0 (2.0, 4.0)	3.0 (2.0, 4.0)	3.0 (2.0, 4.0)	3.0 (2.0, 5.0)
Physical activity	Mean (SD)	2.9 (2.3)	2.7 (2.5)	3.6 (2.3)	3.0 (2.6)
	Median (IQR)	2.0 (1.0, 4.0)	2.0 (1.0, 4.0)	4.0 (2.0, 4.0)	2.0 (1.0, 4.0)
BMI	Mean (SD)	34.9 (6.9)	34.7 (5.3)	31.9 (3.1)	33.8 (4.8)
	Median (IQR)	33.0 (30.3, 36.3)	33.3 (30.5, 37.2)	30.9 (29.9, 33.8)	32.3 (30.5, 35.4)
Waist	Mean (SD)	112.9 (15.2)	109.4 (13.6)	107.4 (10.1)	110.6 (14.6)
	Median (IQR)	110.0 (104.0, 121.0)	108.5 (99.0, 115.5)	107.0 (98.0, 116.0)	110.0 (100.0, 117.0)
Systolic blood pressure	Mean (SD)	130.7 (14.1)	130.6 (14.6)	127.6 (13.0)	131.3 (13.7)
	Median (IQR)	132.0 (121.0, 140.0)	131.0 (120.0, 139.0)	127.0 (120.5, 137.5)	131.5 (120.0, 140.0)
Diastolic blood pressure	Mean (SD)	81.3 (9.1)	79.2 (11.9)	79.4 (8.3)	79.5 (15.7)
	Median (IQR)	81.0 (75.5, 87.5)	80.0 (70.0, 86.0)	79.5 (74.0, 85.0)	79.0 (70.0, 89.5)

Table S1(c): Costs 12 months before and 12 months after enrolment date by control and intervention (excluding outlier)

Outcome	Timepoint	Control			Intervention			Crude DID (95% CI)
		n	Mean (SD)	Mean Diff (95% CI)	n	Mean (SD)	Mean Diff (95% CI)	
GP costs	12m before enrolment	32	\$1,109 (\$485)	Ref	32	\$912 (\$564)	Ref	Ref
	12m after enrolment	32	\$1,088 (\$683)	-\$21 (-\$248, \$207)	32	\$931 (\$579)	\$20 (-\$215, \$254)	-\$40 (-\$353, \$273)
GP & specialist costs	12m before enrolment	32	\$1,268 (\$571)	Ref	32	\$1,158 (\$677)	Ref	Ref
	12m after enrolment	32	\$1,345 (\$1,013)	\$77 (-\$247, \$400)	32	\$1,275 (\$837)	\$116 (-\$220, \$453)	-\$40 (-\$491, \$412)
PBS Costs	12m before enrolment	32	\$315 (\$403)	Ref	32	\$289 (\$366)	Ref	Ref
	12m after enrolment	32	\$328 (\$458)	\$12 (-\$52, \$77)	32	\$320 (\$479)	\$32 (-\$62, \$125)	-\$19 (-\$131, \$93)
GP & PBS costs	12m before enrolment	32	\$1,424 (\$672)	Ref	32	\$1,201 (\$754)	Ref	Ref
	12m after enrolment	32	\$1,416 (\$923)	-\$8 (-\$259, \$243)	32	\$1,252 (\$824)	\$51 (-\$217, \$319)	-\$59 (-\$412, \$293)
GP, specialist & PBS costs	12m before enrolment	32	\$1,583 (\$751)	Ref	32	\$1,447 (\$801)	Ref	Ref
	12m after enrolment	32	\$1,672 (\$1,203)	\$89 (-\$257, \$435)	32	\$1,595 (\$1,037)	\$148 (-\$205, \$502)	-\$59 (-\$535, \$417)
Number of GP visits	12m before enrolment	32	10.9 (0.9)	Ref	32	11.0 (1.1)	Ref	Ref
	12m after enrolment	32	11.3 (1.0)	0.3 (-1.2, 1.9)	32	10.7 (1.0)	-0.3 (-2.5, 2.0)	0.7 (-2.1, 3.4)

Table S1 (d): Costs and number of GP visits 12 months before and 12 months after enrolment date by control and intervention

Outcome	Timepoint	Control			Intervention			Crude DID (95% CI)
		n	Mean (SD)	Mean Diff (95% CI)	n	Mean (SD)	Mean Diff (95% CI)	
GP costs	12m before enrolment	32	\$1,109 (\$485)	Ref	33	\$897 (\$561)	Ref	Ref
	12m after enrolment	32	\$1,088 (\$683)	-\$21 (-\$248, \$207)	33	\$924 (\$571)	\$26 (-\$181, \$234)	-\$47 (-\$367, \$273)
GP & specialist costs	12m before enrolment	32	\$1,268 (\$571)	Ref	33	\$1,149 (\$669)	Ref	Ref
	12m after enrolment	32	\$1,345 (\$1,013)	\$77 (-\$247, \$400)	33	\$1,257 (\$830)	\$108 (-\$192, \$407)	-\$31 (-\$491, \$429)
PBS Costs	12m before enrolment	32	\$315 (\$403)	Ref	33	\$445 (\$969)	Ref	Ref
	12m after enrolment	32	\$328 (\$458)	\$12 (-\$52, \$77)	33	\$348 (\$497)	-\$97 (-\$362, \$167)	\$110 (-\$158, \$378)
GP & PBS costs	12m before enrolment	32	\$1,424 (\$672)	Ref	33	\$1,343 (\$1,103)	Ref	Ref
	12m after enrolment	32	\$1,416 (\$923)	-\$8 (-\$259, \$243)	33	\$1,271 (\$819)	-\$71 (-\$403, \$261)	\$63 (-\$364, \$490)
GP, specialist & PBS costs	12m before enrolment	32	\$1,583 (\$751)	Ref	33	\$1,595 (\$1,157)	Ref	Ref
	12m after enrolment	32	\$1,672 (\$1,203)	\$89 (-\$257, \$435)	33	\$1,605 (\$1,022)	\$10 (-\$397, \$417)	\$79 (-\$472, \$630)
Number of GP visits	12m before enrolment	32	10.9 (0.9)	Ref	33	10.9 (1.0)	Ref	Ref
	12m after enrolment	32	11.3 (1.0)	0.3 (-1.2, 1.9)	33	10.6 (0.9)	-0.3 (-2.5, 2.0)	0.6 (-2.0, 3.2)