

BMJ Open Concerns about and stimuli of COVID-19 vaccination hesitancy among diverse occupational groups in metropolitan areas of China: a cross-sectional study

Zhilan Xie,¹ Yinliang Tan ¹, Yuge Yan,¹ Xiaoqing Gu,² Haiying Chen,² Qian Huang,³ Zhaoxin Wang,¹ Jie Gu ,^{4,5} Jiaoling Huang ¹

To cite: Xie Z, Tan Y, Yan Y, *et al.* Concerns about and stimuli of COVID-19 vaccination hesitancy among diverse occupational groups in metropolitan areas of China: a cross-sectional study. *BMJ Open* 2022;**12**:e062032. doi:10.1136/bmjopen-2022-062032

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-062032>).

ZX and YT contributed equally.

Received 16 February 2022
Accepted 08 November 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Jiaoling Huang;
jiaoling_huang@sina.com

ABSTRACT

Objectives In this study, we aimed to identify concerns and stimuli regarding COVID-19 vaccination acceptance and to compare the findings by occupation.

Methods We conducted a cross-sectional study of individuals vaccinated against COVID-19 between 1 April and 30 June 2021 in four metropolitan areas of China. A total of 20 863 participants completed questionnaires, 20 767 of which were eligible for analysis. We used ordered logistic regression to assess the association of vaccination concerns and stimuli with vaccination hesitancy according to occupation.

Results Farmers were mainly concerned about the quality of vaccines (adjusted OR (aOR): 3.18, 95% CI (CI): 1.83 to 5.54). Among civil servants, media publicity reduced hesitancy (aOR: 0.44, 95% CI: 0.21 to 0.92). Among medical staff, concerns about a short duration of protective effects increased hesitancy (aOR: 8.31, 95% CI: 2.03 to 33.99). For most occupations, concerns about side effects, poor protective effects and health status increased hesitancy. In contrast, protecting oneself and protecting others acted as a stimulus to decrease hesitancy. Interestingly, 'people around me have been vaccinated' was associated with higher vaccination hesitancy among farmers (aOR: 2.19, 95% CI: 1.20 to 4.00).

Conclusion The association of vaccination concerns and stimuli with vaccination hesitancy varied by occupation. The characteristics and concerns of specific target audiences should be considered when designing informational campaigns to promote vaccination against COVID-19.

INTRODUCTION

The emergence of highly transmissible SARS-CoV-2 variants has catalysed new pandemic waves globally. However, the efficacy of vaccines against new variants is more than 70%.¹ Vaccination effectively reduces hospitalisations and deaths,² and is essential in the fight against rapid virus evolution. Cessation of the pandemic requires the achievement of high vaccination coverage in every country.³

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ In this study, we made comparisons among various occupations and included both positive and negative factors of vaccination hesitancy in statistical models.
- ⇒ This was a multicentre and community-based study, conducted in several metropolitan areas of China.
- ⇒ We analysed data collected during the initial stage of a vaccine campaign, with no follow-up observation.
- ⇒ The results may not be generalisable to rural areas or other countries and regions.

The WHO called for a 40% vaccination rate in all countries by the end of 2021. However, some countries, mainly in Africa, had not yet achieved 10% by late November of that year.^{4,5}

In addition to vaccines, countries have called for social distancing measures to reduce infection.⁶ However, this strategy does not apply to individuals in the workplace owing to often unavoidable close contact. Unvaccinated individuals may exacerbate the spread of infection in the workplace,⁷ notably in high-risk public sectors such as healthcare, service and education.^{8–11} Thus, researchers worldwide have investigated the best ways to promote vaccination in occupational settings. Earlier research has focused on specific groups, for example, healthcare workers,¹² but comparisons of vaccination hesitancy according to occupation are limited.

Improved coverage of vaccination in working populations can only be ensured with sufficient public acceptance of COVID-19 vaccines because vaccination is non-compulsory and requires ethical consideration of individual autonomy.^{13,14} Vaccination hesitancy may prolong the pandemic.¹⁵ The SAGE Working Group on Vaccine Hesitancy



defined vaccination hesitancy as ‘delay in acceptance or refusal of vaccination despite the availability of vaccination services’.¹⁶ Most studies on reducing this hesitancy have targeted the general population and reasons for the reluctance to be vaccinated,¹⁷ which stems from concerns regarding vaccine safety and efficacy.¹⁴ Less attention has been paid to increasing vaccination acceptance by explaining that the vaccines offer personal and family protection against COVID-19 infection.¹⁸ Although studies focusing on occupational groups are limited, available studies have been concerned with vaccine efficacy and safety.^{19 20} Considering the varied characteristics of information sources and lifestyles among occupational groups, disparities in attitudes regarding vaccination may be attributed to the job setting. Thus, identification of which professional groups are most vaccine hesitant and whether reasons vary according to occupation may help in developing a more targeted strategy to increase vaccination coverage.

The present study aimed to identify barriers to and stimuli of vaccination acceptance, including concerns about vaccination and reasons behind the decision to receive the vaccine. We then compared these factors according to occupation in large cities of China. The findings of our study can be helpful in designing effective COVID-19 vaccination campaigns.

METHODS

Study design

We conducted a cross-sectional study among individuals vaccinated against COVID-19 between 1 April and 30 June 2021 in four metropolitan areas of China. These cities were selected from three different city tiers, based on China’s City-Tier Classification:²¹ Shanghai and Shenzhen (tier 1), Chengdu (emerging tier 1) and Fuzhou (tier 2). The required sample size was estimated from the proportion of individuals who were highly hesitant to receive vaccination in the previous study, using the following formula.^{22 23}

$$N = \frac{Z_{\alpha/2}^2(1-p)}{\varepsilon^2 p} \times (1 + 20\%) = \frac{1.96^2 \times (1-0.10)}{0.05^2 \times 0.10} \times 1.2 = 16596$$

where N is the minimum sample size, z is the standard normal variate with 5% type I error=1.96, p is the estimate of extreme hesitancy in the population from previous research=0.10, the acceptable margin of error is 0.05 and the estimated ineligibility rate is 20%.

Data collection

We randomly selected 24 community health service centres as investigation sites, with two in each selected district (downtown, suburban and rural) for each city. Nurses and general practitioners there supported the survey. The eligibility criterion was adults receiving COVID-19 vaccination during the research period because vaccination for individuals under age 18 years was not approved at the time of the interviews. Pregnant women were also excluded from vaccination. During the

postvaccination observation period at community health service centres, investigators provided individuals whose vaccination identification number ended in eight with a link to an online questionnaire and informed consent forms. After providing informed consent, participants completed the questionnaires via smartphone, with investigators present to explain the study and answer questions. Investigators assisted participants who could not read or write in completing the questionnaires. Participants without a smartphone were provided with a printed version. Altogether, we recruited 20 863 participants. The complete questionnaire and response options used in this study are provided in online supplemental table 1.

Variables

The primary outcome variable was vaccination hesitancy, separated into five levels. The major explanatory variables included two aspects. The first was concerns about vaccination that were assumed to increase hesitancy, including concerns about side effects, poor or short duration of protective effects and the quality of vaccines, as well as concerns about ill health (poor personal health status) and the risk of infection at vaccination sites. The other aspect was stimuli, that is, reasons that hypothetically contributed to the decision to be vaccinated, including self-protection, protection for others, participants observing that most individuals around them had been vaccinated, participants observing that vaccinated people around them did not experience side effects, media publicity and community mobilisation. Participants’ occupations were mainly categorised according to the Occupational Classification of the People’s Republic of China.²⁴ Categories of education level and household income were developed based on the classification of Chinese official research.²⁵ The following covariates were included in the analysis as potential confounders: age, sex, education level, monthly household income, disability status, vaccination with a first or second dose of COVID-19 vaccine and chronic diseases. Previous research on vaccination hesitancy and influencing factors was referred to in developing the questionnaire.^{26 27} Before the formal survey, we invited experts to evaluate all questions and conducted a pilot study for questionnaire validation.

Patient and public involvement

Patients and the public were not involved in the design, conduct, reporting or dissemination plans of this research.

Statistical analyses

We used IBM SPSS V.24.0 to conduct all the statistical analyses. After removing the records of individuals with uncorrectable errors (N=96), such as age formatting, 20 767 eligible participants were included (eligibility rate: 99.54%). Descriptive indices (mean and SD or percentage) were selected according to the variable type. Charts were used to visually illustrate variations in the proportions of concerns and stimuli according to occupation. ORs and 95% CIs were calculated using ordered

Table 1 Population characteristics and the associations with vaccination hesitancy

Variables	N(%) or mean±SD	OR (95% CI)	P value
Age	30.65±10.64	0.99 (0.99 to 0.99)	<0.001
Gender			
Male	11 948 (57.53)	Ref	
Female	8819 (42.47)	1.41 (1.34 to 1.49)	<0.001
Education			
Illiteracy and primary school	304 (1.46)	Ref	
Junior high school	3645 (17.55)	0.70 (0.57 to 0.87)	0.001
Senior high school	4192 (20.19)	0.75 (0.60 to 0.92)	0.006
University and junior college	11 225 (54.05)	0.77 (0.63 to 0.95)	0.013
Master and above	1401 (6.75)	0.85 (0.68 to 1.06)	0.149
Occupation			
Farmer	550 (2.65)	Ref	
Civil servant	277 (1.33)	1.41 (1.09 to 1.84)	0.010
Teacher/student	6452 (31.07)	1.19 (1.02 to 1.40)	0.031
Medical staff	181 (0.87)	0.87 (0.64 to 1.19)	0.384
White-collar/worker	9471 (45.61)	1.21 (1.03 to 1.41)	0.018
Unemployment and else	3836 (18.47)	1.05 (0.89 to 1.24)	0.537
Monthly household income (¥)			
<5000	6246 (30.08)	Ref	
≥5000 and <10000	6247 (30.08)	0.91 (0.86 to 0.97)	0.005
≥10 000 and <20 000	3499 (16.85)	0.93 (0.86 to 1.00)	0.063
≥2000 and <50 000	2366 (11.39)	0.90 (0.82 to 0.98)	0.015
≥50 000	2409 (11.60)	0.79 (0.72 to 0.86)	<0.001
Disability			
No	20523 (98.83)	Ref	
Yes	244 (1.17)	1.35 (1.07 to 1.69)	0.011
Chronic disease			
No	16949 (81.62)	Ref	
Yes	3818 (18.38)	1.26 (1.18 to 1.34)	<0.001
Shots			
First shot	15 929 (76.70)	Ref	
Second shot	4838 (23.30)	0.66 (0.62 to 0.70)	<0.001

logistic regression to assess the relationship between vaccination hesitancy and related concerns or stimuli, also stratified by occupation. The level of statistical significance was $p < 0.05$ (two sided).

RESULTS

Descriptive statistics

The sample totalled 20 767 participants. Table 1 summarises the characteristics of the participants. The mean age was 30.65 years old (SD 10.64). More men (57.53%) than women were included in the study. More than half of the participants (54.05%) reported graduating from university or junior college and having a monthly household income below ¥10 000 (60.16%). The

main occupations were teacher or student (31.07%) and white-collar worker (45.61%). Additionally, most participants did not have any disability (98.83%) or chronic disease (81.62%).

Figure 1 illustrates the distribution of vaccination hesitancy according to six occupations. The proportion with ‘very low hesitancy’ was highest in five out of the six occupations. The proportion of participants with ‘some-what low’ and ‘medium’ hesitancy ranged from 18.73% to 38.27% and 20.99% to 27.47%, respectively. No medical staff reported ‘very high hesitancy’ for COVID-19 vaccination.

Figure 2 demonstrates the distribution of concerns about and stimuli for vaccination. Self-protection was




Figure 1 Occupational distribution of COVID-19 vaccination hesitancy (%).

the most cited stimulus (>84%) across all six occupations, and ‘people around me have been vaccinated’ was the least cited (19.09%–36.52%). The most common concerns about COVID-19 vaccination were side effects, poor protective effect and short protective effect.

Association between concerns or stimuli and vaccination hesitancy

Table 2 presents multivariate analysis of the concerns and stimuli relevant to vaccination hesitancy. Concerns about side effects, poor protective effects, the quality of vaccines, ill health and risk of infection at vaccination sites all significantly increased vaccination hesitancy. Stimuli, such as self-protection, others’ protection, media publicity and community mobilisation efforts, decreased vaccination hesitancy. Factors that did not diminish vaccine hesitancy included ‘people around me have been vaccinated’ and ‘vaccinated people around me did not experience side effects’.

Table 3 lists the results of multivariate analysis by occupation. Increased vaccination hesitancy was observed among farmers, who had concerns about side effects

Table 2 Adjusted associations between concerns or stimuli and vaccination hesitancy

Variables	aOR	95% CI lower bound	95% CI upper bound
Concerns (ref— not worry)			
Side effect	2.54	2.36	2.74
Poor protective effects	1.30	1.18	1.44
Short protective effects	1.10	1.00	1.21
Quality of vaccines	1.81	1.67	1.98
Ill health status	1.38	1.29	1.48
Risk of infection in vaccination sites	1.29	1.19	1.40
Stimuli (ref— no)			
Self-protection	0.58	0.52	0.64
Others protection	0.62	0.58	0.66
People around have been vaccinated	1.66	1.53	1.80
Vaccinated people around me did not experience side effects	1.11	1.02	1.20
Media publicity	0.75	0.69	0.81
Community mobilisation	0.70	0.65	0.76

(adjusted OR (aOR) 2.62, 95% CI 1.54 to 4.48), the quality of vaccines (aOR 3.18, 95% CI 1.83 to 5.54) and ill health (aOR 2.10, 95% CI 1.33 to 3.32). The prospect of protecting others decreased vaccination hesitancy among farmers (aOR 0.64, 95% CI 0.42 to 0.96), whereas ‘people around me have been vaccinated’ doubled the likelihood of vaccination hesitancy in this group (aOR 2.19, 95% CI 1.20 to 4.00). Among civil servants, concerns about side effects tripled the likelihood of hesitancy (aOR 3.50 95% CI 1.80 to 6.81), whereas the influence of media publicity reduced hesitancy (aOR 0.44, 95% CI 0.21 to 0.92). The significant factors for teachers and students were nearly the same as those for the whole sample population (table 2), except for ‘vaccinated people around me did not experience side effects’. Among medical personnel, concerns about vaccine side effects (aOR 4.19, 95% CI 1.46 to 12.01) and a short protective effect (aOR 8.31, 95% CI 2.03 to 33.99) increased vaccination hesitancy, and self-protection reduced hesitancy (aOR 0.23, 95% CI 0.07 to 0.80). The effects of concerns were the same as those for the total sample population (table 2) among workers, white-collar workers, unemployed participants and those in other occupations.




Figure 2 Occupational distribution of concerns about (blue) and stimuli for (green) COVID-19 vaccination (%).

DISCUSSION

We conducted this cross-sectional survey in several Chinese metropolitan areas during a large-scale COVID-19 vaccination campaign. To our knowledge, our study is the first to characterise vaccination hesitancy and stimuli for

Table 3 Adjusted associations between concerns or stimuli and vaccination hesitancy stratified by occupation

	Farmer	Civil servant	Teacher/ student	Medical staff	Worker/white- collar	Unemployment and else
	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)	aOR (95% CI)
Concerns						
Side effect	2.62 (1.54 to 4.48)	3.50 (1.80 to 6.81)	2.16 (1.88 to 2.47)	4.19 (1.46 to 12.01)	2.80 (2.51 to 3.12)	2.56 (2.14 to 3.06)
Poor protective effects	1.11 (0.62 to 1.99)	0.94 (0.39 to 2.29)	1.38 (1.14 to 1.67)	1.14 (0.25 to 5.23)	1.26 (1.10 to 1.46)	1.40 (1.11 to 1.77)
Short protective effects	0.75 (0.42 to 1.34)	1.53 (0.61 to 3.80)	0.97 (0.80 to 1.18)	8.31 (2.03 to 33.99)	1.14 (1.00 to 1.31)	1.01 (0.8 to 1.27)
Quality of vaccines	3.18 (1.83 to 5.54)	1.46 (0.67 to 3.21)	1.85 (1.57 to 2.17)	0.91 (0.36 to 2.31)	1.80 (1.59 to 2.04)	1.74 (1.43 to 2.12)
Ill health	2.10 (1.33 to 3.32)	1.16 (0.58 to 2.29)	1.49 (1.31 to 1.71)	0.94 (0.36 to 2.46)	1.32 (1.19 to 1.46)	1.39 (1.17 to 1.64)
Risk of infection in vaccination sites	1.23 (0.73 to 2.08)	1.50 (0.73 to 3.11)	1.24 (1.07 to 1.44)	1.16 (0.49 to 2.76)	1.27 (1.13 to 1.41)	1.47 (1.23 to 1.76)
Stimuli						
Self-protection	0.72 (0.44 to 1.17)	0.94 (0.44 to 1.98)	0.48 (0.40 to 0.58)	0.23 (0.07 to 0.80)	0.53 (0.45 to 0.62)	0.81 (0.64 to 1.03)
Others protection	0.64 (0.42 to 0.96)	0.67 (0.38 to 1.19)	0.66 (0.59 to 0.75)	0.91 (0.40 to 2.05)	0.57 (0.52 to 0.63)	0.61 (0.52 to 0.7)
People around have been vaccinated	2.19 (1.20 to 4.00)	1.80 (0.71 to 4.55)	1.64 (1.43 to 1.89)	0.84 (0.27 to 2.61)	1.78 (1.57 to 2.01)	1.55 (1.28 to 1.89)
Vaccinated people around me did not experience side effects	1.16 (0.67 to 2.01)	0.51 (0.22 to 1.17)	1.12 (0.96 to 1.30)	1.08 (0.43 to 2.75)	1.11 (0.99 to 1.25)	1.06 (0.88 to 1.28)
Media publicity	0.68 (0.39 to 1.18)	0.44 (0.21 to 0.92)	0.70 (0.60 to 0.80)	1.83 (0.75 to 4.49)	0.75 (0.67 to 0.84)	0.84 (0.70 to 1.01)
Community mobilisation	0.74 (0.42 to 1.30)	1.07 (0.53 to 2.19)	0.71 (0.61 to 0.82)	0.95 (0.38 to 2.38)	0.69 (0.61 to 0.78)	0.72 (0.60 to 0.86)
<0.4	[0.4, 0.6)	[0.6, 0.8)	[0.8, 1)	Insignificant (1,2)		(3,4)
						>4

Significant factors ($p < 0.05$) are marked in blue if they increased vaccination hesitancy and in orange if they decreased vaccination hesitancy.

vaccination in diverse occupational groups of China. We identified target groups for vaccine promotion, specific concerns and stimuli that increase or decrease hesitancy. This information may be instructive for policy and practices.

The research indicated that COVID-19 vaccination acceptance and hesitancy vary among different occupations.^{28 29} Previous studies have suggested that health workers have lower hesitancy and higher acceptance of vaccination than workers in other fields.^{30 31} Our findings are consistent with this evaluation, perhaps because of the urgency involved, with medical personnel being directly exposed to virus carriers.³² We also examined attitudes among individuals with other occupations. Hesitancy was relatively high among civil servants and office workers

and relatively low among farmers and teachers/students. A descriptive study among adults in the USA found high vaccination hesitancy in the construction or extraction field, and relatively low hesitancy among educators, healthcare practitioners or technicians, as well as computer and mathematics professions.²⁹ A study in Hong Kong showed that clerical, service and sales workers were more likely to refuse the vaccine.¹⁹ Although various occupational categories were adopted by different studies, some general commonalities applied. Individuals with occupational exposure to COVID-19 and those with professional knowledge were associated with less vaccination hesitancy.³³ The differences in vaccination hesitancy among occupations reinforces that vaccination campaigns should tailor their messages according to target audiences.³⁴

Studies in the USA,^{35 36} France, Germany^{37 38} and Egypt³⁹ have reported that vaccine efficacy and safety concerns are associated with a reduced likelihood of being vaccinated. Our study adds new insight into the problem of vaccination hesitancy by identifying the stimuli for receiving vaccination. The most important stimulus for vaccination was the protection of oneself and others. Community mobilisation and media publicity reduced vaccination hesitancy. These findings suggest the importance of publicising vaccine efficacy and safety in health education.⁴⁰ The media⁴¹ and the community⁴² are vital platforms for sharing such information.

The association of concerns and stimuli with vaccination hesitancy also varied by occupation. Concerns about side effects increased hesitancy among participants in all occupations. Previous research has indicated that concerns about side effects are the primary reason for hesitancy regarding vaccines against COVID-19⁴³ and other infectious diseases.^{44 45} Except for civil servants and medical staff, the extent of hesitancy increased with concerns about vaccine quality. Concerns about poor protective effects increased hesitancy among teachers and students, office workers and the unemployed individuals. Concerns about the short duration of protective effects of the vaccine were most significantly associated with hesitancy among medical staff. The importance of vaccine efficacy in hesitancy has been previously noted.⁴⁶ Most concerns focused on the safety or efficacy of vaccines, perhaps because of the accelerated development processes of COVID-19 vaccines.⁴⁷ Thus, information on safety and efficacy should be provided and emphasised when addressing hesitancy.⁴⁸

Concerns about the risk of infection at vaccination sites increased hesitancy among teachers and students, office workers and the unemployed individuals, but not among medical staff. Vaccination sites are usually within a hospital or community health centre, which are patient care facilities; this may lead to the assumption that there is a higher risk of infection than in workplaces or residences.⁴⁹ Therefore, it is necessary to strengthen the protocols for preventing and controlling infections at vaccination sites.⁵⁰ Concerns about infection may be reduced by situating vaccination sites outside patient care facilities, for instance, at universities, factories and community centres. Concerns about personal health also increased hesitancy. For example, a history of allergic reactions to other vaccines may lead to vaccine refusal or hesitancy.⁵¹ Therefore, information about vaccine contraindications should be publicised.

As for stimuli to receive vaccination, participants in most occupations were motivated by protecting themselves and protecting others. Media publicity was most influential in promoting vaccination among civil servants. The characteristics of civil servants may lead them to focus more on the media and their public image. Media influence has been shown to contribute to rapid decisions by government officials.^{52 53} Teachers and students, office workers and unemployed participants were motivated by

community mobilisation. Shanghai and other metropolitan areas are gradually establishing functional communities, which provide exceptional health services according to diverse community characteristics and needs, achieving improvements in healthcare.⁵⁴ Thus, promoting vaccination should involve strategies tailored to the occupation and setting.

In our primary hypothesis, 'people around me have been vaccinated' would be a stimulus for vaccination according to the social network theory.⁵⁵ Interestingly, however, this factor was not a stimulus to vaccination among our participants. We assume these participants believed that vaccination would be unnecessary if enough individuals in their vicinity were vaccinated, suggesting a misunderstanding of the concept of herd immunity.⁵⁶ Another unexpected finding was that 'vaccinated people around me have not experienced side effects' was not associated with hesitancy, as we had assumed, suggesting that our participants did not look to others to dispel their concerns about the side effects of vaccination.

The present study has several limitations that may affect the interpretation of our findings. First, this was a cross-sectional study; therefore, no firm conclusion could be reached regarding concerns or stimuli as a cause of COVID-19 vaccination hesitancy. We merely analysed data collected during the initial stage of a vaccine campaign without follow-up observation. However, participants' attitudes toward COVID-19 and vaccination may change over time. Second, we recruited study participants from four representative metropolitan areas in China. Occupation, concerns and stimuli in rural areas, other regions or countries were not assessed; these may differ from our findings in large cities of China. Moreover, our focus was on factors affecting vaccination hesitancy among broad occupational categories. A more detailed classification of professions could be used in future work to help support the rationale for more targeted interventions. Finally, specific interventions were mentioned but not tested in the present study. Future studies could assess the efficacy of interventions associated with those above or even more factors, such as concerns about side effects and the influence of media, in reducing vaccination hesitancy of among specific occupational groups.

CONCLUSIONS

Promoting COVID-19 vaccination is urgent, and vaccination hesitancy must be addressed. In this study, we found that vaccination hesitancy varied by occupation and was relatively low among medical staff, farmers and teachers/students but relatively high among civil servants and office workers. Concerns about side effects, poor protective effects, risk of infection at vaccination sites, the quality of vaccines and health status increased hesitancy. In contrast, protecting oneself and others acted as a stimulus to receive vaccination in most occupations. Medical staff, in particular, were concerned about short protective effects. The positive influence of media publicity was most

notable among civil servants, and community mobilisation was effective among office workers. Interestingly, ‘people around me have been vaccinated’ and ‘vaccinated people around me have not experienced side effects’ were not factors that promoted vaccination. These findings suggest that informational campaigns encouraging vaccination should use strategies tailored to the target audience.

Author affiliations

¹School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai, China

²Xidu Community Health Service Center of Fengxian District, Shanghai, China

³Weifang Community Health Service Center of Pudong New District, Shanghai, China

⁴Department of General Practition, Zhongshan Hospital Fudan University, Shanghai, China

⁵International Medical Center, Zhong Shan Hospital, Shanghai, China

Acknowledgements The authors thank the research team members and participants for their contribution to this research.

Contributors Supervision: JH. Concept and design: JH and JG. Funding acquisition: JH. Statistical analysis: ZX and YT. Drafting of the manuscript: YT, ZX and JH. Data acquisition: JH, JG, ZX, YT and YY. Critical revision of the manuscript for important intellectual content: JH and JG. Administrative, technical or material support: XG, HC, QH and ZW.

Funding The study was funded by grant nos. 22692192300 and 21692190200 from the Science and Technology Commission of Shanghai Municipality, grant no. WK2102 from Shanghai Jiao Tong University School of Medicine, grant nos. 71904145 and 72274122 from National Natural Science Foundation of China and grant no. 19X100040062 from Shanghai Jiao Tong University.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by Fengxian District Central Hospital medical ethics committee (approval no. 2021-ethic approve-02). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Yinliang Tan <http://orcid.org/0000-0002-3441-2711>

Jie Gu <http://orcid.org/0000-0003-0543-4396>

Jiaoling Huang <http://orcid.org/0000-0003-1975-3937>

REFERENCES

1 Tartof SY, Slezak JM, Fischer H, *et al*. Effectiveness of mRNA BNT162b2 COVID-19 vaccine up to 6 months in a large integrated

- health system in the USA: a retrospective cohort study. *Lancet* 2021;398:1407–16.
- 2 Griffin JB, Haddix M, Danza P, *et al*. SARS-CoV-2 Infections and Hospitalizations Among Persons Aged ≥ 16 Years, by Vaccination Status - Los Angeles County, California, May 1–July 25, 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:1170–6.
- 3 World Health Organization. COVID-19 vaccination financing and budgeting Q&A. Available: <https://www.who.int/news/item/27-04-2021-covid-19-vaccination-financing-and-budgeting-q-a>
- 4 World Health Organization. Who, un set out steps to meet world COVID vaccination targets. Available: <https://www.who.int/news/item/07-10-2021-who-un-set-out-steps-to-meet-world-covid-vaccination-targets>
- 5 World Health Organization. Who coronavirus (COVID-19) Dashboard. Available: <https://covid19.who.int/>
- 6 Honein MA, Christie A, Rose DA, *et al*. Summary of guidance for public health strategies to address high levels of community transmission of SARS-CoV-2 and related deaths, December 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:1860–7.
- 7 Burdorf A, Porru F, Rugulies R. The COVID-19 (Coronavirus) pandemic: consequences for occupational health. *Scand J Work Environ Health* 2020;46:229–30.
- 8 Sim MR. The COVID-19 pandemic: major risks to healthcare and other workers on the front line. *Occup Environ Med* 2020;77:281–2.
- 9 Schwartz J, King C-C, Yen M-Y. Protecting healthcare workers during the coronavirus disease 2019 (COVID-19) outbreak: lessons from Taiwan’s severe acute respiratory syndrome response. *Clin Infect Dis* 2020;71:858–60.
- 10 Kakimoto K, Kamiya H, Yamagishi T, *et al*. Initial Investigation of Transmission of COVID-19 Among Crew Members During Quarantine of a Cruise Ship - Yokohama, Japan, February 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:312–3.
- 11 Lan F-Y, Wei C-F, Hsu Y-T, *et al*. Work-related COVID-19 transmission in six Asian countries/areas: a follow-up study. *PLoS One* 2020;15:e0233588.
- 12 Dror AA, Eisenbach N, Taiber S, *et al*. Vaccine hesitancy: the next challenge in the fight against COVID-19. *Eur J Epidemiol* 2020;35:775–9.
- 13 Adhikari B, Cheah PY. Vaccine hesitancy in the COVID-19 era. *Lancet Infect Dis* 2021;21:1086.
- 14 Chevallier C, Hacquin A-S, Mercier H. COVID-19 vaccine Hesitancy: shortening the last mile. *Trends Cogn Sci* 2021;25:331–3.
- 15 Robinson E, Jones A, Lesser I, *et al*. International estimates of intended uptake and refusal of COVID-19 vaccines: a rapid systematic review and meta-analysis of large nationally representative samples. *Vaccine* 2021;39:2024–34.
- 16 MacDonald NE, SAGE Working Group on Vaccine Hesitancy. Vaccine hesitancy: definition, scope and determinants. *Vaccine* 2015;33:4161–4.
- 17 Joshi A, Kaur M, Kaur R, *et al*. Predictors of COVID-19 vaccine acceptance, intention, and Hesitancy: a scoping review. *Front Public Health* 2021;9:698111.
- 18 Solis Arce JS, Warren SS, Meriggi NF, *et al*. COVID-19 vaccine acceptance and hesitancy in low- and middle-income countries. *Nat Med* 2021;27:1385–94.
- 19 Wang K, Wong EL-Y, Ho K-F, *et al*. Change of willingness to accept COVID-19 vaccine and reasons of vaccine Hesitancy of working people at different waves of local epidemic in Hong Kong, China: repeated cross-sectional surveys. *Vaccines* 2021;9:62.
- 20 Salerno L, Craxi L, Amodio E, *et al*. Factors affecting Hesitancy to mRNA and viral vector COVID-19 vaccines among college students in Italy. *Vaccines* 2021;9:927.
- 21 Wong D. China’s City-Tier Classification: How Does it Work? Available: <https://www.china-briefing.com/news/chinas-city-tier-classification-defined/?hilitte=%27City-Tier%27%2C%27Classification%27>
- 22 Fisher KA, Bloomstone SJ, Walder J, *et al*. Attitudes Toward a Potential SARS-CoV-2 Vaccine : A Survey of U.S. Adults. *Ann Intern Med* 2020;173:964–73.
- 23 Sallam M. COVID-19 vaccine Hesitancy worldwide: a Concise systematic review of vaccine acceptance rates. *Vaccines* 2021;9:160.
- 24 The Ministry of Human Resources and Social Security. Occupational Classification Of the People’s Republic of China (2015 Edition). Available: http://www.jiangmen.gov.cn/jmrsj/gkmlpt/content/2/2334/post_2334805.html#173
- 25 National Bureau of Statistics. Annual household income and consumption expenditure in 2021. Available: http://www.stats.gov.cn/xgsk/sjfb/zxfb2020/202201/t20220117_1826442.html
- 26 Kwok KO, Li K-K, Wei WI, *et al*. Editor’s choice: influenza vaccine uptake, COVID-19 vaccination intention and vaccine hesitancy among nurses: a survey. *Int J Nurs Stud* 2021;114:103854.



- 27 Largent EA, Persad G, Sangenito S, *et al.* Us public attitudes toward COVID-19 vaccine mandates. *JAMA Netw Open* 2020;3:e2033324.
- 28 Jiang T, Zhou X, Wang H, *et al.* COVID-19 vaccination intention and influencing factors among different occupational risk groups: a cross-sectional study. *Hum Vaccin Immunother* 2021;17:3433–40.
- 29 King WC, Rubinstein M, Reinhart A, *et al.* COVID-19 vaccine hesitancy January–May 2021 among 18–64 year old US adults by employment and occupation. *Prev Med Rep* 2021;24:101569.
- 30 Wake AD. The Willingness to Receive COVID-19 Vaccine and Its Associated Factors: "Vaccination Refusal Could Prolong the War of This Pandemic" - A Systematic Review. *Risk Manag Healthc Policy* 2021;14:2609–23.
- 31 Dorman C, Perera A, Condon C, *et al.* Factors associated with willingness to be vaccinated against COVID-19 in a large convenience sample. *J Community Health* 2021;46:1013–9.
- 32 Zigran A, Dror AA, Morozov NG, *et al.* COVID-19 vaccine acceptance among dental professionals based on employment status during the pandemic. *Front Med* 2021;8:618403.
- 33 Dzieciolowska S, Hamel D, Gadio S, *et al.* Covid-19 vaccine acceptance, hesitancy, and refusal among Canadian healthcare workers: a multicenter survey. *Am J Infect Control* 2021;49:1152–7.
- 34 Finney Rutten LJ, Zhu X, Leppin AL, *et al.* Evidence-Based strategies for clinical organizations to address COVID-19 vaccine Hesitancy. *Mayo Clin Proc* 2021;96:699–707.
- 35 Guidry JPD, Laestadius LI, Vraga EK, *et al.* Willingness to get the COVID-19 vaccine with and without emergency use Authorization. *Am J Infect Control* 2021;49:137–42.
- 36 Verger P, Scronias D, Dauby N, *et al.* Attitudes of healthcare workers towards COVID-19 vaccination: a survey in France and French-speaking parts of Belgium and Canada, 2020. *Euro Surveill* 2021;26:2002047.
- 37 Paris C, Bénézit F, Geslin M, *et al.* COVID-19 vaccine hesitancy among healthcare workers. *Infect Dis Now* 2021;51:484–7.
- 38 El-Far Cardo A, Kraus T, Kaifie A. Factors that shape people's attitudes towards the COVID-19 pandemic in Germany-The influence of media, politics and personal characteristics. *Int J Environ Res Public Health* 2021;18:7772.
- 39 Omar DI, Hani BM. Attitudes and intentions towards COVID-19 vaccines and associated factors among Egyptian adults. *J Infect Public Health* 2021;14:1481–8.
- 40 Allen JD, Feng W, Corlin L, *et al.* Why are some people reluctant to be vaccinated for COVID-19? A cross-sectional survey among U.S. adults in May–June 2020. *Prev Med Rep* 2021;24:101494.
- 41 Marco-Franco JE, Pita-Barros P, Vivas-Orts D, *et al.* COVID-19, fake news, and vaccines: should regulation be implemented? *Int J Environ Res Public Health* 2021;18:744.
- 42 Al-Mohaithef M, Padhi BK. Determinants of COVID-19 vaccine acceptance in Saudi Arabia: a web-based national survey. *J Multidiscip Healthc* 2020;13:1657–63.
- 43 Unroe KT, Evans R, Weaver L, *et al.* Willingness of long-term care staff to receive a COVID-19 vaccine: a single state survey. *J Am Geriatr Soc* 2021;69:593–9.
- 44 Riccò M, Vezzosi L, Gualerzi G, *et al.* Knowledge, attitudes and practices (KAP) towards vaccinations in the school settings: an explorative survey. *J Prev Med Hyg* 2017;58:E266–78.
- 45 Alame M, Kaddoura M, Kharroubi S, *et al.* Uptake rates, knowledge, attitudes, and practices toward seasonal influenza vaccination among healthcare workers in Lebanon. *Hum Vaccin Immunother* 2021;17:4623–31.
- 46 Chou W-YS, Budenz A. Considering emotion in COVID-19 vaccine communication: addressing vaccine Hesitancy and fostering vaccine confidence. *Health Commun* 2020;35:1718–22.
- 47 Kashte S, Gulbake A, El-Amin Iii SF, *et al.* COVID-19 vaccines: rapid development, implications, challenges and future prospects. *Hum Cell* 2021;34:711–33.
- 48 Rosenthal SL, Weiss TW, Zimet GD, *et al.* Predictors of HPV vaccine uptake among women aged 19–26: importance of a physician's recommendation. *Vaccine* 2011;29:890–5.
- 49 Nienhaus A, Hod R. COVID-19 among health workers in Germany and Malaysia. *Int J Environ Res Public Health* 2020;17:4881.
- 50 Drohan SE, Levin SA, Grenfell BT, *et al.* Incentivizing hospital infection control. *Proc Natl Acad Sci U S A* 2019;116:6221–5.
- 51 Zhao Y-M, Liu L, Sun J, *et al.* Public willingness and determinants of COVID-19 vaccination at the initial stage of mass vaccination in China. *Vaccines* 2021;9:1172.
- 52 Figenschou TU, Karlsen R, Kolltveit K, *et al.* Mounting media pressure: push and pull forces influencing agendas, resource allocation and decision-making in public bureaucracies. *Eur J Commun* 2019;34:377–94.
- 53 Garland R, Tambini D, Couldry N. Has government been mediatized? A UK perspective. *Media, Culture & Society* 2018;40:496–513.
- 54 Shanghai Municipal Health Commission. Guidelines on promoting community health services in functional communities of the City. Available: <http://wsjkw.sh.gov.cn/jcws2/20210406/146c24803e1342c3b96b6b1824356234.html>
- 55 Bastiampillai T, Allison S, Perry SW, *et al.* Social network theory and rising suicide rates in the USA. *Lancet* 2019;393:1801.
- 56 Frederiksen LSF, Zhang Y, Foged C, *et al.* The long road toward COVID-19 herd immunity: vaccine platform technologies and mass immunization strategies. *Front Immunol* 2020;11:1817.

Appendix Table 1. Questions and responses for concerns and stimuli of COVID-19 vaccination, vaccine hesitancy and other variables.

Category	Questions	Responses
Stimuli	What motivated you to get vaccinated against COVID-19 (Multiple choice)	(1) Self-protection
		(2) Protection of family members
		(3) Protection of neighbors, friends and colleagues
		(4) People around have vaccinated
		(5) Vaccinated people around me did not experience side effects
		(6) Media publicity
		(7) Community mobilization
Concerns	What are your concerns about the COVID-19 vaccination? (Multiple choice)	(1) Side effects
		(2) Poor protective effects
		(3) Short protective effects
		(4) Quality of vaccines
		(5) Ill health status
		(6) Risk of infection in vaccination sites
Vaccine hesitancy	Do you feel hesitant to get COVID-19 vaccination? (Single choice)	(1) Very low
		(2) Somewhat low
		(3) Medium
		(4) Somewhat high
		(5) Very high
Individual factors	Your age in years (Blank)	/
	What is your gender? (Single choice)	(1) Male
		(2) Female
	What is your educational level? (Single choice)	(1) Illiteracy and primary school
		(2) Junior high school
		(3) Senior high school
		(4) University and junior college
		(5) Master and above
	What is your occupation? (Single choice)	(1) Farmer
		(2) Civil servant
(3) Teacher		
(4) Medical staff		
(5) White-collar		
(6) Student		
(7) Worker		
(8) Freelance work		
(9) Housework and unemployment		
(10) Else		
What is your monthly household income(¥)? (Single choice)	(1) <5000	
	(2) >=5000 and <10000	
	(3) >=10000 and <20000	
	(4) >=20000 and <50000	
	(5) >=50000	

	(1) No chronic disease (participants cannot choose other option if he/she chose this one)
	(2) Hypertension
Do you have any chronic disease?(Multiple choice)	(3) Diabetes
	(4) Heart disease (coronary heart disease, heart attack or stroke)
	(5) Cancer
	(6) Else chronic disease
Do you have a disability? (Single choice)	(1) No
	(2) Yes
How many shots have you gotten for COVID-19 vaccination? (Single choice)	(1) One
	(2) Two
