

BMJ Open Mental health and help seeking among trauma-exposed emergency service staff: a qualitative evidence synthesis

Niklas Maximilian Auth ¹, Matthew James Booker,² Jennifer Wild ³, Ruth Riley ^{4,5}

To cite: Auth NM, Booker MJ, Wild J, *et al.* Mental health and help seeking among trauma-exposed emergency service staff: a qualitative evidence synthesis. *BMJ Open* 2022;**12**:e047814. doi:10.1136/bmjopen-2020-047814

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-047814>).

Received 09 December 2020

Accepted 17 January 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

¹College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

²School of Social and Community Medicine, University of Bristol, Bristol, UK

³Experimental Psychology, University of Oxford, Oxford, UK

⁴Social and Community Medicine, University of Bristol, Bristol, UK

⁵Department of Applied Health Research, University of Birmingham, Birmingham, UK

Correspondence to

Niklas Maximilian Auth;
nma676@student.bham.ac.uk

ABSTRACT

Objectives To identify factors and contexts that may contribute to mental health and recovery from psychological difficulties for emergency service workers (ESWs) exposed to occupational trauma, and barriers and facilitators to help-seeking behaviour among trauma-exposed ESWs.

Background ESWs are at greater risk of stressor-related psychopathology than the general population. Exposure to occupational stressors and trauma contribute to the observed rates of post-trauma psychopathology in this occupational group with implications for workforce sustainability. Types of organisational interventions offered to trauma-exposed ESWs are inconsistent across the UK, with uncertainty around how to engage staff.

Design Four databases (OVID MEDLINE, EMBASE, PsycINFO and SCOPUS) were systematically searched from 1 January 1980 to March 2020, with citation tracking and reference chaining. A modified Critical Appraisal Skills Programme tool and quality appraisal prompts were used to identify fatally flawed studies. Qualitative studies of trauma-exposure in front-line ESWs were included, and data were extracted using a customised extraction table. Included studies were analysed using thematic synthesis. **Results** A qualitative evidence synthesis was conducted with 24 qualitative studies meeting inclusion criteria, as defined by the PerSPECTiF framework. Fourteen descriptive themes emerged from this review, categorised into two overarching constructs: (1) factors contributing to mental health (such as the need for downtime, peer support and reassurance) and (2) factors influencing help-seeking behaviour (such as stigma, the content/form/mandatory nature of interventions, and mental health literacy issues including emotional awareness and education).

Conclusion ESWs reported disconnect between the organisations' cultural positioning on trauma-related mental health, the reality of undertaking the role and the perceived applicability and usefulness of trauma interventions. Following traumatic exposure, ESWs identify benefitting from recovery time and informal support from trusted colleagues. A culture which encourages help seeking and open dialogue around mental health may reduce stigma and improve recovery from mental ill health associated with trauma exposure.

Strengths and limitations of this study

- This is the first qualitative evidence synthesis of traumatisation and mental health help-seeking in emergency service workers (ESWs).
- A user advisory group of ambulance management stakeholders and ESWs was involved in the design and purpose of this research.
- Findings are drawn from pre-Covid literature, however, core themes are omnirelevant.
- Study quality varied significantly and there is a predominance of ambulance service literature in included studies.

INTRODUCTION

Emergency service workers (ESWs) consisting of members of the emergency medical services (EMS), the fire service and the police force, consistently experience poorer mental health outcomes when compared with the general population. While subtle differences exist between occupational groups within emergency service organisations (ESOs), ESWs, also referred to as first responders, experience disproportionately higher rates of post-traumatic stress disorder (PTSD), anxiety, depression and psychological distress.^{1–5}

Suicide attempts by ESWs are considerably more prevalent than the estimated rate of 0.5% in the general population.⁶ Exposure to traumatic events accounts for higher PTSD rates in the ESW cohort,² and is the second most commonly reported cause of poor mental health among UK ESWs in a 2019 survey, following excessive workload.⁷ This review will consider the terms 'traumatic incident/event' synonymously with the term 'critical incident', which is defined as: 'any event with sufficient impact to produce significant emotional reactions in people now or later', as described by Mitchell and Everly.⁸ The increased incidence of PTSD in ESW populations^{1 2} is important to appreciate as PTSD is a risk factor for suicidal ideation and

risky behaviours in civilian and military populations,^{9 10} and increases suicidal risk in ESWs.^{11 12} The wider impacts of mental ill health among ESWs include high rates of absenteeism and presenteeism, resulting in significant costs to ESOs.¹³

ESOs employ a variety of programmes to prevent the development of mental ill health in trauma-exposed staff, of which main categories include stress management, psychotherapy and health promotion.¹⁴ Single session critical incident stress debriefing (CISD) following trauma exposure is a psychological intervention with widespread historical and current use in ESOs.^{15–17} However, by the early 2000s, a substantial body of evidence demonstrated that CISD was neutral at best and harmful at worst with respect to preventing PTSD; it appears that CISD interferes with natural recovery.^{15 16 18–20} The following two interventions are emerging among UK ESOs, and are examples of prevention strategies hoped to replace traditional debriefing methods. Trauma Risk Management (TRiM) is a peer support system, delivered by trained volunteers within the organisation,²¹ who assess trauma exposed individuals for risk of mental ill health.^{22 23} The evidence surrounding TRiM's impact on users' mental health outcomes or for improving attitudes to mental ill health is inconclusive.^{23–26} Schwartz rounds, best described as a cultural change initiative, also emerging in the UK,²⁷ allow multidisciplinary healthcare staff to share and discuss non-clinical aspects of their work, such as psychosocial, ethical and emotional issues.²⁸ As of May 2020, Schwartz rounds have been adopted by four UK ambulance trusts.²⁷ Schwartz rounds are reported to improve staff psychological well-being and increase 'empathy and compassion for colleagues',²⁹ although they have not been formally evaluated. Other interventions provided by UK ESOs to support the emotional well-being of ESWs after attending to critical incidents include counselling and 24-hour helplines,^{30–34} 'defusing' programmes³⁵ which require staff who attended a critical incident to discuss facts surrounding the event in a structured group format, and peer support networks.³⁶

Despite the availability of interventions, ESWs experience barriers to mental healthcare with one-third reporting that they experience mental health stigma, a rate that is higher than the general population.³⁷ The

purpose of this review is to identify factors and contexts that may contribute to mental health and recovery from psychological difficulties experienced by ESWs exposed to occupational trauma. We were also interested in identifying barriers and facilitators to help-seeking behaviour among trauma-exposed ESWs.

When discussing methods aimed at improving mental health, 'protection' will be used in the context of trauma-exposed ESWs, while 'recovery' relates to ESWs experiencing psychological distress as a result of trauma-exposure. By identifying important contextual factors which help and hinder staff when they access support, and illuminating benefits and drawbacks of current organisational interventions, this review aims to offer qualitative insights grounded in the perceptions of ESWs, which may help ESOs in decision making about psychological support for their staff following traumatic incident exposure.

METHODS

Methodology

Qualitative evidence synthesis (QES) is a recognised method of integrating primary qualitative research findings in health and social care.³⁸ The methods of this QES are reported using the ENTREQ framework (online supplemental file 1). The research question and final search terms were ratified by a consensus panel of key stakeholders drawn from UK ambulance services and user advisory group of ESWs. This group contributed to the development and refinement of the review questions, search parameters and application of the review findings.

Eligibility criteria

In keeping with qualitative review guidance,³⁸ the PerSPEcTiF framework was used to enhance description of inclusion criteria³⁹ (table 1).

Inclusion criteria

(* indicates further information below)

(1) Study participants were front-line ESWs (studies with mixed populations of eligible participants were included); (2) The study focus was work-related psychological distress*; (3) Data collection included primary

Table 1 The PerSPEcTiF question formulation framework³⁹

Perspective	Setting	Phenomenon of interest/problem	Environment	(Optional comparison, not applicable)	Time/timing	Findings
Emergency service workers	Emergency front-line ambulance, police or fire service work	Factors influencing mental well-being and help-seeking behaviour	Poor mental health outcomes and elevated rates of mental health stigma within the emergency services		Following occupational exposure to traumatic event(s)	Emergency service workers' perceptions and experiences regarding the phenomenon of interest

qualitative interviews, focus groups or observational methods (this included mixed-methods studies with qualitative components); (4) Analysis focused around participant attitudes towards: (A) behaviour aimed at improving or protecting mental health after experiencing a traumatic event or (B) factors which ESWs find helpful or unhelpful for their mental health while experiencing work-related psychological distress; (5) Published in English and peer-reviewed.

No limits were applied to publication date or study location.

Exclusion criteria

1. Due to the unique nature of the traumatic events witnessed in this cohort,⁴⁰ studies investigating a military cohort were excluded.
2. Volunteer ESWs were not included.

*The term 'psychological distress' will be referred to according to Ridner's definition (see online supplemental appendix A).⁴¹

Search strategy

Systematic searches of the following four databases were conducted by the primary author (NA) from 1 January 1980 to March 2020: OVID MEDLINE, EMBASE, PsycINFO and SCOPUS. The first three databases were searched together using the MEDLINE database. In order to avoid duplicates, the SCOPUS database was then searched with an additional filter to exclude MEDLINE results. One reviewer (NA) screened the 13 381 article titles and/or abstracts identified by the search. The 42 full-text articles identified in this process were then independently assessed for their eligibility criteria by two reviewers (NA and RR), with the aid of RefWorks reference management software. During this process, the two reviewers agreed to exclude 17 studies which did not meet the eligibility criteria. In cases of uncertainty surrounding eligibility criteria, there was a group discussion between all authors. One additional study was excluded on the grounds of low methodological study quality; this was agreed on by all authors (figure 1).

To identify articles missed in the electronic database search, the following methods were employed: (1) Using 'related article' feature (when available), (2) Searching the titles of included studies in google scholar for citation tracking purposes and (3) Manual searching of the references of relevant studies (reference mining).

Grey literature was searched during background research for context. Two combinations of search criteria (see online supplemental appendix B) were entered into each database in order to locate relevant literature relating to help seeking and mental health recovery. The search terms were developed during a process of trial and error using qualitative guidance,⁴² other reviews in the field,^{37 43} and virtual consultations with stakeholders and coauthors.

Quality appraisal

One reviewer (NA) assessed study quality using the CASP qualitative checklist.⁴⁴ Studies scoring less than five




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram.¹⁰²

were further appraised independently by one of three reviewers (RR, MJB and JW) to determine whether they should be included in this review based on their conceptual richness, drawing on five quality appraisal prompts as recommended by Dixon-Woods *et al.*⁴⁵ One study was removed during this approach⁴⁶ and in keeping with this approach, 'signal' (the relevance of papers to the review's aims) was prioritised over 'noise' (the inverse of methodological quality).⁴⁵

Data extraction

A customised data extraction template (see online supplemental appendix C) was created using qualitative data extraction guidance⁴² and that of a similar review as a template.⁴³ In keeping with a thematic synthesis approach,⁴⁷ first order constructs (direct participant quotations) as well as the author's interpretations were extracted in separate sections to allow for a more comprehensive review.⁴² Following Thomas and Harden,⁴⁷ all data relevant to the research aims were extracted into templates by one researcher (NA).

Data synthesis

The analytical technique of thematic synthesis is an established method in the field of mental health to investigate barriers to help seeking, and to synthesise qualitative evidence about participant experiences.^{37 43 48} Thematic analysis uses inductive line-by-line coding and is focused around: intervention need, appropriateness, acceptability and effectiveness.⁴⁹ It is therefore appropriate given that the purpose of the review is to inform UK ESO guidelines. The extracted text underwent line-by-line open coding, which allows new codes to emerge from the data, rather than imposing a pre-existing framework

onto the extracted data.⁴⁷ Subsequently, first-level codes were assimilated according to their meanings, similarities and differences. The codes were then arranged in a 'hierarchical tree structure' during which some codes were renamed as new meanings were generated. For example, some quotations initially coded as 'stigma' were later agreed to more appropriately reflect 'macho culture' or 'career concerns', with 'stigma and shame' relating specifically to shaming practices. This inductive process results in 'descriptive themes'. Two reviewers (NA and RR) performed the initial coding process, but grouping into descriptive themes and interpretation of the data involved discussion between all four authors. To enhance transparency, primary quotations used to construct the themes are supplied in online supplemental appendix D. In the final stage of thematic synthesis, the descriptive themes were used to address the review questions. This process involves inferring the meaning behind the data. The final result of this process was the translation of the descriptive themes into implications for ESO well-being policy practice, which involved round table discussion between all four authors.

Patient and public involvement statement

Patients were not involved the design, conduct, dissemination or reporting of this research. However, a user advisory group of ambulance management stakeholders and ESWs was consulted regarding the design and the purpose of the research and advised on questions to include. We believed patient and public involvement to be of reduced relevance as the impetus for this review derived from the Association of Ambulance Chief Executives, who wished to address an evidence gap to inform policy decisions.

RESULTS

Overview of study characteristics

Twenty-four studies were included in this review (table 2). All but two studies^{50 51} employed qualitative methodology only. For these two mixed-methods studies, only data from sections related to their qualitative methodology were extracted. The majority of studies (16 of 24) employed a sample of ambulance personnel,^{50–65} followed by police officers^{66–69} and lastly firefighters.^{70 71} The participants of one study, in addition to ESWs, included participants who did not match the selection criteria,⁷⁰ and two studies included a mix of different types of ESWs.^{72 73} These studies were still eligible as the participant quotations were labelled with an occupational identifier to make it possible to differentiate between eligible and non-eligible participants. The objectives of studies varied widely in terms of relevance to our review aims. Ten of the 24 studies included data relevant to protecting post-traumatic incident mental health,^{51–53 55–58 61 65 73} and one study included data relevant to barriers and facilitators for help-seeking behaviour related to mental health.⁷² The remaining 13 studies^{50 54 59 60 62–64 66–71} included data which were extracted for both of these outcomes.

Study quality

Study quality varied significantly with data collection, data analysis and discussion of findings being adequately explained in the majority of studies. All studies provided adequate research aims, although fewer than half sufficiently justified the research design.^{51 53 55 59 60 65–67 69 71 72} The lowest scoring domains were recruitment and demonstrating reflexivity. In few studies the researchers adequately justified their selection of participants^{51 55 57 67 72 73} or critically examined their relationship with participants.^{58 62 63 66 66 72 73} Four studies failed to present ethical considerations.^{50 51 64 70}

The full Critical Appraisal Skills Programme (CASP) table of all included studies is provided in online supplemental file 2. Following CASP appraisal, six studies^{49 55 57 61 64 70} initially identified as weaker quality underwent further independent quality appraisal by one of three reviewers (RR, MJB and JW), during which process one study was excluded.⁴⁹

This qualitative review identified 14 descriptive themes, which are arranged in groups of higher order themes. In turn, these are grouped into one of the following two overarching constructs: 'factors contributing to mental health' or 'factors influencing mental health help-seeking behaviour'. Table 3 presents a summary of hierarchical thematic structure.

Factors contributing to mental health

The following themes describe factors which participants attribute to having a direct influence on protecting their mental health or facilitating recovery from mental ill health after traumatic incident exposure.

Organisational sources of support

Three themes were identified as being directly linked with systems put in place by the organisations employing ESWs and represent opportunities for mental health support following trauma exposure over which ESOs have control.

Time-out/downtime period

A 'time-out' or 'downtime' period refers to a period of time following a stressful call in which ESWs are temporarily placed off duty, the availability of which was inconsistent across the studies. ESWs working in organisations in which they were offered downtime by their supervisors following certain calls reported that these breaks, ranging from 30 min to 2 hours,^{53 56} were essential in order to allow them to 'decompress' in preparation for the remainder of the shift.^{53 54} ESWs found particular comfort in casual conversation with their colleagues during this time, which did not necessarily relate to the previous call.⁵³ During such discussion, humour could be employed by the group as a method of distraction and off-loading.⁵⁹ While the majority of ESWs preferred to be in the company of colleagues during this time,^{55 59 60 63} some individuals chose to make use of organisation-provided exercise equipment in order to de-stress.⁶⁵

Table 2 Study characteristics					
Title of study (authors, year)	Quality ranking against CASP criteria	Participants	Study aim(s)	Country	Method of data collection
Halpern <i>et al</i> 2009 ⁵²	5/9	N=60 4 supervisors, 54 front-line ambulance workers	To characterise critical incidents as well as elicit suggestions for interventions	Canada	Focus groups/ individual interviews
Evans <i>et al</i> 2013 ⁶⁶	6/9	N=19 Police officers	<ol style="list-style-type: none"> 1. What are police officers' experiences of supportive and unsupportive interactions following potentially traumatic incidents? 2. Do interactions differ on the basis of the context and source of support (ie, at work with colleagues and supervisors, or outside of work with family and friends)? 3. How do supportive/unsupportive interactions facilitate/hinder the processing of traumatic incidents? 	England	Semistructured interviews
Halpern <i>et al</i> 2008 ⁵³	6/9	N=60 4 supervisors, 54 front-line ambulance workers	To explore and describe Emergency Medical Technicians' (EMTs) experiences of critical incidents and views about potential interventions, in order to facilitate development of interventions that take into account EMS culture	Canada	Focus groups/ individual interviews
Jones <i>et al</i> , 2020 ⁷²	9/9	N=32 Twenty-five (78%) of the participants were active firefighters, 15 (47%) were certified EMTs, and 11 (34.4%) were certified EMTs/paramedics.	To explore factors that influenced FRs' perceptions of mental health problems and engagement in mental health services.	USA	Ethnographic individual interviews
Regehr <i>et al</i> , 2002 ⁵⁰	6/9	N=18 Paramedics	This mixed-methods study attempts to better understand factors that lead to higher levels of distress among paramedics within the theoretical framework of emotional and cognitive empathy.	Canada	Semistructured interviews
Jonsson and Segesten 2003 ⁵⁴	5/9	N=362 240 EMTs, 122 registered nurses	The aim of this phenomenological study is to uncover the essence of traumatic events experienced by Swedish ambulance personnel.	Sweden	Written reports
Jonsson and Segesten 2004 ⁵⁵	4/9	N=10 Ambulance nurses and ambulance technicians	The phenomenon approached in this study could be described as 'the way ambulance staff experience and handle traumatic events'.	Sweden	Individual interviews

Continued

Table 2 Continued

Quality ranking against CASP criteria		Participants	Study aim(s)	Country	Method of data collection	Analysis method
Title of study (authors, year)						
Regehr and Millar 2007 ⁵¹	7/9	N=17 Paramedics	This mixed-methods study involving survey design and qualitative interviews seeks further to understand the factors related to these high levels of occupational stress.	Canada	Long interviews	Constant comparative method
Jeruzal et al 2019 ⁵⁶	8/9	N=17 Paramedics and EMTs	This qualitative study was conducted to increase understanding about the difficulties of responding to paediatric calls and to obtain information about how organisations can better support EMS providers in managing potentially difficult calls.	USA	Focus groups	Directed content analysis
Strydom et al 2015 ⁶⁷	5/9	N=40 Police officials	To conduct a qualitative situational analysis by exploring the experience and specific needs with regards to trauma and trauma intervention of police officials within the North-West Province's specialist units.	South Africa	Focus groups	Thematic analysis
Haslam and Mallon 2003 ⁷⁰	4/9	N=31 11 firefighters, 8 station officers, 4 sub officers, 4 leading firefighters, 2 fire control officers, 2 area divisional officers	This preliminary study aimed to conduct an in-depth investigation of symptoms cited by fire service personnel and assess potential risk factors for mental health and PTSD.	England	Individual interviews	Thematic analysis
Fernández-Aedo et al 2017 ⁵⁷	4/9	N=13 7 EMTs, 6 ambulance nurses	To explore the experiences, emotions and coping skills among EMTs and emergency nurses after performing out-of-hospital cardiopulmonary resuscitation manoeuvres resulting in death.	Spain	Semistructured individual interviews and focus groups	Thematic analysis
Svensson and Fridlund 2008 ⁵⁸	7/9	N=25 Ambulance nurses	The purpose of this study was to describe critical incidents in which ambulance nurses experience worry in their professional life and the actions they take in order to prevent and cope with it.	Sweden	Semistructured individual interviews	Structural analysis/critical incident technique
Clompus and Albarran 2016 ⁵⁹	5/9	N=7 Paramedics or EMTs	The aim of this study was to explore the question of how paramedics 'survive' their work within the current healthcare climate.	England	Biographical narrative semistructured interview with all participants	Thematic analysis
Abelsson 2019 ⁷¹	5/9	N=35 Firefighters	The purpose of this paper is to describe firefighters' experiences of first response emergency care	Sweden	Group interviews	Interpretive qualitative content analysis

Continued

Table 2 Continued

Title of study (authors, year)	Quality ranking against CASP criteria	Participants	Study aim(s)	Country	Method of data collection	Analysis method
Douglas <i>et al</i> 2013 ⁶⁰	5/9	N=28 Paramedics	To explore paramedics' experiences and coping strategies with death notification in the field.	Canada	Focus groups	Inductive thematic analysis
Carvello <i>et al</i> 2019 ⁶¹	4/9	N=14 Ambulance nurses	The aim of the study is to explore the experiences, the opinions and feelings of EMS nursing staff in relation to the use of the peer supporting model.	Italy	Semistructured interviews	Not made explicit
Hasselqvist-Ax <i>et al</i> 2019 ⁷³	8/9	N=22 10 police officers, 12 firefighters	The aim of this interview study was to explore firefighters' and police officers' experiences of saving lives in out-of-hospital cardiac arrest in a dual dispatch programme.	Sweden	Individual interviews	Critical incident technique
Oliveira <i>et al</i> 2019 ⁶²	8/9	N=14 Ambulance personnel	The purpose of this paper is to explore, from this group perspective, sources of stress, coping strategies and support measures	Portugal	Semistructured interviews	Thematic analysis
Drewitz-Chesney 2019 ⁶³	7/9	N=8 Paramedics	The study aim was to learn about peer communication and emotional expression between paramedics in the workplace, after they respond to calls.	Canada	Semistructured individual interviews	Constructivist grounded theory
Edwards and Kotera 2020 ⁶⁸	6/9	N=5 Police officers	The study aims to explore institutional negativity and stigma in the police force towards mental ill health	UK	Semistructured individual interviews	Thematic analysis
Gallagher and McGilloway 2007 ⁶⁴	2/9	N=27 21 EMTs, 6 emergency medical controllers/dispatchers	The principal aim of this second stage of the study was to ascertain, using qualitative methods, the impact of CIs on front-line staff by allowing them to tell their own stories.	Ireland	Individual interviews	Thematic analysis
Bullock and Garland 2018 ⁶⁹	5/9	N=59 52 police officers, 2 police staff, 4 police community support officers, one special constable	The processes through which some police officers with mental ill health experience stigmatisation.	England and Wales	Phone interviews	Thematic analysis
Folwell and Kauer 2018 ⁶⁵	6/9	N=25 EMTs	This study explores the lived experience of EMTs involved in unsuccessful paediatric resuscitation efforts and how this experience affects them professionally and personally.	USA	Individual interviews	Constant comparative analysis

EMTs, emergency medical services; PTSD, post-traumatic stress disorder.

Table 3 Summary of themes

Domain	Higher level theme	Descriptive themes
Factors contributing to mental health	Organisational sources of support	<ul style="list-style-type: none"> ▶ Time out/downtime ▶ Supervisor ▶ Official peer support network
	Informal support	<ul style="list-style-type: none"> ▶ Colleagues and family ▶ Regular partner ▶ Reassurance and validation
Factors influencing help-seeking behaviour	Nature of intervention delivery	<ul style="list-style-type: none"> ▶ Mandatory versus non-mandatory ▶ Shared experiences with intervention provider
	Stigma as a barrier to help-seeking	<ul style="list-style-type: none"> ▶ Macho culture ▶ Stigma and shame ▶ Career concerns ▶ Confidentiality concerns
	Mental health literacy	<ul style="list-style-type: none"> ▶ Emotional awareness ▶ Education and stigma

When time-out opportunities were unavailable, ESWs describe rushing into the next call without having psychologically processed the previous call.⁵¹ In such circumstances, paramedics reported difficulty giving their full attention to the next call, limiting their ability to provide life-saving care.⁶⁴ Contrary to the above findings, one ambulance worker preferred to be dispatched to another call immediately following a stressful call, due to the distraction this provided.⁵³

Supervisor

ESWs have supervisors or line managers, whose roles include acting as a point of contact after a traumatic incident. In an ambulance setting, paramedics were appreciative of the 'genuine concern' shown by their managers or supervisors following a traumatic call.⁵² Concern was commonly expressed by asking paramedics how they were feeling, and providing them with an opportunity to talk.^{52 53} Not all ESWs want to be approached by their supervisor immediately following the call, as illustrated by the following quote from a paramedic:

'I don't want you to come up and get in my face and say, are you okay? Just leave me alone. Okay. Ask me in a couple of days, am I okay with the call, sort of thing.'⁵³

Occasionally supervisors were responsible for taking an ambulance crew off duty after a call.⁵³ Even if downtime opportunities were available on individual request, paramedics described not making use of the opportunity unless suggested or requested by the supervisor.⁶⁰ Conversely, a supervisor's influence may also dissuade paramedics from requesting temporary downtime, especially for newer paramedics who were fearful of perceived repercussions linked to an inability to cope.⁶⁰

Paramedics described unsupportive supervisor responses, which could include failing to recognise the traumatising effect of an incident, applying disciplinary

pressure after complicated calls,⁵¹ or showing a lack of concern for paramedics' mental well-being.^{53 62}

Official peer support network

The majority of ambulance nurses taking part in one study were in favour of peer supporters within their organisation whom the nurses described as being able to understand their distress due to their common experiences.⁶¹ Despite the apparent popularity of the service in this organisation, peer support networks in other organisations were rarely used^{59 64} with defusing occurring 'naturally within the halls' instead.⁵⁹ Participants in two studies expressed a hesitancy to make use of peer support opportunities for fear of being judged by colleagues as 'weak'.^{61 64} Other concerns centred around the competence of peer supporters, their ability to maintain confidentiality,^{61 64} and fear of overwhelming colleagues delivering the support.⁶¹

Informal support

In contrast to organisational factors, three themes emerged related to how informal social factors influence the protection of ESW mental health and recovery from mental ill health.

Colleagues and family

ESWs reported that they found it useful to talk with someone in an informal manner.^{53-55 57 59-61 63 64 66 68 70 71} One firefighter described a need to 'vent a backpack', which fills up after each call.⁷¹ Along a similar vein, one study reported suppression to be harmful in the long term since this coping style can obfuscate mental health conditions.⁶⁸ The main providers of such informal support were family members and work colleagues, but there were mixed findings in terms of preferences for support.

Many ESWs reported turning to their family members as a primary source of emotional support following difficult calls,^{50 53 56 59 66 70 72} who were capable of 'selfless

listening' without judgement,⁶⁶ and with whom ESWs felt more comfortable sharing emotional vulnerability compared with colleagues.^{53 66} However, a number of ESWs reported avoiding talking to their family members about stressful calls out of a wish to protect them from the trauma they experienced,^{56 58 59 66 70–72} although this did not apply to family members with a first responder/healthcare background, who were judged to be able to understand ESWs' traumatic experiences.^{57–59} For similar reasons, ESWs were willing to talk to certain colleagues about traumatic calls. The informal sharing of vulnerability was reserved for colleagues with whom ESWs shared a bond of trust^{57 58 63 66} and for those more likely to empathise and understand the emotional impact of the event.^{50 53 55–58} Sharing experiences with trusted colleagues provided an opportunity for reflection and to hear different interpretations of the event. The risks of disclosure included reliving distressing events, and potential for feelings to be invalidated when partners felt differently about the event.⁵⁸

Regular partner

ESWs described how having a regular work partner helped their ability to process traumatic events encountered on the job.^{51 53 63 72} A trusting relationship between partners facilitated comfortable sharing of vulnerability following traumatic calls.^{53 63} Having shared the experience, partners could emotionally support colleagues by allowing them to talk about the call and provide reassurance.^{51 72} Having a regular partner could, however, be a negative influence in the case of an unsympathetic relationship, such as partners who respond insensitively to disclosures of vulnerability.⁶³ Due to the potential stigma arising from the disclosure of vulnerability within earshot of colleagues, the process of 'defusing' between partners, following a call, commonly takes place within the private space of the ambulance, when returning to base and while awaiting the next call.⁶³

Reassurance and validation

Reassurance, provided by colleagues indicating that they would have acted in the same way^{52 54 58} or by receiving praise for their actions from their supervisor,⁵³ were valued by ESWs following traumatic incidents, especially those involving fatalities.⁵³ Reviewing the technical aspects of calls with other ESWs provided reassurance that the final outcome was unavoidable.⁷¹ In cases of suicide, learning about the preceding circumstances could provide closure for some ESWs.⁵³ Following fatal accidents, paramedics also described needing to visit family members in hospital or to attend funerals.⁵²

Factors influencing help-seeking behaviour

The following themes reflect the barriers and facilitators to help seeking following occupational traumatic exposure.

Nature of intervention delivery

Two themes emerged related to how the method in which a formal intervention is offered can influence attitudes towards engagement.

Mandatory versus non-mandatory

The decision to employ optional or mandatory organisational mental health support for ESWs following traumatic calls was raised in several of the studies and often depended on the timing of delivery following the incident. Some participants resisted mandatory organisational mental health support following traumatic calls; police officers expressed a need to 'feel in control of the decision to talk' due to the stigma surrounding any disclosure of vulnerability.⁶⁶ Mandatory interventions for ESWs could lead to a rejection of the intervention,^{53 60} as illustrated by the following paramedic quote:

'My emotions are none of your business and if I wanted to share my emotions with you, I'm going to share [them] with someone I trust...'⁶⁰

Others, however, believed mandatory interventions would reduce stigma associated with their use,^{60 66} and prevent delays to help seeking due to the stigma associated with disclosing vulnerability.⁵³ Police officers who were initially reluctant to participate felt 'calmer' and expressed gratitude after attending a mandatory counselling service.⁶⁶ EMS staff in one study suggested limiting mandatory support to certain types of incident, such as those involving children.⁵³

Shared experiences with intervention provider

Therapists with a background in the emergency services or the military, or trained peers, were preferred by ESWs⁷² because of the belief that they are more likely to understand their problems and experiences.⁶⁰ This finding was observed across the emergency services:

'Many [participants] also approved of a provider that 'knew the job,' either working with multiple FRs in the past, or even as a family member.'⁷²

Stigma as a help-seeking barrier

'Macho' culture

Stigma associated with the disclosure of emotional vulnerability related to traumatic calls, and mental health issues,^{50 53 54 59 60 62–64 66 68 70} was identified across the emergency services. A 'macho' attitude and culture acted as a key barrier to disclosure where there was an expectation to 'deal with it'.^{63 68} Disclosure of vulnerability in such a culture was perceived as a weakness and responders were viewed as unable to cope with the demands of the job as a first responder.^{53 59 60 63 66 69 70} Revealing one's feelings was perceived to be emasculating and prevented ESWs from talking about their feelings and seeking support^{60 66 70} as demonstrated by this quote from a police officer:

I think there's a real element of machismo and masculinity in the police force and it's a bit, sort of a faux

pas to admit that things have really affected you ... If I'd have come out and said 'ah you know, that really affected me badly, let's go and sit down and have a cup of tea and talk about it' I think you're straying into pink and fluffy territory there ... saying that made me feel sad' is a bit too far.⁶⁶

Stigmatising attitudes held by senior organisational members were influential as they prevented ESWs from contacting their supervisors to seek support.^{66 69} Discussing stress in this culture was described as 'taboo'.⁶⁴ ESWs often avoided talking to their colleagues about their emotions following traumatic calls.^{60 62} Police officers described how 'tough' colleagues working in such a culture have died by suicide.⁶⁸ Some organisations described a contrasting culture in which openness about emotional vulnerability was regarded as a strength.⁷¹

Two studies identified a connection between elements of the 'macho' culture described, and the gender of ESWs. Swedish ambulance nurses reported that organisations where there was a higher proportion of women fostered a culture of openness with respect to sharing vulnerability.⁵⁰ Of a small sample of seven paramedics, Clompus and AlbarranC noted that female participants were more likely than male participants to have made use of formal mental health support mechanisms.⁵⁹ The authors attribute the gender bias in accessing support in this study as being related to 'masculinised paramedic culture'.⁵⁹

Stigma and shame

This review identified that the fear of being shamed by being labelled as 'malingerers'⁶⁸ or 'the lazy and the lame'⁶⁹ resulted in presenteeism when officers remained on active duty although mentally unwell. Mental health stigma appears related to the belief that affected individuals are less competent in their responsibilities as an ESW as well as being unreliable.⁶⁹ Police officers described a common belief that 'you're on your own' working a shift with a colleague who has been open about the emotional impact of traumatic calls.⁶⁶ ESWs who have been open about their mental health diagnoses describe being labelled as 'mad'⁶⁷ or a 'crazy guy'.⁷² Such attitudes may result in shame and the avoidance of help seeking.

Career concerns

Four studies report that police officers delay help-seeking for mental illness due to concerns about the perceived impact of disclosure on their careers.⁶⁶⁻⁶⁹ Officers believed that being labelled with a mental health condition would obstruct career progression.⁶⁷⁻⁶⁹ Officers feared being removed from 'public-facing operational roles', and/or feared a reduction in pay which related to being removed from front-line duties.⁶⁸ Fear of involuntary dismissal due to disclosure and help-seeking was also reported by study participants from the fire service.⁷⁰

Confidentiality concerns

Concerns regarding confidentiality were a barrier for ESWs to formal and informal help-seeking behaviour.

Formal support services were viewed with suspicion by police officers⁶⁷ while other officers felt they might be monitored or labelled as 'weak' if they sought a referral to well-being services.⁶⁶ Concerns about confidentiality were also raised by firefighters, who, therefore, requested preference for an anonymous counselling service outside of the brigade.⁷⁰ In the EMSs, emergency medical technicians (EMTs) expressed concerns about loss of confidentiality through the organisation-provided peer support network⁶⁴; similar concerns were expressed by EMTs towards CISD.⁶⁵ There are also perceived risks of confidentiality breaches in informal settings as described by one paramedic:

I think that the stigma is you have to be very careful who you tell that it bothered you or you might get judged as weak or you might get fired.⁷²

Confidentiality concerns could also indirectly influence help-seeking or the provision of support. For example, ambulance supervisors wishing to put their crew on a time-out after a traumatic call could be dissuaded by knowing this information could be disclosed to dispatchers.⁵³

Mental health literacy

Emotional awareness

Participants and authors of the included studies recognised a need for more training and education about mental health related issues for ESWs and family members and supervisors, who may be in a better position to detect behavioural changes associated with mental ill health and could facilitate help-seeking.^{52 53 64 72} Such education should focus on increasing ESWs' ability to detect emotional or behavioural changes within themselves, which could facilitate help-seeking behaviour.^{52 64 64 67 72} ESWs expressed a desire to be informed about the types of emotions which could be triggered by work-related traumatic incidents,^{52 64} which may reduce shame associated with help seeking.⁷² Studies revealed that an inability to recognise milder mental health symptoms acted as a barrier to help seeking with participants writing them off as being 'grumpy',⁶⁸ and not recognising and admitting to emotional distress.⁵² Additionally, participants were sometimes unaware of the support services available to them^{63 67 72} and were unaware of the benefits of seeking help.⁷²

Education and stigma

While stigma may indirectly change through improving general mental health awareness, authors also emphasised the value of education to reduce organisational stigma.⁵³ ESWs recommend such education to be delivered regularly in 'brief and efficient' classes of small groups by a peer from outside of the organisation.⁷² Having an awareness of work-related mental health problems among colleagues appeared to be an important facilitator of help-seeking.^{69 72} Experienced ESWs were regarded as being influential in reducing perceived stigma by giving

permission to other responders to 'open up'.⁷¹ Police officers in one organisation, therefore, advocated for mental health 'champions'; colleagues, preferably leaders, who could model vulnerability by openly disclosing their mental illness and work-related distress.⁶⁹

DISCUSSION

This review synthesised 24 primary studies investigating ESW attitudes to help seeking and protection of mental health and recovery from mental ill health following trauma exposure. The synthesis generated 14 themes relating to 'factors contributing to mental health' and 'factors influencing mental health help-seeking behaviour'. Despite being grouped separately for increased clarity, these overarching constructs are interconnected. Both constructs explore the influence that senior organisational members have on the mental health of ESWs within their organisation. The influence could be positive, such as delivering educational sessions about mental ill health to reduce stigma, or negative, such as the finding that ESWs may not be given downtime after a critical incident. Another interconnecting area relates to the themes of 'macho culture' and 'stigma and shame', as these stigmatising attitudes contribute towards a hesitancy to use official peer support networks.

Help seeking: culture/stigma

The findings of this review support quantitative findings that fears of a breach in confidentiality related to accessing help for mental ill health and associated career repercussions pose significant stigma-related barriers to help-seeking for ESWs.³⁷ This is important as any delays in help seeking can compound or exacerbate mental ill health.^{74 75} Concerns about the impact of mental health disclosure on career progression, professional identity and competence have also been found in military personnel.^{37 43 66 76 77}

Expressing emotional vulnerability, being labelled with a mental health condition and seeking help was equated with a perception of weakness, which contributed to a 'macho' culture. Similar social norms have been described in the military,^{23 77} which is male dominated with high rates of mental health stigma.³⁷ There are, however, many other similarities between the two settings which are likely to be influential, such as 'norms and values that place a premium on self-reliance in the face of obstacles'.³⁷ Furthermore, these findings may reflect evidence that women are more likely to seek help about their mental health,⁷⁸ and corresponds with the literature suggesting that men in male-dominated professions are less likely than women to seek mental health support,⁷⁹ and experience higher rates of suicide.⁸⁰ While women experience a reduced risk of suicide in male-dominated professions, they experience a slightly increased risk in female-dominated professions,⁸⁰ highlighting the complexity of using occupational gender ratios to predict mental health help-seeking behaviours

and mental health outcomes. Occupations dominated by hyper-masculine stereotypes may also disadvantage men who do not identify with these values, therefore discouraging the very demographic who would challenge them.⁸¹ Recruiting a more diverse, emotionally literate and aware workforce, with more women may challenge macho work cultures which prevent ESWs from disclosing their vulnerability.⁸¹

This review's finding of anticipated stigma acting as a barrier to help seeking, is supported by the findings from a systematic review which found that 'stigma can potentially lead to delayed presentation in mental health-care' for ESWs,³⁷ and have an adverse influence on help seeking in civilian populations.⁷⁶ These findings are also consistent with a qualitative review of a military setting.⁴³

In terms of attitudes to mental ill, our findings are consistent with Goffman who argued that stigma is defined in and enacted through socially constructed norms. The norms of what is stigmatising and what isn't are socially constructed within a large variety of contexts and have the potential to shift.⁸² An effective way of challenging stigma associated with mental illness and changing negative perceptions is through a strategy of increasing contact^{23 83–85} in which there is equal status, the opportunity for individuals to get to know each other, information which challenges negative stereotypes, active co-operation and pursuit of a mutual goal. Such approaches may be helpful to tackle the stigma and culture which prevent ESWs from seeking help. Elements of these factors can already be recognised in organisational strategies such as Mental Health Champions and Schwartz rounds.

Help-seeking: education

This review identifies a demand for improving the mental health literacy of ESWs and describes the type of education ESWs believe would be appropriate. Mental health education has been shown to be effective at changing attitudes towards mental health disorders when aimed at large populations, smaller at-risk groups or at the individual level.^{86 87} Antistigma education was introduced in UK emergency services as part of the 'Blue Light Programme',⁷ and demonstrated that achieving antistigma change at the employee level requires sustained education efforts over a number of years.⁷

The literature suggests that certain types of events, such as paediatric fatalities, events involving multiple casualties or suicides, have a higher traumatising potential than others.^{50 70 88} The evidence suggests, however, that the process of developing a post-trauma mental health disorder among ESWs is highly individual, relating to the personal history, situation and perspective of the exposed individual,⁶⁵ and ESWs can experience trauma in different ways depending on how they contextualise the victim.⁵⁰ Questionnaires taking into account individual factors or risk factors, such as the ESW's propensity to dwell,⁸⁹ are therefore a useful resource for predicting post-trauma psychopathology for exposed individuals.⁹⁰

Protecting mental health: social support and downtime

Consistent with the wider literature, review participants identified the importance of social support following exposure to traumatic incidents.^{25 43 91 92} Downtime was positively valued by ESWs, yet is not commonly granted to ESWs in practice.^{92 93} The types of mental health outcome affected by postincident downtime is disputed. Carlier *et al* identified 'insufficient time allowed by the employer to resolve the trauma' was correlated with higher PTSD scores in police officers 3 months post-trauma, but not at 12 months.⁹⁴ Having insufficient recovery time following traumatic calls has been correlated with higher emotional exhaustion⁹³ and psychological distress.⁹⁵ These results are in contrast with the findings of a cross-sectional quantitative survey of the psychological consequences of downtime in 217 ambulance workers, which revealed increasing periods of downtime, up to and including 1 day, to be significantly associated with lower depression scores, but not with symptoms of post-traumatic stress, burnout and stress-related physical symptoms.⁹⁶

Application of review findings to organisational interventions, policy and research

This review was initiated in collaboration with key ambulance management stakeholders who expressed a need for research to guide decisions about well-being interventions for front-line staff, criteria which were expanded by this review to also include fire and police organisations. Our findings have implications for three organisational interventions which are in use or available for use within ESOs.

Trauma risk management

Consistent with qualitative findings of navy personnel regarding the implementation of TRiM,⁹⁷ this review identified a perception among ESWs that peer-support programmes are relevant and suitable for their needs, as well as concerns regarding confidentiality and competence of practitioners of peer support programmes. Peer supporters are not however regarded as possessing the same professional competence/credibility of a professional mental health professional. Of note, while review participants expressed concerns about being judged for being perceived as 'weak' by peers in a peer-support system, such a concern was not detected among naval officers towards TRiM,⁹⁷ despite widespread stigma towards help-seeking and mental health issues in the military.⁴³

Mental health champions

We identified that awareness of colleagues' mental health challenges could be an important facilitator for help seeking. The 'Blue light champions' role⁹⁸ therefore appears to be a potentially effective method for improving attitudes, although quantitative evaluation linking these roles with ESW mental health outcomes and culture change is lacking. Staff satisfaction surveys for these roles reveal a lack of support from management and insufficient time available to dedicate to the role.⁹⁹

Schwartz rounds

In terms of challenging organisational culture which prohibits help seeking due to stigma, Schwartz rounds offer staff the opportunity to disclose their vulnerability while fostering a connectedness to other staff. However, Schwartz Rounds have not been evaluated and it is unclear whether they reduce stigma, facilitate help-seeking behaviour, or alter mental health outcomes. Further evaluation of Schwartz Rounds is required.

A note about COVID-19

This is the first review to qualitatively synthesise the barriers and facilitators to help seeking in the emergency services, and front-line staff's experience of what may protect and what may hinder mental health after traumatic incident exposure. The search was conducted in March 2020, before the emergence of qualitative literature relating to the COVID-19 pandemic, offering a summary of evidence preceding this significant historical benchmark. Qualitative literature relating to ESW mental health in the context of COVID-19 necessitates analysis in its own right, due to the distinct psychological stressors brought on by the pandemic, such as fear of infecting family members.¹⁰⁰ Healthcare workers experience different mental health pressures during pandemic working,¹⁰¹ and should, therefore, be studied in a separate context to police and fire workers.

Strengths and limitations

A core strength of this review is the analysis, which includes multiple perspectives on the topic, comprising of a primary care and ambulance clinician, a medical student and two applied health researchers. A user advisory group including ESWs was consulted regarding the design and purpose of the research and advised on questions to include, broadening its applicability. Limitations include the focus on only English language studies conducted in western countries with the exception of one study.⁶⁷ As such, findings may not generalise to other countries. Study quality varied significantly with research design commonly being inadequately explained. It should be recognised that while the review has applicability to ambulance fire and police services, the majority of studies meeting the eligibility criteria were drawn from an ambulance service background. Only one reviewer performed the initial screen of studies. However, a second reviewer was recruited to independently assess potentially relevant studies against eligibility criteria. Although the included studies are drawn from pre-Covid literature, the findings are omnirelevant to the issues of traumatisation and mental health help seeking in ESWs.

Implications for further research to help inform policy

Following traumatic calls, ESWs will likely benefit from a recovery period during which they may wish to access informal support from their colleagues. ESOs should be aware of the therapeutic effects of informal support in the post-incident setting and facilitate its availability

since ESWs are unlikely to request downtime themselves. It may also be useful for supervisors to consider that the manner in which they approach ESWs for welfare concerns influences their help-seeking behaviours. It may be important to consider that ESWs report trusting professional relationships, such as regular work-partners, as being psychologically protective against the psychological consequences of occupational trauma experiences. ESWs may benefit from education enhancing their ability to recognise pathological emotional and behavioural changes associated with traumatic incident exposure for themselves and colleagues, and to be able to suggest accessing formal organisational resources and interventions as appropriate. ESOs may wish to consider that mental health champion-type roles, providing staff in such roles are adequately supported, are regarded by ESWs as valuable tools for challenging mental health stigma. Our findings may provide insights into how engagement between ESWs and official peer support networks could be improved, by focusing on promoting antistigma and by targeting barriers to help seeking.

CONCLUSION

Our review identified barriers and facilitators to help seeking, which may assist emergency medical organisations in improving staff engagement with organisational well-being interventions. This is in keeping with the organisation's responsibility to dismantle barriers to help seeking and reduce stigma related to mental ill health and vulnerability. Our review identified the importance of organisational cultures in which it was safe to be vulnerable and need for supportive and compassionate leadership.

Twitter Matthew James Booker @MatthewBooker and Jennifer Wild @DrJenWild

Acknowledgements This work was initially written as a dissertation. We would like to thank Justin Waring for being a valued source of support throughout the dissertation. Special thanks to Anna Parry for her vital contributions in the early stages of the project, and involving key ambulance stakeholders during the process of developing the project aims. We would also like to thank Marcus Auth for his assistance provided in the form of advice regarding structure of the paper, and formulating the abstract. Dr Wild's research is supported by MQ, the Wellcome Trust, and the Oxford Health NIHR Biomedical Research Centre.

Contributors NA is the main author involved in all stages of the design, analysis and drafting and writing of the paper. RR was involved in the conception of the project's scope, design, and interpretation of results, providing important supervision and mentorship throughout. RR, MJB and JW assisted with drafting the article and revising it critically for important intellectual content, providing methodological advice, quality appraisal by independently appraising lower CASP scoring studies, and final approval of the version to be published. All authors were involved in addressing comments made during the major revision, and have approved the final version of the paper ahead of publication. As guarantor, RR accepts full responsibility for the conduct of the study, had access to the data, and controlled the decision to publish.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study does not involve human participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information. No additional data are available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Niklas Maximilian Auth <http://orcid.org/0000-0002-1959-6479>

Jennifer Wild <http://orcid.org/0000-0001-5463-1711>

Ruth Riley <http://orcid.org/0000-0001-8774-5344>

REFERENCES

1. Jones S. Describing the Mental Health Profile of First Responders: A Systematic Review [Formula: see text]. *J Am Psychiatr Nurses Assoc* 2017;23:200–14.
2. Lima EdeP, Assunção Ada Ávila. [Prevalence and factors associated with Posttraumatic Stress Disorder (PTSD) in emergency workers: a systematic literature review]. *Rev Bras Epidemiol* 2011;14:217–30.
3. Berger W, Coutinho ESF, Figueira I, et al. Rescuers at risk: a systematic review and meta-regression analysis of the worldwide current prevalence and correlates of PTSD in rescue workers. *Soc Psychiatry Psychiatr Epidemiol* 2012;47:1001–11.
4. Stanley IH, Hom MA, Joiner TE. A systematic review of suicidal thoughts and behaviors among police officers, firefighters, EMTs, and paramedics. *Clin Psychol Rev* 2016;44:25–44.
5. Petrie K, Milligan-Saville J, Gayed A, et al. Prevalence of PTSD and common mental disorders amongst ambulance personnel: a systematic review and meta-analysis. *Soc Psychiatry Psychiatr Epidemiol* 2018;53:897–909.
6. Substance Abuse and Mental Health Services Administration. First Responders: Behavioral Health Concerns, Emergency Response, and Trauma. [internet], 2018. Available: <https://www.samhsa.gov/sites/default/files/dtac/supplementalresearchbulletin-firstresponders-may2018.pdf> [Accessed 14 Oct 2019].
7. Mind. Wellbeing and mental health support in the emergency services. [internet], 2019. Available: https://www.mind.org.uk/media-a/4572/20046_mind-blue-light-programme-legacy-report-v12_online.pdf [Accessed 05 Nov 2019].
8. Mitchell JT, Everly GS. *Critical incident stress debriefings: an operations manual for the prevention of traumatic stress among emergency service and disaster workers*. 2nd ed. Ellicott City, MD: Chevron Publishing Company, 1995.
9. Jakupcak M, Cook J, Imel Z, et al. Posttraumatic stress disorder as a risk factor for suicidal ideation in Iraq and Afghanistan war veterans. *J Trauma Stress* 2009;22:303–6.
10. Tarrier N, Gregg L. Suicide risk in civilian PTSD patients--predictors of suicidal ideation, planning and attempts. *Soc Psychiatry Psychiatr Epidemiol* 2004;39:655–61.
11. Maia DB, Marmar CR, Metzler T, et al. Post-Traumatic stress symptoms in an elite unit of Brazilian police officers: prevalence and impact on psychosocial functioning and on physical and mental health. *J Affect Disord* 2007;97:241–5.
12. et alCase R, Alabakis J, Bowles K-A. High-Risk occupations—Suicide: an evidence check rapid review brokered by the Sax Institute (www.saxinstitute.org.au) for the NSW Ministry of health, 2020
13. Mind. The case for support. [internet], 2017. Available: https://www.mind.org.uk/media-a/4510/blp-blueprint_case-for-support.pdf [Accessed 06 Nov 2019].
14. Antony J, Brar R, Khan PA, et al. Interventions for the prevention and management of occupational stress injury in first responders: a rapid overview of reviews. *Syst Rev* 2020;9:121.

- 15 Smith A, Roberts K. Interventions for post-traumatic stress disorder and psychological distress in emergency ambulance personnel: a review of the literature. *Emerg Med J* 2003;20:75–8.
- 16 Carlier IVE, Lamberts RD, Van Uchelen AJ, *et al.* Disaster-related post-traumatic stress in police officers: a field study of the impact of Debriefing. *Stress Med* 1998;14:143–8.
- 17 Sattler DN, Boyd B, Kirsch J. Trauma-exposed firefighters: relationships among posttraumatic growth, posttraumatic stress, resource availability, coping and critical incident stress Debriefing experience. *Stress Health* 2014;30:356–65.
- 18 Tuckey MR, Scott JE. Group critical incident stress Debriefing with emergency services personnel: a randomized controlled trial. *Anxiety Stress Coping* 2014;27:38–55.
- 19 Regel S. Post-Trauma support in the workplace: the current status and practice of critical incident stress management (CISM) and psychological Debriefing (PD) within organizations in the UK. *Occup Med* 2007;57:411–6.
- 20 Rose SC, Bisson J, Churchill R, *et al.* Psychological Debriefing for preventing post traumatic stress disorder (PTSD). *Cochrane Database Syst Rev* 2002;171:CD000560.
- 21 Whybrow D, Jones N, Greenberg N. Promoting organizational well-being: a comprehensive review of trauma risk management. *Occup Med* 2015;65:331–6.
- 22 Hunt E, Jones N, Hastings V, *et al.* Trim: an organizational response to traumatic events in Cumbria Constabulary. *Occup Med* 2013;63:549–55.
- 23 Gould M, Greenberg N, Hetherington J. Stigma and the military: evaluation of a PTSD psychoeducational program. *J Trauma Stress* 2007;20:505–15.
- 24 Jones N, Burdett H, Green K, *et al.* Trauma risk management (TRIM): promoting help seeking for mental health problems among Combat-Exposed U.K. military personnel. *Psychiatry* 2017;80:236–51.
- 25 Shakespeare-Finch J, Rees A, Armstrong D. Social support, self-efficacy, trauma and well-being in emergency medical Dispatchers. *Soc Indic Res* 2015;123:549–65.
- 26 Greenberg N, Langston V, Everitt B, *et al.* A cluster randomized controlled trial to determine the efficacy of trauma risk management (TRIM) in a military population. *J Trauma Stress* 2010;23:430–7.
- 27 The Point of Care Foundation. List of Schwartz Rounds sites. [internet]. Available: <https://www.pointofcarefoundation.org.uk/our-work/schwartz-rounds/list-of-sites/> [Accessed 19 May 2019].
- 28 Taylor C, Xyrichis A, Leamy MC, *et al.* Can Schwartz center rounds support healthcare staff with emotional challenges at work, and how do they compare with other interventions aimed at providing similar support? A systematic review and scoping reviews. *BMJ Open* 2018;8:e024254.
- 29 Maben J, Taylor C, Dawson J, *et al.* A realist informed mixed-methods evaluation of Schwartz center Rounds® in England. Health Services and Delivery Research, 2018.
- 30 HMICFRS. Fire and rescue- Staffordshire. [internet], 2019. Available: <https://www.justiceinspectorates.gov.uk/hmicfrs/frs-assessment/frs-2018/staffordshire/people/> [Accessed 05 Apr 2020].
- 31 HMICFRS. Fire and rescue- Buckinghamshire. [internet], 2019. Available: <https://www.justiceinspectorates.gov.uk/hmicfrs/frs-assessment/frs-2018/buckinghamshire/> [Accessed 05 Apr 2020].
- 32 HMICFRS. Fire and rescue- Derbyshire. [internet], 2019. Available: <https://www.justiceinspectorates.gov.uk/hmicfrs/frs-assessment/frs-2018/derbyshire/> [Accessed 05 Apr 2020].
- 33 HMICFRS. Police- Dyfed-Powys. [internet], 2019. Available: <https://www.justiceinspectorates.gov.uk/hmicfrs/peel-assessments/peel-2018/dyfed-powys/legitimacy/detailed-findings/#question-3> [Accessed 05 Apr 2020].
- 34 HMICFRS. Police- Cumbria. [internet], 2019. Available: <https://www.justiceinspectorates.gov.uk/hmicfrs/peel-assessments/peel-2018/cumbria/legitimacy/detailed-findings/#question-1> [Accessed 05 Apr 2020].
- 35 Her Majesty's Inspectorate of Constabulary and Fire & Rescue Services (HMICFRS). Fire and rescue- Gloucestershire. [internet], 2019. Available: <https://www.justiceinspectorates.gov.uk/hmicfrs/frs-assessment/frs-2018/gloucestershire/people/> [Accessed 05 Apr 2020].
- 36 UNISON. Linc Peer Support. [internet], 2016. Available: <https://lasunison.com/linc-peer-support> [Accessed 20 Apr 2020].
- 37 Haugen PT, McCrillis AM, Smid GE, *et al.* Mental health stigma and barriers to mental health care for first responders: a systematic review and meta-analysis. *J Psychiatr Res* 2017;94:125–6.
- 38 Flemming K, Noyes J. Qualitative evidence synthesis: where are we at? *Int J Qual Methods* 2021;20:160940692199327.
- 39 Booth A, Noyes J, Flemming K, *et al.* Formulating questions to explore complex interventions within qualitative evidence synthesis. *BMJ Glob Health* 2019;4:e001107.
- 40 Litz BT, Contractor AA, Rhodes C, *et al.* Distinct trauma types in military service members seeking treatment for posttraumatic stress disorder. *J Trauma Stress* 2018;31:286–96.
- 41 Ridner SH. Psychological distress: concept analysis. *J Adv Nurs* 2004;45:536–46.
- 42 Noyes J, Lewin S. Chapter 5: Extracting qualitative evidence. In: Noyes J, Booth A, Hannes K, *et al.*, eds. *Supplementary guidance for inclusion of qualitative research in Cochrane systematic reviews of interventions. version 1 (updated August 2011)*. Cochrane Collaboration Qualitative Methods Group, 2011. <http://cqrmg.cochrane.org/supplemental-handbook-guidance>
- 43 Coleman SJ, Stevelink SAM, Hatch SL, *et al.* Stigma-related barriers and facilitators to help seeking for mental health issues in the armed forces: a systematic review and thematic synthesis of qualitative literature. *Psychol Med* 2017;47:1880–92.
- 44 CASP. Qualitative checklist. [internet], 2018. Available: https://casp-uk.net/wp-content/uploads/2018/03/CASP-Qualitative-Checklist-2018_fillable_form.pdf
- 45 Dixon-Woods M, Cavers D, Agarwal S. *Conducting a critical interpretive review of the literature on access to healthcare by vulnerable groups*, 2006.
- 46 Sommerfeld A, Wagner SL, Harder HG, *et al.* Behavioral health and firefighters: an intervention and interviews with Canadian firefighters. *Journal of Loss and Trauma* 2017;22:307–24.
- 47 Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol* 2008;8:45–6.
- 48 Thomas J, Sutcliffe K, Harden A. *Children and healthy eating: a systematic review of barriers and facilitators*. London: EPPI-Centre, Social Science Research Unit, Institute of Education, University of London, 2003.
- 49 Barnett-Page E, Thomas J. Methods for the synthesis of qualitative research: a critical review. *BMC Med Res Methodol* 2009;9:59.
- 50 Regehr C, Goldberg G, Hughes J. Exposure to human tragedy, empathy, and trauma in ambulance paramedics. *Am J Orthopsychiatry* 2002;72:505–13.
- 51 Regehr C, Millar D. Situation critical: high demand, low control, and low support in paramedic organizations. *Traumatology* 2007;13:49–58.
- 52 Halpern J, Gurevich M, Schwartz B, *et al.* What makes an incident critical for ambulance workers? emotional outcomes and implications for intervention. *Work Stress* 2009;23:173–89.
- 53 Halpern J, Gurevich M, Schwartz B, *et al.* Interventions for critical incident stress in emergency medical services: a qualitative study. *Stress and Health* 2009;25:139–49.
- 54 Jonsson A, Segesten K. The meaning of traumatic events as described by nurses in ambulance service. *Accid Emerg Nurs* 2003;11:141–52.
- 55 Jonsson A, Segesten K. Guilt, shame and need for a container: a study of post-traumatic stress among ambulance personnel. *Accid Emerg Nurs* 2004;12:215–23.
- 56 Jeruzal JN, Boland LL, Frazer MS, *et al.* Emergency medical services provider perspectives on pediatric calls: a qualitative study. *Prehosp Emerg Care* 2019;23:501–9.
- 57 Fernández-Aedo I, Pérez-Urdiales I, Unanue-Arza S, *et al.* A qualitative study about experiences and emotions of emergency medical technicians and out-of-hospital emergency nurses after performing cardiopulmonary resuscitation resulting in death. *Enferm Intensiva* 2017;28:57–63.
- 58 Svensson A, Fridlund B. Experiences of and actions towards worries among ambulance nurses in their professional life: a critical incident study. *Int Emerg Nurs* 2008;16:35–42.
- 59 Clompus SR, Albarran JW. Exploring the nature of resilience in paramedic practice: a psycho-social study. *Int Emerg Nurs* 2016;28:1–7.
- 60 Douglas L, Cheskes S, Feldman M, *et al.* Paramedics' experiences with death notification: a qualitative study. *Journal of Paramedic Practice* 2012;4:533–9.
- 61 Carvello M, Zanotti F, Rubbi I, *et al.* Peer-Support: a coping strategy for nurses working at the emergency ambulance service. *Acta Biomed* 2019;90:29–37.
- 62 Oliveira AC, Neto F, Teixeira F, *et al.* Working in prehospital emergency contexts. *Int J Workplace Health Manag* 2019;12:469–82.
- 63 Drewitz-Chesney C. Exploring paramedic communication and emotional expression in the workplace after responding to emergency calls. *Australasian Journal of Paramedicine* 2019;16.

- 64 Gallagher S, McGilloway S. Living in critical times: The impact of critical incidents on frontline ambulance personnel—a qualitative perspective. *Int J Emerg Ment Health* 2007;9:215–23. -.
- 65 Folwell A, Kauer T. 'You see a baby die and you're not fine:' a case study of stress and coping strategies in volunteer emergency medical technicians. *Journal of Applied Communication Research* 2018;46:723–43.
- 66 Evans R, Pistrang N, Billings J. Police officers' experiences of supportive and unsupportive social interactions following traumatic incidents. *Eur J Psychotraumatol* 2013;4:19696.
- 67 Strydom H, Botha K, Boshoff P. An assessment of the need of police OFFICIALS for trauma intervention programmes – a qualitative approach. *Social Work/Maatskaplike Werk* 2015;51.
- 68 Edwards A-M, Kotera Y. Mental health in the UK police force: a qualitative investigation into the stigma with mental illness. *Int J Ment Health Addict* 2021;19:1116–34.
- 69 Bullock K, Garland J. Police officers, mental (ill-)health and spoiled identity. *Criminology & Criminal Justice* 2018;18:173–89.
- 70 Haslam C, Mallon K. A preliminary investigation of post-traumatic stress symptoms among firefighters. *Work Stress* 2003;17:277–85.
- 71 Abellsson A. First response emergency care – experiences described by firefighters. *International Journal of Emergency Services* 2019;8:247–58.
- 72 Jones S, Agud K, McSweeney J. Barriers and Facilitators to Seeking Mental Health Care Among First Responders: "Removing the Darkness". *J Am Psychiatr Nurses Assoc* 2020;26:43–54.
- 73 Hasselqvist-Ax I, Nordberg P, Svensson L, et al. Experiences among firefighters and police officers of responding to out-of-hospital cardiac arrest in a dual dispatch programme in Sweden: an interview study. *BMJ Open* 2019;9:e030895.
- 74 Fridgen GJ, Aston J, Gschwandtner U, et al. Help-Seeking and pathways to care in the early stages of psychosis. *Soc Psychiatry Psychiatr Epidemiol* 2013;48:1033–43.
- 75 Gronholm PC, Thornicroft G, Laurens KR, et al. Mental health-related stigma and pathways to care for people at risk of psychotic disorders or experiencing first-episode psychosis: a systematic review. *Psychol Med* 2017;47:1867–80.
- 76 Clement S, Schauman O, Graham T, et al. What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. *Psychol Med* 2015;45:11–27.
- 77 Sharp M-L, Fear NT, Rona RJ, et al. Stigma as a barrier to seeking health care among military personnel with mental health problems. *Epidemiol Rev* 2015;37:144–62.
- 78 Australian Bureau of Statistics. *National survey of mental health and wellbeing: summary of results, Australia, 2007. ABS cat. No. 4326.0.* Canberra: ABS, 2008.
- 79 Milner A, Scovelle AJ, King T. Treatment-Seeking differences for mental health problems in male- and non-male-dominated occupations: evidence from the HILDA cohort. *Epidemiol Psychiatr Sci* 2019;28:630–7.
- 80 Milner A, King T. Men's work, women's work and suicide: a retrospective mortality study in Australia. *Aust N Z J Public Health* 2019;43:27–32.
- 81 Peters K, Ryan MK, Haslam SA. Marines, medics, and machismo: lack of fit with masculine occupational stereotypes discourages men's participation. *Br J Psychol* 2015;106:635–55.
- 82 Goffman E. *Stigma: notes on the management of spoiled identity.* Englewood Cliffs, NJ: Prentice-Hall, 1963.
- 83 Dickstein BD, Vogt DS, Handa S, et al. Targeting Self-Stigma in returning military personnel and veterans: a review of intervention strategies. *Military Psychology* 2010;22:224–36.
- 84 Norms, Committee on the Science of Changing Behavioral Health Social, Board on Behavioral C, Education, Division of Behavioral and Social Sciences and, National Academies of Sciences, Engineering. *Approaches to reducing stigma.* National Academies Press (US), 2016.
- 85 Like Minds. THE POWER OF CONTACT- Project to Counter Stigma and Discrimination Associated with Mental Illness. [internet], 2005. Available: <https://www.likeminds.org.nz/assets/National-Plans/1power-of-contact.pdf> [Accessed 06 Apr 2020].
- 86 Jorm AF, Barney LJ, Christensen H, et al. Research on mental health literacy: what we know and what we still need to know. *Aust N Z J Psychiatry* 2006;40:3–5.
- 87 Jorm AF. Mental health literacy: empowering the community to take action for better mental health. *Am Psychol* 2012;67:231–44.
- 88 Jacobsson A, Backteman-Erlanson S, Brulin C, et al. Experiences of critical incidents among female and male firefighters. *Int Emerg Nurs* 2015;23:100–4.
- 89 Wild J, Smith KV, Thompson E, et al. A prospective study of pre-trauma risk factors for post-traumatic stress disorder and depression. *Psychol Med* 2016;46:2571–82.
- 90 Halpern J, Maunder RG, Schwartz B, et al. The critical incident inventory: characteristics of incidents which affect emergency medical technicians and paramedics. *BMC Emerg Med* 2012;12:10.
- 91 Brewin CR, Andrews B, Valentine JD. Meta-Analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *J Consult Clin Psychol* 2000;68:748–66.
- 92 Gouweloos-Trines J, Tyler MP, Giummarra MJ, et al. Perceived support at work after critical incidents and its relation to psychological distress: a survey among prehospital providers. *Emerg Med J* 2017;34:816–22.
- 93 Alexander DA, Klein S. Ambulance personnel and critical incidents: impact of accident and emergency work on mental health and emotional well-being. *Br J Psychiatry* 2001;178:76–81.
- 94 Carlier IV, Lamberts RD, Gersons BP. Risk factors for posttraumatic stress symptomatology in police officers: a prospective analysis. *J Nerv Ment Dis* 1997;185:498–506.
- 95 Mitchell AM, Sakraida TJ, Kameg K. Critical incident stress Debriefing: implications for best practice. *Disaster Manag Response* 2003;1:46–51.
- 96 Halpern J, Maunder RG, Schwartz B, et al. Downtime after critical incidents in emergency medical technicians/paramedics. *Biomed Res Int* 2014;2014:1–7.
- 97 Greenberg N, Langston V, Iversen AC, et al. The acceptability of 'Trauma Risk Management' within the UK Armed Forces. *Occup Med* 2011;61:184–9.
- 98 Mind. Mind Blue light champions. [internet], 2019. Available: <https://www.mind.org.uk/media-a/4843/blp-role-description.pdf> [Accessed 19 Feb 2020].
- 99 Mind. Blue Light Programme Strand 1 (Part 2). [internet], 2016. Available: <https://www.mind.org.uk/media-a/4864/strand-1-part-2.pdf> [Accessed 05 May 2020].
- 100 Cai H, Tu B, Ma J, et al. Psychological Impact and Coping Strategies of Frontline Medical Staff in Hunan Between January and March 2020 During the Outbreak of Coronavirus Disease 2019 (COVID-19) in Hubei, China. *Med Sci Monit* 2020;26:e924171.
- 101 Vagni M, Maiorano T, Giostra V, et al. Coping with COVID-19: emergency stress, secondary trauma and self-efficacy in healthcare and emergency workers in Italy. *Front Psychol* 2020;11:566912.
- 102 Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.

Online supplementary file 1: ENTREQ checklist (Enhancing transparency in reporting the synthesis of qualitative research)

No	Item	Guide and description
1	Aims	-to identify factors and contexts that may contribute to mental health and recovery from psychological difficulties for emergency service workers (ESWs) exposed to occupational trauma -to identify barriers and facilitators to help-seeking behaviour among trauma-exposed ESWs.
2	Synthesis Methodology	Thematic synthesis
3	Approach to searching	Pre-planned comprehensive search strategies, combining database searching with manual search methods.
4	Inclusion criteria	Articles were eligible for inclusion provided they met the following criteria: (1) Study participants were frontline ESWs (studies with mixed populations of eligible participants were included); (2) The study focus was work-related psychological distress*; (3) Data collection included primary qualitative interviews, focus groups or observational methods (this included mixed methods studies with qualitative components); (4) Analysis focussed around participant attitudes towards: a) behaviour aimed at improving or protecting mental health after experiencing a traumatic event OR b) factors which ESWs find helpful or unhelpful for their mental health while experiencing work-related psychological distress; (5) Published in English and peer-reviewed. No limits were applied to publication date or study location. Exclusion criteria: (1) Due to the unique nature of the traumatic events witnesses in this cohort, ⁴⁰ studies investigating a military cohort were excluded; (2) Volunteer ESWs were not included.
5	Data sources	OVID MEDLINE, EMBASE, PsycINFO, SCOPUS. To identify articles missed in the electronic database search, the following

		<p>methods were also employed:</p> <ul style="list-style-type: none"> -Using 'related article' feature (when available). -Searching the titles of included studies in google scholar for citation tracking purposes. -Manual searching of the references of relevant studies (reference chaining). <p>Grey literature was searched during background research for context but not to locate eligible studies.</p>
6	Electronic search strategy	The two combinations of search criteria (see Appendix A) were entered into each database. Figure 1 provides an overview of the database search.
7	Study screening methods	One reviewer (NA) screened the 13381 article titles and/or abstracts identified by the search. The 42 full text articles identified in this process were then independently assessed for their eligibility criteria by two reviewers (NA and RR) (Figure 1).
8	Study characteristics	Study characteristics are presented in Table 1.
9	Study selection results	PRISMA guidance was used to construct a flow diagram displaying the database searching process (Figure 1). Of the 13381 records identified once duplicates were removed, the full text of 42 articles were screened, and 24 studies were included in this qualitative synthesis.
10	Rationale for appraisal	All potentially relevant studies were quality appraised by one reviewer (NA), using the Critical Appraisal Skills Program (CASP) guidelines. Studies meeting less than 5 criteria were then subject to further scrutiny, in the form of five quality appraisal prompts developed by Dixon-Woods et al. (see 'quality appraisal' section). This second stage of appraisal was carried out independently by RR, MB or JW, and decisions relating to the inclusion of these studies in the review were made following thorough communication between reviewers and referring to pre-determined quality prompts.
11	Appraisal terms	Critical Appraisal Skills Program (CASP) guidelines and quality appraisal prompts (see 'quality appraisal' section) were used to quality appraise all included studies.

12	Appraisal process	Quality assessment was carried out primarily by one reviewer (NA), with independent verification from RR, MB or JW for ambiguous studies.
13	Appraisal results	CASP scores of included studies are provided in Table 1. Full study quality assessments are available for review if required.
14	Data extraction	A data extraction template (Appendix B) was created for this review by one reviewer (NA). Sections for first, second and third order constructs are included in the extraction template, which was filled manually by one reviewer (NA).
15	Software	RefWorks reference management software.
16	Number of reviewers	Two reviewers were involved in the coding and analysis (NA, RR).
17	Coding	Two reviewers carried out line-by-line coding, grouping of codes and generation of descriptive themes (NA and RR).
18	Study comparison	During primary readings of the studies, overarching concepts relating to the research aims were noted. The generation of new codes altered pre-existing codes.
19	Derivation of themes	Initial coding was carried out in correspondence between two reviewers (NA and RR), and was grounded in the extracted data. Thematic synthesis of initial codes was an inductive approach, and involved roundtable discussion between all four authors.
20	Quotations	Appendix C provides example literature quotations from included studies which were used to construct themes.
21	Synthesis output	This qualitative review generated 14 descriptive themes. These are grouped into either 'Factors contributing to the protection of mental health' or 'factors influencing mental health help-seeking behaviour'. Table 2 presents a summary of themes. These descriptive themes were applied to the context of UK emergency service organisations to produce implications for practice and research.

Tong A. Enhancing transparency in reporting the synthesis of qualitative research: ENTREQ. BMC Medical Research Methodology ;12(1):181-182.

Manuscript online supplementary files

Online Supplementary File 1: ENTREQ checklist (submitted separately)

Online Supplementary File 2: CASP appraisal table

Online Supplementary File 2: CASP appraisal table

Study title (Authors, year)	Aims & Methods	Research design	Sampling	Data collection	Reflexivity	Ethical issues	Data analysis	Discussion of findings	Value
<i>What makes an incident critical for ambulance workers? Emotional outcomes and implications for intervention.</i> Halpern et al. 2009 ⁴²	Research aim clearly stated, introduction uses relevant literature to successfully explain demand for the research. Qualitative methods are appropriate for investigating complicated emotions, which could not be explored in equal depth via quantitative methods.	Interviews and focus groups seem appropriate methods to address the research aims, but the authors don't specify the reasons for their choice to use both methods.	4 supervisors, 54 front-line ambulance workers. Researchers state that participants were recruited from a specific cohort at a mandatory training conference, but no detail is given about what information was provided to the participants to entice them to enter the study. The authors considered representation of all job levels and both genders	The setting of the data collection away from the workplace was stated but no justification was given for the choice of study setting or methods used. Detail was given about the length of interviews and focus groups, and the main topics of questions asked during interviews and focus groups were stated. The researchers stated they had reached saturation by the end of the study. Data obtained was in the form of	Researchers do not acknowledge how their characteristics may have impacted on the results. The researchers acknowledge that self-selection or reporting bias may have contributed to the discrepancy in findings between men and women.	Ethical approval was obtained from a research ethics board, and participants signed a consent form. No further information was given regarding any ethical safeguards in place. Lack of description of how confidentiality was maintained.	The authors described how ethnographic content analysis was used to generate thematic codes, with a constant comparative method throughout. Three authors were involved in the coding process, although only two of these authors listened to the audiotapes. Initial and final codes are described, and contradictory data are taken into account. Sufficient data are	Findings are explicitly stated and are relevant to the researchers' aims. The only measure of credibility described was the use of three analysts during the coding process. Findings are discussed with reference to wider literature, and implications for interventions for critical incident stress are made.	The authors identify promising post-incident interventions which they believe should be further researched. Wider applicability of the study findings appear to not extend beyond the ambulance cohort. The authors provide implications for interventions for critical incident stress.

			during recruitment.	audio recordings, written transcripts and written notes of non-verbal communication information as observed by the researchers.			presented to support the findings. The authors describe how textual data was not analysed for focus groups.		
<i>Police officers' experiences of supportive and unsupportive social interactions following traumatic incidents.</i> Evans et al. 2013 ⁵⁶	Research aims made clear in objectives section. Overall aim 'was to understand the types of support processes that might promote resilience'. Methods are appropriate as experiences, and other subjective outcomes are best measured qualitatively.	Semi-structured interview design appears reasonable to explore the proposed aims, but choice of qualitative study design not directly justified.	19 Police officers. A snowballing approach was used to recruit officers, which allowed detection of information-rich participants. By limiting participants to those who have served over 2 years, the chance of traumatic incident exposure was increased. Limiting participants to those without diagnosed PTSD was appropriate as the study measured resilience-promoting investigations. Non-participation was not discussed.	An interview schedule was used for the semi-structured interviews, although only the broad focus of the schedule was provided by the authors. Interviews were audio-recorded and transcribed verbatim. No information was given regarding interview setting, and data saturation was not discussed. No justification for choice of methods.	Researchers acknowledge how their role as researchers could have affected the data, giving social desirability bias as an example.	A National Health Service ethics committee granted ethical approval, and participants provided informed consent. No information is provided regarding efforts to maintain confidentiality.	The process of thematic analysis was described in detail. All themes generated can be linked with relevant first order data. Contradictory data is taken into account and present within the themes. Three authors were involved in the analytic process to minimise bias. One author had the main role in the analysis, while two others cross-referenced certain parts of their work in order to increase credibility. All 3 researchers reached a consensus on unclear issues before the final themes were decided upon.	Themes are clearly summarised, with relevant quotations to support each theme. Findings are discussed in relation to the research aims. Three researchers were involved in the analytic process.	The researchers provide detailed recommendations for further quantitative research. Furthermore, the researchers recommend similar qualitative research in a population that has a history of PTSD, so that comparisons can be made.
<i>Interventions for critical incident stress in emergency medical services: a qualitative study</i>	Clear statement of study aims, with overall study purpose 'to facilitate development of interventions that	Explanations were provided for how the focus groups and semi-structured interviews were conducted. E.g	4 supervisors, 54 front-line ambulance workers. Participants were sampled to ensure all job levels and	The setting of the data collection away from the workplace was stated but not explained. Detail was given about	Researchers do not critically examine their role, potential bias and influence of research	Ethical approval was obtained from a research ethics board, and participants signed a consent form. No further	The authors described how ethnographic content analysis was used to generate thematic codes, using a	The findings are clearly presented, using relevant participant quotations to support the authors	The research conclusions appear relevant to all EMS organisations which operate in the format of

Halpern et al. 2008 ⁴³	take into account EMS culture.' Examining experiences adequately requires qualitative methodology.	flexible interview structure 'to permit the elaboration of more in-depth or emotionally significant data' and focus group size of 4-8 members 'to maximize interactive data'. No justification given for choice of interviews and focus groups.	genders were represented. Researchers state that participants were recruited from a specific cohort at a mandatory training conference, but no detail is given about what information was provided to the participants to entice them to enter the study. The sampling process was iterative-preliminary analysis informed subsequent sampling decisions. No further participants were recruited once saturation was reached.	the length of interviews and focus groups, and the main topics of questions asked during interviews and focus groups were stated. The researchers stated they had reached saturation by the end of the study. Data obtained was in the form of audio recordings, written transcripts and written notes of non-verbal communication information as observed by the researchers.	question formulation or data collection.	information was given regarding any ethical safeguards in place.	constant comparative method to categorise themes. Initial and final codes are described. The authors explain why textual data was not analysed for focus groups. Member validation was carried out 6 months post initial data collection. Contradictory data are presented and discussed, and sufficient data are presented to support the findings.	interpretations. 3 analysts were used during the data analysis process. Member validation was carried out 6 months after data was gathered. Contradictory viewpoints are taken into account and discussed.	using supervisors. The authors provide suggestions for interventions which are of relevance to EMS organisations. The authors recommend that the interventions of: 1. Supervisor support 2. Timeout period post-incident should be further researched.
<i>Barriers and Facilitators to Seeking Mental Health Care Among First Responders: "Removing the Darkness"</i> Jones et al, 2020 ⁶²	Aims are clearly stated and the study's importance justified. The authors aims include exploring organisational culture and perceptions of individuals, for which qualitative research is appropriate.	The decision to use individual ethnographic interviews the qualitative method of choice is well explained. The authors also explain why they chose a 'community-based approach'.	32 firefighters and/or EMTs/paramedics. This was a convenience sample. The principal investigator facilitated recruitment by developing relationships with community partners who each recruited participants from	Ethnographic individual interviews were conducted with participants by the principal investigator. An interview guide was used and attached, which was developed by the PI based on the study aim and input from a qualitative methods expert,	The principal investigator acknowledged how being married to a firefighter/paramedic promoted buy in from the community during the recruitment stage.	Ethical board approval was gained. 'Prior to starting the interview, the PI reviewed the study information sheet and completed the consent process with each participant. ' Resources for further emotional support were provided to participants in case of emotional	Content analysis and constant comparison methods were used during the analytic process. Themes which emerged during data analysis were checked with participants in later interviews. Two researchers independently analysed the transcripts, and discussed the process until	Findings are presented as themes, with participant quotations included for each theme to support the author's interpretations. Credibility was enhanced by the use of more than one analyst, and member validation. Findings are discussed in relation to the	The authors make implications for practice, identifying 'need for improving education and awareness regarding duty-related MH problems.' The authors also make implications for nursing practice, and implications for future research.

			their own organisations. They did this by distributing e-mails on behalf of the principal investigator. Reasons for non-participation: 'individuals not answering or returning calls (all were called twice), scheduling conflicts'	along with feedback from community partners to strengthen trustworthiness and validity. Interviews were audio recorded and transcribed. Field notes were also conducted by the authors. Data collection setting was decided on the basis of convenience for participants. Data saturation was reached.		distress experienced during the study.	consensus was reached. (on 80% of top level codes). Exploration of new themes was stopped when saturation was reached. participant quotations are provided throughout to support the author's interpretations. The researchers acknowledge a bias risk due to the principal investigator conducting all the interviews. Sufficient data are presented to support the findings, and contradictory data are reported (such as positive and negative experiences of therapists).	original research question.	
<i>Exposure to human tragedy, empathy, and trauma in ambulance paramedics.</i> Regehr et al, 2002 ⁴⁰	Aims of study are clearly stated, and their importance justified. Qualitative methods are appropriate because the authors want to explore experiences in depth.	Choice of qualitative study design not explicitly justified.	18 paramedics. 'Purposive sampling was used to ensure that participants represented a wide range of experiences in terms of length of time with the service and types of events encountered.' Non participation	A semistructured interview guide was used to conduct interview, and the authors provided examples of guide topics. These interviews were audio-recorded and transcribed, and field notes were taken. The authors acknowledge they	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection. No mention of changing the research design as the study progressed.	No mention of ethical approval, or consent gained from participants. No attempt is made to explain how confidentiality was maintained.	Nvivo was used to aid in data analysis. Open coding was used initially to generate broad categories. Towards the later stages of the process, selective coding was used to develop a meaningful narrative of the experiences of the	Findings are presented in sections headed by themes. Author's comments are supported by first order interpretations. Triangulation with members of emergency service organization was carried out	The authors acknowledge that the study is 'not intended to reflect the views of all paramedics in all organizations; rather, it describes a particular phenomenon experienced and described by one group of paramedics and

			was not discussed.	required a larger than normal sample size to achieve saturation. No information is provided for the setting of data collection. 'Other sources of data included the notes recording the interviewer's impressions.'			workers. Interviewers also recorded field notes. Triangulation with members of emergency service organization was carried out throughout the research, and analysis of data was discussed. Two members of the research team developed the coding tree together. A third member reviewed the open and selective coding. It is not clear how the themes were derived from the data, but sufficient data are presented to support the findings. Contradictory data are included in the findings and discussed.	throughout the research, and analysis of data was discussed. Two members of the research team developed the coding tree together. A third member reviewed the open and selective coding.	points to some interesting avenues for further consideration. These include future attempts to measure cognitive and emotional empathy in paramedics and assess the impact of these strategies on posttraumatic and depressive symptoms and on social supports.'
<i>The meaning of traumatic events as described by nurse sin ambulance service</i> Jonsson et al. 2003 ⁴⁴	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	A phenomenological approach was used, and the reason for its use thoroughly explained. The use of self-reports over the use of interviews was not justified (the authors said that interviews and self-reports are	240 medical technicians and 122 registered ambulance nurses. The authors asked 500 participants of another study to write down their experiences with traumatic events. No detail is given how the sample of the	The question the participants were asked to answer was: 'Write down and describe a situation which you experienced as a traumatic event''. From the data returned by the participants, the authors selected 25% of the written	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection. No mention of changing the research design as the study progressed.	Ethical approval was granted. Consent was implied through willingness to participate in the study, but no information is given regarding a signed consent form. In addition, participants were assured that they could withdraw	Van Kaam's scientific explication was used to analyse the written stories. Detail was provided about the steps of this thematic analytic process. Only 25% of the written stories were further analysed, and	The findings are in the form of themes, supported by first order data and the author's interpretations. The findings are related to the original research question in the discussion. There is no mention of any efforts to increase the	The authors recommend specific areas for further research, limited to the ambulance sector. The organisational interventions recommended by the authors are likewise limited to the ambulance sector.

		the most common methods in empiric-phenomenological studies.	other study was recruited. Non participation is not discussed.	accounts to further analyse. These written accounts were strategically selected based on having the fullest description of a traumatic event. Details of the analytic process are provided. Saturation of data is not discussed.		from the study at any time. The questionnaire was not labelled with the name of the participant. Confidentiality was guaranteed by eliminating names or other identifying characteristics from the essays.	were strategically selected for being more data-rich. There is no information to reveal how many researchers are involved in the analytic process. Sufficient data is presented to support the findings. There is no mention of contradictory data being taken into account. The researcher does not critically appraise their own potential bias.	credibility of the findings.	
<i>Guilt, shame and need for a container: a study of post-traumatic stress among ambulance personnel</i> Jonsson et al. 2004 ⁴⁵	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	Descriptive phenomenology was 'chosen for its potential to grasp the meaning of such phenomena through the description of lived experiences.'	10 Ambulance nurses or ambulance technicians. Insufficient information on how participants were The only relevant information is that participants were strategically selected to obtain a 'variation of sex, age, educational background, and experience as ambulance staff.'	Setting for interviews was described but not justified. Interviews were open in structure, beginning with the question "tell about a traumatic event that you have experienced". 'do you mean?' or "How did you feel about that? ". The interviews, taking about an hour and a half each, were audio-taped and transcribed word by word.' Reason for choice of interview as method of data collection not	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	The study was described, the extent of participation was explained, and potential risks were explored with the participants, who were assured that they had the right to withdraw from the study at any time. The study was approved by the Ethical committee at Göteborg University. To protect participant confidentiality, no quotations were provided.	The authors provide details about the 5 steps of the analytic process, with the latter steps being repeated once the author was familiar with existing philosophy. Due to the authors not using participant quotations (to protect participant confidentiality), it is not possible to determine whether the author's findings are supported by first order interpretations. Contradictory data are discussed, but	Findings are displayed under relevant sub-headings. In the discussion section, there is some debate for and against the researcher's arguments. No attempts made to demonstrate efforts to enhance credibility.	The results can be applied to emergency medical organisations, however caution should be taken due to the small sample size and lack of information about participant demographics. The authors recommend for the insights gained by this study to be 'distributed to all ambulance managers and other relevant personnel categories. ' The authors make specific

				explained. Saturation of data not discussed.			without the direct participant quotations.		recommendations for management.
<i>Situation Critical: High Demand, Low Control, and Low Support in Paramedic Organizations</i> Regehr et al. 2007 ⁴¹	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	Long interview design was chosen 'to explore in detail the experiences of paramedics in their organizations including their roles, the demands placed on them, the control over the working environment that they experience, and the supports that they receive.'	17 paramedics. Paramedics were purposively sampled from the sample of participants partaking in the quantitative component of the study. Purposive sampling was used to ensure that participants represented a wide range of experiences in terms of length of time with the services and types of work experiences encountered.	A long interview method was used to collect data. The interviews followed an interview guide (not provided), were audio recorded and transcribed verbatim. The interviewer made their own notes during the interview. Setting of data collection not provided, saturation of data not discussed.	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	No mention of ethical approval, or consent gained from participants. No attempt is made to explain how confidentiality was maintained.	Data analysis commenced with open coding that captured a broad range of perspectives, whereas axial and selective coding facilitated the structuring of a coding framework. After the initial coding framework was developed, the transcribed interviews were imported into N*VIVO, a computer generated data analysis system, and the interview data were coded by multiple coders and subjected to detailed thematic analysis. In the final stage of analysis, constant comparative method of data analysis was implemented to compare categories and themes across respondents. Furthermore, paramedic organisations were consulted when creating the	Findings are presented in themed subheadings. Authors claim to have enhanced credibility through 'prolonged engagement and persistent observation'. No mention of more than one analyst, and qualitative findings were triangulated with quantitative results.	The authors make recommendations for EMS organisations, based on a mixture of their quantitative and qualitative results.

							research questions, and throughout the research project to guide it with feedback. Sufficient data are presented to support the findings, and contradictory data taken into account.		
<i>Emergency Medical Services Provider Perspectives on Pediatric Calls: A Qualitative Study</i> Jessica et al. 2019 ⁴⁶	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail	The use of focus groups was not explicitly justified.	17 EMS providers. ‘Paramedics and EMTs were recruited for the study through invitations circulated via weekly staff emails, flyers posted at ambulance bases, the agency’s social media page, and e-mails from the EMS chaplain. Interested participants were instructed to contact a single study investigator by phone or email ‘Selection criteria were applied to the participants in order to sample for the most relevant participants- participants had to be working for over a year in order to increase chance of	The process of focus group conduction was thoroughly described. A semi-structured guide was used by the researcher to conduct the focus groups- the same researchers conducted all the focus groups. At the end of the focus group, themes were fed back to participants, who were invited to comment. Focus groups were conducted at a large ambulance service. Audio recordings were transcribed. A second researcher was present during the focus group to take notes. Thematic saturation was reached.	The researchers reflect on the potential biases of the study in the limitations section. They acknowledge that by using discussion prompts, they may have steered the conversation away from topics that the participants would otherwise have talked about. Furthermore, they acknowledge that the fact that a single coder was responsible for coding 80% of the data alone may be a source of bias.	Participants gave informed consent and ethical approval was gained from an ethics committee. Resources were provided to the participants in case of experiencing psychological distress.	Transcripts were analysed using directed content analysis. Two reviewers independently coded 20% of the transcripts, and a single researcher coded the remaining data. Findings were validated through groups consensus. Quotations are provided for all sub-themes. Authors acknowledge the limitations of focus groups, and the potential for bias with one researcher coding 80% of the data. The authors also acknowledge how the discussion guide prompts may have altered the topic of discussion of participants from what they would otherwise have	Findings are presented as sub-themes, with quotations supporting each sub-theme in a separate table. Credibility of findings was enhanced by reviewing the focus group themes with participants at the end of the focus group. Furthermore, some of the data was reviewed by two researchers.	The findings are relevant to EMS organisations, specifically EMS leaders. Although Desired support mechanisms following difficult paediatric calls can be extrapolated to other types of traumatic events.

			traumatic incident exposure. Non participation was not discussed.				talked about. Contradictory data are taken into account.		
<i>An assessment of the need of police officials for trauma intervention programmes - a qualitative approach.</i> Boshoff et al. 2015 ⁵⁷	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	Focus groups were chosen as the method of data collection in order to 'allow the researcher to gain insight into participants' beliefs about and perceptions or accounts of a particular topic'. The researchers state: The focus groups allowed the researcher to interact systematically and simultaneously with several individuals.	40 police officials. Participants were recruited via a purposive sampling method through the health and wellbeing departments, which were used as an intermediary between the researchers and participants. Police officials were purposefully chosen considering the relevance of the topic, specifically referring to their exposure to trauma, resultant symptoms of PTS and their participation in trauma intervention programmes. Non-participation was not discussed, other than reassuring participants that non-participation would not be sanctioned.	Setting for data collection was not outlined. Three focus groups were carried out, and semi-structured interviews were performed within the focus groups. Exact questions asked are provided by the researchers. The sessions were audio recorded, transcribed and analysed. Saturation of the data is not discussed.	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	Ethical approval was gained for this study from the North-West University's ethical committee. Informed consent was taken by the researchers. 'Participants were furthermore encouraged to withdraw from the focus groups at any time should they feel uncomfortable or experience any harm or emotional consequence as a result of participating in the study.'	The authors describe an 8 step method of thematic analysis, but do not explain how they will apply this method. It is not clear how the data is used to arrive at the themes generated. Sufficient data are presented to support the findings. Contradictory data are taken into account.	Findings are presented as sub-themes, supported by sufficient quotations. There is limited discussion for and against the researchers' arguments. No efforts are made to enhance credibility of findings.	The authors give a variety of 'preliminary indicators' for which a 'purposeful psycho-social trauma intervention programme' is indicated. The generalisability of this is limited to police forces.
<i>A preliminary investigation of post-traumatic stress symptoms among firefighters</i>	Aims of study are clearly stated, and their importance justified. Qualitative methodology is	One-to-one interviews were chosen but reasons for their use over focus groups were not	11 Firefighters, 8 Station Officers, 4 Sub Officers, 4 Leading Firefighters, 2 Fire Control	'One-to-one interviews were conducted in a private room with participants. They lasted or up to 90	Researchers do not critically examine their role, potential bias and influence of research	No mention of informed consent, ethical approval or methods to ensure confidentiality.	'The interviews were recorded and fully transcribed and the data were analysed by	Findings are presented as sub-headings of themes, with participant quotations	The authors provide recommendations for fire organisations. These

Haslam et al. 2003 ⁶⁰	appropriate due to nature of the aims, which is to explore subjective experiences in detail.	explained. The interview schedule was developed in conjunction with psychologists working within the fire service, as well using relevant literature.	Officers and 2 Area Divisional Officers. No information provided on how participants were recruited. 'The sample was selected to cover the range of positions in the service and the proportion of respondents in each position broadly reflects the profile of the service.'	minutes. The interview questions were stated by the researchers. Open-ended questions would be used to explore participants' feelings towards incidents. ;The interviews were recorded and fully transcribed and the data were analysed by sorting verbatim material into emergent themes as described by Dey (1993).'' Choice of data collection methods or study setting was not justified. Saturation of data was not discussed.	question formulation or data collection. The researchers acknowledge that their small sample size may not be representative.		sorting verbatim material into emergent themes as described by Dey (1993). A second researcher independently checked the analysis to ensure analysis reliability' There is little transparency to show how the themes were developed from the primary data. Use of a second researcher enhances credibility. Direct participant quotations are sufficiently used to support the authors' interpretations. Contradictory data are taken into account.	supporting the authors' comments. Credibility is enhanced by the use of two analysts.	recommendations are concerned with efforts to improve staff wellbeing.
<i>A qualitative study about experiences and emotions of emergency medical technicians and out-of-hospital emergency nurses after performing cardiopulmonary resuscitation resulting in death</i> Fernández-Aedo et al. 2017 ⁴⁷	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	Both interviews and focus groups were used as methods of data collection, the reason for their use was not explained.	7 EMTs and 6 nurses. Snowball sampling was used to recruit the participants. As a prerequisite for their inclusion in this study, the health workers were required to have carried out at least 5 CPR techniques resulting in death over their entire professional career. No volunteer staff	Questions asked during the interviews were ' developed based on the reviewed literature and experts' opinions', but the authors do not provide a topic guide or examples of other prompts. ' The assignment of participants to' either individual semi-structured interviews or focus groups 'was	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	' The study was approved by the Ethics Committee of the University of the Basque Country.' 'echnique was based on their availability and preferences. Participants gave their written consent to participate in the study and to be recorded in audio and/or video format.'	A detailed description of the analytical process is provided. 'To ensure the quality of the interpretation and guarantee the reliability of the information obtained, a triangulation between all researchers involved in the interviews was also carried out. Any disagreement	Themes are not provided with any participant quotations. 'To ensure the quality of the interpretation and guarantee the reliability of the information obtained, a triangulation between all researchers involved in the interviews was also carried out. Any disagreement	The authors recommend greater training for health professionals when ' notifying bad news and providing psychological support to the family members of the deceased ' No recommendations for research are made.

			was included in the study. ' In order to ensure heterogeneity, the study included healthcare professionals of different ages, both genders and varying years of experience, working at different institutions.' Non participation was not discussed.	based on their availability and preferences.' 'A total of 3 health emergency technicians and 3 nurses were interviewed individually for a total of 11---35 min by two of the researchers involved in the study. The focus group was comprised by 4 EMTs and 3 nurses, and it was used to triangulate the information obtained during the individual interviews. The group session lasted 76 min and was carried out by two researchers, one acting as a moderator and the other as an observer.' Data saturation was reached.			was resolved by consensus.' No participant quotations are provided to support the authors' interpretations. Contradictory data re not taken into account.	was resolved by consensus.' Other than the use of multiple researchers in the analytic process, no other methods to increase credibility are discussed.	
<i>Experiences of and actions towards worries among ambulance nurses in their professional life: A critical incident study</i> Svensson et al. 2008 ⁴⁸	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	There is no justification for the choice of semi-structured interviews as a method of data collection.	25 ambulance nurses. Participants were strategically selected from three ambulance services based on 'socio-demographic and professional characteristics such as age,	The authors explain the structure of the interviews, giving examples of the open-ended questions which they ask to participants at the start of the interview. The authors say the	'As both the researchers and the nurses were familiar with the environment in which the study was conducted, the risk of misunderstanding during the interviews was minimised.'	'The managers of the ambulance service involved gave approval for the study to be conducted in their department.' The authors say ethical board approval is not required in Sweden under	'The interviews were also read through several times before categorizing them into sub-categories, in order to further improve security. The incidents were analysed according to	Findings are arranged into themes and sub-themes, and participant quotations are included for each theme to support the authors' conclusions. Credibility was enhanced via the	The authors recommend further research: 'This phenomenon should be studied more in-depth in order to map how the interaction with colleagues influences the worry among

			education level, sex and years in the profession'. Non participation is not discussed.	interviews took place where it best suited the participants, and three interviews were conducted over the phone. Researchers explain their use of interviews: 'Interviews were chosen as the data collection method, which allowed the respondents to describe their thoughts in more detail with the help of follow-up questions.' The authors also justify the use of the CIT method. Interviews were audio recorded and transcribed verbatim. Saturation of data was not discussed.		certain circumstances. The authors describe how informed consent was taken from participants, including assurances that data will be treated confidentially. 'The study adhered to the principles outlined in the Declaration of Helsinki'	character and content. This step was repeated several times before the end result emerged. The categorizing of incidents into sub-categories was conducted in cooperation with the second researcher, experienced both in theory and practice, which minimised the risk of subjectivity (Andersson and Nilsson, 1964). Direct quotes from the interviews strengthened the accuracy of the study.' Contradictory data were taken into account, and sufficient are presented to support the findings. Saturation of data was not discussed.	use of two researchers when categorising incidents. Authors consider contradictory data in the discussion.	ambulance nurses.' Some recommendations are also made for employers of nurses.
<i>Exploring the nature of resilience in paramedic practice: A psycho-social study</i> Clompus et al. 2016 ⁴⁹	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective	The authors used free association narrative interviewing, a technique which involves a preliminary narrative interview, followed by a semi-structured	'An advert, with a brief study outline, was placed in a regional Paramedic bulletin which was circulated electronically to staff with an invitation to	The researchers justify their use of using Free association narrative interviewing. Saturation was achieved after the 6th interview. Details of the exact two stages	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	' Ethical approval was received from a (NHS) research ethics committee and a university in the SW of England. Confidentiality and anonymity was secured through the use of	Details of the FANI analytic process were given by the researchers. 'Any discrepancies were examined and discussed until consensus was reached. Trustworthiness	Themes are presented as sub-headings, and supported by the direct participant quotations. Member validation was carried out to improve credibility.	The authors relate the findings of the research to the theories of exiting organisational interventions, eg TRiM. The authors conclude that for front-line paramedics, 'applying

	experiences in detail.	interview. 'This enabled a deeper analysis of the affective and often unconscious aspects of paramedics' lives.'	participate.' Participants had to fulfill the following criteria: -Grade of paramedic, technician or emergency care practitioner -Willing to volunteer their time Three out of the 10 individuals became unavailable, but the authors do not explain why.	of the interviewing process were provided by the authors. interviews were audio recorded and transcribed, and carried out at a place of the participants' choosing.		pseudonyms for participants, and all data were kept in a password protected personal computer with access limited to SC. Participants were made aware that they could withdraw at any time and that the anonymised data would be disseminated in various ways. Due to the potential distress that participation could inadvertently provoke, information on how to access counselling services was made available to all participants.'	and data credibility were established by several means including participants feeding into the study's aims, keeping contemporaneous notes, and sending interview transcripts and a summary of findings to each participant for verification'. The authors are not clear how many researchers are involved in the analytic process. For each theme, sufficient participant quotes are provided to support the authors' interpretations. Contradictory data were taken into account and discussed.		interventions and reviewing support mechanisms would seem to be a pressing imperative.'
<i>First response emergency care – experiences described by firefighters</i> Abellsson et al. 2019 ⁶¹	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	The authors justify their choice of group interviews, and why limits are set on group interviews: 'The group sizes of eight and nine participants were considered acceptable to moderate and managed by the researcher.	Authors don't explain how participants were recruited into the study.	35 Firefighters. Setting for group interviews was not given. Participation numbers in the group interviews were provided and discussed. Justification for methods chosen was provided. A rough structure to the interviews was provided by	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	No approval has been sought from an ethics committee, which is in accordance with Swedish law. Consent was implied through participation in the study, following the delivery of verbal 'clear information' by the researchers.	The authors describe their method of text-driven, interpretive qualitative content analysis. This involved repeated readings of the transcribed interviews, and then identification of 'meaning units', then codes were derived, and	Findings are presented as four themes, with consistent use of primary quotations to support the researchers' arguments. The findings are discussed in relation to the original research question.	The researchers contrast their findings with other research in the field, such as in the case of spouses being used as sources of support by firefighters. They also recommend areas of future research, eg involving other strands of

		Interaction in the pre-existing group is key to a successful group interview. In this study, all firefighters worked at the same fire station, which promoted interactions optimal for the research purpose'		giving questions which were asked to participants. There is no mention of saturation of the data. The authors reflected in the 'limitations' section that group interviews could lead to participants being uncomfortable.		Methods taken to protect confidentiality are described.	formed into sub-categories and categories. Sufficient data are present to support the findings. Contradictory data are discussed. It is not clear how many researchers were involved in the analytic process. There is no explicit mention of efforts to improve credibility. Authors acknowledge that 'participants in group interviews may experience a pressure within the group, resulting in similar opinions'.		emergency service workers.
<i>Paramedics' experiences with death notification: a qualitative study</i> Douglas et al. 2013 ⁵⁰	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	The authors used focus groups as their method of data collection, but did not justify their use over other qualitative methods.	28 paramedics. Participants were self-selected. Recruitment messages were delivered via 'departmental e-mail, flyers, and by word-of-mouth'. Authors don't explain why the participants selected were the most appropriate for the study, and non-participation is not discussed.	The authors provide the locations of the focus groups, and give examples of questions asked during the sessions. There is no explicit justification for why focus groups were chosen over other qualitative methods. Methods were modified during the study as follows: 'The question guide was adapted as	The researchers acknowledged that their choice of focus group location, as well as the presence of the supervisor in the focus group may have affected the answers given by participants. The researchers do not critically examine their own role.	Ethical board approval was gained, and written consent obtained. No information is provided on efforts to maintain participant confidentiality.	The authors describe an inductive approach to data analysis. Two researchers were involved in the analysis process. Themes were discussed with a further two authors, until consensus was reached. Credibility was enhanced via the use of member checking the participant responses with	The findings are presented as themes, supported by participant quotations and authors' interpretations. Credibility was enhanced by distributing results from the study to participants for the purpose of generating feedback.	The findings are discussed in relation to existing literature, and the authors make recommendations for practice, specifically regarding paramedic training.

				required by the author and the facilitator after each session to clarify some questions based on the paramedics' responses. ' Focus groups were audiotaped and transcribed. Saturation was discussed: the authors kept recruiting participants until saturation was reached.			participants at the end of focus groups. Contradictory data are taken into account, and sufficient data are presented to support the findings.		
<i>Peer-support: a coping strategy for nurses working at the Emergency Ambulance Service</i> Carvello et al. 2019 ⁵¹	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	The authors used semi-structured interviews. The authors explain their choice of questions result from pre-existing literature, but do not justify the use of interviews over other qualitative methods eg focus groups	'Participants were recruited on a voluntary basis from an emergency medical service in the north of Italy.' The sampling was 'non-probabilistic', according to the following criteria: nurses working at the emergency ambulance service "118" in a hospital in northern Italy, who had experience in the extra-hospital emergency. Non-participation is not discussed.	Data collection was at an Italian emergency service organization called 118. The justification given by the researchers for choosing this location was to choose an environment familiar to the participants so that they would feel at ease. Semi-structured interviews are carried out with participants. The authors explain how they chose their questions, by taking inspiration from similar literature, but choice of	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	Ethical approval was gained from the Bioethical Committee of the University of Bologna. Informed consent was gained from participants before participation. Efforts to protect confidentiality are not discussed.	'The interviews were conducted and analyzed by all the researchers after having been faithfully transcribed on digital text documents, reporting in brackets some relevant non-verbal gestures, and after having evaluated the nodes and relationships generated by the nVivo qualitative research software ¹² .' Although sufficient data are presented to support the findings, it is unclear how the	Findings are presented as themes. Authors discuss arguments for and against certain issues raised. One example of this is the inclusion of quotations of participants who prefer talking to a peer supporter, and those who prefer talking to a professional therapist. No mention is made of efforts to discuss the credibility of findings. Contradictory data are taken into account and discussed.	The authors relate their findings to their aims. They propose that a peer support program should be introduced in the ambulance service. No recommendations for further research are made.

				interviews over other qualitative methods is not explained. 'The interviews were audio-recorded and conducted anonymously'. Saturation of data is not discussed.			themes were generated from the data.		
<i>Experiences among firefighters and police officers of responding to out-of hospital cardiac arrest in a dual dispatch programme in Sweden: an interview study</i> Hasselqvist-Ax et al. 2019 ⁶³	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	The authors don't justify their use of semi-structured interviews over other qualitative methods.	10 police officers, 12 firefighters. Participants were purposively sampled for knowledge of two or more cardiac arrest situations, and to collect as rich descriptions as possible. Three recruitment approaches were used: '1) an invitation letter from the researchers was presented to the main collaboration group for OHCA alarms in Stockholm County; 2) on the police report for cardiac arrest alarms there was a request to contact the researchers for a voluntary interview; 3) fire stations were directly contacted for recruitment of participants.' Non-participation	The authors provide an interview guide with 7 open-ended questions, which form the basis for the semi-structured interviews. Setting for interviews: all but three took place at regular work places (not known where the others took place). Critical interview technique (CIT) was chosen as the method for data collection. The authors don't justify their use of this technique, but they justify their sample size based on the recommended sample size for CIT. Interviews were recorded and transcribed. The authors state no modification of methods during study were necessary. Saturation was	All interviews were conducted by the same author, to increase the chance that they were conducted in a similar way. The authors list the researchers' relevant strengths and past experiences, demonstrating reflexivity.	Ethical approval was obtained. Written informed consent was obtained from the participants and information was given about the possibility to withdraw from the study without any reprisal. Participants were not entitled to financial remuneration or other benefits.	'This was an interview study where data were analysed by using critical incident technique (CIT) and inductive qualitative content analysis.' The authors explain the CIT process step by step, and provide an example with a piece of interview data. To enhance credibility, the researchers discussed the analysis at each step of the process, ensuring all analyses were supported by data. Occasionally the views of participants are summarised without examples of quotations. Contradictory data are taken into account, but saturation of data is not discussed.	The findings are presented as themes. Although participant quotations were provided, the authors didn't provide quotations for all of their comments. Often, the views of participants were summarised without providing participant quotes.	The authors provide recommendations for swedish emergency organizations, such as giving indications for training of paramedics and firefighters.

			was not discussed.	discussed in relation to the critical incident technique- where 20 interviews (of 2-4 CIs per interview) provide sufficient data. The authors included 22 participants.					
<i>Working in prehospital emergency contexts: Stress, coping and support from the perspective of ambulance personnel</i> Oliveira et al. 2019 ⁵²	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	The authors don't explain why semi-structured interviews were chosen over other qualitative methods.	14 ambulance personnel. Participants were recruited from seven institutions, mainly on the basis of 'having an active status on a rescue team and being willing to talk and share their experiences'. Details of how participants were recruited are not provided, and non-participation was not discussed.	Interviews took place at the participants' local professional organisation, and were all conducted by the first author. Semi-structured interviews were conducted with participants following an interview guide, details of which are provided by the authors. The choice of semi-structured interviews was justified as it allowed the authors 'to obtain in-depth information regarding participants' prehospital emergency experiences'. Interviews were audio-recorded and transcribed. Saturation of data was achieved.	The researchers acknowledge the possibility of researcher bias during the coding process, and therefore they discuss their analyses with other researchers.	The research project was approved by the Portuguese Red Cross and by the University's Ethics Committee. Written consent to participate in the study and to audio-record the interview was obtained, as follows: 'We sent a cover letter to all selected structures, explaining the aim of the study, the procedures, the ethical issues guaranteed, the voluntary character of the participation and the possibility to withdraw at any time'.	The authors use Braun and Clark's analytic approach, and outline the process in detail. This includes explaining how themes were derived from the data. Data collection was performed until saturation was reached. Data was discussed between the researchers during coding, enhancing credibility. The researchers made efforts to reduce researcher bias: 'Furthermore, to reduce research bias, there was a concern to engage with other researchers to discuss the process of data analysis and to illustrate themes with verbatim descriptions from participants.' Sufficient data are	Findings are presented as themes. The researchers provide participant quotations to support their findings. Multiple analysts discussed the analytic process to reduce researcher bias.	The researchers make recommendations to ambulance organisations to improve the psychological wellbeing of their staff. The authors also provide multiple avenues of recommended further research.

							presented to support the findings, and contradictory data are taken into account.		
<i>Exploring paramedic communication and emotional expression in the workplace after responding to emergency calls</i> Drewitz-Chesney et al. 2019 ⁵³	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	The authors don't explain why semi-structured interviews were chosen over other qualitative methods.	8 paramedics. The recruitment strategy is explained in detail. Participants were recruited over facebook groups. 'Participants were sampled using convenience, then purposive sampling. Convenience sampling enabled initial recruitment. Snowball and quota sampling were the two forms of purposive sampling used.' Authors are not clear on which attributes are sampled for. The authors justified their choice of participants by saying that 'when sufficient information is gleaned from participants, a smaller sample size is required'	Semi-structured interviews were performed with participants, but the researchers do not justify why interviews are chosen. Interviews were conducted using Skype, and participants could choose to interview via video or audio only. An interview guide is provided. Interviews were audio recorded and transcribed, and data collection was stopped once saturation was reached.	The recruitment messages mention that the researcher previously worked as a paramedic. The researchers acknowledge the possibility of confusion and bias that can arise from this, although this was minimised by the fact that the researcher didn't work with any of the participants.	'This study received ethical approval from the University of Edinburgh's Usher Research Ethics Group. Each participant provided verbal and written consent. To minimise the risk of psychological impact, participants were never asked to recall specific calls or details. Participants were monitored for signs of distress during each interview (none were noted). BCEHS paramedics have access to three services offering support and counselling. These services were listed on their information letters. At the conclusion of each interview, participants were asked if they wanted a referral to any of the services, which	The authors used constructivist grounded theory in their analytic process. The authors are transparent about how themes are generated from the data.' Credibility was enhanced through triangulation, including interviewing the manager of the BCEHS CIS program, Marsha McCall, and Anonymous, a retired BCEHS paramedic, whom substantiated some participant data. Thick descriptions and a diverse sample contributed to credibility and transferability. An audit trail was maintained throughout the research process which enhanced dependability. Journaling and bracketing enriched reflexivity ' Each transcript was read at	Quotations are interspersed among the results which are presented as themes. Credibility was enhanced through the use of triangulation with leading members of the organisation.	The researchers consider their findings in relation to existing literature. They also make various recommendations for emergency service organisations, relating to post-incident organisational interventions that could protect paramedic wellbeing.

						all participants declined.'	least three times to improve accuracy and familiarity'. Sufficient data are presented to support the findings, and contradictory data are taken into account.		
<i>Mental Health in the UK Police Force: a Qualitative Investigation into the Stigma with Mental Illness</i> Edwards et al. 2020 ⁵⁸	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	The authors justify their use of open-ended questions in semi-structured interviews to facilitate meaningful discussion, although the choice of interviews other than qualitative data collection methods is not justified.	Five police officers. 'Participants were recruited through a referral from a charity, personal network, police forum and contact from within the police force.' The authors outline the criteria for participant inclusion in the study. They do not however justify their choice for these criteria. Non-participation was not discussed.	Semi-structured individual interviews were conducted using a topic guide, of which the authors provide details. The authors justify their use of open-ended questions. Interviews were audio recorded and transcribed. Data saturation was not discussed.	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	Ethical board approval was obtained, as well as written consent from the participants. The researchers also had measures in place to support participants who experience distress due to participation in the study.	The authors used Braun and Clark's approach of thematic analysis. They outlined this process in detail, giving examples of how they arrived at themes from the data. It is unclear whether multiple researchers collaborated to compare coding. Participant quotations are consistently embedded within the authors' interpretations, and contradictory data are taken into account. The authors are transparent about how themes were generated.	Findings are presented as themes and sub-themes. The authors include contradictory data, and discuss this. Discussion takes place throughout the study within the results section, but there is also a separate discussion section.	The authors identify areas where further research would be valuable, such as barriers to help-seeking in male dominated professions. Recommendations are also made to police organisations, such as highlighting a need to increase mental health awareness.
<i>Living in Critical Times: The Impact of Critical Incidents on Frontline Ambulance Personnel: A Qualitative Perspective</i>	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to	The authors don't justify why they chose individual interviews over other qualitative methods. They do however justify their use of some closed questions (21 EMTs, 6 EMCs. The qualitative component of this study followed a quantitative Survey. Participants from the earlier study	Setting for data collection is not stated. The authors give examples of the topics discussed during the interview process. The interview	Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	No mention of informed consent, ethical approval or methods to ensure confidentiality.	Thematic analysis was undertaken. 'A random sample of transcripts was read and coded by both authors in order to ensure good reliability and	Findings are presented as themes. For each theme, a range of relevant quotations are provided to support this theme. Credibility	The authors make recommendations to police organisations for post critical incident support for staff.

Gallagher et al. 2007 ⁵⁴	explore subjective experiences in detail.	to elicit background information and to facilitate comparisons across participants.)	were asked if they would be willing to participate in qualitative interviews. Non-participation is not discussed.	schedule used was created with the help of a literature review and the findings from the quantitative component of the study. No justification for the choice interview method is given. The interviews were audio recorded and transcribed.			validity.' It is not clear how the themes were derived from the data. The authors don't critically evaluate their own role in the analytic process. Contradictory data are taken into account- for example when giving examples of participant quotations with conflicting attitudes. The researchers don't critically examine their own role. Sufficient data are presented to support the findings.	of findings is not discussed.	
<i>Police officers, mental (ill-) health and spoiled identity</i> Bullock et al. 2018 ⁵⁹	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	Researchers justified their use of telephone interviews: '. While telephone interviews are often depicted as a less attractive alternative to face-to-face interviewing, telephones may allow respondents to feel relaxed, more able to disclose sensitive information and there is little evidence that they produce lower quality data '	52 police officers, two police staff, four Police Community Support Officers (PCSOs) (four) and one special constable. Participants were recruited from six police constabularies in England and Wales. 'Individual participants were identified by virtue of their contribution to an online survey on the nature of work-related injury in which	The researchers explain in detail how the telephone interviews were conducted. Open questions were asked by the participants. Saturation of data is not discussed. 'All interviews were digitally recorded, professionally transcribed and anonymized. '	The researchers acknowledge that the nature of the interview is sensitive, and that the nature of telephone interviews would make participants more comfortable disclosing such information.	' The nature of the interviewing was inevitably sensitive and mechanisms were put in place to mitigate that; for example, interviewees were provided with the contact details of sources of support.' There is no mention of ethical board approval, informed or methods to protect patient confidentiality.	The authors use Braun and Clark's thematic analysis approach, and give a very brief overview of the approach, but they do not give examples of how themes were derived from the data. The researchers didn't critically examine their own role. Sufficient data are presented to support the findings, and contradictory data are taken into account. Data	The findings are presented as themes. For each theme, the authors use participant quotations as well as relevant literature to discuss the context surrounding the quotations. No efforts to enhance credibility are discussed.	The authors make recommendations to police organisations based on their findings surrounding stigma. No recommendations for research are made.

			they identified themselves as willing to participate in a full-length interview'. This study is part of a wider project. Non-participation was not discussed.				saturation is not discussed.		
<i>'You see a baby die and you're not fine: 'a case study of stress and coping strategies in volunteer emergency medical technicians'</i> Folwell et al. 2018 ⁵⁵	Aims of study are clearly stated, and their importance justified. Qualitative methodology is appropriate due to nature of the aims, which is to explore subjective experiences in detail.	The authors justify their choice of in-depth interviewing technique as they 'hope to gain a deeper understanding of the lived experiences of voluntary EMTs'.	25 EMTs. Participants were recruited from one county in a Western state. It is not clear how the participants were selected, and non-participation is not discussed.	Interviews were conducted in a private space of the participant's fire department. Justification for choice of interview setting was not provided. In-depth interviews were performed with participants, using an interview guide. This was chosen to "gain a deeper understanding of the lived experiences of voluntary EMTs". Saturation was reached by the end of the study. Interviews were audio recorded and transcribed.	'The research team consisted of a female Caucasian professor who holds a doctorate and a male Caucasian undergraduate student with three years' experience as a volunteer EMT. While some of the interviews were conducted by both members of the research team, most interviews were conducted by the volunteer EMT'. Other than providing these details, Researchers do not critically examine their role, potential bias and influence of research question formulation or data collection.	Confidentiality was guaranteed by the authors as the transcripts were anonymised. There is no mention of ethical board approval or informed consent.	The authors describe the process of constant comparison analysis. Two researchers independently read all the transcripts, and collaborated to discuss findings. The authors are very transparent about the analytic process. Contradictory data were discussed, such as in the example of participants discussing the pros and cons of CISD. Sufficient data are presented to support the findings.	Findings are presented as themes and sub-themes. 'To enhance credibility, member validation was performed in which a summary of findings and initial interpretations of data were given to five participants to confirm the researchers accurately depicted viewpoints and experiences. To enrich transferability, we provided detailed descriptions of participants and research sites. To improve dependability, we used the same protocol for each interview and documented the process of data collection and analysis. To	The authors make four detailed recommendations to the specific EMS organisation with which the study was involved. The authors also make recommendations for future research.

								address confirmability, we used direct quotes from participants to support the finding'	
--	--	--	--	--	--	--	--	---	--

Appendix A- Abbreviations and definitions

Abbreviation	Meaning
NHS	National Health Service
PTSD	Post-Traumatic Stress Disorder
ESW	Emergency Service Worker
ESO	Emergency Service Organisation
EMS	Emergency Medical Service
CISD	Critical Incident Stress Debriefing
UK	United Kingdom
TRiM	Trauma Risk Management
GMB Union	General, Municipal, Boilermakers and Allied Trade Union
RTA	Reciprocal Translocation Analysis
CASP	Critical Appraisal Skills Programme
EMT	Emergency Medical Technician

Term	Definition
Psychological distress	'the unique discomforting, emotional state experienced by an individual in response to a specific stressor or demand that results in harm, either temporary or permanent, to the person' ⁴¹
Critical incident	'any event with sufficient impact to produce significant emotional reactions in people now or later'. ⁹
Supervisor	EMS frontline personnel responsible for overseeing a 'crew' of EMTs/paramedics. <i>Brown University- Roles And Responsibilities Of Supervisory Staff. [internet] 2021 [cited 2021 June 1]. Available from: https://www.brown.edu/campus-life/health/ems/roles-and-responsibilities-supervisory-staff#:~:text=EMS%20Supervisors&text=The%20EMS%20Supervisor%20is%20the,and%20appropriate%20emergency%20vehicle%20operation.</i>
Mental health champion	A member of an organisation raising mental health awareness within the workplace ⁹⁹

Type of stigma	Definition
Anticipated stigma (Perceived stigma)	'the extent to which people believe they personally will be viewed or treated in a stigmatizing way if their mental health problem or related help-seeking becomes known' ⁷⁷
Treatment stigma	'the stigma associated with seeking or receiving treatment for mental ill health' ⁷⁶
Internalized stigma	'holding stigmatizing views about oneself' ⁷⁶
Public stigma	'invalidating and unjustified beliefs (i.e., prejudices and endorsed stereotypes) about others' ⁸³

Appendix B: Literature search terms

((((mental AND (health OR well?being)) OR (trauma* OR *stress OR recover* OR PTSD OR post?traumatic?stress OR emotion* OR (critical AND incident) OR (traumatic AND incident) OR (traumatic AND event) OR experience* OR support* OR *support) AND (emergency service* OR emergency medical service* OR EMS OR first responder* OR ambulanc* OR paramedic* OR firefighter* OR fire service* fire department* OR police*) AND (qual* OR mixed?method* OR interview* OR focus?group*) NOT (quality OR emergency?department)

((emergency service* OR emergency medical service* OR EMS OR first responder* OR ambulanc* OR paramedic* OR firefighter* OR fire service* fire department* OR police*) AND (qual* OR mixed?method* OR interview* OR focus?group*) NOT (quality OR emergency?department)) AND ((help-seeking* OR stigma* OR mental* OR barriers OR (MeSH terms: help-seeking behaviour, social stigma, mental health, psychiatry, social support, mental disorders))

Appendix C: Data extraction template

Citation

Reviewer

Country

Aims

Ethics – how ethical issues were addressed

Study setting- e.g. type of organisation

Relevant context to study setting

Socio-demographics of the country / region

Recruitment context (e.g. where people were recruited from)

Sampling- what sampling methods, what were inclusion and exclusion criteria,

Data quality rating

Participants- ‘population described’

Participants- ‘characteristics’- age, sex etc

Theoretical background

Proportion of sample exposed to critical incidents/ definition of critical incident/ anything relevant to the study matching my selection criteria

Definition of critical incidents/ something related

Data collection methods- data collection methods, role of researcher within setting...

Data analysis approach- how many researchers, how did they code, how were findings corroborated,

Themes identified in study (1st order interpretations)- Help-seeking

Themes identified in study (1st order interpretations) – Mental Health recovery

Data extracts related to key themes- Help-seeking

Data extracts related to key themes- Mental health recovery

Author explanation/interpretation of key themes (2nd order interpretations)- Help-seeking

Author explanation/interpretation of key themes (2nd order interpretations)- Mental health recovery

Recommendations made by authors (both outcomes; but specify)

Third order interpretations:

Other potentially relevant information

Appendix D: Example literature quotations used to construct themes

Factors contributing to mental health recovery post traumatic incident exposure	Organisational	Time out/Downtime	<p><i>They prefer support to be offered immediately after the call and find that downtime after a stressful call allows them to decompress and prepare for the rest of their shift: ‘...we knew we weren’t going to get a call right, so we knew we had the two hours, so we watched a funny show and had a nap...But...like for me personally...that’s exactly...what I needed.’ Participant 4 from focus group 4. Douglas et al. 2013</i></p> <p><i>... our supervisor took us out of service for a couple of hours and let us go have lunch, sat down and had lunch and just kind of relax and talk amongst ourselves, not even about the call, just about whatever, just to kind of relax. Before we went back on shift. Certainly we could have booked off the rest of the day, you know, on stress leave or whatever, but we all, found that just having, just being able to have a couple of hours to, kind of, you know, relax a little bit, that helps us a lot.’ (Focus group #520–522)</i></p> <p>Halpern et al. 2008</p> <p><i>I like didn’t want to be like I need to take a couple of hours off because I did not want to look bad in front of the supervisor.-Participant 2 from focus group 2.</i></p> <p>Douglas et al. 2013</p> <p><i>Unfortunately, paramedics often do not have adequate time to talk. Several participants expressed their frustration over ‘calls waiting’. These calls are waiting to be dispatched as no crew is available. When BCEHS paramedics offload their patient(s) at the hospital, dispatch can immediately send them to a waiting call. This can limit paramedics’ ability to discuss CIs. James desired:</i></p> <p><i>...to be given the time uninterrupted, unbothered... Give us our time to decompress.</i></p> <p>Drewitz-Chesney et al. 2019</p>
		Supervisor	<p><i>Depending on who it is ... one supervisor, we took, we did a [critical call], and we took the last hour of the shift off. And he was making us fill out all these forms and, you know, telling us that if we wanted to take the next shift off that we had to go see our doctor and get a note for this and that. And you know, just made it more stress-(Focus group #520–522)</i></p> <p>Halpern et al. 2008</p> <p><i>‘And so our supervisor was really, we have a great supervisor and he’s, you know, asked us all specifically, “are you guys okay”, you know. And the other crew went off on stress for the rest of the shift. We stayed, because we said, you know, we’re okay. We just kind of dealt with the aftermath of everything. It was still a pretty stressful call but at least we had that option. And he had no problems, like, he said, go home. Whatever you guys need. So and that’s a big thing.- (Focus group #520– 522)</i></p> <p>Halpern et al. 2008</p>

			<p><i>Supervisors, however, were perceived differently. Some paramedics reported feeling unsupported when their supervisors questioned why they were reacting to an event that they did not perceive as traumatic:</i> <i>I wasn't really involved in a traumatic event because there was no loss of life, or you know what I mean, um, it, there wasn't really much sympathy the, um, I could, not that I really, um, oh, this wasn't, um, a huge accident, you weren't injured, you know, really why are you complaining?</i> Regehr et al. 2007</p> <p><i>he's [supervisor] quite an old fashioned sort of police officer, not the bloke you would sort of want to go in and have a chat with about a sudden death you'd just been to ... If I went in and said 'Governor, can I have a chat about the sudden death?', he'd look at me as if I'd just asked to kill one of his children!</i> Evans et al. 2013</p>
		Peer support network	<p><i>Nurse 1 "Therefore, talking to a person who has the same skills, abilities and knowledge you have would probably be more meaningful".</i> Carvello et al. 2019</p> <p><i>The majority of nurses is in favour of peer-supporters. The motivation is based on the fact that they recognise the peer-support as someone that can understand what nurses really mean when relating a traumatic event, being one of their colleagues.</i> Carvello et al. 2019</p>
	Informal support	Colleagues and family	<p><i>The more you talk about something, the more it becomes something you've told and your telling becomes part of the memory, as opposed to it being a really shiny, vivid thing inside your head*those images.</i> Evans et al. 2013</p> <p><i>I don't think talking about it to people at work is the release, the escape I need ... it's speaking to people who I care about and who care for me and just having that comfort zone, that's what's important to me. [P16]</i> Evans et al. 2013</p> <p><i>... I never tell my wife that, I would never tell her that because I just think that would have really put the frighteners on her. [P15]</i> Evans et al. 2013</p>
		Regular partner	<p><i>For me it's just always been that partner, because they're right there with you and they'll know what's going on, and you really want somebody that can understand what's happening.</i> Jessica et al. 2009</p> <p><i>...for those of us that have regular partners, regular stations, rely on your partner... They're gonna know if something's up with you.</i> Drewitz-Chesney et al. 2019</p> <p><i>Paramedics working with different partners may be unable to recognise changes as readily as they would in a regular partner. When paramedics are without a regular partner or are uncomfortable speaking with their</i></p>

			<p><i>current partner, they often don't discuss calls. Instead, some participants said they speak with long-time paramedic friends or coworkers with whom they have similarities, ie. age or training level.</i> Drewitz-Chesney et al. 2019</p> <p><i>...if I'm working with my regular partner, those communication lines are very, very open and we can talk about the tough calls, how we're feeling, if there's anything we would have done different [sic] (Sean).</i> Drewitz-Chesney et al. 2019</p> <p><i>I really envy the paramedics that have regular partners that they know, and they trust, and they can talk to (Krista).</i> Drewitz-Chesney et al. 2019</p> <p><i>...I would have a regular partner and that partner would be someone who is just like family to me. We would just talk about everything, without even hesitating...'</i> Drewitz-Chesney et al. 2019</p>
		Reassurance and validation	<p><i>I remember going to the hospital and I remember I was disappointed because the parents weren't there at that time...And I don't know what it was. It was just something that made me feel like I just needed to talk to them. But I never ended up talking to them. So anyway that call definitely stands out as being...(Stuck with you.) Yeah.</i> Halpern et al. 2009</p> <p><i>And it's weird, I looked in the newspaper. I saw his funeral announcement and I went to the funeral. (Oh okay.) I didn't talk to any of the family. I just went for my own, I don't know why ...I, for some reason, I guess I felt I needed to follow up and so I did. And I sent the family a condolence card and then a friend of mine ...met a friend of the family's friend and he had said, oh, his [relative] wants to talk to you ...So she called me and she wanted to know, like, you know, what happened with her [relative] and who was the last one to talk to him ...It's kind of funny because there are so many calls you do and you never think about them again. But this one, I thought about him a lot. I guess because he was so young and again, I really didn't think he was going to die. So I thought, okay, well, you know, he's at the trauma centre now and I had no idea that he was that badly injured. And then he died. So I think that freaked me out. I wasn't prepared for that. And I thought, oh my God, he's so young and his family wasn't there. I felt badly his family wasn't there and then. So that one stuck with me for a while.</i></p> <p><i>Interviewer: "And talking to his [relative], did that help or ...?"</i></p> <p><i>It more, I felt it was helping them so it made me feel better, because she really, really was upset. And I felt she wanted, I think she felt a lot better after talking to me and hearing something about what happened. And so it made me feel better to give them some kind of closure.Pg 181-182</i> Halpern et al. 2009</p>

			<p>... the day after that call, my supervisor kind of sensed that I wanted to talk to him and I don't know why, you know, he came to the hospital. But it was the next day and he was kind of like, so pull up a chair, sit down. Let's just talk. And it was like I wanted somebody just to say to me, okay, this is your time and you can talk right now and I'm not going to judge you. I'm not going to talk about anything. I'm going to let you just have your 10 minutes. And that made a huge difference. I have the utmost respect for my particular supervisor ... Whether it was talking specifically about the call or just having a little bit of a "hey you're okay at your job", type comment. (Participant #122)</p> <p>Halpern et al. 2008</p>
Factors influencing help-seeking behaviour	Nature of intervention delivery	Mandatory vs non-mandatory	<p>We had counselling every six months ... and everybody used to go 'Oh I've got to see the counsellor this week', but I tell you what ... we all quite enjoyed it ... I was so much calmer after speaking to her but it's something I'd never have done had I not been made to do it. [P12]</p> <p>Evans et al. 2013</p> <p>My emotions are none of your business and if I wanted to share my emotions with you, I'm going to share [them] with someone I trust...</p> <p>Participant 1 from focus group 1.</p> <p>Douglas et al. 2013</p>
		Shared experiences with intervention provider	<p>FRs in our study preferred a MH professional with experience as an FR or military veteran. Many also approved of a provider that "knew the job," either working with multiple FRs in the past, or even as a family member.</p> <p>Jones et al. 2020</p> <p>I've been to [therapy] a couple of times. . . . The guy that I got was excellent, but I only think it was excellent because he was prior military (P8, Firefighter ×22 years).</p> <p>Jones et al. 2020 (study 4)</p> <p>...someone I think understands what's going on, and has been through what I've done.- Participant 3 from focus group 4.</p> <p>Douglas et al. 2013</p>

	Stigma as a help-seeking barrier Specific barriers	‘Macho culture’	<p><i>I think there’s a real element of machismo and masculinity in the police force and it’s a bit, sort of a faux pas to admit that things have really affected you ... If I’d have come out and said ‘ah you know, that really affected me badly, let’s go and sit down and have a cup of tea and talk about it’ I think you’re straying into pink and fluffy territory there ... saying ‘that made me feel sad’ is a bit too far. [P3]</i> Evans et al. 2013</p> <p><i>Everyone wants to be tough and strong. Maybe that was my downfall or problem at the time and I didn’t want to admit that I needed any kind of help. I guess I didn’t want to be perceived as weak. (Participant #110)</i> Halpern et al. 2008</p> <p><i>... that’s so ridiculous. You need to harden up if you want to do this job, and old people die.</i> Drewitz-Chesney et al. 2019</p> <p><i>... in metro, there’s a lot more...bravado and joking around about things...If you can get one on one with someone, they’re usually a lot more receptive and a lot more empathetic- (Dennis).</i> Drewitz-Chesney et al. 2019</p> <p><i>Unfortunately there’s a massive stigma [...] one of my sergeants the other week, there was mental health training coming up and his reaction to being put forward to go on the training was, ‘Well what do I want to go and learn how to deal with a load of nutcases for?’</i> Bullock et al. 2018</p> <p><i>We are many women at the station so its easier for us to talk about what happened...You have to talk about what happened otherwise you can’t go on. It’s easier than on a station with only men, it’s not necessary with any “macho style” so I think it’s easier for the men (at our station) to talk about things you must talk about. (A female nurse describes her feelings.)</i> Jonsson et al. 2003</p>
		Stigma and shame	<p>As an institution, they were very very good, in getting me better, at making me understand, but back at work that was a different ball game. You’ve got your colleagues who are still at work, still running around like idiots and they’re like, oh fucking hell, you’ve had three months off, you know, I should have gone off with stress. (Participant 1). Edwards et al. 2020</p> <p><i>There have been officers that are doing the shift that have shown that they can’t deal with situations like that, and been very open about it*and they haven’t got the respect from the shift, because the colleagues go ‘well, you’re on your own if you’re working with her, because she’d back away’ or whatever. So you don’t want to be considered as one of those. [P7]</i> Evans et al. 2013</p>
		Career concerns	<p><i>I think that the stigma is you have to be very careful who you tell that it bothered you or you might get judged as weak or you might get fired” (P2, EMT/paramedic ×20 years).</i> Jones et al. 2020</p> <p>If you’ve a form of mental health illness you will not get on; you will not be promoted, ...people will not want you on their section. (Participant 4). Edwards et al. 2020</p>

			<p>I feel scared to declare anything or do anything about anything because will it bite me later on in life? Will it prevent me from doing something in the police later on? Could it be used against me? ... will it be used in a negative way later on?</p> <p>Bullock et al. 2018</p>
		Confidentiality concerns	<p><i>I know our department's very, very poor at keeping secrets. So if I put a crew out of service, I have to tell the communications center.</i></p> <p><i>Communications says, we're sitting this far apart, Hey X, I just put the 22 car out of service in stress. Every dispatcher in there hears it. Every call receiver in there hears it. So they hear it. I don't know if they say anything. I don't know if they go home and tell all their friends and family. But I don't like that system. There's no quiet way of doing it. (Participant #128)</i></p> <p>Halpern et al. 2008</p> <p><i>I think that the stigma is you have to be very careful who you tell that it bothered you or you might get judged as weak or you might get fired" (P2, EMT/paramedic ×20 years).</i></p> <p>Jones et al. 2020</p> <p><i>.. to whom must I speak in the police if I can't trust anyone. So now I'm seeking professional help outside the police .</i></p> <p>Boshoff et al. 2015</p>
	Mental health literacy	Emotional awareness	<p><i>Some appreciated that difficulties in recognizing and admitting to distress pose significant barriers to accessing support. Recognizing the emotional impact of critical incidents may help to address these barriers.</i></p> <p>Halpern et al. 2009</p> <p><i>I didn't recognise it as what it was; I just thought I was grumpy...you don't see them creeping up, and in the end, the thing that tips you over the edge, the thing that makes your bottle overflow if you like can be something quite small because you've got used to dealing with stuff. (Participant 5).</i></p> <p>Edwards et al. 2020</p>
		Education and stigma	<p><i>I think for, for everybody, is, here's my thought towards your process, is giving them the tools. People are going to be very tough and say, yeah, yeah, fine. But you know if you could somehow identify the emotions that go along with these calls that might be starting to put you on tilt...Then you can teach people to be aware of them and say, hey, you know what, it's okay to say, I need to talk to someone.</i></p> <p>Halpern et al. 2009</p> <p><i>In particular, it would appear from this study that teaching ambulance personnel about the emotional aspects surrounding different types of critical incident may diminish their confusion about which incidents they can expect to impact them.</i></p> <p>Halpern et al. 2009</p> <p><i>I guess that's the main two [barriers], pride and then denial. . . . But, it's just education. Just letting [FRs] know, look, these things are normal. It's going to happen to somebody" (P11, Firefighter ×8.5 years).</i></p> <p>Jones et al. 2020</p>

			<p><i>Knowing that other people are there dealing with that same stuff. You can bounce ideas off each other, see what's worked in their situations and what hasn't . . . you realize you're not crazy (P12, EMT/paramedic ×14 years).</i> Jones et al. 2020</p> <p><i>If you could actually get people in in front of officers saying, 'I was one of those people that didn't believe stress could ever get to that level and it was ridiculous and you just needed to work harder', maybe officers would accept that from another officer more than just somebody standing in front of you training because it's getting officers to accept that actually it's okay, you're only human and your body and your mind can only take so much, and maybe they'd accept it more then.</i> Bullock et al. 2018</p>
--	--	--	--

