BMJ Open Application of primary healthcare principles in national community health worker programmes in low-income and middle-income countries: a scoping review

Shagufta Perveen , ¹ Zohra S Lassi , ² Mohammad Afzal Mahmood , ¹ Henry B Perry , ³ Caroline Laurence , ¹

To cite: Perveen S. Lassi ZS. Mahmood MA. et al. Application of primary healthcare principles in national community health worker programmes in low-income and middle-income countries: a scoping review. BMJ Open 2022;12:e051940. doi:10.1136/ bmjopen-2021-051940

Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (http://dx.doi.org/10.1136/ bmjopen-2021-051940).

Received 06 April 2021 Accepted 05 January 2022



@ Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by

For numbered affiliations see end of article.

Correspondence to

Shagufta Perveen; shagufta.perveen@adelaide. edu.au

ABSTRACT

Objective To identify which primary healthcare (PHC) principles are reflected in the implementation of national community health worker (CHW) programmes and how they may contribute to the outcomes of these programmes in the context of low-income and middle-income countries (LMICs).

Design Scoping review.

Data sources A systematic search was conducted through PubMed, CINAHL, EMBASE and Scopus databases. Eligibility criteria The review considered published primary studies on national programmes, projects or initiatives using the services of CHWs in LMICs focused on maternal and child health. We included only English language studies. Excluded were programmes operated by non-government organisations, study protocols, reviews, commentaries, opinion papers, editorials and conference

Data extraction and synthesis We reviewed the application of four PHC principles (universal health coverage, community participation, intersectoral coordination and appropriateness) in the CHW programme's objectives, implementation and stated outcomes. Data extraction was undertaken systematically in an excel spreadsheet while the findings were synthesised in a narrative manner. The quality appraisal of the selected studies was not performed in this scoping review.

Results From 1280 papers published between 1983 and 2019, 26 met the inclusion criteria. These 26 papers included 14 CHW programmes from 13 LMICs. Universal health coverage and community participation were the two commonly reported PHC principles, while intersectoral coordination was generally missing. Similarly, the cultural acceptability aspect of the principle of appropriateness was present in all programmes as these programmes select CHWs from within the communities. Other aspects, particularly effectiveness, were not evident.

Conclusion The implementation of PHC principles across national CHW programmes in LMICs is patchy. For comprehensiveness and improved health outcomes, programmes need to incorporate all attributes of PHC principles. Future research may focus on how to

Strengths and limitations of this study

- Community health worker programmes in developing and lower-middle-income countries are an essential aspect of the strategy to achieve health for all and sustainable development goals, and this scoping review can be considered as an important step towards reviewing national community health worker programmes in low-income and middle-income countries applying the lens of primary healthcare principles.
- Four bibliographic databases were searched using a basic search strategy that was modified as per the database requirement.
- The studies were heterogeneous in their methods and outcomes assessed and that posed a challenge in comparing primary healthcare principles.
- The generalisability of the results of this study is limited to larger national-level programmes in developing and lower-income and middle-income countries only.

incorporate more attributes of PHC principles while implementing national CHW programmes in LMICs. Better documentation and publications of CHW programme implementation are also needed.

BACKGROUND

Primary healthcare (PHC), as an approach to a reorientation of health services and provision of universal healthcare, has remained the benchmark for most countries' discourse on health since the PHC approach was mobilised by the Alma Ata Health for All (HFA) declaration for comprehensive, evidencebased responses to local health needs with reference to the social context. PHC is a whole-of-society approach to health and aims to attain the highest possible level and distribution of health and well-being by providing



an accessible and wide range of services, including health promotion; disease prevention, treatment and rehabilitation; and palliative care.¹

'HFA' requires that health systems respond to the challenges of a changing world and growing expectations for better performance. PHC includes the key elements needed to improve health security, through a focus on community engagement, preventative collective action, access to good quality medicines, rational prescribing and a core set of essential public health functions, including surveillance and early response. A PHC approach achieves this by strengthening community-based initiatives and building resilience.

Across a wide variety of settings in low-income, middle-income and high-income countries, PHC-oriented health systems have consistently produced better health outcomes, enhanced equity and improved efficiency. In Brazil, for example, enrolment in the family health strategy has been linked to a higher likelihood of regular care, better access to medication and improved patient satisfaction. Hence, PHC has been rightly advocated as the key to achieving HFA and the 2018 Astana Declaration reiterated the importance of this approach for achieving universal health coverage (UHC). ²³

PHC, as an approach to achieve HFA goals,' was built on the principles of equity in access to health services and the right of people to participate in decisions about their own healthcare. These principles that is, 'equity' and 'community empowerment' underpin preventive and promotive health services, appropriate technology and intersectoral collaboration.⁴ Evidence suggests that if countries have explicitly organised their health systems around PHC principles, it has led to improved health outcomes. For example, in Portugal, by 2008, the life expectancy at birth increased 9.2 years more than it was 30 years ago. In Congo, the case-fatality rate after caesarean section dropped from 7% to less than 3% from 1985 to 2000. In, Iran, the under-five child mortality reduced from 80 per 1000 to less than 20 per 1000 in rural areas from 1980 to 2000.⁵

PHC's emphasis on community-based services is an important way to ensure access, in rural, remote areas and for disadvantaged populations. With limited resources and geographical and epidemiological context, it is a challenge for healthcare systems in lowincome and middle-income countries (LMICs) to reach out to the whole population. Therefore, as part of the PHC approach and with a view to its principle of community empowerment, community health worker (CHW) programmes were envisioned as a way to reach a wider population for essential health needs and to achieve HFA. National CHW programmes were implemented by many governments from 1978, operating at the interface between communities and the primary care level of the health system. 6-10 Established under the PHC principles, these programmes were expected to encompass and promote them and in doing so achieve improvements in health outcomes.¹¹

National CHW programmes, as vehicles to incorporate PHC principles into healthcare provision, have contributed significantly in reducing under-5 child mortality in Brazil, 12 Indonesia 12 and Nepal. 13 In Indonesia, immunisation coverage also improved many-fold with an increase in CHWs. These examples demonstrate a clear link and need for incorporating PHC principles when implementing CHW programmes. Over decades of implementation CHW programmes have also faced various challenges including the loss of the PHC movement. 14 15 Though, the PHC principles are evident in the programme design and policies of the CHW programmes in various countries. 16-20 There is not widespread/ comprehensive evidence of the extent to which PHC principles are systematically applied across the national CHW programmes. This study aims to identify the PHC principles in the implementation of these programmes in LMICs and to understand their contribution to the outcomes of those programmes.

METHODS

A systematic scoping review was conducted using a predefined protocol²¹ and reported as per the Preferred Reporting Items for Systematic Reviews and Meta-analyses extension for Scoping Reviews (PRISMA-ScR) guidelines.²² The databases searched in September 2019 were PubMed (MEDLINE), CINAHL (EBSCOhost), EMBASE (Elsevier) and Scopus (Elsevier). The review only considered published primary studies on programmes, projects or initiatives utilising the services of CHWs in LMICs. We focused on the national level CHW programmes defined as any CHW programme that is operated or implemented by the government of a specific country, on multiple sites (jurisdictions/provinces/regions) within a country and has been functional for a minimum of 3 years. We considered national CHW programmes with a maternal and child health (MCH) focus as it is a national priority in the majority of LMICs.

Papers published only in the English language from October 1978 to September 2019 were considered as 1978 was the year of the Alma-Ata declaration that promoted the establishment of national-level CHW programmes under the PHC principles. Excluded were study protocols, narrative reviews, commentaries, text and opinion papers, viewpoints, editorials, conference proceedings/abstracts, correspondences, systematic and scoping reviews and the papers on the CHW programmes operated by a non-government organisations. Papers were also excluded if they involved health professionals other than CHWs such as midwives, nurses and traditional birth attendants. Papers were not excluded based on the unavailability of the abstract.

The search strategy, including all identified keywords and index terms, was adapted for each included database (online supplemental appendix 1—logic grid). The search terms used included 'community health worker', 'Program', 'Maternal and Child Health' and 'Low-and



Middle-Income Countries'. The results of the search are presented in the PRISMA-ScR flow diagram in the results section.

Following the search, all identified records were collated and uploaded into Covidence software²³ and duplicates removed. Two authors (SP and ZL) independently screened titles and abstracts and then matched the full texts selected during screening against the inclusion criteria. The reference lists of relevant papers were also searched for additional studies. Papers meeting the inclusion criteria were included in the review for data charting. In scoping reviews, the data extraction process is referred to as charting the results.24 SP and ZL completed data charting using a pre-developed data charting form. Key attributes of the data charting form included the country of origin, study objective, design and key findings, name of the CHW programme, objective and reflection of PHC principle/s in programme objective, implementation activities, and stated outcomes along with the selection process of CHWs (online supplemental appendix 2). The data charting form was pilot tested and modified accordingly. The operational definition of the PHC principles used as reference in this scoping review are as follows:

- 1. UHC: all people receive the health services they need, including public health services designed to promote better health, prevent illness and to provide treatment, rehabilitation and palliative care of sufficient quality to be effective, while at the same time ensuring that the use of these services does not expose the user to financial hardship.^{2 25}
- 2. Community participation: Active community involvement in defining health problems and needs, developing solutions and implementing and evaluating programmes.²
- 3. Intersectoral coordination: The linkage between health and development.²
- 4. Appropriateness: Services should be effective, culturally acceptable affordable and manageable.²

We examined the included studies in light of all or any of the subattribute of the above listed four PHC principles and reported accordingly. The evidence is reported if it was mentioned explicitly in the article or inferred by the researchers reflecting the implementation of PHC principles even if the evidence was about only one aspect of a principle. The relevant evidence is extracted and reported in the results section.

There was no quality assessment conducted of the included studies. The findings were synthesised in a tabular and narrative manner. The conceptual framework, including definitions of the four principles, for collating and summarising the data is presented in the published protocol. ²¹

Patient and public involvement

We did not involve patients or the public in this scoping review.

RESULTS

Search results

We identified 1280 citations through database searches. After removing duplicates and screening out non-relevant abstracts, we assessed 281 full-text papers for eligibility. 263 of those 281 were excluded as these did not meet the eligibility criteria. In total, 18 papers, ^{17–20} ^{26–39} published from 1983 to 2019 met the eligibility criteria (figure 1). Eight ^{40–47} papers were further included from the reference lists of the included studies, making a total of 26 papers.

Of the 26 papers, two studies were conducted in western Asia, ^{17 35} 12 studies were conducted in South Asia ^{18 27 29 31 33 37 38 40-44} and 1 study in South East Asia. ²⁸ Seven studies were conducted in Africa ranging from the Horn of Africa, ^{19 30 45 46} Central Africa, ²⁰ Western Africa and South Africa. ³⁹ Two studies were conducted in South America, ^{34 47} one in Central America and one study was conducted in the Caribbean. ²⁶ Altogether, these 26 studies covered 14 CHW programmes from 13 LMICs.

Fourteen of the 26 included studies were quantitative $^{19\ 26\ 28\ 31\ 32\ 34-36\ 40\ 42\ 43\ 45-47}$ and 12 studies were qualitative. $^{17\ 18\ 20\ 27\ 29\ 30\ 33\ 37-39\ 41\ 44}$ Online supplemental table 1 provides an overview of the included studies outlining the key objective/s, methods and findings as reported by the authors.

Application of PHC principles

The PHC principles were applied to a varied extent in the objective/s, implementation and outcome of the national CHW programmes reviewed in this study (table 1). The evidence found in the objective, implementation or the outcome of the included studies related to the application of the four PHC principles is organised in online supplemental table 2.

'Universal health coverage' and 'community participation' were the two commonly reflected PHC principles in the national CHW programmes across their objective/s, implementation and outcomes. 'Intersectoral coordination' was only mentioned in the outcome of Iran's Women Health Volunteers programme. ¹⁷ The objective of two CHW programmes not reported in the papers reviewed. ²⁸ ²⁹ In addition, studies from Nepal, ¹⁸ ⁴⁴ Bangladesh ²⁹ and Niger ³² did not report on the outcomes of the CHW programmes.

Universal health coverage

We reviewed the national CHW programmes for the application of this fundamental PHC principle in terms of coverage and access, equity and comprehensiveness. UHC was reflected in the objective of 11 CHW programmes ^{18–20 26 27 32 34–37 39} and in the implementation of 14^{17–20 26–29 32 34–37 39} programmes through the service provision by CHWs in the MCH and family planning domain. These 14 programmes reported improvements in the scope (population coverage) and range (comprehensiveness) of health services provided. For example, an outcome of the CHW programme in Iran




Figure 1 PRISMA flow chart for study selection and inclusion process. CHW, community health worker; IMNCI, integrated management of newborn and childhood illness; MNCH, maternal newborn and child health; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-analyses.

was increased utilisation of MCH care services as a result of the active follow-up by CHWs. The increase in immunisation coverage of children in the rural areas was also attributed to the 'active' approach and vigilance of CHWs and vaccinators serving the PHC network of Iran. In Pakistan, the CHW programme was claimed to be contributing to the increasing utilisation of antenatal care and family planning services. In Rwanda, mHealth was reported as improving communication between CHWs and community members leading to better use of the health services.

The concept of 'care according to need' was reflected in the objective of Pakistan's CHW programme that focuses on the provision of care in underserved areas.²⁷ Service provision to ethnic minorities was one of the focus areas of Nepal's CHW programme.¹⁸

Community participation

Only three^{17–19} of the 14 CHW programmes included in this review incorporated community participation in their programme objective. In terms of implementation, 10 programmes^{17 18 20 27–31 35 36} reflected community participation as they engaged CHWs from within the local communities to provide care to the local population. Moreover, the selection of CHWs from the local community they serve facilitated their access to households, development of good relationships and high acceptability in the community.^{27 30 32} Three programmes^{32 34 39} did not mention the selection process of CHWs while in Jamaica it was not mandatory to select CHWs from within the local community.²⁶

Examples of other activities reflecting the process of community participation² beyond the selection of



Table 1	Application of primary healthcare	principles as reflected in the	National community health worker programmes
Serial no.	Country/CHWP/year commenced	PHC principle/s observed in the CHWP Objective	PHC principle/s observed in the implementation of the CHWP PHC principle/s observed in the stated outcome/ achievement of the CHWP
1.	Iran/Women Health Volunteers Programme/1992 ¹⁷	Community participation	 ▶ Universal health coverage ▶ Community participation* ▶ Community participation ▶ Intersectoral coordination
2.	Iran/Primary Healthcare Network — Expanded Programme on Immunisation/1983 ³⁵	Universal health coverage	 ▶ Universal health coverage ▶ Community participation* ▶ Appropriateness
3.	Pakistan /National Programme for Family Planning and Primary Healthcare/1994 ^{27 33}	Universal health coverage	 ▶ Universal health coverage ▶ Community participation* ▶ Community participation
4.	India/Accredited Social Health Activist Programme/2005 ^{31 37 38}	Universal health coverageAppropriateness	 Universal health coverage Community participation
5.	Bangladesh/ National MCH and Family Planning Programme/1976 ²⁹	Not reported	 Universal health coverage Not reported Community participation*
6.	Nepal/ Female Community Health Volunteer Programme/1988 ¹⁸	Universal health coverageCommunity participation	 Universal health coverage Not reported Community participation*
7.	Cambodia/Village Malaria Worker Project as part of National Malaria Control Programme/2001 ²⁸	Not reported	 ▶ Universal health coverage ▶ Community participation*
8.	Ethiopia/ Health Extension Programme/2003 ^{19 30}	Universal health coverageCommunity participation	 ▶ Universal health coverage ▶ Community participation ▶ Community participation ▶ Appropriateness
9.	Rwanda/RapidSMS programme/2013 ²⁰	Universal health coverageAppropriateness	 Universal health coverage Community participation* Appropriateness Appropriateness
10.	Niger/ Rural Health Improvement Programme/1970s ³²	► Universal health coverage	▶ Universal health coverage Not reported
11.	South Africa/ward-based outreach teams-national CHW programme/2011 ³⁹	► Universal health coverage	 ▶ Universal health coverage ▶ Appropriateness ▶ Community participation
12.	Brazil/ Family Health Programme (Programa de Saude da Familia, PSF)/1994 ³⁴	► Universal health coverage	 Universal health coverage Community participation
13.	El Savador/ Rural Health Aide Programme/1976 ³⁶	► Universal health coverage	 Universal health coverage Community participation*
14.	Jamaica/ Community Health Aide programme/1978 ²⁶	► Universal health coverage	 ▶ Universal health coverage ▶ Community participation

^{*}Community participation consisted of only selection of community health workers from the local community in these programmes. CHWP, Community Health Worker Programme; MCH, maternal and child health; PHC, primary healthcare.

CHWs were reported only in Ethiopia's Health Extension Programme. ³⁰ In this programme the performance of health centres was evaluated by the community quarterly and the CHWs were monitored by the community volunteers. ³⁰

Intersectoral coordination

PHC ought to involve the health sector and all related sectors and aspects of national and community development that have an impact on health.^{2 48} Intersectoral coordination was not reflected in the objective/s or implementation of any CHW programme and only in the outcome of one¹⁷ programme. The WHV Programme of Iran explicitly described the intersectoral link

between health and education sectors for transmitting health messages to the people. ¹⁷ The Accredited Social Health Activist (ASHA) programme from India, while not reporting intersectoral collaboration directly, did report actions to enhance the role of women by creating opportunities by working with other sectors to empower women. ³⁸

Appropriateness

The final PHC principle assessed in this review was appropriateness, that is, services that are effective, culturally acceptable and financially affordable. The included studies reflected one or another of these attributes but none reported all three attributes of appropriateness. For

example, the concept of appropriateness was reflected explicitly in the objective of India's ASHA programme (to provide affordable and quality healthcare) but did not mention cultural appropriateness. The RapidSMS programme of Rwanda reported the cultural acceptability of technology (phone messaging services) and its affordability considering that almost all populations had access to a mobile phone. ²⁰

DISCUSSION

This study has provided insights into the application of PHC principles in the implementation of national CHW programmes. PHC principles do not appear to be applied with the rigour and regularity as one would expect considering the emphasis laid on these during conceptualisation of this significant public health movement called 'PHC'.

Our results show that 'UHC' and 'community participation' were the most common PHC principles reflected in the national CHW programmes. In contrast, intersectoral coordination was stated in the outcome of only 1 of the 14 CHW programs¹⁷ while none of the studies described the programmes with reference to all three attributes of appropriateness (effective, culturally acceptable and financially affordable).

'Enhanced coverage' attribute of UHC was most commonly reflected by the national CHW programmes. There is limited evidence in the reviewed 26 papers on the implementation of other two attributes, that is, coverage on the basis of need (equity) and comprehensiveness. This finding complements the fact that soon after Alma-Ata, selective PHC was proposed as an interim strategy for disease control in LMICs. 49 50 Many vertical programmes utilised CHWs under different names and with different roles⁵¹ resulting in a fragmented and disease-specific approach operating within the context of fragile health systems of LMICs. CHWs however, are not a 'panacea for weak health systems.' They require wellstructured support from the formal health systems with which national CHW programmes are linked. Therefore, achieving UHC requires strengthening of health systems with effective integration of comprehensive CHW programmes in LMICs as PHC can only work when a country has the structures, skills and data to ensure that all people are covered. 15

This review found that the implementation of community participation was patchy, and when it was employed it mainly reflected in the selection of CHWs from the local community. This is not surprising as after the Alma-Ata declaration several governments started CHW programmes as a means for people's participation with local lay people trained to administer basic first-line healthcare in their communities. While CHWs' position as community members themselves may provide a 'natural link' between them and the community, it may also appear to safeguard trust in 30 32 and respect for them from the community side and enhanced self-esteem from the CHW side.

A higher level of community participation where the community is given a stake in the evaluation and redefining of services was evident only in the Ethiopian CHW programme.³⁰ A successful CHW programme requires the support and ownership of the community through their active involvement in the entire process of defining health problems and needs, developing solutions, implementing and evaluating the programme, as well as establishing a supportive social and policy environment for community participation at national, district and local levels.⁵² CHW programmes often struggle to be successful when not part of a broader community engagement process which requires explicit methods for involving individuals and communities, clearly defined roles and responsibilities, training of policymakers and adequate funding. 52 Recent WHO guidelines have explicitly recommended ways to select CHWs, engage and mobilise the community and this can be achieved if there is a supportive social and policy environment.⁵³ With little or no evidence as noted by this scoping review on community involvement in needs assessment, the design of programmes and evaluation may indicate that invoking community participation is a challenge for these programmes. 15 Community participation is a context-dependent, gradual process that is less controllable and less measurable, thereby making it harder to track.⁵⁴ There is a need for robust programme evaluations of community participation activities that measure long-term outcomes and provide support for the CHW programmes to broaden their scope of community participation. Moreover, CHW programmes need to give attention to the experiences of CHWs themselves to address the feelings of powerlessness, and frustrations expressed by CHWs about how organisational processual and relational arrangements hindered them from achieving the desired impact. CHW programmes should systematically identify disempowering organisational arrangements and take steps to remedy these.⁵⁵

The operational problems related to partnerships working (intersectoral, interinstitutional, interdisciplinary and professional/lay partnerships) were highlighted in the early implementation years of CHW programmes in LMICs.⁵⁶ Our review informs that this is still the case.¹⁷ This finding corresponds with the fact that working relationships between partners have often proved difficult, 54 56 as each sector has its priorities.⁵⁴ Though some of the CHW programmes reflect that the CHWs do understand how various actors relate to each other, and where their interests lie and how they 'use this understanding in particular situations to provide an interpretation of the situation and frame courses of action that appeal to existing interests and identities,' inducing cooperation among a range of phenomena.⁵⁷

The PHC literature reports that community participation and intersectoral coordination are the two



most weakly implemented principles. ^{15 54} Our review findings also support this evidence. National CHW programmes ought to view these principles as two pillars that help achieve the UHC of services that are appropriate for the community and their context.

By its nature, the provision of MCH services to women by female CHWs who are also selected from within the local community tends to make it culturally acceptable and meet the principle of appropriateness. However, CHW programmes need to incorporate 'appropriateness' more explicitly in their objectives and then diligently pursue this in programme implementation and outcomes, which may contribute to addressing the current lack of evidence on the effectiveness of these programmes. ⁵⁸

Based on the findings of this scoping review, it can also be inferred that if the CHW programmes follow PHC principles they can be better positioned to help in current pandemic response and prevent future infectious outbreaks/epidemics by increasing access to health products and services, distributing health information, increasing social mobilisation, completing surveillance activities and reducing the burden of formal healthcare system.⁵⁹

The review has a number of limitations. First, it relied solely on the information reported in the papers to assess the application of PHC principles within the programmes. Many papers did not clearly articulate these principles or provide sufficient descriptions of the programme to allow an assessment to be made. As such the authors needed to interpret the evidence about principles in how the programme was implemented. These principles may be delineated elsewhere, for example, programme reports or funding agreements. Therefore, it is likely that we underestimated the application of PHC principles in these programmes. However, the very fact that the research papers that we reviewed failed to document the implementation of those principles, illustrates less than the adequate emphasis on the application of these principles in national CHW programmes.

Second, we reviewed the CHW programmes identified only through the search of peer-reviewed published journal articles and there may be CHW programmes that apply the PHC principles but are not published in peer-reviewed journals in a way to be captured in our search. This scoping review can be considered as a first step towards reviewing national CHW programmes in LMICs applying the lens of PHC principles. Future studies on the analysis of non-peer-reviewed publications or 'grey' literature may produce further evidence on this phenomenon.

CONCLUSION

This scoping review informs that the application of PHC principles across national CHW programmes in LMICs is patchy. For comprehensiveness and improved health outcomes, programmes need to incorporate all attributes of PHC principles. The findings also point to the limited research and published studies on this important topic. Better documentation and publications of programme implementation with reference to PHC principles are needed. Further research is needed to identify reasons for this inadequate emphasis on historic PHC principles, and to find out what other principles are adhered to by the current CHW programmes. Future research may also focus on how to incorporate more attributes of the PHC principles while implementing national CHW programmes in LMICs.

Author affiliations

¹School of Public Health, Faculty of Health and Medical Sciences, The University of Adelaide, Adelaide, South Australia, Australia

²Robinson Research Institute, Adelaide Medical School, Faculty of Health and Medical Sciences, The University of Adelaide, Adelaide, South Australia, Australia ³Department of International Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland, USA

Acknowledgements The authors would like to acknowledge the support of University of Adelaide's librarian Vikki Lington during the process of search strategy development for all included databases.

Contributors SP had the primary responsibility for writing the manuscript and making revisions. SP contributed to the design of the review, designed and conducted the search, adjudicated and appraised studies, charted and analysed data and drafted the manuscript. ZSL was involved in the screening and data charting of the articles and review of the manuscript. CL and MAM were involved in the conceptualisation and design of the scoping review, provided continuous supervision and feedback during the conduct of the scoping review and reviewed all the drafts and provided instrumental feedback to improve subsequent versions by SP. HBP also reviewed the drafts critically and provided feedback. All authors approved the final version. SP is responsible for the overall content of the article.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval The information used in our scoping review was derived from publicly available data sources therefore, a formal ethics approval was not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Shagufta Perveen http://orcid.org/0000-0001-8804-3635 Zohra S Lassi http://orcid.org/0000-0002-5350-6334 Mohammad Afzal Mahmood http://orcid.org/0000-0002-9395-8693



Henry B Perry http://orcid.org/0000-0003-0561-0492 Caroline Laurence http://orcid.org/0000-0002-8506-5238

REFERENCES

- 1 World Health Organization and the United Nations Children's Fund (UNICEF). A vision for primary health care in the 21st century: towards universal health coverage and the sustainable development goals. Geneva, 2018.
- 2 Bryant JH. Ten years after Alma-Ata. World Health, 1988.
- 3 UNICEF WHO. Declaration of Astana global conference on primary health care, Astana, Kazakhstan 2018 October 25-26. Available: https://www.who.int/primary-health/conference-phc [Accessed 02 Feb 2020].
- 4 Rifkin SB. Alma Ata after 40 years: primary health care and health for All-from consensus to complexity. BMJ Glob Health 2018;3:e001188.
- 5 World Health Organization. *The world health report 2008: primary health care now more than ever.* Geneva, 2008.
- 6 World Health Organization. Strengthening the performance of community health workers in primary health care. Report of a WHO Study Group [meeting held in Geneva from Dec 2-9, 1987]. Tech Rep Ser. Geneva, 1989: 1–46.
- 7 Rifkin SB. Paradigms lost: toward a new understanding of community participation in health programmes. *Acta Trop* 1996;61:79–92.
- 8 Lewin S, Babigumira S, Bosch-Capblanch X. Lay health workers in primary and community health care: a systematic review of trials. Geneva: World Health Organization, 2006.
- 9 Rifkin SB. Community health workers. In: Heggenhougen K, Quah S, eds. *International encyclopedia of public health*. San Diego Academic Press, 2008: 773–82.
- 10 Christopher JB, Le May A, Lewin S, et al. Thirty years after Alma-Ata: a systematic review of the impact of community health workers delivering curative interventions against malaria, pneumonia and diarrhoea on child mortality and morbidity in sub-Saharan Africa. Hum Resour Health 2011:9:27.
- 11 Kahssay HM, Taylor ME, Berman P. Community health workers: the way forward. United Kingdom: World Health Organization, 1998.
- 12 Lehmann U, Sanders D. Community health workers: what do we know about them. The state of the evidence on programmes, activities, costs and impact on health outcomes of using community health workers. Geneva: World Health Organization, 2007: 1–42.
- 13 Perry H. Health for the people: national community health worker programs from Afghanistan to Zimbabwe: maternal and child survival program, 2020.
- 14 Perry H. A brief history of community health worker programs. Encyclopedia of public health Berlim. Springer, 2013: 1–12.
- 15 Rifkin SB. Health for All and Primary Health Care, 1978 2018: a historical perspective on policies and programs over 40 years. Oxford Research Encyclopedia of Global Public Health USA: Oxford University Press, 2019.
- 16 Health Department Government of Sindh. Lady health worker program for family planning and primary health care Pakistan. Available: https://www.sindhhealth.gov.pk/LHW-Program [Accessed 13 Sep 2021].
- 17 Damari B, Riazi-Isfahani S. Evaluating the women health volunteers program in Iran- a quarter century experience (1992-2016). Arch Iran Med 2018;21:566-71.
- 18 Panday S, Bissell P, Teijlingen Evan, et al. Perceived barriers to accessing female community health volunteers' (FCHV) services among ethnic minority women in Nepal: a qualitative study. PLoS One 2019;14:e0217070.
- 19 Negussie A, Girma G. Is the role of health extension workers in the delivery of maternal and child health care services a significant attribute? The case of Dale district, southern Ethiopia. BMC Health Serv Res 2017;17:641.
- 20 Musabyimana A, Ruton H, Gaju E, et al. Assessing the perspectives of users and beneficiaries of a community health worker mHealth tracking system for mothers and children in Rwanda. PLoS One 2018;13:e0198725.
- 21 Perveen S, Mahmood MA, Lassi ZS, et al. Application of primary health care principles in national community health worker programs in low- and middle-income countries: a scoping review protocol. JBI Evid Synth 2021;19:270–83.
- 22 Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. Ann Intern Med 2018:169:467–73.
- 23 Covidence systematic review software. Veritas Health Innovation, Melbourne, Australia. Available: www.covidence.org

- 24 Arksey H, O'Malley L. Scoping studies: towards a methodological framework. Int J Soc Res Methodol 2005;8:19–32.
- World Health Organization. World Health organization and international bank for reconstruction and development / the world bank. tracking universal health coverage: 2017 global monitoring report. Geneva, Switzerland, 2017.
- 26 Ennever O, McFarquhar E, White S, et al. Survey of community health aides in Jamaica (1987-1988). West Indian Med J 1990;39:100–8.
- 27 Hafeez A, Mohamud BK, Shiekh MR, et al. Lady health workers programme in Pakistan: challenges, achievements and the way forward. J Pak Med Assoc 2011;61:210.
- 28 Hasegawa A, Yasuoka J, Ly P, et al. Integrating child health services into malaria control services of village malaria workers in remote Cambodia: service utilization and knowledge of malaria management of caregivers. *Malar J* 2013;12:292.
- 29 Koblinsky MA, Brechin SJ, Clark SD, et al. Helping managers to manage: work schedules of field-workers in rural Bangladesh. Stud Fam Plann 1989:20:225–34.
- 30 Kok MC, Kea AZ, Datiko DG, et al. A qualitative assessment of health extension workers' relationships with the community and health sector in Ethiopia: opportunities for enhancing maternal health performance. Hum Resour Health 2015;13:80.
- 31 Kosec K, Avula R, Holtemeyer B, et al. Predictors of essential health and nutrition service delivery in Bihar, India: results from household and frontline worker surveys. Glob Health Sci Pract 2015;3:255–73.
- 32 Magnani RJ, Rice JC, Mock NB, et al. The impact of primary health care services on under-five mortality in rural niger. Int J Epidemiol 1996;25:568–77.
- 33 Memon Z, Zaidi S, Riaz A. Residual barriers for utilization of maternal and child health services: community perceptions from rural Pakistan. *Glob J Health Sci* 2015;8:47–57.
- 34 Mues KE, Resende JC, Santos OCdos, et al. User satisfaction with the family health program in Vespasiano, Minas Gerais, Brazil. Rev Panam Salud Publica 2012;31:454–60.
- 35 Nasseri K, Sadrizadeh B, Malek-Afzali H, et al. Primary health care and immunisation in Iran. Public Health 1991;105:229–38.
- 36 Rubin G, Chen C, de Herrera Y, et al. Primary health care workers: the rural health aide program in El Salvador. Bull Pan Am Health Organ 1983:17:42–50.
- 37 Saprii L, Richards E, Kokho P, et al. Community health workers in rural India: analysing the opportunities and challenges accredited social health activists (ASHAs) face in realising their multiple roles. Hum Resour Health 2015;13:95.
- 38 Ved R, Scott K, Gupta G, et al. How are gender inequalities facing India's one million ASHAs being addressed? policy origins and adaptations for the world's largest all-female community health worker programme. Hum Resour Health 2019;17:3.
- 39 Wilford A, Phakathi S, Haskins L, et al. Exploring the care provided to mothers and children by community health workers in South Africa: missed opportunities to provide comprehensive care. BMC Public Health 2018:18:171.
- 40 Afsar HA, Qureshi AF, Younus M, et al. Factors affecting unsuccessful referral by the lady health workers in Karachi, Pakistan. J Pak Med Assoc 2003;53:521–8.
- 41 Afsar HA, Younus M. Recommendations to strengthen the role of lady health workers in the National program for family planning and primary health care in Pakistan: the health workers perspective. *J Ayub Med Coll Abbottabad* 2005;17:48–53.
- 42 Douthwaite M, Ward P. Increasing contraceptive use in rural Pakistan: an evaluation of the lady health worker programme. *Health Policy Plan* 2005;20:117–23.
- 43 Kohli C, Kishore J, Sharma S, et al. Knowledge and practice of accredited social health activists for maternal healthcare delivery in Delhi. J Family Med Prim Care 2015;4:359–63.
- 44 Panday S, Bissell P, van Teijlingen E, et al. The contribution of female community health volunteers (FCHVs) to maternity care in Nepal: a qualitative study. BMC Health Serv Res 2017;17:623.
- 45 Medhanyie A, Spigt M, Kifle Y, et al. The role of health extension workers in improving utilization of maternal health services in rural areas in Ethiopia: a cross sectional study. BMC Health Serv Res 2012;12:352
- 46 Admassie A, Abebaw D, Woldemichael AD. Impact evaluation of the Ethiopian health services extension programme. J Dev Effect 2009;1:430–49.
- 47 Aquino R, de Oliveira NF, Barreto ML. Impact of the family health program on infant mortality in Brazilian municipalities. Am J Public Health 2009:99:87–93.
- 48 Primary health care: report of the International Conference on primary health care, Alma-Ata, USSR, 6-12 September 1978: WHO, UNICEF 1978:61–3.



- 49 Cueto M. The origins of primary health care and selective primary health care. *Am J Public Health* 2004;94:1864–74.
- 50 Walsh JA, Warren KS. Selective primary health care: an interim strategy for disease control in developing countries. N Engl J Med 1979;301:967–74.
- 51 Tulenko K, Møgedal S, Afzal MM, et al. Community health workers for universal health-care coverage: from fragmentation to synergy. Bull World Health Organ 2013;91:847–52.
- 52 Perry H, Crigler L. Developing and strengthening community health worker programs at scale: a reference guide for program managers and policy makers. Dhaka, Bangladesh: University Press Ltd, 2013.
- 53 World Health Organization. WHO guideline on health policy and systems support to optimize community health worker programmes. Geneva: World Health Organiztion, 2018.
- 54 Lawn JE, Rohde J, Rifkin S, et al. Alma-Ata 30 years on: revolutionary, relevant, and time to revitalise. Lancet 2008;372:917–27.

- 55 Kane S, Kok M, Ormel H, et al. Limits and opportunities to community health worker empowerment: a multi-country comparative study. Soc Sci Med 2016;164:27–34.
- 56 Low A, Ithindi T. Adding value and equity to primary healthcare through partnership working to establish a viable community health workers' programme in Namibia. *Crit Public Health* 2003;13:331–46.
- 57 Kane S, Radkar A, Gadgil M. Community health workers as influential health system actors and not "just another pair of hands". *Int J Health Policy Manag* 2021;10:465–74.
- 58 Gilmore B, McAuliffe E. Effectiveness of community health workers delivering preventive interventions for maternal and child health in low- and middle-income countries: a systematic review. BMC Public Health 2013;13:847.
- 59 Boyce MR, Katz R. Community health workers and pandemic preparedness: current and prospective roles. Front Public Health 2019;7:62.

Supplementary Table 1: Key characteristics of included studies as reported by the authors

Author and year of publication / Country	Key objective of the study	Methods	Main findings
Damari 2018 / IRAN ¹⁷	To evaluate the national Iranian Women Health Volunteers program	Qualitative Document review One FGD Semi-structured questionnaires filled by 44 key informants	Achievements: Increased community participation, increasing health literacy, increased coverage and utilization of health services.
Nasseri 1991 / IRAN ³⁵	To determine the impact of PHC services on immunisation activities in areas where the two services are integrated	Quantitative • Cross-sectional survey	Higher coverage in rural areas is attributed to active approach of CHWs and vaccinators.
Memon 2016 / PAKISTAN ³³	To explore community barriers in accessing MCH services in 10 remote and rural districts of Pakistan	 Qualitative Sixty FGDs with mothers and fathers of children under five and CHWs - 20 each group 	Better awareness was seen among community caregivers for antenatal care and family planning services in the CHW-covered areas.
Hafeez 2011 / PAKISTAN ²⁷	To assess the contribution of the LHWP in enhancing coverage and access of health care services as well as towards improvement of health indicators	Qualitative Document review Interviews, formal and informal interactions and discussions with all the stakeholders Performance validation exercises in the field Feedback from community being served by the program	The LHWP has led to a development of a very well-placed cadre that links first-level care facilities to the community, thus improving the delivery of PHC services. The health indicators are significantly better than the national average in the areas served by the CHWs.
Douthwaite 2005 / PAKISTAN ⁴²	To assess the impact of the LHWP on the uptake of modern contraceptive methods	Quantitative • Secondary data analysis from the 2002 national evaluation of the LHWP	The study provides strong evidence that the LHWP has succeeded in integrating family planning into the doorstep provision of preventive health care and in increasing the use of modern reversible methods in rural areas.
Afsar 2005 / PAKISTAN ⁴¹	To assess the strengths and weaknesses of the LHWP from the Lady Health Workers perspective	Qualitative • 20 key informant interviews with CHWs (n=14), CHW Supervisors (n=4) and 2 medical officers (District	Major strengths: provision of services at the grassroots level, reinforcement of health messages and the community acceptability of workers. Weaknesses: contract-based job, low salaries, irregularity of payment, no career development, and poor logistical support.

Author and year of publication / Country	Key objective of the study	Methods	Main findings
		Coordinator and District Health Education Officer)	
Afsar 2003 / PAKISTAN ⁴⁰	To estimate the proportion of patients who were referred and to identify the factors associated with unsuccessful referral in Karachi, Pakistan	Quantitative • Cross-sectional survey of 347 patients	A high referral rate (55%) by CHWs was found in this study; 76.4% (n=265) were successful and 23.6% (n=82) were unsuccessful referrals. Key factors for unsuccessful referral: never referred before, never visited the referral site before, no knowledge of who to meet at the referral site, and failure of CHW to follow up.
Kohli 2015 / INDIA ⁴³	To assess the knowledge and practices for maternal health care delivery among Accredited Social Health Activist workers in North-East district of Delhi, India	Quantitative • Descriptive cross-sectional study (n = 55)	CHWs' knowledge is good but practices about maternal health were not adequate due to the number of problems faced by them which need to be addressed through skill- based training in terms of good communication and problem solving. Monitoring should be made an integral part of CHW working in the field to ensure that knowledge is converted into practices as well.
Kosec 2015 / INDIA ³¹	To understand predictors of essential health and nutrition service delivery in Bihar, India	Quantitative • Secondary data analysis of a 2012 cross-sectional survey of 6,002 households in 400 randomly selected villages in 1 district of Bihar state • Primary data collection from 382 CHWs	CHWs who maintained records of pregnant women were significantly associated with households receiving such information. Incentivizing frontline workers and helping them organize their work is associated with greater receipt of services by households.
Saprii 2015 / INDIA ³⁷	To explore stakeholders' perceptions and experiences of the CHW scheme in strengthening maternal health	Qualitative (exploratory study) Eighteen in-depth interviews and 3 FGDs with CHWs, key stakeholders and community members	CHWs are valued for their contribution towards maternal health education and for their ability to provide basic biomedical care, but their role as social activists is much less visible as envisioned in the CHW operational guidelines

Author and year of publication / Country	Key objective of the study	Methods	Main findings
Ved 2019 / INDIA ³⁸	To examine how the program is seeking to address gender inequalities facing CHWs, from the program's policy origins to recent adaptations	Qualitative	The value of community embeddedness for CHW programs is widely recognized as a mechanism to ensure program relevance to local needs and secure community ownership, support, and recognition of CHWs
Koblinsky 1989 / BANGLADESH ²⁹	To identify and examine organizational constraints to quality care and to provide a feasible strategy for program managers to overcome those barriers	Qualitative Observations FGDs – number not reported in the study	Only brief, interactions are possible if CHWs are to complete their rounds in the three-month period mandated by the government. The CHWs compensate for the pressure of their workload by skipping visits with some of the women in their area, by visiting even fewer during the monsoon season, and by neglecting to provide valuable information about family planning or health with some of the women they do visit
Panday 2019 / NEPAL ¹⁸	To explore use of MCH care services delivered by CHWs and the reasons for the underutilisation of these services	Qualitative • Interviews and FGDs with 34 CHWs, 26 service users and 11 health workers	Perceived factors that discourage the use of healthcare services by ethnic minority groups are; 1.Lack of knowledge among service users - related to CHWs' inability to communicate health messages; 2.Lack of trust in volunteers; 3.Traditional beliefs and healthcare practices; 4. Low decision-making power of women –

Author and year of publication / Country	Key objective of the study	Methods	Main findings
Panday 2017 / NEPAL ⁴⁴	To explore the role and experience of CHWs in maternal healthcare provision	Qualitative Interviews with 20 CHWs, 26 service users and 11 health workers Four FGDs with 18 CHWs	All study participants acknowledged the contribution of CHWs in basic maternity care in villages With support available to CHWs from the local health centres (regular training and access to medical supplies), CHWs were able to assist with childbirth, distribute medicines, and administer pregnancy tests. Whereas such activities were not reported in the other region where such support was not available to CHWs. Key challenge: lack of monetary incentives
Hasegawa 2013 / CAMBODIA ²⁸	To identify determinants of caregivers' Village Malaria Workers service utilization for childhood illness and caregivers' knowledge of malaria management	Quantitative Cross-sectional survey with CHWs and primary caregivers of children under five years	 Among the caregivers, 23% in M villages (villages with only malaria control services) and 52% in M+C villages (with both malaria and child health services) utilized CHW services for childhood illnesses. Determinants of caregivers' utilization of CHWs in M villages included their VMWs' length of experience (AOR = 11.80, 95% confidence interval [CI] = 4.46-31.19) and CHWs' service quality (AOR = 2.04, CI = 1.01-4.11). In M+C villages, CHWs' length of experience (AOR = 2.44, CI = 1.52-3.94) and caregivers' wealth index (AOR = 0.35, CI = 0.18-0.68) were associated with VMW service utilization. Better service quality of VMWs (AOR = 3.21, CI = 1.34-7.66) and caregivers' literacy (AOR = 9.91, CI = 4.66-21.05) were positively associated with caregivers' knowledge of malaria management.

Author and year of publication / Country	Key objective of the study	Methods	Main findings
Negussie 2017 / ETHIOPIA ¹⁹	To assess the contribution made by the CHWs in MCH care service delivery in Dale district, southern Ethiopia	Quantitative Cross-sectional survey with 613 mothers of reproductive age (15-49), having at least one under-five child	 Overall service coverage of antenatal care (four and more visits), delivery and postnatal care services were low in the district as compared to the national status; and the input from the CHWs, in this regard, was unsatisfactory. The number of home visits was also inadequate for the necessary support of the mothers. Mothers who listen to the radio and who had received information about the MCH services by CHWs were more likely to utilize MCH services.
Kok 2015 / ETHIOPIA ³⁰	To identify facilitators of and barriers to interpersonal relationships between CHWs and actors in the community and health sector	Qualitative • Fourteen FGDs and 44 interviews in 2013 with CHWs, traditional birth attendants, health professionals and community members	CHWs were selected by their communities, which enhanced trust and engagement between them Program design elements facilitating relationships: support for CHWs activities from the community and health sector, monitoring and accountability structures (community and health sector), referral, supervision and training (health sector)
Medhanyie 2012 / ETHIOPIA ⁴⁵	To investigate the role of CHWs in improving utilization of maternal health services by rural women	Quantitative Cross-sectional survey with 725 women with under-five children	CHWs have contributed substantially to the improvement in women's utilization of family planning, antenatal care and HIV testing.
Admassie 2009 / ETHIOPIA ⁴⁶	To evaluate the short-term and intermediate-term effects of the Ethiopian HEP on MCH indicators	Program evaluation using a propensity score matching method and village, facility and household surveys	 HEP has significantly increased the proportion of children fully and individually vaccinated Women in the HEP villages appeared to make their first contact with a skilled health service provider significantly earlier during pregnancy; very little effect is detected on other prenatal and postnatal care services. HEP has not reduced the incidence and duration of diarrhoea and respiratory diseases among under-five children
Musabyimana 2018 / RWANDA ²⁰	To explore perceptions of healthcare officials, providers, and beneficiaries on the impact of the RapidSMS program	Qualitative • 10 FGDs with 93 participants • In-depth interviews with 56 beneficiaries and 36 CHWs	The effectiveness of use of mobile phones to remind of the appointments for improved access to midwifery services at the health facilities was found to be limited. Indirectly, it alerts to the emerging role of contemporary technologies in community health program.

Author and year of publication / Country	Key objective of the study	Methods	Main findings
Magnani 1996 / NIGER ³²	To assess the impact of differential access to health services through the comparison of service use patterns and under-five mortality levels among villages provided different levels of health services	Quantitative • Secondary data analysis of National Morbidity and Mortality Survey – 1985 on 974 women of reproductive age	Children residing in villages proximate to health dispensaries were approximately 32% less likely to have died during the study period than children living further away.
Wilford 2018 / SOUTH AFRICA ³⁹	To explore the quality of CHW household visits providing MCH services	Qualitative • 30 observations [a CHW visit to a mother or pregnant woman was observed by a field worker, followed by an in-depth interview with the participating women and CHWs] • 15 in-depth interviews with mothers/pregnant women and 15 in-depth interviews with CHWs	Mothers receiving the services were satisfied with CHW visits and appreciated that CHWs understood their life experiences and provided relevant and accessible advice and support. CHWs expressed concern of not having the required knowledge to undertake all activities in the household, and requested training and support from supervisors during household visits
Mues 2012 / BRAZIL ³⁴	To assess factors influencing perspectives on Brazil's national family health program and perceptions about PSF accessibility among frequent users (primary caretakers of children under 5)	Ouantitative Cross-sectional household survey of 253 households with at least one child 5 years or younger and covered by the PSF	 Most caretakers of young children were satisfied. However, less than half of the caretakers perceived the PSF unit as being accessible about a quarter of households in the Vespasiano PSF coverage area were not receiving an agent home visit once a month
Aquino 2009 / BRAZIL ⁴⁷	To evaluate the effects of the implementation of the CHW Program on infant mortality rates in Brazilian municipalities from 1996 to 2004	Quantitative – ecological and longitudinal approach • Secondary data analysis from 1991 and 2000 national census and data from Brazilian MoH of 721 municipalities	A statistically significant negative association between CHW program coverage and infant mortality rate was found after controlling for potential confounders.

Author and year of publication / Country	Key objective of the study	Methods	Main findings
Rubin 1983 / EL SAVADOR ³⁶	To evaluate the health service impact of the Rural Health Aide Program in El Salvador	Quantitative • Survey of 363 respondents in cantons served by CHWs for one year and 169 in cantons served by CHWs for two years	Compared to villagers of cantons served by CHWs for one year, those in cantons served by CHWs for 2 years were: -more likely to be visited by their CHW & to visit their CHW -more likely to visit their health centres after referral by their CHW -more likely to have their children vaccinated
Ennever 1990 / JAMAICA ²⁶	 To describe the activities of CHWs currently employed, and their perceptions about supervision and management To describe the current employment status of CHWs who had left the service between 1982 and 1986, and use of the skills they had learned as CHWs. 	Quantitative • Survey of 415 CHWs currently employed and 134 CHWs who had left the service	Currently employed CHWs continued to perform duties in the community & in health centres with emphasis on the MCH services and the management of diabetics and hypertensives. Previously employed CHWs unemployed though many continued to use their skills on a voluntary basis.

CHW = Community Health Worker, FGD = Focus Group Discussion, HEP = Health Extension Program (Ethiopia) LHWP = Lady Health Worker Program (Pakistan), MCH = Maternal and Child Health, PSF = Programa de Saude da Familia (Family Health Program, Brazil)

Supplementary Table 2: Evidence for the application of primary health care principles as reflected in the national community health worker programs

Seri al No.	Country / CHW Program / year commenced	CHW Program Objective	Implementation of the CHW Program	Stated Outcome/ achievement of the CHW Program
1.	IRAN / Women Health Volunteers Program / 1992	Principle observed: - Community Participation as the program aims to increase community involvement in health related activities in order to empower them	Principles observed: - UHC - Community Participation* • The CHWs encouraged and actively followed up on individuals to visit health centres at their required time especially those who needed special care thus contributing to increased service utilisation • CHWs delivering health messages to families and distributing educational materials reflect one aspect of comprehensiveness as part of universal health coverage	Principles observed: - UHC - Community Participation* - Intersectoral coordination - The active follow up by WHV increased utilization of health services – contributing to universal health coverage • The experts and stakeholders believed that CHW program increased people's participation and created self-esteem and self-reliance in people – However, the evidence on how it achieved this is not available in this study
			CHWs are selected from the local community - Community Participation and appropriateness	The WHV network connects MoH, medical universities and health centers to the people – Intersectoral coordination
2.	IRAN / Primary Health Care Network – EPI / 1983 ³⁵	Principle observed: - UHC - As the program aimed to increase immunisation coverage in Iranian children to 90% by their first birthday	Principles observed: - UHC - Community Participation* - CHWS were involved in provision of general preventive services for all the individuals in their coverage area – Comprehensiveness, Universal health coverage - CHWs were also expected to provide basic therapeutic measures for minor illnesses and refer other cases to their immediate Rural Health Centre – universal health coverage - CHWs were selected from the same area in which they work – community participation	Principle observed: - UHC - Appropriateness • Immunisation coverage of children improved significantly in 1987 as compared to 1984 especially for BCG (56.3%) - universal health coverage • Mothers in rural areas with PHC services receive much better MCH care, advice and attention in comparison to mothers in other rural and most urban areas – appropriateness

Seri al No.	Country / CHW Program / year commenced	CHW Program Objective	Implementation of the CHW Program	Stated Outcome/ achievement of the CHW Program
3.	PAKISTAN / National Program for Family Planning & Primary Health Care / 1994 ^{27 33}	Principle observed: - UHC as the program aimed to increase utilisation of promotive, preventive and curative services at the community level particularly for women and children in poor and underserved areas – comprehensivenes s & equity	Principles observed: - UHC - Community Participation* • CHWs were involved in health education and community mobilization along with provision of immunization, family planning services, basic curative care to the community at the doorstep and referral of patients to the appropriate health facility - reflecting universal health coverage	Principles observed: - UHC - Community Participation* • Increased utilisation of antenatal care and family planning - universal health coverage • Improved infant mortality rate, maternal mortality ratio and contraceptive prevalence rate in CHW covered areas as compared to national average - universal health coverage • Cultural acceptability of CHWs, unlimited access to households and free interaction with local women – community participation and appropriateness
4.	INDIA / Accredited Social Health Activist (ASHA) Program / 2005 31 37 38	Principles observed: - UHC through accessible care to rural population especially vulnerable groups - Appropriateness via provision of affordable and quality health care	Principles observed: - UHC via CHWs as 'service extension and link workers' - Community Participation as CHWs are selected from the local communities	Principles observed: - UHC as CHWs were motivating women for antenatal care and hospital delivery through home visits • Women empowerment – as CHWs have reported an increased sense of empowerment and personal growth, in part through their belief in the social value of their work. • Additionally, becoming a CHW enabled rural women to gain knowledge, status as a role model, and exposure beyond the village, as well as to access a limited amount of remuneration
5.	BANGLADESH / National MCH and Family Planning Program / 1976 ²⁹	Not reported	Principles observed: - UHC - Community Participation* • CHWs were utilised for health education and extending immunisation and family planning services at the household level. They also provided referral for antenatal, perinatal, and	Not reported

Seri al No.	Country / CHW Program / year commenced	CHW Program Objective	Implementation of the CHW Program	Stated Outcome/ achievement of the CHW Program
			postnatal care. – comprehensiveness as part of universal health coverage	
6.	NEPAL / Female Community Health Volunteer Program / 1988 18	Principles observed: - UHC via low cost health service provision in remote areas - Community Participation via increase in local women's participation in health promotion	Principles observed: - Community Participation* - UHC via provision of MCH care by CHWs in rural communities	Not reported
7.	CAMBODIA / Village Malaria Worker Project as part of National Malaria Control Program / 2001 ²⁸	Not reported	Principles observed: - UHC - Community Participation* • Malaria prevention, diagnosis and treatment services to remote villages by CHWs — universal health coverage - Management of minor childhood illness, prescribing and providing basic medications, referral and health promotion — comprehensiveness as part of universal health coverage	Principle observed: - UHC - 15,898 children received child health services from village Malaria Workers in 2011
8.	ETHIOPIA / Health Extension Program / 2003 ¹⁹ 30	Principles observed: - UHC - Community Participation - To improve access and utilization of health care particularly for	Principles observed: - UHC - Community Participation • CHWs providing antenatal and postnatal care, family planning and immunization services and conducting clean and safe deliveries - Universal Health Coverage	Principles observed: - UHC - Community Participation • Increased use of health post for antenatal care, family planning, delivery and other illnesses such as diarrhoea – reflecting universal health coverage

Seri al No.	Country / CHW Program / year commenced	CHW Program Objective	Implementation of the CHW Program	Stated Outcome/ achievement of the CHW Program
		children and mothers in rural communities – Universal Health Coverage To improve the health status of families with their full participation, using local technologies & the community's skill & knowledge - Community Participation	Quarterly evaluation of health centers performance by the community during facility or public forums. Monitoring of CHWs by the kebele (lowest administrative unit) administration at the health post level. Need based adjustment of maternal health education — Community Participation	 Statistically significant increase in the proportion of children fully and individually vaccinated against tuberculosis, polio, diphtheria–pertussis–tetanus, and measles in the program villages. Mothers reported that CHWs were available at health posts during their last visit for MCH services Mothers also indicated that they had gotten a complete explanation of their own/child's health condition from the CHWs Moreover, CHWs were understanding, friendly and helpful thus assured a "natural link" between them and the community - appropriateness Community members reported that HEWs being female was important to them, as they prefer to discuss maternal health issues amongst women - appropriateness
9.	RWANDA / RapidSMS program / 2013 ²⁰	Principles observed: - UHC - Appropriateness • To improve access to antenatal, PNC, institutional delivery and emergency obstetric care • To facilitate communication between CHWs and the broader health system, including the ambulance system,	Principles observed: - UHC - Community Participation* - Appropriateness – use of technology • The RapidSMS system sent automatic reminders to CHWs for clinical appointments, delivery, and post-natal care visits, with the intent of increasing timely access and utilization • Provision of a quick link to emergency obstetric care through so-called Red Alerts and creation of a database of clinical records on maternal care delivery – use of technology for increasing access to health care	Principles observed: Appropriateness (use of technology, acceptability) RapidSMS was well accepted by most CHWs and community members — acceptability aspect of appropriateness principle mHealth appeared to have helped improve communication and potentially service use Claims that mHealth has contributed to maternal mortality reduction are not substantiated considering the difficulties that were highlighted by the respondents

Seri al No.	Country / CHW Program / year commenced	CHW Program Objective	Implementation of the CHW Program	Stated Outcome/ achievement of the CHW Program
		health facilities, and MoH officials		
10.	NIGER / Rural Health Improvement Program / 1970s	Principle observed: - UHC – as the program aimed to extend the coverage of PHC services throughout rural Niger	Principle observed: - UHC – By upgrading existing health dispensaries and deploying trained village health teams to unserved villages to deliver PHC services	Not reported
11.	SOUTH AFRICA / ward-based outreach teams (WBOT) - national CHW program / 2011 39	Principle observed: - UHC – via improving health outcomes by providing home and community- based health services	Principle observed: - UHC - Community Participation* • Universal health coverage via CHWs providing treatment support and home-based care in underserved rural areas. Core MCH activities include visiting all mothers during pregnancy, antenatal education and support. Moreover, CHWs are linked in with local PHC clinics	Principle observed: - Appropriateness as CHWs were trusted, accessible and able to understand the mother's situation
12.	BRAZIL / Family Health Program (Programa de Saude da Familia, PSF) / 1994 ³⁴	Principle observed: - UHC – as the organizational principles include universality and equity	Principle observed: - UHC - Community Participation* - Universal health coverage via provision of promotive, preventive and basic curative services by CHWs to mothers and children	Principle observed: UHC – as the growth of the CHW program was associated with a decrease in infant and child mortality rates Caretakers who reported that their agent made at least one home visit per month were significantly more likely to have received care for child diarrhoea from an agent
13.	EL SAVADOR / Rural Health Aide Program / 1976 ³⁶	Principle observed: - UHC – via provision of PHC and family planning services	Principle observed: - UHC - Community Participation* • Health education by CHWs for rural families • Provision of family planning supplies to women	Principle observed: - UHC - Appropriately trained PHC workers promote contact between rural populations and the health care system

Seri al No.	Country / CHW Program / year commenced	CHW Program Objective	Implementation of the CHW Program	Stated Outcome/ achievement of the CHW Program
			 Provision of systematic treatment of minor illnesses; administration of prescribed intramuscular injections; dispensing of antiparasitic medication; and performance of simple first-aid measures Promotion of registration of births and deaths 	To the extent that this improves the health status of the population, particularly in the area of MCH, we might expect to see better health indices in rural populations served by these workers than in populations without them
14.	JAMAICA / Community Health Aide program / 1978 ²⁶	Principle observed: - UHC as the program aimed to train local women to provide basic health care and health education to families.	Principles observed: - UHC – CHWs encouraging for immunization and family planning, weighing babies and testing urine - Community Participation*	Principle observed: - UHC • CHWs have been functioning in both health centre and community, encouraging people to utilize the services and assisting in some of the less technical duties such as weighing babies and testing urine

UHC = Universal Health Coverage

Appendix I: Logic grids for information sources

<u>PubMed</u>

Search	Query	Records retrieved
#1	"community health workers" [mh] OR community health worker* [tiab] OR community health aide* [tiab] OR village health worker* [tiab] OR barefoot doctor* [tiab] OR family planning personnel* [tiab] OR health extension worker* [tiab] OR lady health worker* [tiab] OR community health agent* [tiab] OR Shasthyo Sebika* [tiab] OR community nutrition worker* [tiab] OR maternal health worker* [tiab] OR village malaria worker* [tiab] OR Raedat* [tiab] OR postnatal support worker* [tiab] OR mental health worker* [tiab] OR mother coordinator* [tiab] OR rural health worker* [tiab] OR village health promoter* [tiab] OR accompagnateur* [tiab] OR Saksham Sahaya* [tiab] OR anganwandi worker* [tiab] OR accredited social health activist* [tiab] OR community-based worker* [tiab] OR community health volunteer* [tiab] OR willage health guide* [tiab] OR maternal and child health promotion worker* [tiab] OR maternal child health worker* [tiab] OR kader posyandu* [tiab] OR behvarz* [tiab] OR village health helper* [tiab] OR colaborador voluntario* [tiab] OR nutrition volunteers* [tiab] OR village drug-kit manager* [tiab] OR brigadistas* [tiab] OR female community health volunteer* [tiab] OR Agente Comunitario de Salud* [tiab] OR nutrition worker* [tiab] OR community reproductive health worker* [tiab] OR community drug distributor* [tiab] OR community volunteer* [tiab] OR community health advocate* [tiab] OR lay health visitor* [tiab] OR Promotoras de Salud [tiab]	174984
#2	Program[tiab] OR programs[tiab] OR programme[tiab] OR programmes[tiab] OR initiative*[tiab] OR project[tiab] OR projects[tiab]	959578
#3	"Maternal health"[mh] OR "Maternal Welfare"[mh] OR "child health"[mh] OR "child care"[mh] OR "child welfare"[mh] OR "maternal-child health services"[mh] OR "child health services"[mh] OR "child health services"[mh] OR maternal newborn child health[tiab]	71349

Search	Query	Records retrieved
#4	[(developing country[tw] OR developing countries[tw] OR developing populations[tw] OR developing populations[tw] OR developing populations[tw] OR less developed countries[tw] OR less developed nations[tw] OR less developed nation[tw] OR less developed nation[tw] OR less developed nation[tw] OR less developed nation[tw] OR less developed population[tw] OR less developed populations[tw] OR lesser developed nations[tw] OR lesser developed populations[tw] OR lesser developed nation[tw] OR lesser developed populations[tw] OR lesser developed nation[tw] OR under developed nation[tw] OR under developed countries[tw] OR under developed nation[tw] OR underdeveloped countries[tw] OR underdeveloped countries[tw] OR underdeveloped nation[tw] OR underdeveloped nation[tw] OR underdeveloped nation[tw] OR underdeveloped populations[tw] OR underdeveloped nation[tw] OR underdeveloped nation[tw] OR underdeveloped nation[tw] OR underdeveloped nation[tw] OR underdeveloped countries[tw] OR underdeveloped countries[tw] OR underdeveloped countries[tw] OR underdeveloped nation[tw] OR lower income nation[tw] OR underserved nation[tw] OR underdeveloped nation[tw] OR underdeveloped nation[tw] OR lower income nation[tw] OR deprived nation[tw] OR deprived nation[tw]	1903167

Search	Query	Records retrieved
	Republic(tw) OR East Timor(tw) OR East Timur(tw) OR Timor Leste(tw) OR Ecuador(tw) OR Egypt(tw) OR United Arab Republic(tw) OR El Salvador(tw) OR Eritrea(tw) OR Estionia(tw) OR Ethiopia(tw) OR Fility) OR Gaboniese Republic(tw) OR Gaza(tw) OR Georgian Republic(tw) OR Georgian Republic(tw) OR Gaza(tw) OR Ga	retrieved

Search	Query	Records retrieved
	Cote d'Ivoire[Mesh:noexp] OR Croatia[Mesh:noexp] OR Cuba[Mesh:noexp] OR Cyprus[Mesh:noexp] OR "Democratic Republic of the Congo"[Mesh:noexp] OR Dominica[Mesh:noexp] OR Dominica[Mesh:noexp] OR Dominica[Mesh:noexp] OR East Timor[Mesh:noexp] OR Ecuador[Mesh:noexp] OR Egypt[Mesh:noexp] OR El Salvador[Mesh:noexp] OR Eritrea[Mesh:noexp] OR Estonia[Mesh:noexp] OR Ethiopia[Mesh:noexp] OR Egypt[Mesh:noexp] OR Gabon[Mesh:noexp] OR Gambia[Mesh:noexp] OR Gerece[Mesh:noexp] OR Gambia[Mesh:noexp] OR Gambia[Mesh:noexp] OR Guna[Mesh:noexp] OR Gambia[Mesh:noexp] OR Guna[Mesh:noexp] OR Gambia[Mesh:noexp] OR Guna[Mesh:noexp] OR Guna[Mesh:noexp] OR Guna[Mesh:noexp] OR Guna[Mesh:noexp] OR Hungary[Mesh:noexp] OR Hungary[Mesh:noexp] OR Iraq[Mesh:noexp] OR Hungary[Mesh:noexp] OR Iraq[Mesh:noexp] OR Iraq[Mesh:noexp] OR Hungary[Mesh:noexp] OR Iraq[Mesh:noexp] OR Kosovo[Mesh:noexp] OR Kosovo[Mesh:noexp] OR Lavia[Mesh:noexp] OR Kosovo[Mesh:noexp] OR Kosovo[Mesh:noexp] OR Libra[Mesh:noexp] OR Lavia[Mesh:noexp] OR Lavia[Mesh:noexp] OR Lavia[Mesh:noexp] OR Macadonia[Mesh:noexp] OR Morocco[Mesh:noexp] OR Morocco[Mesh:noexp] OR Morocco[Mesh:noexp] OR Morocco[Mesh:noexp] OR Morocco[Mesh:noexp] OR Morocco[Mesh:noexp] OR Noexp] OR Noexp] OR Noexp] OR Noexp] OR Paraguay[Mesh:noexp] OR Noexp] OR Noexp] OR Noexp] OR Noexp] OR Paraguay[Mesh:noexp] OR Noexp] OR Noexp] OR Noexp] OR Paraguay[Mesh:noexp] OR Paraguay[Mesh:noexp] OR Paraguay[Mesh:noexp] OR Romania[Mesh:noexp] OR Romania[Mesh:noexp] OR Romania[Mesh:noexp] OR Romania[Mesh:noexp] OR Romania[Mesh:noexp] OR Romania[Mesh:noexp] OR Saint Kitts and Nevis"[
#5	#1 AND #2 AND #3 AND #4	956
Limited to	1978 onwards in English language only	863

CINAHL

Community health worker	Program	МСН	LMIC
MH "community health workers" OR MH	TX Program OR	MH "Maternal-	MH "low and middle income countries" OR MH "developing
"rural health personnel" OR TX "community	TX programs OR	Child Health"	countries" OR
health worker*" OR TX "community health"	TX programme	OR TX	TX Afghanistan OR TX Albania OR TX Algeria OR TX Angola OR TX
aide*" OR TX "village health worker*" OR TX	OR TX	"maternal-child	Antigua OR TX Barbuda OR TX Argentina OR TX Armenia OR TX
"barefoot doctor*" OR TX "family planning	programmes OR	health"	Armenian OR TX Aruba OR TX Azerbaijan OR TX Bahrain OR TX
personnel*" OR TX "health extension	TX initiative* OR		Bangladesh OR TX Barbados OR TX Benin OR TX Byelarus OR TX
worker*" OR TX "lady health worker*" OR TX	TX project OR TX		Byelorussian OR TX Belarus OR TX Belorussian OR TX Belorussia
"community health agent*" OR TX "Shasthyo	projects		OR TX Belize OR TX Bhutan OR TX Bolivia OR TX Bosnia OR TX
Sebika*" OR TX "community nutrition			Herzegovina OR TX Hercegovina OR TX Botswana OR TX Brasil OR
worker*" OR TX "maternal health worker*"			TX Brazil OR TX Bulgaria OR TX Burkina Faso OR TX Burkina Fasso
OR TX "voluntary Malaria worker*" OR TX			OR TX Upper Volta OR TX Burundi OR TX Urundi OR TX Cambodia
"village malaria worker*" OR TX "Raedat*"			OR TX Khmer Republic OR TX Kampuchea OR TX Cameroon OR
OR TX "postnatal support worker*" OR TX			TX Cameroons OR TX Cameron OR TX Camerons OR TX Cape
"mental health worker*" OR TX "mother			Verde OR TX "Central African Republic" OR TX Chad OR TX Chile
coordinator*" OR TX "rural health worker*"			OR TX China OR TX Colombia OR TX Comoros OR TX "Comoro
OR TX "village health promoter*" OR TX			Islands" OR TX Comores OR TX Mayotte OR TX Congo OR TX Zaire
accompagnateur* OR TX "Saksham			OR TX "Costa Rica" OR TX "Cote d'Ivoire" OR TX "Ivory Coast" OR
Sahaya*" OR TX "anganwandi worker*" OR			TX Croatia OR TX Cuba OR TX Cyprus OR TX Czechoslovakia OR
TX "accredited social health activist*" OR TX			TX "Czech Republic" OR TX Slovakia OR TX "Slovak Republic" OR
"community-based worker*" OR TX			TX Djibouti OR TX "French Somaliland" OR TX Dominica OR TX
"community health volunteer*" OR TX "village			"Dominican Republic" OR TX "East Timor" OR TX "East Timur" OR
health guide*" OR TX "maternal and child			TX "Timor Leste" OR TX Ecuador OR TX Egypt OR TX "United Arab
health promotion worker*" OR TX "maternal			Republic" OR TX "El Salvador" OR TX Eritrea OR TX Estonia OR TX
child health worker*" OR TX "kader			Ethiopia OR TX Fiji OR TX Gabon OR TX "Gabonese Republic" OR
posyandu*" OR TX behvarz* OR TX "village			TX Gambia OR TX Gaza OR TX "Georgia Republic" OR TX
health helper*" OR TX "colaborador			"Georgian Republic" OR TX Ghana OR TX "Gold Coast" OR TX
voluntario*" OR TX "nutrition volunteers*" OR			Greece OR TX Grenada OR TX Guatemala OR TX Guinea OR TX
TX "village drug-kit manager*" OR TX			Guam OR TX Guiana OR TX Guyana OR TX Haiti OR TX Honduras
brigadistas* OR TX "female community			OR TX Hungary OR TX India OR TX Maldives OR TX Indonesia OR
health volunteer*" OR TX "Agente			TX Iran OR TX Iraq OR TX "Isle of Man" OR TX Jamaica OR TX
Comunitario de Salud*" OR TX "nutrition			Jordan OR TX Kazakhstan OR TX Kazakh OR TX Kenya OR TX
worker*" OR TX "community reproductive			Kiribati OR TX Korea OR TX Kosovo OR TX Kyrgyzstan OR TX
health worker*" OR TX "community drug			Kirghizia OR TX "Kyrgyz Republic" OR TX Kirghiz OR TX Kirgizstan
distributor*" OR TX "community volunteer*"			OR TX "Lao PDR" OR TX Laos OR TX Latvia OR TX Lebanon OR

Community health worker	Program	МСН	LMIC
OR TX "community health advocate*" OR TX "lay health visitor*" OR TX "Promotoras de Salud"			TX Lesotho OR TX Basutoland OR TX Liberia OR TX Libya OR TX Lithuania OR TX Macedonia OR TX Madagascar OR TX "Malagasy Republic" OR TX Malaysia OR TX Malay OR TX Malay OR TX Sabah OR TX Sarawak OR TX Malawi OR TX Nyasaland OR TX Mali OR TX Malta OR TX "Marshall Islands" OR TX Mauritania OR TX Mauritius OR TX "Marshall Islands" OR TX Mexico OR TX Micronesia OR TX "Middle East" OR TX Moldova OR TX Moldovia OR TX Morocco OR TX Ifni OR TX Mozambique OR TX Myanmar OR TX Myanma OR TX Burma OR TX Namibia OR TX Nepal OR TX "Netherlands Antilles" OR TX "New Caledonia" OR TX Nicaragua OR TX Niger OR TX Nigeria OR TX "Northern Mariana Islands" OR TX Oman OR TX Muscat OR TX Pakistan OR TX Palau OR TX Palestine OR TX Panama OR TX Paraguay OR TX Peru OR TX Phillippines OR TX Palau OR TX Rumania OR TX Russian OR TX Saint Lucia" OR TX "St Kitts" OR TX "Saint Vincent" OR TX "St Vincent" OR TX "Saint Vincent" OR TX "Sa Or Tome" OR TX "Saudi Arabia" OR TX "Saint Vincent" OR TX "Sa Or Tome" OR TX Seychelles OR TX "Sierra Leone" OR TX Slovenia OR TX "Saudi Arabia" OR TX "Sierra Leone" OR TX Somania OR TX Suriname OR TX Surinam OR TX Surinam OR TX Surinam OR TX Surinam OR TX Tadzhikistan OR TX Tadjikistan OR TX Tadzhikistan OR TX Tanzania OR TX Trinidad OR TX Tobago OR TX Tunisia OR TX Turkey OR TX Turkmenistan OR TX Tobago OR TX Tunisia OR TX Turkey OR TX Turkmenistan OR TX Turkmen OR TX Uzganda OR TX Uzbekistan OR TX Uzbekistan OR TX Vietnam OR TX "Viet Nam" OR TX "Soviet Union" OR TX "Uzbek OR TX Vietnam OR TX "Viet Nam" OR TX "West Bank" OR TX Yemen OR TX Vietnam OR TX "Viet Nam" OR TX "West Bank" OR TX Yemen OR TX Vietnam OR TX "Viet Nam" OR TX "West Bank" OR TX Yemen OR TX Turgoslavia OR TX Zambia OR TX Zimbabwe OR TX Rhodesia

EMBASE

Community health worker	Program	МСН	LMIC
"Health Auxiliary"/de OR "community health worker*":ti,ab OR "community health aide*":ti,ab OR "village health worker*":ti,ab OR "barefoot doctor*":ti,ab OR "family planning personnel*":ti,ab OR "health extension worker*":ti,ab OR "lady health worker*":ti,ab OR "community health agent*":ti,ab OR "Shasthyo Sebika*":ti,ab OR	Program:ti,ab OR programs:ti,ab OR programme:ti,ab OR programmes:ti,a b OR initiative*:ti,ab OR project:ti,ab	"Maternal child health care"/de OR "Maternal Welfare":ti,ab OR "child health":ti,ab OR "child care":ti,ab OR "child welfare":ti,ab OR "maternal-child	Afghanistan:ti,ab OR Albania:ti,ab OR Algeria:ti,ab OR Angola:ti,ab OR Antigua:ti,ab OR Barbuda:ti,ab OR Argentina:ti,ab OR Armenia:ti,ab OR Armenia:ti,ab OR Armenia:ti,ab OR Bangladesh:ti,ab OR Barbados:ti,ab OR Benin:ti,ab OR Byelarus:ti,ab OR Byelorussian:ti,ab OR Belorussian:ti,ab OR Belorussian:ti,ab OR Belorussia:ti,ab OR Belize:ti,ab OR Bhutan:ti,ab OR Bosnia:ti,ab OR Belorussia:ti,ab OR Belorussia:ti,ab OR Belorussia:ti,ab OR Belorus:ti,ab OR Bosnia:ti,ab OR Belorus:ti,ab OR Bosnia:ti,ab OR Brazil:ti,ab OR Brazil:ti,ab OR Bulgaria:ti,ab OR Burkina Faso:ti,ab OR "Burkina Faso:":ti,ab OR "Upper Volta":ti,ab OR Burundi:ti,ab OR Urundi:ti,ab OR Cambodia:ti,ab OR "Khmer Republic":ti,ab OR Kampuchea:ti,ab OR Cameroon:ti,ab OR
"community nutrition worker*":ti,ab OR "maternal health worker*":ti,ab OR "voluntary Malaria worker*":ti,ab OR "village malaria worker*":ti,ab OR Raedat*:ti,ab OR "postnatal support worker*":ti,ab OR "mental health worker*":ti,ab OR "mother coordinator*":ti,ab OR "rural health worker*":ti,ab OR "rural health worker*":ti,ab OR "saksham Sahaya*":ti,ab OR "anganwandi worker*":ti,ab OR "accredited social health activist*":ti,ab OR "community- based worker*":ti,ab OR "community health volunteer*":ti,ab OR "community health volunteer*":ti,ab OR	OR projects:ti,ab	health services":ti,ab OR "child health services":ti,ab OR "maternal child health":ti,ab OR "maternal newborn child health":ti,ab	Cameroons:ti,ab OR Cameron:ti,ab OR Camerons:ti,ab OR "Cape Verde":ti,ab OR "Central African Republic":ti,ab OR Chad:ti,ab OR Chile:ti,ab OR China:ti,ab OR Colombia:ti,ab OR Comoros:ti,ab OR "Comoro Islands":ti,ab OR Comores:ti,ab OR Mayotte:ti,ab OR Comgo:ti,ab OR Zaire:ti,ab OR "Costa Rica":ti,ab OR "Cote d Ivoire":ti,ab OR "Ivory Coast":ti,ab OR Croatia:ti,ab OR Cuba:ti,ab OR Cyprus:ti,ab OR Czechoslovakia:ti,ab OR "Czech Republic":ti,ab OR Slovakia:ti,ab OR "Slovak Republic":ti,ab OR Djibouti:ti,ab OR "French Somaliland":ti,ab OR Dominica:ti,ab OR "Dominican Republic":ti,ab OR "East Timor":ti,ab OR "El Salvador":ti,ab OR Egypt:ti,ab OR Estonia:ti,ab OR Ethiopia:ti,ab OR Fiji:ti,ab OR Gabon:ti,ab OR "Gabonese Republic":ti,ab OR Gambia:ti,ab OR Gaza:ti,ab OR "Georgia Republic":ti,ab OR "Georgian Republic":ti,ab OR Gold Coast:ti,ab OR Greece:ti,ab OR Guiana:ti,ab OR Guyana:ti,ab OR Haiti:ti,ab OR Honduras:ti,ab OR Hungary:ti,ab OR India:ti,ab OR Maldives:ti,ab OR Indonesia:ti,ab OR Jordan:ti,ab OR Kazakhstan:ti,ab OR Kazakhsti,ab OR Jamaica:ti,ab OR Jordan:ti,ab OR Kazakhstan:ti,ab OR Kazakhsti,ab OR
"maternal and child health promotion worker*":ti,ab OR "maternal child health worker*":ti,ab OR "kader posyandu*":ti,ab OR behvarz*:ti,ab OR "village health helper*":ti,ab OR "colaborador			Kenya:ti,ab OR Kiribati:ti,ab OR Korea:ti,ab OR Kosovo:ti,ab OR Kyrgyzstan:ti,ab OR Kirghizia:ti,ab OR "Kyrgyz Republic":ti,ab OR Kirghiz:ti,ab OR Kirgizstan:ti,ab OR Lao PDR:ti,ab OR Laos:ti,ab OR Latvia:ti,ab OR Lebanon:ti,ab OR Lesotho:ti,ab OR Basutoland:ti,ab OR Liberia:ti,ab OR Libya:ti,ab OR Lithuania:ti,ab OR Macedonia:ti,ab OR Madagascar:ti,ab OR "Malagasy Republic":ti,ab OR Malaysia:ti,ab OR Malaya:ti,ab OR Malay:ti,ab OR

Community health worker	Program	MCH	LMIC
voluntario*":ti,ab OR "nutrition volunteers*":ti,ab OR "village drug-kit manager*":ti,ab OR "female community health volunteer*":ti,ab OR "Agente Comunitario de Salud*":ti,ab OR "nutrition worker*":ti,ab OR "community reproductive health worker*":ti,ab OR "community volunteer*":ti,ab OR "community volunteer*":ti,ab OR "community health advocate*":ti,ab OR "lay health visitor*":ti,ab OR "Promotoras de Salud":ti,ab			Sabah:ti,ab OR Sarawak:ti,ab OR Malawi:ti,ab OR Nyasaland:ti,ab OR Malit:ti,ab OR "Marshall Islands":ti,ab OR Mauritiun:ti,ab OR Mauritiun:ti,ab OR "Agalega Islands":ti,ab OR Mexico:ti,ab OR Micronesia:ti,ab OR "Middle East":ti,ab OR Moldova:ti,ab OR Moldovia:ti,ab OR Moldovia:ti,ab OR Moldovia:ti,ab OR Moldovia:ti,ab OR Mongolia:ti,ab OR Montenegro:ti,ab OR Morocco:ti,ab OR Burma:ti,ab OR Mozambique:ti,ab OR Myanmar:ti,ab OR Myanma:ti,ab OR Burma:ti,ab OR Namibia:ti,ab OR Nepal:ti,ab OR "Netherlands Antilles":ti,ab OR "New Caledonia":ti,ab OR Nicaragua:ti,ab OR Niger:ti,ab OR Nigeria:ti,ab OR "Northern Mariana Islands":ti,ab OR OR OR Denami:ti,ab OR Palau:ti,ab OR Phillipines:ti,ab OR Portugal:ti,ab OR Palau:ti,ab OR Pollipines:ti,ab OR Pollipines:ti,ab OR Phillipines:ti,ab OR Pollipines:ti,ab OR Pollipines:ti,ab OR Pollipines:ti,ab OR Romania:ti,ab OR Romania:ti,ab OR Romania:ti,ab OR Romania:ti,ab OR Romania:ti,ab OR Romania:ti,ab OR Russia:ti,ab OR Russia:ti,ab OR Russia:ti,ab OR Russia:ti,ab OR Russia:ti,ab OR Russia:ti,ab OR "Saint Vincent":ti,ab OR "Saint Lucia":ti,ab OR "St Lucia":ti,ab OR "Saint Vincent":ti,ab OR "Samoa:ti,ab OR "Samoa:ti,ab OR "Samoa:ti,ab OR "Samoan Islands":ti,ab OR "Saudi Arabia":ti,ab OR "Navigator Islands":ti,ab OR Samoa:ti,ab OR Serbia:ti,ab OR Montenegro:ti,ab OR Seychelles:ti,ab OR Senegal:ti,ab OR Serbia:ti,ab OR Montenegro:ti,ab OR Ceylon:ti,ab OR "Sierra Leone":ti,ab OR Sonalia:ti,ab OR Sudan:ti,ab OR Tajikistan:ti,ab OR Tadzhikistan:ti,ab OR Tadzhikistan:ti,ab OR Tadzhikistan:ti,ab OR Toogo:ti,ab OR "Togolese Republic":ti,ab OR Tonga:ti,ab OR Trunkae:ti,ab OR Turkey:ti,ab OR Turkey:ti,ab OR Toogo:ti,ab OR Turkey:ti,ab

SCOPUS

Community health worker	Program	мсн	LMIC
"Health Auxiliary" OR "community health worker*" OR "community health aide*" OR "village health worker*" OR "barefoot doctor*" OR "family planning personnel*" OR "health extension worker*" OR "lady health worker*" OR "community health agent*" OR "Shasthyo Sebika*" OR "community nutrition worker*" OR "maternal health worker*" OR "voluntary Malaria worker*" OR "village malaria worker*" OR "wollage malaria worker*" OR "mental health worker*" OR "mental health worker*" OR "mental health worker*" OR "mother coordinator*" OR "rural health worker*" OR "compagnateur* OR "Saksham Sahaya*" OR "anganwandi worker*" OR "accredited social health activist*" OR "community-based worker*" OR "community-based worker*" OR "community health volunteer*" OR "village health guide*" OR "maternal and child health promotion worker*" OR "maternal child health worker*" OR "colaborador voluntario*" OR behvarz* OR "village health helper*" OR "colaborador voluntario*" OR "nutrition volunteers*" OR "village drug-kit manager*" OR brigadistas* OR "female community health volunteer*" OR "Agente Comunitario de Salud*" OR "nutrition worker*" OR "community health worker*" OR "Genmunity health volunteer*" OR "Agente Comunitario de Salud*" OR "nutrition worker*" OR "community drug distributor*" OR "community	Program OR programs OR programme OR programmes OR initiative* OR project OR projects	"Maternal child health care"/de OR "Maternal Welfare" OR "child care" OR "child welfare" OR "maternal-child health services" OR "child health services" OR "maternal child health" OR "maternal newborn child health"	Afghanistan OR Albania OR Algeria OR Angola OR Antigua OR Barbuda OR Argentina OR Armenia OR Armenian OR Aruba OR Azerbaijan OR Bahrain OR Bangladesh OR Barbados OR Benin OR Byelarus OR Byelorussian OR Belarus OR Belorussian OR Belorussian OR Belorussian OR Belorussian OR Belorussian OR Belorussian OR Bosnia OR Herzegovina OR Hercegovina OR Botswana OR Brasil OR Brazil OR Bulgaria OR Burkina Faso OR "Burkina Fasso" OR "Upper Volta" OR Burundi OR Urundi OR Cambodia OR "Khmer Republic" OR Kampuchea OR Cameroon OR Cameroons OR Cameron OR Camerons OR "Cape Verde" OR "Central African Republic" OR Chad OR Chile OR China OR Colombia OR Comoros OR "Comoro Islands" OR Comores OR Mayotte OR Congo OR Zaire OR "Costa Rica" OR "Cote d'Ivoire" OR "Ivory Coast" OR Croatia OR Cuba OR Cyprus OR Czechoslovakia OR "Czech Republic" OR Slovakia OR "Slovak Republic" OR Djibouti OR "French Somaliland" OR Dominica OR "Dominican Republic" OR "East Timor" OR "East Timur" OR "Timor Leste" OR Ecuador OR Egypt OR "United Arab Republic" OR "El Salvador" OR Eritrea OR Estonia OR Ethiopia OR Fiji OR Gabon OR "Gabonese Republic" OR Gambia OR Gaza OR "Georgia Republic" OR "Georgian Republic" OR Ghana OR Gold Coast OR Greece OR Grenada OR Guatemala OR Guinea OR Guam OR Guiana OR Guyana OR Haiti OR Honduras OR Hungary OR India OR Maldives OR Indonesia OR Iran OR Iraq OR "Isle of Man" OR Jamaica OR Jordan OR Kazakhstan OR Kazakh OR Kenya OR Kiribati OR Korea OR Kosovo OR Kyrgyzstan OR Kirghizia OR "Kyrgyz Republic" OR Kirghiz OR Kirgizstan OR Lao PDR OR Laos OR Latvia OR Lebanon OR Lesotho OR Basutoland OR Liberia OR Libya OR Lithuania OR Macedonia OR Madagascar OR "Malagasy Republic" OR Malaysia OR Malaya OR Malay OR Sabah OR Sarawak OR Malayia OR Nyasaland OR Malay OR Sabah OR Sarawak OR Malawi OR Nyasaland OR Moldovia OR Moldovia OR Moldovia OR Mongolia OR Montenegro OR Morocco OR Ifni OR Mozambique OR Myanmar OR Myanma OR Burma OR Namibia

Community health worker	Program	МСН	LMIC
volunteer*" OR "community health advocate*" OR "lay health visitor*" OR "Promotoras de Salud"			OR Nepal OR "Netherlands Antilles" OR "New Caledonia" OR Nicaragua OR Niger OR Nigeria OR "Northern Mariana Islands" OR Oman OR Muscat OR Pakistan OR Palau OR Palestine OR Panama OR Paraguay OR Peru OR Phillippines OR Phillipines OR Phillippines OR Romania OR Rumania OR Roumania OR Russia OR Russian OR Rwanda OR Ruanda OR "Saint Kitts" OR St Kitts OR Nevis OR "Saint Lucia" OR "St Lucia" OR "Saint Vincent" OR "St Vincent" OR Grenadines OR Samoa OR "Samoan Islands" OR "Navigator Island" OR "Navigator Islands" OR Sao Tome OR "Saudi Arabia" OR Senegal OR Serbia OR Montenegro OR Seychelles OR "Sierra Leone" OR Slovenia OR "Sri Lanka" OR Ceylon OR "Solomon Islands" OR Somalia OR Sudan OR Suriname OR Surinam OR Swaziland OR Syria OR Tajikistan OR Tadzhikistan OR Tadzhik OR Tanzania OR Thailand OR Togo OR "Togolese Republic" OR Tonga OR Trinidad OR Tobago OR Tunisia OR Turkey OR Turkmenistan OR Turkmen OR Uganda OR Ukraine OR Uruguay OR USSR OR "Soviet Union" OR "Union of Soviet Socialist Republics" OR Uzbekistan OR Uzbek OR Vanuatu OR "New Hebrides" OR Venezuela OR Vietnam OR Viet Nam OR West Bank OR Yemen OR Yugoslavia OR Zambia OR Zimbabwe OR Rhodesia OR "Developing Country" OR Africa OR Asia OR Caribbean OR "West Indies" OR "South America" OR "Latin America" OR "Central America" OR "Developing Countr"

Appendix II: Data Charting Form

Scoping Review Title: Application of Primary Health Care Principles in National Community Health Worker Programs in Low- and Middle –Income Countries?	
Data charted by:	9.4 20 4444
Date of data charting:	
Study Details and Characteri	etice
Study citation details (author, year, title, journal, volume, issue, pages)	Sucs
Country of origin	
Study objective / aim	
Type of Study	Qualitative / Quantitative
Methods:	
Study Setting:	
CHW Program Details	
Name of the CHW Program	
Objective of the CHW Program	
Year the program started	
End date	
Implemented by	
Funded by	
Details / Results charted from the Study (in relation to the concept of the scoping	
which PHC principle is reflected in the reported objective of the national program?	 Universal access / Equity Community participation Intersectoral collaboration Appropriateness
How are they implementing the PHC principle (s)?	- Appropriateriess
Stated outcome / achievement of the CHW program with reference to PHC principle (s)	
Key findings of the article	
Characteristics of CHWs	
Key role of CHWs stated	
Nomenclature of CHWs	
Gender	
Employment status	
Pre-service training	
Catchment area	
Additional notes:	