BMJ Open Investigating the effect of an online self-compassion for weight management (SC4WM) intervention on self-compassion, eating behaviour, physical activity and body weight in adults seeking to manage weight: protocol for a randomised controlled trial

Jennifer M Brenton-Peters ¹, Nathan S Consedine ¹, Alana Cavadino ², Rajshri Roy ³, Anna Sofia Serlachius ¹

To cite: Brenton-Peters JM, Consedine NS, Cavadino A, *et al.* Investigating the effect of an online self-compassion for weight management (SC4WM) intervention on self-compassion, eating behaviour, physical activity and body weight in adults seeking to manage weight: protocol for a randomised controlled trial. *BMJ Open* 2022;**12**:e056174. doi:10.1136/bmjopen-2021-056174

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-056174>).

Received 05 August 2021
Accepted 21 December 2021

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Jennifer M Brenton-Peters;
jbre092@aucklanduni.ac.nz

ABSTRACT

Introduction Individual weight management, defined as engaging in behaviours to maintain or lose weight, can improve health and well-being. However, numerous factors influence weight management outcomes, such as genetics, biology, stress, the social and physical environment. Consequently, weight management can be hard. Self-compassion, described as treating oneself kindly in times of failure or distress, has shown promise in improving weight management outcomes. The objectives of this study are twofold: (1) to examine the efficacy of an online self-compassion for weight management (SC4WM) intervention coupled with an online commercial weight management programme (WW Weight Watchers reimagined) with increasing self-compassion and improving weight management outcomes (eating behaviour, physical activity and body weight) in comparison with the WW programme only and (2) to explore whether improvements in weight management outcomes are moderated by eating restraint, weight self-stigma, perceived stress and psychological coping.

Methods and analysis To achieve these objectives, 240 participants seeking to manage their weight were randomised to either an online behavioural commercial weight management programme (WW) or the online WW + SC4 WM intervention. Validated measures of self-compassion, stress, weight self-stigma, eating restraint, psychological coping and weight management outcomes were administered online at baseline, 4 weeks and at a 12-week follow-up.

Ethics and dissemination Ethics has been granted by the University of Auckland Health Research Ethics committee. Results will be communicated in peer-review journals, conferences and a doctoral thesis. If effective in increasing self-compassion and improving weight management outcomes, the intervention could be made more widely available to supplement behavioural weight management programmes.

Strengths and limitations of this study

- This is the first study to test a 100% online, self-compassion for weight management intervention (SC4WM) designed to supplement existing weight management programmes.
- The study uses a randomised controlled trial design with the online version of a commercial behavioural weight management programme as an active control (WW Weight Watchers reimagined).
- This study is limited due to the non-blinding of the first author and self-reporting of outcomes.
- This study investigates a supplemental SC4WM intervention that could increase the efficacy of online behavioural weight management programme by improving self-compassion and weight management outcomes during the global pandemic and beyond.

Trial registration number ACTRN12621000580875; Pre-results.

INTRODUCTION

Successful weight management, defined as losing or maintaining body weight, can improve health outcomes.¹ However, several competing factors influence weight management behaviours (eg, healthy eating and physical activity), including body physiology, biology, psychological well-being, stress management, socioeconomic status and the physical and social environment.^{2 3} Furthermore, ongoing public health restrictions to control the spread of COVID-19 and concerns about exposure mean access to in-person support may be limited and engaging in healthy behaviours can be even

more challenging (eg, accessibility to healthy food, more limited opportunity for exercise and financial stress).^{4 5} Self-compassion, the ability or tendency to respond to oneself with care and kindness in times of suffering and distress,⁶ has shown promise with improving weight management outcomes (healthier eating and reduced or maintained body weight).⁷ However, few studies have investigated whether the changes thought to accompany the development of self-compassion can be successfully taught and practised 100% online as a supplement to an existing online weight management programme.

Healthy eating and physical activity are recommended for everyone to improve health outcomes and manage weight.⁸ For those living with weight-related chronic disease, such as obesity, even a small amount of weight loss (5%–10% of body weight) can have positive physiological effects. For example, modest weight loss can lower blood lipids and improve glycaemic control, resulting in reduced risk for heart disease and diabetes complications.⁹ Appropriate weight management can also support psychological outcomes, including improved emotional well-being, self-esteem, depressive symptoms, body image and health-related quality of life.¹⁰ Behavioural-based weight management interventions have shown some effectiveness with weight loss and maintenance.^{11 12} However, weight management history (eg, previous unsuccessful weight loss attempts and weight cycling) and early weight loss results (eg, in the first 4 weeks) can influence an individual's weight management success.^{13 14} Weight regain is common due to physiological changes and when individuals cannot sustain the behavioural changes needed to maintain weight loss, such as healthy eating, increased physical activity and continued self-monitoring.^{15 16}

Engaging in weight management can be challenging. Body weight is influenced by several factors (eg, biology, stress and environments).² However, many people believe their behaviours *cause* their weight.¹⁷ Therefore, the inevitable ups and downs of body weight may result in feelings of distress, shame, failure and blame.^{18–20} These feelings are exacerbated by weight stigma, which is the negative societal prejudice and stereotypes experienced by people with higher weights.^{21 22} Facing weight stigma is common for adults who are engaging in weight management.²³ When internalised, weight self-stigma can create additional stress, ultimately resulting in less healthy eating and reduced physical activity.^{22 24}

Furthermore, weight fluctuations are associated with stress, regardless of health behaviour.²⁵ The stress response activates the sympathetic nervous system, which can cause a release in hormones (eg, cortisol) and reduce executive function (eg, less mindful regulation of food intake).^{26 27} Stress can trigger binge eating or the consumption of higher fat and higher sugar comfort foods, which may lead to weight gain over time.^{28 29} Eating restraint, defined as engaging in a cognitive effort to reduce caloric intake, can support weight management outcomes.^{30 31} However, those who are more restricted

with their eating behaviours may paradoxically be more prone to overeating in stressful times.³²

Self-compassion has shown the potential to moderate many challenges that plague weight management, including moderating the effects of stress.^{33 34} Self-compassionate people treat themselves with kindness when experiencing stressful situations and appear less likely to use avoidance as a coping mechanism.³⁵ Thus, self-compassion potentially offers a way to cope with stress through the enhanced ability to regulate health-related behaviour such as healthier eating and physical activity.^{36–38} In addition, people with greater self-compassion tend to have lower levels of internalised weight stigma^{39 40} and improved emotional regulation.^{36 41} Consequently, self-compassion could benefit weight management due to its ability to increase an individual's coping and self-management during stressful times.^{39 42 43}

Both mindfulness and self-compassion have been explored in the context of weight management (see recent reviews).^{7 44–47} Mindfulness is typically operationalised as self-regulation of attention, combined with curiosity, openness and acceptance of thoughts and feelings⁴⁸ and is a part of many self-compassion interventions (eg, mindful self-compassion).⁶ Relative to a 'pure' mindfulness intervention, however, self-compassion may be more relevant to the challenges of weight management than mindfulness alone.^{49 50} Specifically, because self-compassion reflects a way of relating to the self in times of failure and distress, it appears directly relevant to effectively managing the inevitable mistakes (eg, diet lapses),⁵¹ challenges (eg, weight stigma)⁵² and stress (eg, physiological and emotional regulation)^{36 53 54} inherent to weight management efforts. Self-compassion interventions encourage participants to develop a more self-compassionate way of thinking (eg, through self-compassionate diaries⁵⁰ or self-compassion meditations),⁵⁵ both self-compassion-specific interventions and self-compassion as part of broader interventions (eg, adding self-compassion to behavioural weight management) increase participant self-compassion in a weight management contexts.⁴⁷

Enhancing self-compassion in the context of weight management is complex.⁵⁶ For example, it is not uncommon when first practising self-compassion to feel some discomfort (eg, people may feel they do not *deserve* kindness).⁵⁷ Individuals engaging in weight management may be particularly vulnerable to feelings of discomfort due to internalised weight bias (negative attitudes associated with higher larger body size).⁵⁸ However, emerging research suggests that greater self-compassion can buffer the effects of weight self-stigma for individuals living with overweight or obesity.⁵² Similarly, individuals may worry that *self-kindness* will undermine their motivation for losing weight.⁵⁷ Self-kindness could easily be interpreted as indulging in a favourite food after a stressful day, as opposed to taking the time to make a healthy meal.⁵⁹ Egan and Mantzios⁵⁹ qualitatively explored perceptions of self-compassion and kindness in weight management and found that people struggled to relate to the

term 'self-kindness', preferring the concept of *caring for their mind*. Thus, while self-compassion does include an element of self-kindness,⁶ it also involves a recognition of experiences as part of common human experiences: *'everyone struggles with weight management at times'*. As well as being mindful of negative thoughts and feelings rather than overidentifying with them: *'I had a diet lapse, but that does not mean I am bad'*.⁶ Viewed in this way, practising self-compassion has the potential to *boost* motivation for self-improvement in face of failures⁶⁰ and promote health behaviours.⁶¹

Although promising, there are still notable gaps in our knowledge of self-compassion in weight management contexts. First, most studies include self-compassion as part of broader interventions (eg, nutrition goal setting, yoga). Second, since this is an emerging area of research, most studies lack robust comparison conditions such as behavioural weight management interventions. Consequently, it is hard to determine if it is self-compassion per se that is contributing to positive outcomes rather than some non-specific element of participation.⁴⁷

Also unclear is how to best deliver such interventions. In-person weight management interventions are often associated with high attrition rates⁶² and unpredictable public health measures to control the spread of COVID-19 may increase attrition and compound the challenges to engaging in health behaviours.^{4 63} Online interventions using cognitive-behavioural therapies have shown good effect sizes with improving health behaviours ($\bar{g}=0.43$, 95% CI 0.27 to 0.59).⁶⁴ Early evidence suggests that online weight management interventions show promise with supporting weight management outcomes, including online compassion-based interventions.^{65–68}

To this point, however, research has not contrasted the efficacy of an online self-compassion intervention tailored to weight management outcomes with a 100% online version of a behavioural weight management programme, which could provide increased scalability.⁴⁷ Therefore, the objective of this study is to assess whether an online, mobile-friendly self-compassion for weight management (SC4WM) intervention can increase participant self-compassion and improve weight management outcomes for those engaging in an online behavioural weight management programme (WW).

METHODS: PARTICIPANTS, INTERVENTIONS AND OUTCOMES

Following Consolidated Standards of Reporting Trials (CONSORT)⁶⁹ and Standard Protocol Items: Recommendations for Interventional Trials⁷⁰ guidelines, a two-armed, randomised controlled study comparing the addition of the SC4WM intervention to a widely available behavioural weight management programme (WW) will be conducted. The proposed study will investigate if the online SC4WM intervention can increase self-reported self-compassion and improve weight management outcomes, defined as eating behaviour, physical activity levels and body weight, for those in the first 4 weeks of

a widely available online commercial programme (WW). In addition, this study will examine potential moderating variables, including perceived stress, weight self-stigma, eating restraint and psychological coping. This study will be delivered 100% online and undertaken over 4 weeks, with a subsequent 12-week follow-up. The randomised controlled trial (RCT) protocol has been prospectively registered with the Australian New Zealand Clinical Trials Registry ACTRN12621000580875 (<https://www.anzctr.org.au/ACTRN12621000580875.aspx>). Protocol amendments will be communicated through the trial registry.

Participants

Participants were New Zealand residents who were seeking to manage their weight. Inclusion criteria for participation included: (1) being an adult (≥ 18 years of age), (2) New Zealand resident and (3) seeking to manage their weight. Exclusion criteria for participation included: (1) being a current WW member, (2) pregnant or expecting to become pregnant in the next 3 months, (3) diagnosed with an active eating disorder (eg, bulimia nervosa or anorexia nervosa) and (4) being prescribed medication for weight management or newly starting a medication that may cause weight gain. All participants were given an access code to a widely available behavioural commercial weight management programme (WW). Participants were randomised into either the WW programme only (active control group) or the WW plus the supplemental SC4WM intervention. Recruitment commenced on 17 June 2021 and finished on 11 October 2021. Participants were able to withdraw at any time during the study. Data analysis is expected to begin in January 2022.

Participant timeline

Participants were recruited via posters, websites (eg, university research site) and national (New Zealand) social media posts and marketing (eg, via community board posts on Facebook and Twitter) with a link to a secure website (REDCap).^{71 72} Potential participants clicked a link to the study eligibility questionnaire. If eligible to participate (eg, not a current WW member and ≥ 18 years of age), participants read and downloaded the Participant Information Sheet. After providing informed consent, participants completed the baseline questionnaires and were randomised using a computer-generated number sequence into either the online WW programme plus the SC4WM intervention or the online WW programme only (active control group). Please see [figure 1](#) for the CONSORT diagram demonstrating the flow of participants through the study.

The intervention group filled out the baseline questionnaires and received an access code and instructions to download the WW app and a separate link to the SC4WM website with a guest code. During the first 4-week period, participants in the intervention group received one automated email at the beginning of each week (via REDCap), reminding them to complete the SC4WM online modules. The control group received an access

Figure 1 CONSORT diagram. The flow of participants through the trial. CONSORT, Consolidated Standards of Reporting Trials; SC4WM, self-compassion for weight management.

code and instructions to download the WW app. They were instructed to use the WW programme as directed and expect subsequent emails at 4 weeks and 12 weeks with follow-up questionnaires.

REDCap sent automated emails to both groups at 4 weeks from baseline (post-intervention) and 12 weeks from baseline (follow-up). After completing the final follow-up questionnaire, both groups were provided with the opportunity to enter a draw for an iPad (see figure 1 CONSORT diagram).

Interventions

Active control group

Participants randomised to the control group partook in the online WW *Weight Watchers reimagined* program (2021). The WW programme is a commercial behavioural weight management programme that aims to support its

members with healthier habits.⁷³ The WW programme uses the Smartpoints food tracking system, which encourages members to make healthier food choices by assigning point values to foods requiring moderation (eg, higher fat and higher sugar foods). In addition, the WW programme promotes exercising, cultivating a positive mindset and tracking body weight with the objective of weight management. The most recent WW programme is the myWW+ programme, which includes a personalised meal plan, tracking and peer-support online. The myWW+ programme provides access to the myWW+ app that offers easy-to-use food and activity tracking, 24/7 live coaching and a supportive network of other WW members through a WW online forum.⁷³ The myWW+ programme may contain elements of self-compassion within its content. However, the SC4WM intervention

Figure 2 SC4WM landing page. SC4WM, self-compassion for weight management.

described further was designed to have a higher, more specific and more concentrated dose of self-compassion.

SC4WM intervention

The SC4WM intervention was designed for the Aotearoa New Zealand population and included consultation with Māori, Indigenous peoples of Aotearoa New Zealand. The SC4WM intervention was delivered online through a mobile-friendly website separate from the WW programme. Participants in the intervention group were also asked to follow the digital myWW+ programme as directed by WW. Participants required data or internet connectivity and an access code to log into the WW and SC4WM website content. The SC4WM intervention incorporated simple, evidence-based techniques (eg, construal journaling,⁵⁰ letter writing and reflections⁵⁵) to deliver a self-compassion intervention tailored to weight management outcomes (ie, eating behaviour, physical activity and weight monitoring). The SC4WM intervention landing page (website) provided an initial definition and background on self-compassion and quick access to each module (see [figure 2 SC4WM landing page](#)). The SC4WM intervention was based on taking the meaning of compassion inward, being mindful of personal suffering or distress

and applying the principles of mindfulness, self-kindness and shared humanity.⁷⁴ Given that previous research has found that people may feel uneasy with the term self-compassion,⁵⁹ care was taken to frame self-compassion as a whole construct, including increasing mindfulness (as opposed to overidentification), common humanity (tempering isolation) and self-kindness as a way to care for the body and mind (attenuating self-judgement).^{6 59}

The SC4WM intervention has four modules designed to specifically target participants' relationship with each weight management outcome, including self-compassion for eating behaviour, self-compassion for physical activity and self-compassion for body weight. Each module includes an adapted construal journal,⁵⁰ meditation and reflection activity⁵⁵ incorporating the principles of self-compassion to eating behaviour, physical activity behaviour and body weight. It was recommended that participants complete one module in sequence per week. The final module unifies all the concepts and encourages the participant to make a plan to include self-compassion in the future. To reduce any discomfort, participants were urged to start slow (eg, not to start with their biggest struggle right away) (see [table 1](#)).

Outcome measures

Completion of the questionnaires at baseline, postintervention (4 weeks) and follow-up (12 weeks) took participants approximately 25 min. REDCap collected demographic data via self-report (eg, sex, age, height, ethnicity, income, pre-existing weight-related chronic diseases and weight loss history, for example, number of previous weight loss attempts). Validated questionnaires were used to assess self-compassion, weight management outcomes and potential moderators at baseline, 4 weeks and 12 weeks after baseline. Since self-compassion is most useful when faced with a struggle or failure, participants reported on the degree of their struggle with weight loss or maintenance at baseline, 4-week and the 12-week follow-up. Adherence to both the WW programme and the SC4WM intervention was self-reported at 4-week and

Table 1 SC4WM modules

Module	Objective
Module 1: eat.	Cultivate a more self-compassionate attitude towards eating behaviours. Includes journaling, meditation and reflection activities designed to create mindfulness of eating behaviour, a feeling of shared humanity and self-kindness to the challenges of eating well.
Module 2: move.	Develop a more self-compassionate attitude towards physical activity behaviours. Includes journaling, meditation and reflection activities designed to create a mindful awareness of physical activity, a feeling of connection and self-kindness to the challenges of engaging in physical activity.
Module 3: weigh.	Foster a more self-compassionate attitude towards body weight. Includes journaling, meditation and reflection activities designed to create a mindful reaction to body weight, an awareness that they are not alone in their weight struggles, and strategies to bring self-kindness to the scale.
Module 4: unify.	Cultivate a more self-compassionate attitude towards weight management as a whole. Includes the selection of activities for the participant to continue with after the intervention.

SC4WM, self-compassion for weight management.

the 12-week follow-up. This study used validated measures that are referenced and described further below.

Primary outcome

To confirm that the intervention increased participant self-compassion compared with the WW programme, we will use the Self-Compassion Scale (SCS⁷⁵) at baseline, and at 4-week and 12-week follow-up. The SCS is a well-validated scale with high internal validity, Cronbach's α between 0.76 and 0.94 in previous studies.⁷⁶ The SCS indexes how participants typically respond towards themselves in times of failure or distress.^{6,77} Using a scale of 1=almost never to 5=almost always, participants are asked questions related to six subscales: self-kindness: '*I am loving towards myself when I am feeling emotional pain*'; self-judgement: '*I am disapproving of my own flaws and inadequacies*'; common humanity: '*When things are going badly for me, I see the difficulties as part of life that everyone goes through*'; isolation: '*When I think about my inadequacies, it tends to make me feel more separate and cut off from the world*'; mindfulness: '*When something upsets me I try to keep my emotions in balance*'; and overidentification: '*When I am feeling down I tend to obsess and fixate on everything that's wrong*'. The scale can be analysed both in terms of specific subscales (ie, self-kindness, mindfulness, common humanity, isolation, overidentification and self-judgement) as well as generating a total SCS score.⁷⁵ The SCS total score at 4 weeks (postintervention) will be used as the primary outcome for this study. The SCS subscales will be analysed to provide a better understanding how self-compassion may support weight management outcomes (eg, increased mindfulness, common humanity or reduced self-judgement, overidentification or isolation).⁷⁷

Secondary outcomes

The Weight Control Strategies Scale (WCSS) will be used to measure changes in participants' eating behaviour and physical activity behaviour.⁷⁸ The WCSS has subscales to measure weight control strategies commonly used in weight management programmes, including dietary choices, physical activity and self-monitoring. The WCSS is a well-validated scale with a Cronbach's $\alpha > 0.79$ and as high as 0.94 in a WW population.³⁰ The WCSS has also been modified to fit the WW programme specifically (eg, low-calorie foods have been changed to low point foods).³⁰ The WCSS asks participants on a scale of 0=never and 4=always to describe the strategies and behaviours they engage in when trying to lose or maintain weight loss over the last month. For example, '*I had several servings of fruits and/or vegetables each day*' and '*I engaged in a moderate-intensity exercise like brisk walking or something similar to brisk walking for at least 30 minutes a day*'. Scale items are totalled and divided by the number of items (total mean score). Separate subscale scores are achieved by summing all items in a subscale and dividing by the number of subscale items (total mean subscale score).⁷⁸

The impact of the SC4WM intervention on body weight outcomes will be determined by self-reported body weight

and height to calculate body mass index (kg/m^2). Self-reported body weight has been seen to strongly correlate with objectively measured body weight.^{30,79}

The overall objective of weight management is to support both physical and psychological health.⁸ Therefore, potential improvements in emotional well-being will be assessed using the WHO-5 Well-being Scale.⁸⁰ The WHO-5 is a brief well-validated scale with good internal reliability ($\alpha=0.84$). Participants will be asked how they have been feeling over the last 2 weeks, with a higher number indicating better emotional well-being. For example, '*I have felt cheerful and in good spirits*' is rated on a scale of 0–5 with 0 meaning 'at no time' and 5 'all the time'. A total score is calculated by summing all items.

Potential moderators

To assess participant eating restraint, the Revised Rigid Restraint Scale (RRRS)⁸¹ will be used. The RRRS is a validated 12-item scale designed to assess participants' thoughts, feelings and behaviours related to restrictive and guilty eating behaviours. The RRRS has good internal reliability with a Cronbach's α of >0.80 in previous studies.⁸¹ Participants are asked to rate how often statements describe their thoughts, feelings or behaviour regarding eating using on a five-item scale, 1 being ('never') to 5 ('always'). For example, '*There are certain unhealthy foods that I try not to eat in any quantity*'. All items are summed to assess dispositional eating restraint. A higher RRRS total score is indicative of higher eating restraint.⁸¹

The Weight Self-Stigma Questionnaire (WSSQ)⁸² is included in this study to measure the extent to which individuals have internalised weight self-stigma. The WSSQ is a 12-item scale used to measure weight self-stigma on a scale of 1 (completely disagree) to 5 (completely agree). For example, '*I don't have enough self-control to maintain a healthy weight*'. All items are summed to calculate a total score, or two subscales, self-devaluation and fear of enacted stigma. The WSSQ was designed to help evaluate if interventions reduce weight self-stigma.⁸² The WSSQ is correlated with the Weight Bias Internalisation Scale and predicts quality of life and self-esteem.⁸³ The WSSQ has good internal reliability with a Cronbach's α of 0.81 in a similar study.⁸⁴

The Perceived Stress Scale (PSS)⁸⁵ will be used to measure stress over the last month. The PSS is a 14-item scale that asks participants to rate feelings and thoughts over the last month related to coping with change, ability to handle personal problems and control irritations on a scale of 0 ('never') to 4 ('very often'). For example, '*In the last month, how often have you felt nervous or "stressed"?*'. The PSS total score is obtained by summing all 14 items. A higher score indicates higher perceived stress in the past month. The PSS is a well-validated scale with high internal reliability Cronbach's alpha of 0.88 in a previous study.⁸⁵

To assess psychological coping, the 28-item brief COPE⁸⁶ will be used. The brief COPE is the most frequently used coping scale; thus, it is a well-validated measure with a median Cronbach's alpha of 0.68 and a range of

0.55–0.92.⁸⁷ The brief COPE has 14 subscales that can be used to evaluate the frequency with which coping strategies are used (eg, self-distraction, active coping, denial, emotional support and behavioural disengagement). Participants use a scale of 1=I have not been doing this at all to 4=I have been doing this a lot. Each subscale item is added together. A subscale of the WCSS will also be used to assess psychological coping. Based on a scale of 0=never and 4=always, participants will be asked to describe how often they engaged in psychological coping strategies in the past month. For example, 'If I regained weight, I thought about my past successes and reminded myself that I could get back on track'.

Adherence

Participants in both groups were asked how often they used the WW app or digital online programme 'How many days did you use the WW program?' at 4 weeks and 12 weeks from baseline. Participants in the intervention group were asked how many modules of the SC4WM intervention they completed and how many days they used the SC4WM activities postintervention (4 weeks) and at follow-up (12 weeks from baseline).

METHODS: DATA COLLECTION, MANAGEMENT AND ANALYSIS

All participant data (eg, demographic data and weight management outcomes) was collected using validated questionnaires through REDCap. No data were collected through or by WW. Participant names only appeared on the online consent form; the remaining data were automatically coded with an alphanumeric participant identification number using REDCap. Potentially identifying information will be stored securely and separately from questionnaire data. Only the research team (JB-P, NC, RR, AS and AC) will have access to participant data.

Hypotheses

It is expected that the SC4WM intervention group will report greater increases in self-compassion, improved eating behaviours (eg, increased fruit and vegetable consumption), physical activity (eg, increased physical activity), body weight outcomes (eg, weight loss or less weight regained) and emotional well-being at the 4-week follow-up compared with the control group and that these improvements will persist at the 12-week follow-up. It is expected greater perceived stress (PSS), eating restraint (RRRS) and weight self-stigma (WSSQ), and decreased psychological coping (WCSS subscale) at baseline will moderate the relationship between the SC4WM intervention and changes in self-compassion and weight management outcomes at 4 and 12 weeks.

Sample size

In order to calculate the required sample size to examine group differences in self-compassion, we used an effect size from a compassion-based intervention study in a population struggling with weight, which found a small

to medium effect size for self-compassion (Cohen's $d=0.38$).⁸⁸ Using GPower,⁸⁹ it was calculated that to detect a small to medium change in self-compassion, using an independent samples t-test (with an alpha of 0.05 and power of 0.90), 240 participants would be required (120 per treatment arm).

Data analysis plan

Using SPSS software version 27, data will be assessed for violations of the parametric assumptions. Generalised linear mixed models will be used to explore within and between-group differences in changes in the primary (SCS total) and secondary outcomes (eating behaviour, physical activity and body weight) at the 4-week follow-up, with random intercepts for participant to account for repeated measures. Primary and secondary outcomes will also be evaluated at a 12-week follow-up. Possible moderators of the relationship between the SC4WM intervention and changes in self-compassion (eg, greater weight self-stigma WSSQ at baseline) and weight management outcomes at 4 and 12 weeks will be explored by adding an interaction term (eg, WSSQ*group*time) to the linear mixed models.

Baseline differences between completers and non-completers in the intervention and control groups will be evaluated using independent sample t-tests. The analyses will be based on both intention-to-treat and complier average causal effect (CACE) analytic methods.⁹⁰ CACE is preferred to per-protocol for this study as it recognises that different individuals may have different intervention needs (eg., differing intervention doses that may be effective for each individual participant).⁹⁰ CACE modelling considers intervention outcomes (μ_i) and proportion of the intervention participants completed (π_i) in four cells defined by treatment use. The mean at each time point for the intervention and control group will be the average of $\mu_1=\pi_1\mu_{c1}+\pi_2\mu_{n1}+\pi_3\mu_{a1}+\pi_4\mu_{d1}$.⁹⁰ For example, mean self-compassion (SCS total) at each time point (eg, 4 weeks from baseline and 12 weeks from baseline) for participants in the SC4WM intervention will be determined by grouping participants based on usage of the SC4WM intervention (eg, 1=completers, 2=half-completers, 3=minimal completers and 4=non-completers based on reported completion of modules of the SC4WM intervention). The self-compassion means for those in the control group will be determined by grouping participants based on usage of the WW programme (eg, 1=completers, 2=half-completers, 3=minimal completers and 4=non-completers based on reported usage of the WW programme). Therefore, using CACE, the mean self-compassion score at each time point (eg, post at 4 weeks from baseline and follow-up, 12 weeks from baseline) for the intervention and control group will be the average of $\mu_1=\pi_1\mu_{c1}+\pi_2\mu_{n2}+\pi_3\mu_{a3}+\pi_4\mu_{d4}$.⁹⁰

Participant open-ended comments on their experience at 4 and 12 weeks will be analysed using directed content analysis.⁹¹

DISCUSSION

This study is designed to investigate whether an SC4WM intervention can improve self-compassion and weight management outcomes for those engaging in a digital, widely available behavioural commercial weight management programme and begin the process of identifying potential moderators of self-compassion efficacy. The findings will be used to refine the SC4WM website and disseminate the SC4WM intervention to a broader audience.

Limitations to this study are expected to be similar to other online weight management interventions and may include low adherence to the intervention and high attrition rates.⁶⁵ We have tried to minimise attrition and improve adherence by keeping the intervention brief (4 weeks), engaging and easy to access.^{64 65} Maintenance of weight management outcomes (eg, weight loss) is a well-established challenge, even with self-compassion interventions.^{47 49} Integrating continued practice of self-compassion for weight management was part of the last module of the SC4WM intervention, with a follow-up at 12 weeks to determine if between group differences in self-compassion and weight management outcomes are maintained. Future research should test the effectiveness of self-compassion based interventions for relapse prevention and weight management in the longer term (6 months+). Finally, other possible limitations are the non-blinding of the first author and participants and relying on self-reported outcomes. Future research avenues include incorporating more objective measures of the intervention usage (eg, Google Analytics), body weight and physiological effects (eg, cortisol and heart rate variability) obtained by physicians or health professionals.

There is an urgent need to increase the scalability of weight management interventions as well as develop flexibility in delivery during the global pandemic and beyond. Self-compassion shows promise in improving weight management outcomes, including healthier eating, physical activity and body weight and could be a useful supplement to behavioural weight management interventions such as WW. However, more research is required to investigate if online self-compassion interventions tailored to weight management outcomes can increase self-reported self-compassion and improve outcomes, especially during the first 4 weeks of engaging in weight management.

It is acknowledged that this intervention uses a fee-based commercial weight management programme that may not be accessible to all. However, testing a widely available commercial weight management programme can support future public health initiatives and enhance current knowledge. For example, an RCT design using a robust behavioural comparator fills a gap in the literature, specifically testing whether self-compassion can enhance weight management outcomes over and above current weight management programmes. Furthermore, if found to be effective, the scalable online SC4WM programme could be made more broadly available to augment weight management programmes in the community or

to increase the efficacy of current public health initiatives targeting weight outcomes.

Author affiliations

¹Department of Psychological Medicine, The University of Auckland, Auckland, New Zealand

²Section of Epidemiology and Biostatistics, Faculty of Medical and Health Sciences, University of Auckland, Auckland, New Zealand

³Department of Nutrition and Dietetics, The University of Auckland, Auckland, New Zealand

Twitter Jennifer M Brenton-Peters @jennbrenton

Acknowledgements The authors would like to acknowledge Dr Chad Rethorst, in his role as WW Director, Clinical Research, Dr Michelle Cardel WW Director, Global Research and Nutrition, and Dr Gary Foster, Chief Scientific Officer WW International, Inc for providing the WW access codes. The authors would also like to thank Kate Wallace-Boyd and David Lim for recording meditations for the SC4WM intervention and Sarah Snow for NZ Māori consultation.

Contributors JB-P: conception and development of the self-compassion for weight management (SC4WM) intervention and study design and wrote the first draft of the protocol manuscript. AS (primary supervisor) and NC (cosupervisor): Supervision of the conception and development of the SC4WM intervention and study design, critical revision of the manuscript, and final approval of the version to be published. AC development and revision of the statistical analysis plan, critical revision of the manuscript and final approval of the version to be published. RR support with study design, critical revision of the manuscript and final approval of the version to be published. All authors read and approved the manuscript for publication.

Funding WW International, Inc. will provide participants 12-week free access to the WW digital program and WW app.

Competing interests This study is supported by WW International, Inc. WW International, Inc. was not involved in the intervention design and will not have access to individual participant data related to this study. Furthermore, WW International, Inc. will not play any role during the analyses, interpretation of the data, or the decision to submit results.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Not applicable.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Jennifer M Brenton-Peters <http://orcid.org/0000-0002-2370-900X>

Nathan S Consedine <http://orcid.org/0000-0002-7691-0938>

Alana Cavadino <http://orcid.org/0000-0002-5709-367X>

Rajshri Roy <http://orcid.org/0000-0002-7939-1790>

Anna Sofia Serlachius <http://orcid.org/0000-0002-4797-8351>

REFERENCES

- 1 Guh DP, Zhang W, Bansback N, *et al*. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. *BMC Public Health* 2009;9:88.
- 2 Froom S, Johnston LM, Matteson CL, *et al*. Obesity, complexity, and the role of the health system. *Curr Obes Rep* 2013;2:320–6.
- 3 Kirk SFL, Penney TL, McHugh T-LF. Characterizing the obesogenic environment: the state of the evidence with directions for future research. *Obes Rev* 2010;11:109–17.
- 4 Bhutani S, Cooper JA. COVID-19-Related home confinement in adults: weight gain risks and opportunities. *Obesity* 2020;28:1576–7.
- 5 Pellegrini CA, Webster J, Hahn KR, *et al*. Relationship between stress and weight management behaviors during the COVID-19 pandemic

- among those enrolled in an Internet program. *Obes Sci Pract* 2021;7:129–34.
- 6 Neff K. Self-Compassion: an alternative conceptualization of a healthy attitude toward Oneself. *Self and Identity* 2003;2:85–101.
 - 7 Rahimi-Ardabili H, Reynolds R, Vartanian LR, et al. A systematic review of the efficacy of interventions that aim to increase self-compassion on nutrition habits, eating behaviours, body weight and body image. *Mindfulness* 2018;9:388–400.
 - 8 Wharton S, Lau DCW, Vallis M, et al. Obesity in adults: a clinical practice guideline. *CMAJ* 2020;192:E875–91.
 - 9 Wing RR, Lang W, Wadden TA, et al. Benefits of modest weight loss in improving cardiovascular risk factors in overweight and obese individuals with type 2 diabetes. *Diabetes Care* 2011;34:1481–6.
 - 10 Lasikiewicz N, Myrissa K, Hoyland A, et al. Psychological benefits of weight loss following behavioural and/or dietary weight loss interventions. A systematic research review. *Appetite* 2014;72:123–37.
 - 11 Anderson JW, Konz EC, Frederich RC. Long-Term weight-loss maintenance: a meta-analysis of US studies. *Am J Clin Nutr* 2001;74:579–84.
 - 12 Hartmann-Boyce J, Johns DJ, Jebb SA, et al. Effect of behavioural techniques and delivery mode on effectiveness of weight management: systematic review, meta-analysis and meta-regression. *Obes Rev* 2014;15:598–609.
 - 13 Teixeira PJ, Goings SB, Houtkooper LB, et al. Pretreatment predictors of attrition and successful weight management in women. *Int J Obes Relat Metab Disord* 2004;28:1124–33.
 - 14 Hadziabdic MO, Mucalo I, Hrabac P, et al. Factors predictive of drop-out and weight loss success in weight management of obese patients. *J Hum Nutr Diet* 2015;28:24–32.
 - 15 Paixão C, Dias CM, Jorge R, et al. Successful weight loss maintenance: a systematic review of weight control registries. *Obes Rev* 2020;21:e13003.
 - 16 Greenway FL. Physiological adaptations to weight loss and factors favouring weight regain. *Int J Obes* 2015;39:1188–96.
 - 17 Wang C, Coups EJ. Causal beliefs about obesity and associated health behaviors: results from a population-based survey. *Int J Behav Nutr Phys Act* 2010;7:19.
 - 18 Kirk SFL, Price SL, Penney TL, et al. Blame, shame, and lack of support: a multilevel study on obesity management. *Qual Health Res* 2014;24:790–800.
 - 19 Rand K, Vallis M, Aston M, et al. "It is not the diet; it is the mental part we need help with." A multilevel analysis of psychological, emotional, and social well-being in obesity. *Int J Qual Stud Health Well-being* 2017;12:1306421.
 - 20 Hartmann-Boyce J, Boylan A-M, Jebb SA, et al. Experiences of self-monitoring in Self-Directed weight loss and weight loss maintenance: systematic review of qualitative studies. *Qual Health Res* 2019;29:124–34.
 - 21 Pearl RL, Puhl RM. Weight bias internalization and health: a systematic review. *Obes Rev* 2018;19:1141–63.
 - 22 Tomiyama AJ. Weight stigma is stressful. A review of evidence for the cyclic Obesity/Weight-Based stigma model. *Appetite* 2014;82:8–15.
 - 23 Puhl RM, Lessard LM, Pearl RL. International comparisons of weight stigma: addressing a void in the field. *Int J Obes* 2021;45:1976–85.
 - 24 O'Brien KS, Latner JD, Puhl RM. The relationship between weight stigma and eating behavior is explained by weight bias internalization and psychological distress. *Appetite* 2016;102:70–6.
 - 25 Serlachius A, Hamer M, Wardle J. Stress and weight change in university students in the United Kingdom. *Physiol Behav* 2007;92:548–53.
 - 26 Dallman MF. Stress-Induced obesity and the emotional nervous system. *Trends Endocrinol Metab* 2010;21:159–65.
 - 27 Yau YHC, Potenza MN. Stress and eating behaviors. *Minerva Endocrinol* 2013;38:255–67.
 - 28 Born JM, Lemmens SGT, Rutters F, et al. Acute stress and food-related reward activation in the brain during food choice during eating in the absence of hunger. *Int J Obes* 2010;34:172–81.
 - 29 Zellner DA, Loaiza S, Gonzalez Z, et al. Food selection changes under stress. *Physiol Behav* 2006;87:789–93.
 - 30 Phelan S, Halfman T, Pinto AM, et al. Behavioral and psychological strategies of Long-Term weight loss Maintainers in a widely available weight management program. *Obesity* 2020;28:421–8.
 - 31 Johnson F, Pratt M, Wardle J. Dietary restraint and self-regulation in eating behavior. *Int J Obes* 2012;36:665–74.
 - 32 Wallis DJ, Hetherington MM. Stress and eating: the effects of ego-threat and cognitive demand on food intake in restrained and emotional eaters. *Appetite* 2004;43:39–46.
 - 33 Rockliff H, Gilbert P, McEwan K. A pilot exploration of heart rate variability and salivary cortisol responses to compassion-focused imagery. *Clinical Neuropsychiatry* 2008;5:132–9.
 - 34 Arch JJ, Brown KW, Dean DJ, et al. Self-compassion training modulates alpha-amylase, heart rate variability, and subjective responses to social evaluative threat in women. *Psychoneuroendocrinology* 2014;42:49–58.
 - 35 Allen AB, Leary MR. Self-Compassion, stress, and coping. *Soc Personal Psychol Compass* 2010;4:107–18.
 - 36 Sirois FM, Kitner R, Hirsch JK. Self-compassion, affect, and health-promoting behaviors. *Health Psychol* 2015;34:661–9.
 - 37 Terry ML, Leary MR. Self-compassion, self-regulation, and health. *Self and Identity* 2011;10:352–62.
 - 38 Ferrari M, Hunt C, Harrysunker A. Self-Compassion interventions and psychosocial outcomes: a meta-analysis of RCTs. *Mindfulness* 2019;10:1455–73.
 - 39 Palmeira L, Cunha M, Pinto-Gouveia J. Processes of change in quality of life, weight self-stigma, body mass index and emotional eating after an acceptance-, mindfulness- and compassion-based group intervention (Kg-Free) for women with overweight and obesity. *J Health Psychol* 2019;24:1056–69.
 - 40 Fekete EM, Herndier RE, Sander AC. Self-Compassion, internalized weight stigma, psychological well-being, and eating behaviors in women. *Mindfulness* 2021;12:1262–71.
 - 41 Kirschner H, Kuyken W, Wright K, et al. Soothing your heart and feeling connected: a new experimental paradigm to study the benefits of Self-Compassion. *Clin Psychol Sci* 2019;7:545–65.
 - 42 Forbes Y, Donovan C. The role of internalised weight stigma and self-compassion in the psychological well-being of overweight and obese women. *Aust Psychol* 2019;54:471–82.
 - 43 Ferreira C, Pinto-Gouveia J, Duarte C. Self-compassion in the face of shame and body image dissatisfaction: implications for eating disorders. *Eat Behav* 2013;14:207–10.
 - 44 Carrière K, Khoury B, Günak MM, et al. Mindfulness-based interventions for weight loss: a systematic review and meta-analysis. *Obes Rev* 2018;19:164–77.
 - 45 Dunn C, Haubenreiser M, Johnson M, et al. Mindfulness approaches and weight loss, weight maintenance, and weight regain. *Curr Obes Rep* 2018;7:37–49.
 - 46 Mantzios M, Wilson JC, Mindfulness WJ. Mindfulness, eating behaviours, and obesity: a review and reflection on current findings. *Curr Obes Rep* 2015;4:141–6.
 - 47 Brenton-Peters J, Consedine NS, Boggiss A, et al. Self-compassion in weight management: a systematic review. *J Psychosom Res* 2021;150:110617.
 - 48 Bishop SR, Lau M, Shapiro S, et al. Mindfulness: a proposed operational definition. *Clinical Psychology: Science and Practice* 2004;11:230–41.
 - 49 Mantzios M, Wilson JC. Exploring mindfulness and mindfulness with self-compassion-centered interventions to assist weight loss: theoretical considerations and preliminary results of a randomized pilot study. *Mindfulness* 2015;6:824–35.
 - 50 Mantzios M, Wilson JC. Making concrete construals mindful: a novel approach for developing mindfulness and self-compassion to assist weight loss. *Psychol Health* 2014;29:422–41.
 - 51 Thøgersen-Ntoumani C, Dodos LA, Stenling A, et al. Does self-compassion help to deal with dietary lapses among overweight and obese adults who pursue weight-loss goals? *Br J Health Psychol* 2021;26:767–88.
 - 52 Hilbert A, Braehler E, Schmidt R, et al. Self-Compassion as a resource in the Self-Stigma process of overweight and obese individuals. *Obes Facts* 2015;8:293–301.
 - 53 Luo X, Qiao L, Che X. Self-compassion modulates heart rate variability and negative affect to experimentally induced stress. *Mindfulness* 2018;9:1522–8.
 - 54 Ferrari M, Hunt C, Harrysunker A, et al. Self-compassion interventions and psychosocial outcomes: a meta-analysis of RCTs. *Mindfulness* 2019;10:1455–73.
 - 55 Neff KD, Germer CK. A pilot study and randomized controlled trial of the mindful self-compassion program. *J Clin Psychol* 2013;69:28–44.
 - 56 Mantzios M, Egan HH. On the role of self-compassion and self-kindness in weight regulation and health behavior change. *Front Psychol* 2017;8:229.
 - 57 Germer CK, Neff KD. Self-compassion in clinical practice. *J Clin Psychol* 2013;69:856–67.
 - 58 Carels RA, Wott CB, Young KM, et al. Implicit, explicit, and internalized weight bias and psychosocial maladjustment among treatment-seeking adults. *Eat Behav* 2010;11:180–5.
 - 59 Egan H, Mantzios M. A qualitative exploration of self-kindness and "treating oneself" in contexts of eating, weight regulation and other health behaviors: Implications for mindfulness-based eating programs. *Front Psychol* 2018;9:880.
 - 60 Breines JG, Chen S. Self-compassion increases self-improvement motivation. *Pers Soc Psychol Bull* 2012;38:1133–43.

- 61 Phillips WJ, Hine DW. Self-compassion, physical health, and health behaviour: a meta-analysis. *Health Psychol Rev* 2021;15:113–39.
- 62 Miller BM, Brennan L. Measuring and reporting attrition from obesity treatment programs: A call to action! *Obes Res Clin Pract* 2015;9:187–202.
- 63 Yeo C, Ahmed S, Oo AM, *et al.* COVID-19 and Obesity-the management of pre- and Post-bariatric patients amidst the COVID-19 pandemic. *Obes Surg* 2020;30:3607–9.
- 64 van Genugten L, Dusseldorp E, Webb TL, *et al.* Which combinations of techniques and modes of delivery in Internet-based interventions effectively change health behavior? A meta-analysis. *J Med Internet Res* 2016;18:e155.
- 65 van Beurden SB, Simmons SI, Tang JCH, *et al.* Informing the development of online weight management interventions: a qualitative investigation of primary care patient perceptions. *BMC Obes* 2018;5:7.
- 66 Caverio-Redondo I, Martinez-Vizcaino V, Fernandez-Rodriguez R, *et al.* Effect of behavioral weight management interventions using lifestyle mHealth self-monitoring on weight loss: a systematic review and meta-analysis. *Nutrients* 2020;12. doi:10.3390/nu12071977. [Epub ahead of print: 03 Jul 2020].
- 67 Duarte C, Gilbert P, Stalker C, *et al.* Effect of adding a compassion-focused intervention on emotion, eating and weight outcomes in a commercial weight management programme. *J Health Psychol* 2019;26:1700–15. doi:10.1177/1359105319890019
- 68 Schnepfer R, Reichenberger J, Blechert J. Being My Own Companion in Times of Social Isolation - A 14-Day Mobile Self-Compassion Intervention Improves Stress Levels and Eating Behavior. *Front Psychol* 2020;11:595806.
- 69 Schulz KF, Altman DG, Moher D. Statement: updated guidelines for reporting parallel group randomized trials. *Annals of Internal Medicine* 2010;2010:726–32.
- 70 Chan A-W, Tetzlaff JM, Altman DG, *et al.* Spirit 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;158:200–7.
- 71 Harris PA, Taylor R, Minor BL, *et al.* The REDCap Consortium: building an international community of software platform partners. *J Biomed Inform* 2019;95:103208.
- 72 Harris PA, Taylor R, Thielke R, *et al.* Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.
- 73 Ww. introducing myWW+. WW weight Watchers reimaged, 2020. Available: <https://www.weightwatchers.com/nz/ourprogram> [Accessed November 30, 2020].
- 74 Goetz JL, Keltner D, Simon-Thomas E. Compassion: an evolutionary analysis and empirical review. *Psychol Bull* 2010;136:351–74.
- 75 Neff KD. The development and validation of a scale to measure Self-Compassion. *Self and Identity* 2003;2:223–50.
- 76 Sirois FM. The association between self-compassion and self-rated health in 26 samples. *BMC Public Health* 2020;20:74.
- 77 Neff KD, Tóth-Király I, Yarnell LM, *et al.* Examining the factor structure of the Self-Compassion scale in 20 diverse samples: support for use of a total score and six subscale scores. *Psychol Assess* 2019;31:27–45.
- 78 Pinto AM, Fava JL, Raynor HA, *et al.* Development and validation of the weight control strategies scale. *Obesity* 2013;21:2429–36.
- 79 McGuire MT, Wing RR, Klem ML, *et al.* What predicts weight regain in a group of successful weight losers? *J Consult Clin Psychol* 1999;67:177–85.
- 80 Bech P, Olsen LR, Kjoller M, *et al.* Measuring well-being rather than the absence of distress symptoms: a comparison of the SF-36 mental health subscale and the WHO-Five well-being scale. *Int J Methods Psychiatr Res* 2003;12:85–91.
- 81 Adams CE, Leary MR. Promoting Self-Compassionate attitudes toward eating among restrictive and guilty Eaters. *J Soc Clin Psychol* 2007;26:1120–44.
- 82 Lillis J, Luoma JB, Levin ME. Measuring weight Self-stigma: the weight Self-stigma questionnaire. *Obesity* 2012;18.
- 83 Durso LE, Latner JD. Understanding self-directed stigma: development of the weight bias internalization scale. *Obesity* 2008;16 Suppl 2:S80–6.
- 84 Levin ME, Potts S, Haeger J, *et al.* Delivering acceptance and commitment therapy for weight Self-Stigma through guided self-help: results from an open pilot trial. *Cogn Behav Pract* 2018;25:87–104.
- 85 Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;24:385–96.
- 86 Carver CS. You want to measure coping but your protocol's too long: consider the brief cope. *Int J Behav Med* 1997;4:92–100.
- 87 Kato T. Frequently used coping scales: a meta-analysis. *Stress Health* 2015;31:315–23.
- 88 Palmeira L, Pinto-Gouveia J, Cunha M. Exploring the efficacy of an acceptance, mindfulness & compassionate-based group intervention for women struggling with their weight (Kg-Free): A randomized controlled trial. *Appetite* 2017;112:107–16.
- 89 Faul F, Erdfelder E, Lang A-G, *et al.* G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007;39:175–91.
- 90 Connell AM. Employing complier average causal effect analytic methods to examine effects of randomized encouragement trials. *Am J Drug Alcohol Abuse* 2009;35:253–9.
- 91 Hsieh H-F, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res* 2005;15:1277–88.