BMJ Open Definitions and measurement of health literacy in health and medicine research: a systematic review

Kristin Hjorthaug Urstad ^{1,2}, Marit Helen Andersen,^{3,4} Marie Hamilton Larsen,^{5,6} Christine Råheim Borge,^{3,7} Sølvi Helseth,^{8,9} Astrid Klopstad Wahl^{3,4}

To cite: Urstad KH, Andersen MH, Larsen MH, *et al.* Definitions and measurement of health literacy in health and medicine research: a systematic review. *BMJ Open* 2022;**12**:e056294. doi:10.1136/bmjopen-2021-056294

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-056294>).

Received 11 August 2021
Accepted 15 December 2021

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Kristin Hjorthaug Urstad;
kristin.h.urstad@uis.no

ABSTRACT

Objectives The way health literacy is understood (conceptualised) should be closely linked to how it is measured (operationalised). This study aimed to gain insights into how health literacy is defined and measured in current health literacy research and to examine the relationship between health literacy definitions and instruments.

Design Systematic review in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement.

Data sources The MEDLINE, PsycINFO, ERIC and CINAHL databases were searched for articles published during two randomly selected months (March and October) in 2019.

Eligibility criteria We included articles with a quantitative design that measured health literacy, were peer-reviewed and original, were published in the English language and included a study population older than 16 years.

Data extraction and synthesis Six researchers screened the articles for eligibility and extracted the data independently. All health literacy definitions and instruments were considered in relation to category 1 (describing basic reading and writing skills, disease-specific knowledge and practical skills) and category 2 (social health literacy competence and the ability to interpret and critically assess health information). The categories were inspired by Nutbeam's descriptions of the different health literacy levels.

Results 120 articles were included in the review: 60 within public health and 60 within clinical health. The majority of the articles (n=77) used instruments from category 1. In total, 79 of the studies provided a health literacy definition; of these, 71 were in category 2 and 8 were in category 1. In almost half of the studies (n=38), health literacy was defined in a broad perspective (category 2) but measured with a more narrow focus (category 1).

Conclusion Due to the high degree of inconsistency between health literacy definitions and instruments in current health literacy research, there is a risk of missing important information about health literacy considered to be important to the initial understanding of the concept recognised in the studies.

PROSPERO registration number CRD42020179699.

INTRODUCTION

Health literacy is usually understood as cognitive and social skills that determine the motivation and ability to understand and use

Strengths and limitations of this study

- To our knowledge, this is the first systematic review to investigate connections between health literacy definitions and instruments used in current health literacy research.
- The health literacy definitions and instruments were categorised based on a health literacy model described by Nutbeam.
- The initial plan to assess health literacy definitions and instruments for three categories was changed to two due to difficulties in distinguishing between two of the categories.
- Searches were limited to two randomly selected months.

health information,¹ and adequate health literacy is seen as a prerequisite for healthy behaviours. Researchers have increasingly worked to identify challenges associated with health literacy and investigate the role they play in an individual's ability to comprehend self-care information and its relationship to health outcomes.² Empirical studies have reported that low health literacy is associated with poor health-related outcomes, such as high hospital admission rates,^{3 4} low participation in preventive activities,^{5–9} poor self-management of chronic conditions,¹⁰ poor disease outcomes and high mortality.^{11–13}

The concept of health literacy emerged in the 1970s when health education was viewed as social policy.¹⁴ A topic-specific query in the PubMed tools reveals a recent exponential growth of articles about health literacy, with 129 references between 1986 and 1990 increasing to more than 8000 in the past 5 years. Today, health literacy is seen as a global goal for enhancing health promotion through improved education and communication strategies to improve health outcomes.¹⁵

Health literacy is defined in numerous ways.^{14 16–18} In a systematic review by Sorensen

*et al.*¹⁴ 17 different definitions and 12 conceptual models were reported on health literacy. Another systematic review found that 34 research articles between 2010 and 2015 had an explicit objective to define the concept of health literacy.¹⁷ Moreover, the literature has reported the use and development of more than 150 health literacy instruments over the last decade.^{19–21} Traditionally, health literacy approaches have focused on individual skill deficits and health education based on the communication of factual information regarding health risks as well as how to use the health system,²² with the majority of health literacy research having used instruments measuring reading and numerical skills.²³ However, in recent years, more multidimensional perspectives and instruments measuring health literacy have been introduced, such as the Health Literacy Questionnaire²⁴ and the Health Literacy Survey European Questionnaire 47.²⁵

The current and sometimes confusing use of various and inconsistent interpretations of health literacy is a challenge for the development of valid and reliable measurements.¹⁶ In 2000, Nutbeam proposed a health literacy model that is now widely cited in the health literacy literature and is seen by many health literacy researchers as useful in analysing health literacy abilities required in various health situations.²² According to Nutbeam, health literacy contains three different levels, progressing from basic skills in reading and writing (functional health literacy), to the ability to derive meaning from different forms of communication and apply new information to changing situations (interactive health literacy) and to more advanced cognitive skills which, together with social skills, can be applied to critically analyse information and

to achieve policy and organisational changes (critical health literacy).

Different understandings of health literacy and different measurement tools may be useful as they complement each other and provide different perspectives. However, the way health literacy is understood (conceptualised) should be closely linked to how it is measured (operationalised) in each study context.²³ Nguyen *et al* has described this as a ‘conceptual stumbling block’ that needs to be resolved for the field to progress.²³ A first useful step for addressing this might be to systematically explore how it appears in current health literacy research. Hence, by performing a systematic review, our aim was to gain insights into how health literacy is defined and measured in current health literacy research. In particular, we will examine the relationship between health literacy definitions and instruments. This review may increase our understanding of potential conceptual and methodological challenges or gaps that need to be addressed in future research.

METHODS

This systematic review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement²⁶ and registered in PROSPERO (https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=179699). The review was designed with a time frame limited to two randomly selected months in 2019 (March and October). Due to a high number of health literacy articles published every year, random selection was chosen in order to reflect current health literacy

Table 1 Overview of study populations in clinical and public health studies

Clinical health studies		Public health studies	
Populations classified by diseases and related health problems	n	Type of study population	n
Diseases in the circulatory system	11	General population	19
Endocrine diseases	9	Students	13
Mental illness	6	Parents/caregivers	6
Cancer	6	Elderly people	5
Diseases in the respiratory system	4	Immigrants	3
Diseases in the urine and genital organs	4	Work-related populations	3
Infectious and parasite diseases	3	Young adults	2
Diseases in the ear	2	Veterans	2
Diseases in the musculoskeletal system and connective tissue	2	Health personnel	2
Diseases in the nervous system	1	Men	1
Diseases in the digestive system	1	Women	1
		Pet owners	1
Pregnancy, birth, postnatal period	1	Relatives of patients with cancer	1
Others: general chronic conditions (n=2), chronic pain (n=3), patients in primary care (n=2), next of kin (n=2), patients in specialist care (n=1)	10	Churchgoers	1
Total	60	Total	60

Figure 1 Flow diagram shows the study selection process. HL, health literacy.

Figure 2 Overview of study designs of included studies. RCT, randomised controlled trial.

Figure 3 Studies' continent of origin.

research. To ensure that March and October were not unique in terms of the number of articles published, we performed the same search strategy using the same databases for one other randomly selected month in 2019. This search yielded nearly the same number of articles.

Search strategy

Systematic literature searches were conducted in collaboration with a trained librarian (4 March 2020). The MEDLINE, PsycINFO, ERIC and CINAHL databases were searched for the term 'health literacy'. For all databases except for ERIC (where this was not possible), the search was automatically restricted to two randomly selected months in 2019: March and October. Citations in ERIC were manually assessed for articles published in March 2019 and October 2019.

The search yielded a total number of 1038 citations. Endnote V.X9 was used to manage the generated research articles. After removing duplicates, 951 citations

remained. All records not published for the first time in 2019 were removed, leaving 414 articles for screening (see online supplemental appendix 1 for the search history).

Selection criteria

The inclusion and exclusion criteria were developed a priori. The 414 published articles were distributed among six researchers (KHU, AKW, MHA, CRhB, SH and MHL), who worked in pairs. The articles were included if they fulfilled the following inclusion criteria: (1) 'health literacy' was mentioned in the title or abstract; (2) the article was peer reviewed; (3) the research was original; (4) it used a quantitative design; (5) it was published in the English language; and (6) the study population included individuals older than 16 years of age.

To reduce selection bias, the researchers independently screened the articles for eligibility according to the criteria. They then met in pairs to compare their results, resolve any conflicting opinions and decide whether to include each article. Conflicting opinions in pairs were presented and solved by the whole team.

Data extraction (selection and coding)

To achieve consistency in the data extracted from the included articles, an Excel V. 2019 spreadsheet was created. The initial question in this spreadsheet was whether health literacy was measured. If the answer was no, the article was excluded. The data extracted from the articles included information about the study design and context, such as country of origin and whether the study was conducted in a public health or clinical health setting. For clinical studies that included participants with health problems, the different types of diseases were categorised according to the International Statistical Classification of Diseases and Related Health Problems.²⁷ For the public health studies, we categorised type of study populations

Table 2 Most frequently used references for health literacy definitions in the included studies

Reference	Definition	Studies (n)
Category 1		
Jorm (1997) Jorm ³⁶	'Knowledge about appropriate treatment options; and <i>attitudes</i> that facilitate recognition and treatment-seeking' ' <i>Knowledge and beliefs</i> about mental disorders which aid their recognition, management or prevention'	5
American Medical Association (1999) ³⁹	'The <i>constellation of skills, including the ability</i> to perform basic reading and numerical tasks required to function in the healthcare environment'	1
Category 2		
Nutbeam ^{22 32–34}	'The <i>cognitive and social skills</i> which determine the motivation and ability of individuals to gain access to understand and use information in ways which promote and maintain good health.'	19
Sørensen et al ¹⁴	'Health literacy entails people's knowledge, motivation and competences to access, <i>understand, appraise, and apply</i> health information in order to make judgments and take decisions in everyday life concerning healthcare, disease prevention and health promotion to maintain or improve quality of life during the life course.'	14
The Institute of Medicine (2004) ⁴⁰	'The individuals' capacity to <i>obtain, process</i> and understand basic health information and services needed to make appropriate health decisions.'	11

Figure 4 Studies categorised with the same level of HL definitions and HL instruments. *Category 1 includes basic reading and writing skills, disease-specific knowledge and practical skill competences. **Category 2 includes skills to communicate and interact with healthcare providers and the ability to interpret and critically analyse health information. HL, health literacy.

. Furthermore, the data included information about whether the instruments were generic or disease specific, whether it was used in combination with other health literacy instruments or if a reference was provided for the instrument.

The published articles were analysed with respect to which health literacy definitions and instruments were used. Before the data extraction, a pilot assessment of three studies was performed to determine the feasibility of the data extraction tool. The initial plan was to assess health literacy definitions and instruments for three categories: (1) functional, (2) interactive and (3) critical health literacy, inspired by Nutbeam's description of different health literacy levels.²² As a result of the pilot testing, the number of categories was changed due to difficulties in distinguishing between the two latter levels. All health literacy definitions and health literacy instruments were therefore considered in relation to two categories: category 1 included definitions and instruments describing basic reading and writing skills, disease-specific knowledge and practical skill competencies needed to function in everyday situations; and category 2 included definitions and instruments that also described health literacy as skills to communicate and interact with healthcare providers as well as the ability to interpret and critically analyse health information (online supplemental appendix 2 illustrates the coding for the data extractions).

The researchers first extracted data and considered independently the definitions and instruments related to the two categories. They then met in pairs to compare their results and resolve any conflicting opinions. The remaining conflicting opinions within pairs were presented and solved by the whole team in a group meeting.

Patient and public involvement

No patient was involved.

RESULTS

The search yielded 1038 articles in total (see the online supplemental appendix 1 for the search history). After

removing duplicates, 951 articles remained. Of the 163 articles that were read in full text, 1 was a duplicate, and 39 were excluded because they did not measure health literacy. Ultimately, 120 articles were included (see figure 1).

Characteristics of included studies

Sixty articles described studies with a clinical health focus, and 60 were conducted in public health settings. The clinical studies included a great variety of diseases, but the most frequent diseases were those in the circulatory system (n=11) and endocrine diseases (n=9) followed by mental illness (n=6) and cancer diseases (n=6). In the public health studies, the most frequent study group was the general population (n=19), looking into aspects such as vaccination programmes, oral care and mental health. Furthermore, a major part of the public health studies included students (n=13), mainly represented by health professional students. Six studies focused on parents/caregivers, while five studies focused on the elderly population. (table 1)

The majority of the studies used a cross-sectional design (n=90). Nine studies used a randomised controlled design, and 10 studies were quasi-experimental. Four of the studies used a longitudinal design, while one was a case-control study. Figure 2 presents an overview of all study designs.

As figure 3 indicates, the majority of studies were conducted in North America (n=56) and Asia (n=31). The fewest studies were performed in Africa (n=4) and South America (n=3).

Health literacy definitions and instruments

Out of the 120 included studies, 88 used generic health literacy instruments, while 32 were context-specific. Eleven studies used a combination of health literacy instruments. A total of 77 studies used instruments from category 1 (describing basic reading and writing skills, disease-specific knowledge and practical skills), whereas 43 instruments were from category 2 (describing communication and interaction skills and/or the ability to interpret and critically analyse health information). The most frequently used health literacy instruments in category 1 were Newest Vital Sign²⁸ (n=19) and Test of Functional Health Literacy²⁹ (n=13). The most frequently used health literacy instruments in category 2 were The European Health Literacy Survey Questionnaire³⁰ and The eHealth Literacy Scale³¹ (n=10).

Of the 120 articles, 79 provided a health literacy definition. Of these, 46 were public health studies and 33 were clinical health studies. Only eight studies provided a definition addressed to category 1. Five out of these eight studies focused on mental health challenges.

The most frequently used reference in category 2 was Nutbeam.^{22 32 33 34} The most frequently used reference in category 1 was a definition provided by Jorm and colleagues.^{35 36} Table 2 gives an overview of the text in

Table 3 Studies categorised with mutual levels of health literacy definitions and instruments (n=41)

Study reference	Study context	Design	Health literacy definition reference	Category	Health literacy instrument	Category
Category 1						
Cehuen Neto <i>et al</i> 2019 ⁴¹	Clinical	Cross-sectional	American Medical Association Ad Hoc Committee on Health, 1999 ³⁹	1	Short Test of Functional Health Literacy in Adults (Parker <i>et al</i>) ²⁹	1
Clough <i>et al</i> 2019 ⁴²	Public health	RCT	Jorm <i>et al</i> ³⁵	1	Mental Health Literacy Scale (O'Connor and Casey, 2015) ⁴³	1
Clough <i>et al</i> 2019 ⁴⁴	Public health	Cross-sectional	Jorm <i>et al</i> ³⁵	1	Mental Health Literacy Scale (O'Connor and Casey, 2015) ⁴³	1
Joncho <i>et al</i> 2019 ⁴⁵	Clinical	Cross-sectional	Jorm <i>et al</i> ³⁵	1	Developed within the study context: Mental Health Literacy Questionnaire	1
Lichtveld <i>et al</i> 2019	Public health	Cross-sectional	Finn and O'Fallon, 2019 Gray, 2018 ⁴⁶	1	Developing and validating a questionnaire measuring environmental health literacy: measuring media-specific knowledge, attitudes, and behaviours	1
Nye <i>et al</i> 2019 ⁴⁷	Public health	Longitudinal	US Department of Health and Human Services, 2010 ⁴⁸	1	The Upper Peninsula Oral Health Assessment Survey	1
Waldmann <i>et al</i> 2020 ⁴⁹	Clinical	Cross-sectional	Jorm ³⁶	1	Mental Health Knowledge Schedule. Depression Literacy Scale (Wilson, 2015) ⁵⁰	1
Williston <i>et al</i> 2020 ⁵¹	Public health	Longitudinal	Jorm <i>et al</i> ³⁵	1	Mental Health Literacy Scale (O'Connor and Casey, 2015) ⁴³	1
Category 2						
An <i>et al</i> 2019 ⁵²	Public health	Cross-sectional	Nutbeam ²²	2	The Health Literacy Scale (Chew, 2004) ⁵³	2
Azizi <i>et al</i> 2019 ⁵⁴	Public health	Mixed method	Nutbeam ²²	2	Health Literacy Scale for Workers ⁵⁴	2
Brandstetter <i>et al</i> 2019 ⁵⁵	Public health	Cross-sectional	Sørensen <i>et al</i> ¹⁴	2	The European Health Literacy Survey (HLS-EU) (Sørensen <i>et al</i>) ²⁵	2
Degan <i>et al</i> 2019 ⁵⁶	Clinical	Cross-sectional	WHO, 1998 ¹	2	The Health Literacy Questionnaire (Osborn <i>et al</i>) ²⁴	2
Eo <i>et al</i> 2019 ⁵⁷	Public health	Cross-sectional	Nutbeam ²²	2	The Health Literacy Assessment Scale for Asian Immigrant Women (And and Yang, 2015) ⁵⁸	2
Ernsting <i>et al</i> 2019 ⁵⁹	Clinical	Cross-sectional	Sørensen <i>et al</i> ¹⁴	2	European Health Literacy Survey Questionnaire—short form (Sørensen <i>et al</i> 2013) and The eHealth Literacy Scale (Norman and Skinner) ³¹	2
Erunal <i>et al</i> 2019 ⁶⁰	Public health	RCT	Nutbeam ²²	2	Turkish Health Literacy Scale-32 (Okay <i>et al</i> 2016) ⁶¹	2

Continued

Table 3 Continued

Study reference	Study context	Design	Health literacy definition reference	Category	Health literacy instrument	Category
Evans <i>et al</i> 2019 ⁶²	Public health	Cross-sectional	Sørensen <i>et al</i> ¹⁴	2	European Health Literacy Survey Questionnaire (Sørensen <i>et al</i> ²⁵)	2
Fernandez Gutierrez <i>et al</i> 2019 ⁶³	Public health	Quasi-experimental	Nutbeam ²²	2	HLS EU-16 (Sørensen <i>et al</i> ²⁵) Health Literacy App Questionnaire+five practical tests	2
Goto <i>et al</i> 2019 ⁶⁴	Public health	Cross-sectional	Nutbeam, 1998 ³³	2	The Functional Communicative and Critical HL Tool (Ishikawa <i>et al</i> 2008)	2
Guclu <i>et al</i> 2019 ⁶⁵	Clinical	Cross-sectional	WHO, 1998	2	European Health Literacy Survey Questionnaire (Sørensen <i>et al</i> 2013) ²⁵	2
Güner <i>et al</i> 2019 ⁶⁶	Public health	Cross-sectional	Nutbeam, 2000 ²²	2	Questionnaire developed within the study context (consisting of 21 questions about HL knowledge, practice and attitudes)	2
Hu <i>et al</i> 2019 ⁶⁷	Public health	Cross-sectional	Nutbeam, 2000 ²²	2	Health Literacy Management Scale (Jordan, 2013) ⁶⁸	2
Indino <i>et al</i> 2019 ⁶⁹	Public health	Cross-sectional	Nutbeam, 2015 ³²	2	The Functional Communicative and Critical Health Literacy Tool (Ishikawa <i>et al</i> 2008)	2
Kaper <i>et al</i> , 2019 ⁷⁰	Public health	Quasi-experimental	Kwan, 2006 ⁷¹	2	Questionnaire developed within the study context by combining subscales from other questionnaires	2
Kobayashi <i>et al</i> 2019 ⁷²	Public health	Mixed method	Sørensen <i>et al</i> ¹⁴	2	European Health Literacy Survey Questionnaire (Sørensen <i>et al</i> ²⁵)	2
Lim <i>et al</i> 2019 ⁷³	Clinical	Cross-sectional	Osborne <i>et al</i> ²⁴	2	Health Literacy Questionnaire (Osborne <i>et al</i> ²⁴)	2
Logullo <i>et al</i> 2019 ⁷⁴	Public health	Cross-sectional	Sørensen <i>et al</i> ¹⁴	2	Short Assessment of Health Literacy for Portuguese Speaking Adults (Apolonario <i>et al</i> 2012) ⁷⁵	2
Lu <i>et al</i> 2020 ⁷⁶	Clinical	Cross-sectional	Sørensen, 2012 ¹⁴	2	HLS EU-16 (Sørensen <i>et al</i> 2013) ²⁵	2
Lu <i>et al</i> 2019 ⁷⁷	Clinical	Cross-sectional	Sørensen, 2012 ¹⁴	2	HLS EU-16 (Sørensen <i>et al</i> 2013) ²⁵	2
Ma <i>et al</i> 2019 ⁷⁸	Public health	Cross-sectional	Norman and Skinner ³¹	2	The eHealth Literacy Scale (Norman and Skinner ³¹)	2
Nokes <i>et al</i> 2019 ⁷⁹	Clinical	Quasi-experimental	Norman and Skinner ³¹ and 2011	2	The eHealth Literacy Scale (Norman and Skinner ³¹)	2
Oh <i>et al</i> 2019 ⁸⁰	Public health	Cross-sectional	WHO, 1998 ¹	2	The eHealth Literacy Scale (Norman and Skinner ³¹)	2
Oliffe <i>et al</i> 2020 ⁸¹	Public health	Cross-sectional	Peerson and Saunders, 2009, ⁸² O'Brian <i>et al</i> 2005	2	The Health Literacy Questionnaire (Osborn <i>et al</i> 2013) ²⁴	2

Continued

Table 3 Continued

Study reference	Study context	Design	Health literacy definition reference	Category	Health literacy instrument	Category
Pobhirun and Pinitsoontorn, 2019 ⁸³	Public health	Cross-sectional	Nutbeam, 1998 ³³	2	A questionnaire developed within the study context including six dimensions: cognitive skills, access, communication skills, self-management, media literacy and decision-making skills	2
Rababah ⁸⁴ et al 2019	Public health	Cross-sectional	WHO, 2013, Sørensen et al ¹⁴	2	The Health Literacy Questionnaire (Osborn et al 2013) ²⁴	2
Ruëgg and Abel, 2019 ⁸⁵	Public health	Cross-sectional	Selden et al 2000 ⁸⁶	2	Short survey tool for public health and health promotion research (Abel et al, 2014) ⁸⁷	2
Solhjoo et al 2019 ⁸⁸	Public health	Mixed method	Sørensen, 2012 ¹⁴	2	The eHealth Literacy Scale (Norman and Skinner ³¹)	2
Stømer et al 2019 ⁸⁹	Clinical	Cross-sectional	WHO, 1998 ¹	2	The Health Literacy Questionnaire ²⁴ (Osborn et al 2013)	2
Uysal et al 20 ⁹⁰ 20	Public health	Cross-sectional	Sørensen, 2012 ¹⁴	2	European Health Literacy Survey Questionnaire (Sørensen et al ²⁵)	2
Wang et al 2019 ⁹¹	Clinical	Cross-sectional	Kutner et al 2003 ⁹²	2	Chinese Health Literacy Scale for Diabetes (Leung, 2013) ⁹³	2
Wang et al 2019 ⁹⁴	Clinical	Cross-sectional	Nutbeam, 2009 ⁹⁵	2	Diabetes Health Literacy Scale (Lee et al, 2018) ⁹⁶	2
Zhang et al 20 ⁹⁷ 20	Clinical	Cross-sectional	Sørensen, 2012 ¹⁴	2	Heart Failure-specific Health Literacy Scale (Matsuka et al 2018) ⁹⁸	2

HLS EU, European Health Literacy Survey Questionnaire ; RCT, randomised controlled trial.

the most cited health literacy definitions. Words in italics are examples of key words considered important in the decision to add the definition to category 1 or category 2. For instance, definitions that described health literacy with terms such as ‘knowledge’, ‘skills’ and ‘attitude’ were referred to category 1, while definitions with terms like ‘appraise’ and ‘social skills’ were referred to category 2.

Regarding the connection between health literacy definitions and instruments in the 79 relevant studies, 41 articles used health literacy definitions and instruments characterised at the same level. Thirty-three of these were in category 2, and 8 were in category 1 (see [figure 4](#) and [table 3](#)).

In the remaining 38 studies, there was a disconnect between levels of health literacy definitions and instruments. In all of these, health literacy definitions were from category 2, and all instruments were in category 1 (see [table 4](#)).

DISCUSSION

This systematic review aimed to gain insights into how current research defines and measures health literacy and, in particular, whether studies consistently used definitions and instruments. The high number of articles published in the defined time frame shows that health literacy is of high research interest in both public health and the clinical field in large parts of the world.

We found a large variety of instruments used, and the majority of the included studies (79 out of 120) presented a health literacy definition as part of the study’s theoretical background. However, there seems to be an inconsistency between the definitions and the instruments in a significant number of the studies. In nearly half of the studies, health literacy was defined in a broad perspective (including aspects such as social health literacy competence and the ability to process and appraise health information) while using instruments with a more narrow focus (measuring basic skills and knowledge). As a result, almost half of the articles in our review lacked data on the participants’ ability to critically appraise health information and their social health literacy competence despite the fact that the authors had stated such aspects to be health literacy. This concern has previously been addressed. Numerous systematic reviews have reported on the diversity of understandings of health literacy and the various use of instruments not aligned to the definitions in current research.^{16–21} However, the current study is, to the best of our knowledge, the first review to systematically investigate the relationship between health literacy definitions and instruments from the perspective of a health literacy model.

The use of instruments that focused on functional health literacy (addressed to category 1), such as Newest Vital Sign and Test of Functional Health Literacy, were predominant in our review.^{28 29} This finding corresponds with previous findings.^{19 20} The widespread use of Newest Vital Sign and Test of Functional Health Literacy is

justified by the fact that they are screening tools—that is, they are quick, available in an ever-increasing number of languages and can be adapted to different settings.³⁷ However, a broad range of skills and tasks covering functional, interactive and critical domains needs to be included in measures of health literacy in order to capture the health literacy definitions used.³² In recent years, an increasing number of multidimensional instruments have been developed. In our review, The European Health Literacy Survey Questionnaire and The eHealth Literacy Scale, both addressed to category 2, were the multidimensional instruments most often used.^{30 31} In light of the increased focus on a more multidimensional perspective of health literacy, and the fact that the current review only includes studies from 2019, one could expect a higher use of multidimensional instruments. Instead, they represented only 43 out of 120 instruments.

Clearly, our study highlights the discrepancy between definitions and measurements as well as the narrow focus of health literacy in a large amount of empirical research. An important question to raise is therefore how the findings from this specific study can guide future research strategies to overcome the identified inconsistency. In other words, as most health literacy researchers seem to base their research on a broad understanding of health literacy, what can be done to facilitate an increased use of broad measurements? A first step should be to make researchers aware about the existing mismatch in current research. Furthermore, it seems necessary to develop more instruments that can answer the research questions posed. Despite the high number of instruments, there still seems to be a need for questionnaires in the field of health literacy that capture more multidimensional dimensions besides the functional aspects.

Nutbeam’s description of three levels of health literacy inspired the categorisation of definitions and instruments.²² Initially, we planned to distinguish among the three levels. However, the line between levels 2 and 3 was challenging to distinguish. Therefore, it was decided to merge the two latter categories. In a recent publication from 2020, Nutbeam has provided a more thorough description of the three levels with a more detailed explanation of how the levels should be understood.³⁸ Perhaps, these descriptions would have contributed to a clearer guidance in our work with the categorisation. However, this material was not available at the time of our work and, in general, this situation illustrates the challenge of adapting a theoretical model into practise.

The interpretation of terms used in the definitions and instruments that guided the choice of category also represented some challenges. These were resolved through discussions both in pairs and as a research team. For instance, definitions describing health literacy using terms like knowledge and ‘beliefs’ were included in category 1, while definitions using terms such as ‘appraise’ and ‘understand and process’ were included in category 2. An interesting finding is that, among the eight studies providing a health literacy definition from category 1, the

Table 4 Studies categorised with conflicting levels of health literacy definitions and instruments (n=38)

Study reference	Study context	Design	Health literacy definition reference	Category	Health literacy instrument	Category
Anderson <i>et al</i> 20 ⁹⁹	Clinical	Cross-sectional	Defined within the study context (no reference provided)	2	S-TOFHLA (Parker <i>et al</i> ²⁹)	1
Avci <i>et al</i> 2019 ¹⁰⁰	Public health	Cross-sectional	US Department of Health and Human Services, 2000 ⁴⁸	2	NVS (Weiss <i>et al</i> ²⁸) and REALM (Davis, 1993) ¹⁰¹	1
Bonaccorsi ¹⁰² <i>et al</i> 2019	Public health	Cross-sectional	Sørensen <i>et al</i> ¹⁴	2	NVS (Weiss <i>et al</i> ²⁸)	1
Bonaccorsi <i>et al</i> 2019 II ¹⁰³	Public health	Cross-sectional	Sørensen <i>et al</i> ¹⁴	2	NVS (Weiss <i>et al</i> ²⁸)	1
Carducci <i>et al</i> 2019 ¹⁰⁴	Public health	Cross-sectional	Ratzan and Parker, 2000 ²²	2	TOFHLA (Parker, 1995) ²⁹	1
Chen <i>et al</i> 2019 ¹⁰⁵	Clinical	Cross-sectional	Nutbeam, 2000 ²²	2	Health Literacy Scale for Diabetes (Lee <i>et al</i> 2016) ¹⁰⁶	1
de Melo <i>et al</i> 2019 ¹⁰⁷	Clinical	Cross-sectional	Cavanaugh, 2011 ¹⁰⁸	2	Short Test of Functional Health Literacy in Adults (Parker <i>et al</i> ²⁹)	1
Flynn <i>et al</i> 2019 ¹⁰⁹	Public health	Cross-sectional	American Dental Association Council, 20 ¹¹⁰ 09	2	Oral Health Literacy Adults Questionnaire (Sistani <i>et al</i> 2014) ¹¹¹	1
Gaikwad, 2019 ¹¹²	Public health	Cross-sectional	Ratzan and Parker, 2000 ¹¹³	2	Rapid Estimation of Adult Literacy in Dentistry 30-word version (Lee <i>et al</i> 2007) ¹¹⁴	1
Güner <i>et al</i> 2019 ¹¹⁵	Public health	Cross-sectional	Nutbeam ²²	2	Developed within the context of the study	1
Han <i>et al</i> 2019 ¹¹⁶	Public health	Cross-sectional	Sørensen <i>et al</i> ¹⁴ Nutbeam ²²	2	Assessment of Health Literacy in Cancer Screening (Han <i>et al</i> 2014) ¹¹⁷	1
Himes <i>et al</i> 2019 ¹¹⁸	Public health	Cross-sectional	Institute of Medicine, US, Committee HL, 2004 ⁴⁰	2	Chew <i>et al</i> 's (2004) set of brief questions ⁵³	1
Irvin <i>et al</i> 2019 ¹¹⁹	Public health	Cross-sectional	Peters <i>et al</i> 2012 ¹²⁰	2	The Water Environmental Literacy Level Scale (Irvin <i>et al</i> 2019) ¹¹⁹	1
Kaur <i>et al</i> 2019 ¹²¹	Public health	RCT	Healthy People, 2010 (Oral health) ¹²²	2	TS-REALD Two-Stage Rapid Estimate of Adult Literacy (Stucky <i>et al</i> 2011) ¹²³	1
Kim <i>et al</i> 2019 ¹²⁴	Public health	Cross-sectional	Ratzan and Parker, 2000 ¹¹³	2	Three questions, each addressing oral, listening, and written literacies (no reference)	1
Kim <i>et al</i> 2019 ¹²⁵	Public health	Quasi-experimental	Joint Committee on National Education Standards, 1995 ¹²⁶	2	Knowledge questions developed in the study context	1
Kim <i>et al</i> 2019 ¹²⁷	Clinical	RCT	Institute of Medicine, US, Committee HL, 2004 ⁴⁰	2	S-TOFHLA (Parker <i>et al</i> ²⁹)	1
Kino <i>et al</i> 2020 ¹²⁸	Public health	Cross-sectional	Inst of Medicine, 2004 ⁴⁰	2	Three indicators of HL (Haun <i>et al</i> ²⁰)	1

Continued

Table 4 Continued

Study reference	Study context	Design	Health literacy definition reference	Category	Health literacy instrument	Category
Lin <i>et al</i> 2019 ¹²⁹	Public health	Quasi-experimental	Nutbeam, 2000 ²²	2	Mandarin Health Literacy Scale (Lee <i>et al</i> 2011) ¹³⁰	1
Lindahl <i>et al</i> 2020 ¹³¹	Clinical	Cross-sectional	Nutbeam and Kickbusch, 2000 ¹³²	2	The three-item Brief Health Literacy Screen (Cavanaugh <i>et al</i> 2015) ¹³³	1
Mackert <i>et al</i> 2019 ¹³⁴	Public health	Cross-sectional	Berkman <i>et al</i> 2010 ¹³⁵	2	NVS (Weiss <i>et al</i> ²⁸)	1
Mayer <i>et al</i> 2019 ¹³⁶	Clinical	Cross-sectional	Nutbeam ³³	2	Developed within the context of the study	1
Meyers <i>et al</i> 2019 ¹³⁷	Clinical	Cross-sectional	Institute of Medicine, 2004 ⁴⁰	2	NVS (Weiss <i>et al</i> ²⁸)	1
Miranda, 2019 ¹³⁸	Clinical	Cross-sectional	Sørensen <i>et al</i> ¹⁴	2	REALM-D (Davis, 1993)	1
Mock <i>et al</i> 2019 ¹³⁹	Clinical	Cross-sectional	Institution of Medicine, 2004 ⁴⁰	2	Single Item Literacy Screening (Morris, 2006), S-TOFHLA (Parker <i>et al</i> ²⁹)	1
Mora-Pinzon <i>et al</i> 2019 ¹⁴⁰	Clinical	Cross-sectional	Institute of Medicine, 2004 ⁴⁰	2	S-TOFHLA (Parker, 1995) ²⁹ REALM-D (Davis, 1993) ¹⁰¹	1
Noback <i>et al</i> 2019 ¹⁴¹	Clinical	Cross-sectional	Paasche-Orlow <i>et al</i> 2007, ¹⁴² Berkman <i>et al</i> 2010 ¹³⁵ ,	2	NVS (Weiss <i>et al</i> 2005) ²⁸ Musculoskeletal specific literacy survey (LiMP questionnaire) developed within the study context	1
O'Connor <i>et al</i> 2019 ¹⁴³	Clinical	Cross-sectional	Institute of Medicine, 2004 ⁴⁰	2	S-TOFHLA (Parker <i>et al</i> ²⁹)	1
Penaloza <i>et al</i> 2019 ¹⁴⁴	Clinical	Cross-sectional	Nutbeam ³³ and Institute of Medicine, 2004 ⁴⁰	2	Short Assessment of Health Literacy Spanish (Lee <i>et al</i> 2006) ¹⁴⁵	1
Rafferty <i>et al</i> 2019 ¹⁴⁶	Clinical	Cross-sectional	Institute of Medicine, 2004 ⁴⁰ and Sørensen <i>et al</i> ¹⁴	2	Questions focusing on health advice and information-seeking skills, oral literacy, and print literacy	1
Tavakoly Sany <i>et al</i> 2019 ¹⁴⁷	Clinical	Quasi-experimental	US Health Resources and Services Administration	2	TOFHLA (Parker <i>et al</i> ²⁹)	1
Scrivner <i>et al</i> 2019 ¹⁴⁸	Clinical	Cross-sectional	Nutbeam, 2008 ³⁴	2	Three questions assessing health literacy (no reference)	1
Tucker <i>et al</i> 2019 ¹⁴⁹	Public health	Quasi-experimental	Ratzan and Parker, 2000 ¹¹³	2	NVS (Weiss <i>et al</i> 2005) ²⁸	1
Van Wormer <i>et al</i> 2019 ¹⁵⁰	Public health	Cross-sectional	Institute of Medicine, 2004 ⁴⁰	2	Oral Health Literacy in Adults Questionnaire (Sistani <i>et al</i> 2013) ¹⁵¹	1
Weaver <i>et al</i> 2019 ¹⁵²	Public health	Cross-sectional	Ratzan and Parker, 2000 ²²	2	REALM (Davis, 1993) ¹⁰¹	1

Continued

Table 4 Continued

Study reference	Study context	Design	Health literacy definition reference	Category	Health literacy instrument	Category
Zhong <i>et al</i> 2020 ¹⁵³	Clinical	Cross-sectional	Kindig, 2004 ¹⁵⁴	2	NVS (Weiss <i>et al</i> 2005) ²⁸	1
Williams <i>et al</i> 2019 ¹⁵⁵	Clinical	Cross-sectional	Institute of Medicine, 2004 ⁴⁰	2	NVS (Weiss <i>et al</i> 2005) ²⁸	1
Winokur <i>et al</i> 2019 ¹⁵⁶	Clinical	Cross-sectional	Morrison <i>et al</i> 2013 ¹⁵⁷	2	NVS (Weiss <i>et al</i> 2005) ²⁸	1

NVS, Newest Vital Sign; RCT, randomised controlled trial; REALM, Rapid Estimate of Adult Literacy in Medicine; REALM-D, Rapid Estimate of Adult Literacy, Dutch Version; S-TOFHLA, Short Test of Functional Health Literacy in Adults; TOFHLA, Test of Functional Health Literacy in Adults.

majority (n=5) were in the context of mental health. The predominant reference in these studies was the definition provided by Jorm and colleagues, who, unlike others, defined health literacy as being linked to beliefs and 'attitudes'.^{35 36} Whether these terms are more closely linked to mental health literacy challenges, compared with other more 'physical' health literacy issues, is not clear but would be interesting for further investigation.

Another aspect worth noting is that we discovered that many publications did not cite the primary source of the health literacy definitions but instead referred to secondary sources (other researchers presenting definitions of health literacy). Referring to the original sources should be the first choice and would perhaps make it easier for readers to recognise the definition's affiliation.

The present study has some limitations. First, this study was designed to analyse and describe health literacy research in two randomly selected months. This period may not be representative of health literacy research in general. However, a large number of health literacy studies are published every year. A random selection can therefore give a good picture of health literacy research. Second, we did not conduct a quality assessment screening of the included studies. This was considered less relevant for the current study as the aim of the current study was to explore connections between health literacy definitions and instruments rather than to assess methodology. Furthermore, searches were limited to the English language only. It is possible that similar studies may have been published in languages other than English.

The current review included only quantitative measurements. However, qualitative approaches might provide valuable and more in-depth insights into the field. For future research, it would be interesting to also explore how qualitative research links health literacy definitions to the research questions posed.

CONCLUSION

There seems to be an inconsistency between the definitions and the instruments used in a significant part of current health literacy research. This situation raises the risk of missing information about health literacy that was considered

to be important in the initial understanding of the concept recognised in the studies. This gap should be taken into consideration in future health literacy research. We hope our work contributes to making explicit where the problem might be rooted and that it can be useful in the discussion about strategies for moving forward to better align health literacy measurement with definitions of health literacy.

Author affiliations

¹Department of Quality and Health Technology, University of Stavanger, Stavanger, Norway

²Faculty of Health Sciences, VID Specialized University, Oslo, Akershus, Norway

³Department of Interdisciplinary Health Sciences, University of Oslo, Oslo, Norway

⁴Department of Transplantation Medicine, Oslo University Hospital Rikshospitalet, Oslo, Norway

⁵Department of Behavioural Sciences in Medicine, University of Oslo, Oslo, Norway

⁶Lovisenberg Diaconal University College, Oslo, Akershus, Norway

⁷Lovisenberg Diakonale Hospital, Oslo, Norway

⁸Faculty of Health Sciences, Oslo Metropolitan University, Oslo, Norway

⁹Department of Health and Nursing Sciences, University of Agder, Kristiansand, Norway

Contributors KHU, MHA and AKW initiated the project and wrote and revised the manuscript. KHU led the project. KHU, MAH, AKW, SH, CRB and MHL designed the study, selected the articles and extracted the data. All authors contributed to drafting the manuscript and met authorship criteria. KHU had full responsibility for the work and the conduct of the study, had access to the data, and controlled the decision to publish.

Funding The Centre for Advanced Study in Oslo, Norway, funded and hosted our research project, The Body in Translation: Historicising and Reinventing Medical Humanities and Knowledge Translation, during the 2019/2020 academic year.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study does not involve human participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which

permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID ID

Kristin Hjorthaug Urstad <http://orcid.org/0000-0002-8830-4564>

REFERENCES

- WHO. *Health promotion glossary*. Geneva: WHO, 1998.
- Visscher BB, Steunenberg B, Heijmans M, et al. Evidence on the effectiveness of health literacy interventions in the EU: a systematic review. *BMC Public Health* 2018;18:1414.
- Mitchell SE, Sadikova E, Jack BW, et al. Health literacy and 30-day postdischarge Hospital utilization. *J Health Commun* 2012;17:325–38.
- Baker DW, Gazmararian JA, Williams MV, et al. Functional health literacy and the risk of hospital admission among Medicare managed care enrollees. *Am J Public Health* 2002;92:1278–83.
- Adams RJ, Piantadosi C, Ettridge K, et al. Functional health literacy mediates the relationship between socio-economic status, perceptions and lifestyle behaviors related to cancer risk in an Australian population. *Patient Educ Couns* 2013;91:206–12.
- Thomson MD, Hoffman-Goetz L. Application of the health literacy framework to diet-related cancer prevention conversations of older immigrant women to Canada. *Health Promot Int* 2012;27:33–44.
- Jovanić M, Zdravković M, Stanisavljević D, et al. Exploring the importance of health literacy for the quality of life in patients with heart failure. *Int J Environ Res Public Health* 2018;15:1761.
- O'Hara J, McPhee C, Dodson S, et al. Barriers to breast cancer screening among diverse cultural groups in Melbourne, Australia. *Int J Environ Res Public Health* 2018;15:1677.
- Chen J-Z, Hsu H-C, Tung H-J, et al. Effects of health literacy to self-efficacy and preventive care utilization among older adults. *Geriatr Gerontol Int* 2013;13:70–6.
- Olesen K, F Reynders AL, Joensen L, et al. Higher health literacy is associated with better glycemic control in adults with type 1 diabetes: a cohort study among 1399 Danes. *BMJ Open Diabetes Res Care* 2017;5:e000437.
- Bostock S, Steptoe A. Association between low functional health literacy and mortality in older adults: longitudinal cohort study. *BMJ* 2012;344:e1602.
- Peterson PN, Shetterly SM, Clarke CL. Low health literacy is associated with increased risk of mortality in patients with heart failure. *Circulation* 2009;120:749.
- Peterson PN, Shetterly SM, Clarke CL, et al. Health literacy and outcomes among patients with heart failure. *JAMA* 2011;305:1695–701.
- Sørensen K, Van den Broucke S, Fullam J, et al. Health literacy and public health: a systematic review and integration of definitions and models. *BMC Public Health* 2012;12:80.
- Parker RM, Ratzan S. Re-enforce, not re-define health literacy—moving forward with health literacy 2.0. *J Health Commun* 2019;24:923–5.
- Malloy-Weir LJ, Charles C, Gafni A, et al. A review of health literacy: definitions, interpretations, and implications for policy initiatives. *J Public Health Policy* 2016;37:334–52.
- Liu C, Wang D, Liu C, et al. What is the meaning of health literacy? A systematic review and qualitative synthesis. *Fam Med Community Health* 2020;8:e000351.
- Parnell TA, Stichler JF, Barton AJ, et al. A concept analysis of health literacy. *Nurs Forum* 2019;54:315–27.
- O'Neill B, Gonçalves D, Ricci-Cabello I, et al. An overview of self-administered health literacy instruments. *PLoS One* 2014;9:e109110.
- Haun JN, Valerio MA, McCormack LA, et al. Health literacy measurement: an inventory and descriptive summary of 51 instruments. *J Health Commun* 2014;19:302–33.
- Nguyen TH, Park H, Han H-R, et al. State of the science of health literacy measures: validity implications for minority populations. *Patient Educ Couns* 2015;1512 doi:10.1016/j.pec.2015.07.013
- Nutbeam D. Health literacy as a public health goal: a challenge for contemporary health education and communication strategies into the 21st century. *Health Promot Int* 2000;15:259–67.
- Nguyen TH, Paasche-Orlow MK, McCormack LA. The state of the science of health literacy measurement. *Stud Health Technol Inform* 2019;240:189–203.
- Osborne RH, Batterham RW, Elsworth GR, et al. The grounded psychometric development and initial validation of the health literacy questionnaire (HLQ). *BMC Public Health* 2013;13:658.
- Sørensen K, Van den Broucke S, Pelikan JM, et al. Measuring health literacy in populations: illuminating the design and development process of the European health literacy survey questionnaire (HLS-EU-Q). *BMC Public Health* 2013;13:948.
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
- Cataloguing-in-Publication Data International statistical classification of diseases and related health problems. - 10th revision, edition 2010. 3 v.
- Weiss BD, Mays MZ, Martz W, et al. Quick assessment of literacy in primary care: the newest vital sign. *Ann Fam Med* 2005;3:514–22.
- Parker RM, Baker DW, Williams MV, et al. The test of functional health literacy in adults: a new instrument for measuring patients' literacy skills. *J Gen Intern Med* 1995;10:537–41.
- Sørensen K, Van den Broucke S, Pelikan JM, et al. Measuring health literacy in populations: illuminating the design and development process of the European health literacy survey questionnaire (HLS-EU-Q). *BMC Public Health* 2013;13:948.
- Norman CD, Skinner HA. eHEALS: the eHealth literacy scale. *J Med Internet Res* 2006;8:e27.
- Nutbeam D. Defining, measuring and improving health literacy. *Health evaluation and promotion* 2015;42:450–6.
- Nutbeam D. Health promotion glossary. *Health Promot Int* 1998;13:349–64. Volume.
- Nutbeam D. The evolving concept of health literacy. *Soc Sci Med* 2008;67:2072–8.
- Jorm AF, Korten AE, Jacomb PA, et al. "Mental health literacy": a survey of the public's ability to recognise mental disorders and their beliefs about the effectiveness of treatment. *Med J Aust* 1997;166:182–6.
- Jorm AF. Mental health literacy: empowering the community to take action for better mental health. *Am Psychol* 2012;67:231–43.
- Marques SRL, Lemos SMA. Instrumentos de avaliação do letramento em saúde: revisão de literatura. *Audiology - Communication Research* 2017;22:e1757.
- Nutbeam D, Muscat DM. Advancing health literacy interventions. *Stud Health Technol Inform* 2020;269:115–27.
- American Medical Association. Health literacy: report of the Council on scientific affairs. AD hoc Committee on health literacy for the Council on scientific Affairs, American Medical association. *JAMA* 1999;281:552–7.
- Institute of Medicine. *Health literacy: a prescription to end confusion*. Washington DC: National Academies Press, 2004.
- Chehuen Neto JA, Costa LA, Estevanin GM, et al. Functional health literacy in chronic cardiovascular patients. *Cien Saude Colet* 2019;24:1121–32.
- Clough BA, Nazareth SM, Casey LM. Making the grade: a pilot investigation of an e-intervention to increase mental health literacy and help-seeking intentions among international university students. *Br J Guid Couns* 2020;48:347–59.
- O'Connor M, Casey L. The mental health literacy scale (MHLS): a new scale-based measure of mental health literacy. *Psychiatry Res* 2015;229:511–6.
- Clough BA, Nazareth SM, Day JJ, et al. A comparison of mental health literacy, attitudes, and help-seeking intentions among domestic and international tertiary students. *Br J Guid Couns* 2019;47:123–35.
- Johnco C, Salloum A, McBride NM, et al. Mental health literacy, treatment preferences, and barriers in Salvadorian parents. *Int J Ment Health* 2019;48:139–64.
- Gray KM. From content knowledge to community change: a review of representations of environmental health literacy. *Int J Environ Res Public Health* 2018;15:466.
- Nye R, Robinia K. The effect of a community based educational intervention on oral health knowledge levels of adult caregivers of young children in the rural upper Peninsula of Michigan. *J Community Health Nurs* 2019;36:188–98.
- Department of Health and Human Services. Office of disease prevention and health promotion. No. 21 oral health: terminology. *Healthy People* 2010;2010. [Epub ahead of print: Retrieved from] <https://healthypeople.gov/2010/Document/HTML/Volume2/21Oral.htm>
- Waldmann T, Staiger T, Oexle N, et al. Mental health literacy and help-seeking among unemployed people with mental health problems. *J Ment Health* 2020;29:270–6.
- Wilson CJ, Deane FP, Ciarrochi J, et al. Measuring help-seeking intentions: properties of the general help-seeking questionnaire. *Canadian Journal of Counselling and Psychotherapy* 2005;39:15–28.
- Williston SK, Bramande EA, Vogt DS, et al. An examination of the roles of mental health literacy, treatment-seeking stigma, and

- perceived need for care in female veterans' service use. *Psychiatr Serv* 2020;71:144–50.
- 52 An J-Y, Moon H, Cha S. Health literacy, depression, and stress among Chinese immigrants in South Korea. *Public Health Nurs* 2019;36:603–14.
 - 53 Chew LD, Bradley KA, Boyko EJ. Brief questions to identify patients with inadequate health literacy. *Fam Med* 2004;36:588–94.
 - 54 Azizi N, Karimy M, Abedini R, et al. Development and validation of the health literacy scale for workers. *Int J Occup Environ Med* 2019;10:30–9.
 - 55 Brandstetter S, Atzendorf J, Seelbach-Göbel B, et al. Sociodemographic factors associated with health literacy in a large sample of mothers of newborn children: cross-sectional findings from the KUNO-Kids birth cohort study. *Eur J Pediatr* 2020;179:165–9.
 - 56 Degan TJ, Kelly PJ, Robinson LD, et al. Health literacy in people living with mental illness: a latent profile analysis. *Psychiatry Res* 2019;280:112499 <https://www.sciencedirect.com/science/article/pii/S0165178119307140>
 - 57 Eo Y-S, Kim J-S. Associations of health belief and health literacy with Pap smear practice among Asian immigrant women. *Eur J Oncol Nurs* 2019;42:63–8.
 - 58 An J, Yang SJ. Development of a health literacy assessment scale for Asian immigrant women in South Korea. *J Korean Acad Community Health Nurs* 2015;26:330–41.
 - 59 Ernsting C, Stühmann LM, Dombrowski SU, et al. Associations of health APP use and perceived effectiveness in people with cardiovascular diseases and diabetes: population-based survey. *JMIR Mhealth Uhealth* 2019;7:e12179.
 - 60 Erunal M, Ozkaya B, Mert H, et al. Investigation of health literacy levels of nursing students and affecting factors. *International Journal of Caring Sciences* 2019;12:270–9.
 - 6160 Okyay P, Abacigil F, Harlak H. *Turkish health literacy scales reliability and validity study. 1. edition*. Ankara: Saglik Bakanligi Yayin No, 2016: 43–60.
 - 62 Evans A-Y, Anthony E, Gabriel G. Comprehensive health literacy among undergraduates: a Ghanaian University-Based cross-sectional study. *Health Lit Res Pract* 2019;3:e227–37.
 - 63 Fernández-Gutiérrez M, Bas-Sarmiento P, Poza-Méndez M. Effect of an mHealth intervention to improve health literacy in immigrant populations: a quasi-experimental study. *Comput Inform Nurs* 2019;37:142–50.
 - 64 Goto E, Ishikawa H, Okuhara T, et al. Relationship of health literacy with utilization of health-care services in a general Japanese population. *Prev Med Rep* 2019;14:100811.
 - 65 Guclu OA, Demirci H, Ocakoglu G, et al. Relationship of pneumococcal and influenza vaccination frequency with health literacy in the rural population in turkey. *Vaccine* 2019;37:6617–23.
 - 66 Güner MD, Ekmekci PE. A survey study evaluating and comparing the health literacy knowledge and communication skills used by nurses and physicians. *INQUIRY* 2019;56:004695801986583.
 - 67 Hu A, Yang XY. Is College education an equalizer for social disparities in health literacy? A case study in Shanghai, China. *Glob Health Promot* 2019;26:62–70.
 - 68 Jordan JE, Buchbinder R, Briggs AM, et al. The health literacy management scale (HeLMS): a measure of an individual's capacity to seek, understand and use health information within the healthcare setting. *Patient Educ Couns* 2013;91:228–35.
 - 69 Indino K, Sharp R, Esterman A. The effect of health literacy on treatment adherence in maintenance haemodialysis patients: a cross-sectional study. *Renal Society of Australasia Journal* 2019;15:11–18 <https://search.informit.org/doi/>
 - 70 Kaper MS, Winter AFde, Bevilacqua R, et al. Positive outcomes of a comprehensive health literacy communication training for health professionals in three European countries: a multi-centre pre-post intervention study. *Int J Environ Res Public Health* 2019;16. doi:10.3390/ijerph16203923. [Epub ahead of print: 15 10 2019].
 - 71 Kwan B, Frankish J, Rootman I, et al. The Development and Validation of Measures of "Health Literacy" in Different Populations. *UBC Institute of Health Promotion Research and University of Victoria Community Health Promotion Research*;2006.
 - 72 Kobayashi R, Ishizaki M. Examining the interaction between medical information seeking online and understanding: exploratory study. *JMIR Cancer* 2019;5:e13240.
 - 73 Lim BT, Huang Y-J, Shepherd HL, et al. Health literacy and cancer care coordination in Chinese migrant patients and their carers: a cross-sectional survey. *Psychooncology* 2019;28:1048–55.
 - 74 Logullo P, Torloni MR, de O C Latorraca C, et al. The Brazilian Portuguese version of the discern instrument: translation procedures and psychometric properties. *Value Health Reg Issues* 2019;20:172–9.
 - 75 Apolinario D, Braga RdeCOP, Magaldi RM, et al. Short assessment of health literacy for Portuguese-speaking adults. *Rev Saude Publica* 2012;46:702–11.
 - 76 Lu M, Xia H, Ma J, et al. Relationship between adherence to secondary prevention and health literacy, self-efficacy and disease knowledge among patients with coronary artery disease in China. *Eur J Cardiovasc Nurs* 2020;19:230–7.
 - 77 Lu M, Ma J, Lin Y, et al. Relationship between patient's health literacy and adherence to coronary heart disease secondary prevention measures. *J Clin Nurs* 2019;28:2833–43.
 - 78 Ma Z, Wu M. The psychometric properties of the Chinese eHealth literacy scale (C-eHEALS) in a Chinese rural population: cross-sectional validation study. *J Med Internet Res* 2019;21:e15720.
 - 79 Nokes KM, Reyes DM. Do brief educational sessions increase electronic health literacy of low-income persons living with HIV/AIDS? *Comput Inform Nurs* 2019;37:315–20.
 - 80 Oh HJ, Lee H. When do people verify and share health Rumors on social media? the effects of message importance, health anxiety, and health literacy. *J Health Commun* 2019;24:837–47.
 - 81 Oliffe JL, McCreary DR, Black N, et al. Canadian men's health literacy: a nationally representative study. *Health Promot Pract* 2020;21:993–1003.
 - 82 Peerson A, Saunders M. In search of a gender lens.. *International Journal of Men's Health*. 2011;2.
 - 83 Pobhirun T, Pinitsoontorn S. The association between health literacy and pesticide use behaviors among sweet corn farmers in the PAK Chong district of Thailand: a cross-sectional study. *F1000Res* 2019;8:448.
 - 84 Rababah JA, Al-Hammouri MM, Drew BL, et al. Health literacy: exploring disparities among college students. *BMC Public Health* 2019;19:1401.
 - 85 Rüegg R, Abel T. The relationship between health literacy and health outcomes among male young adults: exploring confounding effects using decomposition analysis. *Int J Public Health* 2019;64:535–45.
 - 86 Selden CR, Zorn M, Ratzan S, et al/Selden CR, ed. *Health literacy*. Bethesda: National Library of Medicine, 2000.
 - 87 Abel T, Hofmann K, Ackermann S, et al. Health literacy among young adults: a short survey tool for public health and health promotion research. *Health Promot Int* 2015;30:725–35.
 - 88 Solhjoo N, Naghshineh N, Fahimnia F. Veterinarians' information Prescription and Clients' eHealth Literacy. *J Can Health Libr Assoc* 2019;40:32–44.
 - 89 Stømer UE, Göransson LG, Wahl AK, et al. A cross-sectional study of health literacy in patients with chronic kidney disease: associations with demographic and clinical variables. *Nurs Open* 2019;6:1481–90.
 - 90 Uysal N, Ceylan E, Koç A. Health literacy level and influencing factors in university students. *Health Soc Care Community* 2020;28:505–11.
 - 91 Wang X, Gao H, Xu H. Cluster analysis of unhealthy lifestyles among elderly adults with prediabetes: a cross-sectional study in rural China. *Diabetes Ther* 2019;10:1935–48.
 - 92 Kutner ME, Greenberg E, Jin Y, et al. The health literacy of America's adults: results from the 2003 national assessment of adult literacy (NCES 2006–483). *National Cent Educ Stat* 2006;39:685–7.
 - 93 Leung AYM, Lou VWQ, Cheung MKT, et al. Development and validation of Chinese health literacy scale for diabetes. *J Clin Nurs* 2013;22:2090–9.
 - 94 Wang R-H, Hsu H-C, Chen S-Y, et al. Modeling patient empowerment and health literacy to glycemic control in insulin-treated patients: a prospective study. *Patient Educ Couns* 2019;102:1336–41.
 - 95 Nutbeam D. Defining and measuring health literacy: what can we learn from literacy studies? *Int J Public Health* 2009;54:303–5.
 - 96 Lee E-H, Lee YW, Lee K-W, et al. A new comprehensive diabetes health literacy scale: development and psychometric evaluation. *Int J Nurs Stud* 2018;88:1–8.
 - 97 Zhang J, Gilmour S, Liu Y, et al. Effect of health literacy on quality of life among patients with chronic heart failure in China. *Qual Life Res* 2020;29:453–61.
 - 98 Matsuoka S, Kato N, Kayane T, et al. Development and validation of a heart Failure-Specific health literacy scale. *J Cardiovasc Nurs* 2016;31:131–9.
 - 99 Anderson AN, Haardörfer R, Holstad MM, et al. A path analysis of patient and Social-Level factors on health literacy and retention in care among African Americans living with HIV. *AIDS Behav* 2020;24:1124–32.
 - 100 Avci G, Kordovski VM, Woods SP. A preliminary study of health literacy in an ethnically diverse university sample. *J Racial Ethn Health Disparities* 2019;6:182–8.

- 101 Davis TC, Long SW, Jackson RH, *et al.* Rapid estimate of adult literacy in medicine: a shortened screening instrument. *Fam Med* 1993;25:391–6.
- 102 Bonaccorsi G, Pieralli F, Innocenti M, *et al.* Non-Familial paid caregivers as potential flu carriers and cause of spread: the primary prevention of flu measured through their adherence to flu vaccination campaigns-A Florentine experience. *Hum Vaccin Immunother* 2019;15:2416–22.
- 103 Bonaccorsi G, Pieralli F, Innocenti M, *et al.* Health literacy among non-familial caregivers of older adults: a study conducted in Tuscany (Italy). *Int J Environ Res Public Health* 2019;16:3771.
- 104 Carducci A, Fiore M, Azara A, *et al.* Environment and health: risk perception and its determinants among Italian university students. *Sci Total Environ* 2019;691:1162–72.
- 105 Chen S-Y, Hsu H-C, Wang R-H, *et al.* Glycemic control in insulin-treated patients with type 2 diabetes: Empowerment perceptions and diabetes distress as important determinants. *Biol Res Nurs* 2019;21:182–9.
- 106 Lee Y-J, Shin S-J, Wang R-H. *Pathways of empowerment perceptions, health literacy, self-efficacy, and self-care behaviors to glycemic control in patients with type 2 diabetes mellitus, patient education and counseling, volume 99*, 2016: Pages 287–294, ISSN-3991. <https://www.sciencedirect.com/science/article/pii/S0738399115300586>
- 107 de Melo Bezerra JN, de Oliveira Lessa SR, Francisco do Ó M, *et al.* De Oliveira Tito Borba AK. health literacy of individuals undergoing dialysis therapy. *Texto & Contexto Enfermagem* 2019;28:1–13.
- 108 Cavanaugh KL. Health literacy in diabetes care: explanation, evidence and equipment. *Diabetes Manag* 2011;1:191–199.
- 109 Flynn PM, John MT, Sistani MM. Confirmation of the unidimensional structure of the oral health literacy adults questionnaire. *Int Dent J* 2019;69:207–13.
- 110 American Dental Association. *Health literacy in dentistry action plan: 2010–2015*. Chicago IL: ADA, 2009.
- 111 Naghibi Sistani MM, Montazeri A, Yazdani R, *et al.* New oral health literacy instrument for public health: development and pilot testing. *J Invest Clin Dent* 2014;5:313–21.
- 112 Gaikwad RN. Quantifying OHIP in the context with oral health literacy in rural and urban population. *J Contemp Dent Pract* 2019;20:834–7.
- 113 Ratzan SC, Parker RM. Introduction. In: Selden CR, Zorn M, Ratzan SC, eds. *National library of medicine current bibliographies in medicine:Health literacy*. NLM Pub. No. CBM. Bethesda: MD, 2000: 2000–1.
- 114 Lee JY, Rozier RG, Lee S-YD, *et al.* Development of a word recognition instrument to test health literacy in dentistry: the REALD-30—a brief communication. *J Public Health Dent* 2007;67:94–8.
- 115 Güner MD, Ekmekci PE. A survey study evaluating and comparing the health literacy knowledge and communication skills used by nurses and physicians. *Inquiry* 2019;56:004695801986583.
- 116 Han H-R, Kim K, Cudjoe J, *et al.* Familiarity, navigation, and comprehension: key dimensions of health literacy in Pap test use among Korean American women. *J Health Commun* 2019;24:585–91.
- 117 Han H-R, Huh B, Kim MT, *et al.* Development and validation of the assessment of health literacy in breast and cervical cancer screening. *J Health Commun* 2014;19 Suppl 2:267–84.
- 118 Himes DO, Davis SH, Lassetter JH, *et al.* Does family communication matter? exploring knowledge of breast cancer genetics in cancer families. *J Community Genet* 2019;10:481–7.
- 119 Irvin VL, Rohlman D, Vaughan A, *et al.* Development and validation of an environmental health literacy assessment screening tool for domestic well owners: the water environmental literacy level scale (wells). *Int J Environ Res Public Health* 2019;16. doi:10.3390/ijerph16050881. [Epub ahead of print: 11 03 2019].
- 120 Peters E. Beyond comprehension: the role of numeracy in judgments and decisions. *Current Directions in Psychological Science* 2012;21:31–5.
- 121 Kaur N, Kandelman D, Potvin L. Effectiveness of "Safeguard Your Smile," an oral health literacy intervention, on oral hygiene self-care behaviour among Punjabi immigrants:A randomized controlled trial. *Can J Dent Hyg* 2019;53:23–32.
- 122 people H. Vol. II: *Oral Health*. Washington, DC:US Government Printing Office 2010;2000.
- 123 Stucky BD, Lee JY, Lee S-YD, *et al.* Development of the two-stage rapid estimate of adult literacy in dentistry. *Community Dent Oral Epidemiol* 2011;39:474–80.
- 124 Kim K, Han H-R. The association between health literacy and breast and cervical cancer screening behaviors: findings from the behavioral risk factor surveillance system. *Nurs Res* 2019;68:177–88.
- 125 Kim S-W, Park W-Y, Jhon M, *et al.* Physical health literacy and health-related behaviors in patients with psychosis. *Clin Psychopharmacol Neurosci* 2019;17:279–87.
- 126 Joint Committee on National Health Education Standards. National health education standards: achieving health literacy. Reston, VA: Association for the advancement of health education; 1995.
- 127 Kim SH, Utz S. Effectiveness of a social Media-Based, health Literacy-Sensitive diabetes self-management intervention: a randomized controlled trial. *J Nurs Scholarsh* 2019;51:661–9.
- 128 Kino S, Kawachi I. Can health literacy boost health services utilization in the context of expanded access to health insurance? *Health Educ Behav* 2020;47:134–42.
- 129 Lin S-C, Chen I-J, Yu W-R, *et al.* Effect of a community-based participatory health literacy program on health behaviors and health empowerment among community-dwelling older adults: a quasi-experimental study. *Geriatr Nurs* 2019;40:494–501.
- 130 Tsai T-I, Lee S-YD, Tsai Y-W, *et al.* Methodology and validation of health literacy scale development in Taiwan. *J Health Commun* 2011;16:50–61.
- 131 Lindahl B, Norberg M, Johansson H, *et al.* Health literacy is independently and inversely associated with carotid artery plaques and cardiovascular risk. *Eur J Prev Cardiol* 2020;27:209–15.
- 132 Nutbeam D, Don Nutbeam IK. Advancing health literacy: a global challenge for the 21st century. *Health Promot Int* 2000;15:183–4.
- 133 Cavanaugh KL, Osborn CY, Tentori F, *et al.* Performance of a brief survey to assess health literacy in patients receiving hemodialysis. *Clin Kidney J* 2015;8:462–8.
- 134 Mackert M, Mabry-Flynn A, Donovan EE, *et al.* Health literacy and perceptions of stigma. *J Health Commun* 2019;24:856–64.
- 135 Berkman ND, Davis TC, McCormack L. Health literacy: what is it? *J Health Commun* 2010;15 Suppl 2:9–19.
- 136 Mayer S, Spickschen J, Stein KV, *et al.* The societal costs of chronic pain and its determinants: the case of Austria. *PLoS One* 2019;14:e0213889.
- 137 Meyers N, Glick AF, Mendelsohn AL, *et al.* Parents' use of technologies for health management: a health literacy perspective. *Acad Pediatr* 2020;20:23–30.
- 138 Miranda R, Meeks KAC, Snijder MB, *et al.* Health literacy and hypertension outcomes in a multi-ethnic population: the HELIUS study. *Eur J Public Health* 2020;30:516–21.
- 139 Mock MS, Sethares KA. Concurrent validity and acceptability of health literacy measures of adults hospitalized with heart failure. *Appl Nurs Res* 2019;46:50–6.
- 140 Mora-Pinzon MC, Chrischilles EA, Greenlee RT, *et al.* Variation in coordination of care reported by breast cancer patients according to health literacy. *Support Care Cancer* 2019;27:857–65.
- 141 Noback PC, Seetharaman M, Tantigate D, *et al.* Musculoskeletal health literacy, pain Catastrophization, and sleep quality: effect on functional disability scores. *J Surg Orthop Adv* 2019;28:144–9.
- 142 Paasche-Orlow MK, Wolf MS. The causal pathways linking health literacy to health outcomes. *Am J Health Behav* 2007;31 Suppl 1:19–26.
- 143 O'Connor R, Muellers K, Arvanitis M, *et al.* Effects of health literacy and cognitive abilities on COPD self-management behaviors: a prospective cohort study. *Respir Med* 2019;160:105630.
- 144 Penalosa R, Navarro JI, Jolly PE, *et al.* Health literacy and knowledge related to tuberculosis among outpatients at a referral hospital in Lima, Peru. *Res Rep Trop Med* 2019;10:1–10.
- 145 Lee S-YD, Bender DE, Ruiz RE, *et al.* Development of an easy-to-use Spanish health literacy test. *Health Serv Res* 2006;41:060720074824046–???
- 146 Rafferty AP, Winterbauer NL, Luo H, *et al.* Diabetes self-care and clinical care among adults with low health literacy. *J Public Health Manag Pract* 2021;27:144–53.
- 147 Tavakoly Sany SB, Peyman N, Zadehahmad Z, *et al.* Effect of educational interventions on health literacy in patients with heart failure. *Int J Health Promot Educ* 2019;57:23–36.
- 148 Scrivner B, Szaflarski M, Baker EH, *et al.* Health literacy and quality of life in patients with treatment-resistant epilepsy. *Epilepsy Behav* 2019;99:106480.
- 149 Tucker CM, Kang S, Ukonu NA, *et al.* A culturally sensitive Church-Based Health-Smart intervention for increasing health literacy and health-promoting behaviors among black adult Churchgoers. *J Health Care Poor Underserved* 2019;30:80–101.
- 150 VanWormer JJ, Tambe SR, Acharya A. Oral health literacy and outcomes in rural Wisconsin adults. *J Rural Health* 2019;35:12–21.
- 151 Naghibi Sistani MM, Montazeri A, Yazdani R, *et al.* New oral health literacy instrument for public health: development and pilot testing. *J Invest Clin Dent* 2014;5:313–21.

- 152 Weaver SB, Maneno M, Chijioke P, *et al.* Factors associated with health literacy and medication adherence in an urban community. *Sr Care Pharm* 2019;34:604–12.
- 153 Zhong Y, Patel N, Ferris M, *et al.* Health literacy, nutrition knowledge, and health care transition readiness in youth with chronic kidney disease or hypertension: a cross-sectional study. *J Child Health Care* 2020;24:246–59.
- 154 Kindig DA, Panzer AM, Nielsen-Bohlman L, *et al.* *Health literacy: a prescription to end confusion*, 2004.
- 155 Williams TA, Wolf MS, Parker RM, *et al.* Parent dosing tool use, beliefs, and access: a health literacy perspective. *J Pediatr* 2019;215:244–51.
- 156 Winokur EJ, Rutledge DN, McGowan JJ. A picture is worth a thousand words: Pictographs to improve understanding of discharge instructions. *J Emerg Nurs* 2019;45:531–7.
- 157 Morrison AK, Myrvik MP, Brousseau DC, *et al.* The relationship between parent health literacy and pediatric emergency department utilization: a systematic review. *Acad Pediatr* 2013;13:421–9.