

BMJ Open Loneliness and mental well-being in the Polish population during the COVID-19 pandemic: a cross-sectional study

Ewa Kobos ,^{1,2} Barbara Knoff,² Beata Dziedzic,² Rafał Maciąg,² Anna Idzik²

To cite: Kobos E, Knoff B, Dziedzic B, *et al.* Loneliness and mental well-being in the Polish population during the COVID-19 pandemic: a cross-sectional study. *BMJ Open* 2022;**12**:e056368. doi:10.1136/bmjopen-2021-056368

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-056368>).

Received 12 August 2021
Accepted 10 January 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Development of Nursing, Social and Medical Sciences, Faculty of Health Sciences, Medical University of Warsaw, Warsaw, Poland

²Medical University of Warsaw, Warsaw, Poland

Correspondence to

Dr Ewa Kobos;
ekobos@wum.edu.pl

ABSTRACT

Objective There is a growing concern that the restrictions imposed due to the COVID-19 pandemic could lead to increased loneliness and mental disorders, which are considered a major public health problem. The aim of the study was to assess loneliness, anxiety, depression and irritability in the Polish population during the COVID-19 pandemic.

Design A cross-sectional study. The study participants completed an online questionnaire using the computer-assisted web interview technique. Data were collected using Revised UCLA Loneliness Scale and Hospital Anxiety and Depression Scale.

Setting Poland.

Participants This study was conducted between 6 October and 12 October 2020, in a representative sample of 890 Polish residents.

Primary outcome measures Self-reported loneliness; sociodemographic and COVID-19 pandemic factors associated with loneliness, anxiety, depression and irritability.

Results The analyses showed a moderately high degree of loneliness in 22%, symptoms of anxiety in 27%, depression in 14% and irritability in 33% of the respondents. The increasing severity of anxiety, depression and irritation in the study group was accompanied by higher loneliness. Generally, younger people, both tested positive for SARS-CoV-2 and those who experienced home quarantine, scored higher in both scales.

Conclusions It is necessary to identify those most vulnerable to loneliness, anxiety and depression during a crisis to assess health needs and proactively allocate resources during and after the pandemic. Loneliness, anxiety, depression and irritability are important factors to consider in a population of younger, disadvantaged people, who tested positive for SARS-CoV-2, people who were quarantined at home, and people who believe that their physical and mental health is worse than in the pre-pandemic period. It is important to cater for the mental health of individuals during the COVID-19 pandemic and to promote psychological interventions to improve mental well-being in potentially vulnerable social groups.

BACKGROUND

In March 2020, governments of many countries began to implement relatively harsh pandemic-related restrictions and most citizens were confined to their homes to limit

Strengths and limitations of this study

- The study was conducted in a representative sample of Polish residents.
- The current data were collected through online surveys, which may have excluded potential respondents who lack internet access or are not knowledgeable about modern technologies.
- The cross-sectional survey design precluded conclusions about causality.

the spread of the SARS-CoV-2 virus. The first case of SARS-CoV-2 infection in Poland was confirmed on 4 March 2020.^{1 2} Full lockdown was implemented on 25 March, with illegal gatherings of more than two people and restrictions on leaving home, including a recommendation to work remotely.³ This meant that most of the employees started working from home and that people stayed at home most of the time. The restrictions involved the protection of mouth and nose in public places, social distancing, cancellation of public events, limiting the number of customers in stores, closing of shopping centres, ban on international travel and closed airports. In the spring of 2020, the COVID-19 epidemic was suppressed by implementing massive restrictions on social contacts.⁴ As of 23 December 2021, the global number of reported COVID-19 cases reached 276 436 619, including 5 374 744 deaths, and it continues to grow.⁵ In Poland, from 3 January 2020 to 23 December 2021, there were 4 017 420 confirmed cases of COVID-19 with 93 445 deaths.⁵ In the period from 6 to 12 October 2020, when this study was conducted, a total of 2236–5300 daily new cases and from 32 to 75 deaths per day were recorded in Poland, and the total number of cases was 104 316 on 6 October 2020 and 130 210 on 12 October 2020.⁵

Loneliness can be defined as a distressing feeling resulting from a dissatisfaction with the quality and quantity of one's social



relationships.⁶ Loneliness has long been recognised as a risk factor for poor mental health; however, since the outbreak of the COVID-19 pandemic, it has emerged as a critical public health issue.^{7–9} Loneliness is a strong risk factor for many diseases, for example, cardiovascular diseases and is associated with an increased risk of mortality.^{7 10–13} Chronic loneliness is characterised by deficits in attention, affect, cognition, and behaviour that affect genetic, neural, and hormonal mechanisms, and thus contribute to higher morbidity and mortality rates.¹⁴ Apart from its detrimental influence on physical health, COVID-19 also affects people's mental well-being, which may lead to numerous psychological problems.^{15–20} People are more likely to experience different levels of fear of illness or death, a sense of helplessness and stigma after being infected with SARS-CoV-2.²¹ Loneliness may become an alarming global problem due to the pandemic-related social and physical distancing. Current research points to two important factors contributing to poor emotional well-being in lockdown: an increase in loneliness and more screen time.²² A study by Groarke *et al* demonstrated the incidence rate of loneliness among UK citizens to be 27% in lockdown, with one in four respondents admitting to feeling lonely 'in the previous 2 weeks'.²³ Increased loneliness during the pandemic was found in adult Norwegians, as compared with the pre-pandemic period.²⁴ In Canada, 8.4% of respondents reported feeling lonely for at least 5 days in the week prior to the study.²⁵ There are reports that show a positive correlation between the pandemic-related necessity to stay at home and the perceived level of loneliness. Tull *et al* found that although staying at home was associated with increased loneliness, the perceived impact of COVID-19 alone was negatively correlated with loneliness.¹⁷ Those suffering from COVID-19 received even greater support from their immediate environment. These findings correspond with suggestions that widespread shared COVID-19 experiences can increase social closeness and cohesion.²⁶

As far as mental health, loneliness and social isolation are concerned, not only do they strongly correlate with depression, but they may also increase the risk of future depressive episodes.^{27–30} The incidence rates of anxiety and depressive disorders in a systematic review of studies on stress, anxiety and depression in the general population during the COVID-19 pandemic were 31.9% and 33.7%, respectively.¹⁶ The highest prevalence of anxiety (32.9%) and depression (35.3%) was found in Asia, and the highest prevalence of stress was observed in Europe (31.9%).¹⁶ The general data on anxiety, depression and psychological stress during the pandemic show that the mean prevalence rates of anxiety and depression are 38.12% and 34.31%, respectively.³¹ Luo *et al* showed a combined incidence of anxiety and depressive disorders in 32% and 27% of the general population, respectively.³² Castaldelli-Maia *et al* estimated the worldwide pandemic-related prevalence of both depression and anxiety at 24.0% and 21.3%, respectively.³³

Latest research seems to confirm that people in isolation or quarantine are more likely to show significant levels of anxiety, anger, confusion and stress.³⁴ Research data from the COVID-19 pandemic demonstrate that people affected by quarantine present with stress, emotional anxiety, depression, mood swings, aggression, anger, attention deficits, hyperactivity, insomnia and post-traumatic stress.^{35–43} Contrary to the reviews of cross-sectional studies, a meta-analysis of longitudinal studies and natural experiments assessing the link between the COVID-19 lockdown and mental health indicates that the initial impact of the lockdown on mental health is relatively small. Research findings suggest high mental toughness of people in response to the necessity to stay at home, lockouts and restrictions that were likewise implemented nationally or regionally around the world in response to the COVID-19 pandemic.⁴⁴ Since these findings concern the first lockdowns during the COVID-19 pandemic, which were introduced between January and June 2020, the impact of repeated or prolonged lockdown on mental well-being requires further studies. The exact impact of the COVID-19 pandemic on mental health may vary across social groups, as well as across contexts and countries.

This study, which is representative of the Polish population, is one of the few studies conducted in Poland that focused on the impact of the COVID-19 pandemic on the mental health of Polish citizens. A study by Szczesniak *et al* demonstrated that apart from preventing the spread of COVID-19, the requirements on the use of face masks may have contributed to the perceived level of self-protection and promotion of social solidarity, which, in turn, may have improved the overall mental health.⁴⁵ However, it should be noted that these data were collected from 16 March to 26 April 2020, that is, in the first week of the regulations on the use of face masks in Poland. A study by Hamer and Baran on the impact of the pandemic on the mental well-being of Polish population showed increased loneliness and greater distress in the age group of 18–24 years old. The study was repeated four times between March and December 2020.⁴⁶ The research conducted in Poland in the group of students (20–26 April 2020, n=2170; mean age 22.1±2.2 years) showed moderate to very severe depression and anxiety in 43.4% and 27.3% of the respondents, respectively.⁴⁷ A mean Revised UCLA Loneliness Scale (R-UCLA) score of 42.041 out of a maximum of 80 was obtained in a study conducted 1.5 weeks after the introduction of home isolation in Poland (3 April 2020), in a group of 471 respondents (students accounted for 61.6% of the group; mean age 25.5±2 years).⁴⁸ It was shown that the criterion for minor mental disorders (≥24 points) in the General Health Questionnaire (GHQ) scale was met by 59.2% of respondents (n=2155) at the peak of the second wave of the pandemic in Poland (November 2021).⁴⁹

Early detection of mental disorders allows for more effective intervention strategies. Global health crises, for example, such as the COVID-19 pandemic, affect not

only healthcare workers, but also the general population, and result in undesirable psychological changes induced by fear, anxiety, depression or insecurity.⁵⁰ The Lancet published a consensus statement calling for research on the impact of the COVID-19 pandemic on mental health in the whole population as well as in vulnerable groups.⁵¹ It is therefore necessary to assess and supervise the mental health of various population groups during crises such as a pandemic. The aim of the study was to assess loneliness, anxiety, depression and irritability in the Polish population during the COVID-19 pandemic.

METHODS

Patient and public involvement

No patient involved.

Participants and data collection

The study was conducted from 6 October to 12 October 2020 in a representative sample of Polish residents (N=890), which was selected using the random-quota method. A two-stage random sampling was performed, in which places of residents were randomly selected first, and then a quota sample was made. The quotas were selected for gender, age, education, size of the place of residence and province. The above-mentioned layers were used to select study participants. Fulfilment of the general population representativeness criteria by the study participants was the study criterion. Only those with access to the internet and ability to use a computer or other devices that enable answering questions via an internet panel were included in the study. After sending an inquiry regarding confirmation of consent to participate in the study, 40% of respondents refused to participate in the study. The size of the study group was supplemented with the assumed amount (890 participants in relation to the National Census of the Polish Population) in accordance with the above-mentioned amounts.

The respondents completed an online questionnaire using the computer-assisted web interview technique. The study was conducted by Kantar, which is one of the global research companies, using a website dedicated to this study. In compliance with the International Chamber of Commerce (ICC) /European Society for Opinion and Marketing Research (ESOMAR) International Code, personal data protection requirements and other applicable laws and principles set out in the national code of good research practice, the recruitment of study participants was conducted using the database of the research company responsible for conducting our study, in compliance with all the representativeness of the study group.⁵² Each of the potential respondents was asked for consent to conduct the study. It was only after obtaining the consent that the respondent received a link to the questionnaire. The answers provided by the respondents were stored in the Information Technology system of the company conducting the study. This was followed by database creation.

Measurement

Sociodemographic and COVID-19-related data

The questionnaire used to collect sociodemographic data of the study participants enquired about age, sex, place of residence, education, marital status, professional status, type of accommodation, number of people per household, income level, financial situation, and type of work before and during the pandemic. Questionnaire data related to the pandemic period included: a diagnostic test for SARS-CoV-2; home quarantine during the coronavirus pandemic and currently; home/hospital isolation; having family members/friends who were tested positive for SARS-CoV-2; and those in home quarantine or home/hospital isolation.

Revised UCLA Loneliness Scale

We used the Polish adaptation of the R-UCLA Loneliness Scale to assess loneliness.⁵³ The scales are characterised by good reliability: total score $\alpha=0.92$, intimate others $\alpha=0.90$, social others $\alpha=0.83$, belonging and affiliation $\alpha=0.80$. The original scale, that is, UCLA Loneliness Scale, was first designed by Russell, Peplau and Cutrona.⁵⁴ It comprises 20 items; the respondents indicate how often each statement is true for them on a 4-point scale (1=I never feel this way, 4=I often feel this way). The maximum score is 80. The total score is the sum of scores from three subscales: belonging and affiliation, intimate others and social others.⁵⁵ Based on Perry's classification, the following degrees of loneliness were set for the entire scale: 65–80, a high degree; 50–64, a moderately high degree; 35–49, a moderate degree; 20–34, a low degree of loneliness.⁵⁶ In this study, the reliability coefficient was $\alpha=0.91$ for the entire questionnaire, $\alpha=0.90$ for intimate others (subscale); $\alpha=0.86$ for social others (subscale), and $\alpha=0.69$ for belonging and affiliation (subscale).

Hospital Anxiety and Depression Scale

The Polish version of Hospital Anxiety and Depression Scale (HADS-M) comprises 16 questions, with 2 additional questions on irritability.⁵⁷ It is a modified version of HADS (the Hospital Anxiety and Depression Scale) by Zigmond and Snaith.⁵⁸ The HADS-M includes three independent subscales: depression, anxiety and irritability; consisting of a total of 16 questions (each can be scored 0–3): depression subscale (seven questions) with the maximum score of 21; anxiety subscale (seven questions) with the maximum score of 21; irritability subscale (two questions) with the maximum score of 6. A higher score in the subscales indicates a higher level of the feature assessed. Based on the original scale, the following interpretation of the anxiety and depression subscales was used to analyse the prevalence of symptoms: no disorders (0–7), borderline states (8–10) and disorders (>10). The interpretation of the irritability subscale was as follows: 0–2: no disorders, 3: borderline states, 4–6: disorders. The maximum score was 48. For the consolidated evaluation, individual states can be defined as: 0–16, no disorders; 17–22, border states; 23–48, disorders.

**Table 1** Subjects' R-UCLA scores (n=890)

R-UCLA Scale	Degrees of loneliness (%)					
	Subscales	M	SD	Low	Moderate	Moderately high
Intimate others	22.40	6.77	36	40	22	2
Social others	8.46	3.18				
Belonging and affiliation	9.52	2.95				
Total score	40.40	10.79				

M, mean difference; R-UCLA, Revised UCLA Loneliness Scale.

In most analyses, the two subscales of depression and irritability are combined to present the results more clearly. In this study, the total reliability was $\alpha=0.92$; $\alpha=0.87$ for the anxiety, $\alpha=0.84$ for the depression and $\alpha=0.88$ for the irritability subscale.

Data analysis

The Shapiro-Wilk test was used to verify the normality of distributions and the Levene test was used to assess the homogeneity of variance. The following tests were used to assess the differences between the groups: for more than two compared groups, one-way analysis of variance or (when the assumption of equal variance was not met) the Welch's test along with the Tukey's post-hoc tests; under special circumstances (small and very diverse numbers between groups), the Kruskal-Wallis test was used along with the Dunn post-hoc test; for two compared groups, Student's t-test for independent variables or the Cochran-Cox test (when the assumption of equal variance was not met). Correlation analyses were performed using the Pearson linear correlation coefficient or the Spearman correlation coefficient (r). Statistica V.10.0 was used for calculations. The results of the analyses were considered statistically significant at $p \leq 0.05$.

RESULTS

Characteristics of the study group

The study included 49.2% men and 50.8% women. The mean age of respondents was 44.3 ± 16.10 years. Secondary education was declared by 51.5% of respondents, there were 38% of rural respondents, 52% of professionally active respondents and 56.4% of married respondents. A

total of 51.5% of the respondents lived in a single-family house, 59.4% of the respondents lived with their family and 10.3% lived alone (see online supplemental file 1).

Loneliness

Table 1 summarises data on the levels of loneliness in the study group. Overall, the mean R-UCLA score was 40.40 ± 10.79 . Moderately high and high degree of loneliness were found in 22% and 2% of respondents, respectively.

Anxiety, depression and irritability

Table 2 shows the data obtained in the HADS-M. The mean score obtained by the respondents was 15 ± 9.56 ; disorders in anxiety and depression subscales were found in 27% and 14% of respondents, respectively.

Anxiety, depression, irritability and the feeling of loneliness

Table 3 shows positive statistically significant correlations between HADS-M and R-UCLA scores. The data show that the rate of loneliness increased with an increase in anxiety, depression and irritability.

Health self-assessment and anxiety, depression, irritability and loneliness

Negative, statistically significant correlations between health status self-assessment and the results in both scales are presented in online supplemental file 2. The self-rating of general physical and mental health decreased with increasing scores obtained by the respondents on the scale of loneliness, anxiety, depression and irritability ($p=0.000$). The strength of the relationship was higher for mental health self-assessment ($R=-0.552$; $R=-0.340$).

Table 2 Subjects' HADS-M scores (n=890)

HADS-M subscales	M	SD	Presence of disorders		
			No disorders	Borderline abnormal	Presence of disorders
			%	%	%
Anxiety	7.22	4.71	57	16	27
Depression	5.30	4.29	68	18	14
Irritability	2.47	1.75	56	11	33
Depression and irritability	7.77	5.32	53	15	32
Total score	15.00	9.56	58	17	25

HADS-M, Modified Hospital Anxiety and Depression Scale; M, mean difference.

Table 3 Correlations between sense of loneliness and anxiety, depression and irritability (n=890)

HADS-M subscales		R-UCLA subscales			
		Intimate others	Social others	Belonging and affiliation	Total score
Anxiety	R	0.4298	0.3547	0.3129	0.4597
	P value	***	***	***	***
Depression	R	0.4324	0.4465	0.3796	0.5067
	P value	***	***	***	***
Depression and Irritability	R	0.4374	0.4212	0.3515	0.4946
	P value	***	***	***	***
Total score	R	0.4552	0.4092	0.3498	0.5018
	P value	***	***	***	***

***P<0.001, correlations were statistically significant.

HADS-M, Modified Hospital Anxiety and Depression Scale; R, Spearman's correlation coefficient; R-UCLA, Revised UCLA Loneliness Scale.

Respondents who reported deteriorated health compared with the pre-pandemic period also scored higher for loneliness, anxiety, depression and irritability.

Sociodemographic variables and anxiety, depression and loneliness

Table 4 indicates that women show significantly higher levels of anxiety, depression and irritability (16.05) compared with men (13.91). As far as the depression and loneliness subscale is concerned, the results did not differ significantly depending on gender. The analyses showed statistically significant negative correlations between R-UCLA and HADS-M scores and the age of respondents.

Younger age correlated with higher anxiety, depression and irritability, as well as higher loneliness. Correlations are presented in table 5.

Overall, the R-UCLA scale (44.45) and the intimate others subscale (25.84) showed a significantly higher loneliness in respondents with the lowest level of education (see online supplemental file 3).

No significant correlation was found between the feeling of loneliness and the place of residence of respondents (p=0.856). Anxiety, depression and irritability

increased with the decreasing size of the place of residence in the total study group. This is presented in online supplemental file 4.

The differences in the severity of loneliness, anxiety, depression and irritability depending on the professional status of respondents are reported in online supplemental file 5. Significantly, higher loneliness was shown in pupils and students for the entire R-UCLA scale (44.10), as well as the intimate others (24.72) and social others subscales (9.38).

The study showed no significant differences in the severity of anxiety, depression and irritability depending on the marital status of respondents (p=0.322). Single respondents showed significantly higher loneliness (43.85) both for total and individual subscale scores (see online supplemental file 6).

Respondents living in single-family houses showed significantly higher levels of anxiety, depression and irritability on the HADS-M (15.80) and its subscales. This is presented in online supplemental file 7.

A significant positive correlation was found between the intensity of anxiety, depression and irritability and

Table 4 HADS-M and R-UCLA scores with regard to the subjects' gender

HADS-M; R-UCLA	Male (n=438)		Female (n=452)		t	P value
	M	SD	M	SD		
Anxiety	6.51	4.55	7.90	4.76	-4.443	***
Depression	5.14	4.29	5.47	4.28	-1.146	
Depression and irritability	7.39	5.26	8.14	5.36	-2.099	*
HADS-M—total score	13.91	9.35	16.05	9.66	-3.351	***
Intimate others	22.32	6.61	22.48	6.93	-0.342	
Social others	8.59	3.30	8.34	3.05	1.144	
Belonging and affiliation	9.58	3.03	9.46	2.86	0.594	
R-UCLA—total score	40.50	10.78	40.29	10.82	0.284	

*P<0.05, ***p<0.001, correlations were statistically significant.

HADS-M, Modified Hospital Anxiety and Depression Scale; R-UCLA, Revised UCLA Loneliness Scale; t, Student's t-test.

**Table 5** HADS-M and R-UCLA scores with regard to the subjects' age

	Age	
	r	P value
HADS-M		
Anxiety	-0.1580	***
Depression	-0.1167	***
Depression and irritability	-0.1634	***
Total score	-0.1688	***
R-UCLA		
Intimate others	-0.1640	***
Social others	-0.1950	
Belonging and affiliation	-0.0977	**
Total score	-0.1870	***

P<0.01, *p<0.001, correlations were statistically significant. HADS-M, Modified Hospital Anxiety and Depression Scale; r, Pearson correlation coefficient; R-UCLA, Revised UCLA Loneliness Scale.

the number of people per household (see online supplemental file 8).

The feeling of loneliness was significantly higher in the respondents living alone (see online supplemental file 9).

Correlations between the financial situation/income and loneliness, as well as the severity of anxiety, depression and irritability in the study group are shown in online supplemental file 10. Positive, statistically significant correlations were found between these variables. The intensity of anxiety, depression and irritability increased with the deterioration of the financial situation (p=0.000) and lower income (p=0.005). Higher loneliness was accompanied by a significantly worse financial situation (p=0.000) and lower income (p=0.022).

Variables related to the COVID-19 pandemic and anxiety, depression, irritability and loneliness

Table 6 shows the intensity of anxiety, depression and irritability as well as loneliness depending on the respondent's testing for SARS-CoV-2. Generally, a significantly higher level of anxiety, depression, irritability (23.14) and loneliness (48.42) was found in the group of patients with a positive test result.

Table 7 containing data on the respondent's experience of home quarantine shows that anxiety, depression, irritability (19.16) and loneliness (43.37) were significantly higher in those who experienced home quarantine.

Respondents who did not know any people with a positive test result scored significantly higher on the loneliness scale (40.78) (see online supplemental file 11).

Data on the correlation between respondents' knowledge of people who were/are under home quarantine or home/hospital isolation and HADS-M/R-UCLA scores are illustrated in online supplemental file 12. A significantly higher level of loneliness (41.63) was found among respondents who did not know such individuals.

Our data show that respondents who worked from home before the pandemic scored significantly higher on the individual subscales of the HADS-M and overall (19.33) on the entire scale (see online supplemental file 13).

People working from home during the pandemic scored higher on the anxiety subscale (p=0.041) and on the social others subscale (p=0.042) of loneliness. This is presented in online supplemental file 14.

DISCUSSION

The aim of the study was to assess loneliness, anxiety, depression and irritability in the Polish population during the COVID-19 pandemic. The unexpected pandemic has affected the majority of the world's population, changing and complicating the everyday life of many societies.

Table 6 Diagnostic test for SARS-CoV-2 and the HADS-M and R-UCLA scores

HADS-M; R-UCLA	No (n=780)		Yes, negative result (n=74)		Yes, positive result (n=14)		H	P value
	M	SD	M	SD	M	SD		
Anxiety	7.05	4.71	8.14	4.79	10.28	2.72	12.287	**
Depression	5.13	4.28	5.98	4.24	9.78	2.66	19.212	***
Depression and irritability	7.55	5.30	8.68	5.53	12.85	3.43	16.897	***
HADS-M—total score	14.61	9.55	16.83	9.91	23.14	4.62	16.157	***
Intimate others	22.30	6.65	21.74	7.25	25.85	7.28	5.797	
Social others	8.40	3.09	8.58	3.54	11.35	4.60	4.268	
Belonging and affiliation	9.52	2.90	9.08	3.06	11.21	3.94	4.862	
R-UCLA—total score	40.23	10.72	39.40	11.79	48.42	10.02	8.762	*

*P<0.05, **p<0.01, ***p<0.001, correlations were statistically significant.

H, Kruskal-Wallis test; HADS-M, Modified Hospital Anxiety and Depression Scale; M, mean difference; R-UCLA, Revised UCLA Loneliness Scale.

Table 7 Home quarantine and the HADS-M and R-UCLA scores

HADS-M; R-UCLA	No (n=819)		Yes (n=48)		t	P value
	M	SD	M	SD		
Anxiety	7.08	4.67	8.70	5.26	-2.322	*
Depression	5.16	4.24	6.93	4.82	-2.781	**
Depression and irritability	7.55	5.24	10.45	6.21	-3.688	***
HADS-M—total score	14.63	9.43	19.16	11.11	-3.197	**
Intimate others	22.14	6.69	24.70	6.62	-2.576	*
Social others	8.42	3.15	9.33	3.37	-1.937	*
Belonging and affiliation	9.53	2.95	9.33	2.79	0.464	
R-UCLA—total score	40.10	10.84	43.37	10.06	-2.033	*

*P<0.05, **p<0.01, ***p<0.001, correlations were statistically significant.

HADS-M, Modified Hospital Anxiety and Depression Scale; M, mean difference; R-UCLA, Revised UCLA Loneliness Scale; t, Student's t-test.

Despite certain cultural differences in social perception, it can be assumed that experiencing a pandemic is a collective, global experience, and it affects mental health regardless of the latitude, causing justified anxiety in the vast majority of humans.⁵⁹ In total, the respondents had a mean R-UCLA score of 40.4±10.79, which is lower (42.04±11.78) compared with Bartoszek *et al*⁴⁸ who used the same research tool in Poland. However, it should be noted that the compared study involved mainly students (61.6%), with a mean age of 25.5 years, who were not a representative group for Poland. A Turkish study using social networking sites, which was conducted in the general population (n=860, March 2020), showed a significantly higher level of loneliness (57.94±8.26) compared with Poland.⁶⁰ Although the pandemic affects everyone, the intensity and severity of the response may vary depending on the latitude. Such comparisons between three different countries were presented by Emodi-Perlman *et al*.⁶¹ They found that Polish participants showed the second largest concern for mental health and relationships with relatives and friends.

Research suggests that certain demographic groups are at an increased risk of loneliness during COVID-19 self-isolation. In this study, the greatest level of loneliness was found in the age group of 16–29 years, which is consistent with the findings of Hamer and Baran,⁴⁶ who showed the highest level of loneliness in the youngest group (18–24 years) at each stage of the study. In the group of Turkish adolescents aged 12–18 years, the total R-UCLA score was 41.89±9.81⁶² and was lower compared with that obtained in this study in the group of 15–19 years old (44.05±11.01). This was also confirmed in a Canadian study by Wickens *et al*²⁵ who found the highest scores in the group of young adults aged 18–29 years, and the lowest scores among people aged ≥60 years. In our study, the growing levels of loneliness were accompanied by the younger age of the respondents, which is in line with the results of other authors, who found that during the COVID-19 pandemic, it is younger people who are more likely to suffer from loneliness, as well as the symptoms

of anxiety, depression or stress.^{63–66} This study also found the severity of anxiety, depression and irritability to be higher in younger respondents. Research confirms that the prevalence of psychological distress in the younger population is higher than in other age groups.^{46 67–69} The overall average HADS-M score in the Polish population aged ≥60 years was 65.18±4.06 out of possible 80 during the second wave (October 2020) of the pandemic.⁷⁰

Since the pandemic affects people's daily activities, its consequences will most probably be more severe for younger age groups than for older people (including the retired), who do not normally work and have far fewer other responsibilities outside the home. Greater loneliness perceived by younger adults may be a consequence of limited social interactions and reduced size of social networks as a result of social distancing. In comparison with the younger generations, older adults may attach more value to the quality of their interactions and social networks, rather than quantity.^{71–74} Furthermore, the pandemic may have affected the nature of social interaction. Limited direct personal contact with other young adults leads to greater loneliness.⁷⁵ In this study, the highest levels of loneliness were found in students who had been home schooled since the first lockdown. During a pandemic, this may be due to the fact that students and adolescents, just like adults, are concerned about their future, their own health and the health of their family and friends. Isolation and limited direct social interactions both inside and outside school can increase the feeling of loneliness.⁷⁶

Our study found no significant gender-related differences in the severity of loneliness, despite the fact that the data collected during the COVID-19 pandemic in other countries have shown that loneliness is more common in women than in men.^{25 48 63 77} Researchers imply that this may be related to the fact that when faced with a stressor, women rely on social support and emotion-focused coping more than men and these may be compromised in a situation of limited access to family and friends during a pandemic.^{78–80} This may be further supported by the fact



that women scored significantly higher for anxiety and depression than men both in this study and in the study by Hamer and Baran,⁴⁶ and Babicki *et al.*⁴⁹ McQuaid *et al* also report the highest rates of loneliness, depression and anxiety in young women.⁸¹

When comparing the results of this study with the one conducted in Poland by Bartoszek *et al.*⁴⁸ with a division into people living alone (47.10±11.28) and those living with a partner or family members (41.36±14.18), people living alone (42.66±12.00) as well as those living with a spouse/partner (38.68±9.45) and with family members (40.55±11) scored lower in our study. Single people scored significantly higher compared with those living with a spouse/partner, including the intimate others subscale, which may be due to the unpleasantness of being both emotionally and physically alone.¹⁴ Wickens *et al* also showed a higher level of loneliness in the group of single, unmarried, widowed and divorced people.²⁵ The analysis of potentially protective social factors showed that living with others protected against loneliness, with a 75% lower risk of the highest rate of loneliness in people living with others to those living alone.⁶³ On the other hand, this study showed that people living with a family or with a larger number of people per household experienced significantly higher levels of anxiety, depression and irritability. On the one hand, it can be assumed that living with a family is associated with a greater possibility of receiving help and support; on the other hand, it may give rise to additional fear for the health of other family members and increase the severity of mental health problems. Furthermore, functioning in a large community in one place of residence, especially during a lockdown, may be difficult. Sharing a household with other family members, who often work or learn remotely, for a long time may lead to organisational difficulties, conflicts, and thus contribute to deteriorated mental health.⁶⁹ For large families, the fear of losing a job and financial liquidity, as well as not being able to support one's own family, may intensify during a pandemic. Shahriarirad *et al* found a higher number of individuals per household to be a risk factor for anxiety.⁸²

According to Bu *et al*, life in a rural community can also be a protective factor, with a 24% lower risk of the highest rate of loneliness. People with a larger circle of close friends had a 42% lower chance of experiencing the highest rate of loneliness.⁶³ In this study, higher values on the scale and subscales of loneliness were shown in single people (43.85) compared with those in a relationship (38.72); however, no correlation between loneliness and the size of the place of residence or the number of people per household was confirmed. These data confirm previous research results implying a relationship between social network size and the perceived social support and a lower risk of loneliness.^{83–86} It is worth noting that the preference for face-to-face contact before the pandemic was not a risk factor for higher loneliness during lockdown, suggesting that a sudden change in social behaviour itself is not an independent predictor of loneliness.⁶³ In this

study, a larger place of residence was a protective factor for against anxiety, depression and irritability, which was not confirmed in Polish population by Babicki *et al* and Gambin *et al.*^{49 69} A study by Cao *et al* conducted among students showed living in urban areas as a protective factor for anxiety.⁸⁷

Studies, like those by Bu *et al*, did not confirm the relationship between lower education and higher loneliness, as noted by Wickens *et al* or Hamer and Baran.^{25 46 63} Lower education was a predictor of increased anxiety, depression and irritability in our study; depression in the study by Passavanti *et al.*^{75 88} and anxiety in the study by Babicki *et al* and Shahriarirad *et al.*^{49 82} Respondents with lower education may lack adequate health awareness, for example, about the use of personal protective equipment, the safety of vaccinations, they may not know how to act during a crisis, distrust experts and thus show a greater fear of contracting COVID-19. This group of respondents may also experience greater fear of losing a job and a source of income. People with lower education levels may also have fewer opportunities to maintain or find employment.

Social distancing during the COVID-19 pandemic has resulted in changes in the working environment and has contributed to job insecurity. Disruptions to work can be one of the main sources of stress during the pandemic.⁸⁹ A comparison of studies analysing the differences in mental health between employed and unemployed individuals during social distancing implementation has shown that the study participants who were employed had better mental health than those unemployed on all measures of mental health.⁹⁰ In this study, people working from home showed the highest levels of loneliness. They scored significantly higher for the social other factor, which refers, among other things, to the perceived lack of support from one's social network and a lack of a sense of closeness with other people (in terms of their availability). People who experienced redundancy or unemployment due to the pandemic reported loneliness more often, as shown by Wickens *et al.*²⁵ In a study by Alzueta *et al*, significant predictors of the severity of depression symptoms included difficulty to take care of others, and problems with transition to working from home.⁹¹ A study by Choi *et al.*⁹² revealed higher levels of depression and anxiety in those respondents who expressed concern about not being able to work from home during the pandemic. These findings correspond with a previously established fact that employment is a protective factor for loneliness in middle-aged adults.^{93 94}

In our study, we confirmed that people who considered their financial situation to be worse and had lower income were significantly more likely to experience loneliness, which corresponds to findings by Wickens *et al.*²⁵ and McQuaid *et al.*⁸¹ Anxiety, depression and irritability increased with the decrease in income. Studies showed that respondents with low income reported anxiety and psychological stress at the early stage of the pandemic more often.⁶⁸ Greater depression and anxiety symptoms

were noted in respondents whose income was negatively impacted by social restrictions and restriction on earning opportunities.^{49 95}

The occurrence and rapid spread of COVID-19 has led to a worldwide exacerbation of anxiety symptoms, which resulted in a higher incidence of severity of emotional distress. Since the outbreak of the pandemic, anxiety, symptoms of psychosis, trauma, suicidal thoughts and panic attacks have been reported in literature.⁹⁶ Recent research has shown that COVID-19 triggers anxiety, depression and post-traumatic stress symptoms.^{65 97} Research in Europe assessing the consequences of quarantine showed that anxiety, depression and stress occurred in 23.8%, 32.4% and 31.9% of the study population, respectively.³⁷ In our study, which used the HADS-M, disorders on the anxiety subscale and the depression subscale (including irritability) were shown in 27% and 32% of respondents, respectively. A Turkish study conducted on 14–16 April 2020 using HADS showed that 23.6% and 45.1% of the population scored above the depression and anxiety cut-off point, respectively.⁹⁸ Compared with our findings, the scores in the anxiety subscale were much higher, probably due to the fact that the study was conducted in the initial period of the pandemic.

Research confirms that people whose family members, relatives or friends developed COVID-19 show significantly higher severity of anxiety.^{23 49 99 100} During the outbreak of SARS-CoV-2 in Taiwan, higher levels of depression were observed among people quarantined along with their families or friends and suspected of having a SARS-CoV-2 infection.¹⁰¹ A COVID-19-positive patient in the family or near one's family increased state anxiety scores in adolescents.⁶² Knowing a person who developed COVID-19 was a crucial factor influencing the stress and anxiety subscales.⁸⁸ The reason for this may be that a family member tested positive for COVID-19 requires isolation from their family. Separation from one's family may lead to an increased risk of the development of mental disorders in young people.¹⁰² In this study, the severity of anxiety and depression was significantly higher among quarantined versus non-quarantined respondents, which corresponds with the findings by Babicki *et al*, Tang *et al* and Mazza *et al*.^{49 103 104} As shown by Babicki *et al*, who conducted their study in November 2020, at the peak of the second wave of the pandemic in Poland, experiences with COVID-19 resulting from an illness of one's own or a family member, or death of a loved one were associated with a higher score in the GHQ. The study showed that 59.2% of the respondents met the GHQ criterion for minor mental disorders.⁴⁹

This study showed positive correlations between the severity of anxiety and depression and the feeling of loneliness. Increased loneliness rates along with an increase in the severity of depression and anxiety symptoms during the pandemic were also confirmed in studies in the Canadian population.⁸¹ The relationship between loneliness and depressive symptoms was demonstrated by Kiggore *et al*.⁹ A study using the three-item UCLA scale

showed that more than 50% of respondents aged ≥ 60 years complained of greater pandemic-related loneliness, which was associated with increased depression (62% vs 9%) and anxiety (57% vs 9%).¹⁰⁵ In a group of adolescents, a positive correlation was found between loneliness and state anxiety, and the trait anxiety.⁶²

LIMITATIONS

Our study has several limitations. This was a cross-sectional study, which may not reflect shifts in the mental state of the population over time. However, mental states change both with time and due to fluctuations in the environment around us. The research results cited in the paper were obtained at different time points during the pandemic. Therefore, there is a need to present the psychological effects of the COVID-19 pandemic over a longer and more forward-looking period of time. The results of several studies cited in the discussion section were obtained with the use of different scales assessing anxiety, depression and loneliness. The current data were collected through online surveys, which may have excluded potential respondents who lack internet access or are not knowledgeable about modern technologies. Also, 'no response' error may have occurred. However, there is evidence that online surveys have higher rates of self-disclosure on sensitive items when compared with computer-assisted telephone surveys¹⁰⁶ and a higher data reliability, which is probably due to reduced privacy concerns.¹⁰⁷

CONCLUSIONS

This study shows that the COVID-19 pandemic affects mental well-being of individuals as well as entire communities. It is necessary to identify those most vulnerable to loneliness, anxiety and depression during a crisis to assess health needs and proactively allocate resources during and after the pandemic. Our findings clearly demonstrated that loneliness, anxiety, depression and irritability are important factors that need to be taken into account in people tested positive for SARS-CoV-2, those under home quarantine, and those who consider their physical and mental health worse than in the pre-pandemic period. Younger age and a worse financial situation were the most important sociodemographic factors associated with higher loneliness, anxiety, depression and irritability. Loneliness was correlated with depression, anxiety and irritability scores. It is important to cater for the mental health of individuals during the COVID-19 pandemic and to promote psychological interventions to improve mental well-being in potentially vulnerable social groups.

Acknowledgements The study authors thank the participants who completed the survey.

Contributors Conception and design of the work—EK, AI and BD. Acquisition, analysis and interpretation of data for the work—EK, BK, BD and AI. Drafting the work and revising it critically for important intellectual content—EK, BD and RM. Final approval of the version to be published—EK, BK, BD, RM and AI. Agreement

to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved—EK, BK, BD, RM and AI. Guarantor - EK.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval It was voluntary for the respondents to answer the questionnaire, and they had the right to withdraw their undertaking of participation at any time. The study was approved by the Bioethics Committee of the Medical University of Warsaw (AKBE/232/2020). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Ewa Kobos <http://orcid.org/0000-0001-7231-8411>

REFERENCES

- 1 usnews.com. Poland reports first coronavirus case – health Minister. Available: <https://www.gov.pl/web/zdrowie/pierwszy-przypadek-koronawirusa-w-polsce> [Accessed 17 Dec 2020].
- 2 Ministry of Health (in Polish). Pierwszy przypadek koronawirusa w Polsce [First case of coronavirus in Poland]. Available: <https://www.reuters.com/article/health-coronavirus-poland-idINKBN20R0X7> [Accessed 17 Dec 2020].
- 3 Coronavirus: what are the lockdown measures across Europe? Available: <https://www.dw.com/en/coronavirus-what-are-the-lockdown-measures-across-europe/a-52905137> [Accessed 17 Dec 2020].
- 4 Wojtyński B, Goryński P. *Health status of Polish population and its determinants 2020*. Warsaw: National Institute of Public Health National Institute of Hygiene, 2020. <https://www.pzh.gov.pl/>
- 5 World Health Organization. WHO coronavirus disease (COVID-19) dashboard, 2021. Available: <https://covid19.who.int/> [Accessed 27 Dec 2021].
- 6 Peplau LA, Perlman D. *Loneliness: a sourcebook of current theory, research, and therapy*. New York: Wiley, 1982: 1–18.
- 7 Leigh-Hunt N, Bagguley D, Bash K, et al. An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public Health* 2017;152:157–71.
- 8 Cacioppo JT, Cacioppo S. The growing problem of loneliness. *Lancet* 2018;391:426.
- 9 Killgore WDS, Cloonan SA, Taylor EC, et al. Loneliness: a signature mental health concern in the era of COVID-19. *Psychiatry Res* 2020;290:113117.
- 10 Paul E, Bu F, Fancourt D. Loneliness and risk for cardiovascular disease: mechanisms and future directions. *Curr Cardiol Rep* 2021;23:68.
- 11 Valtorta NK, Kanaan M, Gilbody S, et al. Loneliness, social isolation and risk of cardiovascular disease in the English longitudinal study of ageing. *Eur J Prev Cardiol* 2018;25:1387–96.
- 12 Rico-Uribe LA, Caballero FF, Martín-María N, et al. Association of loneliness with all-cause mortality: a meta-analysis. *PLoS One* 2018;13:e0190033.
- 13 Holt-Lunstad J, Smith TB, Baker M, et al. Loneliness and social isolation as risk factors for mortality: a meta-analytic review. *Perspect Psychol Sci* 2015;10:227–37.
- 14 Hawkey LC, Cacioppo JT. Loneliness matters: a theoretical and empirical review of consequences and mechanisms. *Ann Behav Med* 2010;40:218–27.
- 15 Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res* 2020;288:112954.
- 16 Salari N, Hosseini-Far A, Jalali R, et al. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Global Health* 2020;16:57.
- 17 Tull MT, Edmonds KA, Scamaldo KM, et al. Psychological outcomes associated with Stay-at-Home orders and the perceived impact of COVID-19 on daily life. *Psychiatry Res* 2020;289:113098.
- 18 Xiang Y-T, Yang Y, Li W, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry* 2020;7:228–9.
- 19 Petzold MB, Plag J, Ströhle A. Umgang MIT psychischer Belastung bei Gesundheitsfachkräften Im Rahmen Der Covid-19-Pandemie. *Nervenarzt* 2020;91:417–21.
- 20 Wu W, Zhang Y, Wang P, et al. Psychological stress of medical staffs during outbreak of COVID-19 and adjustment strategy. *J Med Virol* 2020;92:1962–70.
- 21 Lei L, Huang X, Zhang S, et al. Comparison of prevalence and associated factors of anxiety and depression among people affected by versus people unaffected by quarantine during the COVID-19 epidemic in southwestern China. *Medical Science Monitor* 2020;26:e924609.
- 22 Stieger S, Lewetz D, Swami V. Emotional well-being under conditions of Lockdown: an experience sampling study in Austria during the COVID-19 pandemic. *J Happiness Stud* 2021;2:1–18.
- 23 Groarke JM, Berry E, Graham-Wisener L, et al. Loneliness in the UK during the COVID-19 pandemic: cross-sectional results from the COVID-19 psychological wellbeing study. *PLoS One* 2020;15:e0239698.
- 24 Hoffart A, Johnson SU, Ebrahimi OV. Loneliness and social distancing during the COVID-19 pandemic: risk factors and associations with psychopathology. *Front Psychiatry* 2020;11:589127.
- 25 Wickens CM, McDonald AJ, Elton-Marshall T, et al. Loneliness in the COVID-19 pandemic: associations with age, gender and their interaction. *J Psychiatr Res* 2021;136:103–8.
- 26 Courtet P, Olié E, Debien C, et al. Keep socially (but not physically) connected and carry on: preventing suicide in the age of COVID-19. *J Clin Psychiatry* 2020;81:20com13370.
- 27 Pitman A, Mann F, Johnson S. Advancing our understanding of loneliness and mental health problems in young people. *Lancet Psychiatry* 2018;5:955–6.
- 28 Erzen E, Çikrikci Özkan. The effect of loneliness on depression: a meta-analysis. *Int J Soc Psychiatry* 2018;64:427–35.
- 29 Martín-María N, Caballero FF, Lara E, et al. Effects of transient and chronic loneliness on major depression in older adults: a longitudinal study. *Int J Geriatr Psychiatry* 2021;36:76–85.
- 30 Cruwys T, Dingle GA, Haslam C, et al. Social group memberships protect against future depression, alleviate depression symptoms and prevent depression relapse. *Soc Sci Med* 2013;98:179–86.
- 31 Necho M, Tsehay M, Birkie M, et al. Prevalence of anxiety, depression, and psychological distress among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Int J Soc Psychiatry* 2021;67:892–906.
- 32 Luo M, Guo L, Yu M, et al. The psychological and mental impact of coronavirus disease 2019 (COVID-19) on medical staff and general public - A systematic review and meta-analysis. *Psychiatry Res* 2020;291:113190.
- 33 Castaldelli-Maia JM, Marziali ME, Lu Z, et al. Investigating the effect of national government physical distancing measures on depression and anxiety during the COVID-19 pandemic through meta-analysis and meta-regression. *Psychol Med* 2021;51:1–13.
- 34 Fan FC, Zhang SY, Cheng Y. Incidence of psychological illness after coronavirus outbreak: a meta-analysis study. *J Epidemiol Community Health* 2021;75:jech-2020-215927.
- 35 Bliźniowska-Kowalska KM, Halariś A, Wang S-C, et al. A review of the global impact of the COVID-19 pandemic on public mental health, with a comparison between the USA, Australia, and Poland with Taiwan and Thailand. *Med Sci Monit* 2021;27:e932220.
- 36 Kołodziejczyk A, Misiak B, Szcześniak D, et al. Coping styles, mental health, and the COVID-19 quarantine: a nationwide survey in Poland. *Front Psychiatry* 2021;12:625355.

- 37 Brooks SK, Webster RK, Smith LE, *et al.* The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020;395:912–20.
- 38 Yuksel D, McKee GB, Perrin PB, *et al.* Sleeping when the world locks down: correlates of sleep health during the COVID-19 pandemic across 59 countries. *Sleep Health* 2021;7:134–42.
- 39 Costi S, Paltrinieri S, Bressi B, *et al.* Poor sleep during the first peak of the SARS-CoV-2 pandemic: a cross-sectional study. *Int J Environ Res Public Health* 2021;18:306.
- 40 Rubin GJ, Wessely S. The psychological effects of quarantining a City. *BMJ* 2020;368:m313.
- 41 Casagrande M, Favieri F, Tambelli R, *et al.* The enemy who sealed the world: effects quarantine due to the COVID-19 on sleep quality, anxiety, and psychological distress in the Italian population. *Sleep Med* 2020;75:12–20.
- 42 Panda PK, Gupta J, Chowdhury SR, *et al.* Psychological and behavioral impact of Lockdown and quarantine measures for COVID-19 pandemic on children, adolescents and caregivers: a systematic review and meta-analysis. *J Trop Pediatr* 2021;67:fmaa122.
- 43 Li Y, Qin Q, Sun Q, *et al.* Insomnia and psychological reactions during the COVID-19 outbreak in China. *J Clin Sleep Med* 2020;16:1417–8.
- 44 Prati G, Mancini AD. The psychological impact of COVID-19 pandemic lockdowns: a review and meta-analysis of longitudinal studies and natural experiments. *Psychol Med* 2021;51:201–11.
- 45 Szczesniak D, Ciulkowicz M, Maciaszek J, *et al.* Psychopathological responses and face mask restrictions during the COVID-19 outbreak: results from a nationwide survey. *Brain Behav Immun* 2020;87:161–2.
- 46 Hamer K, Baran M. *The influence of COVID19 pandemic on emotions, attitudes and behaviors of Poles - a summary of 10 months of the pandemic.* Warsaw: Institute of Psychology at Polish Academy of Sciences, 2021: 1–20.
- 47 Juchnowicz D, Baj J, Forma A, *et al.* The outbreak of SARS-CoV-2 pandemic and the well-being of Polish students: the risk factors of the emotional distress during COVID-19 Lockdown. *J Clin Med* 2021;10:944.
- 48 Bartoszek A, Walkowiak D, Bartoszek A, *et al.* Mental well-being (depression, loneliness, insomnia, daily life fatigue) during COVID-19 related Home-Confinement-A study from Poland. *Int J Environ Res Public Health* 2020;17:7417.
- 49 Babicki M, Szewczykowska I, Mastalerz-Migas A. Mental health in the era of the second wave of SARS-CoV-2: a cross-sectional study based on an online survey among online Respondents in Poland. *Int J Environ Res Public Health* 2021;18:2522.
- 50 Zhang J, Lu H, Zeng H, *et al.* The differential psychological distress of populations affected by the COVID-19 pandemic. *Brain Behav Immun* 2020;87:49–50.
- 51 Holmes EA, O'Connor RC, Perry VH, *et al.* Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry* 2020;7:547–60.
- 52 ICC/ESOMAR (the world association for social, opinion and market research) International Code on market, opinion and social research. Available: <http://www.ofbor.pl/public/File/Miedzynaro> [Accessed 18 Dec 2021].
- 53 Kwiatkowska M, Rogoza R, Kwiatkowska K. Analysis of the psychometric properties of the revised UCLA loneliness scale in a Polish adolescent sample. *Current Issues in Personality Psychology* 2018;6:164–70.
- 54 Russell D, Peplau LA, Cutrona CE. The revised UCLA loneliness scale: concurrent and discriminant validity evidence. *J Pers Soc Psychol* 1980;39:472–80.
- 55 Austin BA. Factorial structure of the UCLA loneliness scale. *Psychol Rep* 1983;53:883–9.
- 56 Perry GR. Loneliness and coping among tertiary-level adult cancer patients in the home. *Cancer Nurs* 1990;13:293–302.
- 57 Majkowicz M. Practical assessment of palliative care efficacy: selected research techniques. In: de Walden-Galuszko K, Majkowicz M, eds. *Evaluation of palliative care quality in theory and practice.* Gdańsk, DC: Medical Academy, 2000: 26–31.
- 58 Zigmund AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- 59 Sokół-Szałowka M. Mental health impact of quarantine during the COVID-19 pandemic. *Psychiatry* 2021;18:57–62.
- 60 Yıldırım M, Kızılgöç M, Seçer İsmail, *et al.* Meaning in Life, Religious Coping, and Loneliness During the Coronavirus Health Crisis in Turkey. *J Relig Health* 2021;60:2371–85.
- 61 Emodi-Perlman A, Eli I, Uziel N, *et al.* Public concerns during the COVID-19 Lockdown: a multicultural cross-sectional study among Internet survey Respondents in three countries. *J Clin Med* 2021;10:1577.
- 62 Kılınçel Şenay, Kılınçel O, Muratdağı G, *et al.* Factors affecting the anxiety levels of adolescents in home-quarantine during COVID-19 pandemic in turkey. *Asia Pac Psychiatry* 2021;13:e12406.
- 63 Bu F, Steptoe A, Fancourt D. Loneliness during a strict lockdown: trajectories and predictors during the COVID-19 pandemic in 38,217 United Kingdom adults. *Soc Sci Med* 2020;265:113521.
- 64 Losada-Baltar A, Martínez-Huertas José Ángel, Jiménez-Gonzalo L, *et al.* Longitudinal correlates of loneliness and psychological distress during the lockdown situation due to COVID-19. Effects of age and self-perceptions of aging. *J Gerontol B Psychol Sci Soc Sci* 2021;gbab012.
- 65 Ahmed MZ, Ahmed O, Aibao Z, *et al.* Epidemic of COVID-19 in China and associated psychological problems. *Asian J Psychiatr* 2020;51:102092.
- 66 Qiu J, Shen B, Zhao M, *et al.* A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. *Gen Psychiatr* 2020;33:e100213.
- 67 Shanahan L, Steinhoff A, Bechtiger L, *et al.* Emotional distress in young adults during the COVID-19 pandemic: evidence of risk and resilience from a longitudinal cohort study. *Psychol Med* 2020:1–10.
- 68 Nagasu M, Muto K, Yamamoto I. Impacts of anxiety and socioeconomic factors on mental health in the early phases of the COVID-19 pandemic in the general population in Japan: a web-based survey. *PLoS One* 2021;16:e0247705.
- 69 Gambin M, Sękowski M, Woźniak-Prus M, *et al.* Generalized anxiety and depressive symptoms in various age groups during the COVID-19 lockdown in Poland. specific predictors and differences in symptoms severity. *Compr Psychiatry* 2021;105:152222.
- 70 Dziędzic B, Idzik A, Kobos E, *et al.* Loneliness and mental health among the elderly in Poland during the COVID-19 pandemic. *BMC Public Health* 2021;21:1976.
- 71 Victor CR, Yang K. The prevalence of loneliness among adults: a case study of the United Kingdom. *J Psychol* 2012;146:85–104.
- 72 Carmichael CL, Reis HT, Duberstein PR. In your 20S it's quantity, in your 30S it's quality: the prognostic value of social activity across 30 years of adulthood. *Psychol Aging* 2015;30:95–105.
- 73 Luhmann M, Hawkey LC. Age differences in loneliness from late adolescence to oldest old age. *Dev Psychol* 2016;52:943–59.
- 74 Nicolaisen M, Thorsen K. What are friends for? friendships and loneliness over the Lifespan-From 18 to 79 years. *Int J Aging Hum Dev* 2017;84:126–58.
- 75 Child ST, Lawton L. Loneliness and social isolation among young and late middle-age adults: associations with personal networks and social participation. *Aging Ment Health* 2019;23:196–204.
- 76 Elmer T, Mepham K, Stadfeld C. Students under lockdown: comparisons of students' social networks and mental health before and during the COVID-19 crisis in Switzerland. *PLoS One* 2020;15:e0236337.
- 77 Losada-Baltar A, Jiménez-Gonzalo L, Gallego-Alberto L, *et al.* "We Are Staying at Home." Association of Self-perceptions of Aging, Personal and Family Resources, and Loneliness With Psychological Distress During the Lock-Down Period of COVID-19. *J Gerontol B Psychol Sci Soc Sci* 2021;76:e10–16.
- 78 Matud MP. Gender differences in stress and coping styles. *Personality and Individual Differences* 2004;37:1401–15.
- 79 Matud MP, Bethencourt JM, Ibañez I. Gender differences in psychological distress in Spain. *Int J Soc Psychiatry* 2015;61:560–8.
- 80 Ptacek JT, Smith RE, Dodge KL. Gender differences in coping with stress: when stressor and appraisals do not differ. *Personality and Social Psychology Bulletin* 1994;20:421–30.
- 81 McQuaid RJ, Cox SML, Ogunlana A, *et al.* The burden of loneliness: implications of the social determinants of health during COVID-19. *Psychiatry Res* 2021;296:113648.
- 82 Shahriarirad R, Erfani A, Ranjbar K, *et al.* The mental health impact of COVID-19 outbreak: a nationwide survey in Iran. *Int J Ment Health Syst* 2021;15:19.
- 83 Kemperman A, van den Berg P, Weijs-Perrée M, *et al.* Loneliness of older adults: social network and the living environment. *Int J Environ Res Public Health* 2019;16:406.
- 84 Jaspal R, Breakwell GM. Socio-Economic inequalities in social network, loneliness and mental health during the COVID-19 pandemic. *Int J Soc Psychiatry* 2020:20764020976694.
- 85 Tomaz SA, Coffee P, Ryde GC, *et al.* Loneliness, wellbeing, and social activity in Scottish older adults resulting from social distancing during the COVID-19 pandemic. *Int J Environ Res Public Health* 2021;18:4517.
- 86 Lee C-YS, Goldstein SE. Loneliness, stress, and social support in young adulthood: does the source of support matter? *J Youth Adolesc* 2016;45:568–80.



- 87 Cao W, Fang Z, Hou G, *et al*. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res* 2020;287:112934.
- 88 Passavanti M, Argentieri A, Barbieri DM, *et al*. The psychological impact of COVID-19 and restrictive measures in the world. *J Affect Disord* 2021;283:36–51.
- 89 Zheng J, Morstead T, Sin N, *et al*. Psychological distress in North America during COVID-19: the role of pandemic-related stressors. *Soc Sci Med* 2021;270:113687.
- 90 Geirdal Amy Østertun, Ruffolo M, Leung J, *et al*. Mental health, quality of life, wellbeing, loneliness and use of social media in a time of social distancing during the COVID-19 outbreak. A cross-country comparative study. *J Ment Health* 2021;30:148–55.
- 91 Alzueta E, Perrin P, Baker FC, *et al*. How the COVID-19 pandemic has changed our lives: a study of psychological correlates across 59 countries. *J Clin Psychol* 2021;77:556–70.
- 92 Choi EPH, Hui BPH, Wan EYF. Depression and anxiety in Hong Kong during COVID-19. *Int J Environ Res Public Health* 2020;17:3740.
- 93 Franssen T, Stijnen M, Hamers F, *et al*. Age differences in demographic, social and health-related factors associated with loneliness across the adult life span (19–65 years): a cross-sectional study in the Netherlands. *BMC Public Health* 2020;20:1118.
- 94 Matthews T, Danese A, Caspi A, *et al*. Lonely young adults in modern Britain: findings from an epidemiological cohort study. *Psychol Med* 2019;49:268–77.
- 95 Feter N, Caputo EL, Doring IR, *et al*. Sharp increase in depression and anxiety among Brazilian adults during the COVID-19 pandemic: findings from the PAMPA cohort. *Public Health* 2021;190:101–7.
- 96 World Health Organization. Mental health and psychosocial considerations during the COVID-19 outbreak, 18 March 2020. World Health organization, 2020. Available: <https://apps.who.int/iris/handle/10665/331490> [Accessed 09 Aug 2021].
- 97 Zhou S-J, Zhang L-G, Wang L-L, *et al*. Prevalence and socio-demographic correlates of psychological health problems in Chinese adolescents during the outbreak of COVID-19. *Eur Child Adolesc Psychiatry* 2020;29:749–58.
- 98 Özdin S, Bayrak Özdin Şükriye. Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish Society: the importance of gender. *Int J Soc Psychiatry* 2020;66:504–11.
- 99 Moghanibashi-Mansourieh A. Assessing the anxiety level of Iranian general population during COVID-19 outbreak. *Asian J Psychiatr* 2020;51:102076.
- 100 Wang C, Pan R, Wan X, *et al*. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int J Environ Res Public Health* 2020;17:1729.
- 101 Ko C-H, Yen C-F, Yen J-Y, *et al*. Psychosocial impact among the public of the severe acute respiratory syndrome epidemic in Taiwan. *Psychiatry Clin Neurosci* 2006;60:397–403.
- 102 Golberstein E, Wen H, Miller BF. Coronavirus disease 2019 (COVID-19) and mental health for children and adolescents. *JAMA Pediatr* 2020;174:819–20.
- 103 Tang F, Liang J, Zhang H, *et al*. COVID-19 related depression and anxiety among quarantined respondents. *Psychol Health* 2021;36:164–78.
- 104 Mazza C, Ricci E, Biondi S, *et al*. A nationwide survey of psychological distress among Italian people during the COVID-19 pandemic: immediate psychological responses and associated factors. *Int J Environ Res Public Health* 2020;17:3165.
- 105 Kotwal AA, Holt-Lunstad J, Newmark RL, *et al*. Social isolation and loneliness among San Francisco Bay area older adults during the COVID-19 Shelter-in-Place orders. *J Am Geriatr Soc* 2021;69:20–9.
- 106 Milton AC, Ellis LA, Davenport TA, *et al*. Comparison of self-reported telephone interviewing and web-based survey responses: findings from the second Australian young and well national survey. *JMIR Ment Health* 2017;4:e37.
- 107 Braunsberger K, Wybenga H, Gates R. A comparison of reliability between telephone and web-based surveys. *Journal of Business Research* 2007;60:758–64.

Supplementary file 1. Study participant characteristics

Variable		n	%
Gender	Male	438	49.2
	Female	452	50.8
Education	Primary/junior high school	51	5.7
	Vocational	235	26.4
	Secondary	458	51.5
	Higher	146	16.4
Place of residence	Village	338	38.0
	City with less than 100,000 population	291	32.7
	City with more than 100,000 population	261	29.3
Employment status	Employed	463	52.0
	Retiree/pensioner	212	23.8
	Student	68	7.7
	Unemployed	79	8.9
	No response	68	7.6
Marital status	Unmarried	192	21.6
	Married	502	56.4
	In a relationship	100	11.2
	Divorced/separated	55	6.2
	Widow/widower	35	3.9
	No response	6	0.7
Type of accommodation	Single-family house, detached/semi-detached/terraced	458	51.5

	Multi-family house (block of flats)	418	46.9
	None of the above	14	1.6
Number of people per household	1	90	10.1
	2	252	28.3
	3	214	24.0
	4	186	20.9
	≥5	140	17.7
	None of the above	4	0.4
Person with whom one currently resides	Alone	92	10.3
	With spouse/partner only	246	27.7
	With family (children, relatives)	529	59.4
	No response	23	2.6
Type of pre-pandemic job	N/A, unemployed	255	28.6
	Work at the place of employment	446	50.1
	Remote work	57	6.4
	Remote work and work at the place of employment	80	9.0
	None of the above	52	5.9
Type of job performed during the pandemic	N/A, unemployed	310	34.8
	Work at the place of employment	398	44.7
	Remote work	62	7.0
	Remote work and work at the place of employment	93	10.5
	None of the above	27	3.0

Income	No income	59	6.6
	≤ 2000	213	24.0
	2001-3000	217	24.4
	3001-400	152	17.1
	4001-5000	79	8.8
	5001-6000	32	3.6
	6001-7000	29	3.3
	7001-10000	24	2.1
	No response	85	9,1
	Financial situation rating	Very good	40
Quite good		317	35.6
Neither good nor bad		392	44.0
Quite bad		106	11.9
Very bad		35	4.0

Supplementary file 2. HADS-M and R-UCLA scores with regard to health self-rating

Self-rated health		n	%	HADS-M		R-UCLA	
				R	p	R	p
Subjective rating of health during the pandemic							
Physical health	Very good	128	14.3	-.270	***	-.155	***
	Good	395	44.4				
	Neither good nor bad	264	29.7				
	Bad	89	10.0				
	Very bad	14	1.6				
Mental health	Very good	213	23.9	-.552	***	-.340	***
	Good	402	45.1				
	Neither good nor bad	183	21.0				
	Bad	65	7.4				
	Very bad	23	2.6				
Self-assessment of health compared to the pre-pandemic period							
Physical health	It has definitely improved	51	5.7	-.092	**	-.035	
	It has somewhat improved	107	12.0				
	It has not changed	598	67.3				
	It has somewhat deteriorated	116	13.0				
	It has definitely deteriorated	18	2.0				

Mental health	It has definitely improved	40	4.5	-.294	***	-.135	***
	It has somewhat improved	74	8.3				
	It has not changed	602	67.7				
	It has somewhat deteriorated	138	15.5				
	It has definitely deteriorated	36	4.0				
R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; R, Spearman test; p, statistical significance; **p < .01, ***p < .001, correlations were statistically significant							

Supplementary file 3. HADS-M and R-UCLA scores with regard to the subjects' education

HADS-M; R-UCLA	Primary/junior high school (n=51)		Vocational (n=235)		Secondary (n=458)		Higher (n=146)		F	p
	M	SD	M	SD	M	SD	M	SD		
Anxiety	8.58	5.08	7.47	4.61	6.97	4.74	7.12	4.58	2.096	
Depression	5.70	4.44	5.76	4.08	5.02	4.41	5.32	4.12	1.685	
Depression and Irritability	8.86	5.53	8.15	5.13	7.47	5.38	7.75	5.36	1.587	
HADS-M - total score	17.45	10.24	15.62	9.19	14.45	9.67	14.8	9.48	1.965	
Intimate others	25.84	7.61	22.14	6.51	22.38	6.85	21.71	6.37	5.066	**
Social others	8.94	2.96	8.56	3.28	8.33	3.11	8.58	3.27	.793	
Belonging and affiliation	9.66	3.04	9.61	2.99	9.48	2.92	9.43	2.94	.188	
R-UCLA - total score	44.45	10.70	40.32	10.59	40.20	10.82	39.73	10.88	2.648	*
<p>R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; F, Welcha test; p, statistical significance *p < .05, **p < .01, correlations were statistically significant</p>										

Supplementary file 4. HADS-M and R-UCLA scores with regard to the subjects' place of residence (n=890)

HADS-M; R-UCLA	R Spearman	p
Anxiety	-.056	
Depression	-.090	**
Depression and Irritability	-.102	**
HADS-M – total score	-.085	*
Intimate others	.019	
Social others	-.053	
Belonging and affiliation	-.016	
R-UCLA - total score	-.006	
<p>R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; R, Spearman test; p, statistical significance; *p < .05, **p < .01, correlations were statistically significant</p>		

Supplementary file 5. HADS-M and R-UCLA scores with regard to the subjects' employment status

HADS-M; R-UCLA	Employed (n=463)		Retiree/ pensioner (n=212)		Student (n=68)		Unemployed (n=79)		F	p
	M	SD	M	SD	M	SD	M	SD		
Anxiety	7.17	4.14	6.83	4.87	7.91	4.79	7.70	4.73	1.195	
Depression	5.24	5.20	5.06	4.70	5.08	4.18	5.75	4.01	.561	
Depression and Irritability	7.70	9.37	7.25	5.82	7.85	4.86	8.40	4.92	.976	
HADS-M – total score	14.87	6.90	14.08	10.32	15.76	9.14	16.11	9.15	1.098	
Intimate others	22.20	3.29	22.22	6.38	24.72	7.06	22.16	7.16	2.639	*
Social others	8.49	3.03	7.90	2.78	9.38	3.31	8.31	2.97	4.287	**
Belonging and affiliation	9.33	10.93	9.48	2.73	10.00	3.01	9.68	2.77	1.143	
R-UCLA - total score	40.03	4.14	39.61	10.22	44.10	11.32	40.16	10.92	2.938	*
<p>R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; F, Welcha test; p, statistical significance *p < .05, **p < .01, correlations were statistically significant</p>										

Supplementary file 6. HADS-M and R-UCLA scores with regard to the subjects' marital status

HADS-M; R-UCLA	In a relationship (n=602)		Single (n=282)		t	p
	M	SD	M	SD		
Anxiety	7.02	4.66	7.60	4.71	-1.740	
Depression	5.26	4.29	5.37	4.24	-.355	
Depression and Irritability	7.73	5.37	7.82	5.17	-.239	
HADS-M – total score	14.75	9.58	15.43	9.38	-.989	
Intimate others	21.29	6.41	24.70	6.95	-7.177	***
Social others	8.13	3.07	9.17	3.32	-4.537	***
Belonging and affiliation	9.29	2.87	9.97	3.07	-3.220	**
R-UCLA - total score	38.72	10.02	43.85	11.55	-6.745	***
R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; t, student's t-test; p, statistical significance; **p < .01, ***p < .001, correlations were statistically significant						

Supplementary file 7. Type of accommodation and the HADS and R-UCLA scores

HADS-M; R-UCLA	Multi-family house (block of flats) (n=418)		Single-family house, detached/semi- detached/terraced (n=458)		t	p
	M	SD	M	SD		
Anxiety	6.83	4.47	7.55	4.90	-2.247	*
Depression	4.98	4.21	5.58	4.35	-2.043	*
Depression and Irritability	7.21	5.19	8.25	5.40	-2.878	**
HADS-M – total score	14.05	9.16	15.80	9.88	-2.706	**
Intimate others	22.30	6.78	22.44	6.80	-.302	
Social others	8.44	3.23	8.48	3.13	-.172	
Belonging and affiliation	9.72	3.00	9.34	2.90	1.912	
R-UCLA - total score	40.48	11.01	40.27	10.67	.280	
R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; t, student's t-test; p, statistical significance; *p < .05, **p < .01, correlations were statistically significant						

Supplementary file 8. The number of people per household and the HADS and R-UCLA scores (n=886)

HADS-M; R-UCLA	R Spearman	p
Anxiety	.098	**
Depression	.064	
Depression and Irritability	.105	**
HADS-M – total score	.100	**
Intimate others	-.024	
Social others	-.004	
Belonging and affiliation	-.043	
R-UCLA - total score	-.015	
R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; R, Spearman test; p, statistical significance; **p < .01, correlations were statistically significant		

Supplementary file 9. People with whom the respondent lives and the HADS and R-UCLA scores

HADS-M; R-UCLA	With spouse/partner only (n=246)		With family (children, relatives) (n=529)		Alone (n=92)		F	p
	M	SD	M	SD	M	SD		
Anxiety	6.87	4.56	7.45	4.76	6.77	4.80	1.822	
Depression	5.18	4.42	5.41	4.23	5.01	4.37	.485	
Depression and Irritability	7.30	5.42	8.13	5.29	7.04	5.31	2.993	*
HADS-M – total score	14.15	9.54	15.58	9.55	13.81	9.72	2.632	
Intimate others	20.95	5.67	22.60	6.99	24.50	7.38	11.140	***
Social others	8.20	3.08	8.51	3.19	8.46	3.23	.840	
Belonging and affiliation	9.52	2.79	9.43	3.00	9.69	3.01	.315	
R-UCLA - total score	38.68	9.45	40.55	11.00	42.66	12.00	5.276	**
R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; F, Welcha test; p, statistical significance; *p < .05, **p < .01, ***p < .001, correlations were statistically significant								

Supplementary file 10. Financial situation/income and the HADS and R-UCLA scores

HADS-M; R-UCLA	Assessment of the financial situation (n=883)		Income (n=805)	
	R	p	R	p
Anxiety	-.221	.000***	-.117	**
Depression	-.251	.000***	-.069	*
Depression and Irritability	-.255	.000***	-.074	*
HADS-M – total score	-.249	.000***	-.096	**
Intimate others	-.162	.000***	-.080	*
Social others	-.137	.000***	-.067	
Belonging and affiliation	-.118	.000***	-.105	**
R-UCLA - total score	-.163	.000***	-.080	*

R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; R, Spearman test; p, statistical significance; *p < .05, **p < .01, ***p < .001, correlations were statistically significant

Supplementary file 11. Having family members/friends who were tested positive for SARS-CoV-2 and the HADS and R-UCLA scores

HADS-M; R-UCLA	No (n= 662)		Yes (n=210)		t	p
	M	SD	M	SD		
Anxiety	7.09	4.74	7.41	4.65	-.861	
Depression	5.30	4.38	5.08	3.98	.655	
Depression and Irritability	7.70	5.39	7.76	5.16	-.144	
HADS-M – total score	14.79	9.67	15.18	9.33	-.504	
Intimate others	22.52	6.77	21.69	6.45	1.570	
Social others	8.55	3.18	8.10	3.02	1.790	
Belonging and affiliation	9.70	2.91	8.91	2.92	3.411	***
R-UCLA - total score	40.78	10.74	38.71	10.83	2.425	*

R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; t, student's t-test; p, statistical significance; *p < .05, **p < .01, ***p < .001, correlations were statistically significant

Supplementary file 12. Having family members/friends who were/are in home quarantine or home/hospital isolation and the HADS and R-UCLA scores

HADS-M; R-UCLA	Yes (n=370)		No (n=501)		t	p
	M	SD	M	SD		
Anxiety	7.08	4.70	7.22	4.73	-.435	
Depression	4.88	4.22	5.52	4.32	-2.206	*
Depression and Irritability	7.38	5.34	7.94	5.32	-1.507	
HADS-M – total score	14.47	9.59	15.16	9.56	-1.053	
Intimate others	21.49	6.34	22.95	6.92	-3.183	**
Social others	7.98	2.98	8.80	3.21	-3.827	***
Belonging and affiliation	9.02	2.88	9.88	2.90	-4.360	***
R-UCLA - total score	38.49	10.71	41.63	10.74	-4.269	***
R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; t, student's t-test; p, statistical significance; *p < .05, **p < .01, ***p < .001, correlations were statistically significant						

Supplementary file 13. The type of pre-pandemic job and the HADS and R-UCLA scores

HADS-M; R-UCLA	Work at the place of employment (n=446)		N/A, unemployed (n=255)		Remote work (n=57)		Remote work and work at the place of employment (n=80)		F	p
	M	SD	M	SD	M	SD	M	SD		
Anxiety	7.29	4.87	6.80	4.50	9.12	4.31	7.22	4.46	4.388	**
Depression	5.22	4.23	5.01	4.38	7.21	4.28	5.70	4.30	4.332	**
Depression and Irritability	7.63	5.22	7.39	5.50	10.21	5.29	8.23	5.28	4.642	**
HADS-M – total score	14.96	9.62	14.20	9.57	19.33	9.19	15.46	9.24	4.811	**
Intimate others	21.99	6.80	22.43	6.62	23.07	6.22	23.78	7.41	1.663	
Social others	8.44	3.26	8.23	2.98	9.68	3.71	8.30	2.91	2.547	
Belonging and affiliation	9.33	2.99	9.80	2.91	10.12	3.21	9.28	2.73	2.214	
R-UCLA - total score	39.77	10.81	40.47	11.03	42.87	10.25	41.37	10.23	1.838	
<p>R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; F, Welch test; p, statistical significance; *p < .05, **p < .01, ***p < .001, correlations were statistically significant</p>										

Supplementary file 14. The type of job performed during the pandemic and the HADS and R-UCLA scores

HADS-M; R-UCLA	Work at the place of employment (n=398)		N/A, unemployed (n=310)		Remote work (n=62)		Remote work and work at the place of employment (n=93)		F	p
	M	SD	M	SD	M	SD	M	SD		
Anxiety	6.85	4.49	7.24	4.80	8.19	4.75	8.17	5.11	2.787	*
Depression	4.98	3.97	5.39	4.58	6.00	4.46	5.84	4.37	1.822	
Depression and Irritability	7.42	4.95	7.83	5.65	8.90	5.36	8.23	5.54	1.758	
HADS-M – total score	14.28	8.93	15.07	9.99	17.09	9.70	16.40	10.24	2.370	
Intimate others	21.88	6.69	22.64	6.66	23.58	6.69	22.88	7.63	1.678	
Social others	8.46	3.27	8.26	2.97	9.64	3.59	8.24	3.06	2.777	*
Belonging and affiliation	9.31	3.05	9.75	2.80	10.04	2.89	9.26	3.00	2.153	
R-UCLA - total score	39.67	10.68	40.66	10.93	43.27	10.20	40.39	11.18	2.301	

R-UCLA, Revised UCLA Loneliness Scale; HADS-M, Hospital Anxiety and Depression Scale; M, mean difference; SD, standard deviation; F, Welch test; p, statistical significance; *p < .05, correlations were statistically significant