



# BMJ Open Barriers to healthcare and self-reported adverse outcomes for autistic adults: a cross-sectional study

Mary Doherty <sup>1</sup>, Stuart Neilson,<sup>2</sup> Jane O'Sullivan,<sup>3</sup> Laura Carravallah,<sup>4</sup> Mona Johnson,<sup>5</sup> Walter Cullen,<sup>6</sup> Sebastian C K Shaw <sup>7</sup>

**To cite:** Doherty M, Neilson S, O'Sullivan J, *et al*. Barriers to healthcare and self-reported adverse outcomes for autistic adults: a cross-sectional study. *BMJ Open* 2022;**12**:e056904. doi:10.1136/bmjopen-2021-056904

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-056904>).

Received 29 August 2021  
Accepted 13 January 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Anaesthesia, Our Lady's Hospital, Navan, Meath, Ireland

<sup>2</sup>Independent Researcher, Cork, Ireland

<sup>3</sup>Department of Anaesthesia, Mater Private Hospital, Dublin, Leinster, Ireland

<sup>4</sup>Pediatrics and Human Development, and Medicine, Michigan State University-College of Human Medicine, East Lansing, Michigan, USA

<sup>5</sup>Patient Safety, NHS Digital, Leeds, UK

<sup>6</sup>Department of General Practice, School of Medicine, UCD, Dublin, Ireland

<sup>7</sup>Department of Medical Education, Brighton and Sussex Medical School, Brighton, UK

## Correspondence to

Dr Mary Doherty;  
drmdoherty@gmail.com

## ABSTRACT

**Objectives** Autistic people experience poor physical and mental health along with reduced life expectancy compared with non-autistic people. Our aim was to identify self-reported barriers to primary care access by autistic adults compared with non-autistic adults and to link these barriers to self-reported adverse health consequences.

**Design** Following consultation with the autistic community at an autistic conference, *Autescape*, we developed a self-report survey, which we administered online through social media platforms.

**Setting** A 52-item, international, online survey.

**Participants** 507 autistic adults and 157 non-autistic adults.

**Primary and secondary outcome measures** Self-reported barriers to accessing healthcare and associated adverse health outcomes.

**Results** Eighty per cent of autistic adults and 37% of non-autistic respondents reported difficulty visiting a general practitioner (GP). The highest-rated barriers by autistic adults were deciding if symptoms warrant a GP visit (72%), difficulty making appointments by telephone (62%), not feeling understood (56%), difficulty communicating with their doctor (53%) and the waiting room environment (51%). Autistic adults reported a preference for online or text-based appointment booking, facility to email in advance the reason for consultation, the first or last clinic appointment and a quiet place to wait. Self-reported adverse health outcomes experienced by autistic adults were associated with barriers to accessing healthcare. Adverse outcomes included untreated physical and mental health conditions, not attending specialist referral or screening programmes, requiring more extensive treatment or surgery due to late presentations and untreated potentially life-threatening conditions. There were no significant differences in difficulty attending, barriers experienced or adverse outcomes between formally diagnosed and self-identified autistic respondents. **Conclusions** Reduction of healthcare inequalities for autistic people requires that healthcare providers understand autistic perspectives, communication needs and sensory sensitivities. Adjustments for autism-specific needs are as necessary as ramps for wheelchair users.

## INTRODUCTION

Autism is a common neurodevelopmental condition affecting 1%–2% of the

## Strengths and limitations of this study

- Our study arose from a community-identified need to develop autism awareness training for healthcare providers and benefited from an autistic-led research team, including autistic medical doctors, using participatory methods.
- To date, this large cross-sectional study is the first to explore the associations between barriers to accessing healthcare and self-reported adverse health outcomes for autistic adults.
- As we used a convenience sample and self-report survey, generalisability of the data may be limited.
- As the initial pilot questionnaire was undertaken in the UK, we did not include issues specific to other healthcare systems, such as cost or insurance, in this study.

population.<sup>1</sup> While autism is lifelong and heterogeneous in presentation, most autistic people are adult, do not have intellectual disability and are likely to be undiagnosed.<sup>2</sup> Doctors may underestimate the number of autistic patients under their care.<sup>3 4</sup> Autistic adults have poor physical and mental health compared with the general population.<sup>5</sup> Most medical conditions are more prevalent in the autistic population,<sup>6 7</sup> including diabetes, hypertension and obesity.<sup>8</sup> Autistic people experience premature mortality.<sup>9–11</sup> Life expectancy is potentially reduced by 16–30 years, with increased mortality across almost all diagnostic categories.<sup>9</sup> In-hospital mortality is also increased.<sup>12</sup> Autistic people are over two times as likely to use emergency departments<sup>13</sup> and to die after attending emergency care and three times as likely to require inpatient admission.<sup>14</sup>

Alongside increased health needs, autistic people report a greater likelihood that these needs are unmet.<sup>13</sup> Pervasive, multi-factorial barriers to healthcare access are experienced.<sup>15</sup> Some are shared by other disabled people, but autistic patients experience additional autism-specific barriers.<sup>16</sup>

Patient–provider communication, sensory sensitivities, executive functioning/planning difficulties and prior negative experiences with healthcare providers are important barriers.<sup>17 18</sup>

In response to primary legislation<sup>19</sup> and statutory guidance,<sup>20</sup> the Royal College of General Practitioners developed an Autism Patient Charter.<sup>21</sup> This recommended: staff awareness and training; autism friendly environment; reasonable adjustments following disclosure or clinical suspicion of autism; patient-tailored communications and behaviour-sensitive accommodations.<sup>21</sup> Despite efforts to champion autism, proposals to formalise autism training<sup>18 22</sup> and specific awareness-raising interventions,<sup>21</sup> almost 40% of general practitioners (GPs) report no formal training in autism.<sup>22</sup> They also report limited confidence in managing autistic patients.<sup>22</sup> Greater autism awareness exists where GPs have personal knowledge of autism, either through a relative or friend on the autistic spectrum, or because they themselves are autistic.<sup>22</sup> Communication skills training for healthcare providers may be the most pressing need.<sup>4</sup> GPs<sup>22</sup> and hospital specialists<sup>3</sup> report difficulties communicating with autistic patients. Only 25% of primary healthcare providers reported high confidence in communicating with autistic adult patients or identifying and making necessary accommodations.<sup>4</sup>

This study primarily aimed to identify self-reported barriers to accessing primary healthcare faced by autistic adults with a focus on autism-specific communication, sensory issues and procedural considerations. Secondary aims included capturing self-reported adverse health outcomes and the associations between these and reported healthcare access barriers, adding a narrative frame to the existing evidence base around health disparities. This is to our knowledge the largest study of primary healthcare access barriers to date and benefits from a high degree of participatory design by the autistic community.

## METHODS

### Conception and design

Here, we present part of a larger cross-sectional study. This work was inspired by a quality improvement project designed to inform autism training for local healthcare providers as part of an ‘Autism Friendly Town’ initiative by AsIAM, Ireland’s National Autism Charity.<sup>23 24</sup> In 2018, MD attended *Autescape*,<sup>25</sup> an annual conference by and for autistic people. Participants of all ages are welcome at *Autescape*, including those who are non-speaking, have high support needs or require full-time care, although the majority of attendees typically have low to moderate support needs. While there, MD distributed a qualitative questionnaire entitled ‘What do you wish your GP knew about autism?’ MD reviewed the 75 responses and grouped these under broad themes. That project formed the inspiration and basis for the study reported in this paper. Using the data gathered at *Autescape*, MD developed an online survey to investigate barriers to primary

healthcare in a larger sample of autistic adults, compared with a non-autistic adult comparison group. Nine autistic adults assisted with refining the survey. The resulting survey contained a mix of quantitative questions and free comment boxes. Quantitative questions included yes–no responses, single-item and multiple-item selections from a list and Likert scales. We asked about specific barriers encountered accessing healthcare, reasons for delaying or avoiding a visit and difficulties booking, planning or waiting for a GP visit. We explored the challenges during a consultation, including communication, sensory and organisation issues as well as available social supports. We also explored the impact of such barriers including self-reported consequences of failure to access healthcare and the reasonable adjustments to standard care which facilitate access. We used Google Forms to host the survey.

### Piloting and refinement

We piloted the survey in 2018. Preliminary analysis revealed a recurring theme of total non-engagement with healthcare providers, despite expressed healthcare needs. Consequently, we altered the survey to add response options applicable to non-attenders. Our research team, comprising autistic and non-autistic GPs, experienced academics and other autistic individuals, adapted and refined the survey into its final 52-item form.

### Sampling, recruitment and data collection

Autistic adults were recruited using a convenience sampling approach, through Twitter, Facebook and the AsIAM website. We recruited non-autistic controls (without autistic children) through personal and professional contacts of research team members, local area groups and parenting groups on social media. Recruitment took place in August 2019. We provided participant information, with informed consent implied through subsequent completion of the questionnaire. We asked respondents, particularly those who were parents, to respond specifically about seeking healthcare for themselves. For those identifying as autistic, we asked if they were formally diagnosed or self-identified.

### Data analysis

We used the statistical package ‘R’ to assess significance of between-group associations using a test of proportions and a Wilcoxon-Mann-Whitney U test. Participants who skipped questions were omitted from the analyses of those questions. We intend to present our qualitative results elsewhere.

### Patient and public involvement

Our study was conducted by an autistic-led research team including autistic medical doctors, using participatory methods. In addition, nine autistic individuals assisted with developing and refining the survey into its final form.

**Table 1** Participant data

	Autistic	Non-autistic
Participants (n)	507	157
Age		
Median (range)	38 (17–73)	38 (18–70)
Gender		
Male	99 (20%)	16 (10%)
Female	311 (62%)	132 (85%)
Non-binary	83 (17%)	7 (5%)
Prefer not to say	9 (2%)	1 (1%)
Location		
UK	330 (65%)	67 (43%)
Ireland	77 (15%)	63 (40%)
North America	44 (9%)	20 (13%)
Other	56 (11%)	7 (4%)
Formal diagnosis of autism	77%	
By psychiatrist	25%	
By clinical psychologist	48%	
By multidisciplinary team	26%	
Age at diagnosis		
Median (range)	33 (2–67)	

## RESULTS

### Participants

We are reporting 664 responses to the online survey: 507 autistic adults and 157 non-autistic adults (table 1). Unless otherwise specified, results relate to primary care.

### Barriers to access

The most common reason for a GP visit was a physical condition or illness in both groups (86% vs 92%, n.s.). Autistic individuals were more likely to attend for mental health difficulties (61% vs 27%, difference 34%, 95% CI (25.2% to 42.3%),  $p < 0.001$ ). Twenty-two per cent of the autistic respondents usually attended for issues directly related to autism. Compared with 37% of non-autistic respondents, 80% of autistic respondents reported difficulty visiting a GP when needed (difference 43%, 95% CI (34.4% to 51.9%),  $p < 0.001$ ). While difficulty deciding if symptoms warrant a visit was a barrier for both groups (72% vs 65%, n.s.), the most notable difference related to difficulties using the telephone to book an appointment (62% vs 16%). Not feeling understood was a reason to avoid or delay for 56% of autistic respondents compared with 13% of non-autistic respondents. Difficulty communicating with the doctor during the appointment was a barrier for 53% of the autistic group but only 6% of non-autistic respondents. See online supplemental table 1 for specific barriers in order of frequency.

### Communication

Alongside difficulty using the telephone, not feeling understood and difficulty communicating with the

doctor, autistic respondents reported difficulty communicating with reception staff more often than non-autistic respondents (46% vs 8%, difference 38%, 95% CI (31.5% to 44.6%),  $p < 0.001$ ). Fifty nine per cent of autistic respondents reported difficulty communicating during a consultation ‘all the time’ or ‘frequently’ compared with 12% of non-autistic respondents ( $p < 0.001$ ). Seventy eight per cent of autistic adults reported that ‘anxiety makes it harder to communicate’

Autistic respondents reported avoiding the telephone (78%), voicemail (61%) and face-to-face verbal communication (30%). Forty one per cent reported that it is ‘easier for me to communicate in writing’ (table 2).

### Sensory processing

The waiting room environment was a barrier for 51% of autistic respondents, but only 8% of non-autistic respondents. Specific sensory barriers are detailed in table 3. Sensory issues made communication more difficult for 31% of the autistic group (see table 2). Only 10% of autistic respondents marked ‘none of the above’ to sensory questions compared with 71% of non-autistic respondents.

### Perceived stigma

Only 3% of autistic respondents stated they did not feel anxious going to the doctor, compared with 33% of non-autistic respondents (difference 30%, 95% CI (–37.7% to –21.8%)  $p < 0.001$ ). Autistic respondents reported being ‘concerned I won’t be taken seriously when I describe my symptoms’ (67%); worried about ‘wasting the doctor’s time’ (66%) and ‘being considered a hypochondriac’ (65%). They also reported difficulty ‘asking for help’ (63%) and ‘discussing mental health’ (59%). Autistic respondents reported that unusual behaviour or stimming elicited negative reactions from other patients (15%), reception staff (9%) or medical staff (7%) (online supplemental table 2).

### Planning and organising

Autistic respondents reported difficulties with summarising when describing medical problems, with 66% noting the ‘need to give the whole story and not leave anything out’ compared with 12% of non-autistic respondents (difference 54%, 95% CI (47.1% to 60.9%),  $p < 0.001$ ). Autistic respondents reported difficulties with organisation and planning for healthcare, including difficulties ‘making an appointment in advance’ (59%), ‘prioritising my health issues’ (58%) and ‘making changes to my lifestyle or habits’ (56%). Forty five per cent reported forgetting a medical appointment and 30% had attended on the wrong day (online supplemental table 3).

### Predictability and control

Autistic respondents reported more difficulty with uncertainty than non-autistic respondents. Particular difficulties included not knowing the wait duration (70% vs 30%, difference 40%, 95% CI (31.5% to 48.7%),  $p < 0.001$ ), what would happen during the consultation (63% vs 16%,

**Table 2** Communication barriers

	Autistic n (%)	Non-autistic n (%)	Difference (95% CI)	P value
<b>Reasons to avoid or delay GP visit (communication)</b>				
Difficulty using the telephone to book an appointment	314 (62)	25 (16)	46% (38.5% to 53.5%)	<0.001
Not feeling understood	283 (56)	21 (13)	42% (35.2% to 49.7%)	<0.001
Difficulty communicating with the doctor during the appointment	269 (53)	10 (6)	47% (40.5% to 52.9%)	<0.001
Difficulty communicating with the reception staff	235 (46)	13 (8)	38% (31.5% to 44.6%)	<0.001
<b>Communication preferences</b>				
Telephone generally avoided where possible	395 (78)	59 (38)	40% (31.5% to 49.1%)	<0.001
Voicemail generally avoided where possible	311 (61)	61 (39)	23% (13.3% to 31.6%)	<0.001
Verbal, face-to-face communication generally avoided where possible	152 (30)	12 (8)	22% (16.2% to 28.5%)	<0.001
It is easier for me to communicate in writing	208 (41)	13 (8)	33% (26.3% to 39.2%)	<0.001
<b>Communication challenges</b>				
Anxiety makes it harder to communicate	395 (78)	42 (23)	51% (42.9% to 59.4%)	<0.001
Sensory issues make communication more difficult	156 (31)	2 (1)	30% (24.7% to 34.3%)	<0.001
I need extra time to process what is being said	286 (56)	12 (8)	49% (42.4% to 55.2%)	<0.001
I can't describe my pain or symptoms accurately	272 (54)	36 (23)	31% (22.4% to 39.0%)	<0.001
Verbal communication is difficult	234 (46)	11 (7)	39% (32.8% to 45.5%)	<0.001
I express emotions differently for example, I can appear angry when I am afraid or in pain	227 (45)	6 (4)	41% (35.3% to 46.6%)	<0.001
I have difficulty prioritising when describing medical symptoms	333 (66)	34 (22)	44% (36.0% to 52.1%)	<0.001
I need to give the whole story and not leave anything out	332 (66)	18 (12)	54% (47.1% to 60.9%)	<0.001
None of the above (communication)	11 (2)	58 (37)	-35% (-42.8% to -26.7%)	<0.001

GP, general practitioner.

**Table 3** Sensory barriers

	Autistic n (%)	Non-autistic n (%)	Difference (95% CI)	P value
<b>Reasons to avoid or delay GP visit (sensory)</b>				
The waiting room environment	256 (51)	12 (8)	42.8% (36.4% to 49.3%)	<0.001
<b>Specific sensory challenges</b>				
Noise in the waiting room from other patients	319 (63)	19 (12)	51% (43.8% to 57.8%)	<0.001
Crowded waiting area	299 (59)	22 (14)	45% (37.6% to 52.3%)	<0.001
Bright or fluorescent lights	268 (53)	14 (9)	44% (37.3% to 50.6%)	<0.001
Uncomfortable furniture	195 (39)	11 (7)	32% (25.2% to 37.7%)	<0.001
Unexpected touch	193 (38)	9 (6)	32% (26.3% to 38.3%)	<0.001
Music playing in the waiting room	172 (34)	9 (6)	28% (22.3 to 34.1%)	<0.001
Smells in the waiting room	171 (34)	8 (5)	29% (22.9% to 34.4%)	<0.001
Touch during examination	160 (32)	11 (7)	25% (18.5% to 30.7%)	<0.001
Noise from the reception desk	140 (28)	4 (3)	25% (20.0% to 30.1%)	<0.001
Smells in the doctor's office	104 (21)	6 (4)	17% (11.7% to 21.7%)	<0.001
None of the above (sensory)	51 (10)	112 (71)	-61% (-69.2% to -53.3%)	<0.001

GP, general practitioner.

difference 47%, 95% CI (39.7% to 54.7%),  $p < 0.001$ ), which doctor they would see (58% vs 24%, difference 33%, 95% CI (25.0% to 41.8%),  $p < 0.001$ ) and the consultation length (40% vs 8%, difference 32%, 95% CI (25.6% to 38.4%)  $p < 0.001$ ).

### Support needs

Autistic adults reported physical mobility needs (16%) and unmet support needs in primary care, for example, 'needing a support person to come with me' (21%). This extended to secondary care: 17% had no one to support unexpected hospital admission, collection from hospital (20%) or home care following discharge (26%) (online supplemental table 4).

### Adverse consequences


Autistic respondents reported adverse consequences more frequently than non-autistic respondents, including untreated mental (69%) and physical (63%) health conditions. Notably, 60% were told they 'should have seen a doctor sooner' and 47% 'did not attend referral to a specialist'. Thirty-six per cent 'required more extensive treatment or surgery' and 34% did not access treatment for a 'potentially serious or life-threatening condition'. Additionally, they were less likely to 'attend on schedule for screening programmes' than the non-autistic respondents (39% vs 21%, difference 18%, 95% CI (9.8% to 26.2%),  $p < 0.001$ ) (figure 1).

Compared with autistic respondents who had no difficulty visiting a doctor, those who experienced difficulty (80%) reported more untreated mental and physical health conditions ( $p < 0.001$ ). They were also more likely to not attend specialist referral ( $p < 0.001$ ), to need more extensive treatment ( $p = 0.009$ ), to experience untreated life-threatening conditions ( $p = 0.006$ ) and to not attend screening ( $p = 0.028$ ) (figure 2). Four per cent of autistic respondents reported no access to primary healthcare


and did not attend any doctor at all. This group differed from the non-autistic respondents who reported no access to primary healthcare (5%) in two areas: all had difficulty visiting the doctor when needed, compared with 50% of non-attending non-autistic respondents ( $p = 0.002$ ); and 95% of autistic non-attenders had experienced at least one delayed treatment outcome, compared with 43% of non-attending, non-autistic respondents ( $p = 0.01$ ). There were no significant differences in difficulty attending a GP, the barriers experienced or adverse outcomes between formally diagnosed and self-identified autistic respondents.

### Facilitators

While most respondents (67% vs 65%) reported booking an appointment online would facilitate access, autistic patients selected a need to 'email my doctor in advance with a description of the issue I need to discuss' (62%), 'wait in a quiet place or outside until my turn' (56%) and 'book an appointment by text' (41%). Some autistic



**Figure 1** Adverse healthcare outcomes. For all comparisons between autistic and non-autistic groups  $p < 0.001$ .



**Figure 2** Adverse outcomes according to difficulty attending a GP. \*\* $p < 0.001$  \* $p < 0.05$ . y-axis=N. GP, general practitioner.

individuals would benefit if they ‘could book the first or last appointment’ (41%) or had a ‘sensory box available in the waiting room’ (16%) (online supplemental table 5).

Despite the outlined difficulties of visiting their doctor, autistic individuals felt that their relationship with their GP was ‘very important’ or ‘important’ significantly more than non-autistic respondents (70% vs 56%,  $p = 0.001$ ), but only 33% of autistic respondents reported a good relationship with their doctor ( $p < 0.001$ ). Only 62% of autistic individuals reported that their doctor knew they were autistic. Twenty two per cent were unsure, whereas 16% had not disclosed their diagnosis. Autistic respondents appreciated GPs who ask direct questions, give clear explanations, are honest about not understanding autism but know that autism is not a mental illness.

### Geographical variations

Some barriers to access for autistic respondents varied by geographical location (UK vs elsewhere in the world). Autistic respondents from the UK had more difficulty using the telephone to book appointments (66% vs 54%,  $p = 0.012$ ), more difficulty communicating with reception staff (52% vs 37%,  $p = 0.002$ ) and were less likely to experience no barriers to access at all (0.3% vs 2.8%,  $p = 0.038$ ). Autistic respondents from the UK also found it harder to see a known or preferred doctor (58% vs 29%,  $p < 0.001$ ), reported longer waits to get appointments (59% vs 33%,  $p < 0.01$ ) and found their online appointment booking systems more confusing (26% vs 10%,  $p < 0.001$ ). However, they were less likely to report no access to online booking systems compared with those from elsewhere in the world (26% vs 42%,  $p < 0.001$ ). There were no other significant differences for autistic respondents by geographical location.

There was only one significant geographical variation in self-reported adverse outcomes. For autistic respondents, those in the UK were less likely than those from elsewhere in the world to miss/not attend specialist referrals (43% vs 55%,  $p = 0.019$ ).

**Table 4** Barriers and outcomes associations for autistic respondents

	Untreated mental health condition	Untreated physical health condition	Specialist referral missed	Told they should have presented sooner	More extensive treatment or surgery required	Serious or life-threatening condition
Difficulty using the telephone to book an appointment	$p < 0.001$	$p < 0.001$	$p = 0.036$	$p = 0.015$	n.s.	$p = 0.007$
Difficulty with advance planning	$p = 0.020$	$p = 0.032$	$p < 0.001$	$p = 0.030$	n.s.	n.s.
Difficulty communicating with reception staff	$p < 0.001$	$p < 0.001$	$p = 0.008$	$p < 0.001$	$p = 0.041$	$p = 0.004$
Difficulty communicating with the doctor during the appointment	$p < 0.001$	$p < 0.001$	$p < 0.001$	$p = 0.003$	$p = 0.003$	$p = 0.002$
The waiting room environment	$p = 0.007$	$p < 0.001$	n.s.	$p = 0.010$	$p < 0.001$	n.s.
Inability to see a known or preferred doctor	$p < 0.001$	$p = 0.003$	n.s.	$p = 0.027$	n.s.	n.s.
Waiting to see the doctor is too difficult	$p = 0.018$	$p = 0.001$	n.s.	n.s.	$p = 0.026$	n.s.
Not feeling understood	$p < 0.001$	$p < 0.001$	$p < 0.001$	n.s.	n.s.	$p = 0.026$
Needing a support person to come with me	n.s.	$p = 0.002$	$p = 0.004$	n.s.	$p = 0.009$	$p = 0.002$

n.s., not significant.

## Associations between barriers and outcomes for autistic respondents

Table 4 outlines associations between reported barriers and outcomes for autistic respondents. There were no significant associations between any adverse outcomes and difficulty deciding if symptoms warrant a GP visit, not having an online booking system, having a confusing online booking system, having a long wait to get an appointment or having enough time to visit a doctor. In contrast, difficulty communicating with reception staff and difficulty communicating with the doctor during appointments were both significantly associated with all adverse outcomes.

Difficulty using the telephone to book an appointment was significantly associated with all adverse outcomes apart from having to undergo more extensive treatment or surgery than if they had attended sooner. Challenges with the waiting room environment were significantly associated with all adverse outcomes apart from missing specialist referrals and having a potentially serious or life-threatening condition for which they did not access treatment. Difficulty planning an appointment in advance was significantly associated with all adverse outcomes apart from having a potentially serious or life-threatening condition for which they did not access treatment, and having to undergo more extensive treatment or surgery than if they had attended sooner. Needing a support person to attend appointments was significantly associated with all adverse outcomes apart from having had a mental health condition remain untreated due to difficulties accessing healthcare and being told they should have presented sooner. Not feeling understood was significantly associated with all adverse outcomes apart from being told they should have presented sooner and having to undergo more extensive treatment or surgery than if they had attended sooner.

The inability to see a known or preferred doctor was significantly associated with having both untreated mental and physical health conditions. It was also significantly associated with being told they should have presented sooner. Finding waiting to see a doctor too difficult was significantly associated with having both untreated mental and physical health conditions. It was also significantly associated with having to undergo more extensive treatment or surgery than if they had attended sooner.

Reporting no barriers to access healthcare had no significant associations with any of the adverse outcomes.

## DISCUSSION

Our study describes the results of a survey of autistic adults and compares their experiences with non-autistic adults. It highlights barriers faced by autistic people accessing and engaging with primary healthcare. In our study, these included greater difficulties deciding when to seek care, reluctance to bother their GP, difficulties planning appointments and greater communication difficulties—with particular emphasis

on telephone use. Communication was also impaired by anxiety and sensory issues. We linked those barriers to self-reported adverse outcomes. Our data indicated that autistic people may present for healthcare later in the natural course of an illness. Autistic participants reported reduced attendance for screening, late presentations, missed opportunities for early detection and more extensive therapy being required. They also delayed or avoided healthcare because they did not feel understood by their doctors. Furthermore, a substantial minority of autistic adults did not disclose their autism diagnosis, which may impede identification of their autism-specific needs. These barriers may have real consequences, as evidenced in reduced life expectancy, and higher levels of physical and mental health conditions among autistic people.


## Comparison with the existing literature

This study confirms the findings of Vohra,<sup>14</sup> Raymaker<sup>16</sup> and several recent reviews,<sup>15 17 26</sup> which all identified three groups of barriers: (1) patient-level factors; (2) provider-level factors and (3) system-level factors. Our study stratifies individual barriers from the perspective of autistic individuals. We couple these barriers to self-reported adverse consequences, highlighting factors that may lead to excess morbidity and mortality in the autistic population.

## Strengths and limitations

Our study arose from a community-identified need to develop autism awareness training for healthcare providers. It benefited from an autistic-led research team, including autistic doctors, using participatory methods. Our study provided a unique picture of autistic adults' healthcare experiences, including those entirely excluded from healthcare due to access barriers. In particular, we highlighted the difficulties with using the telephone which is a distilled, concentrated essence of verbal communication.

As we used a convenience sample and self-report survey, generalisability of the data may be limited. Respondents required the ability to complete the survey, which excluded those with reduced ability to self-report. While we did not set out to create a validated tool, our survey may have benefited from some validity and reliability testing. As the initial quality improvement questionnaire was undertaken in the UK, we did not include issues specific to other healthcare systems, such as cost or insurance. Our analyses did not account for potential confounding factors, such as ethnicity or socioeconomic status. Female participants were over-represented in both groups, which is not unusual for online surveys, but it is interesting given the higher rate of autism diagnosis in men. While we noted significant gender differences in relation to non-binary participants, these participants were almost all autistic and we were, therefore, unable to attribute differences to gender identity or autism with



**Figure 3** Autism friendly general practice.

any degree of certainty. Furthermore, as this is a cross-sectional study, while we can identify associations, we cannot confirm causality.

### Implications for research

Our study suggests a need for personalised approaches to healthcare access. A prior study investigated using a previsit telephone call to identify individualised accommodations.<sup>27</sup> Our data suggest that this could be problematic for autistic adults. The AASPIRE Healthcare Toolkit<sup>28</sup> includes a publicly available online programme, which generates a computerised report of required healthcare accommodations. Adaptation of such a toolkit in NHS General Practice should be considered and researched. Social care interventions and healthcare facilitators in general practice have shown benefit with a vulnerable population,<sup>29</sup> similar approaches could benefit an autistic population. The significant difficulties among the small number of autistic people not registered with any GP indicate a need for further research into this group.

### Implications for clinical practice

Based on online supplemental table 5 and the lived experience of the autistic members of our research team, figure 3 outlines our proposed elements of an autism friendly practice. Such adjustments may minimise anxiety, manage sensory issues and ensure mutual understanding—promoting clear, unambiguous communication. Autism friendly practices should employ a personalised approach, with a healthcare access needs assessment and, where possible, a specialist liaison nurse or facilitator.

### Implications for policy

Given the identified barriers, the extension of annual health checks to autistic adults<sup>30 31</sup> and the recently announced Oliver McGowan Mandatory Training in Learning Disability and Autism<sup>32</sup> are welcome. These will likely bring important benefits provided they are informed by the autistic community and autistic healthcare providers. Autism registers in GP practice have been recommended.<sup>33 34</sup> The success of such initiatives will likely depend on greater awareness by medical practitioners of autistic culture and communication needs. Specific training for GPs during core training and continuing professional development may be beneficial. GPs with a special interest in autism should be facilitated to develop their skills, but management of general health needs and co-occurring conditions fall within the remit of every GP. Implementing existing autism legislation or development where lacking is required in order to reduce health inequities for autistic people.

### CONCLUSIONS

Autistic people face barriers accessing the healthcare system, followed by difficulties interacting with healthcare providers, which may contribute to known healthcare disparities, including increased morbidity and mortality. Our study has highlighted a variety of specific barriers to accessing primary healthcare for autistic adults, including use of the telephone to book appointments, not feeling understood, and difficulty communicating with doctors as well as sensory and organisational issues, which impede healthcare access. We identified a variety of significant associations between self-reported barriers to healthcare access and adverse outcomes for autistic respondents. One of our most impactful findings was the lack of any significant differences between formally diagnosed and self-identified autistic respondents in difficulty attending a GP, barriers experienced or self-reported adverse healthcare outcomes. Progress towards eliminating healthcare disparities for autistic people may be achieved by understanding the healthcare experiences and access barriers for this vulnerable patient group. These barriers represent not so much a failure to deliver or to avail of healthcare, but a lack of intersection between the communication patterns of autistic healthcare users and non-autistic providers. Reasonable accommodations are legally<sup>35</sup> and morally required. Adjustments for communication needs are as necessary for autistic people as ramps for wheelchair users.

**Twitter** Mary Doherty @AutisticDoctor and Sebastian C K Shaw @Autistic\_Doc

**Acknowledgements** We are indebted to Professor Louise Gallagher for her guidance during the early stages of this project. We acknowledge the input received from the autistic adult community recruited via local groups and online contacts during the development of the online survey. Assistance with content, structure and proofreading of the surveys was received from nine autistic adults in Ireland and the UK. We received assistance from members of peer support group 'Autistic Doctors International'. We also thank Dr David Hillebrandt, Dr Natalie Teasdale, Elaine McGoldrick and Karen Leneh Buckle for their assistance during this project.



**Contributors** Conception and design of study: MD, SN, JO'S, with contributions from those listed in acknowledgements. Acquisition of data: MD, SN, JO'S. Analysis of data: SN, MD, SCKS. Interpretation of data: MD, SN, JO'S, LC, MJ, WC, SCKS. Drafting and revising the manuscript: MD, SN, JO'S, LC, MJ, WC, SCKS. Approval of the version of the manuscript to be published: MD, SN, JO'S, LC, MJ, WC, SCKS. MD is the Guarantor of this work.

**Funding** We are grateful to AsIAM, Ireland's National Autism Charity and Scally's SuperValu, Clonakilty, for the funding to enable open access publication.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was approved by SJH/TUH Joint Research and Ethics Committee REC: 2019-06 Chairman's Action (17). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. All data relevant to the study are included in the article or uploaded as supplementary information.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Mary Doherty <http://orcid.org/0000-0002-6995-3745>

Sebastian C K Shaw <http://orcid.org/0000-0001-9597-7436>

## REFERENCES

- 1 Data CDC. Statistics. autism spectrum disorder. Resource document. Available: <https://www.cdc.gov/ncbddd/autism/data.html>
- 2 Royal College of Psychiatrists. The psychiatric management of autism in adults (CR228). Available: <https://www.rcpsych.ac.uk/improving-care/campaigning-for-better-mental-health-policy/college-reports/2020-college-reports/cr228> [Accessed 24 Jul 2021].
- 3 Zerbo O, Massolo ML, Qian Y, et al. A study of physician knowledge and experience with autism in adults in a large integrated healthcare system. *J Autism Dev Disord* 2015;45:4002–14.
- 4 Nicolaidis C, Schnider G, Lee J, et al. Development and psychometric testing of the AASPIRE adult autism healthcare provider self-efficacy scale. *Autism* 2021;25:767–73.
- 5 Ryzewska E, Hughes-McCormack LA, Gillberg C, et al. General health of adults with autism spectrum disorders – a whole country population cross-sectional study. *Res Autism Spectr Disord* 2019;60:59–66.
- 6 Croen LA, Zerbo O, Qian Y, et al. The health status of adults on the autism spectrum. *Autism* 2015;19:814–23.
- 7 Xue Ming, Brimacombe M, Chaaban J, et al. Autism spectrum disorders: concurrent clinical disorders. *J Child Neurol* 2008;23:6–13.
- 8 Flygare Wallén E, Ljunggren G, Carlsson AC, et al. High prevalence of diabetes mellitus, hypertension and obesity among persons with a recorded diagnosis of intellectual disability or autism spectrum disorder. *J Intellect Disabil Res* 2018;62:269–80.
- 9 Hirvikoski T, Mittendorf-Rutz E, Boman M, et al. Premature mortality in autism spectrum disorder. *Br J Psychiatry* 2016;208:232–8.
- 10 Hwang YJ, Srasuebku P, Foley K-R, et al. Mortality and cause of death of Australians on the autism spectrum. *Autism Res* 2019;12:806–15.
- 11 Bilder D, Botts EL, Smith KR, et al. Excess mortality and causes of death in autism spectrum disorders: a follow up of the 1980s Utah/UCLA autism epidemiologic study. *J Autism Dev Disord* 2013;43:1196–204.
- 12 Akobirshoev I, Mitra M, Dembo R, et al. In-hospital mortality among adults with autism spectrum disorder in the United States: a retrospective analysis of US hospital discharge data. *Autism* 2020;24:177–89.
- 13 Nicolaidis C, Raymaker D, McDonald K, et al. Comparison of healthcare experiences in autistic and non-autistic adults: a cross-sectional online survey facilitated by an academic-community partnership. *J Gen Intern Med* 2013;28:761–9.
- 14 Vohra R, Madhavan S, Sambamoorthi U. Emergency department use among adults with autism spectrum disorders (ASD). *J Autism Dev Disord* 2016;46:1441–54.
- 15 Walsh C, Lydon S, O'Dowd E, et al. Barriers to healthcare for persons with autism: a systematic review of the literature and development of a taxonomy. *Dev Neurorehabil* 2020;8:1–8.
- 16 Raymaker DM, McDonald KE, Ashkenazy E, et al. Barriers to healthcare: instrument development and comparison between autistic adults and adults with and without other disabilities. *Autism* 2017;21:972–84.
- 17 Mason D, Ingham B, Urbanowicz A, et al. A systematic review of what barriers and facilitators prevent and enable physical healthcare services access for autistic adults. *J Autism Dev Disord* 2019;49:3387–400.
- 18 Vogan V, Lake JK, Tint A, et al. Tracking health care service use and the experiences of adults with autism spectrum disorder without intellectual disability: a longitudinal study of service rates, barriers and satisfaction. *Disabil Health J* 2017;10:264–70.
- 19 Autism Act UK. HM Government, 2009. Available: <https://www.legislation.gov.uk/ukpga/2009/15/contents>
- 20 Department of Health. *Implementing fulfilling and rewarding lives. Statutory guidance for local authorities and NHS organisation to support implementation of the autism strategy*. London: Department of Health, 2010.
- 21 Buckley C. Making your practice autism friendly. *InnovAiT* 2017;10:327–31.
- 22 Unigwe S, Buckley C, Crane L, et al. GPs' confidence in caring for their patients on the autism spectrum: an online self-report study. *Br J Gen Pract* 2017;67:e445–52.
- 23 AsIAM. A first for Ireland with Clonakilty becoming Ireland's first autism friendly town, 2016. Available: <https://asiam.ie/clonakilty-autism-friendly-town/>
- 24 AsIAM. Are you ready to make your Clonakilty commitment for Autism?, 2020. Available: <https://asiam.ie/asiam-public-sector-training/autism-friendly-communities/>
- 25 The Autscope Organisation. Autscope 2018: exploring inclusion. Tonbridge, Kent, 2018. Available: <http://www.autscope.org/2018/>
- 26 Bradshaw P, Pellicano E, van Driel M, et al. How can we support the healthcare needs of autistic adults without intellectual disability? *Curr Dev Disord Rep* 2019;6:45–56.
- 27 Saqr Y, Braun E, Porter K, et al. Addressing medical needs of adolescents and adults with autism spectrum disorders in a primary care setting. *Autism* 2018;22:51–61.
- 28 Nicolaidis C, Raymaker D, McDonald K, et al. The development and evaluation of an online healthcare toolkit for autistic adults and their primary care providers. *J Gen Intern Med* 2016;31:1180–9.
- 29 Abel J, Kingston H, Scally A, et al. Reducing emergency hospital admissions: a population health complex intervention of an enhanced model of primary care and compassionate communities. *Br J Gen Pract* 2018;68:e803–10.
- 30 National Institute for Health and Care Excellence. Chapter 3: Learning disability and Autism. In: *Long term plan [London]*, 2019. <https://www.longtermplan.nhs.uk/online-version/chapter-3-further-progress-on-care-quality-and-outcomes/a-strong-start-in-life-for-children-and-young-people/learning-disability-and-autism/>
- 31 Harper G, Smith E, Parr J. Autistica action briefing: health checks. Available: <https://www.autistica.org.uk/downloads/files/Autistica-Action-Briefing-Health-Checks.pdf>
- 32 Health Education England. Partners announced to deliver the Oliver McGowan mandatory learning disability and autism training for all health and social care staff, 2020. Available: <https://www.hee.nhs.uk/news-blogs-events/news/partners-announced-deliver-oliver-mcgowan-mandatory-learning-disability-autism-training-all-health>
- 33 National Institute for Health and Care Excellence. The practice establishes and maintains a register of all patients with a diagnosis of autism [London] ([NM153]), 2017. Available: <https://www.nice.org.uk/standards-and-indicators/qofindicators/the-practice-establishes-and-maintains-a-register-of-all-patients-with-a-diagnosis-of-autism>



34 Westminster Commission on Autism. A spectrum of obstacles an inquiry into access to healthcare for autistic people. Available:

[https://westminsterautismcommission.files.wordpress.com/2016/03/ar1011\\_ncg-autism-report-july-2016.pdf](https://westminsterautismcommission.files.wordpress.com/2016/03/ar1011_ncg-autism-report-july-2016.pdf)

35 Great Britain. *Equality act 2010*. London: Stationary Office, 2010.

Table S1 Access Barriers to Healthcare

**Which of the following would cause you to delay or avoid seeing your doctor when you need to?**

	<b>Autistic n (%)</b>	<b>Non-autistic n (%)</b>	<b>Difference (95% Confidence Interval)</b>	<b>p value</b>
Difficulty deciding if symptoms warrant a GP visit	366 (72%)	102 (65%)	7% (CI -1.6%,16.1%)	p=0.102 ns
Difficulty using the telephone to book appointment	314 (62%)	25 (16%)	46% (CI 38.5%,53.5%)	p<0.001
Not feeling understood	283 (56%)	21 (13%)	42% (CI 35.2%,49.7%)	p<0.001
Difficulty communicating with the doctor during the appointment	269 (53%)	10 (6%)	47% (CI 40.5, 52.9%)	p<0.001
The waiting room environment	256 (51%)	12 (8%)	43% (CI 36.4%,49.3%)	p<0.001
Long wait to get an appointment	251 (50%)	70 (45%)	5% (CI -4.4%,14.2%)	p=0.324 ns
Difficulty planning an appointment in advance	243 (48%)	51 (33%)	15% (CI 6.5%,24.4%)	p<0.001
Inability to see a known or preferred doctor	241 (48%)	35 (23%)	25% (CI 17.0%,33.5%)	p<0.001
Difficulty communicating with the reception staff	235 (46%)	13 (8%)	38% (CI 31.5%,44.6%)	p<0.001
Not having enough time to visit the doctor	174 (34%)	61 (39%)	-5% (CI -13.6%,4.6%)	p=0.346 ns
No online booking system	160 (32%)	35 (23%)	9% (CI 1.2%,17.3%)	p=0.033
Waiting to see the doctor is too difficult	114 (23%)	7 (5%)	18% (CI 12.7%,23.3%)	p<0.001
Needing a support person to come with me	106 (21%)	7 (5%)	16% (CI 11.2%,21.7%)	p<0.001
There is an online booking system but it's confusing	102 (20%)	9 (6%)	14% (CI 8.9%,19.8%)	p<0.001
Not having anyone to look after my child	66 (13%)	18 (12%)	2% (CI -4.6%,7.8%)	p=0.708 ns
None of the above	6 (1%)	17 (11%)	-10% (CI -15.0%,-4.3%)	p<0.001

**Table S2 Perceived Stigma**

<b><u>Going to the doctor, I am anxious :</u></b>	<b>Autistic n (%)</b>	<b>Non-autistic n (%)</b>	<b>Difference (95% Confidence Interval)</b>	<b>p value</b>
I won't be taken seriously when I describe my symptoms	341 (67%)	54 (34%)	32.9% (CI 24.0%,41.8%)	p<0.001
I might be wasting the doctor's time	333 (66%)	53 (34%)	31.9% (CI 23.0%,40.8%)	p<0.001
I might be considered a hypochondriac about asking for help	330 (65%)	42 (27%)	38.3% (CI 29.8%,46.8%)	p<0.001
about discussing mental health	318 (63%)	28 (18%)	44.9% (CI 37.2%,52.6%)	p<0.001
that there might be something wrong	301 (59%)	37 (24%)	35.8% (CI 27.5%,44.1%)	p<0.001
I don't feel anxious going to the doctor	217 (43%)	55 (35%)	7.8% (CI -1.3%,16.8%)	p=0.102 ns
	17 (3%)	52 (33%)	-30% (CI -37.7%,-21.8%)	p<0.001

<b><u>Is stimming* a problem for you at the doctors' office?</u></b>	<b>Autistic n (%)</b>	<b>Non-autistic n (%)</b>	<b>Difference (95% Confidence Interval)</b>	<b>p value</b>
Unusual behaviours or stimming elicit negative reactions from other patients	74 (14.6%)	4 (2.5%)	12% (CI 7.7%,16.4%)	p<0.001
Unusual behaviours or stimming elicit negative reactions from reception staff	44 (8.7%)	1 (0.6%)	8% (CI 4.9%,11.2%)	p=0.001
Unusual behaviours or stimming elicit negative reactions from medical staff	37 (7.3%)	2 (1.3%)	6% (CI 2.7%,9.3%)	p=0.009
I feel comfortable with stimming at the doctors' office	101 (19.9%)	74 (47.1%)	-27% (CI -36.2%,-18.2%)	p<0.001
I do not feel comfortable with stimming at the doctors' office	4 (0.8%)	5 (3.2%)	-2% (CI -5.7%,0.9%)	p=0.061 ns
No need to stim at the doctors' office	133 (26.2%)	0 (0.0%)	26% (CI 22.0%,30.5%)	p<0.001
I don't understand the term "stimming"	29 (5.7%)	62 (39.5%)	-34% (CI -42.1%,-25.4%)	p<0.001

\*Stimming refers to repetitive motor movements or vocalisations which are commonly used by autistic people as a self-regulatory mechanism.

**Table S3 Planning and Organising**

	<b>Autistic n (%)</b>	<b>Non-autistic n (%)</b>	<b>Difference (95% Confidence Interval)</b>	<b>p value</b>
I find it difficult to prioritise when describing my medical problems	333 (66%)	34 (22%)	44% (CI 36.0%,52.1%)	p<0.001
I need to give the whole story and not leave anything out	332 (66%)	18 (12%)	54% (CI 47.1%,60.9%)	p<0.001
I find it difficult to make appointments in advance	300 (59%)	45 (29%)	31% (CI 21.8%,39.2%)	p<0.001
Making changes to my lifestyle or habits is difficult for me	282 (56%)	36 (23%)	33% (CI 24.4%,41.0%)	p<0.001
I have forgotten to attend a medical appointment	230 (45%)	34 (22%)	24% (CI 15.5%,31.9%)	p<0.001
I need to write things down	227 (45%)	17 (11%)	34% (CI 27.0%,40.9%)	p<0.001
I find waiting difficult	221 (44%)	15 (10%)	34% (CI 27.3%,40.8%)	p<0.001
I have difficulty making decisions about my health	222 (44%)	20 (13%)	31% (CI 23.9%,38.2%)	p<0.001
I have turned up for a medical appointment on the wrong day	151 (30%)	16 (10%)	20% (CI 13.0%,26.2%)	p<0.001
It is difficult to arrange someone to come with me	103 (20%)	2 (1%)	19% (CI 14.7%,23.4%)	p<0.001
I have forgotten why I made the appointment	54 (11%)	1 (1%)	10% (CI 6.6%,13.4%)	p<0.001
None of the above	13 (3%)	64 (41%)	-38% (CI -46.4%,-30.0%)	p<0.001

**Table S4 Support Needs**

<b><u>Do you visit your doctor:</u></b>	<b>Autistic n (%)</b>	<b>Non-autistic n (%)</b>	<b>Difference (95% Confidence Interval)</b>	<b>p value</b>
Alone, by choice	306 (60%)	134 (85%)	-25% (CI -32.4%,-17.6%)	p<0.001
Alone, but would prefer to have a support person	165 (33%)	10 (6%)	26% (CI 20.2%,32.2%)	p<0.001
With a parent, partner or support person	137 (27%)	9 (6%)	21% (CI 15.6%,27.0%)	p<0.001
With a parent, partner or support person but I would prefer to go alone	12 (2%)	0 (0%)	2% (CI 0.6%,4.1%)	p=0.109 ns
With a support animal	5 (1%)	0 (0%)	1% (CI -0.3%,2.3%)	p=0.471 ns
To support an autistic adult	26 (5%)	0 (0%)	5% (CI 2.8%,7.5%)	p=0.008
As a parent to access healthcare for my child	78 (15%)	30 (19%)	-4% (CI -11.0%,3.6%)	p=0.327 ns
With my child, but I would prefer to go alone	6 (1%)	12 (8%)	-7% (CI -11.1%,-1.8%)	p<0.001

<b><u>If you were suddenly admitted to hospital, who would be able to bring your personal belongings to you?</u></b>	<b>Autistic n (%)</b>	<b>Non-autistic n (%)</b>	<b>Difference (95% Confidence Interval)</b>	<b>p value</b>
Spouse or partner	234 (46%)	106 (68%)	-21% (CI -30.3%,-12.4%)	p<0.001
Parent	168 (33%)	70 (45%)	-11% (CI -20.7%,-2.2%)	p=0.012
Other family member	96 (19%)	72 (46%)	-27% (CI -35.8%,-18.0%)	p<0.001
Friend	118 (23%)	57 (36%)	-13% (CI -21.8%,-4.2%)	p=0.002
Neighbour	17 (3%)	14 (9%)	-6% (CI -10.7%,-0.4%)	p=0.008
Paid support person or carer	15 (3%)	1 (1%)	2% (CI -0.0%,4.7%)	p=0.174 ns
Volunteer support person or carer	6 (1%)	0 (0%)	1% (CI -0.2%,2.5%)	p=0.375 ns
Nobody available	88 (17%)	5 (3%)	14% (CI 9.5%,18.9%)	p<0.001

<b><u>If you were admitted to hospital for a day case surgical procedure, who would be available to collect you afterwards?</u></b>	<b>Autistic n (%)</b>	<b>Non-autistic n (%)</b>	<b>Difference (95% Confidence Interval)</b>	<b>p value</b>
Spouse or partner	208 (41%)	100 (64%)	-23% (CI -31.7%,-13.6%)	p<0.001

Parent	176 (35%)	73 (47%)	-12% (CI -21.0%,-2.5%)	p=0.010
Other family member	96 (19%)	75 (48%)	-29% (CI -37.8%,-19.9%)	p<0.001
Friend	116 (23%)	67 (43%)	-29% (CI -37.8%,-19.9%)	p<0.001
Neighbour	17 (3%)	8 (5%)	-2% (CI -5.9%,2.5%)	p=0.446 ns
Paid support person or carer	17 (3%)	0 (0%)	3% (CI 1.4%,5.3%)	p=0.042
Volunteer support person or carer	3 (1%)	0 (0%)	1% (CI -0.5%,1.7%)	p=0.776 ns
Nobody available	99 (20%)	3 (2%)	18% (CI 13.1%,22.1%)	p<0.001

**If you needed assistance at home after an operation, who would be available to provide that care?**

	Autistic n (%)	Non-autistic n (%)	Difference (95% Confidence Interval)	p value
Spouse or partner	219 (43%)	101 (64%)	-21% (CI -30.2%,-12.1%)	p<0.001
Parent	152 (30%)	74 (47%)	-17% (CI -26.3%,-8.0%)	p<0.001
Other family member	83 (16%)	61 (39%)	-23% (CI -31.2%,-13.8%)	p<0.001
Friend	74 (15%)	43 (27%)	-13% (CI -20.8%,-4.8%)	p<0.001
Neighbour	14 (3%)	8 (5%)	-2% (CI -6.5%,1.8%)	p=0.241 ns
Paid support person or carer	30 (6%)	3 (2%)	4% (CI 0.6%,7.4%)	p=0.071 ns
Volunteer support person or carer	4 (1%)	0 (0%)	1% (CI -0.4%,2.0%)	p=0.599 ns
Nobody available	131 (26%)	13 (8%)	18% (CI 11.4%,23.7%)	p<0.001

**If you are a parent and you were unable to care for your child due to illness, who would be available to provide that care to your child?**

	Autistic n (%)	Non-autistic n (%)	Difference (95% Confidence Interval)	p value
Spouse or partner	110 (22%)	64 (41%)	-19% (CI -28.0%,-10.2%)	p<0.001
Parent	37 (7%)	40 (26%)	-18% (CI -25.8%,-10.6%)	p<0.001
Other family member	40 (8%)	40 (26%)	-18% (CI -25.2%,-10.0%)	p<0.001
Friend	21 (4%)	27 (17%)	-13% (CI -19.6%,-6.5%)	p<0.001
Neighbour	4 (1%)	8 (5%)	-5% (CI -8.2%,-0.4%)	p<0.001

Paid support person or carer	5 (1%)	9 (6%)	-5% (CI -8.9%,-0.6%)	p<0.001
Volunteer support person or carer	1 (1%)	0 (0%)	0% (CI -0.4%,0.8%)	p=1.000 ns
Nobody available	34 (7%)	6 (4%)	3% (CI -1.2%,7.0%)	p=0.256 ns
I don't have a child requiring care	274 (54%)	78 (50%)	4% (CI -5.0%,13.7%)	p=0.387 ns



**Table S5 Facilitators**

<b>Visits to my doctor would be easier if:</b>	<b>Autistic n (%)</b>	<b>Non-autistic n (%)</b>	<b>Difference (95% Confidence Interval)</b>	<b>p value</b>
I could book an appointment online	339 (67%)	102 (65%)	2% (CI -7.0%,10.8%)	p=0.732 ns
I could email in advance with a description of the issue I need to discuss	316 (62%)	35 (22%)	40% (CI 31.9%,48.2%)	p<0.001
I could wait in a quiet place or outside until it was my turn	284 (56%)	13 (8%)	48% (CI 41.2%,54.3%)	p<0.001
I could book the first or last appointment of the day	210 (41%)	38 (24%)	17% (CI 8.8%,25.6%)	p<0.001
I could book an appointment by text	209 (41%)	44 (28%)	13% (CI 4.6%,21.8%)	p=0.004
There was a sensory box* available in the waiting room	80 (16%)	4 (3%)	13% (CI 8.8%,17.7%)	p<0.001
None of the above	16 (3%)	34 (22%)	-19% (CI -25.5%,-11.5%)	p<0.001

\*A sensory box contains items to manage sensory stress, such as sunglasses, ear defenders or ear plugs, items providing tactile sensory input such as fidget tools and olfactory input such as scented items.

## **Supplement 1**

### Barriers to accessing healthcare for autistic adults: online survey

1. Are you autistic?

Includes Autism, Autistic Spectrum Conditions, Asperger Syndrome, self-diagnosed or formally diagnosed.

- Yes  
 No

2. Do you usually attend the same medical practice?

- Yes  
 No  
 I don't attend any medical practice

3. Do you usually see the same doctor?

- Yes  
 No  
 I don't attend any doctor

4. Do you have difficulty visiting your doctor when you need to?

- Yes  
 No

5. Which of the following would cause you to delay or avoid seeing your doctor when you need to?

Please check ALL THAT APPLY

- Difficulty deciding if symptoms warrant a GP visit  
 Difficulty using the telephone to book appointment  
 No online booking system  
 There is an online booking system but it's confusing  
 Difficulty planning an appointment in advance  
 Difficulty communicating with the reception staff  
 Difficulty communicating with the doctor during the appointment  
 The waiting room environment  
 Inability to see a known or preferred doctor  
 Long wait to get an appointment  
 Waiting to see the doctor is too difficult  
 Not feeling understood  
 Not having enough time to visit the doctor  
 Needing a support person to come with me  
 Not having anyone to look after my child  
 None of the above

6. Which of the following is the most difficult part of booking an appointment?

Please choose the one issue which is MOST DIFFICULT for you

- Using the telephone to book an appointment
- No online booking system
- There is an online booking system but it's confusing
- None of the above

7. Which of the following is the most difficult part of planning a visit?

Please choose the one issue which is MOST DIFFICULT for you

- Deciding if your symptoms warrant a GP visit
- Long wait to get an appointment
- Planning an appointment in advance
- Inability to see a known or preferred doctor
- Not having enough time to visit the doctor
- Arranging for someone to come with me
- If you are a parent, not having anyone to look after your child
- None of the above

8. Which of the following is most difficult during an appointment?

Please choose the one issue which is MOST DIFFICULT for you

- Communicating with the reception staff
- Communicating with the doctor during the appointment
- The waiting room environment
- Not feeling understood
- Waiting to see the doctor is difficult
- None of the above

9. Do you visit your doctor...

Check ALL THAT APPLY

- Alone, by choice
- Alone, but would prefer to have a support person
- With a parent, partner or support person
- With a parent, partner or support person but I would prefer to go alone
- With a support animal
- To support an autistic adult
- As a parent with my child

10. Why do you usually visit your doctor?

Check ALL THAT APPLY

- Physical condition or illness
- Mental health difficulties
- Issues directly related to autism
- Other

11. How much do you value your relationship with your GP?

- It is very important to me
- It is important to me
- It is neither important nor unimportant to me
- It is not very important to me

It is not at all important to me

12. Do you have a good relationship with your doctor?

Yes

Sometimes

No

Not currently, but had good relationship with GP in past

13. Do you find it difficult to communicate during a consultation?

All the time

Frequently

Sometimes

Rarely

Not at all

14. Which of the following communication issues cause you problems during a consultation?

Check ALL THAT APPLY

Verbal communication is difficult

Sensory issues make it harder to communicate

Anxiety makes it harder to communicate

It is easier for me to communicate in writing

I'm scared of the receptionist

I have difficulty asking for help

I have difficulty prioritising my health issues

I am concerned I might be labelled a hypochondriac or malingerer

Vague or open ended questions are difficult

I can't describe my pain or symptoms accurately

I express emotions differently (e.g. I can appear to be angry when I am afraid or in pain)

I need extra time to process what is being said

I am concerned I won't be taken seriously when I describe my symptoms

None of the above

Other

15. Which communication issue causes you the MOST problems during a consultation?

Please choose the one issue which is MOST DIFFICULT for you

Verbal communication is difficult

Sensory issues make it harder to communicate

Anxiety makes it harder to communicate

It is easier for me to communicate in writing

I'm scared of the receptionist

I have difficulty asking for help

I have difficulty prioritising my health issues

I am concerned I might be labelled a hypochondriac or malingerer

Vague or open ended questions are difficult

I can't describe my pain or symptoms accurately

I express emotions differently (e.g. I can appear to be angry when I am afraid or in pain)

I need extra time to process what is being said

- I am concerned I won't be taken seriously when I describe my symptoms
- None of the above
- Other

16. Is stimming a problem for you at the doctors' office?

Check ALL THAT APPLY

- My unusual behaviours or stimming elicit negative reactions from other patients
- My unusual behaviours or stimming elicit negative reactions from reception staff
- My unusual behaviours or stimming elicit negative reactions from medical staff
- I feel comfortable stimming at the doctors' office
- I do not feel comfortable stimming at the doctors' office
- I don't feel a need to stim at the doctors' office
- I don't understand the term "stimming"

17. Do you experience any of the following?

Check ALL THAT APPLY

- I find it difficult to make appointments in advance
- I have turned up for a medical appointment on the wrong day
- I have forgotten to attend a medical appointment
- I find it difficult to prioritise when describing my medical problems
- I need to give the whole story and not leave anything out
- I have forgotten why I made the appointment
- I find waiting difficult
- It is difficult to arrange someone to come with me
- I need to write things down
- I have difficulty making decisions about my health
- Making changes to my lifestyle or habits is difficult for me
- None of the above

18. Do you experience sensory issues which make it difficult to visit your doctor?

Check ALL THAT APPLY

- Bright or fluorescent lights
- Noise in the waiting room from other patients
- Music playing in the waiting room
- Noise from the reception desk
- Smells in the waiting room
- Smells in the doctor's office
- Crowded waiting area
- Uncomfortable furniture
- Touch, such as during examination
- Unexpected touch
- None of the above
- Other

19. How would you describe your pain threshold?

- Very high
- High

- Neither particularly high nor low
- Low
- Very low
- I don't know

20. What communication methods do you use?

Check ALL THAT APPLY

- Verbal, face-to-face
- Verbal, telephone
- Voicemail
- Text messaging
- Online messaging
- Email
- Other online method
- Written
- Alternative communication device
- Sign language
- Other

21. What communication methods do you AVOID if possible?

Check ALL THAT APPLY

- Verbal, face-to-face
- Verbal, telephone
- Voicemail
- Text messaging
- Online messaging
- Other online method
- Email
- Written
- Alternative communication device
- Sign language
- Other

22. If your GP offered options for making an appointment, which would you be most likely to use?

- Verbal, face-to-face
- Verbal, telephone
- Voicemail
- Text messaging
- Email
- Online booking system
- Alternative communication device
- Other

23. My doctors are really good for me because they are...

Check ALL THAT APPLY

- Patient

- Good listeners
- Understanding
- Supportive
- Kind
- Friendly
- Efficient
- Familiar
- Not condescending
- Aware of my needs
- Honest about not understanding autism
- None of the above

24. My doctors are really good for me because they...

Check ALL THAT APPLY

- Ask direct questions
- Give clear explanations
- Accept written or email communication
- Use printed information and diagrams
- Value my opinion
- Accept my right to make decisions regarding my health
- Know that autism is not a mental health condition
- Do home visits
- Have reception staff who are trained regarding autism
- Remind me to make my next appointment before leaving
- None of the above

25. Do you find it difficult not knowing...

Check ALL THAT APPLY

- Which doctor you will see
- How long you will wait
- How long the consultation will last
- What will happen during the consultation
- None of the above

26. When going to the doctor, what causes you to feel anxious?

Check ALL THAT APPLY

- asking for help
- discussing mental health
- that there might be something wrong
- that I might be considered a hypochondriac
- that I might be wasting the doctor's time
- I don't feel anxious going to the doctor

27. Do you have mobility needs which make visiting your doctor difficult?

- Yes
- No

28. My doctor is knowledgeable about physical conditions:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

29. My doctor is knowledgeable about mental health conditions:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

30. My doctor is knowledgeable about autism:

- Strongly agree
- Agree
- Neutral
- Disagree
- Strongly disagree

31. Visits to my doctor would be easier if...

Check ALL THAT APPLY

- I could book an appointment online
- I could book an appointment by text
- I could book the first or last appointment of the day
- I could wait in a quiet place or outside until it was my turn
- I could email my doctor in advance with a description of the issue I need to discuss
- There was a sensory box available in the waiting room
- None of the above
- Other

32. Have you ever had a mental health condition remain untreated due to difficulties accessing healthcare?

- Yes
- No
- Possibly
- Other

33. Have you ever had a physical health condition remain untreated due to difficulties accessing healthcare?

- Yes
- No
- Possibly
- Other



34. Have you even been referred to a specialist but did not make an appointment or did not attend?

- Yes  
 No

35. Have you ever been told you should have seen a doctor sooner?

- Yes  
 No

36. Have you ever had to undergo more extensive treatment or surgery than if you had attended sooner?

- Yes  
 No

37. Have you ever had a potentially serious or life threatening condition for which you did not access treatment?

- Yes  
 No

38. If you answered yes to any of the last 6 questions, would you like to give more details?

39. Do you attend on schedule for screening programmes applicable to you?

Includes cervical screening, sexual health screening, breast check, colon screening etc.

- Yes  
 No  
 Sometimes

40. If you were suddenly admitted to hospital, who would be able to bring your personal belongings to you?

Check ALL THAT APPLY

- Spouse or partner  
 Parent  
 Other family member  
 Friend  
 Neighbour  
 Paid support person or carer  
 Volunteer support person or carer  
 Nobody available  
 Other

41. If you were admitted to hospital for a day case surgical procedure, who would be available to collect you afterwards?

Check ALL THAT APPLY

- Spouse or partner  
 Parent  
 Other family member  
 Friend

- Neighbour
- Paid support person or carer
- Volunteer support person or carer
- Nobody available
- Other

42. If you needed assistance at home after an operation, who would be available to provide that care?

Check ALL THAT APPLY

- Spouse or partner
- Parent
- Other family member
- Friend
- Neighbour
- Paid support person or carer
- Volunteer support person or carer
- Nobody available
- Other

43. If you are a parent and you were unable to care for your child due to illness, who would be available to provide that care to your child?

Check ALL THAT APPLY

- Spouse or partner
- Parent
- Other family member
- Friend
- Neighbour
- Paid support person or carer
- Volunteer support person or carer
- Nobody available
- I don't have a child requiring care
- Other

44. Do you identify as autistic?

Includes Autism, Autistic Spectrum Conditions, Asperger Syndrome, self-diagnosed or formally diagnosed.

- Yes
- No
- Possibly

45. Do you have a formal diagnosis? \*

- Yes
- No

46. If you have a formal diagnosis, was it provided by:

- Psychiatrist
- Clinical Psychologist

- Multidisciplinary Team
- Other

47. Does your doctor know you are autistic? \*

- Yes
- No
- I don't know

48. What is your age?

49. What age were you when you were diagnosed?

If applicable

50. Which country do you live in?

- England
- Scotland
- Wales
- Northern Ireland
- Republic of Ireland
- United States
- Canada
- Other:

51. What is your gender?

- Male
- Female
- Non-binary
- Prefer not to say

52. Please give any further information or suggestions here.