


# BMJ Open Comparison and determination of factors associated with smoking status, smoking knowledge, attitude and practice (S-KAP) between smoke-free and non-smoke-free campuses in public universities in Malaysia: a cross-sectional study

Shazwani Mohmad <sup>1,2</sup>, Aniza Ismail,<sup>2</sup> Hayati KS,<sup>3</sup> Noraryana Hassan,<sup>4</sup> Arfah Mahani Imran,<sup>5</sup> Nur Faezah Hamzah,<sup>5</sup> Wan Azrin Izani Wan Mohd Zain<sup>5</sup>

**To cite:** Mohmad S, Ismail A, KS H, *et al.* Comparison and determination of factors associated with smoking status, smoking knowledge, attitude and practice (S-KAP) between smoke-free and non-smoke-free campuses in public universities in Malaysia: a cross-sectional study. *BMJ Open* 2022;**12**:e052275. doi:10.1136/bmjopen-2021-052275

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-052275>).

Received 12 April 2021  
Accepted 15 February 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Professor Aniza Ismail;  
aniza@ppukm.ukm.edu.my

## ABSTRACT

**Objectives** We compared the smoking prevalence, smoking knowledge, attitudes and practices (S-KAP) between smoke-free campuses (SFCs) and non-SFCs (NSFCs) and determined the associated factors of smoking status and S-KAP.

**Design**

Cross-sectional study.

**Settings** The research was conducted in four public universities in Malaysia; two SFCs and two NSFCs.

### Participants

Students and staff from SFCs (n=1063) and NSFCs (n=1040).

**Main outcome measures** Compared the smoking prevalence, S-KAP between SFCs and NSFCs and determined the associated factors.

**Results** The prevalence of smokers among the SFC and NSFC respondents was 5.2% and 6.7%, respectively. University type and smoking attitude were significantly related (p=0.02). At the SFCs, the factors associated with becoming a smoker were male gender (p<0.001), monthly income ≥RM3000 (p=0.02), positive smoking attitude (p=0.003) and positive smoking practice (p<0.001); at NSFCs, the associated factors were male gender (p<0.001), low smoking knowledge (p=0.004), positive smoking attitude (p=0.001) and practice (p<0.001). The factors associated with good smoking knowledge were female gender (SFCs: p=0.001; NSFCs: p=0.004), and monthly income ≥RM3000 (NSFCs: p=0.02). Male respondents were likely to have positive smoking attitudes (SFCs: p<0.001; NSFCs: p<0.001) and negative smoking practices (SFCs: p<0.001; NSFCs: p<0.001).

**Conclusion** Overall, smoking prevalence, knowledge and practice were not much different between SFCs and NSFCs. However, there was a significant relationship between university type and smoking attitude due to the existence of the SFC policy likely rendering smoking a less acceptable social norm. Continual education programmes on smoking harms and smoking cessation strategies are

## Strengths and limitations of this study

- This study has a high response rate (84.1%), contributing to the quality of the study.
- The use of anonymous self-reported questionnaires may reduce social desirability bias.
- Smoking status was assessed using self-reporting and may have resulted in self-reported bias.
- The respondents were volunteers rather than randomly sampled, which may have introduced selection bias.
- The cross-sectional nature of the data precludes any causal effects.

highly recommended to aid the SFC policy in preventing secondhand smoke in universities.

## INTRODUCTION

Globally, over 1.1 billion people, predominantly men, are smokers.<sup>1</sup> Tobacco control is indeed a major challenge to public health, especially for the Southeast Asian region, which is home to 246 million smokers and 290 million passive smokers.<sup>2</sup> Despite declines in overall prevalence in some countries, there are alarming tobacco use trends among young adults.<sup>1</sup> As young adults, university students are a very vulnerable group for risky behaviour initiation, including smoking and drug abuse.<sup>3 4</sup> Smoking prevalence among university students worldwide varies greatly, for example, 15.5% in the UK,<sup>5</sup> 11.1% in New Zealand,<sup>6</sup> 60.2% in Bangladesh,<sup>7</sup> 20.7% in Syria,<sup>8</sup> 3.1% in Hong Kong<sup>9</sup> and 23.4%

in China.<sup>9</sup> In Malaysia, smoking is prevalent among 12%–22.4% of university students.<sup>10–13</sup>

According to the National Health and Morbidity Survey 2019,<sup>14</sup> about 21.3% of the Malaysian population aged  $\geq 15$  years were smokers, and most of them were men. An estimated 30.5% of smokers smoked  $\geq 15$  cigarettes daily.<sup>14</sup> Although the number of smokers had decreased compared with 23.1% in 2011 and 22.8% in 2015,<sup>15 16</sup> the reduction was insufficiently substantial. Therefore, more awareness and efforts should be made to support smoking cessation. The adverse effects and health consequences of smoking are known and have been discussed for decades.<sup>17–19</sup> Smoking not only affects the smoker but also has many harmful effects on non-smokers through secondhand smoke. Millions of people worldwide are affected by secondhand smoke, which contributes to the increased incidence of chronic conditions such as lung cancer, heart disease and respiratory problems<sup>20</sup> and has resulted in legislative action to reduce tobacco consumption among the public. Article 8 of the WHO Framework Convention on Tobacco Control<sup>21</sup> encourages countries to protect people from tobacco smoke in indoor workplaces, public transport and other public places as appropriate. The main objective of a smoke-free area is to reduce smoking, which primarily benefits non-smokers.

Restricting smoking in universities can curb the number of new smokers and reduce smoking prevalence throughout adulthood. The implementation of the smoke-free campus (SFC) policy in universities has been increasing by the year worldwide. By July 2020, at least 2511 US colleges and universities were 100% SFC.<sup>22</sup> Effective SFC policies may lower the prevalence of tobacco use, increase knowledge on smoking harms, limit the visibility of smoking in universities and reduce smoking initiation and daily use.<sup>23–25</sup> Furthermore, a university campus-based smoke-free policy also can lead to smoke-free norms in the university's communities, thus restricting the occurrence of smoking, discouraging initiation and supporting smoking cessation.<sup>26 27</sup>

Malaysia has been a signatory of the WHO FCTC since 2005<sup>15</sup> and has gazetted 23 categories of locations, including universities, where smoking is prohibited under the Control of Tobacco Product Regulations 2004.<sup>28</sup> Although the SFC policy in Western countries, especially in the USA, has been implemented actively, most universities in Malaysia still struggle to implement a 100% SFC policy. The difficulties are due to a lack of standardised policy resources for implementation and a limited workforce for enforcement. Despite these challenges, many university residents support implementing an SFC policy in Malaysia.<sup>29–31</sup> Given the slowly growing number of universities implementing this policy in Malaysia, the impact of the SFC policy has not been evaluated.

Therefore, the purpose of the present study was to compare the smoking prevalence, smoking knowledge, attitudes and practices (S-KAP) between public universities in Malaysia with an SFC policy (SFCs) and those without one (non-SFCs, NSFCs). Second, we wanted to

determine the factors associated with smoking status and S-KAP at SFCs and NSFCs.

## MATERIALS AND METHODS

### Study design and participants

This was a cross-sectional study using baseline data obtained via a self-administered online questionnaire. Invitations to complete the web-based questionnaire were distributed via email and during university events and activities, staff and student electronic newsletters and promotion by representative bodies at the university's main page or social media ads. Data were collected from September 2019 to November 2019. The sample size of 2500 respondents was stratified by the number of students and staff based on the entire population figures from each university.

The inclusion criteria were Malaysian nationality, enrolled student and employed staff at the participating Malaysian public universities and age  $\geq 18$  years. The two SFCs (located on the southwestern and northwestern coast of Peninsular Malaysia, respectively) selected are the earliest public universities in Malaysia to have implemented the SFC policy, that is, since 2013, and have been awarded a Blue Ribbon certificate by the Malaysian Health Promotion Board (MySihat).<sup>15 32</sup> The two NSFCs (located on the west and northwestern coast of Peninsular Malaysia, respectively) were selected by convenience sampling, but have yet to enforce the SFC policy and receive a Blue Ribbon certificate.

### Questionnaire development

The questionnaire was developed using items adapted from the Global Adult Tobacco Survey<sup>33</sup> and the KAP Towards Smoking Survey questionnaire of International Islamic University Malaysia.<sup>34</sup> Experts validated the questionnaire's content, and a pilot study was conducted for the questionnaire's face validity to assess its acceptability and feasibility, and it was subsequently modified before implementation in the present study.

The developed questionnaire consisted of four parts: sociodemographic characteristics and S-KAP studies were conducted in two universities, one with SFC policy implemented and the other without SFC policy. Both universities are located on the west coast of Peninsular Malaysia. There were 136 respondents involved in the pilot studies. Cronbach's alpha value was 0.823 and the exploratory factor analysis (EFA), that is, Kaiser-Meyer-Olkin (KMO) test, was 0.626, and the Bartlett test was  $< 0.0001$ . The strength of correlations between questionnaire items was measured with Cronbach's alpha test ( $> 0.7$ ) while internal reliability was measured with EFA (KMO  $> 0.5$ ; Bartlett test  $< 0.05$ ). These results show that the questionnaire has good internal reliability and that there is a significant correlation between the data.

The final four parts of the questionnaire included:

- Sociodemographic characteristics: gender, age, education level, campus role, monthly income and

smoking status. Smoking status was either non-smoker (including ex-smoker) or smoker (current smoker).

- ▶ **Smoking knowledge:** refers to the university community's awareness on general knowledge of smoking and facts on smoking. The items of the knowledge were: (1) a person who does not smoke but inhales the cigarette smoke is called a secondhand smoker, (2) cigarette smoke contains about 7000 toxic chemicals, (3) quitting smoking can improve the smoker's health, (4) smoking is a major contributor to premature death, (5) nicotine is not addictive, (6) smoking is not a major risk factor for sudden infant death syndrome, (7) smoking increases diabetes risk, (8) smoking causes impotence, (9) no safe level of exposure to secondhand smoke, (10) cigarette smoke particles can remain on any surface for 6 months, (11) cigarette smoke does not deplete the ozone layer, (12) the particle size of cigarette smoke is larger than the particle size of haze. The answer to each question was yes/no/do not know. The correct answer was scored 1 point. Smoking knowledge was measured by calculating the score of correctly answered questions per total items and was categorised as good (if respondents scored >median score/50% of the total score) or poor (if respondents scored ≤median score/50% of the total score).<sup>35</sup>
- ▶ **Smoking attitude:** refers to the level of agreement and beliefs regarding smoking norms. The items of smoking attitude were: (1) smoking contributes to many adverse effects on my health, (2) cigarette smoke is a serious threat to my health, (3) quitting smoking can improve smokers' health, (4) I feel better if I so not smoke, (5) I feel uncomfortable when I get close to people who smoke, (6) smoking shows bad behaviour, (7) smoking in universities should be completely banned, (8) I am easily accepted by friends if I smoke, (9) Smoking describes maturity. Smoking was measured using a 5-point Likert scale and with a score of 1 to 5. Smoking attitude was measured by calculating the score of the answered questions and was categorised as negative (if respondents scored >60% of the maximum score) and positive (if respondents scored ≤60% of the maximum score).
- ▶ **Smoking practice:** refers to the consistency of the university's community involvement in controlling smoking habits. The items of the smoking practice were: (1) I reprimand anyone who smokes on university grounds, (2) I have never advised my friends to quit smoking, (3) I cover my nose when someone smokes around me, (4) I avoid spending time with friends who smoke, (5) I avoid going to the smoking area to breathe fresh air, (6) I live a healthy lifestyle instead of smoking. Smoking practice was measured using a 5-point Likert scale and with a score of 1 to 5. Smoking practice was measured by calculating the score of the answered questions and was categorised as negative (if respondents scored >60% of the

maximum score) or positive (if participants respondents ≤60% of the maximum score).

### Data analysis

The data were divided into two categories: SFC and NSFC. The descriptive statistics of the sample were computed and showed the percentage distributions for all response categories, among all respondents, and smoking status. The association between university type with smoking status and S-KAP was determined using the  $\chi^2$  test for independence. Univariate and multivariate logistic regression was used to measure the relationships between independent variables (gender, age, education level, employment status, monthly income) and the OR of smoking status and S-KAP. The collected data were analysed using IBM SPSS Statistics V.21.

## RESULTS

A total of 2103 out of 2500 respondents completed and returned the questionnaire, yielding an overall response rate of 84.1%. For both university types, most respondents were women (SFCs, 64%; NSFCs, 65.9%) and from the 19–40 year age group (SFCs, 95.7%; NSFCs, 95.3%). Up to 29% (SFCs) and 31.6% (NSFCs) of the respondents had secondary education, 65% (SFCs) and 63.1% (NSFCs) had undergraduate education and 6% (SFCs) and 5.3% (NSFCs) had postgraduate education.

The majority of respondents were students (SFCs, 87.9%; NSFCs, 85.5%). Among the students and staff, most of them had a monthly income of <RM3000 (SFCs, 90.4%; NSFCs, 89.8%). The majority of respondents were non-smokers (SFCs, 94.8%; NSFCs, 93.3%). Only 5.2% and 6.7% of the SFC and NSFC respondents, respectively, were smokers. [Table 1](#) shows a detailed summary of the respondents' sociodemographic characteristic.

### Association between university type with smoking status and S-KAP

Descriptively, the SFCs had better smoking prevalence and S-KAP than the NSFCs. The  $\chi^2$  test for independence showed that the relationship between university type and smoking status ( $p=0.131$ ), level of smoking knowledge ( $p=0.17$ ) and smoking practices ( $p=0.177$ ) was not significant ([table 2](#)). However, there was a significant relationship between university type and smoking attitude ( $p=0.02$ ). Backward likelihood ratio multiple logistic regression was applied ([table 3](#)) and the Hosmer-Lemeshow test was >0.05 ( $p=0.92$ ). There was no significant difference between the observed and expected probability. Thus, the assumption of model fitness was met. [Table 4](#) shows that there was a significant interaction between smoking status×income ( $p=0.013$ ), smoking status×smoking knowledge ( $p=0.009$ ), smoking status×smoking attitude ( $p=0.012$ ) and smoking attitude×gender ( $p=0.006$ ).

**Table 1** Sociodemographic characteristics of respondents at SFC and NSFC

Sociodemographic characteristic		Type of university			
		SFC (n=1063)		NSFC (n=1040)	
		n	Percentage	n	Percentage
Gender	Male	383	36.0	355	34.1
	Female	680	64.0	685	65.9
Age	19–40	1017	95.7	991	95.3
	41 and above	46	4.3	49	4.7
Education level	Secondary education	308	29.0	329	31.6
	Undergraduate	691	65.0	656	63.1
	Postgraduate	64	6.0	55	5.3
Campus role	Student	934	87.9	889	85.5
	Staff	129	12.1	151	14.5
Monthly income	<RM3000	961	90.4	934	89.8
	RM3000 and above	102	9.6	106	10.2
Smoking status	Non-smoker	1008	94.8	970	93.3
	Smoker	55	5.2	70	6.7

NSFC, non-SFC; SFC, smoke-free campus.

**Table 2** Association between university type with respondents' characteristics, smoking status and S-KAP

Domain		Type of university		X <sup>2</sup> statistic (df)	P * value
		SFC n (%)	NSFC n (%)		
Gender	Male	383 (36.0)	355 (34.1)	0.829 (1)	0.36
	Female	680 (64.0)	685 (65.9)		
Age	19–40	1017 (95.7)	991 (95.3)	0.18 (1)	0.67
	≥41	46 (4.3)	49 (4.7)		
Education level	Secondary	308 (29.0)	329 (31.6)	2.031 (2)	0.36
	Undergraduate	691 (65.0)	656 (63.1)		
	Postgraduate	64 (6.0)	55 (5.3)		
Campus role	Student	934 (87.9)	889 (85.5)	2.588 (1)	0.11
	Staff	129 (12.1)	151 (14.5)		
Monthly income	<RM3000	961 (90.4)	934 (89.8)	0.210 (1)	0.647
	≥RM3000	102 (9.6)	106 (10.2)		
Smoking status	Non-smoker	1008 (94.8)	970 (93.3)	2.279 (1)	0.131
	Smoker	55 (5.2)	70 (6.7)		
Smoking knowledge	0–8 (≤50%)	334 (31.4)	356 (34.2)	1.883 (1)	0.17
	9–16 (>50%)	729 (68.6)	684 (65.8)		
Smoking attitude	Negative	1018 (95.8)	973 (93.6)	5.088 (1)	0.02
	Positive	45 (4.2)	67 (6.4)		
Smoking practice	Negative	953 (89.7)	913 (87.8)	1.826 (1)	0.177
	Positive	110 (10.3)	127 (12.2)		

\*Chi-square test for independence; n=frequency; df=degree of freedom.

NSFC, non-SFC; SFC, smoke-free campus; S-KAP, smoking knowledge, attitudes and practices.

**Table 3** Logistic regression model of university type

Variable	Type of university				Adjusted OR (95% CI)	P value
	SFC		NSFC			
	Mean (SD)	n (%)	Mean (SD)	n (%)		
Gender	Male	383 (36.0)	355 (34.1)	0.87 (0.72 to 1.04)	0.13	
	Female	680 (64.0)	685 (65.9)	1		
Age		23.8 (6.45)	24.0 (6.91)	1.0 (0.98 to 1.03)	0.97	
Education level	Secondary school	308 (29.0)	329 (31.6)	1	0.13	
	Undergraduate	691 (65.0)	656 (63.1)	0.88 (0.73 to 1.06)		
	Postgraduate	64 (6.0)	55 (5.3)	0.66 (0.43 to 1.02)		
Campus role	Student	934 (87.9)	889 (85.5)	0.81 (0.63 to 1.05)	0.11	
	Staff	129 (12.1)	151 (14.5)	1		
Monthly income	<RM3000	961 (90.4)	934 (89.8)	1	0.97	
	≥RM3000	102 (9.6)	106 (10.2)	0.99 (0.61 to 1.6)		
Smoking status	Non-smoker	1008 (94.8)	970 (93.3)	1	0.33	
	Smoker	55 (5.2)	70 (6.7)	0.91 (0.76 to 1.1)		
Smoking knowledge	0–8 (≤50%)	334 (31.4)	356 (34.2)	1	0.30	
	9–16 (>50%)	729 (68.6)	684 (65.8)	0.91 (0.75 to 1.09)		
Smoking attitude	Negative	1018 (95.8)	973 (93.6)	1.56 (1.06 to 2.3)	0.03	
	Positive	45 (4.2)	67 (6.4)	1		
Smoking practice	Negative	953 (89.7)	913 (87.8)	1.5 (0.76 to 1.44)	0.78	
	Positive	110 (10.3)	127 (12.2)	1		

Backward likelihood ratio multiple logistic regression was applied. Hosmer-Lemeshow test  $p=0.92$  was accepted to check model fitness. NSKC, non-SKC; SKC, smoke-free campus.

### Factors associated with smoking status

Multiple logistic regression analyses of the SFCs showed that the determinants for a person to become a smoker were gender, monthly income, smoking attitude and smoking practice (table 5). This analysis was interpreted as: (1) male respondents had seven times higher odds than female respondents of becoming a smoker (95% CI 3.2 to 16.04,  $p<0.001$ ), (2) respondents with a monthly income  $\geq$ RM3000 had two times higher odds than those earning <RM3000 of becoming smokers (95% CI 1.16 to 5.34,  $p=0.02$ ), (3) negative smoking attitude was less likely to be a smoker's attitude (95% CI 0.12 to 0.66,  $p=0.003$ ), (4) negative smoking practice was less likely to be a smoker's practice (95% CI 0.09 to 0.33,  $p<0.001$ ).

Multiple logistic regression analyses of the NSFCs showed that the determinants for a person to become a

smoker were gender and S-KAP (table 5). This analysis was interpreted as: (1) male respondents had 10 times higher odds than female respondents of becoming a smoker (95% CI 4.7 to 23.55,  $p<0.001$ ), (2) >50% correct answers for smoking knowledge was less likely to be a smoker's knowledge score (95% CI 0.21 to 0.74,  $p=0.004$ ), (3) negative smoking attitude was less likely to be a smoker's attitude (95% CI 0.12 to 0.57,  $p=0.001$ ), (4) negative smoking practice was less likely to be a smoker's practice (95% CI 0.07 to 0.28,  $p<0.001$ ).

### Factors associated with smoking knowledge

The multiple logistic regression analysis showed that, for SFCs, the determinant of >50% correct answers for smoking knowledge was gender (table 6). This analysis can be interpreted as: (1) male respondents were 29% less likely than female respondents to have >50% correct answers on smoking knowledge (95% CI 0.54 to 0.92,  $p=0.01$ ).

For the NSFCs, multiple logistic regression analysis showed that the determinants of >50% correct answers for smoking knowledge were gender and monthly income (table 6). This analysis was interpreted as: (1) male respondents were 33% less likely than female respondents to have >50% correct answers on smoking knowledge (95% CI 0.51 to 0.88,  $p=0.004$ ); (2) respondents earning  $\geq$ RM3000 had 1.8 times higher odds than

**Table 4** Significant interaction between variables

Variable	Type of university, adjusted OR (95% CI)	P value
Smoking Status $\times$ Income	0.24 (0.08 to 0.74)	0.013
Smoking Status $\times$ Smoking Knowledge	0.4 (0.18 to 0.79)	0.009
Smoking Status $\times$ Smoking Attitude	2.22 (1.19 to 4.12)	0.012
Smoking Attitude $\times$ Gender	1.96 (1.2 to 3.18)	0.006



**Table 5** Multiple logistic regression analysis for factors related to smoking status

Type of university										
SFC (n=1063)										
Smoking status										
Non-smoker										
Smoker										
Variables	Mean (SD)	n (%)	Mean (SD)	n (%)	Adjusted OR (95% CI)	P value	Mean (SD)	n (%)	Adjusted OR (95% CI)	P value
NSFC (n=1040)										
Smoking status										
Non-smoker										
Smoker										
Gender						<0.001				<0.001
Male	336 (87.7)	47 (12.3)	7.24 (3.2 to 16.04)	293 (82.5)	10.5 (4.70 to 23.55)		62 (17.5)			
Female	672 (98.8)	8 (1.2)	1	677 (98.8)	1		8 (1.2)			
Age	23.6 (6.37)	26.0 (7.42)	0.98 (0.92 to 1.05)	23.9 (6.9)	1.04 (0.96 to 1.13)	0.68	25.3 (6.99)			0.30
Campus role										
Student	893 (95.6)	41 (4.4)	0.86 (0.14 to 5.37)	835 (93.9)	0.82 (0.17 to 3.97)	0.87	54 (6.1)			0.80
Staff	115 (89.1)	14 (10.9)	1	135 (89.4)	1		16 (10.6)			
Monthly income										
<RM3000	918 (95.5)	43 (4.5)	1	869 (93.0)	1	0.02	65 (7.0)			0.39
≥RM3000	90 (88.2)	12 (11.8)	2.48 (1.16 to 5.34)	101 (95.3)	0.66 (0.26 to 1.68)		5 (4.7)			
Smoking knowledge										
0–8 (≤50%)	310 (92.8)	24 (7.2)	1	309 (86.8)	1	0.64	47 (13.2)			0.004
9–16 (>50%)	698 (95.7)	31 (4.3)	0.8 (0.45 to 1.62)	661 (96.6)	0.4 (0.21,0.74)		23 (3.4)			
Smoking attitude										
Negative	978 (96.1)	40 (3.9)	0.28 (0.12 to 0.66)	935 (96.1)	0.26 (0.12 to 0.57)	0.003	38 (3.9)			0.001
Positive	30 (66.7)	15 (33.3)	1	35 (52.2)	1		32 (47.8)			
Smoking practice										
Negative	928 (97.4)	25 (2.6)	0.17 (0.09 to 0.33)	889 (97.4)	0.14 (0.07 to 0.28)	<0.001	24 (2.6)			<0.001
Positive	80 (72.7)	30 (27.3)	1	81 (63.8)	1		46 (36.2)			

NSFC, non-SFC; SFC, smoke-free campus.

**Table 6** Multiple logistic regression analysis for factors related to smoking knowledge

Variables	Type of university								
	SFC (n=1063)				NSFC (n=1040)				
	Smoking knowledge		Adjusted OR (95% CI)	P value	Smoking knowledge		Adjusted OR (95% CI)	P value	
	0-8 (≤50%) n (%)	9-16 (>50%) n (%)			0-8 (≤50%) n (%)	9-16 (>50%) n (%)			
Gender								0.01	0.004
Male	139 (36.3)	244 (63.7)	0.71 (0.54 to 0.92)		142 (40.0)	213 (60.0)	0.67 (0.51 to 0.88)		
Female	195 (28.7)	485 (71.3)	1		214 (31.2)	471 (68.8)	1		
Educational level								0.07	0.40
Secondary Education level	113 (36.7)	195 (63.3)	1		122 (37.1)	207 (62.9)	1		
Undergraduate level	205 (29.7)	486 (70.3)	1.33 (1.0 to 1.77)		223 (34)	433 (66)	1.08 (0.82 to 1.44)		
Postgraduate level	16 (25.0)	48 (75.0)	1.75 (0.94 to 3.2)		11 (20)	44 (80)	1.69 (0.78 to 3.65)		
Monthly income								0.69	0.02
<RM3000	307 (31.9)	654 (68.1)	1		331 (35.4)	603 (64.6)	1		
≥RM3000	27 (26.5)	75 (73.5)	1.12 (0.65 to 1.04)		25 (23.6)	81 (76.4)	1.76 (1.09 to 2.83)		

NSFC, non-SFC; SFC, smoke-free campus.

those earning <RM3000 to have >50% correct answers on smoking knowledge (95% CI 1.09 to 2.83,  $p=0.02$ ).

### Factors associated with smoking attitudes and smoking practices

For both SFCs and NSFCs, multiple logistic regression analyses showed that the determinant of negative smoking attitudes (table 7) and negative smoking practice (table 8) was gender. Male respondents were less likely than female respondents to have negative smoking attitudes (SFCs: 95% CI 0.1 to 0.29,  $p<0.001$ ; NSFCs: 95% CI 0.16 to 0.55,  $p<0.001$ ) and negative smoking practices (SFCs: 95% CI 0.14 to 0.33,  $p<0.001$ ; NSFCs: 95% CI 0.17 to 0.36,  $p<0.001$ ).

## DISCUSSION

### Smoking status

We found that the smoking prevalence at SFCs (5.2%) and NSFCs (6.7%) was not much different. Nonetheless, similar to another study in Kazan city that compared SFCs

and NSFCs,<sup>36</sup> SFCs had a lower smoking prevalence than NSFCs descriptively. Other studies have also reported low smoking prevalence in universities that implement SFC policies, for example, 9.2% in the USA<sup>37</sup> and 8.4% in Thailand,<sup>24</sup> due to decreases in smoking behaviour. Other studies<sup>38-39</sup> have reported that SFCs are expected to have lower smoking prevalence than NSFCs because students and staff have a greater chance of being exposed to awareness campaigns and knowledge on smoking risk. SFCs also have supportive programmes for smoking cessation and have a policy strictly banning smoking on campus grounds.

However, we found no significant association between university type and smoking status ( $p=0.131$ ). This is probably because all universities in Malaysia have been gazetted as non-smoking areas under Regulation 11 (1) of the Tobacco Products Control Regulations (Amendment) 2018.<sup>40</sup> Therefore, although there is no specific SFC policy, all universities are expected to advertise smoke-free campaigns and help initiate a smoke-free environment, thus helping smokers stop smoking.

**Table 7** Multiple logistic regression analysis for factors related to smoking attitude

Variables	Type of university								
	SFC (n=1063)				NSFC (n=1040)				
	Smoking attitude		Adjusted OR (95% CI)	P value	Smoking attitude		Adjusted OR (95% CI)	P value	
	Negative n (%)	Positive n (%)			Negative n (%)	Positive n (%)			
Gender								<0.001	<0.001
Male	354 (92.4)	29 (7.6)	0.17 (0.1 to 0.29)		306 (86.2)	49 (13.8)	0.29 (0.16 to 0.55)		
Female	664 (97.6)	16 (2.4)	1		667 (97.4)	18 (2.6)	1		

NSFC, non-SFC; SFC, smoke-free campus.

**Table 8** Multiple logistic regression analysis for factors related to smoking practice

Variables	Type of university		Adjusted OR (95% CI)	P value	NSFC (n=1040)		Adjusted OR (95% CI)	P value
	SFC (n=1063)				Smoking practice			
	Negative	Positive			Negative	Positive		
	n (%)	n (%)			n (%)	n (%)		
Gender				<0.001				<0.001
Male	307 (80.2)	76 (19.8)	0.21 (0.14 to 0.33)		274 (77.2)	81 (22.8)	0.24 (0.17 to 0.36)	
Female	646 (95)	34 (5)	1		639 (93.3)	46 (6.7)	1	

NSFC, non-SFC; SFC, smoke-free campus.

In the present study, we demonstrated, for both university types, males were more likely to be smokers. Others have reported a similar result<sup>41–43</sup> as well as that for smoking prevalence worldwide.<sup>44</sup> In addition, there was a significant interaction between smoking status and gender. The main possible reason is social norms. Some believe that social acceptability of smoking differs between men and women. Women are more likely to believe that society disapproves of smoking, and men tend to believe that smoking will help them appear more mature and more attractive, and that it helps to control body weight.<sup>45</sup>

In the present study, we found that, for SFCs, a monthly income of  $\geq$ RM3000 is associated with becoming a smoker. Besides, there was a significant interaction between smoking status and monthly income. Other studies have also reported a strong relationship between personal income and smoking.<sup>46,47</sup> An increase in income may lead to higher cigarette consumption and higher smoking prevalence rates among adolescents with a higher personal income.<sup>47</sup> Income status is a powerful barrier against smoking. Moreover, cigarettes are now becoming more expensive due to high taxes, one of the WHO FCTC initiatives through MPOWER.<sup>21</sup> Increasing the retail price of tobacco products through higher taxes is the single most effective means of decreasing consumption and encouraging smokers to quit.<sup>48</sup> However, monthly income was a predictor of being a smoker for only SFCs and not NSFCs, which requires further exploration.

We also found that, in the NSFCs, the factor associated with becoming a smoker was poor knowledge of the adverse effects of smoking. Surveys from Iraq<sup>49</sup> and India have reported similar results<sup>18</sup>; smokers have only a moderate level of awareness of the health risk effects of smoking, especially on non-smokers, and this might be one reason they became smokers. In contrast, SFCs present the chance to gain awareness of the danger of smoking through their smoke-free policies, and campus campaigns may provide the university community with good knowledge of smoking harms and thus prevent them from taking up smoking.<sup>50</sup>

On the other hand, positive smoking attitude and positive smoking practice were associated with becoming a smoker for both SFCs and NSFCs. The interaction between the variables also indicates that there was an

interaction between smoking status and smoking attitude. Smokers tended to agree that smoking norms are acceptable (positive smoking attitudes) and tended to not be involved in the control of smoking activity on campus (positive smoking practices). Such actions mean that there are no smoking-limiting behaviours, thus it is easier for such respondents to become smokers.

### Smoking knowledge

In the present study, the descriptive data show that SFC respondents had better smoking knowledge compared with NSFC respondents. Understanding of the harmful effects of smoking increased after the implementation of a smoke-free policy.<sup>51</sup> However, a  $\chi^2$  test for independence indicated no significant association between university type and smoking knowledge ( $p=0.17$ ). This is probably because information on the harm of tobacco is not only obtained from the SFC policy but is mostly at our fingertips, namely, on social media and TV ads. The factors associated with good smoking knowledge among the SFC respondents were gender and for the NSFCs were gender and monthly income. Female respondents were more likely to have adequate smoking knowledge than male respondents because they were more aware of smoking diseases. They were also more attentive and focused on antismoking messages through various channels such as social media ads, TV ads and printed materials.<sup>52,53</sup> Female respondents were less likely to become smokers because of their high awareness of the harm of smoking. Following the results linking monthly income and smoking knowledge, we showed that respondents with an income of  $>$ RM3000 have a good understanding of smoking compared with those with an income of  $<$ RM3000. This result is similar to that of another study that reported that people with low education, low-level occupations and low income have less access to adequate healthcare information.<sup>54</sup>

### Smoking attitude

There was a significant association between smoking attitude and university type ( $p=0.02$ ). Our findings are consistent with the previous studies reported that the SFC policy directly impacts general attitude towards smoking and that respondents are more likely to see the ban as



having a positive effect on their quality of life.<sup>51 55</sup> At Indiana University, the implementation of the SFC policy significantly reduced the number of people who agreed to allow smoking on campus.<sup>38</sup> The SFC policy might change the perceived tobacco-related norms in a community, making smoking less socially acceptable.<sup>27 38</sup> In the present study, we found that, for both university types, the factor associated with negative smoking attitude was female gender. In addition, there was a significant interaction between smoking attitude and gender. These findings are consistent with other studies examining the role of gender in attitude regarding smoking.<sup>56–58</sup> Women are more likely to have a negative attitude towards smoking. They are more likely to be bothered by tobacco smoke and have a stronger belief in the danger of smoking.

### Smoking practice

We found no significant association between smoking practice and university type ( $p=0.177$ ). However, the SFCs had a higher percentage of negative smoking practice (51.1%) than the NSFCs (48.95%). Consistent with other studies, smoking bans decrease smoking visibility in public areas and are useful for reducing smoking behaviour.<sup>26 27 59</sup> The SFC policy restricts where smoking occurs on campus; later, it may discourage smoking initiation and support tobacco cessation. Consequently, it influences smokers' behaviour, thus they comply with the rules and have negative smoking practice. Here, the multivariate analysis showed that female respondents tended to have negative smoking practices compared with male respondents for both university types. According to Chinwong *et al*,<sup>43</sup> women are significantly more likely than men to have health concerns and perceive the risk of dying from smoking.

### Study limitations and strengths

The limitations of this study are mostly due to the use of the self-administered questionnaire. Smoking status was assessed using self-report and may have resulted in self-reported bias. Furthermore, the low proportion of smokers who participated in our survey may have been due to selective non-reporting or under-reporting and the low response rate. Besides, this study is based on convenience sampling. The respondents were volunteers rather than randomly sampled, which may have introduced selection bias. Finally, the cross-sectional nature of the data precludes any causal effects.

Despite the limitations, this is the first study of public universities in Malaysia that provides valuable novel findings in the comparison of smoking prevalence and S-KAP between SFCs and NSFCs. Although convenience sampling was used in this study, we minimised bias by using it together with probability sampling (stratified sampling). Prior to convenience sampling, the sample size was stratified by the number of students and staff based on the entire population figures from each university. For non-response bias, we minimised the bias by achieving a high response rate (>80%). The use of anonymous

self-reported questionnaires may have reduced social desirability bias.

### CONCLUSIONS

In general, there is not much difference in the smoking prevalence, knowledge and practice between SFCs and NSFCs in public universities in Malaysia. This is probably because all universities in Malaysia have been gazetted as non-smoking areas under Regulation 11 (1) of the Tobacco Products Control (Amendment) Regulations 2018<sup>40</sup> and are expected to advertise smoke-free campaigns and help initiate a smoke-free environment. However, there was a significant relationship between university type and smoking attitude. The existence of the SFC policy is likely to create new norms related to smoking and render smoking less acceptable as a social norm.<sup>27</sup> Even so, our findings showed that most of the factors associated with smoking status and S-KAP were related to gender. Different approaches between men and women need to be emphasised, so that the impact of the SFC policy is balanced between the genders. We hope that our study can aid policymakers' understanding of the S-KAP among university residents in different environments and enable future evaluation of the policy to improve efforts in effective implementation and make an important contribution to creating non-smoking norms, thus reducing secondhand smoke in universities.

### Author affiliations

<sup>1</sup>Centre for Leadership & Professional Development, Institute for Health Management, Shah Alam, Selangor, Malaysia

<sup>2</sup>Community Health, Universiti Kebangsaan Malaysia, Cheras, Malaysia

<sup>3</sup>Community Health, Universiti Putra Malaysia, Serdang, Malaysia

<sup>4</sup>Disease Control Division, Ministry of Health, Malaysia, Putrajaya, Malaysia

<sup>5</sup>Health Education Division, Ministry of Health Malaysia, Putrajaya, Malaysia

**Contributors** SM—conceptualisation, methodology, software, formal analysis, investigation, data curation, writing—original draft preparation, visualisation and project administration. AI—conceptualisation, methodology, validation, formal analysis, data curation, writing—original draft preparation, writing—review and editing, visualisation and supervision, project administration. HKS—validation, formal analysis, writing—original draft preparation, writing—review and editing and visualisation. NH—resources, supervision, and project administration. AMI—resources, project administration and funding acquisition. NFH—resources, project administration, and funding acquisition. WAIWMZ—resources, project administration and funding acquisition.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was approved by Ethics Committee—Universiti Kebangsaan Malaysia (UKM) Medical Centre Reference number—Ref: UKM PPI/111/8/JEP-2019-153. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. The data presented in this study are available on request from the corresponding author. The data are not publicly available to protect confidentiality of the research participants.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content

includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Shazwani Mohamad <http://orcid.org/0000-0003-0811-9498>

#### REFERENCES

- World Health Organization. World Health organization global report on trends in prevalence of tobacco smoking 2015 2015.
- Southeast Asia Tobacco Control Alliance. *Addicted to change SEATCA*. Bangkok, Thailand: SEATCA, 2015.
- Creamer MR, Wang TW, Babb S. Tobacco product use and cessation indicators among adults — United States, 2018 2018.
- Wu Y, Fan H, Guo Z, et al. Factors associated with smoking intentions among Chinese college students. *Am J Mens Health* 2019;13:1557988318818285.
- Bartington SE, Wootton R, Hawkins P, et al. Smoking behaviours and attitudes towards campus-wide tobacco control policies among staff and students: a cross-sectional survey at the University of Birmingham. *BMC Public Health* 2020;20:252.
- Wamamili B, Wallace-Bell M, Richardson A, et al. Cigarette smoking among university students aged 18-24 years in New Zealand: results of the first (baseline) of two national surveys. *BMJ Open* 2019;9:e032590.
- Hossain S, Hossain S, Ahmed F, et al. Prevalence of tobacco smoking and factors associated with the initiation of smoking among university students in Dhaka, Bangladesh. *Cent Asian J Glob Health* 2017;6:244.
- Al-Kubaisy W, Abdullah NN, Al-Nuaimy H, et al. Factors associated with smoking behaviour among university students in Syria. *Procedia Soc Behav Sci* 2012;38:59-65.
- Seo D-C, Torabi MR, Kim N, et al. Smoking among East Asian college students: prevalence and correlates. *Am J Health Behav* 2013;37:199-207.
- Prevalence and Associated Factors of Smoking Among Male Undergraduate Students in International Islamic University Malaysia. *Proceedings of the 2nd Syedza Saintika International Conference on Nursing, Midwifery, Medical Laboratory Technology, Public Health, and Health Information Management (SeSICNiMPH 2021)*, 2021.
- Elkalmi RM, Alkoudmani RM, Elsayed TM, et al. Effect of religious beliefs on the smoking behaviour of university students: quantitative findings from Malaysia. *J Relig Health* 2016;55:1869-75.
- Chirkitsakul BJ, Kadir Rohana, Abdul R. Nordin, Amer Siddiq Amer, Azizi, Ridza. prevalence of and factors associated with smoking among University of Malaya students in Malaysia. *Southeast Asian J Trop Med Public Health* 2019;50:1202-11.
- Saravanan C, Heidhy I. Psychological problems and psychosocial predictors of cigarette smoking behavior among undergraduate students in Malaysia. *Asian Pac J Cancer Prev* 2014;15:7629-34.
- National Institutes of Health MoH. *National health and morbidity survey 2019: non-communicable diseases, healthcare demand and health literacy*. Shah Alam, Selangor: Institute for Public Health, National Institutes of Health, Ministry of Health, 2019.
- Malaysia KK. *Pelan Strategik Kebangsaan Bagi Kawalan Tembaku 2015 - 2020*. Kementerian Kesihatan Malaysia, 2015.
- Institute for Public Health. *National Health And Morbidity Survey 2015 - Report On Smoking Status Among Malaysian Adults Institute for Public Health*. Malaysia: National Institutes of Health, Ministry of Health, 2015.
- Center for Disease Control and Prevention. Health effects of Secondhand smoke center for disease control and prevention website, 2017. Available: [https://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/secondhand\\_smoke/health\\_effects/index.htm](https://www.cdc.gov/tobacco/data_statistics/fact_sheets/secondhand_smoke/health_effects/index.htm)
- Centers for Disease Control and Prevention. Fast facts of smoking website: centers for disease control and prevention, 2017. Available: [https://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/fast\\_facts/index.htm](https://www.cdc.gov/tobacco/data_statistics/fact_sheets/fast_facts/index.htm)
- Malek AM, Cushman M, Lackland DT, et al. Secondhand smoke exposure and stroke. *Am J Prev Med* 2015;49:e89-97.
- Lubick N. Global estimate of shs burden. *Environ Health Perspect* 2011;119:A66-7.
- World Health Organization. *Who framework convention tobacco control*. 2005. Geneva, Switzerland: World Health Organization, 2005.
- Foundation TANRA. Smokefree colleges and universities continue to grow in popularity.
- Services USDoHaH. *Preventing tobacco use among youth and young adults: a report of the surgeon General*. Atlanta, GA: Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2012.
- Sookaneknun S, Sookaneknun P, Seesin T. Outcomes of a two-year smoke-free university and organizational policy management. *Malays J Public Health Med* 2018;18:109-22.
- Blake KD, Klein AL, Walpert L, Laura Walpert LC, et al. Smoke-Free and tobacco-free colleges and universities in the United States. *Tab Control* 2020;29:289-294.
- Procter-Scherdtel A, Collins D. Social norms and smoking bans on campus: interactions in the Canadian university context. *Health Educ Res* 2013;28:101-12.
- Rogers CJ, Barrington-Trimis JL, Unger JB, et al. Changes in smoking prevalence and perception of smoking on campus before and after a smoke-free university campus policy. *J Am Coll Health* 2020:1-5.
- Kementerian Kesihatan Malaysia. Kenyataan Akhbar Kementerian Kesihatan Malaysia. In: Malaysia KPK, ed. *Taman Awam DAN Taman Negara Sebagai Tempat Dilarang Merokok DAN Pemansuhan Peraturan 21 PPKHT 2004 Kementerian Kesihatan Malaysia*, 2017.
- Mohd Nor NA, Ismail N, Ibrahim F. Tobacco use and attitudes towards tobacco control activities of Malaysian dental students. *Archiv Orofac Sci* 2018;13:6-15.
- Kuang Hock L, Hui Li L, Chien Huey T, et al. Support for smoke-free policy among Malaysian adults: findings from a population-based study. *BMJ Open* 2019;9:e020304.
- Yasin SM, Zubillah A, Selamat MI, et al. Tobacco-free policy support among students in University Campuses. *E-BPJ* 2020;5:119-26.
- Alliance SATC. Malaysia: blue ribbon Award: Southeast Asia tobacco control alliance, 2020. Available: <https://smokefreeasean.seatca.org/awards/> [Accessed 20.12.2020].
- Group GATSC. *Tobacco questions for survey: a subset of key questions from the global adult tobacco survey (GATS)*. 2nd Edition. Centers for Disease Control and Prevention, World Health Organization, 2011.
- Ah N, Ar A A, Rahman DNI. Knowledge, attitude and practice towards smoking among international Islamic university Malaysia Kuantan communities. *Int Med J Malays* 2016;15:2016.
- Alzahrani SH. Levels and factors of knowledge about the related health risks of exposure to secondhand smoke among medical students: a cross-sectional study in Jeddah, Saudi Arabia. *Tab Induc Dis* 2020;18:1-14.
- Andreeva TI, Ananjeva GA, Daminova NA, et al. Smoke-Free universities help students avoid establishing smoking by means of facilitating quitting. *Health Promot Perspect* 2015;5:241-9.
- Gatto A, Powell SE, Walters EF, et al. A mixed-methods assessment of a Peer-Enforced Tobacco- and smoke-free policy at a large urban university. *J Community Health* 2019;44:365-76.
- Seo D-C, Macy JT, Torabi MR, et al. The effect of a smoke-free campus policy on college students' smoking behaviors and attitudes. *Prev Med* 2011;53:347-52.
- Lechner WV, Meier E, Miller MB, et al. Changes in smoking prevalence, attitudes, and beliefs over 4 years following a campus-wide anti-tobacco intervention. *J Am Coll Health* 2012;60:505-11.
- Government M. *Tobacco products control regulations (Amendment) 2018*. Malaysia: Malaysian Government, 2018: 9-10.
- Mandil A, BinSaeed A, Ahmad S, et al. Smoking among university students: a gender analysis. *J Infect Public Health* 2010;3:179-87.
- Karadoğan D, Önal Özgür, Kanbay Y. Prevalence and determinants of smoking status among university students: Artvin Çoruh university sample. *PLoS One* 2018;13:e0200671-e71.
- Chinwong D, Mookmanee N, Chongpornchai J, et al. A comparison of gender differences in smoking behaviors, intention to quit, and nicotine dependence among Thai university students. *J Addict* 2018;2018:8081670-70.
- Ritchie H. Who smokes more, men or women? : Our World In Data, 2019. Available: <https://ourworldindata.org/who-smokes-more-men-or-women> [Accessed 15.12.2020].
- Parkinson CM, Hammond D, Fong GT, et al. Smoking beliefs and behavior among youth in Malaysia and Thailand. *Am J Health Behav* 2009;33:366-75.

- 46 Savvides ECG, Christophi CA, Paisi M, *et al.* Factors associated with intent to quit tobacco use in Cyprus adolescents. *Prev Med* 2014;60:83–7.
- 47 Perelman J, Alves J, Pfoertner T-K, *et al.* The association between personal income and smoking among adolescents: a study in six European cities. *Addiction* 2017;112:2248–56.
- 48 WHO. Raise taxes on tobacco: World Health Organization, 2020. Available: [https://www.who.int/tobacco/mpower/raise\\_taxes/en/](https://www.who.int/tobacco/mpower/raise_taxes/en/) [Accessed 18 Dec 2020].
- 49 Dawood OT, Rashan MAA, Hassali MA, *et al.* Knowledge and perception about health risks of cigarette smoking among Iraqi smokers. *J Pharm Bioallied Sci* 2016;8:146–51.
- 50 Schoenaker DAJM, Brennan E, Wakefield MA, *et al.* Anti-Smoking social norms are associated with increased cessation behaviours among lower and higher socioeconomic status smokers: a population-based cohort study. *PLoS One* 2018;13:e0208950.
- 51 Bich NN, Cook M, Johnstone K, *et al.* Students' knowledge and attitudes towards smoke-free universities: changes since enactment of Vietnamese tobacco control legislation. *Asian Pac J Cancer Prev* 2016;17:65–70.
- 52 Yasin SM, Ismail N, Noor NM, *et al.* Gender differences in responses towards anti-smoking messages and policy implementation among future doctors in Malaysia. *Asian Pac J Cancer Prev* 2013;14:303–8.
- 53 Obtel M, Slama K, Abda N. Nurses' Knowledge, Attitudes and Practice in Anti-Smoking Campaigns in Morocco. *OALib* 2014;01:1–8.
- 54 El Fakir S, Serhier Z, Berraho M, *et al.* Knowledge and perceptions of smoking according to income level in Morocco. *Am J Health Promot* 2011;25:387–91.
- 55 Berg MB, Lin L. How effective are campus-wide smoking bans? A comparison of two small colleges. *J Am Coll Health* 2020:1–8.
- 56 Loukas A, Garcia MR, Gottlieb NH. Texas college students' opinions of no-smoking policies, secondhand smoke, and smoking in public places. *J Am Coll Health* 2006;55:27–32.
- 57 Hyland A, Higbee C, Borland R, *et al.* Attitudes and beliefs about secondhand smoke and smoke-free policies in four countries: findings from the International tobacco control four country survey. *Nicotine Tob Res* 2009;11:642–9.
- 58 Hall ME, Williams RD, Hunt B. Assessment of attitudes regarding tobacco-free campus policy and Secondhand smoke exposure among college students, faculty, and staff. *Am J Health Educ* 2015;46:48–57.
- 59 Harris KJ, Stearns JN, Kovach RG, *et al.* Enforcing an outdoor smoking ban on a College campus: effects of a multicomponent approach. *J Am Coll Health* 2009;58:121–6.