BMJ Open Variations of urinary N-acetyl- β -D-glucosaminidase levels and its performance in detecting acute kidney injury under different thyroid hormones levels: a prospectively recruited, observational study

Silin Liang,^{1,2} Dandong Luo,¹ Linhui Hu,^{3,4} Miaoxian Fang,¹ Jiaxin Li,¹ Jia Deng,^{1,2} Heng Fang,^{1,2} Huidan Zhang,^{1,2} Linling He,^{1,2} Jing Xu,^{1,2} Yufan Liang,^{3,4} Chunbo Chen ^{1,2,5}

To cite: Liang S, Luo D, Hu L, et al. Variations of urinary N-acetyl- β -D-glucosaminidase levels and its performance in detecting acute kidney injury under different thyroid hormones levels: a prospectively recruited, observational study. *BMJ Open* 2022;**12**:e055787. doi:10.1136/bmjopen-2021-055787

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-055787>).

SL, DL and LH are joint first authors.

Received 24 July 2021

Accepted 09 February 2022

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Chunbo Chen;
gghccm@163.com

ABSTRACT

Objective Changes in thyroid function will be accompanied by changes in urinary N-acetyl- β -D-glucosaminidase (uNAG) levels. Therefore, whether thyroid hormones interfere the ability of uNAG in detecting acute kidney injury (AKI) has raised concern in patients with critical illness.

Design A prospectively recruited, observational study was performed.

Setting Adults admitted to the intensive care unit of a grade A tertiary hospital in China.

Participants A total of 1919 critically ill patients were enrolled in the study.

Main outcome measures To investigate the variations of the ability of uNAG to detect AKI in patients with critical illness under different thyroid hormones levels (differences in area under the curve (AUC) for uNAG diagnosis and prediction of AKI with different thyroid hormones levels).

Results The bivariate correlation analysis revealed that FT3 and TT3 levels were independently associated with uNAG levels ($p < 0.001$). FT3 and uNAG also showed correlation in multivariable linear regression analysis ($p < 0.001$). After stratification according to the levels of FT3 or TT3, significant variation was observed in the uNAG levels with different quartiles ($p < 0.05$). However, in patients with varying FT3 and TT3 levels, no significant difference was found in the AUCs of uNAG to detect AKI ($p > 0.05$).

Conclusions Even if uNAG levels varied with FT3 and TT3 levels, these hormones did not interfere with uNAG's ability to detect AKI in patients with critical illness.

INTRODUCTION

Acute kidney injury (AKI) is a growing burden in critically ill patients.^{1 2} Early effective treatments can alleviate burden due to high morbidity and mortality of AKI, especially in the intensive care unit (ICU).^{3–6} However, individual differences in serum creatinine

Strengths and limitations of this study

- This is the first study to investigate the influences of thyroid hormones for the performance of urinary N-acetyl- β -D-glucosaminidase (uNAG) in detecting acute kidney injury (AKI).
- This prospectively recruited, observational study was of a long duration and had an adequate sample size.
- Intensive care unit patients were divided into four groups according to quartiles of thyroid hormones levels to represent the population with different thyroid function status.
- We evaluated the ability of uNAG to diagnose and predict AKI primarily by calculating area under the curve.
- A limitation of this study was that the relationship between uNAG and thyroid hormones could not be dynamically assessed because we only measured uNAG and thyroid hormones levels at the time of admission.

(SCr) levels and clinical inconveniences in recording hourly urine output usually lead to delayed initiation of treatments.^{7–9} Therefore, confronting the challenges of AKI management demands a biomarker that can predict AKI onset as early as possible.¹⁰

Urinary N-acetyl- β -D-glucosaminidase (uNAG) is a lysosomal enzyme that is secreted predominantly by the proximal renal tubules.^{11–13} Its macromolecular characteristics prevent it from being filtered by the glomerulus, so high levels of uNAG are likely derived from the kidney.^{14–16} As acute tubular necrosis is one of the histopathological types in AKI,^{17 18} uNAG is considered to

be a sensitive tubular biomarker for AKI.^{11 12} uNAG is valuable in the early detection of AKI, as demonstrated in previous studies.^{19–23} Besides, recent studies have indicated that a combination of functional markers and renal tubular damage markers could improve AKI detection in patients with critical illness,^{24 25} once again proving the diagnostic value of uNAG.

Thyroid disease was thought to be associated with renal disorders and significant changes in renal markers such as creatinine and cystatin C.^{26 27} Many studies have suggested that uNAG levels would increase in patients with hyperthyroidism.^{28–31} High levels of uNAG in patients with hyperthyroidism might be caused by glomerular hyperfiltration, hyperactive tubular secretory function, and proteinuria.^{28 32 33} Abnormal thyroid metabolism is usually present in patients with critical illness.^{34 35} Thus, the question about the effect of fluctuations in thyroid hormones levels on uNAG in patients with critical illness is worth investigating. However, most previous studies have analysed patients with thyroid disease, and the influence of thyroid hormones levels on the performance of uNAG for detecting AKI was not explored.

Therefore, this prospectively recruited, observational study was performed in a large cohort of patients with critical illness to explore which types of thyroid hormones are associated to uNAG levels, and elucidate the accuracy of uNAG in the identification of AKI under different levels of correlated thyroid hormones.

MATERIALS AND METHODS

Study design and participants

This prospectively recruited, observational study was undertaken in the ICU of Guangdong Provincial People's Hospital. All adult patients (≥18 years) admitted between October 2014 and December 2017 were assessed for inclusion. Patients were excluded if they were (1) suffering from hyperthyroidism or hypothyroidism, (2) had preexisting end-stage renal disease or undergoing renal replacement therapy before ICU admission, (3) had a history of renal transplantation or nephrectomy and (4) had missing clinical data or refused consent.

Samples and data collection

Blood samples used to determine thyroid hormones, thyroid-stimulating hormone (TSH), serum albumin, and urine samples used to measure uNAG levels were obtained once from patients within 1 hour after ICU admission. SCr was measured at least once a day as part of routine clinical care to monitor AKI onset. Then, we recorded the following data of patients during their hospital stay: occurrence of AKI within 1 week after ICU admission, AKI grade, length of ICU stay, length of hospital stay, ICU mortality and in-hospital mortality. Comprehensive baseline clinical data were collected prospectively, including age, sex, body mass index, preexisting clinical conditions, admission type, baseline SCr, baseline-estimated glomerular filtration rate (eGFR), SCr at admission, serum

albumin, Acute Physiology and Chronic Health Evaluation II (APACHE II) score, free triiodothyronine (FT3), free thyroxine (FT4), total triiodothyronine (TT3), total thyroxine (TT4), TSH and uNAG. We calculated the eGFR based on the Chronic Kidney Disease (CKD) Epidemiology Collaborative creatinine equation.³⁶

Definitions

In the light of the Kidney Disease Improving Global Outcomes (KDIGO) criteria, AKI was diagnosed if there was an increase in SCr level by 0.3 mg/dL (26.5 mmol/L) within 48 hours, increase in SCr to 1.5 times of the baseline level within 1 week, or urine output <0.5 mL/kg/h for 6 hours.³⁷ Because recording hourly urine output was inconvenient in clinical practice and insensitive after diuretic administration, AKI was diagnosed by detecting changes in SCr.³⁸ A patient's baseline SCr was determined in the following order of preference³⁹: (1) the most recent pre-ICU value between 30 and 365 days before ICU admission, (2) a stable pre-ICU value >365 days for patients aged <40 years (stable defined as within 15% of the lowest ICU measurement) before ICU admission, (3) pre-ICU value >365 days before ICU admission and lower than the initial SCr value at ICU admission, (4) a pre-ICU value (between 3 and 39 days before ICU admission) less than or equal to the initial on-admission SCr value to ICU and not distinctly during AKI, and (5) the lowest SCr on initial admission to the ICU, last ICU value, or minimum value at follow-up to 365 days. AKI diagnosed at ICU admission or within 1 week after ICU admission was defined as established AKI or later-onset AKI, respectively. Mild AKI was defined as KDIGO stage 1 and severe AKI was defined as KDIGO stage 2 or 3.^{37 40} The primary outcome was the presence of AKI, and secondary outcomes included length of ICU and hospital stay, as well as ICU and hospital mortality.

Patient groups

Based on the results of the bivariate correlation analysis between uNAG and thyroid hormones, patients were stratified into four groups according to the quartiles of relevant thyroid hormones levels. Baseline characteristics, outcome and performance of uNAG in determining AKI were compared among these groups. Besides, to shut off the influences of renal function, we also divided patients into two subgroups, namely, AKI group and non-AKI group, and these subgroups were also split into four quartiles according to relevant thyroid hormones.

Laboratory methods

All specimen assays were accomplished by the central laboratory of Guangdong Provincial People's Hospital within 24 hours after collection. SCr, uNAG and serum albumin were analysed using the UniCel DxC 800 Synchron System (Beckman Coulter, Brea, California, USA). Values of uNAG were normalised to that of the urinary creatinine concentrations. The coefficients of inter-assay and intra-assay variations for uNAG were both ≤10%. Thyroid

hormones were measured by chemiluminescent immunoassay using Unicel DxI800 Synchron System (Beckman Coulter). The normal value ranges of TT3, TT4, FT3, and FT4 are 1.34–2.73 nmol/L, 78.40–158.40 nmol/L, 3.80–6.00 pmol/L and 7.50–21.10 pmol/L, respectively.

Patient and public involvement

Patients were not involved in the study.

Statistical analysis

A two-tailed $p < 0.05$ was regarded as significant. Continuous variables were summarised as median (IQR, IQR) depending on their abnormal distribution and were compared using the non-parametric tests. Categorical variables were reported as frequency (percentage) and compared using the χ^2 test or Fisher's exact test. Bivariate correlation analysis was used to examine the association between two variables to confirm representative indicators of thyroid function for further analysis. Multivariable linear regression analysis with a stepwise variable selection was also used to assess the relationship between uNAG and other variables. Receiver operating characteristic curve analysis and area under the curve (AUC), as assessed using the Hanley-McNeil method, were used to evaluate the ability of uNAG to detect AKI. The optimal cut-off value for AKI detection was determined with Youden's index. All statistical analyses were performed using SPSS V.23.0 (SPSS) and MedCalc V.18.2.1 (MedCalc Software, Ostend, Belgium) software programs.

RESULTS

Participants

In strict accordance with the selection criteria, 138 patients were excluded from among 2057 patients; finally, 1919 patients with critical illness were included in the

Figure 1 Recruitment of patients into the study. Established AKI indicated the diagnosis of AKI at ICU admission. Later-onset AKI was defined as no AKI diagnosis at ICU admission but reaching the KDIGO criteria within 1 week after admission. AKI, acute kidney injury; ESRD, end-stage renal disease; ICU, Intensive care unit; KDIGO, Kidney Disease Improving Global Outcomes; RRT, renal replacement therapy.

analysis (figure 1). The total number of patients with AKI was 454, of which 202 had established AKI and 252 had later-onset AKI. The incidence of total AKI was approximately 23.7%.

Factors related to uNAG

The bivariate correlation between TT3, TT4, FT3 and FT4 and uNAG were determined to identify indicators for thyroid function that influenced uNAG (online supplemental table 1). The results revealed that both FT3 and TT3 were significantly related to uNAG ($p < 0.001$). FT3 and TT3 showed consistent negative correlation with uNAG, so we could conclude that uNAG levels decreased at higher FT3 and TT3 levels. However, FT4, TT4 and TSH levels were not related to uNAG ($p > 0.05$). According to the results of the bivariate correlation analysis, the factors that are correlated with uNAG were included in the multivariable linear regression analysis. In multivariable linear regression analysis (online supplemental table 2), we found uNAG was negatively correlated with FT3, even though the correlation was weak (standardised $\beta = -2.893$, $p = 0.016$). Based on the above results, we divided the patients into four quartile groups according to FT3 or TT3 levels.

Patient's characteristics and outcomes under different FT3 or TT3 levels

Table 1 shows the characteristics and outcomes of patients under different FT3 levels. From the table, patients with higher FT3 levels were younger, had fewer complications, were less likely to require emergency surgery, had a lower APACHE II score, had a higher THS levels and had a better basic renal function. Essentially, high thyroid hormones levels help the body respond better to critical situations. Patient outcomes were also in line with this explanation. Patients with high FT3 levels had a lower incidence of AKI. Even if AKI occurs, it has a milder form. A downward trend was shown in uNAG levels with the increase of FT3 and TT3 levels. Table 2 indicates that patients with different TT3 levels demonstrate comparable characteristics and outcomes.

Variations of uNAG levels under different FT3 or TT3 levels

Although the variations of uNAG levels were moderate, a significant difference could be demonstrated among different FT3 levels (figure 2A). Under different TT3 levels, uNAG levels also showed significant differences, but changes in uNAG levels in quartiles III and IV were not statistically significant (figure 2B). In this study, as the level of FT3 or TT3 increases, the level of uNAG gradually decreases. We additionally analysed the changes in the levels of uNAG in the quartiles of FT4, TT4 and TSH. It was found that the level of uNAG had no obvious regularity in different levels of FT4, TT4 and TSH (online supplemental figure 1).

Variations of uNAG levels in the non-AKI group and AKI group under different FT3 or TT3 levels

To eliminate influences of changes in renal function, the patients were also divided into non-AKI and AKI subgroups and then categorised into four quartiles according to the

Table 1 Characteristics and outcomes of patients according to quartiles of FT3

Variables	Quartile I	Quartile II	Quartile III	Quartile IV	P value
FT3	2.98 (2.63–3.21)	3.66 (3.52–3.80)	4.13 (4.00–4.25)	4.76 (4.54–5.07)	<0.001
No	486	482	473	478	
Baseline characteristics					
Age, years	59 (47–70)	54 (43–65)	53 (40–63)	49 (36–59)	<0.001
Male sex, n (%)	262 (53.9)	253 (52.5)	244 (51.6)	251 (52.5)	0.912
BMI, kg/m ²	22.20 (20.82–24.20)	22.20 (20.52–24.00)	22.46 (21.05–24.76)	22.36 (20.57–24.89)	0.085
Preexisting clinical conditions					
Hypertension, n (%)	101 (20.8)	91 (18.9)	89 (18.8)	72 (15.1)	0.138
DM, n (%)	52 (10.7)	40 (8.3)	35 (7.4)	26 (5.4)	0.025
CKD, n (%)	43 (8.8)	13 (2.7)	6 (1.3)	4 (0.8)	<0.001
CHD, n (%)	23 (4.7)	13 (2.7)	11 (2.3)	10 (2.1)	0.062
Stroke, n (%)	90 (18.5)	68 (14.1)	52 (11.0)	41 (8.6)	<0.001
CHF, n (%)	24 (4.9)	11 (2.3)	5 (1.1)	2 (0.4)	<0.001
Malignancy, n (%)	88 (18.1)	59 (12.2)	51 (10.8)	41 (8.6)	<0.001
COPD, n (%)	23 (4.7)	5 (1.0)	3 (0.6)	7 (1.5)	<0.001
Chronic liver disease, n (%)	11 (2.3)	3 (0.6)	3 (0.6)	0 (0.0)	0.001
Sepsis, n (%)	145 (29.8)	51 (10.6)	21 (4.4)	18 (3.8)	<0.001
Admission type					
Elective surgical, n (%)	304 (62.6)	407 (84.4)	415 (87.7)	418 (87.4)	<0.001
Emergency surgical, n (%)	79 (16.3)	28 (5.8)	20 (4.2)	13 (2.7)	
Medical, n (%)	103 (21.2)	47 (9.8)	38 (8.0)	47 (9.8)	
uNAG, U/g Cr	34.09 (20.78–58.62)	27.01 (15.39–41.56)	23.61 (15.30–36.25)	21.04 (13.97–31.92)	<0.001
Baseline eGFR, ml/min/1.73 m ²	98.35 (79.56–110.61)	103.31 (99.89–113.39)	103.10 (90.37–115.23)	104.67 (92.29–117.55)	<0.001
Baseline SCr, umol/L	64.34 (52.20–80.75)	61.85 (51.28–75.33)	64.00 (53.54–75.30)	65.00 (54.88–77.43)	0.091
Scr at ICU admission umol/L	78.02 (61.88–105.06)	70.00 (58.95–85.68)	71.96 (59.90–87.00)	71.80 (59.86–88.50)	<0.001
Serum albumin, g/L	28.70 (25.00–32.45)	30.90 (27.49–34.20)	32.40 (29.20–34.90)	34.55 (31.90–37.80)	<0.001
TSH uIU/ml	1.00 (0.52–2.05)	1.22 (0.64–2.10)	1.33 (0.78–2.48)	1.38 (0.74–2.42)	<0.001
APACHE II score	12 (8–20)	9 (6–12)	8 (5–12)	6 (4–10)	<0.001
Primary outcomes					
Total AKI, n (%)	214 (44.0)	99 (20.5)	81 (17.1)	60 (12.6)	<0.001
Established AKI, n (%)	123 (25.3)	41 (8.5)	23 (4.9)	15 (3.1)	<0.001
Later-onset AKI, n (%)	91 (18.7)	58 (12.0)	58 (12.3)	45 (9.4)	<0.001

Continued

Table 1 Continued

Variables	Quartile I	Quartile II	Quartile III	Quartile IV	P value
Grade of AKI					<0.001
Non-AKI, n (%)	272 (56.0)	383 (79.5)	392 (82.9)	418 (87.4)	
Mild AKI, n (%)	132 (27.2)	75 (15.6)	62 (13.1)	53 (11.1)	
Severe AKI, n (%)	82 (16.9)	24 (5.0)	19 (4.0)	7 (1.5)	

The non-normally distributed continuous variables are expressed as median (25–75th percentile [IQR]). Categorical variables are expressed as n (%). Established AKI, defined as diagnosis of AKI at ICU admission; later-onset AKI, indicated no AKI diagnosis at ICU admission but reaching the KDIGO criteria within 1 week after admission; mild-AKI: defined as reaching KDIGO stage 1 diagnostic criteria of AKI; severe-AKI, defined as reaching KDIGO stage 2 or stage 3 diagnostic criteria of AKI. P value for global comparisons among groups by rank sum test and χ^2 test for continuous and categorical variables, respectively. AKI, acute kidney injury; APACHE II score, Acute Physiology and Chronic Health Evaluation II score; BMI, body mass index; CHD, coronary heart disease; CHF, chronic heart failure; CKD, chronic kidney disease; COPD, chronic obstructive pulmonary disease; DM, diabetes mellitus; eGFR, estimated glomerular filtration rate; FT3, free triiodothyronine; ICU, intensive care unit; KDIGO, Kidney Disease: Improving Global Outcomes; n, sample size; SCr, serum creatinine; TSH, thyroid-stimulating hormone; uNAG, urinary N-acetyl- β -D-glucosaminidase.

levels of FT3 or TT3 (online supplemental figure 2). The trend of uNAG in the non-AKI and AKI groups was consistent, which decreased with the increase in FT3 or TT3 levels. This showed that thyroid hormones independently affect the levels of uNAG.

Performance of uNAG in detecting AKI in patients with different FT3 or TT3 levels at ICU admission

We quantified the performance of uNAG to detect AKI based on AUC values (table 3). In the entire cohort, uNAG detected total AKI with an AUC of 0.682, diagnosed established AKI with an AUC of 0.700, and predicted later-onset AKI with an AUC of 0.668.

To explore the effects of thyroid hormones on the efficacy of uNAG to detect AKI, the patients were divided into four quartiles according to FT3 levels and TT3 levels at ICU admission, and the AUC values of uNAG were calculated and compared with detect AKI between different groups (table 3). From the results in table 3, the performance of uNAG to detect total AKI, diagnose established AKI, and predict later-onset AKI was not distinct under different thyroid hormones levels. No significant difference in the AUC was observed between any two groups, whether divided into four quartiles by FT3 levels or by TT3 levels. On the contrary, the optimal cut-off value of uNAG for detecting total AKI, diagnosing established AKI, and predicting later-onset AKI showed fair discrimination in different quartiles based on FT3 or TT3 levels. However, changes in the cut-off values of uNAG were not as simple as uNAG increasing with higher FT3 or TT3 level.

DISCUSSION

Recently, combining functional and renal tubular injury biomarkers as a new strategy to detect AKI has attracted research attention.^{24 41} uNAG was considered an important biomarker of renal tubular damage in combination with AKI biomarkers.²⁴ This prospectively recruited cohort study investigated the variations of the ability of uNAG to detect AKI in patients with critical illness under different thyroid hormones levels. Patients were divided into four quartile groups according to FT3 and TT3 levels based on the results of the bivariate correlation analysis and multivariate linear regression analysis between uNAG and thyroid hormones, and the AUC values of uNAG in diagnosing and predicting AKI were calculated for each group. In this study, significant differences were found uNAG levels of patients between different quartiles, but the performance of uNAG in detecting AKI was not different among patients under different quartiles.

As a sensitive marker of renal tubular damage,^{11 12} elevated uNAG levels were considered an early sign of AKI.^{19–22} However, uNAG levels were also influenced by non-renal factors as reported previously.^{23 29–31 33 42–45} The animal experiment results of Lapointe *et al* showed

Table 2 Characteristics and outcomes of patients according to quartiles of TT3

Variables	Quartile I	Quartile II	Quartile III	Quartile IV	P value
TT3	0.63 (0.51–0.72)	0.90 (0.85–0.95)	1.09 (1.04–1.15)	1.38 (1.29–1.51)	<0.001
No	484	482	486	467	
Baseline characteristics					
Age, years	59 (46–69)	54 (42–65)	53 (41–63)	50 (37–60)	<0.001
Male sex, n (%)	285 (58.9)	238 (49.4)	236 (48.6)	251 (53.7)	0.004
BMI, kg/m ²	22.19 (20.56–24.34)	22.39 (20.83–24.57)	22.41 (20.81–24.60)	22.32 (20.50–24.18)	0.206
Preexisting clinical conditions					
Hypertension, n (%)	114 (23.6)	86 (17.8)	73 (15.0)	80 (17.1)	0.005
DM, n (%)	57 (11.8)	45 (9.3)	27 (5.6)	24 (5.1)	<0.001
CKD, n (%)	35 (7.2)	17 (3.5)	8 (1.6)	6 (1.3)	<0.001
CHD, n (%)	25 (5.2)	11 (2.3)	9 (1.9)	12 (2.6)	0.010
Stroke, n (%)	94 (19.4)	59 (12.2)	61 (12.6)	37 (7.9)	<0.001
CHF, n (%)	24 (5.0)	5 (1.0)	10 (2.1)	3 (0.6)	<0.001
Malignancy, n (%)	84 (17.4)	65 (13.5)	54 (11.1)	36 (7.7)	<0.001
COPD, n (%)	20 (4.1)	8 (1.7)	6 (1.2)	4 (0.9)	0.001
Chronic liver disease, n (%)	9 (1.9)	6 (1.2)	2 (0.4)	0 (0.0)	0.002
Sepsis, n (%)	151 (31.2)	40 (8.3)	27 (5.6)	17 (3.6)	<0.001
Admission type					
Elective surgical, n (%)	279 (57.6)	419 (86.9)	426 (87.7)	420 (89.9)	
Emergency surgical, n (%)	86 (17.8)	21 (4.4)	16 (3.3)	17 (3.6)	
Medical, n (%)	119 (24.6)	42 (8.7)	44 (9.1)	30 (6.4)	
uNAG, U/g Cr	34.75 (20.69–57.78)	25.86 (16.80–42.18)	23.48 (14.94–35.99)	21.13 (13.88–33.73)	<0.001
Baseline eGFR, ml/min/1.73 m ²	99.05 (82.74–113.07)	100.03 (89.21–112.97)	103.72 (91.64–114.05)	105.94 (92.32–117.21)	<0.001
Baseline SCr, umol/L	64.00 (52.25–79.90)	63.50 (53.98–78.00)	63.40 (51.95–75.20)	64.00 (54.00–76.70)	0.447
SCr at ICU admission umol/L	78.05 (62.93–102.00)	71.00 (59.33–89.47)	71.00 (58.68–86.47)	71.50 (59.50–87.00)	<0.001
Serum albumin, g/L	28.45 (24.93–32.25)	30.80 (27.50–34.20)	32.65 (29.50–35.24)	34.60 (31.60–37.10)	<0.001
TSH uIU/ml	0.91 (0.49–1.81)	1.21 (0.64–2.12)	1.30 (0.78–2.27)	1.50 (0.79–2.91)	<0.001
APACHE II score	13 (8–20)	9 (6–12)	8 (5–12)	7 (4–10)	<0.001
Primary outcomes					
Total AKI, n (%)	221 (45.7)	101 (21.0)	69 (14.2)	63 (13.5)	<0.001
Established AKI, n (%)	119 (24.6)	41 (8.5)	27 (5.6)	15 (3.2)	<0.001
Later-onset AKI, n (%)	102 (21.1)	60 (12.4)	42 (8.6)	48 (10.3)	<0.001

Continued

Table 2 Continued				
Variables	Quartile I	Quartile II	Quartile III	Quartile IV
Grade of AKI, n (%)				
Non-AKI, n (%)	263 (54.3)	381 (79.0)	417 (85.8)	404 (86.5)
Mild AKI, n (%)	140 (28.9)	75 (15.6)	52 (10.7)	55 (11.8)
Severe AKI, n (%)	81 (16.7)	26 (5.4)	17 (3.5)	8 (1.7)

The non-normally distributed continuous variables are expressed as median (25–75th percentile (IQR)). Categorical variables are expressed as n (%). Established AKI, defined as diagnosis of AKI at ICU admission; later-onset AKI, indicated no AKI diagnosis at ICU admission but reaching the KDIGO criteria within 1 week after admission; mild-AKI: defined as reaching KDIGO stage 1 diagnostic criteria of AKI; severe-AKI, defined as reaching KDIGO stage 2 or stage 3 diagnostic criteria of AKI. P value for global comparisons among groups by rank sum test and χ^2 test for continuous and categorical variables, respectively. AKI, acute kidney injury; APACHE II score, Acute Physiology and Chronic Health Evaluation II score; BMI, body mass index; CHD, coronary heart disease; CKD, chronic kidney disease; COPD, chronic obstructive pulmonary disease; DM, diabetes mellitus; eGFR, estimated glomerular filtration rate; ICU, intensive care unit; KDIGO, Kidney Disease: Improving Global Outcomes; n, sample size; SCr, serum creatinine; TSH, thyroid-stimulating hormone; TT3, total triiodothyronine; uNAG, urinary N-acetyl- β -D-glucosaminidase.

Figure 2 The median of uNAG levels in patients with different FT3 or TT3 levels. (A) FT3: I vs II $p < 0.05$, I vs III $p < 0.05$, I vs IV $p < 0.05$; II vs III $p < 0.05$, II vs IV $p < 0.05$; III vs IV $p < 0.05$; (B) TT3: I vs II $p < 0.05$, I vs III $p < 0.05$, I vs IV $p < 0.05$; II vs III $p < 0.05$, II vs IV $p < 0.05$; III vs IV $p > 0.05$. FT3, free triiodothyronine; TT3, total triiodothyronine; uNAG, urinary N-acetyl- β -D-glucosaminidase.

that the uNAG levels of hyperthyroid cats without CKD and hyperthyroid cats with CKD were higher than those of healthy cats and the uNAG level of hyperthyroid cats without CKD decreased after treatment with methimazole.³³ Tominaga *et al* also found similar situation in patients with hyperthyroidism.³¹ Their results suggested that patients with hyperthyroidism had higher uNAG levels than normal people and patients with diabetes without kidney damage. Another study further proved that elevated uNAG level not only occurred in Graves' disease but also in subacute thyroiditis and silent thyroiditis.³⁰ Undoubtedly, the aforementioned studies indicated that excessive thyroid hormones level could cause an increase in uNAG levels.

uNAG is a lysosomal hydrolase that is derived from the lysosome of the proximal tubule cells of the kidney.^{11–13} Thyroid hormones could increase uNAG levels by changing the lysosomal function or affecting kidney function. Lysosomal enzyme activity is dependent on the state of thyroid function.^{46 47} The possible hypothetical mechanism was that thyroid hormones stimulated the biosynthesis of lysosomal enzymes,⁴⁸ whereas excessive thyroid hormones could lead to lysosomal enzyme leak because changes in lysosomal

Table 3 Detection of AKI using uNAG by different quartiles

	Total AKI				Established AKI				Later-onset AKI			
	AUC-ROC	95% CI	Cut-off	P value	AUC-ROC	95% CI	Cut-off	P value	AUC-ROC	95% CI	Cut-off	P value
Total	0.682±0.015	0.661 to 0.703	41.11	<0.001	0.700±0.021	0.677 to 0.722	41.11	<0.001	0.668±0.019	0.645 to 0.690	22.02	<0.001
FT3												
Quartiles I	0.650±0.025	0.606 to 0.692	32.27	<0.001	0.663±0.031	0.614 to 0.709	32.27	<0.001	0.632±0.033	0.580 to 0.682	23.17	<0.001
Quartiles II	0.637±0.032	0.592 to 0.680	53.23	<0.001	0.651±0.050	0.604 to 0.697	76.22	0.003	0.627±0.039	0.580 to 0.672	21.93	0.001
Quartiles III	0.660±0.035	0.615 to 0.703	48.36	<0.001	0.649±0.059	0.601 to 0.695	20.34	0.012	0.664±0.042	0.618 to 0.708	48.36	<0.001
Quartiles IV	0.668±0.038	0.624 to 0.711	40.23	<0.001	0.652±0.051	0.606 to 0.697	20.34	0.003	0.674±0.047	0.629 to 0.716	33.67	<0.001
TT3												
Quartiles I	0.636±0.025	0.592 to 0.679	65.78	<0.001	0.663±0.032	0.613 to 0.710	65.78	<0.001	0.605±0.033	0.553 to 0.656	65.78	0.002
Quartiles II	0.654±0.031	0.610 to 0.697	21.22	<0.001	0.668±0.048	0.620 to 0.712	58.53	<0.001	0.645±0.037	0.599 to 0.690	21.93	<0.001
Quartiles III	0.656±0.038	0.612 to 0.698	40.23	<0.001	0.664±0.054	0.618±0.708	24.14	0.002	0.651±0.050	0.605 to 0.694	40.23	0.003
Quartiles IV	0.693±0.036	0.649 to 0.735	45.00	<0.001	0.651±0.061	0.603 to 0.697	20.34	0.014	0.706±0.043	0.662 to 0.748	45.00	<0.001

Established AKI, defined as diagnosis of AKI at ICU admission; later-onset AKI, indicated no AKI diagnosis at ICU admission but reaching the KDIGO criteria within 1 week after admission.

FT3: total AKI: I vs II: Z=0.320, p=0.749; I vs III: Z=-0.232, p=0.816; I vs IV: Z=-0.396, p=0.692; II vs III: Z=0.485, p=0.628; II vs IV: Z=0.624, p=0.533; III vs IV: Z=0.155, p=0.877; established AKI: I vs II: Z=0.204, p=0.838; I vs III: Z=0.210, p=0.834; I vs IV: Z=0.184, p=0.854; II vs III: Z=0.026, p=0.979; II vs IV: Z=-0.014, p=0.989; III vs IV: Z=-0.039, p=0.969; later-onset AKI: I vs II: Z=0.098, p=0.922; I vs III: Z=-0.599, p=0.549; I vs IV: Z=-0.731, p=0.465; II vs III: Z=-0.646, p=0.519; II vs IV: Z=-0.770, p=0.442; III vs IV: Z=-0.159, p=0.874.

TT3: total AKI: I vs II: Z=-0.452, p=0.651; I vs III: Z=-0.440, p=0.660; I vs IV: Z=-1.301, p=0.193; II vs III: Z=0.055, p=0.956; II vs IV: Z=0.219, p=0.827; III vs IV: Z=0.160, p=0.873; later-onset AKI: I vs II: Z=-0.087, p=0.931; I vs III: Z=-0.016, p=0.987; I vs IV: Z=0.174, p=0.862; II vs III: Z=0.055, p=0.956; II vs IV: Z=0.219, p=0.827; III vs IV: Z=0.160, p=0.873; later-onset AKI: I vs II: Z=-0.807, p=0.294; I vs III: Z=-0.768, p=0.443; I vs IV: Z=-1.863, p=0.062; II vs III: Z=-0.097, p=0.923; II vs IV: Z=-1.075, p=0.253; III vs IV: Z=-0.834, p=0.4.

AKI, acute kidney injury; AUC-ROC, area under the receiver operating characteristic curve; FT3, free triiodothyronine; ICU, intensive care unit; KDIGO, Kidney Disease: Improving Global Outcomes; TT3, total triiodothyronine; uNAG, urinary N-acetyl-β-D-glucosaminidase.

membrane permeability resulted from the increase in the production of reactive oxygen species.^{49–51} Thyroid hormones might cause changes in uNAG levels by affecting kidney function and damaging kidney structure. In case of hyperthyroidism, hyperfiltration, proteinuria, hypertrophy and hyperplasia altogether damage the glomerular basement membrane and renal tubules.^{28 32 33} Animal experiments have found that the increased activity of angiotensin II will lead to interstitial fibrosis and matrix protein accumulation by inducing an increase in the production of transforming growth factor β 1 and platelet derived growth factor and ultimately resulting in tubulointerstitial nephritis.^{52–54} An increase in angiotensin II levels should also have the same influence in patients with hyperthyroidism.⁵⁵

However, compared with previous studies, this study presented opposite results in exploring the relationship between thyroid hormones and uNAG. Our results indicated that FT3 and TT3 levels were negatively correlated with uNAG in the bivariate correlation analysis. In the multivariate linear regression analysis, similar conclusions were drawn in the correlation between FT3 and uNAG. This finding could be explained by the fact that triiodothyronine was the main physiologically active substance. Triiodothyronine has been shown more biologically active than tetraiodothyronine in terms of pathophysiological effects.⁵⁶ Even in critically ill patients without thyroid disease, thyroid hormones levels might be abnormal, and the incidence in AKI patients was as high as 80%.⁵⁷ Among them, the reduction of triiodothyronine was the most common.⁵⁸ Low thyroid hormones levels have been shown to be a risk factor for poor prognosis in critically ill patients,⁵⁸ which is consistent with this study finding that critically ill patients with low thyroid hormones levels had higher APACHE II score. uNAG levels decreased with the increase in FT3 and TT3 levels, and significant differences were noted in uNAG levels between different FT3 and TT3 levels. Even if the study patients were divided into AKI and non-AKI groups for analysis, we still obtained the same results. The most important reason for the contrasting results between our study and previous studies might be the difference in research objects. Previous studies have mainly explored the effect of excessive thyroid hormones on uNAG levels caused by thyroid diseases. However, the study excluded patients with thyroid disease to explore changes in uNAG levels in patients with critical illness under different thyroid function states. Our analysis found that patients whose thyroid function remained intact despite having a severe disease had lower APACHE II score and incidence of AKI. This might explain our study finding that patients with higher thyroid hormones levels had lower uNAG levels.

This study did not display a significant variable difference in the AUC values of uNAG in discovering

AKI among patients with different FT3 or TT3 levels. In the absence of thyroid disease, the results suggested that uNAG levels in patients with critical illness were negatively correlated with FT3 and TT3 levels, and the ability of uNAG to diagnose and predict AKI was not interfered by thyroid hormones.

This study still has some limitations. First, we used thyroid hormones levels at admission rather than continuous thyroid hormones variability to test our hypotheses. Second, our study has not verified the influences of thyroid hormones on the ability of uNAG to detect AKI in patients with thyroid diseases. We thought a large sample size of patients with thyroid diseases was needed to reveal this influence.

CONCLUSION

Although FT3 and TT3 affected the level of uNAG, these hormones did not affect the performance of uNAG to recognise AKI in patients with critical illness.

Author affiliations

¹Department of Intensive Care Unit of Cardiovascular Surgery, Guangdong Cardiovascular Institute, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, 96 Dongchuan Road, Guangzhou 510080, Guangdong Province, People's Republic of China

²Department of Critical Care Medicine, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, 106 Zhongshan Er Road, Guangzhou 510080, Guangdong Province, People's Republic of China

³Department of Critical Care Medicine, Maoming People's Hospital, 101 Weimin Road, Maoming 525000, Guangdong Province, People's Republic of China

⁴Center of Scientific Research, Maoming People's Hospital, 101 Weimin Road, Maoming 525000, Guangdong Province, People's Republic of China

⁵The Second School of Clinical Medicine, Southern Medical University, Guangzhou, Guangdong Province, People's Republic of China

Acknowledgements The authors thank all the doctors, nurses, technicians and patients at the Guangdong Provincial People's Hospital for their dedication in the study.

Contributors CBC is the guarantor of this work. CBC, SLL, DDL and LHH equally contributed to the design of the research. SLL and LHH equally contributed to analyze and interpret the data. CBC and DDL contributed to the conception of the research and critically revised the manuscript. SLL, DDL, LHH, MXF, JXL, JD, HF, HDZ, LLH, JX and YFL performed the research and collected data. All authors contributed to the acquisition and analysis of the data, drafted the manuscript, and agreed to be fully accountable for ensuring the integrity and accuracy of the work. All authors read and approved the final manuscript.

Funding The study was supported by Chunbo Chen's research grants from the Science and Technology Planning Project of Guangdong Province (Key Programme) (2014B020212023), the National Natural Science Foundation of China (82172162), the Guangzhou Livelihood Science and Technology Programme (201803010058) and the Major Program of Summit Project, Guangdong Province High-level Hospital Construction Project of Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences (DFJH2020028). Author Linhui Hu is currently receiving a grant (zx2020017) from the High-level Hospital Construction Research Project of Maoming People's Hospital.

Competing interests The authors have no conflicts of interest to disclose.

Patient and public involvement statement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval The current study received the approval of ethics committee of the Guangdong Provincial People's Hospital (No.GDREC2015396H(R1)). The ethics

committee of the Guangdong Provincial People's Hospital supervised the study, including the study design, protocol, ethical issue and data and sample collection. Written informed consent was obtained from each patient or from the appropriate guardian. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. The datasets generated and/or analysed during this study are not publicly available, owing to currently ongoing research studies, but the data are available from the corresponding author on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Chunbo Chen <http://orcid.org/0000-0001-5662-497X>

REFERENCES

- Arora T, Martin M, Grimshaw A, *et al.* Prediction of outcomes after acute kidney injury in hospitalised patients: protocol for a systematic review. *BMJ Open* 2020;10:e042035.
- Sako K, Furuichi K, Yamamura Y, *et al.* Association between the recurrence period of acute kidney injury and mortality: a single-centre retrospective observational study in Japan. *BMJ Open* 2019;9:e023259.
- Thakar CV, Christianson A, Freyberg R, *et al.* Incidence and outcomes of acute kidney injury in intensive care units: a Veterans administration study. *Crit Care Med* 2009;37:2552–8.
- Zarbock A, Kellum JA, Schmidt C, *et al.* Effect of early vs delayed initiation of renal replacement therapy on mortality in critically ill patients with acute kidney injury: the ELAIN randomized clinical trial. *JAMA* 2016;315:2190–9.
- Xu X, Nie S, Liu Z, *et al.* Epidemiology and clinical correlates of AKI in Chinese hospitalized adults. *Clin J Am Soc Nephrol* 2015;10:1510–8.
- Hou FF, Yang X. Advances in the management of acute cardiorenal syndrome in China: biomarkers for predicting development and outcomes. *Kidney Dis* 2017;2:145–50.
- Jo SK, Rosner MH, Okusa MD. Pharmacologic treatment of acute kidney injury: why drugs haven't worked and what is on the horizon. *Clin J Am Soc Nephrol* 2007;2:356–65.
- Martins CB, De Bels D, Honore PM, *et al.* Early prediction of acute kidney injury by machine learning: should we add the urine output criterion to improve this new tool? *J Transl Int Med* 2020;8:201–2.
- Xu X, Wang L, Xiong M, *et al.* New criterion to evaluate acute-on-chronic kidney injury based on the creatinine reference change. *Am J Nephrol* 2020;51:453–62.
- Yang Y, Ge B, Liu Y, *et al.* The efficacy of biomarkers in the diagnosis of acute kidney injury secondary to liver cirrhosis. *Medicine* 2021;100:e25411.
- Charlton JR, Portilla D, Okusa MD. A basic science view of acute kidney injury biomarkers. *Nephrol Dial Transplant* 2014;29:1301–11.
- Ostermann M, Joannidis M. Acute kidney injury 2016: diagnosis and diagnostic workup. *Crit Care* 2016;20:299.
- Vaidya VS, Waikar SS, Ferguson MA, *et al.* Urinary biomarkers for sensitive and specific detection of acute kidney injury in humans. *Clin Transl Sci* 2008;1:200–8.
- Sprengle P, Russo P. Molecular markers for ischemia, do we have something better than creatinine and glomerular filtration rate? *Arch Esp Urol* 2013;66:99–114.
- Vaidya VS, Ferguson MA, Bonventre JV. Biomarkers of acute kidney injury. *Annu Rev Pharmacol Toxicol* 2008;48:463–93.
- Yoo J-J, Kwon JH, Kim YS, *et al.* The role of urinary N-acetyl-β-D-glucosaminidase in cirrhotic patients with acute kidney injury: multicenter, prospective cohort study. *J Clin Med* 2021;10:4328.
- Ricci Z, Cruz DN, Ronco C. Classification and staging of acute kidney injury: beyond the rifle and akin criteria. *Nat Rev Nephrol* 2011;7:201–8.
- Wen Y, Yang C, Menez SP, *et al.* A systematic review of clinical characteristics and histologic descriptions of acute tubular injury. *Kidney Int Rep* 2020;5:1993–2001.
- Cheng B, Jin Y, Liu G, *et al.* Urinary N-acetyl-beta-D-glucosaminidase as an early marker for acute kidney injury in full-term newborns with neonatal hyperbilirubinemia. *Dis Markers* 2014;2014:1–6.
- Doi K, Negishi K, Ishizu T, *et al.* Evaluation of new acute kidney injury biomarkers in a mixed intensive care unit. *Crit Care Med* 2011;39:2464–9.
- Liangos O, Perianayagam MC, Vaidya VS, *et al.* Urinary N-acetyl-beta-(D)-glucosaminidase activity and kidney injury molecule-1 level are associated with adverse outcomes in acute renal failure. *J Am Soc Nephrol* 2007;18:904–12.
- Peng Z-Y. The biomarkers for acute kidney injury: a clear road ahead? *J Transl Int Med* 2016;4:95–8.
- Wang L, Deng Y, Zhai Y, *et al.* Impact of blood glucose levels on the accuracy of urinary N-acetyl-β-D-glucosaminidase for acute kidney injury detection in critically ill adults: a multicenter, prospective, observational study. *BMC Nephrol* 2019;20:186.
- Deng Y, Chi R, Chen S, *et al.* Evaluation of clinically available renal biomarkers in critically ill adults: a prospective multicenter observational study. *Crit Care* 2017;21:46.
- Deng Y, Ma J, Hou Y, *et al.* Combining serum cystatin C and urinary N-acetyl-beta-D-glucosaminidase improves the precision for acute kidney injury diagnosis after resection of intracranial space-occupying lesions. *Kidney Blood Press Res* 2020;45:142–56.
- Simeoni M, Cerantonio A, Pastore I, *et al.* The correct renal function evaluation in patients with thyroid dysfunction. *J Endocrinol Invest* 2016;39:495–507.
- Grecu M, Foti DP, Aversa A, *et al.* Cystatin C, a controversial biomarker in hypothyroid patients under levothyroxine therapy: THYRenal, a pilot cohort observational study. *J Clin Med* 2020;9. doi:10.3390/jcm9092958. [Epub ahead of print: 13 09 2020].
- Basu G, Mohapatra A. Interactions between thyroid disorders and kidney disease. *Indian J Endocrinol Metab* 2012;16:204–13.
- Komosińska-Vashev K, Olczyk K, Kozma EM, *et al.* Graves' disease-associated changes in the serum lysosomal glycosidases activity and the glycosaminoglycan content. *Clin Chim Acta* 2003;331:97–102.
- Nakamura S, Ishiyama M, Kosaka J, *et al.* Urinary N-acetyl-beta-D-glucosaminidase (Nag) activity in patients with Graves' disease, subacute thyroiditis, and silent thyroiditis: a longitudinal study. *Endocrinol Jpn* 1991;38:303–8.
- Tominaga M, Fujiyama K, Hoshino T, *et al.* Urinary N-acetyl-beta-D-glucosaminidase in the patients with hyperthyroidism. *Horm Metab Res* 1989;21:438–40.
- Iglesias P, Bajo MA, Selgas R, *et al.* Thyroid dysfunction and kidney disease: an update. *Rev Endocr Metab Disord* 2017;18:131–44.
- Lapointe C, Bélanger M-C, Dunn M, *et al.* N-Acetyl-Beta-D-Glucosaminidase index as an early biomarker for chronic kidney disease in cats with hyperthyroidism. *J Vet Intern Med* 2008;22:1103–10.
- Economidou F, Douka E, Tzanela M, *et al.* Thyroid function during critical illness. *Hormones* 2011;10:117–24.
- Zaloga GP, Chernow B, Smallridge RC, *et al.* A longitudinal evaluation of thyroid function in critically ill surgical patients. *Ann Surg* 1985;201:456–64.
- Levey AS, Stevens LA, Schmid CH, *et al.* A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;150:604–12.
- Khwaja A. KDIGO clinical practice guidelines for acute kidney injury. *Nephron Clin Pract* 2012;120:c179–84.
- McCullough PA, Shaw AD, Haase M, *et al.* Diagnosis of acute kidney injury using functional and injury biomarkers: Workgroup statements from the tenth acute dialysis quality initiative consensus conference. *Contrib Nephrol* 2013;182:13–29.
- Endre ZH, Walker RJ, Pickering JW, *et al.* Early intervention with erythropoietin does not affect the outcome of acute kidney injury (the EARLYARF trial). *Kidney Int* 2010;77:1020–30.
- Basu RK, Wang Y, Wong HR, *et al.* Incorporation of biomarkers with the renal anemia index for prediction of severe AKI in critically ill children. *Clin J Am Soc Nephrol* 2014;9:654–62.
- Basu RK, Wong HR, Krawczeski CD, *et al.* Combining functional and tubular damage biomarkers improves diagnostic precision for acute kidney injury after cardiac surgery. *J Am Coll Cardiol* 2014;64:2753–62.

- 42 Alderman MH, Melcher L, Drayer DE, *et al.* Increased excretion of urinary N-acetyl-beta-glucosaminidase in essential hypertension and its decline with antihypertensive therapy. *N Engl J Med* 1983;309:1213–7.
- 43 Kohno M, Kanayama Y, Yasunari K, *et al.* Significance of the measurement of urinary alanine aminopeptidase and N-acetyl-beta-D-glucosaminidase activity in evaluating patients with essential hypertension. *Clin Exp Hypertens A* 1985;7:1347–60.
- 44 Ouchi M, Suzuki T, Hashimoto M, *et al.* Urinary N-acetyl-β-D-glucosaminidase levels are positively correlated with 2-hr plasma glucose levels during oral glucose tolerance testing in prediabetes. *J Clin Lab Anal* 2012;26:473–80.
- 45 Agirbasli M, Radhakrishnamurthy B, Jiang X, *et al.* Urinary N-acetyl-beta-D-glucosaminidase changes in relation to age, sex, race, and diastolic and systolic blood pressure in a young adult biracial population. The Bogalusa heart study. *Am J Hypertens* 1996;9:157–61.
- 46 Guillou H, David V, Lorcy Y, *et al.* Serum lysosomal acid hydrolase activities in Graves' disease. *Clin Chim Acta* 1982;120:219–24.
- 47 Wład H, Suszka B, Lacka K, *et al.* Enzymes depolymerizing glycosaminoglycans in blood serum of patients with HyPer and hypothyroidism. *Endokrynol Pol* 1985;36:215–20.
- 48 Ullrich K, Gieselmann V, Mersmann G, *et al.* Endocytosis of lysosomal enzymes by non-parenchymal rat liver cells. Comparative study of lysosomal-enzyme uptake by hepatocytes and non-parenchymal liver cells. *Biochem J* 1979;182:329–35.
- 49 Mano T, Sinohara R, Sawai Y, *et al.* Effects of thyroid hormone on coenzyme Q and other free radical scavengers in rat heart muscle. *J Endocrinol* 1995;145:131–6.
- 50 Pehowich DJ. Thyroid hormone status and membrane n-3 fatty acid content influence mitochondrial proton leak. *Biochim Biophys Acta* 1999;1411:192–200.
- 51 Venditti P, De Rosa R, Di Meo S. Effect of thyroid state on susceptibility to oxidants and swelling of mitochondria from rat tissues. *Free Radic Biol Med* 2003;35:485–94.
- 52 Gilbert RE, Wu LL, Kelly DJ, *et al.* Pathological expression of renin and angiotensin II in the renal tubule after subtotal nephrectomy. Implications for the pathogenesis of tubulointerstitial fibrosis. *Am J Pathol* 1999;155:429–40.
- 53 D'Amico G, Bazzi C. Pathophysiology of proteinuria. *Kidney Int* 2003;63:809–25.
- 54 Tryggvason K, Pettersson E. Causes and consequences of proteinuria: the kidney filtration barrier and progressive renal failure. *J Intern Med* 2003;254:216–24.
- 55 Klein I, Ojamaa K. Thyroid hormone and the cardiovascular system. *N Engl J Med* 2001;344:501–9.
- 56 Surks MI, Oppenheimer JH. Concentration of L-thyroxine and L-triiodothyronine specifically bound to nuclear receptors in rat liver and kidney. Quantitative evidence favoring a major role of T3 in thyroid hormone action. *J Clin Invest* 1977;60:555–62.
- 57 Iglesias P, Olea T, Vega-Cabrera C, *et al.* Thyroid function tests in acute kidney injury. *J Nephrol* 2013;26:164–72.
- 58 Plikat K, Langgartner J, Buettner R, *et al.* Frequency and outcome of patients with nonthyroidal illness syndrome in a medical intensive care unit. *Metabolism* 2007;56:239–44.