



# BMJ Open Quality of refractive error care (Q.REC) in Cambodia, Malaysia and Pakistan: protocol for a cross-sectional unannounced standardised patient study

Anthea M Burnett <sup>1,2</sup>, Ling Lee,<sup>1,2</sup> Myra McGuinness <sup>3,4</sup>, Beatrice Varga,<sup>1</sup> Yadira Perez Hazel,<sup>1</sup> Suit May Ho<sup>1</sup>

**To cite:** Burnett AM, Lee L, McGuinness M, *et al*. Quality of refractive error care (Q.REC) in Cambodia, Malaysia and Pakistan: protocol for a cross-sectional unannounced standardised patient study. *BMJ Open* 2022;**12**:e057594. doi:10.1136/bmjopen-2021-057594

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-057594>).

Received 21 September 2021  
Accepted 03 February 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>The Fred Hollows Foundation, Melbourne, Victoria, Australia

<sup>2</sup>School of Optometry and Vision Science, University of New South Wales, Sydney, New South Wales, Australia

<sup>3</sup>Melbourne School of Population and Global Health, University of Melbourne, Melbourne, Victoria, Australia

<sup>4</sup>Biostatistics, Centre for Eye Research Australia, Melbourne, Victoria, Australia

**Correspondence to**  
Dr Anthea M Burnett;  
[aburnett@hollows.org](mailto:aburnett@hollows.org)

## ABSTRACT

**Introduction** There are 161 million people living with vision impairment, due to uncorrected refractive errors. A further 510 million people are living with near-vision impairment. There is a need for clearly defined indicators that capture the quality of refractive error service outputs and outcomes and provide insights to shape, change and stimulate action. This study aims to evaluate the quality of refractive error care (Q.REC) in Cambodia, Malaysia and Pakistan, by using unannounced standardised patients (USPs) to identify the proportion of prescribed and dispensed spectacles appropriate for people's refractive error needs and pinpoint/detail opportunities for quality improvement.

**Method and analysis** A cross-sectional Q.REC study will be conducted in randomly selected optical services in Cambodia (180 services, 900 USP visits), the Klang Valley in Malaysia (66 services, 198 USP visits) and in Jhang, Sahiwal and Khanewal districts of Punjab region/state in Pakistan (64 services, 256 USP visits). USPs will receive baseline refractions by three skilled study optometrists/refractionists trained in the Q.REC protocol. USPs will then visit individual optical services, undergo a refraction, purchase spectacles or lenses (if recommended) and record observations about which elements of a refraction and dispensing were conducted. The study optometrist/refractionist will assess each pair of dispensed spectacles by examining the USP's aided visual acuity and visual comfort at distance and/or near and compare the lens prescription to the averaged baseline refraction.

**Ethics and dissemination** This study has been approved by the University of New South Wales Human Research Ethics Committee (HC210102), the National Ethics Committee for Health Research in Cambodia (043 NECHR), National Medical Research Registry and the Medical Research and Ethics Committee (NMRR-21-689-59279) in Malaysia and the College of Ophthalmology & Allied Vision Sciences Ethical Review Board (COAVS 545/2021) in Pakistan. Written informed consent will be obtained from USPs. Service owners will have the opportunity to opt-out verbally or in writing. Results will be disseminated locally through workshops including the relevant local ministry of health personnel and stakeholders, published in

## Strengths and limitations of this study

- This multisite study evaluates the 'real-world' quality of refractive error care and identifies specific opportunities for quality improvement.
- The 'gold standard' for evaluating clinical practice quality will be used, employing unannounced standardised patients (USPs) with a range of refractive error profiles.
- As all USPs will be adults, the findings may not be applicable to the quality of children's refractive error care.
- As only an immediate assessment of comfort with spectacles is included, any subsequent positive or negative adaptation cannot be assessed.
- Although USPs are trained to present their eye health history in a clear way, USPs may inadvertently mislead optical service staff and give an unclear version of their symptoms.

peer-reviewed publications and presented at national and international conferences.

## INTRODUCTION

Uncorrected refractive errors are the leading global cause of vision impairment, with 161 million people living with distance-vision impairment and an additional 510 million people living with near-vision impairment.<sup>1</sup> The WHO World Report on Vision states that integrated people-centred eye care, and a commitment to universal health coverage, is the model of choice for quality and affordable eye care services and quality eye care services need to be provided according to population needs.<sup>2</sup> The Lancet Global Commission on Global Eye Health argues that universal health coverage is not universal without affordable, high-quality, equitable eye care.<sup>3</sup> In 2021, the United Nations General Assembly adopted a

resolution committing the international community to provide eye health for the 1.1 billion people living with vision impairment by 2030.<sup>4</sup> To be able to monitor progress towards universal health coverage and the quality of eye care services, a broader set of measurement indicators are required to monitor progress and drive change. Indicators should provide insights to shape change and stimulate action, track outcome progress and the quality of an intervention.<sup>3</sup>

Indicators have recently been established,<sup>5</sup> to assess the ‘real-world’ quality of refractive care. These quality of refractive error care (Q.REC) criteria provide information on the proportion of prescribed spectacles that are clinically optimal for patients, by comparing a baseline prescription with dispensed spectacles. The Q.REC criteria were tested in Vietnam, in 93 optical stores and 480 individual visits, and demonstrated a strong association between the criteria for optimally prescribed spectacles and both vision and comfort.

This Q.REC indicator can also be used to assess core dimensions of quality, such as whether refractive error services are effective, equitable, safe and people centred, and have the potential to identify particular aspects of clinical practice that require improvement or further training.

A Q.REC study employs simulated patients—the gold standard for evaluating quality in clinical practice<sup>6</sup>—and can be easily administered in urban settings. Simulated patients, or unannounced standardised patients (USPs), are ‘actors’ who are trained to act covertly as patients in a standardised fashion, while observing clinical techniques and services provided. USPs have been used extensively in low-income and middle-income countries, often in evaluating family planning, pharmaceutical dispensing patterns and clinical prescribing patterns.<sup>7</sup> Studies employing USPs have also previously been conducted to evaluate refractive error outcomes.<sup>5 8 9</sup> If executed well, the distinct advantage of this method is that observation bias is minimised, as care providers are likely to modify their behaviours if they feel that they are being observed or examined.

A pilot Q.REC study in Vietnam in 2018 found that out of 417 pairs of spectacles purchased from 93 optical outlets, only 44% of spectacles resulted in both good vision and comfort for patients.<sup>5</sup> This highlights a significant gap in current models of care and represents an opportunity for improving quality outcomes within the existing infrastructure.

It is anticipated that a Q.REC study will identify the proportion of people who are prescribed and dispensed spectacles appropriate for their refractive error needs and identify specific opportunities for quality improvement which can then be translated into policy changes or quality improvement initiatives. Q.REC studies can also be used to monitor ongoing delivery of quality refractive error care services, within the context of integrated people-centred eye care.

The overall objective of this study is to evaluate the Q.REC in Cambodia, Malaysia and Pakistan. The

primary outcome of interest is the proportion of spectacles dispensed from optical service providers that meet ‘optimal’ quality standards, as defined by Lee *et al.*<sup>5</sup> Secondary outcomes of interest are the association between ‘optimal’ quality spectacles and refraction/dispensing techniques, subjective visual acuity and USP’s comfort while wearing dispensed spectacles.

## METHODS AND ANALYSIS

### Study setting

In each location we have drawn a representative sample of eligible optical services using simple random sampling. The sampling frame is comprised of all optical services within provinces with more than five optical services in Cambodia, all optical services within the Klang Valley in Malaysia and all optical services within the Jhang, Sahiwal and Khanewal districts of Punjab in Pakistan. It is anticipated that enrolment will commence in October 2021 and will be completed by June 2022.

### Study population

The inclusion criteria for USPs are adults aged 18 years or above, fluent native speaker of the primary language of the district, good ocular health and a refractive error profile of interest. This includes adults who need spectacles, as well as adults who have emmetropic eyes and do not need spectacles. We will exclude USPs who have had prior refractive eye surgery, eye surgery within the prior 3 months, USPs with manifest or intermittent strabismus, USPs with amblyopia and USPs who have any ocular or health conditions that can cause variable spectacle prescription. Each USP will order glasses from each service and may visit multiple services.

Eligible optical services are those that provide refractive error services in Cambodia, Malaysia (Klang Valley) and Pakistan (Jhang, Sahiwal and Khanewal districts of the Punjab province). Optical services that are part of a franchise or chain will be considered a different service. Optical services will be excluded from the study if the optical service is personally known to the USP or if it is identified that the optical service has staff working across multiple services that have already been selected. The participant information statement and withdrawal form will be hand delivered or sent via registered post, in the local language, at least 1 month in advance advising that the services might be visited for research purposes. During this time, the optical service owner/manager will be asked to read the form, consider whether they would like to opt-out of the study and identify whether study feedback is desired. If the randomly selected services do not return the form for withdrawal of participation by the due date and the first USP attending says they are unlikely to be detected, the services are considered successfully enrolled. As the public (potential optical service clients/patients) have a right to understand the quality of the services that they might expect to receive in each location, an opt-out approach will be used to ensure

**Table 1** Parameters for sample size estimates

	Total number eligible services	Anticipated percentage optimal	Spectacles from each service	Total number required	
				Spectacles	Services
Cambodia	265	50	5	900*	180
Malaysia	319	70	3	198	66
Pakistan	90	50	4	256	64

Intraclass correlation=0.1, design effect=1.2.

\*The number of spectacles in Cambodia has been increased to provide an update on the precision of the intraclass coefficient.

that there is a high participation rate. Also, the research is likely to be compromised if optical stores are aware that they are providing optical services to a USP due to the Hawthorne effect (where clinicians modify their behaviour in response to being observed). The quality of each individual store will not be published in any way so the privacy of each store will be maintained.

### Sampling strategy

Lists of all eligible services were compiled by consulting with relevant ministries of health, optical councils, optometry associations and cross-referencing with Google Maps. Services will be selected by computer-generated simple random sampling by study coordinators after the due date to withdraw from participation.

### Sample size

The anticipated proportion of spectacles of optimal quality ( $p$ ) was chosen based on previous studies and local knowledge (see [table 1](#)).<sup>5-9</sup> Malaysia has a more developed and regulated optometry industry than Cambodia or Pakistan, so it was anticipated that there the proportion of spectacles that are of optimal quality was higher. The margin of error for the 95% CI was chosen to provide sufficient precision with the available resources, as each study was funded separately. For Malaysia and Pakistan, the desired margin of error was set at 7%. In Cambodia, margin of error was set at 4% as the study budget provided the opportunity to maximise the sample size, so as to reduce the margin of error and make additional learnings on the precision of the intraclass coefficient. The intraclass correlation was estimated to be 0.1, and the design effect was estimated to be 1.2. Based on the parameters presented in [table 1](#), a total of 1354 spectacles will be purchased.

It is anticipated that up to 40 USPs will be employed and each USP will be able to visit three to four services per day.

### Development of USPs

#### Recruitment of USPs

To recruit USPs, study flyers have been advertised in ophthalmology outpatient clinics and at adult education institutions in local languages. Participating USPs will be reimbursed for participation, plus any travel, accommodation, meal or spectacle purchase expenses.

### Training

All USPs will be briefed/trained in person on the tasks involved by a study coordinator and study optometrist, and will be allowed sufficient time to practice acting as a patient. Training packages have been developed to provide USPs with a 'script' that they can use during optical service visits. Post-training observations by a study optometrist will be conducted to identify whether the USPs can accurately identify elements of refraction and dispensing techniques. If USPs are unable to accurately identify elements of refraction and dispensing techniques, further training will be provided.

### Baseline refraction

Baseline refractions of each USP will be conducted by three skilled study optometrists/refractionists who will be recruited within each country to conduct individual clinical refractions, in order to establish the USP's average baseline prescription and their best-corrected visual acuity. All USPs will undergo three baseline refractions by the study optometrists/refractionists. For USPs with significant refractive error where a modified prescription might be required for adaptation, the study optometrists/refractionists will agree on level of modification that is to be considered for the baseline refraction, otherwise full prescription will be considered for baseline. If the refraction components (spherical and cylindrical powers) are not within 0.75 D of each other, a fourth refraction is required. The most senior optometrist will then decide which three refraction results will be used for the averaged baseline refraction for that USP.

### Procedures

#### USP optical service visit

To assess the quality of the spectacles prescribed and dispensed, USPs will visit individual optical services, undergo a refraction, purchase spectacles (if recommended) and record observations about which elements of a refraction and dispensing were conducted (online supplemental file 1). Although a schedule will be developed for each USP to attend selected optical services, no prior appointment will be made unless it is standard practice for the optical service. Once the eye examination is complete, the USP will place an order for new spectacles/lenses. After the visit, the USP will find a suitable location

**Table 2** Criteria for optimally prescribed spectacles

Spectacle component	Tolerance limits*
Spherical power (most plus power)	±0.50 D
Cylindrical power	±0.50 D
Cylindrical axis (if baseline cylinder power ≤ -0.50 DC)	±7°
Cylindrical axis (if baseline > -0.50 DC to ≤ -1.50 DC)	±5°
Cylindrical axis (if baseline > -1.50 DC)	±2°
Horizontal prism	<1 prism dioptre (in/out direction)
Vertical prism	<0.50 prism dioptre (up/down direction)

\*Development and testing of the tolerance limits are described in Lee *et al.*<sup>5</sup>

to complete the optical services visit checklist electronically. The USP will return to each service to collect the purchased spectacles/lenses and obtain a copy of the spectacle prescription.

### Spectacle quality

Each pair of spectacles dispensed will be assessed by the study optometrist/refractionist to determine whether the prescribed spectacles pass or fail the criteria for optimally prescribed spectacles (table 2) and assess the visual acuity and vision comfort at distance and near.

Evidence-based spectacle quality criteria were developed from published mean subjective intolerance to spectacles for lens power and induced prism<sup>10 11</sup> and published dispensing standards for tolerance to cylindrical axis<sup>12</sup> (table 2). Achieving all criteria in both eyes with a pair of spectacles is defined as 'optimally prescribed spectacles'.

### Analysis

#### Spectacle characteristics

The distribution of spectacle characteristics will be compared between and across regions using Pearson's  $\chi^2$  test. Spectacle characteristic will include lens type (single vision near, single vision distance) and refractive error type (binary variable for each of presbyopia, astigmatism, myopia and hypermetropia).

#### Primary outcome

The unit of analysis will be one pair of spectacles. The proportion of spectacles that are 'optimal' within each country and district will be estimated using sampling weights, and logit-transformed 95% CIs will be estimated using robust standard errors to account for intraservice correlation.

#### Secondary outcomes

Service characteristics (refraction techniques and refraction equipment used), USP comfort and USP-corrected

visual acuity while wearing dispensed spectacles (dichotomised as achieving best-corrected binocular visual acuity vs not achieving (worse than 1.5 lines on a logMAR chart)) will be compared between spectacles of optimal and suboptimal quality via logistic regression, adjusting for sampling weights and intraservice correlation.

Analyses will be conducted using Stata/BE V.17.0 (StataCorp LLC, College Station, Texas) or equivalent.

### Database management

Study data will be collected, managed and securely stored using Research Electronic Data Capture hosted at The Fred Hollows Foundation.<sup>13 14</sup> Study data will be entered into the database by the USPs and study optometrists/refractionists. Data quality will be assured by conducting daily queries to identify and resolve discrepancies and by using data quality rules.

### Patient and public involvement

There was no patient or public involvement in the design of this protocol, although all coauthors have lived experience with refractive errors.

All participating USPs and service owners will have the opportunity to receive a summary of the results in their preferred language.

### ETHICS AND DISSEMINATION

As specified above, enrolment of services in this study will be on an opt-out basis, with written information on the nature of the study being provided to the service owners. Services owners will be offered the opportunity to decline participation, verbally or in writing. Written informed consent will also be obtained from the USPs after the nature of the study and potential harms have been explained. Ethics approvals have been obtained from the following ethics committees:

- ▶ Cambodia: National Ethics Committee for Health Research, Cambodia (043 NECHR).
- ▶ Malaysia: National Medical Research Registry and the Medical Research and Ethics Committee (NMRR-21-689-59279).
- ▶ Pakistan: College of Ophthalmology & Allied Vision Sciences Ethical Review Board (COAVS 545/2021).
- ▶ Australia: University of New South Wales Human Research Ethics Committee (HC210102).

Results will be disseminated to local government partners through workshops to discuss potential study and policy implications. In addition, the results will be published in peer-reviewed publications and presented at national and international conferences.

### DISCUSSION

The Q.REC study aims to provide a much-needed methodology for assessing the Q.REC, as experienced by patients. This study will identify the proportion of spectacles obtained from optical services that provide quality vision and comfort outcomes and the associations

between spectacle quality and refraction and dispensing techniques. The results of this study will provide valuable information on the Q.REC and identify specific opportunities for quality improvement in Cambodia, Malaysia and Pakistan. This information will provide much-needed empirical evidence to aid policy-makers, education institutions and service providers on how to ensure that refractive error services provide quality care. The findings can be translated into policy changes or quality improvement initiatives and additional Q.REC studies can be used to monitor progress towards delivering quality refractive error care.

The Lancet Global Health Commission on Global Eye Health stated that concerted efforts are needed to improve quality eye health outcomes and reliable survey and service data should be available and used by implementers and policy-makers to monitor progress.<sup>3</sup> A Q.REC study provides clearly defined indicators that capture refractive error service outputs and outcomes, along with ‘insights to shape change and stimulate action’.<sup>3</sup> In addition, a Q.REC study includes broader dimensions of refractive error quality. Using visual acuity alone as the indicator of refractive error quality does not take into account components of spectacle quality that can exacerbate binocular vision disorders or contribute to visual discomfort. Nor does visual acuity alone take into account timeliness, visual comfort, effectiveness, safety (eg, spectacle prescriptions that result in unnecessarily reduced vision) or any assessment of the extent to which services are people centred. These additional aspects of quality care are a core component of advancing universal health coverage through eye care.

There are several limitations to this study that should be acknowledged. First, as all USPs will be adults, the findings may not be applicable to the quality of children’s refractive error care. Second, USPs may inadvertently mislead optical service staff and give an unclear version of their symptoms. To reduce this potential bias training will be provided to ensure USPs are confident at presenting symptoms and history in a clear manner. Third, it could be argued that by only including an immediate assessment of comfort, the possibility of any positive or negative adaptation that might occur with extended wear is ignored. However, this methodology is not designed to assess spectacle compliance, which is likely to be heavily impacted by adaptation that occurs during extended wear. Although prior studies have indicated that spectacle wear compliance rates can be as low as 40%,<sup>15</sup> which infers that as many as 60% of dispensed spectacles remain unworn, this study may provide evidence on the proportion of spectacles that are likely to be unsuitable for population needs. These insights can be used to stimulate action and provide refractive error services according to population needs—resulting in higher rates of spectacle wear.

Currently, the Q.REC indicators are focused on whether they are people centred, effective, equitable and safe. Additional questions and components could be added to assess other dimensions of quality, such as whether they

are timely, integrated and efficient. Future trials could also include more aspects of dispensing techniques, such as frame fit and comfort.

This study will produce a set of validated tools for assessing the quality of spectacles obtained from optical services, resulting in solution-focused contextually relevant research that will encourage high quality and universal eye health for all.

**Contributors** AMB and LL led the design of the study and developed the protocol. AMB applied for ethical approvals and drafted the manuscript. MM contributed to design of statistical analysis methods, sample size calculations and drafting the protocol. BV, YPH and SMH provided advice on key study issues and feedback on the protocol. All authors read, revised and approved the final manuscript.

**Funding** This work was funded by The Fred Hollows Foundation with support from the Australia Government through the Australian NGO Cooperation Program.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Anthea M Burnett <http://orcid.org/0000-0003-3772-6625>

Myra McGuinness <http://orcid.org/0000-0002-5422-040X>

#### REFERENCES

- Adelson J, Bourne RRA, Briant PS. Causes of blindness and vision impairment in 2020 and trends over 30 years : evaluating the prevalence of avoidable blindness in relation to VISION 2020: the Right to Sight. *Lancet Global Health* 2020.
- World Health Organization (WHO). *World report on vision*. Geneva: WHO, 2019.
- Burton MJ, Ramke J, Marques AP, *et al*. The Lancet global health Commission on global eye health: vision beyond 2020. *The Lancet Global Health* 2021;9:e489–551.
- United Nations General Assembly. Vision for everyone: accelerating action to achieve the sustainable development goals, 2021. Available: <https://undocs.org/en/A/RES/75/310> [Accessed 3 Sep 2021].
- Lee L, Burnett AM, D’Esposito F, *et al*. Indicators for assessing the quality of refractive error care. *Optom Vis Sci* 2021;98:24–31.
- Rethans J-J, Gorter S, Bokken L, *et al*. Unannounced standardised patients in real practice: a systematic literature review. *Med Educ* 2007;41:537–49.
- Wiseman V, Lagarde M, Kovacs R, *et al*. Using unannounced standardised patients to obtain data on quality of care in low-income and middle-income countries: key challenges and opportunities. *BMJ Glob Health* 2019;4:1–3.
- Shah R, Ctori I, Edgar DF, *et al*. Use of standardised patients in optometry training. *Clin Exp Optom* 2021;104:848–53.
- Nie J, Zhang L, Gao J, *et al*. Using incognito standardised patients to evaluate quality of eye care in China. *Br J Ophthalmol* 2021;105:bjophthalmol-2019-315103.
- du Toit R, Ramke J, Brian G. Tolerance to prism induced by readymade spectacles: setting and using a standard. *Optom Vis Sci* 2007;84:1053–9.



- 11 Freeman CE, Evans BJW. Investigation of the causes of non-tolerance to optometric prescriptions for spectacles. *Ophthalmic Physiol Opt* 2010;30:1–11.
- 12 Brown WL. Revisions to tolerances in cylinder axis and in progressive addition lens power in ANSI Z80.1-2005. *Optometry* 2006;77:343–9.
- 13 Harris PA, Taylor R, Thielke R, *et al*. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.
- 14 Harris PA, Taylor R, Minor BL, *et al*. The REDCap Consortium: building an international community of software platform partners. *J Biomed Inform* 2019;95:103208.
- 15 Dhirar N, Dudeja S, Duggal M, *et al*. Compliance to spectacle use in children with refractive errors- a systematic review and meta-analysis. *BMC Ophthalmol* 2020;20:71.



**The Fred Hollows  
Foundation**



**UNSW  
AUSTRALIA**

## Q.REC Study: Record Forms

### Form A: Baseline USP Data Record Form

This form is to be filled out by the research optometrist / refractionist.

Optometrist or

Refractionist name: .....

USP ID: .....

USP Gender: .....

USP Age: .....

	Right lens	Left lens
Sph		
Cyl		
Axis		
Monocular distance VA*		
Binocular distance VA*		
Near addition		
Monocular near VA*		
Binocular near VA*		
Pupillary distance in mm (dist/near)		

\* Visual acuity (VA) should be in 6/6 or 20/20 format including incomplete lines, e.g. 6/7.5<sup>+2</sup>

With the amount of refractive error, would you prescribe spectacles?

Yes

No

What types of spectacles would be suitable for this USP? (tick all that apply)

Distance

Near

Bifocal

Multifocal / Progressive



**The Fred Hollows  
Foundation**



**UNSW  
AUSTRALIA**

## Form B: Post-training Assessment

This form is to be filled out by the research optometrist / refractionist.

1. Conduct a refraction with the USP including a spectacle recommendation *Or* Attend an optical service with a USP to observe.
2. Observe the USP completing the 'Optical service visit checklist' immediately after the visit.

Checklist	Yes, recorded or observed	Additional training needs
Was the USP able to correctly identify and record if focimetry was or was not performed?		
Was the USP able to correctly identify and record if distance visual acuity was or was not measured?		
Was the USP able to correctly identify and record if pinhole visual acuity was or was not measured?		
Was the USP able to correctly identify and record if near visual acuity was or was not measured?		
Was the USP able to correctly identify and record if retinoscopy was or was not performed?		
Was the USP able to correctly identify and record if auto-refraction was or was not performed?		
Was the USP able to correctly identify and record what type of subjective refraction was or was not performed?		
Was the USP able to correctly identify and record if the <i>spherical component</i> of subjective refraction was or was not performed?		
Was the USP able to correctly identify and record if the <i>cylindrical component</i> of subjective refraction was or was not performed?		
Was the USP able to correctly identify and record if the subjective near refraction was or was not performed?		
Was the USP able to correctly identify and record if the pupillary distance was or was not measured?		
Did the USP request a written prescription?		
Did the USP assess the visual acuity chart distance?		
Did the USP stick to the script?		





**The Fred Hollows  
Foundation**



**UNSW  
AUSTRALIA**

## Form C: USP Optical service visit checklist

This form is to be filled out by the USP immediately after each optical service visit.

USP ID: \_\_\_\_\_

Optical Service ID: \_\_\_\_\_

Q No.	Question	Response (Code)	Go to
1	Do you feel the optical service has identified you as an unannounced standardised patient (USP)?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	If yes: Q2 If no or unsure: Q4
2	When do you think you were identified?	<input type="checkbox"/> Before the eye test (1) <input type="checkbox"/> During the eye test (2) <input type="checkbox"/> At the end of the eye test (3) <input type="checkbox"/> When I went to pick up the glasses (4)	Q3
3	Why do you think you were or might have been identified as a USP?	<i>[Open comment]</i>	Q4
4	Did you bring your glasses with you?	<input type="checkbox"/> Yes, I wore them (1) <input type="checkbox"/> Yes, I had them with me (2) <input type="checkbox"/> No (0)	If yes: Q5 If no: Q6
5	Did any of the staff perform focimetry on your glasses?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q6
6	Did the clinician check distance visual acuity before refraction?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q7
7	Did the clinician check distance visual acuity with a pinhole?	<input type="checkbox"/> Yes, (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q8



**The Fred Hollows  
Foundation**



Q No.	Question	Response (Code)	Go to
8	Did the clinician check near visual acuity at the beginning of the eye test?	<input type="checkbox"/> Yes(1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q9
9	Did the clinician perform auto-refraction?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q10
10	Did the clinician perform retinoscopy?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q11
11	Did the clinician perform a distance vision refraction?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q12 If no: Q17
12	Did the clinician use a phoropter during the refraction?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q13
13	Did the clinician use a trial frame during the refraction?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q14
14	Did the clinician test the <b>spherical element</b> during the refraction?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q15
15	Did the clinician test the <b>cylindrical element</b> during the refraction?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q16
16	Did the clinician check your near visual acuity with the distance lenses?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q17
17	Did the clinician perform a near vision refraction?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q18



**The Fred Hollows  
Foundation**



Q No.	Question	Response (Code)	Go to
18	Did the clinician recommend new glasses?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	If yes: Q19 If no: Q22
19	Why did the clinician say you need new glasses?	<input type="checkbox"/> My prescription has changed a lot (1) <input type="checkbox"/> My current glasses are poor quality (2) <input type="checkbox"/> No reason provided (3) <input type="checkbox"/> Other. Please describe (4) _____ <input type="checkbox"/> Unsure (88)	Q20
20	What type of glasses were you recommended? (select all that apply)	<input type="checkbox"/> Distance only (1) <input type="checkbox"/> Near only (2) <input type="checkbox"/> Both distance and near as separate glasses (3) <input type="checkbox"/> Bifocals (4) <input type="checkbox"/> Multifocals (5) <input type="checkbox"/> Unsure (88)	Q21
21	Did any of the staff measure pupil distance?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q23
22	Why did the clinician say you DO NOT need new glasses?	<input type="checkbox"/> My prescription has not changed, my current glasses are still good (1) <input type="checkbox"/> I do not need to wear glasses (2) <input type="checkbox"/> No reason was provided. (3) <input type="checkbox"/> Unsure (88)	Q23
23	Did the clinician recommend a referral to secondary/tertiary eye care services?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q24
24	Did the clinician recommend a follow up eye examination?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q25
25	Did the clinician and staff members communicate to you clearly:		



**The Fred Hollows  
Foundation**



Q No.	Question	Response (Code)	Go to
25a	-During the eye test	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q25b
25b	-With the outcomes of the eye test	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q25c
25c	-When explaining which glasses (lenses or frames) you need OR why you don't need new glasses	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q26
26	Gender of clinician	<input type="checkbox"/> Female (1) <input type="checkbox"/> Male (2) <input type="checkbox"/> Unsure (88)	Q27
27	Were there any qualifications or registration certificate(s) observed in the store?	<input type="checkbox"/> Yes (1) <input type="checkbox"/> No (0) <input type="checkbox"/> Unsure (88)	Q28
28	What type of distance visual acuity chart was used?	<input type="checkbox"/> Paper chart directly in front of me (1) <input type="checkbox"/> Paper chart placed behind me but could be seen in the mirror in front of me (2) <input type="checkbox"/> Internally lit chart in front of me (3) <input type="checkbox"/> Internally lit chart placed behind me but could be seen in the mirror in front of me (4) <input type="checkbox"/> Computer chart in front of me (5) <input type="checkbox"/> Computer chart placed behind me but could be seen in the mirror in front of me (6) <input type="checkbox"/> Other. Please describe: (7) _____ <input type="checkbox"/> Unsure (88)	Q29



**The Fred Hollows  
Foundation**



**UNSW**  
AUSTRALIA

Q No.	Question	Response (Code)	Go to
29	What was the approximate distance between you and the distance visual acuity chart?	_____ (in metres)	Complete



**The Fred Hollows  
Foundation**



**UNSW  
AUSTRALIA**

## Form D: Assessment of Spectacles/Glasses

This form is to be filled out by the research optometrist / refractionist.

USP ID: \_\_\_\_\_ Optical Service ID: \_\_\_\_\_

Type of spectacles:

- Distance  
 Near  
 Bifocal  
 Multifocal

Visual acuity (6/6 or 20/20 format including incomplete lines, e.g. 6/7.5<sup>+2</sup>)

	Right eye	Left eye	Both eyes
Distance			
Near			

Near viewing distance (in cm): : \_\_\_\_\_

With the spectacles, does the USP experience any eye strain or discomfort with:

Distance vision	<input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> NA (near spectacles only)
Near vision	<input type="checkbox"/> Yes <input type="checkbox"/> No

### Focimetry

	Right lens	Left lens
Sph		
Cyl		
Axis		
Near addition		
Vertical prism ( $\Delta$ and direction)		

	Distance prescription	Near prescription
Lens centration distance (mm)		



**The Fred Hollows  
Foundation**



**UNSW  
AUSTRALIA**

Written prescription (can be scanned and uploaded to database)

	Right lens	Left lens
Sph		
Cyl		
Axis		
Near addition		
Horizontal prism ( $\Delta$ and direction)		
Vertical prism ( $\Delta$ and direction)		

	Distance	Near
Pupil distance (mm)		

Additional notes (e.g. specifying lens type): \_\_\_\_\_

Does the written prescription have any information about the optical services' registration?

- Yes – a registration number
- Yes – other indication Please describe: \_\_\_\_\_
- No
- Unsure