

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Hypertension, its correlates, and differences in access to healthcare services by gender among rural Zambian residents: a cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-055668
Article Type:	Original research
Date Submitted by the Author:	20-Jul-2021
Complete List of Authors:	Tateyama, Yukiko; Kyoto University, Health Service; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research Techasrivichien, Teeranee ; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research; International Institute of Socio-epidemiology Musumari, Patou Masika ; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research; International Institute of Socio-epidemiology Suguimoto, S. Pilar; International Institute of Socio-epidemiology; National Institute of Health Ongosi, Anita; International Institute of Socio-epidemiology Zulu, Richard; Center for Primary Care Research, University of Zambia School of Medicine Dube, Christopher; Ndola District Health Office Ono-Kihara, Masako; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research; International Institute of Socio-epidemiology Kihara, Masahiro; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research; International Institute of Socio-epidemiology
Keywords:	Hypertension < CARDIOLOGY, PUBLIC HEALTH, International health services < HEALTH SERVICES ADMINISTRATION & MANAGEMENT

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Hypertension, its correlates, and differences in access to healthcare services by gender among rural Zambian residents: a cross-sectional study

Author information

Yukiko Tateyama,^{1,2,3*} Teeranee Techasrivichien,^{2,3} Patou Masika Musumari,^{2,3} S. Pilar Suguimoto,^{3,6} Anita Ongosi,³ Richard Zulu,⁴ Christopher Dube,⁵ Masako Ono-Kihara,^{2,3} Masahiro Kihara^{2,3}

Author affiliations

1. Kyoto University Health Service, Kyoto, Japan
2. Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research, Kyoto University, Kyoto, Japan
3. International Institute of Socio-epidemiology, Kyoto, Japan
4. Center for Primary Care Research, University of Zambia School of Medicine, Lusaka, Zambia
5. Ndola District Health Office, Ndola, Zambia
6. National Institute of Health, Lima, Peru

*Corresponding author

Postal address: Yoshida-honmachi, Sakyo-ku, Kyoto, Japan 606-8501

Email: tateyama.yukiko.57c@kyoto-u.jp

Word Count: 3999 words

32 ABSTRACT

33 **Objectives:** To examine the prevalence of hypertension and access to related healthcare
34 services among rural residents of Mumbwa district in Zambia.

35 **Design:** Cross-sectional study with probability cluster sampling.

36 **Setting:** Rural Zambia.

37 **Participants:** We recruited 690 residents from Mumbwa district aged 25–64 years who had
38 been living in the study area for ≥ 6 months and had adopted the lifestyle of the study area.
39 Pregnant women and women who had given birth in the last 6 months were excluded. The data
40 collection - questionnaire survey and anthropometric and biological measurements - was
41 conducted between May and July 2016.

42 **Results:** In the overall sample, 39.7% of the men and 33.5% of the women had hypertension
43 (SBP ≥ 140 or DBP ≥ 90 mmHg), respectively. Among the participants without a previous
44 diagnosis of hypertension, 30.3% presented with hypertension at the time of measurement. In
45 the multivariable analysis, alcohol intake and urban residence in men, and older age group,
46 higher education, and body mass index ≥ 25 in women were significantly associated with
47 hypertension. Among the 21.8% who never had their blood pressure (BP) measured, 83.8%
48 were men; in this group of men, older age (AOR, 0.43; 95% CI, 0.25–0.73) and HIV positive
49 status (AOR, 0.37; 95% CI, 0.14–0.97) were negatively associated while current smoker status
50 (AOR, 2.09; 95%CI, 1.19–3.66) was positively associated with the lack of BP measurements.

51 **Conclusion:** We found that hypertension is prevalent in the target rural area. However, many
52 were not aware of their hypertension status and many never had their BP measured, indicating
53 a serious gap in cardiovascular disease prevention services in Zambia. There is an urgent need
54 for health promotion and screening for hypertension, especially in the primary health services
55 of rural Zambia. Particular attention should be paid to issues related to healthcare accessibility
56 in men.

58 KEY WORDS

59 Hypertension, blood pressure, rural, cluster sampling, access, health promotion

61 Strengths and limitations of this study

- 62 • This study assessed the prevalence and factors associated with hypertension stratified by
63 gender to understand the current hypertension status among rural residents of Zambia.

- 64 • We employed multi-stage cluster random sampling and obtained a relatively high response
- 65 rate, which helped ensure that the results are representative of the target population.
- 66 • Socially desirable responses due to face-to-face interviews might have affected the results.

67

68 INTRODUCTION

69 Hypertension is a major global health concern; currently, there are 17.9 million cases of
70 mortality each year due to coronary heart disease and stroke worldwide.[1] The burden of
71 hypertension has globally increased during the past quarter century and accounts for 7% of
72 disability-adjusted life years.[2] It has been reported that if no action is taken to control
73 hypertension, economic losses will outstrip public healthcare spending.[3]

74 It is difficult to be aware of hypertension without assessment during the early stages
75 because it is asymptomatic.[3] To address the increasing prevalence of hypertension, early
76 detection and awareness of hypertension is important, particularly at the primary healthcare
77 level. However, many people with hypertension in sub-Saharan Africa (SSA) may remain
78 undiagnosed, untreated, or uncontrolled because of an inadequate healthcare system.[4] In fact,
79 a systematic review of SSA studies reported that only 22.5% of people with hypertension were
80 aware of their hypertension status.[2] Additionally, a South African study found that among
81 people with hypertension, 51% ever had their blood pressure (BP) measured, of which nearly
82 half had not been told that they had high BP.[5] This indicates a lack of necessary health
83 services for prevention and screening of hypertension, particularly in developing countries.

84 SSA has been reporting rising rates of hypertension, with the highest prevalence rate in the
85 world (46% of adults aged 25 and older),[6] and the prevalence remains high in Zambia as well
86 (19.0% in 2017).[7] A recent study in Zambia found that the prevalence of hypertension in
87 people aged over 25 in rural settings was 23.1%.[8] However, this information was collected
88 from the clinical visit records at primary healthcare facilities and did not include people without
89 access to health facilities.

90 The aforementioned evidence underscores the importance of strengthening the assessment
91 and treatment of hypertension. However, research on screening and diagnosis of hypertension
92 has been limited in Zambia.[8-10] Therefore, we aimed to find the prevalence of hypertension
93 including undiagnosed cases to understand the current status and access to healthcare service
94 for hypertension among rural residents in Zambia. We also examined the correlation of
95 demographic, behavioural, and biological factors with hypertension.

96

97 **METHODS**

98 **Design and settings**

99 This was a cross-sectional study conducted between May and July 2016 in Zambia. We selected
100 Mumbwa district in Central Province as our study area because it is a typical rural area
101 experiencing urbanization and economic growth while maintaining traditional culture. The
102 district is located 150 km west of the capital Lusaka city and is home to approximately 210,847
103 inhabitants – 15% in urban areas and 85% in rural areas.[11] The target population included
104 residents aged 25–64 years. Since the objective of this study was to investigate lifestyle-related
105 risk factors, we only included residents who had been living in the study area for ≥ 6 months
106 and had adopted the prevalent lifestyle of the study area. Pregnant women and women who had
107 given birth in the last 6 months were excluded because of potentially different dietary habits
108 and lifestyles and the fact that prepartum and postpartum weight could affect their
109 anthropometric and biological data.

111 **Sampling**

112 We employed a three-stage probability proportional to size (PPS) cluster sampling. The sample
113 size calculation was based on the recommendations of the WHO STEPwise approach to
114 surveillance (STEPS),[12] assuming 95% confidence level, 5% margin of error (e_2), and 30%
115 prevalence of hypertension in rural areas.[12] The minimum sample size required was 167
116 subjects, which was increased to 800 to address design effects (loss of sampling efficiency due
117 to cluster sampling), an assumed 20% non-response rate, and planned subgroup and
118 multivariate analyses.

119 The Central Statistical Office (CSO) of Zambia provided the list of study sampling clusters and
120 Standard Enumeration Areas (SEAs). In the first stage, we selected 32 SEAs through PPS
121 sampling without replacement using the sampling frame of the Zambia Population and Housing
122 Census 2010.[11] In the second stage, within each selected SEA, field staff consisting of
123 mappers from CSO and research assistants mapped the area and listed all households and their
124 eligible members. A total of 25 households in each SEA were selected through systematic
125 sampling. In the third stage, from each selected household, only one individual was selected
126 using the Kish Household Coversheet based on the WHO STEPS.[12] We scheduled a date
127 and place to administer the questionnaire survey and take anthropometric and biological
128 measurements as per the participants' convenience. We met with all recruited individuals (if
129 absent, their family members or closest neighbours) 1–2 days before testing to request them to

1
2 130 start fasting at 8:00 pm on the day prior to the biological measurements and to visit the testing
3 131 venue on the scheduled date.

4
5 132

6 7 133 **Data collection**

8
9 134 The questionnaire was developed in English and three local languages based on the review of
10 135 Zambian and international literature[9,13] and the results of an earlier qualitative study. [14]
11 136 A pilot study was conducted to resolve language discrepancies, to assess the face validity of
12 137 the questionnaire and test-retest reliability, and confirm the feasibility of anthropometric and
13 138 biological measurements. Face-to-face interviews were carried out by field staff at venues such
14 139 as the participant's home, community meeting places, or schools. Additionally, licensed nurses
15 140 were recruited and trained to collect anthropometric measurements and biological samples.

16
17 141

18 19 142 **Measurements**

20 143 BP was measured using electronic equipment (Omron HEM-7130-HP, Omron Corporation,
21 144 Kyoto, Japan). Three measurements were taken from the participants at three-minute intervals
22 145 while they were seated after 15 minutes of rest, and the average of the last two readings was
23 146 recorded. Weight was measured while the participants were barefoot and wearing light clothing
24 147 using an electronic scale (Omron HBF-223-G, Omron Corporation, Kyoto, Japan). Glycated
25 148 haemoglobin (HbA1c) and blood lipids (total cholesterol, LDL-cholesterol, HDL-cholesterol,
26 149 and triglycerides) were measured using point-of-care testing device (Cobas b 101, Roche
27 150 Diagnostics K.K., Tokyo, Japan). Other variables used in the analysis included socio-
28 151 demographic characteristics, food security using the Household Food Insecurity Access
29 152 Scale,[15] medical history and current medications, psychological distress using the Kessler-6
30 153 scale,[16] and lifestyle-related variables (tobacco, alcohol, physical activity, and dietary habits).
31 154 The results of the measurements were explained by local nurses and given to each participant.
32 155 Those who had extremely abnormal results were encouraged to visit the nearest health facilities
33 156 with the reports.

34
35 157

36 37 158 **Patient and public involvement**

38 159 Participants were not involved in the design, conduct, reporting, and dissemination plans of our
39 160 research.

40
41 161

42 43 162 **Statistical analysis**

1
2 163 We analysed the data using the Complex Sample module in IBM SPSS Statistics version 21
3 164 (IBM Corp., Armonk, NY, USA) to adjust for the effects of multistage sampling, clustering,
4 165 and weighting. Sample weights accounted for different selection probabilities at each sampling
5 166 stage, non-response rate in each SEA, and post-stratification adjustments to correct for
6 167 differences between our sample and the district population estimates based on the 2010 census.
7 168 Total weights were standardised as the final weight. Bivariate analyses were performed to
8 169 determine statistically significant associations between independent variables and high BP
9 170 (systolic blood pressure (SBP) ≥ 140 or diastolic blood pressure (DBP) ≥ 90 mmHg) using
10 171 logistic regression. Variables that showed significant associations with high BP ($p < 0.10$) in
11 172 the bivariate analysis were entered into the multiple logistic regression models stratified by
12 173 gender.
13 174

175 **Ethical considerations**

176 This study was approved by the Ethics Committee of the Graduate School and Faculty of
177 Medicine of Kyoto University, Japan (R0403) and ERES Converge, Zambia (No. 2016-Jan-
178 003) for the pilot phase. The University of Zambia Biomedical Research Ethics Committee,
179 Zambia (No. 011-02-16) and the National Health Research Authority, Zambia (MH/101/23/10-
180 1) granted approval for the main survey. All participants provided written informed consent
181 prior to their participation.
182

183 **RESULTS**

184 Of the 800 targeted subjects, 712 agreed to participate. We excluded 22 participants from the
185 analyses due to missing interviews or anthropometric/biological data. The final valid response
186 rate was 86.3%. Table 1 shows the weighted characteristics of the study population by gender.
187 The sample consisted of 48.6% of men, and the mean age was 41.9 years (SE 0.6). Most of the
188 participants were married (%), had only primary education (%), and were self-employed (%).
189 Nearly one-half had an income of 50 USD or less (Zambia's minimum wage); one-quarter were
190 living with severe food insecurity. For the self-reported medical history, 10.4% had with human
191 immunodeficiency virus (HIV) and were receiving antiretroviral therapy (ART). Only 8% and
192 0.7% of participants had been diagnosed with hypertension and diabetes, respectively. More
193 than 50% of both men and women had family members or relatives who had hypertension, and
194 about 20% reported having family members or relatives who had experienced a stroke.
195

196

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Sociodemographic characteristics and related medical histories among overall participants in the Mumbwa district, Central Province of Zambia, 2016

	Overall n (%)	Male n (%)	Female n (%)
Number			
Unweighted	690	332	358
Weighted	689 (100)	335 (48.6)	354 (51.4)
Age, years [SE]	41.9 [0.6]	42.7 [0.8]	41.1 [0.7]
Residential area of the district			
Urban area	87 (10.4)	35 (14.7)	52 (12.6)
Rural area	602 (89.6)	300 (85.3)	302 (87.4)
Marital Status			
Not married	27 (3.9)	20 (5.8)	7 (2.0)
Married	557 (80.8)	300 (89.4)	257 (72.6)
Divorced/widow/widower	106 (15.4)	16 (4.8)	90 (25.4)
Education			
Primary	513 (74.3)	229 (68.3)	284 (80.1)
Secondary	127 (18.5)	77 (23.0)	50 (14.2)
Tertiary	49 (7.2)	29 (8.7)	20 (5.7)
Monthly income (USD)			
≤50	326 (47.4)	157 (46.9)	169 (47.8)
>50	362 (52.6)	178 (53.1)	185 (52.2)
Work Status			
Employed	85 (12.4)	58 (17.2)	28 (7.8)
Self-employed	481 (69.7)	255 (76.2)	225 (63.6)
Unemployed/Retired	123 (17.9)	22 (6.6)	101 (28.6)
Food security			
Secure	192 (27.9)	107 (32.1)	85 (23.9)
Mildly insecure	45 (6.6)	25 (7.4)	21 (5.8)
Moderately insecure	261 (37.9)	132 (39.4)	129 (36.4)
Severely insecure	191 (27.7)	71 (21.2)	120 (33.8)
Medical history (Self-reported)			
HIV positive*	71 (10.4)	28 (8.4)	43 (12.2)
Hypertension	55 (8.0)	18 (5.4)	37 (10.4)
Diabetes	5 (0.7)	3 (0.9)	2 (0.6)
Past history within family and relatives (Self-reported)			
Hypertension	381 (55.3)	174 (52.0)	207 (58.5)
Stroke	140 (20.3)	68 (20.4)	71 (20.2)
Heart disease	64 (9.3)	25 (7.5)	39 (11.0)
Diabetes	123 (17.9)	52 (15.5)	71 (20.2)

Data are presented as numbers (%)

SE, standard error

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

*All have been receiving antiretroviral treatment.

Table 2 shows the prevalence of hypertension in each stage and the current status of access to health services for hypertension stratified by gender. The prevalence of hypertension (Stage 2 and hypertensive crisis) was 36.6% in the overall sample and was greater in men than in women but without statistical significance (39.7% vs. 33.5%, $p = 0.10$). In contrast, the prevalence of hypertensive crisis, which refers to severe BP elevation, was slightly higher in women than in men (5.1% vs. 3.2%) ($p = 0.32$). Prehypertension (SBP, 120–139 or DBP, 80–89 mmHg [Elevated and Stage 1]), which is the risk of developing future hypertension and cardiovascular disease, was found in 39.9% of the men and 30.6% of the women, and the difference was statistically significant ($p = 0.02$). There was a significant association between the stage of hypertension and age in both men and women (men, $p = 0.02$; women, $p < 0.01$). Area of residence in the district had a significant association with hypertension in men ($p = 0.02$) but not in women ($p = 0.82$). Regarding access to healthcare services for hypertension, the

218 prevalence of hypertension was higher among men than among women, and the proportion of
 219 men who had “never had their BP measured” was significantly higher than that of women
 220 (37.3% [125/335] vs. 6.8% [24/354]) ($p < 0.01$). The proportion of participants who had a
 221 previous diagnosis of hypertension was 5.4% [18/335] for men and 10.5% [37/354] for women,
 222 and 2.4% [8/335] of the men and 4.5% [16/354] of the women received antihypertensive
 223 treatment ($p = 0.09, p = 0.25$).

224

225

226 **Table 2. Stage of hypertension relative to demographics and access to care and services among all**
 227 **participants in the Mumbwa district, Central Province of Zambia, 2016**

228

229

230

	Male									
	Overall	Normal	Prehypertension				Hypertension			p value
			Elevated	Stage 1		Stage 2		Hypertensive crisis		
Total n(%)	335	68 (20.4)	33 (9.8)	101 (30.1)	122 (36.5)	11 (3.2)				
Age										
25-44	203	45 (22.0)	26 (12.7)	62 (30.6)	68 (33.4)	3 (1.3)			0.02	
45-64	132	24 (17.9)	7 (5.4)	39 (29.2)	54 (41.2)	8 (6.2)				
Residential area of the district										
Urban area	35	5 (13.2)	1 (2.9)	8 (22.1)	17 (48.5)	5 (13.2)			0.02	
Rural area	300	64 (21.2)	32 (10.6)	93 (31.0)	105 (35.1)	6 (2.1)				
Access to hypertension care and services										
Have never blood pressure measured	125	27 (21.3)	8 (6.1)	35 (27.9)	55 (43.9)	1 (0.8)			0.11	
Diagnosed as hypertensive	18	2 (11.4)	0 (0.0)	2 (11.4)	8 (45.7)	6 (31.4)			<0.01	
On treatment	8	0 (0.0)	0 (0.0)	1 (13.3)	2 (26.7)	5 (60.0)			<0.01	
Medical history (Self-reported)										
HIV positive	28	8 (29.1)	2 (7.3)	9 (30.9)	9 (32.7)	0 (0.0)			0.69	
Diabetic	3	0 (0.0)	0 (0.0)	1 (33.3)	2 (66.7)	0 (0.0)			0.80	

	Female									
	Overall	Normal	Prehypertension				Hypertension			p value
			Elevated	Stage 1		Stage 2		Hypertensive crisis		
Total n(%)	354	127 (35.8)	12 (3.3)	97 (27.3)	101 (28.4)	18 (5.1)				
Age										
25-44	226	105 (46.4)	5 (2.3)	63 (27.7)	50 (22.0)	4 (1.6)			<0.01	
45-64	128	22 (17.3)	7 (5.2)	34 (26.5)	51 (39.8)	14 (11.2)				
Residential area of the district										
Urban area	52	16 (31.7)	2 (3.0)	17 (32.7)	15 (28.7)	2 (4.0)			0.82	
Rural area	302	111 (36.6)	10 (3.4)	80 (26.4)	86 (28.4)	16 (5.3)				
Access to hypertension care and services										
Have never blood pressure measured	24	8 (31.9)	1 (2.1)	9 (38.3)	7 (27.7)	0 (0.0)			0.53	
Diagnosed as hypertensive	37	2 (4.2)	0 (0.0)	7 (19.4)	15 (40.3)	13 (36.1)			0.00	
On treatment	16	0 (0.0)	0 (0.0)	3 (18.8)	6 (34.4)	8 (46.9)			0.00	
Medical history (Self-reported)										
HIV positive	43	19 (44.0)	3 (6.0)	11 (26.2)	10 (23.8)	0 (0.0)			0.36	
Diabetic	2	1 (50.0)	0 (0.0)	0 (0.0)	1 (50.0)	0 (0.0)			0.90	

231

232

233

234

235

236

237

238

239

240

241

242

243

244

Data are number (%)HIV, human immunodeficiency virus
 Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.
 Blood pressure category: Normal (SBP<120 and DBP<80), Elevated (SBP 120-129 and DBP < 80), Stage 1 (SBP 130-139 or DBP 80-89), Stage 2 (SBP≥140 or DBP≥90), Hypertensive crisis (SBP>180 and/or DBP>120)

234

235

236

237

238

239 The present status of hypertension screening and diagnosis is shown in Figure 1. We see that
 240 21.8% (150/689) never had their BP measured, and the main reasons given were ‘do not know
 241 where to obtain the service’ (41.6%), ‘do not have the time or opportunity to check’ (24.8%),
 242 and ‘I think it is not important or I am healthy’ (18.8%). Among the participants who never
 243 had their BP measured, 41.9% (63/150) presented with hypertension at the time of
 244 measurement in this study. Among participants who had their BP measured previously, 89.8%

1
2 245 (485/539) had not been diagnosed with hypertension but 30.3% (147/485) presented with
3 246 hypertension. Among the participants already diagnosed with hypertension, 56.4% (31/55)
4 247 were not taking antihypertensive medication, of which 71.0% (22/31) presented with
5 248 hypertension. Furthermore, most participants taking antihypertensive medication (20/24)
6 249 presented with hypertension, indicating poor BP control.

10 250

12 251

13 252 **Figure 1. Status of screening and diagnosis of hypertension among all participants in the Mumbwa district,**
14 253 **Central Province of Zambia, 2016 (weighted)**

16 254

18 255

19 256 Table 3 shows the proportion of hypertension in relation to each covariate and the association
20 257 of each covariate with hypertension by multivariable analysis (adjusted for the variables that
21 258 showed an association of $p < 0.10$ in the bivariate analysis) in the overall sample analysis and
22 259 the analysis stratified by gender. In the overall sample, older age group (45–64 years) (adjusted
23 260 odds ratio [AOR] = 1.95, 95%CI, 1.35–2.80), higher education (\geq college) (2.00, 95%CI, 1.04–
24 261 3.82), alcohol intake (a few times/week or everyday) (2.14, 95%CI, 1.28–3.58), and BMI \geq 25
25 262 (1.83, 95%CI, 1.24–2.71) were positively associated, while the presence of HIV infection was
26 263 negatively associated with hypertension (0.53, 95%CI, 0.29–0.96). Gender, marital status, food
27 264 insecurity, smoking, physical activity, cooking oil intake, sugar intake, and HbA1c were not
28 265 associated with hypertension. There was a significant association between hypertension and
29 266 alcohol intake in both genders (men \geq a few times/week or everyday: 2.28, 95%CI, 1.24–4.17;
30 267 women \leq a few times/month: 1.79, 95%CI, 1.01–3.19), but the association with urban residence
31 268 was significant only in men (2.46, 95%CI, 1.09–5.56). Older age (45–64 years) (2.68, 95%CI,
32 269 1.56–4.63), higher education (\geq college) (3.39, 95%CI, 1.19–9.64), low-level alcohol intake (\leq a
33 270 few times/month) (1.79, 95%CI, 1.01–3.19), and BMI \geq 25 (1.98, 95%CI, 1.18–3.29) showed
34 271 significant association with hypertension only in women.

46 272

Table 3. Multivariate correlates of hypertension among all participants in the Mumbwa district, Central Province of Zambia, 2016

	Overall				Male				Female			
	Total	Hypertension n(%)	Adjusted OR (95%CI)	p value	Total	Hypertension n (%)	Adjusted OR (95%CI)	p value	Total	Hypertension n(%)	Adjusted OR (95%CI)	p value
Socio-demographic characteristics												
Gender												
Male	335	133 (39.7)	–		–	–	N/A		–	–	N/A	
Female	354	119 (33.5)	–		–	–			–	–		
Age												
25-44	429	124 (28.9)	1 (Reference)		203	70 (34.7)	1 (Reference)		226	53 (23.6)	1 (Reference)	
45-64	260	128 (49.2)	1.95 (1.35-2.80)	0.00	132	63 (47.5)	1.57 (0.95-2.58)	0.07	128	65 (51.0)	2.68 (1.56-4.63)	0.00
Marital Status												
Not married	27	8 (30.8)	1 (Reference)		20	7 (34.2)	–		7	2 (21.4)	–	
Married	557	200 (36.0)	1.57 (0.63-3.90)	0.33	300	120 (40.0)	–		257	81 (31.4)	–	
Divorced/Widow/Widowed	106	43 (40.8)	1.83 (0.68-4.97)	0.23	16	7 (41.9)	–		90	36 (40.6)	–	
Education												
≤primary	513	185 (36.1)	1 (Reference)		229	90 (39.6)	–		284	95 (33.3)	1 (Reference)	
Secondary	127	44 (34.7)	1.03 (0.66-1.61)	0.89	77	32 (41.3)	–		50	12 (24.5)	0.68 (0.32-1.45)	0.32
≥college	49	23 (45.8)	2.00 (1.04-3.82)	0.04	29	11 (36.8)	–		20	12 (59.0)	3.39 (1.19-9.64)	0.02
Work Status												
Employed	85	29 (33.7)	–		58	19 (33.0)	1 (Reference)		28	10 (35.2)	–	
Self-employed	481	178 (37.1)	–		255	102 (39.8)	1.43 (0.74-2.78)	0.29	225	77 (34.0)	–	
Unemployed/Retired	123	45 (36.3)	–		22	12 (55.8)	1.95 (0.66-5.75)	0.22	101	32 (32.0)	–	
Monthly income (USD)												
≤50	326	112 (34.3)	–		157	61 (38.9)	–		169	51 (30.1)	–	
>50	363	140 (38.5)	–		178	72 (40.5)	–		185	68 (36.7)	–	
Residential area of the district												
Urban area	87	39 (44.4)	–		35	22 (61.8)	2.46 (1.09-5.56)	0.03	52	17 (32.7)	–	
Rural area	602	213 (35.4)	–		300	112 (37.2)	1 (Reference)		302	102 (33.7)	–	
Food security												
Secure	192	58 (30.2)	1 (Reference)		107	33 (31.1)	–		85	25 (29.1)	–	
Insecure	497	194 (39.0)	1.46 (0.99-2.16)	0.06	228	100 (43.8)	–		269	94 (34.9)	–	
Family planning												
Not used	–	–	–		–	–	–		249	96 (38.6)	1 (Reference)	
Used	–	–	–		–	–	–		101	96 (21.8)	0.61 (0.32-1.153)	0.13
No data	–	–	–		–	–	–		4	1 (14.3)	–	
Behavioral and psychological characteristics												
Smoking												
Never	525	179 (34.1)	1 (Reference)		185	66 (35.8)	–		340	113 (33.1)	–	
Ex-smoker	86	37 (43.1)	1.26 (0.73-2.18)	0.41	74	33 (44.4)	–		12	4 (34.8)	–	
Current smoker	79	36 (45.8)	1.27 (0.71-2.29)	0.42	76	34 (44.6)	–		3	2 (80.0)	–	
Alcohol												
Never	354	106 (30.0)	1 (Reference)		102	30 (29.3)	1 (Reference)		252	77 (30.3)	1 (Reference)	
≤a few times/month	206	80 (39.0)	1.49 (0.98-2.27)	0.06	123	49 (39.6)	1.43 (0.78-2.59)	0.24	82	31 (38.1)	1.79 (1.01-3.19)	0.046
≥a few times/week or everyday	130	65 (50.4)	2.14 (1.28-3.58)	0.00	110	54 (49.5)	2.28 (1.24-4.17)	0.01	20	11 (55.3)	1.61 (0.58-4.49)	0.37
Physical activity (activities of daily life and sports ≥once a week)												
Neither	54	25 (46.2)	1 (Reference)		16	8 (46.9)	–		38	17 (45.9)	1 (Reference)	
Either	497	182 (36.6)	0.62 (0.34-1.15)	0.13	215	96 (44.7)	–		282	86 (30.4)	0.51 (0.24-1.08)	0.08
Both	138	45 (32.5)	0.66 (0.32-1.35)	0.25	104	29 (28.2)	–		34	15 (45.5)	1.60 (0.54-4.71)	0.40
Cooking oil intake												
Low <20.83ml	510	198 (38.8)	1 (Reference)		244	103 (42.2)	–		266	95 (35.7)	–	

6/bmjopen-2021-006668 on 8 April 2022. Downloaded from http://bmjopen.bmj.com/ on April 19, 2024 by guest. Protected by copyright.

High ≥ 20.83 ml	177	54	(30.5)	0.71	(0.48-1.07)	0.10	92	30	(33.1)	–	85	24	(27.7)	–		
Don't know	1	0	(0.0)	–			–	–		–	1	0	(0.0)	–		
No data	2	0	(0.0)	–			–	–		–	2	0	(0.0)	–		
Sugar intake																
Low < 28.0 g	353	136	(38.6)	1	(Reference)		168	70	(42.0)	–	185	66	(35.6)	–		
High ≥ 28.0 g	335	114	(33.9)	0.79	(0.56-1.12)	0.18	167	62	(37.0)	–	168	52	(30.9)	–		
No data	2	2	(100.0)	–			1	1	(100.0)	–	1	1	(100.0)	–		
Clinical characteristics																
Body mass index (kg/m²)																
Normal (< 25)	505	167	(33.1)	1	(Reference)		281	106	(37.9)	1	(Reference)	224	61	(27.1)	1	(Reference)
Overweight/Obese (≥ 25)	185	85	(46.0)	1.83	(1.24-2.71)	0.00	54	27	(49.1)	1.73	(0.91-3.29)	130	58	(44.7)	1.98	(1.18-3.29)
HbA1c																
Normal (< 5.7)	404	144	(35.6)	1	(Reference)		222	89	(40.1)	–	182	55	(30.1)	1	(Reference)	
High risk (5.7-6.4)	266	96	(36.1)	1.05	(0.73-1.50)	0.80	111	42	(38.1)	–	156	54	(34.7)	1.02	(0.61-1.71)	
Diabetes (≥ 6.5)	18	12	(65.7)	2.69	(0.91-7.96)	0.07	3	2	(66.7)	–	15	10	(65.5)	2.33	(0.68-7.96)	
No data	1	0	(0.00)	–			0	0	(0.0)	–	1	0	(0.0)	–		
Medical history (self-reported)																
HIV infection (Self-reported)																
No	618	232	(37.6)	1	(Reference)		307	124	(40.4)	–	311	108	(34.9)	1	(Reference)	
Yes	71	20	(27.3)	0.53	(0.29-0.96)	0.04	28	9.25	(32.7)	–	43	10	(23.8)	0.54	(0.24-1.24)	

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

"No data" and "Don't know" were excluded from statistical test.

Hypertension is defined as SBP $140 \geq$ mmHg or DBP ≥ 90 mmHg.

280 Table 4 shows factors associated with ‘never had BP measured’ among men, as 83.8%
 281 (125/150) of participants who never had their BP measured were men. In the multivariable
 282 analysis, older age (0.43, 95%CI, 0.25–0.73) and HIV positive status (0.37, 95%CI, 0.14–0.97)
 283 were negatively associated, while being current smoker status was positively associated with
 284 ‘never had BP measured’ (2.09, 95%CI, 1.19–3.66). In contrast, in women, though not shown
 285 in the table, older age was positively associated with ‘never had BP measured’ (4.53, 95%CI,
 286 1.81–11.4).

287

288

289 **Table 4. Bivariate and multivariate correlates of "never had blood pressure measured" (Men only)**

290

	Male (n=335)	Never had blood pressure measured (n=125)				
	Total	n of total (%)	Crude OR (95%CI)	p value	Adjusted OR (95%CI)	p value
Age						
25-44	203	83 (41.0)	1 (reference)		1 (reference)	
45-64	132	42 (31.9)	0.48 (0.30-0.77)	0.00	0.43 (0.25-0.73)	0.00
Residential area of the district						
Urban area	35	6 (16.2)	1 (reference)		1 (reference)	
Rural area	300	120 (39.9)	3.60 (1.35-9.61)	0.01	2.79 (0.98-7.93)	0.06
Education						
Primary	229	96 (42.0)	1 (reference)		1 (reference)	
≥Secondary	106	29 (27.5)	0.62 (0.38-1.01)	0.05	0.84 (0.48-1.45)	0.53
Work Status						
Employed	58	13 (23.2)	1 (reference)		1 (reference)	
Unemployed/Retired	278	112 (40.4)	2.04 (1.08-3.83)	0.03	1.86 (0.92-3.76)	0.09
HIV infection						
No	307	119 (38.7)	1 (reference)		1 (reference)	
Yes	28	7 (23.6)	0.40 (0.16-1.02)	0.06	0.37 (0.14-0.97)	0.04
Smoking						
Never, Ex-smoker	259	87 (33.7)	1 (reference)		1 (reference)	
Current smoker	76	38 (50.0)	2.01 (1.19-3.38)	0.01	2.09 (1.19-3.66)	0.01
Alcohol						
Never or a few times/month	225	85 (37.9)	1 (reference)		—	
≥a few times/week or everyday	110	40 (36.4)	1.03 (0.64-1.66)	0.91	—	
Body mass index (kg/m²)^c						
Normal (<25)	281	111 (39.6)	1 (reference)		1 (reference)	
Overweight/Obese (≥25)	54	14 (26.4)	0.43 (0.22-0.85)	0.02	0.66 (0.32-1.40)	0.28

Data are number (%).

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

OR: Odds ratio

291

292

293

294

295

296

297 **DISCUSSION**

298 This study assessed the prevalence and the risk factors of hypertension among the genders to
299 understand the current situation of hypertension among rural residents in Zambia. We also
300 explored the status of screening and diagnosis of hypertension and their correlates to evaluate
301 the situation of access to healthcare services for hypertension.

302 In this study, we found that more than 35% of the participants had hypertension, and the
303 profile of hypertension correlates was different between men and women. More than 80% of
304 the people with high BP measurements had never been previously diagnosed with hypertension,
305 and over 40% of them had never had their BP measured, suggesting the lack of access to or
306 availability of healthcare services for BP control among the studied population.

307

308 **The prevalence of hypertension in rural areas**

309 The prevalence of hypertension among the targeted rural residents of this study in 2016 was
310 39.7% in men and 33.5% in women, respectively, both being much higher than the national
311 averages found in the Zambia STEPS Survey of 2017 (20.5% in men and 17.6% in women).[7]
312 Previous research has reported mixed findings regarding the prevalence of hypertension in rural
313 areas of Zambia. While similar prevalence of hypertension (46.9%) was reported among people
314 attending health check-up in other rural area of Zambia,[17] it was only 23.1% in a primary
315 healthcare-based study conducted in several rural districts between 2011–2014.[18] Comparing
316 our results with those of previous studies is, however, difficult due to methodological
317 differences. For example, previous studies were based on convenient samples with potential
318 selection bias, while our study was based on probability sample of the whole area. Studies
319 using probability sampling are needed for documenting the accurate status of blood pressure
320 among Zambian rural populations. The prevalence of prehypertension and hypertension was
321 slightly higher in men than in women in our study, a tendency that has been observed
322 throughout the African region.[6]

323

324 **Gender differences in factors associated with hypertension**

325 In this study, a difference in gender was found not only in the prevalence of hypertension, but
326 also in the profile of the correlates of hypertension. In men, residence in the urban area of the
327 district and high frequency of alcohol intake were significantly associated with hypertension.
328 While in women, older age, higher education level, low frequency of alcohol intake, and BMI
329 ≥ 25 were associated with hypertension, suggesting the different mechanism(s) involved in the
330 development of high blood pressure between the genders. This implies that different pathways

1
2 331 for hypertension including behavioural and socio-cultural factors exist in men and women,
3 332 which could affect prevention strategies [19]

4 333 Alcohol consumption was the only factor moderately associated with hypertension in both
5 334 genders, which is in line with well-established findings worldwide.[20] Although the exact
6 335 mechanism is unclear, it can be caused directly through the chronic effect of alcohol and/or
7 336 indirectly through related socioeconomic status and lifestyles among the study population.[20]
8 337 Regardless of the mechanism, however, it is important to follow the trend of alcohol intake
9 338 over time with special attention to the type, amount, and pattern since it may rapidly change in
10 339 both quantity and quality with future economic growth.

11 340 Living in the urban area of the district was significantly associated with hypertension only
12 341 in men. Although the study region was “rural” in general (neighbouring the capital city,
13 342 Lusaka), there are some areas with relatively easy access to the capital city. Men living in such
14 343 areas may be involved in urbanized lifestyles, probably in relation to their jobs, in terms of
15 344 eating habits and lifestyles, including high calorie diets and lack of exercise. Studies in
16 345 Cameroon and Mali have shown a similar tendency with higher prevalence of hypertension
17 346 among men in ‘urban areas’ than in rural areas.[21,22]

18 347 The relationship between age and hypertension has been reported in SSA countries [23-
19 348 25]. In our study, a significant association with age was observed only in women, reflecting
20 349 the age-related distribution of hypertension between the genders, where the difference of the
21 350 proportions of hypertension between younger (25–44 years) and older (45–64 years) age
22 351 groups was large (23.6% vs. 51.0%, respectively) in women, but small in men (34.7% and
23 352 47.4%, respectively). Similar age disparities in the proportion of hypertension by gender have
24 353 been reported in previous studies of Zambia and Senegal.[18,25] This may suggest that men
25 354 are more likely to develop hypertension at a younger age than women. The reasons for this age
26 355 disparity by gender should be one of the focus points in future research.

27 356 An association between hypertension and education level was observed only among women.
28 357 Slightly high odds of hypertension in people with higher levels of education were also observed
29 358 in the study in Malawi [21]. This may suggest that in SSA countries that experienced rapid
30 359 economic growth in recent years, the risk of hypertension has increased among people with
31 360 higher levels of education due to spread of urbanized eating habits and lifestyles (over-nutrition
32 361 and physically inactive).[26] The reason why the association was detected only in women in
33 362 our study is unclear but higher education may be related to urbanized eating habits and
34 363 lifestyles more in women than in men.

35 364 The association between ‘overweight and obesity’ (BMI \geq 25) and hypertension has been
36 365 reported in SSA countries including Zambia, with its tendency being stronger in women than

1
2 366 in men.[27] Similarly in our study, although the association was observed both in both genders,
3 367 it was significant only in women. This may be related to biological factors such as an increase
4
5 368 in obesity with age in women in African societies and their cultural preferences. In men,
6
7 369 behavioural factors such as alcohol consumption and psychological stress may be more likely
8
9 370 to be associated with developing hypertension than obesity.

10 371

11 12 372 **Status of Hypertension Management**

13
14 373 In this study, only 16.7% of the participants who presented with hypertension had previously
15
16 374 been diagnosed with hypertension. Among the participants with documented hypertension but
17
18 375 no previous diagnosis, 30% never had their BP measured. Our results concur with findings
19
20 376 from a systematic review of hypertension in the SSA indicating that only 22.5% of people with
21
22 377 hypertension had already been diagnosed with hypertension.[18] This indicates the need to
23
24 378 strengthen screening and diagnosis of hypertension particularly at the primary healthcare level
25
26 379 which is the entry level to health care systems in most SSA countries.

27
28 380 Moreover, only 8% of the participants in this study reported having been previously
29
30 381 diagnosed with hypertension, which was much lower than the actual proportion presenting with
31
32 382 hypertension in our study. In addition, only fewer than half of the participants diagnosed with
33
34 383 hypertension were taking antihypertensive medications, and of them, many presented with
35
36 384 hypertension at the time of the measurement, indicating challenges in accessing treatment and
37
38 385 management of hypertension. A previous study in Zambia reported that 18% of people who
39
40 386 presented with hypertension at the time of the study had been prescribed antihypertensive
41
42 387 medication at a health centre.[18] In our study, only 7.9% of the participants with hypertension
43
44 388 had been prescribed antihypertensive medication. Furthermore, about 83% of the participants
45
46 389 who reported taking antihypertensive medication in our study presented hypertension at the
47
48 390 time of measurement. This was consistent with the results of a previous study in Zambia where
49
50 391 nearly 90% had poorly controlled hypertension,[18] and other reports from the entire SSA
51
52 392 region.[2] These results indicate that there are various challenges in the management of
53
54 393 hypertension in the rural areas of Zambia, as in other SSAs, in terms of ‘difficulties in accessing
55
56 394 appropriate treatment and health services including hypertension’, ‘lack of screening and
57
58 395 diagnostic opportunities for hypertension’, and ‘lack of awareness of the importance of BP
59
60 396 control’.

397

398

399

400 **Access to healthcare services related to hypertension**

401 In this study, we also assessed the differences in access to healthcare services related to
402 hypertension between the genders. Identifying the management status of hypertension (care
403 cascade) is important evidence that can contribute to health policy and interventions.[28] We
404 specifically focused on the 'history of BP measurement' as it relates to the awareness of having
405 hypertension. In our study, more than 20% of the participants reported have never having had
406 their BP measured previously, suggesting the difficulties in accessing screening and diagnostic
407 services for hypertension care. In particular, despite the higher prevalence of hypertension
408 among men than among women, the proportion of men who 'never had their BP measured'
409 was 37.4%, which was 5.5 times higher than that of women. Men also tended to be less likely
410 to have been diagnosed with and treated for hypertension.

411 There was a significant positive association between smoking and 'never had their BP
412 measured' in men. While this finding requires further assessment in future research, it may
413 suggest that people who engage in high-risk health behaviours such as smoking tend to be less
414 concerned about their health and less likely to engage in health seeking behaviours than those
415 who do not engage in such behaviour. In this study, we also included self-reported HIV status
416 in the analysis as a factor affecting access to healthcare services. Men in older age groups and
417 men with HIV-positive status were less likely to have 'never had their BP measured before',
418 suggesting that they were likely to be aware of their BP. The association with 'older age group'
419 may be due to the fact that they were likely to receive medical care during their lifetime.
420 Regarding the association with 'HIV-positive', all HIV-positive individuals were receiving
421 HIV treatment, so regular medical consultations at a healthcare facility may have been the
422 important opportunity for BP measurement.

423 Men have fewer opportunities to access healthcare services other than for illness or injury,
424 than women who visit healthcare facilities for maternal and child health services. Patients with
425 asymptomatic conditions like hypertension may not receive the required healthcare services
426 due to psychological and geographical barriers, e.g., low level of attention to health or distance
427 to healthcare facilities. Therefore, along with strengthening the screening for hypertension, we
428 suggest that the use of existing mobile health services, such as vaccination campaigns, mobile
429 voluntary counselling and testing services (VCTs), and cooperation with community health
430 workers may be advantageous in treating many people.[29] For women, although the number
431 of people who never had their BP measured was too few, the odds of never having BP measured
432 were significantly higher in the older age group. This gender difference will need to be
433 examined in further research with a large sample size.

434

435 **Strengths and limitations**

436 The strength of our study is that we used multi-stage cluster random sampling and obtained a
437 relatively high response rate. Thus, our results are representative of patients at risk of CVD in
438 the target population in the rural area. In terms of limitations, the recorded BP may have been
439 higher than usual due to white coat hypertension. Socially desirable responses due to face-to-
440 face interviews could also have affected the results, even though we trained interviewers before
441 the study. Unmeasured factors may have affected some of the associations found in our study.

443 **CONCLUSION**

444 We found that more than one-third of the participants in a rural district in Zambia had
445 hypertension. Among them, most were not diagnosed with hypertension yet and one-quarter of
446 them never had their BP measured. These results indicate a serious lack of CVD prevention
447 services, including access to and availability of healthcare services for hypertension, among
448 rural residents in Zambia. Therefore, health promotion and screening strategies for
449 hypertension are urgently required, especially in primary healthcare settings in rural areas.
450 Particular attention should be paid to healthcare access, specifically among men.

452 **Acknowledgments**

453 We are thankful to all the participants, the staff at the Institute of Economic and Social Research,
454 the University of Zambia, and the members of the Mumbwa District Medical Office for the
455 local arrangement and support to this study. We gratefully acknowledge the local nurses and
456 research assistants for their assistance with the data collection. Lastly, we extend our gratitude
457 to the late Mubiana Macwan'gi for contributing to the supervision of the research in Zambia.

459 **Contributors**

460 YT, TT, MOK and MK contributed to study conception and design. YT, RZ and CD
461 contributed to the data collection. YT, TT and MK contributed to data analysis and drafted the
462 manuscript. YT, PPM, SPS, OA, RZ and CD revised the manuscript. MOK and MK supervised
463 the study. All authors read and approved the final manuscript.

465 **Funding**

1
2 466 This study was supported by the United Nations University, Institute for the Advanced Study
3 467 of Sustainability (Global Leadership Training Program in Africa) and the Inter-Graduate
4 468 School Program for Sustainable Development and Survivable Societies, Kyoto University. It
5 469 was also supported by the 2016 Kyoto University School of Public Health Super Global
6 470 Course's travel scholarship.

471

472 **Competing interests**

473 None declared.

474

475 **Patient consent for publication**

476 Not required.

477

478 **Data sharing statement**

479 Data are available upon reasonable request.

480

481 **References**

- 482 1. World Health Organization. Fact sheets. Cardiovascular diseases (CVDs).
483 [https://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
484 [Accessed May 2021]
- 485 2. Ataklte F, Erqou S, Kaptoge S, et al. Burden of undiagnosed hypertension in sub-Saharan Africa: a
486 systematic review and meta-analysis. *Hypertension*. 2015;65:291–98.
487 doi:10.1161/HYPERTENSIONAHA.114.04394
- 488 3. Dzudie A, Rayner B, Ojji D, et al. Roadmap to Achieve 25% Hypertension Control in Africa by 2025.
489 *Glob Heart*. 2018;13:45–59. doi: 10.1016/j.ghheart.2017.06.001.
- 490 4. Boateng D, Wekesah F, Browne JL, et al. Knowledge and awareness of and perception towards
491 cardiovascular disease risk in sub-Saharan Africa: A systematic review. *PLoS One*. 2017; 12:e0189264.
492 doi: 10.1371/journal.pone.0189264.
- 493 5. Berry KM, Parker WA, Mchiza ZJ, et al. Quantifying unmet need for hypertension care in South Africa
494 through a care cascade: evidence from the SANHANES, 2011-2012. *BMJ Glob Health*. 2017;
495 16;2:e000348. doi: 10.1136/bmjgh-2017-000348.
- 496 6. World Health Organization. Global status report on noncommunicable diseases 2014.
497 https://apps.who.int/iris/bitstream/handle/10665/148114/9789241564854_eng.pdf.
498 [Accessed May 2021]
- 499 7. World Health Organization. Zambia STEPS Survey 2017 Factsheet
500 <https://www.afro.who.int/publications/zambia-steps-survey-2017-factsheet>. [Accessed

- 1 501 May 2021]
- 2 501
- 3 502 8. Yan, L.D., Chi, B.H., Sindano, N, et al. Prevalence of hypertension and its treatment among adults
- 4 503 presenting to primary health clinics in rural Zambia: analysis of an observational database. *BMC Public*
- 5 504 *Health* 2015;15: 933. <https://doi.org/10.1186/s12889-015-2258-4>
- 6 504
- 7 505 9. Mulenga D, Siziya S, Rudatsikira E, et al. District specific correlates for hypertension in Kaoma and
- 8 506 Kasama rural districts of Zambia. *Rural Remote Health*. 2013;13:2345.
- 9 506
- 10 507 10. Yan LD, Chi BH, Sindano N, et al. Prevalence of hypertension and its treatment among adults
- 11 508 presenting to primary health clinics in rural Zambia: analysis of an observational database. *BMC Public*
- 12 508 *Health*. 2015;15:933. doi: 10.1186/s12889-015-2258-4.
- 13 509
- 14 510 11. Central Statistical Office Zambia. 2010 Census of Population and Housing—Volume 11 National
- 15 510 Descriptive Tables. 2012. Available from:
- 16 511 <http://catalog.ihnsn.org/index.php/catalog/4124/download/55313>. [Accessed May 2021].
- 17 511
- 18 512 12. World Health Organization. WHO STEPS Surveillance Manual. 2017. Available from:
- 19 512 http://www.who.int/ncds/surveillance/steps/STEPS_Manual.pdf?ua=1. [Accessed May 2021].
- 20 513
- 21 514 13. BeLue R, Okoror TA, Iwelunmor J, et al. An overview of cardiovascular risk factor burden in sub-
- 22 514 Saharan African countries: a socio-cultural perspective. *Glob Health* 2009;5:10. doi: 10.1186/1744-
- 23 515 8603-5-10.
- 24 516
- 25 517 14. Tateyama Y, Musumari PM, Techasrivichien T, et al. Dietary habits, body image, and health service
- 26 517 access related to cardiovascular diseases in rural Zambia: A qualitative study. *PLoS One*. 2019;
- 27 518 14:e0212739. doi: 10.1371/journal.pone.0212739
- 28 518
- 29 519 15. Coates J, Anne S, Paula B. Household Food Insecurity Access Scale (HFIAS) for Measurement of
- 30 519 Household Food Access: Indicator Guide (v. 3). 2007. Available from:
- 31 520 http://www.fao.org/fileadmin/user_upload/eufao-fsi4dm/doc-training/hfias.pdf Cited February 9 2018
- 32 521 [Accessed May 2021]
- 33 522
- 34 523 16. Kessler RC, Green JG, Gruber MJ, et al. Screening for serious mental illness in the general population
- 35 523 with the K6 screening scale: results from the WHO World Mental Health (WMH) survey initiative. *Int*
- 36 524 *J Methods Psychiatr Res*. 2010;19 Suppl 1:4–22. doi:10.1002/mpr.310
- 37 524
- 38 525 17. Rush KL, Goma FM, Barker JA, et al. Hypertension prevalence and risk factors in rural and urban
- 39 525 Zambian adults in western province: a cross-sectional study. *Pan Afr Med J*. 2018;30:97. doi:
- 40 526 10.11604/pamj.2018.30.97.14717.
- 41 527
- 42 528 18. Yan LD, Chi BH, Sindano N, et al. Prevalence of hypertension and its treatment among adults
- 43 528 presenting to primary health clinics in rural Zambia: analysis of an observational database. *BMC Public*
- 44 529 *Health*. 2015;15:933. <https://doi.org/10.1186/s12889-015-2258-4>
- 45 530
- 46 531 19. Ghosh S, Mukhopadhyay S, Barik A. Sex differences in the risk profile of hypertension: a cross-
- 47 531 sectional study. *BMJ Open*. 2016;6:e010085. doi: 10.1136/bmjopen-2015-010085.
- 48 532
- 49 533 20. Beilin LJ, Puddey IB. Alcohol and hypertension: an update. *Hypertension*. 2006;47:1035–38.
- 50 533 doi:10.1161/01.HYP.0000218586.21932.3c
- 51 534
- 52 535 21. Cooper R, Rotimi C, Ataman S, et al. The prevalence of hypertension in seven populations of west
- 53 535 African origin. *Am J Public Health*. 1997; 87:160–8. doi: 10.2105/ajph.87.2.160.
- 54 536
- 55 537 22. Bâ HO, Camara Y, Menta I, et al. Hypertension and Associated Factors in Rural and Urban Areas Mali:
- 56 537
- 57 538
- 58 539
- 59 540
- 60 540

- 1
2 541 Data from the STEP 2013 Survey. *Int J Hypertens*. 2018;2018:6959165. doi:10.1155/2018/6959165
- 3 542 23. Berry KM, Parker WA, Mchiza ZJ, et al. Quantifying unmet need for hypertension care in South Africa
4 543 through a care cascade: evidence from the SANHANES, 2011-2012. *BMJ Glob Health*.
5 544 2017;2:e000348. doi:10.1136/bmjgh-2017-000348
- 6
7 545 24. Price AJ, Crampin AC, Amberbir A, et al. Prevalence of obesity, hypertension, and diabetes, and
8 546 cascade of care in sub-Saharan Africa: a cross-sectional, population-based study in rural and urban
9 547 Malawi. *Lancet Diabetes Endocrinol*. 2018;6:208–22. doi:10.1016/S2213-8587(17)30432-1
- 10
11 548 25. Kwarisiima D, Balzer L, Heller D, et al. Population-Based Assessment of Hypertension Epidemiology
12 549 and Risk Factors among HIV-Positive and General Populations in Rural Uganda. *PLoS One*.
13 550 2016;11:e0156309. doi:10.1371/journal.pone.0156309
- 14
15 551 26. Macia E, Gueye L, Duboz P. Hypertension and Obesity in Dakar, Senegal. *PLoS One*.
16 552 2016;11:e0161544. doi:10.1371/journal.pone.0161544
- 17
18 553 27. Akpa OM, Made F, Ojo A, et al. Regional Patterns and Association Between Obesity and Hypertension
19 554 in Africa: Evidence from the H3Africa CHAIR Study. *Hypertension*. 2020;75:1167–78.
20 555 doi:10.1161/HYPERTENSIONAHA.119.14147
- 21
22 556 28. Geldsetzer P, Manne-Goehler J, Marcus ME, et al. The state of hypertension care in 44 low-income and
23 557 middle-income countries: a cross-sectional study of nationally representative individual-level data from
24 558 1·1 million adults. *Lancet*. 2019;394:652–62. doi:10.1016/S0140-6736(19)30955-9
- 25
26 559 29. Opie LH, Seedat YK. Hypertension in sub-Saharan African populations. *Circulation*.
27 560 2005;112:3562-3568. doi:10.1161/CIRCULATIONAHA.105.539569
- 28
29
30
31
32 561
33
34 562

563 Appendix

564

565 Table 1: Bivariate and multivariate correlates of 'never had blood pressure measured' (Women)

	Female (n=354)		Never had blood pressure measured (n=24)			
	Total	n of total (%)	Crude OR (95CI)	p value	Adjusted OR (95CI)	p value
Age						
25-44	226	6 (2.5)	1 (reference)		1 (reference)	
45-64	128	19 (14.5)	5.33 (2.18-13.06)	0.00	4.53 (1.81-11.35)	0.00
Residential area						
Urban	52	4 (6.9)	1 (reference)		—	
Rural	302	21 (6.8)	1.39 (0.40-4.81)	0.60	—	
Education						
<=primary	284	19 (6.7)	1 (reference)		—	
>=Secondary	70	5 (7.3)	0.997 (0.36-2.75)	0.995	—	
Work Status						
Employed	28	3 (9.3)	1 (reference)		—	
Unemployed/Retired	326	22 (6.6)	0.71 (0.20-2.50)	0.59	—	
HIV infection						
No	311	23 (7.3)	1 (reference)		—	
Yes	43	2 (3.6)	0.83 (0.24-2.89)	0.77	—	
Smoking						
Never, Ex-smoker	352	24 (6.7)	1 (reference)		—	
Current smoker	3	1 (20.0)	6.60 (0.58-75.32)	0.13	—	
Alcohol						
Never or a few times/m	335	20 (6.0)	1 (reference)		1 (reference)	
≥a few times/w or everyday	20	4 (21.1)	3.37 (1.04-10.88)	0.04	2.19 (0.65-7.43)	0.21
Body mass index (kg/m²)						
Normal (<25)	224	11 (4.8)	1 (reference)		1 (reference)	
Overweight/Obese (25 and over)	130	13 (10.3)	2.12 (0.95-4.72)	0.07	1.71 (0.74-3.92)	0.21

Data are number (%).

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

OR: Odds ratio

566

567

568

569

570

Table 2: Bivariate and multivariate correlates of 'never had blood pressure measured' (Overall)

	Overall (n=689)		Never had blood pressure measured (n=150)			
	Total	n of total (%)	Crude OR (95CI)	p value	Adjusted OR (95CI)	p value
Gender						
Male	335	125 (37.4)	7.71 (4.88-12.18)	0.00	6.27 (3.84-10.23)	0.00
Female	354	24 (6.8)	1 (reference)		1 (reference)	
Age						
25-44	429	89 (20.7)	1 (reference)		—	
45-64	260	61 (23.3)	0.90 (0.61-1.31)	0.57	—	
Residential area						
Urban	87	9 (10.7)	1 (reference)		1 (reference)	
Rural	602	140 (23.3)	3.02 (1.43-6.41)	0.00	2.70 (1.22-5.98)	0.01
Education						
<=primary	513	115 (22.5)	1 (reference)		—	
>=Secondary	177	34 (19.5)	0.96 (0.63-1.45)	0.84	—	
Work Status						
Employed	85	16 (18.7)	1 (reference)		—	
Unemployed/Retired	604	134 (22.1)	1.16 (0.67-2.01)	0.60	—	
HIV infection						
No	618	141 (22.9)	1 (reference)		1 (reference)	
Yes	71	8 (11.5)	0.44 (0.21-0.90)	0.03	0.46 (0.21-0.995)	0.049
Smoking						
Never, Ex-smoker	611	111 (18.2)	1 (reference)		1 (reference)	
Current smoker	79	39 (49.0)	4.46 (2.74-7.28)	0.00	2.11 (1.19-3.73)	0.01
Alcohol						
Never or a few times/m	560	105 (18.8)	1 (reference)		1 (reference)	
≥a few times/w or everyday	130	44 (34.1)	2.29 (1.50-3.50)	0.00	0.93 (0.55-1.55)	0.77
Body mass index (kg/m²)						
Normal (<25)	505	122 (24.1)	1 (reference)		1 (reference)	
Overweight/Obese (25 and over)	185	28 (15.0)	0.50 (0.31-0.79)	0.00	0.91 (0.55-1.53)	0.73

Data are number (%).

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

OR: Odds ratio

571

572

573

574

575


576

577

578

579

580



2021-055668 on 8 April 2022. Downloaded from <http://bmjopen.bmj.com/> on April 19, 2024 by guest. Protected by copyright.

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	3
Objectives	3	State specific objectives, including any prespecified hypotheses	3
Methods			
Study design	4	Present key elements of study design early in the paper	4
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	4
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	4
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5
Bias	9	Describe any efforts to address potential sources of bias	5
Study size	10	Explain how the study size was arrived at	4
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	5
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	5
		(b) Describe any methods used to examine subgroups and interactions	5
	(c) Explain how missing data were addressed	6	
	(d) If applicable, describe analytical methods taking account of sampling strategy	4	
	(e) Describe any sensitivity analyses	6	
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6
		(b) Give reasons for non-participation at each stage	6
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	6,7
		(b) Indicate number of participants with missing data for each variable of interest	NA
Outcome data	15*	Report numbers of outcome events or summary measures	7,8
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	9,10

		(b) Report category boundaries when continuous variables were categorized	8,11
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	12
Discussion			
Key results	18	Summarise key results with reference to study objectives	13
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	17
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	17
Generalisability	21	Discuss the generalisability (external validity) of the study results	17
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	17

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Hypertension, its correlates, and differences in access to healthcare services by gender among rural Zambian residents: a cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-055668.R1
Article Type:	Original research
Date Submitted by the Author:	08-Dec-2021
Complete List of Authors:	Tateyama, Yukiko; Kyoto University, Health Service; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research Techasrivichien, Teeranee ; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research; International Institute of Socio-epidemiology Musumari, Patou Masika ; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research; International Institute of Socio-epidemiology Sugimoto, S. Pilar; International Institute of Socio-epidemiology; National Institute of Health Ongosi, Anita; International Institute of Socio-epidemiology Zulu, Richard; Center for Primary Care Research, University of Zambia School of Medicine Dube, Christopher; Ndola District Health Office Ono-Kihara, Masako; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research; International Institute of Socio-epidemiology Kihara, Masahiro; Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research; International Institute of Socio-epidemiology
Primary Subject Heading:	Global health
Secondary Subject Heading:	Public health
Keywords:	Hypertension < CARDIOLOGY, International health services < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, PRIMARY CARE, PUBLIC HEALTH, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Hypertension, its correlates, and differences in access to healthcare services by gender among rural Zambian residents: a cross-sectional study

Author information

Yukiko Tateyama,^{1,2,3*} Teeranee Techasrivichien,^{2,3} Patou Masika Musumari,^{2,3} S. Pilar Suguimoto,^{3,6} Anita Ongosi,³ Richard Zulu,⁴ Christopher Dube,⁵ Masako Ono-Kihara,^{2,3} Masahiro Kihara^{2,3}

Author affiliations

1. Kyoto University Health Service, Kyoto, Japan
2. Kyoto University, Global Health Interdisciplinary Unit, Center for the Promotion of Interdisciplinary Education and Research, Kyoto University, Kyoto, Japan
3. International Institute of Socio-epidemiology, Kyoto, Japan
4. Center for Primary Care Research, University of Zambia School of Medicine, Lusaka, Zambia
5. Ndola District Health Office, Ndola, Zambia
6. National Institute of Health, Lima, Peru

*Corresponding author

Postal address: Yoshida-honmachi, Sakyo-ku, Kyoto, Japan 606-8501

Email: tateyama.yukiko.57c@kyoto-u.jp

Word Count: 4088 words

33 ABSTRACT

34 **Objectives:** To examine the prevalence of hypertension and access to related healthcare
35 services among rural residents of Mumbwa district in Zambia.

36 **Design:** Cross-sectional study with probability cluster sampling.

37 **Setting:** Rural Zambia.

38 **Participants:** We recruited 690 residents from Mumbwa district aged 25–64 years who had
39 been living in the study area for ≥ 6 months and had adopted the lifestyle of the study area.
40 Pregnant women and women who had given birth in the past 6 months were excluded. The data
41 collection—questionnaire survey and anthropometric and biological measurements—was
42 conducted between May and July 2016.

43 **Results:** In the overall sample, 39.7% and 33.5% of the men and women had hypertension
44 (SBP ≥ 140 or DBP ≥ 90 mmHg), respectively. Among the participants without a previous
45 diagnosis of hypertension, 30.3% presented with hypertension at the time of measurement. In
46 the multivariable analysis, alcohol intake and urban residence in men, and older age group,
47 higher education, and body mass index ≥ 25 kg/m² in women were significantly associated with
48 hypertension. Among the 21.8% who never had their blood pressure (BP) measured, 83.8%
49 were men; among these men, older age (adjusted odds ratio [AOR], 0.43; 95% confidence
50 interval [CI], 0.25–0.73) and HIV positive status (AOR, 0.37; 95%CI, 0.14–0.97) were
51 negatively associated, while current smoker status (AOR, 2.09; 95%CI, 1.19–3.66) was
52 positively associated with the lack of BP measurements.

53 **Conclusion:** We found that hypertension is prevalent in the target rural area. However, many
54 were not aware of their hypertension status and many never had their BP measured, indicating
55 a serious gap in cardiovascular disease prevention services in Zambia. There is an urgent need
56 for health promotion and screening for hypertension, especially in the primary health services
57 of rural Zambia. Issues related to healthcare accessibility in men require particular attention.

59 KEY WORDS

60 Hypertension, blood pressure, rural, cluster sampling, access, health promotion

62 Strengths and limitations of this study

- 63 • This study assessed the prevalence and factors associated with hypertension stratified by
64 gender, to understand the current hypertension status among rural residents of Zambia.

- 65 • We employed a multi-stage cluster random sampling method and obtained a relatively
66 high response rate, which helped ensure that the results are representative of the target
67 population.
- 68 • Socially desirable responses due to face-to-face interviews might have affected the
69 results.

70

71 INTRODUCTION

72 Hypertension is a major global health concern; currently, 17.9 million mortality cases are
73 reported yearly due to coronary heart disease and stroke worldwide.[1] The burden of
74 hypertension has increased globally during the past quarter century and accounts for 7% of
75 disability-adjusted life years.[2] It has been reported that if no action is taken to control
76 hypertension, economic losses will outstrip public healthcare spending.[3]

77 It is difficult to be aware of hypertension without assessment during the early stages
78 because it is asymptomatic.[3] To address the increasing prevalence of hypertension, early
79 detection and awareness are important, particularly at the primary healthcare level. However,
80 many people with hypertension in sub-Saharan Africa (SSA) may remain undiagnosed,
81 untreated, or uncontrolled because of an inadequate healthcare system.[4] In fact, a systematic
82 review of SSA studies reported that only 22.5% of people with hypertension were aware of
83 their status.[2] Additionally, a South African study found that among people with hypertension,
84 51% ever had their blood pressure (BP) measured, of which nearly half had not been informed
85 of their high BP.[5] This indicates a lack of necessary health services for prevention and
86 screening of hypertension, particularly in developing countries.

87 SSA has been reporting rising rates of hypertension, with the highest prevalence rate
88 worldwide (46% of adults aged ≥ 25 years),[6] and the prevalence remains high in Zambia as
89 well (19.0% in 2017).[7] A recent study in Zambia found that the prevalence of hypertension
90 in people aged over 25 years in rural settings was 23.1%.[8] However, this information was
91 obtained from the clinical visit records at primary healthcare facilities and did not include
92 people without access to health facilities.

93 The aforementioned evidence underscores the importance of strengthening the assessment
94 and treatment of hypertension. However, research on screening and diagnosis of hypertension
95 has been limited in Zambia.[8-10] Therefore, we aimed to investigate the prevalence of
96 hypertension including undiagnosed cases to understand the current status and access to

1
2 97 healthcare service for hypertension among rural residents in Zambia. We also examined the
3 98 correlation of demographic, behavioural, and biological factors with hypertension.
4
5
6 99

8 100 **METHODS**

10 111 **Design and settings**

12 102 This was a cross-sectional study conducted between May and July 2016 in Zambia. We selected
13 103 Mumbwa district in Central Province as our study area because it is a typical rural area
14 104 experiencing urbanization and economic growth while maintaining traditional culture. The
15 105 district is located 150 km west of the capital Lusaka city and is home to approximately 210,847
16 106 inhabitants—15% in urban areas and 85% in rural areas.[11] The target population included
17 107 residents aged 25–64 years. Since the objective of this study was to investigate lifestyle-related
18 108 risk factors, we only included residents who had been living in the study area for ≥ 6 months
19 109 and had adopted the prevalent lifestyle of the study area. Pregnant women and women who had
20 110 given birth in the last 6 months were excluded because of potentially different dietary habits
21 111 and lifestyles and the fact that prepertum and postpartum weight could affect their
22 112 anthropometric and biological data.
23
24
25
26
27
28
29
30
31

32 113 33 34 114 **Sampling**

35 115 We employed a three-stage probability proportional to size (PPS) cluster sampling. The sample
36 116 size calculation was based on the recommendations of the WHO STEPwise approach to
37 117 surveillance (STEPS),[12] assuming a 95% confidence level, 5% margin of error (e2), and 30%
38 118 prevalence of hypertension in rural areas.[12] The minimum sample size required was 167
39 119 subjects, which was increased to 800 to address design effects (loss of sampling efficiency due
40 120 to cluster sampling), an assumed 20% non-response rate, and planned subgroup and
41 121 multivariate analyses.
42
43
44
45
46
47

48 122 The Central Statistical Office (CSO) of Zambia provided the list of study sampling clusters and
49 123 Standard Enumeration Areas (SEAs). In the first stage, we selected 32 SEAs through PPS
50 124 sampling without replacement using the sampling frame of the Zambia Population and Housing
51 125 Census 2010.[11] In the second stage, mappers from CSO and research assistants mapped each
52 126 selected SEA and listed all households and their eligible members. Then, using the list of each
53 127 SEA created, a total of 25 households in each SEA were selected through systematic sampling,
54 128 which uses a random starting point and a sampling interval calculated by dividing the total
55 129 number of households in each SEA. In the third stage, from each selected household, only one

1
2 130 individual was selected using the Kish Household Coversheet based on the WHO STEPS.[12]
3 131 We scheduled a date and place to administer the questionnaire survey and take anthropometric
4 132 and biological measurements as per the participants' convenience. We met with all recruited
5 133 individuals (if absent, their family members or closest neighbours) 1–2 days before testing to
6 134 request them to start fasting at 8:00 pm on the day prior to the biological measurements and to
7 135 visit the testing venue on the scheduled date.
8
9
10
11
12
13

14 137 **Data collection**

16 138 The questionnaire was developed in English and three local languages based on the review of
17 139 Zambian and international literature[9,13] and the results of an earlier qualitative study. [14]
18 140 A pilot study was conducted to resolve language discrepancies, to assess the face validity of
19 141 the questionnaire and test-retest reliability, and confirm the feasibility of anthropometric and
20 142 biological measurements. Face-to-face interviews were carried out by field staff at venues such
21 143 as the participant's home, community meeting places, or schools. Additionally, licensed nurses
22 144 were recruited and trained to collect anthropometric measurements and biological samples. A
23 145 nurse explained the results of blood and urine tests to the participants following their
24 146 cooperation with the study, then soap and washing paste were given as rewards for participation.
25
26
27
28
29
30
31
32

33 148 **Measurements**

35 149 BP was measured using electronic equipment (Omron HEM-7130-HP, Omron Corporation,
36 150 Kyoto, Japan). Three measurements were taken from the participants at three-minute intervals
37 151 while they were seated after 15 minutes of rest, and the average of the last two readings was
38 152 recorded. Weight was measured while the participants were barefoot and wearing light clothing
39 153 using an electronic scale (Omron HBF-223-G, Omron Corporation, Kyoto, Japan). Glycated
40 154 haemoglobin (HbA1c) and blood lipids (total cholesterol, LDL-cholesterol, HDL-cholesterol,
41 155 and triglycerides) were measured using point-of-care testing device (Cobas b 101, Roche
42 156 Diagnostics K.K., Tokyo, Japan). Other variables used in the analysis included socio-
43 157 demographic characteristics, food security using the Household Food Insecurity Access
44 158 Scale,[15] medical history and current medications, psychological distress using the Kessler-6
45 159 scale,[16] and lifestyle-related variables (tobacco, alcohol, physical activity, and dietary habits).
46 160 The results of the measurements were explained by local nurses and given to each participant.
47 161 Those who had extremely abnormal results were encouraged to visit the nearest health facilities
48 162 with the reports.
49
50
51
52
53
54
55
56
57
58
59
60

164 **Patient and public involvement**

165 Participants were not involved in the design, conduct, reporting, and dissemination plans of our
166 research.

168 **Statistical analysis**

169 We analysed the data using the Complex Sample module in IBM SPSS Statistics version 21
170 (IBM Corp., Armonk, NY, USA) to adjust for the effects of multistage sampling, clustering,
171 and weighting. Sample weights accounted for different selection probabilities at each sampling
172 stage, non-response rate in each SEA, and post-stratification adjustments to correct for
173 differences between our sample and the district population estimates based on the 2010 census.
174 Total weights were standardised as the final weight. Bivariate analyses were performed to
175 determine statistically significant associations between independent variables and high BP
176 (systolic BP (SBP) ≥ 140 or diastolic BP (DBP) ≥ 90 mmHg) using logistic regression. Variables
177 that showed significant associations with high BP ($p < 0.10$) in the bivariate analysis were
178 entered into the multiple logistic regression models stratified by gender.

180 **Ethical considerations**

181 This study was approved by the Ethics Committee of the Graduate School and Faculty of
182 Medicine of Kyoto University, Japan (R0403) and ERES Converge, Zambia (No. 2016-Jan-
183 003) for the pilot phase. The University of Zambia Biomedical Research Ethics Committee,
184 Zambia (No. 011-02-16) and the National Health Research Authority, Zambia (MH/101/23/10-
185 1) granted approval for the main survey. All participants provided written informed consent
186 prior to their participation.

188 **RESULTS**

189 Of the 800 targeted subjects, 712 agreed to participate. We excluded 22 participants from the
190 analyses due to missing interviews or anthropometric/biological data. The final valid response
191 rate was 86.3%. Table 1 shows the weighted characteristics of the study population by gender.
192 The proportion of men was 48.6%, and the mean age was 41.9 years (Standard error [SE] 0.6).
193 Most of the participants were married (80.8%), had only primary education (74.3%), and were
194 self-employed (69.7%). Nearly one-half had a monthly income of 50 USD or less (Zambia's
195 minimum wage), and one-quarter were living with severe food insecurity. For the self-reported

1
2 196 medical history, 10.4% had human immunodeficiency virus (HIV) and were receiving
3 197 antiretroviral therapy. Only 8% and 0.7% of participants had been diagnosed with hypertension
4 and diabetes, respectively. More than 50% of both men and women had family members or
5 198 relatives who had hypertension, and about 20% reported having family members or relatives
6 who had experienced a stroke.
7 199
8 200

10 201
11 202
12 203

13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

204 **Table 1. Sociodemographic characteristics and related medical histories among overall participants**
 205 **in the Mumbwa district, Central Province of Zambia, 2016**
 206

	Overall n (%)	Male n (%)	Female n (%)
Number			
Unweighted	690	332	358
Weighted	689 (100)	335 (48.6)	354 (51.4)
Age, years [SE]	41.9 [0.6]	42.7 [0.8]	41.1 [0.7]
Residential area of the district			
Urban area	87 (10.4)	35 (14.7)	52 (12.6)
Rural area	602 (89.6)	300 (85.3)	302 (87.4)
Marital Status			
Not married	27 (3.9)	20 (5.8)	7 (2.0)
Married	557 (80.8)	300 (89.4)	257 (72.6)
Divorced/widow/widower	106 (15.4)	16 (4.8)	90 (25.4)
Education			
Primary	513 (74.3)	229 (68.3)	284 (80.1)
Secondary	127 (18.5)	77 (23.0)	50 (14.2)
Tertiary	49 (7.2)	29 (8.7)	20 (5.7)
Monthly income (USD)			
≤50	326 (47.4)	157 (46.9)	169 (47.8)
>50	362 (52.6)	178 (53.1)	185 (52.2)
Work Status			
Employed	85 (12.4)	58 (17.2)	28 (7.8)
Self-employed	481 (69.7)	255 (76.2)	225 (63.6)
Unemployed/Retired	123 (17.9)	22 (6.6)	101 (28.6)
Food security			
Secure	192 (27.9)	107 (32.1)	85 (23.9)
Mildly insecure	45 (6.6)	25 (7.4)	21 (5.8)
Moderately insecure	261 (37.9)	132 (39.4)	129 (36.4)
Severely insecure	191 (27.7)	71 (21.2)	120 (33.8)
Medical history (Self-reported)			
HIV positive*	71 (10.4)	28 (8.4)	43 (12.2)
Hypertension	55 (8.0)	18 (5.4)	37 (10.4)
Diabetes	5 (0.7)	3 (0.9)	2 (0.6)
Past history within family and relatives (Self-reported)			
Hypertension	381 (55.3)	174 (52.0)	207 (58.5)
Stroke	140 (20.3)	68 (20.4)	71 (20.2)
Heart disease	64 (9.3)	25 (7.5)	39 (11.0)
Diabetes	123 (17.9)	52 (15.5)	71 (20.2)

Data are numbers (%)

SE, standard error

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

*All have been receiving antiretroviral treatment.

207
 208
 209
 210
 211
 212
 213 Table 2 shows the prevalence of hypertension in each stage and the current status of access to
 214 health services for hypertension stratified by gender. The prevalence of hypertension (Stage 2
 215 and hypertensive crisis) was 36.6% in the overall sample and was greater in men than in women
 216 but without statistical significance (39.7% vs. 33.5%, $p=0.10$). In contrast, the prevalence of
 217 hypertensive crisis, which refers to severe BP elevation, was slightly higher in women than in
 218 men (5.1% vs. 3.2%) ($p=0.32$). Prehypertension (SBP, 120–139 or DBP, 80–89 mmHg
 219 [Elevated and Stage 1]), which is the risk of developing future hypertension and cardiovascular
 220 disease, was found in 39.9% of the men and 30.6% of the women, and the difference was
 221 statistically significant ($p=0.02$). There was a significant association between the stage of
 222 hypertension and age in both men and women (men, $p=0.02$; women, $p<0.01$). Area of
 223 residence in the district had a significant association with hypertension in men ($p=0.02$) but not
 224 in women ($p=0.82$). Regarding access to healthcare services for hypertension, the prevalence

of hypertension was higher among men than among women, and the proportion of men who had “never had their BP measured” was significantly higher than that of women (37.3% [125/335] vs. 6.8% [24/354]; $p<0.01$). The proportion of participants who were previously diagnosed with hypertension was 5.4% [18/335] for men and 10.5% [37/354] for women, and 2.4% [8/335] and 4.5% [16/354] received antihypertensive treatment, respectively ($p=0.09$, $p=0.25$).

231

232

Table 2. Stage of hypertension relative to demographics and access to care and services among all participants in the Mumbwa district, Central Province of Zambia, 2016

233
234
235
236

Male												
	Overall	Normal		Prehypertension				Hypertension		p value		
				Elevated	Stage 1		Stage 2	Hypertensive crisis				
Total n(%)	335	68	(20.4)	33	(9.8)	101	(30.1)	122	(36.5)	11	(3.2)	
Age												
25-44	203	45	(22.0)	26	(12.7)	62	(30.6)	68	(33.4)	3	(1.3)	0.02
45-64	132	24	(17.9)	7	(5.4)	39	(29.2)	54	(41.2)	8	(6.2)	
Residential area of the district												0.02
Urban area	35	5	(13.2)	1	(2.9)	8	(22.1)	17	(48.5)	5	(13.2)	
Rural area	300	64	(21.2)	32	(10.6)	93	(31.0)	105	(35.1)	6	(2.1)	
Access to hypertension care and services												<0.01
Have never blood pressure measured	125	27	(21.3)	8	(6.1)	35	(27.9)	55	(43.9)	1	(0.8)	
Diagnosed as hypertensive	18	2	(11.4)	0	(0.0)	2	(11.4)	8	(45.7)	6	(31.4)	
On treatment	8	0	(0.0)	0	(0.0)	1	(13.3)	2	(26.7)	5	(60.0)	
Medical history (Self-reported)												0.69
HIV positive	28	8	(29.1)	2	(7.3)	9	(30.9)	9	(32.7)	0	(0.0)	
Diabetic	3	0	(0.0)	0	(0.0)	1	(33.3)	2	(66.7)	0	(0.0)	

237
238

Female												
	Overall	Normal		Prehypertension				Hypertension		p value		
				Elevated	Stage 1		Stage 2	Hypertensive crisis				
Total n(%)	354	127	(35.8)	12	(3.3)	97	(27.3)	101	(28.4)	18	(5.1)	
Age												<0.01
25-44	226	105	(46.4)	5	(2.3)	63	(27.7)	50	(22.0)	4	(1.6)	
45-64	128	22	(17.3)	7	(5.2)	34	(26.5)	51	(39.8)	14	(11.2)	
Residential area of the district												0.82
Urban area	52	16	(31.7)	2	(3.0)	17	(32.7)	15	(28.7)	2	(4.0)	
Rural area	302	111	(36.6)	10	(3.4)	80	(26.4)	86	(28.4)	16	(5.3)	
Access to hypertension care and services												0.00
Have never blood pressure measured	24	8	(31.9)	1	(2.1)	9	(38.3)	7	(27.7)	0	(0.0)	
Diagnosed as hypertensive	37	2	(4.2)	0	(0.0)	7	(19.4)	15	(40.3)	13	(36.1)	
On treatment	16	0	(0.0)	0	(0.0)	3	(18.8)	6	(34.4)	8	(46.9)	
Medical history (Self-reported)												0.36
HIV positive	43	19	(44.0)	3	(6.0)	11	(26.2)	10	(23.8)	0	(0.0)	
Diabetic	2	1	(50.0)	0	(0.0)	0	(0.0)	1	(50.0)	0	(0.0)	

239

240

241

242

243

244

245

246

247

248

249

250

251

252

Data are number (%)
HIV, human immunodeficiency virus
Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.
Blood pressure category: Normal (SBP<120 and DBP<80), Elevated (SBP 120-129 and DBP < 80), Stage 1 (SBP 130-139 or DBP 80-89), Stage 2 (SBP≥140 or DBP≥90), Hypertensive crisis (SBP>180 and/or DBP>120)

The present status of hypertension screening and diagnosis is shown in Figure 1. Among the residents, 21.8% (150/689) never had their BP measured, and the main reasons given were ‘do not know where to obtain the service’ (41.6%), ‘do not have the time or opportunity to check’ (24.8%), and ‘I think it is not important or I am healthy’ (18.8%). Among the participants who never had their BP measured, 41.9% (63/150) presented with hypertension at the time of measurement in this study. Among participants who had their BP measured previously, 89.8% (485/539) had not been diagnosed with hypertension but 30.3% (147/485) presented with

1
2 253 hypertension. Among the participants already diagnosed with hypertension, 56.4% (31/55)
3 254 were not using antihypertensive medication, of which 71.0% (22/31) presented with
4 255 hypertension. Furthermore, most participants using antihypertensive medication (20/24)
5 256 presented with hypertension, indicating poor BP control. Among individuals with HIV-positive
6 257 status (N=71), the distribution of hypertension and its stages showed no difference from the
7 258 overall distribution, but subgroup analysis showed that the proportion of individuals who had
8 259 never had their BP measured was lower than the overall proportion for both men and women
9 260 (Supplementary Table 1).

15 261
16 262 Table 3 shows the prevalence of hypertension in relation to each covariate and the association
17 263 of each covariate with hypertension by multivariable analysis (adjusted for the variables with
18 264 $p < 0.10$ in the bivariate analysis) in the overall sample analysis and the analysis stratified by
19 265 gender (Supplementary Table 2). In the overall sample, older age group (45–64 years) (adjusted
20 266 odds ratio [AOR] = 1.95, 95% confidence interval [CI], 1.35–2.80), higher education
21 267 (\geq college) (2.00, 95%CI, 1.04–3.82), alcohol intake (a few times/week or everyday) (2.14,
22 268 95%CI, 1.28–3.58), and body mass index (BMI) ≥ 25 kg/m² (1.83, 95%CI, 1.24–2.71) were
23 269 positively associated, while HIV-positive status was negatively associated with hypertension
24 270 (0.53, 95%CI, 0.29–0.96). Gender, marital status, food insecurity, smoking, physical activity,
25 271 cooking oil intake, sugar intake, and HbA_{1c} were not associated with hypertension. There was
26 272 a significant association between hypertension and alcohol intake in both genders (men \geq a few
27 273 times/week or everyday: 2.28, 95%CI, 1.24–4.17; women \leq a few times/month: 1.79, 95%CI,
28 274 1.01–3.19), but the association with urban residence was significant only in men (2.46, 95%CI,
29 275 1.09–5.56). Older age (45–64 years) (2.68, 95%CI, 1.56–4.63), higher education (\geq college)
30 276 (3.39, 95%CI, 1.19–9.64), low-level alcohol intake (\leq a few times/month) (1.79, 95%CI, 1.01–
31 277 3.19), and BMI ≥ 25 kg/m² (1.98, 95%CI, 1.18–3.29) showed significant association with
32 278 hypertension only in women.

33 279
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Multivariate correlates of hypertension among all participants in the Mumbwa district, Central Province of Zambia, 2016

	Overall				Male				Female			
	Total	Hypertension n (%)	Adjusted OR (95%CI)	p value	Total	Hypertension n (%)	Adjusted OR (95%CI)	p value	Total	Hypertension n(%)	Adjusted OR (95%CI)	p value
Socio-demographic characteristics												
Gender												
Male	335	133 (39.7)	–		–	–	N/A		–	–	N/A	
Female	354	119 (33.5)	–		–	–			–	–		
Age												
25-44	429	124 (28.9)	1 (Reference)		203	70 (34.7)	1 (Reference)		226	53 (23.6)	1 (Reference)	
45-64	260	128 (49.2)	1.95 (1.35-2.80)	0.00	132	63 (47.5)	1.57 (0.95-2.58)	0.07	128	65 (51.0)	2.68 (1.56-4.63)	0.00
Marital Status												
Not married	27	8 (30.8)	1 (Reference)		20	7 (34.2)	–		7	2 (21.4)	–	
Married	557	200 (36.0)	1.57 (0.63-3.90)	0.33	300	120 (40.0)	–		257	81 (31.4)	–	
Divorced/Widow/Widowed	106	43 (40.8)	1.83 (0.68-4.97)	0.23	16	7 (41.9)	–		90	36 (40.6)	–	
Education												
≤primary	513	185 (36.1)	1 (Reference)		229	90 (39.6)	–		284	95 (33.3)	1 (Reference)	
Secondary	127	44 (34.7)	1.03 (0.66-1.61)	0.89	77	32 (41.3)	–		50	12 (24.5)	0.68 (0.32-1.45)	0.32
≥college	49	23 (45.8)	2.00 (1.04-3.82)	0.04	29	11 (36.8)	–		20	12 (59.0)	3.39 (1.19-9.64)	0.02
Work Status												
Employed	85	29 (33.7)	–		58	19 (33.0)	1 (Reference)		28	10 (35.2)	–	
Self-employed	481	178 (37.1)	–		255	102 (39.8)	1.43 (0.74-2.78)	0.29	225	77 (34.0)	–	
Unemployed/Retired	123	45 (36.3)	–		22	12 (55.8)	1.95 (0.66-5.75)	0.22	101	32 (32.0)	–	
Residential area of the district												
Urban area	87	39 (44.4)	–		35	22 (61.8)	2.46 (1.09-5.56)	0.03	52	17 (32.7)	–	
Rural area	602	213 (35.4)	–		300	112 (37.2)	1 (Reference)		302	102 (33.7)	–	
Food security												
Secure	192	58 (30.2)	1 (Reference)		107	33 (31.1)	–		85	25 (29.1)	–	
Insecure	497	194 (39.0)	1.46 (0.99-2.16)	0.06	228	100 (43.8)	–		269	94 (34.9)	–	
Family planning												
Not used	–	–	–		–	–	–		249	96 (38.6)	1 (Reference)	
Used	–	–	–		–	–	–		101	96 (21.8)	0.61 (0.32-1.153)	0.13
No data	–	–	–		–	–	–		4	1 (14.3)	–	
Behavioral and psychological characteristics												
Smoking												
Never	525	179 (34.1)	1 (Reference)		185	66 (35.8)	–		340	113 (33.1)	–	
Ex-smoker	86	37 (43.1)	1.26 (0.73-2.18)	0.41	74	33 (44.4)	–		12	4 (34.8)	–	
Current smoker	79	36 (45.8)	1.27 (0.71-2.29)	0.42	76	34 (44.6)	–		3	2 (80.0)	–	
Alcohol												
Never	354	106 (30.0)	1 (Reference)		102	30 (29.3)	1 (Reference)		252	77 (30.3)	1 (Reference)	
≤a few times/month	206	80 (39.0)	1.49 (0.98-2.27)	0.06	123	49 (39.6)	1.43 (0.78-2.59)	0.24	82	31 (38.1)	1.79 (1.01-3.19)	0.046
≥a few times/week or everyday	130	65 (50.4)	2.14 (1.28-3.58)	0.00	110	54 (49.5)	2.28 (1.24-4.17)	0.01	20	11 (55.3)	1.61 (0.58-4.49)	0.37
Physical activity (activities of daily life and sports ≥once a week)												
Neither	54	25 (46.2)	1 (Reference)		16	8 (46.9)	–		38	17 (45.9)	1 (Reference)	
Either	497	182 (36.6)	0.62 (0.34-1.15)	0.13	215	96 (44.7)	–		282	86 (30.4)	0.51 (0.24-1.08)	0.08
Both	138	45 (32.5)	0.66 (0.32-1.35)	0.25	104	29 (28.2)	–		34	15 (45.5)	1.60 (0.54-4.71)	0.40
Cooking oil intake												
Low <20.83ml	510	198 (38.8)	1 (Reference)		244	103 (42.2)	–		266	95 (35.7)	–	
High ≥20.83ml	177	54 (30.5)	0.71 (0.48-1.07)	0.10	92	30 (33.1)	–		85	24 (27.7)	–	
Don't know	1	0 (0.0)	–		–	–	–		1	0 (0.0)	–	
No data	2	0 (0.0)	–		–	–	–		2	0 (0.0)	–	
Sugar intake												

6/bmjopen-2021-006668 on 8 April 2022. Downloaded from http://bmjopen.bmj.com/ on April 19, 2024 by guest. Protected by copyright.

	Low <28.0g	353	136	(38.6)	1	(Reference)		168	70	(42.0)	–		185	66	(35.6)	–		
	High ≥28.0g	335	114	(33.9)	0.79	(0.56-1.12)	0.18	167	62	(37.0)	–		168	52	(30.9)	–		
	No data	2	2	(100.0)	–			1	1	(100.0)	–		1	1	(100.0)	–		
	Clinical characteristics																	
	Body mass index (kg/m ²)																	
	Normal (<25)	505	167	(33.1)	1	(Reference)		281	106	(37.9)	1	(Reference)	224	61	(27.1)	1	(Reference)	
	Overweight/Obese (≥25)	185	85	(46.0)	1.83	(1.24-2.71)	0.00	54	27	(49.1)	1.73	(0.91-3.29)	130	58	(44.7)	1.98	(1.18-3.29)	0.01
	HbA1c																	
	Normal (<5.7)	404	144	(35.6)	1	(Reference)		222	89	(40.1)	–		182	55	(30.1)	1	(Reference)	
	High risk (5.7-6.4)	266	96	(36.1)	1.05	(0.73-1.50)	0.80	111	42	(38.1)	–		156	54	(34.7)	1.02	(0.61-1.71)	0.93
	Diabetes (≥6.5)	18	12	(65.7)	2.69	(0.91-7.96)	0.07	3	2	(66.7)	–		15	10	(65.5)	2.33	(0.68-7.96)	0.18
	No data	1	0	(0.00)	–			0	0	(0.0)	–		1	0	(0.0)	–		
	Medical history (self-reported)																	
	HIV status (Self-reported)																	
	Negative	618	232	(37.6)	1	(Reference)		307	124	(40.4)	–		311	108	(34.9)	1	(Reference)	
	Positive	71	20	(27.3)	0.53	(0.29-0.96)	0.04	28	9.25	(32.7)	–		43	10	(23.8)	0.54	(0.24-1.24)	0.15

Data are number (%)

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

"No data" and "Don't know" were excluded from statistical test.

Hypertension is defined as SBP 140 ≥ mmHg or DBP ≥90 mmHg.

OR: Odds ratio

288 Table 4 shows factors associated with ‘never had BP measured’ among men, as 83.8%
 289 (125/150) of participants who never had their BP measured were men. In the multivariable
 290 analysis, older age (0.43, 95%CI, 0.25–0.73) and HIV positive status (0.37, 95%CI, 0.14–0.97)
 291 were negatively associated, while being a current smoker was positively associated with ‘never
 292 had BP measured’ (2.09, 95%CI, 1.19–3.66). In contrast, in women, though not shown in the
 293 table, older age was positively associated with ‘never had BP measured’ (4.53, 95%CI, 1.81–
 294 11.4) (Supplementary Table 3,4).

295

296

297 **Table 4. Bivariate and multivariate correlates of "never had blood pressure measured" (Men only)**

298

	Male (n=335)	Never had blood pressure measured (n=125)						
	Total	n of total (%)	Crude OR (95%CI)		p value	Adjusted OR (95%CI)		p value
Age								
25-44	203	83 (41.0)	1	(reference)		1	(reference)	
45-64	132	42 (31.9)	0.48	(0.30-0.77)	0.00	0.43	(0.25-0.73)	0.00
Residential area of the district								
Urban area	35	6 (16.2)	1	(reference)		1	(reference)	
Rural area	300	120 (39.9)	3.60	(1.35-9.61)	0.01	2.79	(0.98-7.93)	0.06
Education								
Primary	229	96 (42.0)	1	(reference)		1	(reference)	
≥Secondary	106	29 (27.5)	0.62	(0.38-1.01)	0.05	0.84	(0.48-1.45)	0.53
Work Status								
Employed	58	13 (23.2)	1	(reference)		1	(reference)	
Unemployed/Retired	278	112 (40.4)	2.04	(1.08-3.83)	0.03	1.86	(0.92-3.76)	0.09
HIV infection								
No	307	119 (38.7)	1	(reference)		1	(reference)	
Yes	28	7 (23.6)	0.40	(0.16-1.02)	0.06	0.37	(0.14-0.97)	0.04
Smoking								
Never, Ex-smoker	259	87 (33.7)	1	(reference)		1	(reference)	
Current smoker	76	38 (50.0)	2.01	(1.19-3.38)	0.01	2.09	(1.19-3.66)	0.01
Alcohol								
Never or a few times/month	225	85 (37.9)	1	(reference)		—		
≥a few times/week or everyday	110	40 (36.4)	1.03	(0.64-1.66)	0.91	—		
Body mass index (kg/m²)^c								
Normal (<25)	281	111 (39.6)	1	(reference)		1	(reference)	
Overweight/Obese (≥25)	54	14 (26.4)	0.43	(0.22-0.85)	0.02	0.66	(0.32-1.40)	0.28

Data are number (%).

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

OR: Odds ratio

299

300

301

302

303

304

DISCUSSION

In this study, we assessed the prevalence and the risk factors for hypertension by gender to understand the current situation of hypertension among rural residents in Zambia. We also explored the status of screening and diagnosis of hypertension and its correlates to evaluate the situation of access to healthcare services for hypertension.

We found that more than 35% of the participants had hypertension, and the profile of hypertension correlates was different between men and women. More than 80% of the people with high BP measurements had never been previously diagnosed with hypertension, and 30% of them had never had their BP measured, suggesting the lack of access to or availability of healthcare services for BP control.

The prevalence of hypertension in rural areas

The prevalence of hypertension among the targeted rural residents of this study in 2016 was 39.7% in men and 33.5% in women, both being much higher than the national averages found in the Zambia STEPS Survey of 2017 (20.5% and 17.6%, respectively).[7] Previous research has reported mixed findings regarding the prevalence of hypertension in rural areas of Zambia. While a similar rate (46.9%) was reported among people attending health check-ups in other rural areas of Zambia,[17] 23.1% was reported in a primary healthcare-based study conducted in several rural districts between 2011–2014.[18] However, comparing our results with those of previous studies is difficult due to methodological differences. For example, previous studies used convenient sampling with potential selection bias, while we conducted probability sampling of the whole area. Studies using probability sampling are required for documenting the accurate BP status among Zambian rural populations. The prevalence rates of prehypertension and hypertension were slightly higher in men than in women in our study, a tendency that has been observed throughout the African region.[6]

Gender differences in factors associated with hypertension

In this study, a gender difference was found not only in the prevalence of hypertension, but also in the profile of the correlates of hypertension. In men, residence in the urban area of the district and high frequency of alcohol intake were significantly associated with hypertension. While in women, older age, higher education level, low frequency of alcohol intake, and BMI ≥ 25 kg/m² were associated with hypertension, suggesting the different mechanism(s) involved in the development of high BP between the genders. This implies that different pathways for

1
2 338 hypertension including behavioural and socio-cultural factors exist between men and women,
3 339 which could affect prevention strategies [19]

4 340 Alcohol consumption was the only factor moderately associated with hypertension in both
5 341 genders, which is in line with well-established findings worldwide.[20] Although the exact
6 342 mechanism is unclear, it can be caused directly through the chronic effect of alcohol and/or
7 343 indirectly through related socioeconomic status and lifestyles among the study population.[20]
8 344 Regardless of the mechanism, it is important to follow the trend of alcohol intake over time
9 345 with special attention to the type, amount, and pattern since it may rapidly change in both
10 346 quantity and quality with future economic growth.

11 347 Living in the urban area of the district was significantly associated with hypertension only
12 348 in men. Although the study region was “rural” in general (neighbouring the capital city,
13 349 Lusaka), there are some areas with relatively easy access to the capital city. Men living in such
14 350 areas may be involved in urbanized lifestyles, probably in relation to their jobs, in terms of
15 351 eating habits and lifestyles, including high calorie diets and lack of exercise. Studies in
16 352 Cameroon and Mali have shown a similar tendency with higher prevalence of hypertension
17 353 among men in “urban areas” than in rural areas.[21,22]

18 354 The relationship between age and hypertension has been reported in SSA countries.[23-25]
19 355 In our study, a significant association with age was observed only in women, reflecting the age-
20 356 related distribution of hypertension between the genders, where the difference in prevalence
21 357 between younger (25–44 years) and older (45–64 years) age groups was large (23.6% vs. 51.0%,
22 358 respectively) in women, but small in men (34.7% and 47.4%, respectively). Similar age
23 359 disparities in the prevalence of hypertension by gender have been reported in previous studies
24 360 of Zambia and Senegal.[18,25] This may suggest that men are more likely to develop
25 361 hypertension at a younger age than women. The reasons for this age disparity by gender should
26 362 be one of the focus points in future research.

27 363 An association between hypertension and education level was observed only among women.
28 364 Slightly high odds of hypertension in people with higher levels of education were also observed
29 365 in a study in Malawi [21]. This may suggest that in SSA countries that experienced rapid
30 366 economic growth in recent years, the risk of hypertension has increased among people with
31 367 higher levels of education due to spread of urbanized eating habits and lifestyles (over-nutrition
32 368 and physical inactivity).[26] The reason why the association was detected only in women in
33 369 our study is unclear; however, higher education may be related to urbanized eating habits and
34 370 lifestyles more in women than in men.

35 371 The association of overweight and obesity ($BMI \geq 25 \text{ kg/m}^2$) with hypertension has been
36 372 reported in SSA countries including Zambia, with its tendency being stronger in women than

1
2 373 in men.[27] Similarly in our study, although the association was observed in both genders, it
3 374 was significant only in women. This may be related to biological factors such as an increase in
4 375 obesity with age in women in African societies and their cultural preferences. In men,
5 376 behavioural factors such as alcohol consumption and psychological stress are more likely to be
6 377 associated with developing hypertension than obesity.
7
8
9

10 378

11 379 **Status of Hypertension Management**

12 380 In this study, only 16.7% of the participants who presented with hypertension had previously
13 381 been diagnosed. Among the participants with documented hypertension but no previous
14 382 diagnosis, 30% never had their BP measured. Our results concur with findings from a
15 383 systematic review of hypertension in SSA indicating that only 22.5% of people with
16 384 hypertension had already been diagnosed.[18] This indicates the need to strengthen
17 385 hypertension screening and diagnosis, particularly at the primary healthcare level which is the
18 386 entry level to health care systems in most SSA countries.

19 387 Moreover, only 8% of the participants in this study reported having been previously
20 388 diagnosed with hypertension, which was much lower than the actual proportion presenting with
21 389 hypertension. In addition, only fewer than half of the participants diagnosed with hypertension
22 390 were using antihypertensive medications, and many of them presented with hypertension at the
23 391 time of the measurement, indicating challenges in accessing treatment for hypertension. A
24 392 previous study in Zambia reported that 18% of people who presented with hypertension at the
25 393 time of the study had been prescribed antihypertensive medication at a health centre.[18] In
26 394 our study, only 7.9% of the participants with hypertension had been prescribed
27 395 antihypertensive medication. Furthermore, about 83% of the participants who reported using
28 396 antihypertensive medication in our study presented with hypertension at the time of
29 397 measurement. This was consistent with the results of a previous study in Zambia where nearly
30 398 90% had poorly controlled hypertension,[18] and other reports from the entire SSA region.[2]
31 399 These results indicate that there are various challenges in the management of hypertension in
32 400 the rural areas of Zambia, as in other SSAs, in terms of ‘difficulties in accessing appropriate
33 401 treatment and health services including hypertension’, ‘lack of screening and diagnostic
34 402 opportunities for hypertension’, and ‘lack of awareness of the importance of BP control’.

35 403

36 404 **Access to healthcare services related to hypertension**

37 405 We also assessed the differences in access to healthcare services related to hypertension
38 406 between the genders. Identifying the management status of hypertension (care cascade) can

1
2 407 contribute to health policy and interventions.[28] We focused on the history of BP
3 408 measurement as it relates to the awareness of hypertension status. In our study, more than 20%
4
5 409 of the participants had never had their BP measured previously, suggesting the difficulties in
6
7 410 accessing screening and diagnostic services for hypertension care. Despite the higher
8
9 411 prevalence of hypertension among men than among women, the proportion of men who 'never
10
11 412 had their BP measured' was 37.4%, which was 5.5 times higher than that of women. Men also
12
13 413 tended to be less likely to have been diagnosed with and treated for hypertension.

14 414 There was a significant positive association between smoking and 'never had their BP
15
16 415 measured' in men. While this finding requires further assessment in future, it may suggest that
17
18 416 people who engage in high-risk health behaviours such as smoking tend to be less concerned
19
20 417 about their health and less likely to engage in health seeking behaviours than those who do not
21
22 418 engage in such behaviour. In this study, we also included self-reported HIV status in the
23
24 419 analysis as a factor affecting access to healthcare services. Men in older age groups and men
25
26 420 with HIV-positive status were less likely to have 'never had their BP measured', suggesting
27
28 421 that they were likely to be aware of their BP. The association with 'older age group' may be
29
30 422 because they were likely to receive medical care during their lifetime. Regarding the
31
32 423 association with HIV-positive status, all HIV-positive individuals were receiving HIV
33
34 424 treatment, so regular medical consultations at a healthcare facility may have provided the
35
36 425 opportunity for BP measurement.

37 426 Men have fewer opportunities to access healthcare services besides illness or injury,
38
39 427 compared to women who visit for maternal and child health services. Patients with
40
41 428 asymptomatic conditions like hypertension may not receive the required healthcare services
42
43 429 due to psychological and geographical barriers, e.g., low level of attention to health or distance
44
45 430 to healthcare facilities. In Zambia, access to quality essential healthcare services remains
46
47 431 limited due to weak health systems including workforce shortage. For instance, the proportion
48
49 432 of medical doctors per 10,000 population was 0.93 in 2016 and the universal health coverage
50
51 433 service coverage index in 2017 was lower than the global average.[29] Therefore, along with
52
53 434 strengthening hypertension screening, we suggest that the use of existing mobile health services,
54
55 435 such as vaccination campaigns, mobile voluntary counselling and testing services, and
56
57 436 cooperation with community health workers may be advantageous in treating many people.[30]
58
59 437 For women, although only a few never had their BP measured, the odds of never having BP
60
438 measured were significantly higher in the older age group. This gender difference should be
439
440 examined in further research with a large sample size.

441 **Strengths and limitations**

1
2 442 The strength of our study is that we used multi-stage cluster random sampling and obtained a
3 443 relatively high response rate. Thus, our results are representative of patients at risk of CVD in
4 444 the target population in the rural area. Regarding limitations, the recorded BP may have been
5 445 higher than usual due to white coat hypertension. A previous study that used the same
6 446 hypertension criteria as our study reported that the prevalence of white coat hypertension (false-
7 447 positive) was 13%, masked hypertension (false-negative) 14%, and correctly classified
8 448 hypertension 73%.[31] Therefore, data on the prevalence of hypertension in this study should
9 449 be interpreted cautiously. Socially desirable responses due to face-to-face interviews could also
10 450 have affected the results, even though we trained the interviewers before the study.
11 451 Unmeasured factors may have affected some of the associations found in our study.
12
13
14
15
16
17
18
19
20

21 453 **CONCLUSION**

22
23 454 More than one-third of the participants in a rural district in Zambia had hypertension; most
24 455 were not diagnosed yet and one-quarter of them never had their BP measured. These results
25 456 indicate a lack of CVD prevention services, including access to and availability of healthcare
26 457 services for hypertension, among rural residents in Zambia. Therefore, health promotion and
27 458 screening strategies for hypertension are urgently required, especially in primary healthcare
28 459 settings in rural areas. Particular attention should be paid to healthcare access, specifically
29 460 among men.
30
31
32
33
34
35
36

37 462 **Acknowledgments**

38
39 463 We are thankful to all the participants, the staff at the Institute of Economic and Social Research,
40 464 the University of Zambia, and the members of the Mumbwa District Medical Office for the
41 465 local arrangement and support to this study. We gratefully acknowledge the local nurses and
42 466 research assistants for their assistance with the data collection. Lastly, we extend our gratitude
43 467 to the late Mubiana Macwan'gi for contributing to the supervision of the research in Zambia.
44
45
46
47
48
49

50 469 **Contributors**

51
52 470 YT, TT, MOK and MK contributed to study conception and design. YT, RZ and CD
53 471 contributed to the data collection. YT, TT and MK contributed to data analysis and drafted the
54 472 manuscript. YT, PMM, SPS, AO, RZ and CD revised the manuscript. MOK and MK
55 473 supervised the study. All authors read and approved the final manuscript.
56
57
58
59
60

475 **Funding**

476 This study was supported by the United Nations University, Institute for the Advanced Study
477 of Sustainability (Global Leadership Training Program in Africa) and the Inter-Graduate
478 School Program for Sustainable Development and Survivable Societies, Kyoto University. It
479 was also supported by the 2016 Kyoto University School of Public Health Super Global
480 Course's travel scholarship.

482 **Competing interests**

483 None declared.

485 **Patient consent for publication**

486 Not required.

488 **Data sharing statement**

489 Data are available upon reasonable request.


491 **References**

- 492 1. World Health Organization. Fact sheets. Cardiovascular diseases (CVDs).
493 [https://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
494 (accessed May 2021).
- 495 2. Ataklte F, Erqou S, Kaptoge S, et al. Burden of undiagnosed hypertension in sub-Saharan
496 Africa: a systematic review and meta-analysis. *Hypertension*. 2015;65:291–98.
497 doi:10.1161/HYPERTENSIONAHA.114.04394
- 498 3. Dzudie A, Rayner B, Ojji D, et al. Roadmap to Achieve 25% Hypertension Control in
499 Africa by 2025. *Glob Heart*. 2018;13:45–59. doi: 10.1016/j.gheart.2017.06.001
- 500 4. Boateng D, Wekesah F, Browne JL, et al. Knowledge and awareness of and perception
501 towards cardiovascular disease risk in sub-Saharan Africa: A systematic review. *PLoS*
502 *One*. 2017; 12:e0189264. doi: 10.1371/journal.pone.0189264.
- 503 5. Berry KM, Parker WA, Mchiza ZJ, et al. Quantifying unmet need for hypertension care
504 in South Africa through a care cascade: evidence from the SANHANES, 2011-2012.
505 *BMJ Glob Health*. 2017; 16;2:e000348. doi: 10.1136/bmjgh-2017-000348

- 1
2 506 6. World Health Organization. Global status report on noncommunicable diseases 2014.
3 507 https://apps.who.int/iris/bitstream/handle/10665/148114/9789241564854_eng.pdf
4 508 (accessed May 2021).
5
6
7 509 7. World Health Organization. Zambia STEPS Survey 2017 Factsheet
8 510 <https://www.afro.who.int/publications/zambia-steps-survey-2017-factsheet> (accessed
9 511 May 2021).
10
11
12 512 8. Yan, LD, Chi BH, Sindano N, et al. Prevalence of hypertension and its treatment among
13 513 adults presenting to primary health clinics in rural Zambia: analysis of an observational
14 514 database. *BMC Public Health* 2015;15:933. doi:10.1186/s12889-015-2258-4
15
16 515 9. Mulenga D, Siziya S, Rudatsikira E, et al. District specific correlates for hypertension in
17 516 Kaoma and Kasama rural districts of Zambia. *Rural Remote Health*. 2013;13:2345.
18
19 517 10. Yan LD, Chi BH, Sindano N, et al. Prevalence of hypertension and its treatment among
20 518 adults presenting to primary health clinics in rural Zambia: analysis of an observational
21 519 database. *BMC Public Health*. 2015;15:933. doi: 10.1186/s12889-015-2258-4
22
23 520 11. Central Statistical Office Zambia. 2010 Census of Population and Housing—Volume 11
24 521 National Descriptive Tables. 2012.
25 522 <http://catalog.ihnsn.org/index.php/catalog/4124/download/55313> (accessed May 2021).
26
27 523 12. World Health Organization. WHO STEPS Surveillance Manual. 2017.
28 524 http://www.who.int/ncds/surveillance/steps/STEPS_Manual.pdf?ua=1 (accessed May
29 525 2021).
30
31 526 13. BeLue R, Okoror TA, Iwelunmor J, et al. An overview of cardiovascular risk factor
32 527 burden in sub-Saharan African countries: a socio-cultural perspective. *Glob Health*
33 528 2009;5:10. doi: 10.1186/1744-8603-5-10
34
35 529 14. Tateyama Y, Musumari PM, Techasrivichien T, et al. Dietary habits, body image, and
36 530 health service access related to cardiovascular diseases in rural Zambia: A qualitative
37 531 study. *PLoS One*. 2019; 14:e0212739. doi: 10.1371/journal.pone.0212739
38
39 532 15. Coates J, Anne S, Paula B. Household Food Insecurity Access Scale (HFIAS) for
40 533 Measurement of Household Food Access: Indicator Guide (v. 3). 2007. Available from:
41 534 http://www.fao.org/fileadmin/user_upload/eufao-fsi4dm/doc-training/hfias.pdf (accessed
42 535 May 2021).
43
44 536 16. Kessler RC, Green JG, Gruber MJ, et al. Screening for serious mental illness in the
45 537 general population with the K6 screening scale: results from the WHO World Mental
46 538 Health (WMH) survey initiative. *Int J Methods Psychiatr Res*. 2010;19 Suppl 1:4–22.
47 539 doi:10.1002/mpr.310
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2 540 17. Rush KL, Goma FM, Barker JA, et al. Hypertension prevalence and risk factors in rural
3 541 and urban Zambian adults in western province: a cross-sectional study. *Pan Afr Med J*.
4 542 2018;30:97. doi: 10.11604/pamj.2018.30.97.14717
5
6
7 543 18. Yan LD, Chi BH, Sindano N, et al. Prevalence of hypertension and its treatment among
8 544 adults presenting to primary health clinics in rural Zambia: analysis of an observational
9 545 database. *BMC Public Health*. 2015;15:933. doi:10.1186/s12889-015-2258-4
10
11
12 546 19. Ghosh S, Mukhopadhyay S, Barik A. Sex differences in the risk profile of hypertension:
13 547 a cross-sectional study. *BMJ Open*. 2016;6:e010085. doi: 10.1136/bmjopen-2015-
14 548 010085.
15
16
17 549 20. Beilin LJ, Puddey IB. Alcohol and hypertension: an update. *Hypertension*.
18 550 2006;47:1035–38. doi:10.1161/01.HYP.0000218586.21932.3c
19
20
21 551 21. Cooper R, Rotimi C, Ataman S, et al. The prevalence of hypertension in seven
22 552 populations of west African origin. *Am J Public Health*. 1997;87:160–8.
23 553 doi:10.2105/ajph.87.2.160
24
25
26 554 22. Bâ HO, Camara Y, Menta I, et al. Hypertension and Associated Factors in Rural and
27 555 Urban Areas Mali: Data from the STEP 2013 Survey. *Int J Hypertens*.
28 556 2018;2018:6959165. doi:10.1155/2018/6959165
29
30
31 557 23. Berry KM, Parker WA, Mchiza ZJ, et al. Quantifying unmet need for hypertension care
32 558 in South Africa through a care cascade: evidence from the SANHANES, 2011-2012.
33 559 *BMJ Glob Health*. 2017;2:e000348. doi:10.1136/bmjgh-2017-000348
34
35
36 560 24. Price AJ, Crampin AC, Amberbir A, et al. Prevalence of obesity, hypertension, and
37 561 diabetes, and cascade of care in sub-Saharan Africa: a cross-sectional, population-based
38 562 study in rural and urban Malawi. *Lancet Diabetes Endocrinol*. 2018;6:208–22.
39 563 doi:10.1016/S2213-8587(17)30432-1
40
41
42
43 564 25. Kwarisiima D, Balzer L, Heller D, et al. Population-Based Assessment of Hypertension
44 565 Epidemiology and Risk Factors among HIV-Positive and General Populations in Rural
45 566 Uganda. *PLoS One*. 2016;11:e0156309. doi:10.1371/journal.pone.0156309
46
47
48 567 26. Macia E, Gueye L, Duboz P. Hypertension and Obesity in Dakar, Senegal. *PLoS One*.
49 568 2016;11:e0161544. doi:10.1371/journal.pone.0161544
50
51
52 569 27. Akpa OM, Made F, Ojo A, et al. Regional Patterns and Association Between Obesity and
53 570 Hypertension in Africa: Evidence from the H3Africa CHAIR Study. *Hypertension*.
54 571 2020;75:1167–78. doi:10.1161/HYPERTENSIONAHA.119.14147
55
56
57 572 28. Geldsetzer P, Manne-Goehler J, Marcus ME, et al. The state of hypertension care in 44
58 573 low-income and middle-income countries: a cross-sectional study of nationally
59
60

- 1
2 574 representative individual-level data from 1·1 million adults. *Lancet*. 2019;394:652–62.
3 575 doi:10.1016/S0140-6736(19)30955-9
4
5 576 29. World Health Organization. Global Health Observatory data repository.
6
7 577 <https://apps.who.int/gho/data/node.main.HWFGRP?lang=en> (accessed November 2021).
8
9 578 30. Opie LH, Seedat YK. Hypertension in sub-Saharan African populations. *Circulation*.
10 579 2005;112:3562–8. doi:10.1161/CIRCULATIONAHA.105.539569
11
12 580 31. Sheppard JP, Stevens R, Gill P, et al. Predicting Out-of-Office Blood Pressure in the
13 581 Clinic (PROOF-BP): Derivation and Validation of a Tool to Improve the Accuracy of
14 582 Blood Pressure Measurement in Clinical Practice. *Hypertension*. 2016;67(5):941-950.
15 583 doi:10.1161/HYPERTENSIONAHA.115.07108
16
17 584
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



2021-055668 on 8 April 2022. Downloaded from <http://bmjopen.bmj.com/> on April 19, 2024 by guest. Protected by copyright.

Supplementary Tables

Table 1. Access to hypertension care and services among People Living with HIV (N=71)

	Overall n (%)	Male n (%)	Female n (%)
Age			
25-44	49 (68.3)	15 (52.7)	34 (78.6)
45-64	23 (31.7)	13 (47.3)	9 (21.4)
Residential area of the district			
Urban area	3 (4.3)	0 (0.0)	3 (7.1)
Rural area	68 (95.7)	28 (100.0)	40 (92.9)
Blood pressure			
High (hypertension)	20 (27.3)	9 (32.7)	10 (23.8)
Access to hypertension care and services			
Have never blood pressure measured	8 (11.5)	7 (23.6)	2 (3.6)
Diagnosed as hypertensive	5 (7.2)	4 (12.7)	2 (3.6)
On treatment	0 (0.0)	0 (0.0)	0 (0.0)

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer. Hypertension is defined as SBP ≥ 140 mmHg or DBP ≥ 90 mmHg.

Table 2. Bivariate and Multivariate correlates of hypertension among all participants in the Mumbwa district, Central Province of Zambia, 2016 (by gender)

	Male						Female					
	Hypertension						Hypertension					
	n	n(%)	Crude OR (95%CI)	p value	Adjusted OR (95%CI)	p value	n	n(%)	Crude OR (95%CI)	p value	Adjusted OR (95%CI)	p value
Socio-demographic characteristics												
Age												
25-44	203	70 (34.7)	1 (Reference)		1 (Reference)		226	53 (23.6)	1 (Reference)		1 (Reference)	
45-64	132	63 (47.5)	1.05 (1.250-3.133)	0.004	1.57 (0.95-2.58)	0.07	128	65 (51.0)	3.54 (2.236-5.616)	0.000	2.68 (1.56-4.63)	0.00
Marital Status												
Not married	20	7 (34.2)	1 (Reference)		–		7	2 (21.4)	1 (Reference)		–	
Married	300	120 (40.0)	1.78 (0.628-5.025)	0.278	–		257	81 (31.4)	2.01 (0.424-9.34)	0.379	–	
Divorced/Widow/Widowed	16	7 (41.9)	1.75 (0.442-6.928)	0.425	–		90	36 (40.6)	3.34 (0.686-16.232)	0.136	–	
Education												
≤primary	229	90 (39.6)	1 (Reference)		–		284	95 (33.3)	1 (Reference)		1 (Reference)	
Secondary	77	32 (41.3)	1.09 (0.644-1.860)	0.739	–		50	12 (24.5)	0.83 (0.426-1.617)	0.583	0.68 (0.32-1.45)	0.32
≥college	29	11 (36.8)	1.03 (0.469-2.283)	0.932	–		20	12 (59.0)	3.85 (1.488-9.74)	0.005	3.39 (1.19-9.64)	0.02
Work Status												
Employed	58	19 (33.0)	1 (Reference)		1 (Reference)		28	10 (35.2)	1 (Reference)		–	
Self-employed	255	102 (39.8)	1.34 (0.726-2.456)	0.352	1.43 (0.74-2.78)	0.29	225	77 (34.0)	0.84 (0.388-1.920)	0.659	–	
Unemployed/Retired	22	12 (55.8)	2.80 (1.025-7.646)	0.045	1.95 (0.66-5.75)	0.22	101	32 (32.0)	0.69 (0.299-1.96)	0.387	–	
Monthly income (USD)												
≤50	157	61 (38.9)	1 (Reference)		–		169	51 (30.1)	1 (Reference)		–	
>50	178	72 (40.5)	1.02 (0.652-1.599)	0.927	–		185	68 (36.7)	1.29 (0.832-2.011)	0.252	–	
Residential area												
Rural	300	112 (37.2)	1 (Reference)		1 (Reference)		302	102 (33.7)	1 (Reference)		–	
Urban	35	22 (61.8)	2.46 (1.176-5.144)	0.017	2.46 (1.09-5.56)	0.03	52	17 (32.7)	0.98 (0.528-1.802)	0.937	–	
Food security												
Secure	107	33 (31.1)	1 (Reference)		–		85	25 (29.1)	1 (Reference)		–	
Insecure	228	100 (43.8)	1.47 (0.901-2.399)	0.122	–		269	94 (34.9)	1.39 (0.822-2.353)	0.220	–	
Family planning												
Not used	–	–	–		–		249	96 (38.6)	1 (Reference)		1 (Reference)	
Used	–	–	–		–		101	96 (21.8)	0.42 (0.248-0.724)	0.002	0.61 (0.32-1.153)	0.13
No data	–	–	–		–		4	1 (14.3)	–		–	
Behavioral and psychological characteristic												
Smoking												
Never	185	66 (35.8)	1 (Reference)		–		340	113 (33.1)	1 (Reference)		–	
Ex-smoker	74	33 (44.4)	1.42 (0.816-2.463)	0.216	–		12	4 (34.8)	0.74 (0.192-2.931)	0.657	–	
Current smoker	76	34 (44.6)	1.42 (0.889-2.724)	0.113	–		3	2 (80.0)	3.93 (0.353-4.805)	0.266	–	
Alcohol												
Never	102	30 (29.3)	1 (Reference)		1 (Reference)		252	77 (30.3)	1 (Reference)		1 (Reference)	
≤a few times/m	123	49 (39.6)	1.66 (0.938-2.937)	0.082	1.43 (0.78-2.59)	0.24	82	31 (38.1)	1.40 (0.844-2.320)	0.193	1.79 (1.01-3.19)	0.046
≥a few times/w or everyday	110	54 (49.5)	2.37 (1.323-4.248)	0.004	2.28 (1.24-4.17)	0.01	20	11 (55.3)	2.49 (1.014-6.10)	0.046	1.61 (0.58-4.49)	0.37
Fruit and vegetable intake (fruits ≥once a week, vegetables daily)												
Neither	50	20 (40.2)	1 (Reference)		–		51	14 (27.0)	1 (Reference)		–	
Either	159	66 (41.3)	1.06 (0.548-2.015)	0.862	–		160	53 (33.3)	1.18 (0.570-2.422)	0.661	–	
Both	126	47 (37.6)	1.15 (0.585-2.253)	0.689	–		142	51 (36.1)	1.44 (0.697-2.975)	0.324	–	
Physical activity (activities of daily life and sports ≥once a week)												
Neither	16	8 (46.9)	1 (Reference)		–		38	17 (45.9)	1 (Reference)		1 (Reference)	
Either	215	96 (44.7)	0.99 (0.361-2.696)	0.980	–		282	86 (30.4)	0.52 (0.266-1.032)	0.062	0.51 (0.24-1.08)	0.08
Both	104	29 (28.2)	0.53 (0.185-1.527)	0.240	–		34	15 (45.5)	0.91 (0.354-2.324)	0.840	1.60 (0.54-4.71)	0.40
Salt intake (per day)												
<5g	228	89 (39.2)	1 (Reference)		–		295	102 (34.5)	1 (Reference)		–	
≥5g	90	33 (36.4)	0.82 (0.495-1.365)	0.449	–		58	16 (27.4)	0.63 (0.347-1.156)	0.137	–	

1	Don't know	16	11	(65.6)	–	–	–	–	–	1	1	(100.0)	–	–	–	–		
2	Cooking oil intake																	
3	Low <20.83ml	244	103	(42.2)	1	(Reference)	–	–	–	266	95	(35.7)	1	(Reference)	–	–		
4	High ≥20.83ml	92	30	(33.1)	0.72	(0.429-1.218)	0.223	–	–	85	24	(27.7)	0.71	(0.420-1.003)	0.204	–		
5	Don't know	–	–	–	–	–	–	–	–	1	0	(0.0)	–	–	–	–		
6	No data	–	–	–	–	–	–	–	–	2	0	(0.0)	–	–	–	–		
7	Sugar intake																	
8	Low <28.0g	168	70	(42.0)	1	(Reference)	–	–	–	185	66	(35.6)	1	(Reference)	–	–		
9	High ≥28.0g	167	62	(37.0)	0.73	(0.464-1.139)	0.164	–	–	168	52	(30.9)	0.73	(0.471-1.040)	0.168	–		
10	No data	1	1	(100.0)	–	–	–	–	–	1	1	(100.0)	–	–	–	–		
11	Psychological distress (K6)																	
12	Low	166	58	(35.0)	1	(Reference)	–	–	–	169	77	(45.3)	1	(Reference)	–	–		
13	High (10 and over)	169	56	(32.8)	0.73	(0.463-1.137)	0.162	–	–	184	94	(51.1)	0.97	(0.627-1.408)	0.901	–		
14	Clinical characteristics																	
15	Body mass index (kg/m ²) ^c																	
16	Normal (<25)	281	106	(37.9)	1	(Reference)	–	–	–	224	61	(27.1)	1	(Reference)	–	1 (Reference)		
17	Overweight/Obese (≥25)	54	27	(49.1)	2.04	(1.13-3.70)	0.02	1.73	(0.91-3.29)	130	58	(44.7)	2.51	(1.59-3.91)	0.00	1.98	(1.18-3.29)	0.01
18	Blood lipids																	
19	Normal	70	27	(39.0)	1	(Reference)	–	–	–	42	9	(22.2)	1	(Reference)	–	–		
20	Abnormal	263	106	(40.3)	1.09	(0.630-1.887)	0.757	–	–	310	109	(35.2)	1.56	(0.758-3.220)	0.227	–		
21	No data	3	0	(0.0)	–	–	–	–	–	3	1	(20.0)	–	–	–	–		
22	HbA1c																	
23	Normal (<5.7)	222	89	(40.1)	1	(Reference)	–	–	–	182	55	(30.1)	1	(Reference)	–	1 (Reference)		
24	High risk (5.7-6.4)	111	42	(38.1)	0.81	(0.502-1.320)	0.404	–	–	156	54	(34.7)	1.38	(0.879-2.179)	0.161	1.02	(0.61-1.71)	0.93
25	Diabetes (≥6.5)	3	2	(66.7)	3.30	(0.295-36.973)	0.333	–	–	15	10	(65.5)	4.87	(1.588-15.921)	0.006	2.33	(0.68-7.96)	0.18
26	No data	0	0	(0.0)	–	–	–	–	–	1	0	(0.0)	–	–	–	–		
27	Medical history (self-reported)																	
28	HIV infection (Self-reported)																	
29	No	307	124	(40.4)	1	(Reference)	–	–	–	311	108	(34.9)	1	(Reference)	–	1 (Reference)		
30	Yes	28	9	(32.7)	0.77	(0.338-1.743)	0.527	–	–	43	10	(23.8)	0.47	(0.226-0.983)	0.045	0.54	(0.24-1.24)	0.15

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

"No data" and "Don't know" excludes from statistical test

Hypertension is defined as SBP 140 ≥ mmHg or DBP ≥90 mmHg.

Abnormal blood lipid includes any abnormal measurements in total cholesterol, triglyceride, LDL-cholesterol, or HDL-cholesterol

Cooking oil intake is defined as low and high by median (20.83ml).

Sugar intake is defined as low and high by median (28.0g).

Table 3: Bivariate and multivariate correlates of ‘never had blood pressure measured’ (Women)

	Female (n=354)	Never had blood pressure measured (n=24)				
	Total	n of total (%)	Crude OR (95CI)	p value	Adjusted OR (95CI)	p value
Age						
25-44	226	6 (2.5)	1 (reference)		1 (reference)	
45-64	128	19 (14.5)	5.33 (2.18-13.06)	0.00	4.53 (1.81-11.35)	0.00
Residential area						
Urban	52	4 (6.9)	1 (reference)		—	
Rural	302	21 (6.8)	1.39 (0.40-4.81)	0.60	—	
Education						
<=primary	284	19 (6.7)	1 (reference)		—	
>=Secondary	70	5 (7.3)	0.997 (0.36-2.75)	0.995	—	
Work Status						
Employed	28	3 (9.3)	1 (reference)		—	
Unemployed/Retired	326	22 (6.6)	0.71 (0.20-2.50)	0.59	—	
HIV infection						
No	311	23 (7.3)	1 (reference)		—	
Yes	43	2 (3.6)	0.83 (0.24-2.89)	0.77	—	
Smoking						
Never, Ex-smoker	352	24 (6.7)	1 (reference)		—	
Current smoker	3	1 (20.0)	6.60 (0.58-75.32)	0.13	—	
Alcohol						
Never or a few times/m	335	20 (6.0)	1 (reference)		1 (reference)	
≥a few times/w or everyday	20	4 (21.1)	3.37 (1.04-10.88)	0.04	2.19 (0.65-7.43)	0.21
Body mass index (kg/m²)						
Normal (<25)	224	11 (4.8)	1 (reference)		1 (reference)	
Overweight/Obese (25 and over)	130	13 (10.3)	2.12 (0.95-4.72)	0.07	1.71 (0.74-3.92)	0.21

Data are number (%).

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

OR: Odds ratio

Table 4: Bivariate and multivariate correlates of ‘never had blood pressure measured’ (Overall)

	Overall (n=689)	Never had blood pressure measured (n=150)				
	Total	n of total (%)	Crude OR (95CI)	p value	Adjusted OR (95CI)	p value
Gender						
Male	335	125 (37.4)	7.71 (4.88-12.18)	0.00	6.27 (3.84-10.23)	0.00
Female	354	24 (6.8)	1 (reference)		1 (reference)	
Age						
25-44	429	89 (20.7)	1 (reference)		—	
45-64	260	61 (23.3)	0.90 (0.61-1.31)	0.57	—	
Residential area						
Urban	87	9 (10.7)	1 (reference)		1 (reference)	
Rural	602	140 (23.3)	3.02 (1.43-6.41)	0.00	2.70 (1.22-5.98)	0.01
Education						
<=primary	513	115 (22.5)	1 (reference)		—	
>=Secondary	177	34 (19.5)	0.96 (0.63-1.45)	0.84	—	
Work Status						
Employed	85	16 (18.7)	1 (reference)		—	
Unemployed/Retired	604	134 (22.1)	1.16 (0.67-2.01)	0.60	—	
HIV infection						
No	618	141 (22.9)	1 (reference)		1 (reference)	
Yes	71	8 (11.5)	0.44 (0.21-0.90)	0.03	0.46 (0.21-0.995)	0.049
Smoking						
Never, Ex-smoker	611	111 (18.2)	1 (reference)		1 (reference)	
Current smoker	79	39 (49.0)	4.46 (2.74-7.28)	0.00	2.11 (1.19-3.73)	0.01
Alcohol						
Never or a few times/m	560	105 (18.8)	1 (reference)		1 (reference)	
≥a few times/w or everyday	130	44 (34.1)	2.29 (1.50-3.50)	0.00	0.93 (0.55-1.55)	0.77
Body mass index (kg/m²)						
Normal (<25)	505	122 (24.1)	1 (reference)		1 (reference)	
Overweight/Obese (25 and over)	185	28 (15.0)	0.50 (0.31-0.79)	0.00	0.91 (0.55-1.53)	0.73

Data are number (%).

Totals of percentages may differ from 100 due to rounding. Weighted values are rounded to the nearest integer.

OR: Odds ratio

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	3
Objectives	3	State specific objectives, including any prespecified hypotheses	3
Methods			
Study design	4	Present key elements of study design early in the paper	4
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	4
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	4
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5
Bias	9	Describe any efforts to address potential sources of bias	5
Study size	10	Explain how the study size was arrived at	4
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	5
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	5
		(b) Describe any methods used to examine subgroups and interactions	5
		(c) Explain how missing data were addressed	6
		(d) If applicable, describe analytical methods taking account of sampling strategy	4
		(e) Describe any sensitivity analyses	6
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6
		(b) Give reasons for non-participation at each stage	6
		(c) Consider use of a flow diagram	NA
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	6,7
		(b) Indicate number of participants with missing data for each variable of interest	NA
Outcome data	15*	Report numbers of outcome events or summary measures	7,8

1			
2	Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included
3			11,12,13
4			
5			
6			(b) Report category boundaries when continuous variables were categorized
7			9,12
8			
9			(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period
10			NA
11	Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses
12			13
13			
14	Discussion		
15	Key results	18	Summarise key results with reference to study objectives
16	Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
17			18
18			
19			
20	Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
21			18
22			
23			
24	Generalisability	21	Discuss the generalisability (external validity) of the study results
25			18
26	Other information		
27	Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based
28			19
29			
30			

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.