




BMJ Open A pilot multicentre cluster randomised trial to compare the effect of trauma life support training programmes on patient and provider outcomes

Martin Gerdin Wärnberg ^{1,2}, Johanna Berg ^{1,3}, Prashant Bhandarkar,^{4,5} Anirban Chatterjee,⁶ Shamita Chatterjee,⁷ Chintamani Chintamani,⁸ Li Felländer-Tsai,^{9,10} Anita Gadgil,⁵ Geeta Ghag,¹¹ Marie Hasselberg,¹ Catherine Juillard,¹² Monty Khajanchi,¹³ Deepa Kizhakke Veetil,¹⁴ Vineet Kumar,¹⁵ Debabrata Kundu,¹⁶ Anurag Mishra,¹⁷ Priti Patil,^{5,18} Nobhojit Roy ^{1,5}, Amit Roy,¹⁹ Siddarth David,^{1,20} Rajdeep Singh,¹⁷ Harris Solomon,²¹ Kapil Dev Soni,²² Lovisa Strömmer,^{23,24} Megha Tandon,⁸ Trauma life support training Effectiveness Research Network (TERN) collaborators

To cite: Gerdin Wärnberg M, Berg J, Bhandarkar P, *et al.* A pilot multicentre cluster randomised trial to compare the effect of trauma life support training programmes on patient and provider outcomes. *BMJ Open* 2022;**12**:e057504. doi:10.1136/bmjopen-2021-057504

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-057504>).

Received 17 September 2021
Accepted 28 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Martin Gerdin Wärnberg; martin.gerdin@ki.se

ABSTRACT

Introduction Trauma accounts for nearly 10% of the global burden of disease. Several trauma life support programmes aim to improve trauma outcomes. There is no evidence from controlled trials to show the effect of these programmes on patient outcomes. We describe the protocol of a pilot study that aims to assess the feasibility of conducting a cluster randomised controlled trial comparing advanced trauma life support (ATLS) and primary trauma care (PTC) with standard care.

Methods and analysis We will pilot a pragmatic three-armed parallel, cluster randomised controlled trial in India, where neither of these programmes are routinely taught. We will recruit tertiary hospitals and include trauma patients and residents managing these patients. Two hospitals will be randomised to ATLS, two to PTC and two to standard care. The primary outcome will be all-cause mortality at 30 days from the time of arrival to the emergency department. Our secondary outcomes will include patient, provider and process measures. All outcomes except time-to-event outcomes will be measured both as final values as well as change from baseline. We will compare outcomes in three combinations of trial arms: ATLS versus PTC, ATLS versus standard care and PTC versus standard care using absolute and relative differences along with associated CIs. We will conduct subgroup analyses across the clinical subgroups men, women, blunt multisystem trauma, penetrating trauma, shock, severe traumatic brain injury and elderly. In parallel to the pilot study, we will conduct community consultations to inform the planning of the full-scale trial.

Ethics and dissemination We will apply for ethics approvals to the local institutional review board in each hospital. The protocol will be published to Clinical Trials Registry—India and ClinicalTrials.gov. The results will be published and the anonymised data and code for analysis will be released publicly.

Strengths and limitations of this study

- Cluster randomised controlled trial comparing the effect of advanced trauma life support and primary trauma care and standard care on patient and provider outcomes.
- Prospective data collection with direct observations by dedicated project officers.
- Participating centres' heterogeneity may affect the study estimates and bias the results.

INTRODUCTION

Trauma, defined as the clinical entity composed of physical injury and the body's associated response, causes 4.5 millions deaths every year.¹ Almost 10% of the global burden of disease is due to trauma and trauma is the top contributor to the burden of disease in children and adults aged 10–49 years.²

Trauma care is time sensitive and early management of life-threatening or limb-threatening condition is crucial. Several trauma life support training programmes have been developed to improve the early management of patients as they arrive at hospital by providing a structured framework to assessment and treatment.^{3–5}

The proprietary advanced trauma life support (ATLS) is the most established trauma life support training programme and >1 million doctors in over 80 countries have been trained in the programme.⁶ Uptake in low-income and middle-income countries (LMIC) has been slow, potentially due to high costs.⁵

The free primary trauma care (PTC) programme is the most widely spread alternative programme. The goal of PTC is to improve trauma care in LMIC.⁷ Like ATLS, doctors in over 80 countries have been trained in PTC, and the programme has been endorsed by WHO, among other international organisations including several professional societies.⁷

Despite the widespread use of these training programmes, there are no controlled trials showing that they impact patient outcomes.³⁻⁵ But there is level 1 evidence that these programmes improve provider skills and practices,^{8,9} and observational data suggesting that they also improve patient outcomes.¹⁰

We will perform a pilot study that aims to assess the feasibility of conducting a cluster randomised controlled trial comparing ATLS and PTC with standard care. Recent methodological guidelines indicate that the design of efficient cluster randomised controlled trials requires data on probable or target effect sizes, proportion of participants with the outcome (if binary) and the intracluster correlation coefficient.¹¹ The objectives of this pilot study will be to:

- ▶ Estimate probable effect sizes on patient outcomes associated with ATLS and PTC compared with standard care, estimate the proportion of participants with the outcome (if binary) and estimate the intracluster correlation coefficient, as a basis for future sample size calculations.
- ▶ Assess the feasibility of recruiting participants and collecting data on primary and secondary outcomes, such as mortality, in-hospital complications, length of stay and quality of life.
- ▶ Assess how the effect sizes and directions of these effects of ATLS and PTC may differ across clinically important subgroups.

METHODS

Trial design

This study will pilot a pragmatic three-armed parallel, cluster randomised controlled trial, by the Trauma life support training Effectiveness Research Network (TERN, www.tern.network). There will be two intervention arms, ATLS and PTC training, and one control arm, standard care. We will collect data for 4 months in all three arms, first during a 1-month observation phase and then during a 3-month intervention phase (or continued observation in the control arm). This design will allow us to assess outcomes both as final values and as change from baseline. Our study is a pilot study because its objectives involve estimating quantities, such as the probable effect sizes, proportion of participants with the outcome (if binary) and the intracluster correlation coefficient, needed for the sample size calculations of a full-scale trial.¹¹ The full-scale trial will be planned regardless of the effect sizes identified in this pilot study. This pilot study will also establish how many participants that can be enrolled, as well as likely dropout rates, and the feasibility of collecting primary and secondary outcomes.

Study setting

We will conduct this pilot study in Indian tertiary hospitals, where neither ATLS, PTC nor any other trauma life support training programme is routinely taught. India is the world's second most populous country and has 20% of the world's trauma deaths. The trauma system is still developing, with limited prehospital care, and the in hospital trauma mortality as well as the proportion of preventable deaths remain high. Lack of standard trauma training for healthcare providers, limited hospital resources, inadequate processes of care, overcrowding emergency departments are some of the factors that contribute to the high mortality and morbidity. During recent years, efforts have been made to improve hospital trauma care, through capacity building for trained trauma care providers, augmenting facilities and developing care protocols within the hospitals. Our pilot study is planned to start during 2022.

Eligibility criteria for participants and clusters

There will be two groups of participants: patients and resident doctors.

Patient participants

Adults (aged 15 years or older) who present to the emergency department at participating hospitals with a history of trauma. History of trauma is here defined as having any of the external causes of morbidity and mortality listed in block V01-Y36, chapter XX of the International Classification of Disease version 10 (ICD-10) codebook as reason for presenting. We will explore intervention effects across the following clinical subgroups: men, women, blunt multisystem trauma, penetrating trauma, shock, severe traumatic brain injury and elderly, as defined by Hornor *et al.*¹² The consent form for patients are available in online supplemental material 1.

Resident doctor participants

Resident doctors doing their specialty training in surgery or emergency medicine, who manage trauma patients in the emergency department, and who are expected to remain in the participating hospitals for at least 1 year. To facilitate administration each clinical department is divided into units, which manages the outpatient department, emergency department, operating rooms, etc on different days each week. One or two, out of typically six, units' residents will be selected from each hospital. One unit consists of at least 3 faculty and 3–12 residents.

To be eligible, units should have a maximum of 25% of the doctors trained in either ATLS, PTC or similar training programmes before the start of the pilot (hospitals that have so far agreed to participate have no or single current residents trained in any programme). Those residents who have received training in the last 5 years will be considered as trained. The figure of 25% was decided through consensus in the research team, to balance feasibility and contamination of results. We will select the units by conducting a prior survey to ascertain

this criteria. Consent will be sought from the residents in each of the intervention groups before they undergo the ATLS or PTC training. The consent form for residents are available in online supplemental material 2. We will not ask for consent from residents at the units in the control hospitals as their practice will not be affected by this pilot study and we will not collect any personal identifiable data on them. This is in line with ethical regulations in the study setting.

Clusters

Indian tertiary care hospitals that admit 400–800 adult patients with trauma each year. We randomise on the cluster (hospital) level to avoid contamination between intervention and control arms. To be eligible for inclusion, hospitals have to provide the following services round the clock: operation theatres, X-ray, CT and ultrasound facilities, and blood bank. In addition, the baseline admission rate should be >35 adult patients with major trauma per month.

Interventions

In each intervention arm, one or two units', out of typically six, residents per hospital providing emergency care to trauma patients will be trained in either ATLS or PTC. For the purpose of this pilot study, we will target to train a minimum of 75% of residents in each unit. If residents drop out or change units after training but before data collection is completed, we will conduct additional training if needed to meet the 75% criterion. We will not train the units' faculty, as they are typically not involved in the initial management of trauma patients.

The ATLS training will be conducted in the nearest ATLS certified training centre in India according to the standard ATLS curriculum.⁶ The PTC training will be arranged in hospitals randomised to the PTC arm, according to the standard PTC curriculum.⁷ These courses will be conducted over a period of 2.5–3 days. The residents certified 'pass' will be considered as trained in respective courses.

The control group provides standard care with no intervention.

Modifications

Both ATLS and PTC are standard training programmes with fixed curricula.^{6,7} We will not modify the delivery or content of these programmes during this pilot.

Adherence

The intervention is the training in either ATLS or PTC. Participants are required to adhere to, that is, participate in, the training, to be eligible for passing. We will not consider adherence to training contents during care delivery as adherence to the trial intervention, but rather as a provider-level outcome.

Concomitant care

Baseline training

The care provided by all participating hospitals at baseline is based on the training curriculum formulated by The National Medical Council of India for postgraduation in general surgery.¹³ Regarding trauma, these guidelines state that the student should:

1. have knowledge about response to trauma; burns: causes, prevention and management; wounds of scalp and its management; recognition, diagnosis and monitoring of patients with head injury, Glasgow coma scale;
2. be able to provide and coordinate emergency resuscitative measures in acute surgical situations including trauma;
3. choose, perform and interpret appropriate imaging in trauma—ultrasound focused abdominal sonography in trauma (FAST);
4. undergo advanced trauma and cardiac life support course (certified) before appearing in final examination;
5. undergo clinical posting in emergency and trauma;
6. present or discuss cases of blunt abdominal trauma.

Although training in an ATLS course is part of the curriculum, it is optional and not doing this training does not result in failure to obtain postgraduation completion.

Standard of care

At most medical colleges in India trauma patients present to the emergency department, where they are assessed by a doctor and referred to the surgical bay for further management. In the surgical bay, a second-year or third-year general surgery resident sees all the major trauma and provide the initial care, including initiating treatment and investigations. This resident informs the consultant on call who is generally an Assistant Professor. Most procedures like intercostal drainage, open wound suturing, intubation, etc would be done in the surgical resuscitation area, by the surgical resident.

Compared with other settings where a trauma team approach is adopted, nurses and other healthcare professionals are involved to a limited extent during the initial management. Their roles include assisting during intubation and other bedside procedures, charting the vitals (not recording) and giving injections. They also accompany the resident during transfers of serious patients.

After completing the assessment and starting initial resuscitation, the resident decides to send the patient for imaging (X-rays/FAST/CT scan) or to the operation room in consultation with or after assessment by the on-call consultant. A portable X-ray and an ultrasound machine to conduct FAST may or may not be available in the surgical bay. The patients who are operated, managed conservatively, not intubated or with minor trauma will be sent to the surgical ward. Those who need increased monitoring or mechanical ventilation remain in the surgical bay or in the intensive care unit (ICU) depending on the availability of ICU beds. The further treatment continues in the respective

ward or ICU and patients are finally discharged from the ward.

Outcomes

Our pilot study include both participant and feasibility outcomes. Prior to deciding on these participant outcomes, we searched the Core Outcome Measures in Effectiveness Trials Initiative's database but were unable to identify appropriate core outcome sets for our populations of participants.

The primary participant outcome will be all-cause mortality within 30 days from the time of arrival to the emergency department. The primary outcome and most secondary outcome will be assessed and compared both as final values and as change from baseline. All outcomes that pertain to the individual participant level are detailed in online supplemental material 3. We decided to include a large number of outcomes, including some more exploratory, so that we can test their feasibility and relevance. We may remove secondary participant outcomes during the course of the pilot study, if they prove to be too difficult to collect. If we remove outcomes, we will document the reasons for doing so.

We will also assess the following feasibility outcomes, which pertain both to overall study population as well as to the individual cluster level:

- ▶ Recruitment rate. For both patients and residents, this will equal the proportion of participants enrolled, out of the total number of eligible participants, over the course of the pilot study.
- ▶ Lost to follow-up rate. This will apply only to patients and equal the proportion of patients that do not complete 30-day follow-up, out of all enrolled patients, over the course of the pilot study.
- ▶ Pass rate. This will apply only to residents in the intervention arms and equal the proportion of residents that pass the training programme, out of the total number of trained residents, over the course of the pilot study.
- ▶ Missing data rate. This will apply to each outcome and variable and equal the proportion of missing data, over the course of the pilot study.
- ▶ Differences in distributions of observed and extracted data. This will apply to each outcome and variable and will compare the distributions of data collected by observations versus extracted from hospital records. For quantitative variables, this will be the difference in means, SD, medians, IQRs and ranges. For qualitative variables, this will be the differences in absolute counts and percentages, across categories.

Participant timeline

Patients

Patients will be screened for eligibility as they arrive at the emergency department. Eligible patients will be approached in the emergency department to consent to follow-up, if they are conscious. If they are unconscious, a patient representative will be approached to consent to

follow-up. Once the patient is conscious, we will approach the patient to affirm the patient representative's consent. We will follow-up patients at discharge, at 24 hours after arrival at the emergency department, and at 30 days after arrival at the emergency department.

Residents

Surgical units will be screened for eligibility once hospitals confirm their participation. All residents in eligible units will be approached to consent to training if their hospital is randomised to either of the intervention arms. Training will be conducted as soon as possible after the study starts. Resident participants will be followed up 30 days after training, if they are in the intervention arms, or 30 days after the study started, if they are in the control arm.

Sample size

Given budget and time constraints, including the rotation of units in Indian hospitals (which often happen on a 6-month basis), the feasible data collection period is 4 months. Each of the units see two to four trauma patients per week. If we select a minimum of one unit per hospital then each hospital will enrol 8–16 patients per month and 32–64 patients during the 4 months of this pilot study. With a 20% attrition rate we expect each hospital to enrol 26–51 patients, coming to a total sample size of between 156 and 306 patients for this pilot study.

Recruitment

To ensure adequate recruitment, we only approach hospitals with trauma volumes high enough to allow us to reach the sample size goals detailed above. Patients will be enrolled by a dedicated project officer as they arrive at the emergency department. The recruitment period will be 4 months. Recruitment will be monitored weekly through online conferences. No financial or non-financial incentives will be provided to trial investigators or participants for enrolment.

Allocation

Sequence generation

We will use simple randomisation to allocate sites to trial arms. We will prepare six sealed envelopes of which one representative from each pilot site will draw one. The content of the envelope will dictate what trial arm (ATLS, PTC or standard care) the hospital will be in. There will be two hospitals in each trial arm.

Concealment mechanism

We will not conceal the sequence, see 'Sequence generation' section.

Implementation

The random allocation sequence will be generated by the project's core group, who also enrol clusters. Patient participants will be included if they present during the project officers shift. Resident participants are enrolled if they are in the units selected for training. We will use

simple random sampling to select units if there are more than two eligible units in a hospital. For patient participants, consent for follow-up is sought after randomisation from patients or patient relatives as appropriate. For resident participants, consent is sought before randomisation. If residents in a unit decline to participate, so that the target of training 75% of residents in a given unit cannot be met, another unit will be selected for participation.

Blinding

It will not be possible to blind investigators or participants to interventions. We will not blind the data analysts during this pilot, but we plan to blind the data analysts during the full-scale trial.

Data collection

Data collection will start 1 month before the training is delivered, to establish a baseline. A variability of 3 months of the date when data collection is started between hospitals will be accepted. Each participating hospital will have a dedicated project officer to collect data. The project officers will have a masters in a health science field and should have experience in data collection.

Because participating residents are assigned designated days for trauma care for a period of 6 months, data will be collected during those particular days and shifts when these trained doctors are in the emergency department. The project officers will collect data both by observing the care delivered and by interviewing the participants, and by extracting data from hospital records.

Data collection will continue for a minimum of 3 months after training. The research officers will collect data of all patients, who present with trauma in the surgical bay during their duty hours. Those patients who are admitted will be followed up for complications and other in-hospital outcome measures, for example, length of stay. Patients who are not admitted will be followed up telephonically for mortality outcomes and quality of life outcomes. The follow-up period will be 30 days. The project officers will make at most three attempts to reach a participant or participant representative telephonically, after which the data will be recorded as missing.

The project officer will administer the study information and informed consent (consent will only be sought for data collection including follow-up) to the patient, or the patient's representative as appropriate, once the patient is stabilised. They will continue to collect data once they have received the consent.

Details of data of those patients/relatives not willing to give consent will be removed from the analysis. The number of patients who opt out from data collection will be collected, as well as limited data on their age and sex. Patients will be followed up in the ward regularly for the various outcome variables. They will also be followed up telephonically after they have been discharged.

Variables

The project officers will collect data on demographics, time of injury to arrival at the participating hospital, time to recording vital signs, vital signs and times to and management details including imaging and surgery. Details of any injury sustained will be collected and coded using ICD-10 and the Abbreviated Injury Scale (AIS). For ICD-10, coders will undergo the WHO online ICD-10 training module and for AIS they will be accredited. Based on AIS, we will calculate the Injury Severity Score (ISS) and the New ISS. Online supplemental material 4 contains a full variable list, with definitions.

Patient and public involvement

In this study, we will conduct community consultations to collect inputs from patients, their caregivers, patient groups and resident doctors to be used in the selection of outcome measures and implementation of the full-scale trial, following the Guidance for Reporting Involvement of Patients and the Public 2.¹⁴

During the pilot study, interviews will be conducted with postdischarge trauma patients and their caregivers to identify outcomes most relevant to them. These patients will be identified through the medical registers of the participating hospitals, contacted through telephone and after receiving their consent be interviewed as per their convenience. Their consent form is available as online supplemental material 5. Additionally, members from non-government organisations working with trauma patients and the hospital social service section will also be contacted for their views on contextual patient-centred outcomes for trauma patients. Their consent form is available as online supplemental material 6. For feasibility, these interviews will be held in each of the cities where the participating centres are located. The most common patient-centred outcomes reported across all the locations will be incorporated into the evaluation of the effects of the different training programmes and standardised care on patient outcomes.

Similarly, the inputs of resident doctor participants at each participating centre will be collected during the pilot study. A discussion and periodic surveys will be conducted to document any challenges or suggestions they may have in the scheduling or implementation of the training programmes. These inputs will be incorporated in the final study.

A summary of the findings of the study as well as their inputs will be shared with those who participated in the interviews and surveys. A meeting will be held with the patient participants, at each city, where the changes in the measured patient-centred outcomes would be presented to them. Another meeting will be held with the resident doctors at each hospital to present the confidence of the residents after being trained. Any suggestions and reflections from the participants during the meetings will be used as inputs for planning the final study.



Data management

We will supply an online data collection tool, accessible only over a virtual private network, for each participating hospital to upload pseudonymised data to secure servers. Data validation techniques like restricted values or values of a specific range will be used to avoid ambiguous data entries and ensure the validity of the data. Ambiguous responses, data errors, if any, will be resolved after discussion with the core team during weekly meetings. An instruction manual or codebook for data variables will be prepared to ensure consistency in data entry. This manual will be referred to during the project data collection and variable descriptions are visible for each variable in the online data collection tool. Pseudonymised data will be stored at the centralised server. The data will be accessible by the project's principal investigator or by delegation of the project principal investigator only.

Data monitoring

Weekly meetings with the core team and project officers will take place and for this meeting a data status report will be automatically generated highlighting missing data and number of patients awaiting follow-up. Cluster-specific interim analysis will take place after 2 months. The results of this will be presented to the core team, this team will decide if the pilot should be terminated. Although we will not have formal termination criteria because of the short duration of the study, reasons not to continue could include that the collection of key variables, such as mortality outcomes, is unfeasible or that patients are not consenting to be included in the data collection. A data monitoring committee will not be used in the pilot study due to its limited scope.

Statistical methods

We will analyse all pilot data using descriptive statistics. Quantitative variables will be summarised as mean±SD, median, IQR and range. Qualitative variables will be presented as absolute numbers and percentages. Feasibility outcomes will be summarised both on the overall sample level as well as on the individual cluster level. We will use an empty generalised linear mixed model to estimate the intracluster correlation coefficient.

We will compare participant outcomes in three combinations of trial arms: ATLS versus PTC, ATLS versus standard care and PTC versus standard care. In each combination, we will compare both differences in final values and differences in change from baseline. For example, for the primary participant outcome of all-cause mortality within 30 days from the time of arrival to the emergency department, comparing ATLS versus PTC, we will compare both the difference in mortality between the ATLS and PTC arms as well as the difference in the change from baseline in mortality between the ATLS and PTC arms.

For the intervention arms, the change from baseline will be calculated as the difference between the 1-month period of data collection before the training was

undertaken and the 3-month period after the training. For the control arm, the data collection period will be 4 months and the difference from baseline will be calculated as the difference between the first 1 month and the following 3 months.

Within each combination of trial arms, we will conduct subgroup analyses of men, women, blunt multisystem trauma, penetrating trauma, shock, severe traumatic brain injury and elderly. Table S7.1 in online supplemental material 7 shows which outcomes will be assessed in which subgroups, decided through consensus in the research team. We will further compare the results of all subgroups with the results in the whole cohort, and compare the results in the female subgroup with the male subgroup, and the results in the blunt multisystem trauma subgroup with the penetrating trauma subgroup. We are aware that the numbers in some of these subgroups are likely to be small, but we include them to help guide the formulation of the statistical analysis plan for the full-scale trial.

We will calculate both absolute and relative differences for each comparison, along with 75%, 85% and 95% CIs. We will use an empirical bootstrap procedure with 1000 draws to estimate these CIs. We will not perform any formal hypothesis tests during the analysis of this pilot's data.¹⁵ We will also compare the data collected through observations and interviews with the data collected from hospital records, to assess the feasibility of collecting data from hospital records in the full-scale trial.

ETHICS AND DISSEMINATION

We will apply for research ethics approval at local clusters in India to the local institutional review board committees. The protocol will be submitted for journal publication as well as to Clinical Trials Registry—India and ClinicalTrials.gov. Amendments to the protocol after this will be determined by the core research group and updated on Clinical Trials Registry—India and ClinicalTrials.gov. Substantial amendments, such as modifications to the eligibility criteria or outcomes will also be resubmitted to the journal. Declaration of interest will be submitted from all participating researchers both in the core team and at each site. The final anonymised dataset and code for analysis will be released publicly. The results will be submitted for publication in peer-reviewed open access journals. Authorship will follow the International Committee of Medical Journal Editors guidelines.

Author affiliations

¹Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden

²Function Perioperative Medicine and Intensive Care, Karolinska University Hospital, Stockholm, Sweden

³Emergency Medicine, Department of Internal and Emergency Medicine, Skåne University Hospital, Malmö, Sweden

⁴Tata Institute of Social Sciences School of Health Systems Studies, Deonar, Maharashtra, India

⁵World Health Organization Collaborating Center for Research in Surgical Care Delivery in Low-and-Middle Income Countries, Mumbai, India

⁶Department of Orthopaedic Sciences, Medica Superspecialty Hospital, Kolkata, India

⁷Department of Surgery, Seth Sukhlal Karnani Memorial Hospital, Kolkata, West Bengal, India

⁸Department of Surgery, Vardhman Mahavir Medical College and Safdarjung Hospital, New Delhi, Delhi, India

⁹Division of Orthopaedics and Biotechnology, Department of Clinical Science Intervention and Technology (CLINTEC), Karolinska Institutet, Stockholm, Sweden

¹⁰Department of Reconstructive Orthopedics, Karolinska University Hospital, Stockholm, Sweden

¹¹Department of Surgery, HBT Medical College and Dr R N Cooper Municipal General Hospital, Mumbai, India

¹²Division of General Surgery, Department of Surgery, David Geffen School of Medicine at UCLA, Los Angeles, UK

¹³Department of Surgery, Seth Gowardhandas Sunderdas Medical College and King Edward Memorial Hospital, Mumbai, India

¹⁴Department of Surgery, Manipal Hospital Dwarka, New Delhi, India

¹⁵Department of Surgery, Lokmanya Tilak Municipal Medical College and General Hospital, Mumbai, Maharashtra, India

¹⁶Department of Surgery, Medical College Kolkata, Kolkata, India

¹⁷Department of Surgery, Maulana Azad Medical College, New Delhi, Delhi, India

¹⁸Department of Statistics, Bhabha Atomic Research Centre Medical Division, Mumbai, Maharashtra, India

¹⁹Department of Surgery, Sir Nil Ratan Sircar Medical College & Hospital, Kolkata, India

²⁰Doctors For You, Mumbai, India

²¹Department of Cultural Anthropology and the Duke Global Health Institute, Duke University, Durham, North Carolina, USA

²²Critical Care, All India Institute of Medical Sciences, New Delhi, Delhi, India

²³Department of Surgery, Capio S:t Görans Hospital, Stockholm, Sweden

²⁴Division of Surgery, Department of Clinical Science, Intervention and Technology (CLINTEC), Karolinska Institutet, Stockholm, Sweden

Twitter Nobhohit Roy @#nobsroy

Contributors MGW conceived of the study. AG, AM, CJ, DKV, HS, JB, KDS, LF-T, LS, MH, MGW, MK, NR, PB, PP, RS, DS and VK contributed to the design of the study. DKV, KDS, MK and MGW drafted the first version of the protocol. AG, HS and DS drafted the first version of the patient and public involvement activities. JB and PP drafted the first versions of the data management sections and wrote the data management plan. PB and PP drafted the first versions of the statistical analysis section. AG, AM, CJ, DKV, HS, JB, KDS, LF-T, LS, MH, MGW, MK, NR, PB, PP, RS, SC, DS and VK contributed to the refinement of the protocol. AR, AC, C, DK, GG, MK, MT and VK are representatives of prospective participating hospitals.

Funding Doctors for You through grants awarded to Karolinska Institutet by the Swedish Research Council (grant number 2020-03779) and the Laerdal Foundation (grant number 2021-0048).

Competing interests Several authors are Advanced Trauma Life Support instructors.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those

of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Martin Gerdin Wärnberg <http://orcid.org/0000-0001-6069-4794>

Johanna Berg <http://orcid.org/0000-0001-7553-7337>

Nobhohit Roy <http://orcid.org/0000-0003-2022-7416>

REFERENCES

- Naghavi M, Abajobir AA, Abbafati C, *et al*. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the global burden of disease study 2016. *The Lancet* 2017;390:1151–210.
- Vos T, Lim SS, Abbafati C, *et al*. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the global burden of disease study 2019. *The Lancet* 2020;396:1204–22.
- Mohammad A, Branicki F, Abu-Zidan FM. Educational and clinical impact of advanced trauma life support (atls) courses: a systematic review. *World J Surg* 2014;38:322–9.
- Jayaraman S, Sethi D, Chinnock P, *et al*. Advanced trauma life support training for hospital staff. *Cochrane Database Syst Rev* 2014:CD004173.
- Kadhun M, Sinclair P, Lavy C. Are primary trauma care (PTC) courses beneficial in low- and middle-income countries - A systematic review. *Injury* 2020;51:136–41.
- American College of Surgeons. *Advanced trauma life support® student course manual*, 2018.
- Primary Trauma Care Foundation. *Primary trauma care course manual for trauma management in locations with limited resources*, 2018.
- Ali I, Cohen R, Reznick R. Demonstration of acquisition of trauma management skills by senior medical students completing the ATLS program. *J Trauma* 1995;38:687–91.
- Ali J, Cohen R, Adam R, *et al*. Teaching effectiveness of the advanced trauma life support program as demonstrated by an objective structured clinical examination for practicing physicians. *World J Surg* 1996;20:1121–6.
- Qin L-J, Shi X-P, Chang Y-X, *et al*. Systemic analysis of pre-hospital trauma emergency treatment in Zhengzhou. *J Acute Dis* 2019;8:34.
- Hemming K, Eldridge S, Forbes G, *et al*. How to design efficient cluster randomised trials. *BMJ* 2017;358:j3064.
- Hornor MA, Hoeft C, Nathens AB. Quality benchmarking in trauma: from the ntbd to tqip. *Current Trauma Reports* 2018;4:160–9.
- Medical Council of India. *Guidelines for competency based postgraduate training programme for MS in general surgery*, 2019.
- Staniszewska S, Brett J, Simeria I, *et al*. GRIPP2 reporting checklists: tools to improve reporting of patient and public involvement in research. *BMJ* 2017;358:j3453;j3453.
- Lee EC, Whitehead AL, Jacques RM, *et al*. The statistical interpretation of pilot trials: should significance thresholds be reconsidered? *BMC Med Res Methodol* 2014;14:41.

Supplemental Material 1

1.1. Study Enrolment: Patient Information Sheet

Patient Information Sheet

You are being invited to participate in a research study. Before your data can be included in this data bank the purpose of the data collection must be explained to you, and you must be given the chance to ask questions. Please read carefully the information provided here. If you agree to participate, please sign the informed consent form. You will be given a copy of this document to take home with you.

Protocol Title: Trauma Training Effectiveness Research Network

Principal Investigator: Name

PURPOSE OF THE RESEARCH STUDY

We are currently conducting research in this hospital to study the feasibility of assessing the effect of trauma life support training programs on care and outcome of patients with injury. We ask you to participate in this study because you presented to this hospital after having an injury.

STUDY PROCEDURES AND VISIT SCHEDULE

If you agree to participate, we will:

- Store health data registered in your hospital records and vital signs recordings from the emergency department
- Contact you in person or by telephone for follow ups to obtain information about your health status at the following times:
 - On arrival at the emergency department and wards till you are discharged
 - On hospital discharge
 - 24 hours after arrival to the hospital
 - 30 days after arrival to the hospital

When you arrived at the hospital, we recorded some basic parameters such as your age, gender, and how you were injured. We also recorded health data such as blood pressure, heart rate, oxygen levels, respiratory rate, surgical care and treatment provided. During your stay, we will record periodically health data, the investigations and treatment that you have undergone. During the follow-up calls, we may ask you details about your health and general information on returning back to your normal life and experience of the injury in your life. If you want complete information regarding all the parameters that were recorded, please do not hesitate to ask, and we will be happy to inform you.

Should you wish that your data is deleted from the study, you may please tell us now or contact us using the contact information provided below. The results of the study may be used for research that can be published as scientific articles; however, it will not be possible to identify you by reading any article that may result from this data bank. Further, data from this project will be combined with data from other hospitals that use the same system and shared for other researchers and individuals to use, but it will not be possible to identify you using that data. Research on the data without identifiers may seek to answer other questions than those stated above.

WITHDRAWAL FROM STUDY

Participation in this study is completely voluntary. Even if you agree to participate now you are free to withdraw at any time without giving any reason for doing so. Withdrawing will not affect your ordinary treatment or the care given to you. To withdraw contact any of the contact persons using the contact information provided. Note that we can only delete data from the data collected in this hospital. We cannot delete data once it has been de-identified because we will not be able to tell from whom the data came.

POSSIBLE RISKS, DISCOMFORTS AND INCONVENIENCES

We have not been able to identify any major risks associated with participating in this study. Even if you agree to participate now you are free to withdraw at any time without giving any reason for doing so. You are free to withdraw at that point, or at any time using the contact information provided below.

POTENTIAL BENEFITS

This research may help to improve the care of injured patients. Although this study will not affect the care you are given in this hospital at this time, your participation will contribute to medical knowledge, and may improve care for you if you are injured again in the future care for others that are injured.

SUBJECT'S RIGHTS

Your participation in this study is entirely voluntary. Your questions will be answered clearly and to your satisfaction. In the event of any new information becoming available that may be relevant to your willingness to continue in this study, you will be informed in a timely manner by the Principal Investigator or his/her representative.

CONFIDENTIALITY OF STUDY AND MEDICAL RECORDS

The results of this research may be published as a scientific article; however, it will not be possible to identify you by reading any article that may result from this work. Further, data from this project will be combined with data from other hospitals that use the same system and shared for other researchers and individuals to use, but it will not be possible to identify you using that data.

Also, Regulatory Agencies, Institutional Review Board and Ministry of Health will be granted direct access to your original medical records to check study procedures and data, without

making any of your information public. By signing the Informed Consent Form attached, you are authorizing such access to your study and medical records.

COSTS OF PARTICIPATION

No charge will be levied on you if you take part in this study. You will not receive any compensation for participating in this study.

RESEARCH RELATED INJURY AND COMPENSATION

Due to the observational nature of this project, it is unlikely to cause any research related injury. The hospital will provide medical care for any problems that may arise during this study.

WHOM TO CONTACT IF YOU HAVE QUESTIONS

If you have questions regarding this study and your rights, or in the case of any injuries sustained during this project, you may contact the Principal Investigator:

Name

Designation, Department

Phone Number

Email

1.2 Study Enrolment: Patient Consent Form

Consent Form**Protocol Title: Trauma Training Effectiveness Research Network****Patient Details**

Name: _____ NRIC/PNR/SSN No.: _____

Address: _____

Date of birth _____ Phone No: _____
dd/mm/yyyy

Phone number(s) of your relatives or friends that you agree we may contact, in case you do not answer your phone:

Part I - to be filled by the patient

I, _____ (Pt ID No. _____) **agree / do not agree** to participate in the project as described and, on the terms detailed in the Patient Information Sheet. The nature of my participation in the proposed project has been explained to me in _____ by Dr/Mr/Ms _____. I have fully discussed and understood the purpose and procedures of this project. I have been given the Patient Information Sheet and the opportunity to ask questions about this project and have received satisfactory answers and information. I understand that my participation is voluntary and that I am free to withdraw at any time, without giving any reasons and without my medical care being affected. I also give permission for information in my medical records to be used for this project. In any event of publication and sharing of the data with other researchers and individuals, I understand that this information will not bear my name or other identifiers and that due care will be taken to preserve the confidentiality of this information.

(Signature/Thumbprint (Right / Left) of Subject)_____
(Date of signing)**Part II - to be filled by parent / legal guardian, where applicable**

I, _____ hereby give consent for the above patient to participate in the proposed project. The nature, risks and benefits of the study have been explained clearly to me and I fully understand them.

(Signature/Thumbprint (Right / Left)

(Date of signing)

of parent /legal guardian]

Part III - to be filled witness, where applicable

An impartial witness should be present during the entire informed consent discussion if a subject or the subject's legally acceptable representative is unable to read. After the written informed consent form and any written information to be provided to subjects, is read and explained to the subject or the subject's legally acceptable representative, and after the subject or the subject's legally representative has orally consented to the subject's participation in the study and, if capable of doing so, has signed and personally dated the consent form, the witness should sign and personally date the consent form.

(Name of witness)

(Designation of witness)

(Signature of witness)

(Date of signing)

4.3 Study Enrolment: Patient Assent Form

Patient Assent Form
(15- to 17-year-old participants)

Protocol Title: Trauma Training Effectiveness Research Network

Principal Investigator: Name

1. What we wish to tell you?

I, Dr/Mr/Ms _____ wanted to tell you something we are doing called a research study. A research study is when doctors collect a lot of information to learn something about health and diseases.

You are being invited to be part of a research study. We will explain about the study to you and we will ask you if you would like to be part of the study. You will be given a copy of this document to take home with you.

2. Why are we doing this study?

We want to find out if a training program for doctors on trauma life support will help patients with injury. For this we will collect information from people who are injured like you.

3. What will happen to you if you are in this study?

If you agree to be part of the study, we collect data about your health available in the hospital. We will also contact you in person or by telephone to collect information about your health at the following times:

- On arrival at the emergency department and wards till you are discharged
- On hospital discharge
- 24 hours after arrival to the hospital
- 30 days after arrival to the hospital

During this time, we will also ask you details about your health and general information on returning back to your normal life and experience of the injury in your life.

4. Is this bad or hurtful for you to be part of this study?

No, there will be no pain or risk involved in participating in this study.

5. How will this research be useful to you?

This study will not make you get well. But the doctors may find out something about the how the training program can improve the care of other injured patients like you or if you are injured again.

6. Will everybody come to know about your health condition?

We will not tell other people that you are in this research and we will also not share information about you to anyone who does not work in the research study. We will combine the information from all the patients who agree to be part of this study and no one will be able to identify you from the combined information.

7. Do you get anything for being in the research?

No, you will not receive anything for being part of this study

8. Will you tell me the results?

The information we collect from you and other patients will be combined and studied by doctors and other researchers who are part of the study. We will publish the results in medical and scientific journals so that the knowledge from this study can help injured patients across India and the world. But no one will be able to identify you or your information in these published results.

9. Do you have any questions?

You can ask questions at any time. You can ask now or can ask later. You can talk to me or you can talk to someone else. I have attached the details of the person supervising the study at this hospital in case you want to contact us.

10. Do you have to be in this study?

No, you do not. No one will be force you if you don't want to do this. And remember, you can say yes now and change your mind later. It is up to you. You can do that at any time using the contact information provided below. This will not in any way affect your treatment in this hospital.

11. Who can you talk to or ask questions to?

You can also talk to anyone you want to about this like a family member, friend, or teacher. Doctor. I have attached the details of the person supervising the study at this hospital in case you want to contact us.

Name
Designation, Department
Phone Number
Email

You can also talk to anyone you want to about this like a family member, friend, or teacher. Doctor.

12. Signature of Person Conducting Assent Discussion

I have explained the study to _____ in language
he/she can understand, and he/she has **agreed/not agreed** to be in the study.

(Name of the person conducting assent discussion)

(Signature of the person conducting assent discussion)

(Date of signing)

Supplemental Material 2

2.1 Study Enrolment: Resident Information Sheet

Participant Information Sheet

You are being invited to participate in a research study being conducted at your hospital. Please read carefully the information provided here. If you agree to participate, please sign the informed consent form. You will be given a copy of this document to keep with you.

Protocol Title: Trauma Training Effectiveness Research Network

Principal Investigator: Name

PURPOSE OF THE RESEARCH STUDY

We are currently conducting research in this hospital to study the feasibility of assessing the effect of trauma life support training programs on care and outcome of patients with injury. Life support training, which involves skills in how to take care of injured patients when they come to hospital, may improve how well patients recover from their injuries and we are studying if, and to what extent, that is true. To better measure the outcomes of the training program at your hospital on surgical residents undergoing, we would want to measure your knowledge and confidence during the course of the study. We ask you to participate in this study because you trained at this hospital as part of the study.

STUDY PROCEDURES

If you agree to participate you will be provided training in a specific trauma training program as per the randomization process. This could be Advanced Trauma Life Support (ATLS) and the Primary Trauma Care (PTC) course or standard care. We will collect information related to your demography, academic background and training, as well as measure your perception of improvement in knowledge, skills, and confidence at specific points before, during, and after the training. The data collected will be confidential and anonymous.

The results of the study may be used for research that can be published as scientific articles; however, it will not be possible to identify you by reading any article that may result from this data bank. Further, data from this project will be combined with data from other hospitals that use the same system and shared for other researchers and individuals to use, but it will not be possible to identify you using that data. Research on the data without identifiers may seek to answer other questions than those stated above.

WITHDRAWAL FROM STUDY

Participation in this study is completely voluntary. Even if you agree to participate now you are free to withdraw at any time without giving any reason for doing so. To withdraw you contact any of the study contact persons on the numbers or emails listed below.

POSSIBLE RISKS, DISCOMFORTS AND INCONVENIENCES

We have not been able to identify any major risks associated with participating in this study. If you would at that point, or any other point of time, wish to withdraw from participating in the study, you are free to do so.

POTENTIAL BENEFITS

This research may help to improve the implementation of trauma life support training as well as improve the care of injured patients. Although you will not directly benefit from this study, your participation will contribute to medical knowledge about the effect of training surgical residents in trauma life support training programs to improve trauma care management in India.

CONFIDENTIALITY OF STUDY AND MEDICAL RECORDS

The results of this research may be published as a scientific article; however, it will not be possible to identify you by reading any article that may result from this work. Further, data from this project will be combined with data from other hospitals that use the same system and shared for other researchers and individuals to use, but it will not be possible to identify you using that data.

Also, Regulatory Agencies, Institutional Review Board and Ministry of Health will be granted direct access to the data collected as part of this study to check study procedures, without making any of the data public. By signing the Informed Consent Form attached, you are authorizing such access.

COSTS OF PARTICIPATION

No charge will be levied on you if you take part in this study. You will not receive any compensation for participating in this study.

RESEARCH RELATED INJURY AND COMPENSATION

Due to the observational nature of this study, it is unlikely to cause any research related injury.

WHOM TO CONTACT IF YOU HAVE QUESTIONS

If you have questions about this research study and your rights during the course of this study, you may contact:

Name

Department

Phone Number

Email

2.2 Study Enrolment: Resident Consent Form

Consent Form

Protocol Title: Trauma Training Effectiveness Research Network

Subject Details

Name: _____ NRIC/PNR/SSN No.: _____

Address: _____

Date of birth _____ Phone No: _____

dd/mm/yyyy

I, _____ **agree / do not agree** to participate in the project as described and, on the terms detailed in the Participant Information Sheet. The nature of my participation in the proposed project has been explained to me in _____ by Dr/Mr/Ms _____. I have fully discussed and understood the purpose and procedures of this project. I have been given the Participant Information Sheet and the opportunity to ask questions about this project and have received satisfactory answers and information. I understand that my participation is voluntary and that I am free to withdraw at any time, without giving any reasons. I also give permission to the data collected from me to be used for this project. In any event of publication and sharing of the data with other researchers and individuals, I understand that this information will not bear my name or other identifiers and that due care will be taken to preserve the confidentiality of this information.

(Signature/Thumbprint (Right / Left) of Subject)

(Date of signing)

Supplemental Material 3

Table S3.1: Participant outcomes

Type	Outcome	Specific measurement variable	Analysis metric	Method of aggregation	Time point
Primary	All cause mortality within 30 days from the time of arrival to the emergency department	Death as reported in patient records or by a patient contact person during telephone follow up.	Final value / Change from baseline	Proportion / Difference in proportions between baseline and after intervention.	30 days from the time of arrival to the emergency department.
Secondary	All cause mortality within 24 hours from the time of arrival to the emergency department	Death as reported in patient records or by a patient contact person during telephone follow up.	Final value / Change from baseline	Proportion / Difference in proportions between baseline and after intervention.	24 hours from the time of arrival to the emergency department.
Secondary	Time to all cause mortality during follow up	Time to (in days) death as reported in patient records or by a patient contact person during telephone follow up.	Final value / Change from baseline	Survival analysis, hazard.	End of follow up.
Secondary	Cause-specific in-hospital mortality	Categorical presumed cause of death as judged by the treating physician. Recorded by asking the physician.	Final value / Change from baseline	Proportion / Difference in proportions between baseline and after intervention.	End of follow up.
Secondary	Adherence to the WHO trauma care checklist*	The number of items in the WHO trauma care checklist* that is adhered to during initial management. Recorded through observation.	Final value / Change from baseline	Mean / median.	On first encounter with surgical unit in the emergency department.

Table S3.1: Participant outcomes (*continued*)

Type	Outcome	Specific measurement variable	Analysis metric	Method of aggregation	Time point
Secondary	Fluids for resuscitation in first one hour in patients	The type of fluids, i.e. crystalloids, colloids or blood products, used in the first hour after arrival to the emergency department. Recorded through observation.	Final value / Change from baseline	Proportion.	During the first hour after the patient arrived at the emergency department.
Secondary	Massive transfusion, defined as four or more units of packed red blood cells, plasma or platelets transfused within the first 24 hours after arrival to the emergency department	The number of units of packed red blood cells, plasma or platelets transfused during the first 24 hours after the patient arrived at the emergency department. Extracted from patient records.	Final value / Change from baseline.	Proportion.	24 hours from the time of arrival to the emergency department.
Secondary	Time to first surgery	The time, in hours, from the patient's first encounter with the surgical unit, to start of surgery. Extracted from patient records.	Time to event.	Survival analysis, hazard.	24 hours from the time of arrival to the emergency department.
Secondary	Time to first intubation	The time, in hours, from the patient's first encounter with the surgical unit, to intubation. Recorded through observation.	Time to event.	Survival analysis, hazard.	24 hours from the time of arrival to the emergency department.

Table S3.1: Participant outcomes (*continued*)

Type	Outcome	Specific measurement variable	Analysis metric	Method of aggregation	Time point
Secondary	Time to CT scan	The time, in hours, from the patient's first encounter with the surgical unit, to CT scan. Extracted from patient records.	Time to event.	Survival analysis, hazard.	24 hours from the time of arrival to the emergency department.
Secondary	Ventilator free days	The number of days, out of the total length of hospital stay, that the patient is not mechanically ventilated. Extracted from patient records.	Final value / Change from baseline	Mean / median.	At patient discharge.
Secondary	ICU free days	The number of days, out of the total length of hospital stay, that the patient is not admitted to the ICU. Extracted from patient records.	Final value / Change from baseline	Mean / median.	On patient discharge.
Secondary	Pulmonary complications	Measured by identifying new infiltrates/consolidations on X-ray chest or CT-scan chest or diagnosed by a clinician or reintubated after initially extubated.	Final value / Change from baseline.	Proportion.	On patient discharge or 30 days from the time of arrival to the emergency department, whichever occurs first.

Table S3.1: Participant outcomes (*continued*)

Type	Outcome	Specific measurement variable	Analysis metric	Method of aggregation	Time point
Secondary	Septic shock	Measured by recognizing patients needing vasopressor support beyond the first 48 hours or new initiation of vasopressors in the absence of bleeding or diagnosed by a clinician.	Final value / Change from baseline.	Proportion.	On patient discharge or 30 days from the time of arrival to the emergency department, whichever occurs first.
Secondary	Renal failure	Measured by identifying a patient on dialysis or diagnosed by a clinician.	Final value / Change from baseline.	Proportion.	On patient discharge or 30 days from the time of arrival to the emergency department, whichever occurs first.
Secondary	Coagulopathy	Measured by transfusion of plasma /platelets	Final value / Change from baseline.	Proportion.	On patient discharge or 30 days from the time of arrival to the emergency department, whichever occurs first.
Secondary	Length of stay	The number of days that the patient is admitted to the hospital. Extracted from patient records.	Final value / Change from baseline	Mean / median.	On patient discharge.

Table S3.1: Participant outcomes (*continued*)

Type	Outcome	Specific measurement variable	Analysis metric	Method of aggregation	Time point
Secondary	Quality of life	Measured using the appropriate translation of EQ5D3L. Recorded through interview or telephone follow up.	Final value / Change from baseline	Proportion or mean/median depending on domain.	30 days from the time of arrival to the emergency department.
Secondary	Number of hospitalizations after the index admission during the follow up period	The number of hospitalizations after the first (index) admission. Recorded from patient or patient contact person during telephone follow up.	Final value / Change from baseline	Mean / median.	30 days from the time of arrival to the emergency department.
Secondary	Return to work	Return to any form of work (including house work), as yes or no. Recorded through interview or telephone follow up.	Final value / Change from baseline	Proportion.	30 days from the time of arrival to the emergency department.
Secondary	Need for unplanned re-exploration	New unplanned surgery for a previously operated injury during the index admission. Extracted from patient records.	Final value / Change from baseline	Proportion.	30 days from the time of arrival to the emergency department.

Table S3.1: Participant outcomes (*continued*)

Type	Outcome	Specific measurement variable	Analysis metric	Method of aggregation	Time point
Secondary	Failure of non-operative management	Surgery for initially non-operatively treated conditions, for example liver or splenic injury in stable patients. Extracted from patient records.	Final value / Change from baseline	Proportion.	48 hours.
Secondary	Patient satisfaction	Patient satisfaction measured in Likert scale of 1-5 from Not satisfied to Satisfied completely about their hospital experience that includes healthcare person's behaviour and care received (from Harris et al. 2007)].	Final value / Change from baseline	Median	Prior to discharge
Secondary	Out-of-pocket expenditure	Direct out-of-pocket expenditure (in Indian Rupees, INR) on medicines, diagnostics, medical equipment, and follow-up treatment recorded through interview or telephone follow-up.	Final value / Change from baseline	Mean/Median	At patient discharge and 30 days from the time of arrival to the emergency department

Table S3.1: Participant outcomes (*continued*)

Type	Outcome	Specific measurement variable	Analysis metric	Method of aggregation	Time point
Secondary	Self-ambulatory	Whether or not the patient can walk unassisted. Recorded through interview or telephone follow up.	Final value / Change from baseline	Proportion.	30 days from the time of arrival to the emergency department.
Secondary	Residents' confidence in managing trauma patients	Visual Analogue Scale. Recorded through interview.	Final value / Change from baseline.	Median.	30 days from the time of training, or study start.

Note:

* The World Health Organization's trauma care checklist is available from <https://www.who.int/publications/i/item/trauma-care-checklist> and its implementation was published as Lashoher A, Schneider EB, Juillard C, Stevens K, Colantuoni E, Berry WR, Bloem C, Chadbunchachai W, Dharap S, Dy SM, Dziekan G, Gruen RL, Henry JA, Huwer C, Joshipura M, Kelley E, Krug E, Kumar V, Kyamanywa P, Mefire AC, Musafir M, Nathens AB, Ngendahayo E, Nguyen TS, Roy N, Pronovost PJ, Khan IQ, Razzak JA, Rubiano AM, Turner JA, Varghese M, Zakirova R, Mock C. Implementation of the World Health Organization Trauma Care Checklist Program in 11 Centers Across Multiple Economic Strata: Effect on Care Process Measures. *World J Surg.* 2017 Apr;41(4):954-962. doi: 10.1007/s00268-016-3759-8. PMID: 27800590.

Supplemental Material 4

Table S4.1: Variable list.

Sr No	Variable	Description
1	patient_id	Increment ID in local data base, created by reg_hospital_id, user_id and incremental value
2	local_patient_id	Hospitals patient ID to track the patient in local charts, not uploaded.
3	reg_hospital_id	Assigned ID for the hospital where registration is taking place
4	referral	If patient was referred from another hospital
5	ref_hospital_code	Type of hospital referred from Public or Private or charitable
6	pt_age	Age of the patient
7	pt_gender	Gender of patient
8	moi	As defined in ICD-10 codes
9	dominating_injury_type	Indication of the type of injury produced by the trauma.
10	arrival_by	How did the patient arrive to hospital? (Walking, private car, EMS?)
11	ed_hr	Heart rate recorded in the emergency department.
12	ed_sbp	Systolic blood pressure in the emergency department.
13	ed_dbp	Diastolic blood pressure in the emergency department.
14	ed_gcs	Total GCS score recorded in the emergency department.
15	ed_rr	Respiratory rate recorded in the emergency department.
16	ed_sat	Saturation recorded in the emergency department.
17	ed_temperature	Temperature of patient in the emergency department.
18	ed_pupils	Pupillary response (Unilateral response, bilateral response, non-responsive unilateral or bilateral)
19	ed_shock_index	hr divided by sbp in the emergency department
20	ed_initial_serum_lactate	Serum lactate as measured in the emergency department(from ABG)
21	ed_intial_be	BE as measured in the emergency department (from ABG)
22	intubation	Endotracheal intubation done Before Arrival/After arrival/No
23	time_of_intubation	Date + time of intubation (Set to 0000-00-00 if unknown, null if not intubated)
24	time_mechanical_ventilation_started	Timestamp, coded 0000-00-00 if unknown time, null if patient was not on mechanical ventilation
25	time_mechanical_ventilated_stopped	Timestamp, coded 0000-00-00 if unknown time, null if patient was not on mechanical ventilation
26	chest_tube	Time of insertion of Intercostal drain Before Arrival/After arrival/No
27	time_of_chest_tube	Date + time of intubation (Set to 0000-00-00 if unknown, null if not placed)
28	vasopressors	Yes/No

Table S4.1: Variable list. (continued)

Sr No	Variable	Description
29	time_of_vasopressors	Date + time of intubation (Set to 0000-00-00 if unknown, null if not placed)
30	num_blood transfusion_within_24h	Number of transfusions given within first 24h
31	fluids_within_24h	Quantity of fluids in the first 24 hrs (other than blood)
32	intervention	Other intervention, not defined, free-text (surgical airway, packing of wound, central line, closed reduction)
33	data_of_injury	Date and time when the accident occurred.
34	date_of_transport	Date and time when the EMS service started transportation from the scene, if applicable.
35	date_of_arrival	Date and time of arrival to the emergency department.
36	admitted	If the patient was admitted to hospital
37	date_of_admission	Date and time of admission to the emergency department.
38	surgery_during_stay	Yes/No
39	date_of_surgery	Date and time when the patient was taken to surgery
40	date_of_admission_icu	The time the patient was admitted to the ICU
41	date_of_admission_ward	The time the patient was admitted to the ward
42	date_of_discharge_icu	The time the patient was discharged from the ICU
43	date_of_discharge_ward	The time the patient was discharged from the ward
44	dialysis_within_30_days	Did the patient undergo dialysis during the visit
45	discharge_alive	Yes/No
46	alive_after_30_days	If a 30 day follow up is done, this can be added.
47	time_of_death	Time of death if the patient died in ED or during hospital stay or after discharge
48	type_of_initial_surgery	Free text about the surgery
49	sbp_at_start_of_surgery	First sbp recorded at start of surgery
50	time_surgery_start	Time and date when surgery started
51	time_surgery_end	Time and date when surgery ended
52	findings_or	Findings of during surgery, free text
53	injury_first_or_icd10	ICD10 codes for found injuries on first surgery
54	initial_xray_findings	First X-ray findings
55	time_fast	Time and date when FAST was done
56	findings_fast	Findings on FAST
57	time_first_ct	Time and date when the first CT was done
58	type_first_ct	Type of CT, head, abdomen
59	findings_first_ct	Findings on first CT scan
60	injury_first_ct_icd10	ICD10 code of the injuries found on first CT
61	time_second_ct	Time and date when the second CT was done
62	type_second_ct	Type of CT, head, abdomen
63	findings_second_ct	Findings on second CT scan
64	injury_second_ct_icd10	ICD10 code of the injuries found on second CT
65	findings_additional_ct	Collected findings on following CTs

Table S4.1: Variable list. (continued)

Sr No	Variable	Description
66	injury_following_ct	ICD10 code of the injuries found on following CTs
67	injury_external_1	Description of found external injuries
68	injury_external_1_icd10	ICD10 codes for external injuries
69	body_surface_burn	Burns over body in percent
70	inhalation_injury	If there are any inhalational burnsYes/no
71	co_morbidity_index	CCI Charlson Comorbidity Index
72	occupation	Indicate patient's usual or principal work or business to earn a living
73	prior_facility_interventions	Interventions (procedures, medications, diagnostics) administered in a facility prior to arrival at current facility
74	number_of_serious_injuries	Total number of serious injuries as judged by provider
75	physician_likely_cause_death	Likely cause of death as per the treating doctor
76	cause_of_death	Patient's official (legal) cause of death
77	complication_pulmonary	Measured by identifying new infiltrates/consolidations on X-ray chest or CT scan report suggestive of ARDS, Pneumonia or PTE or diagnosed by a clinician. Patients on mechanical ventilation: Increased FIo2, PEEP
78	pulmonary_complication_reason	
79	complication_septic_shock	Patients needing inotropic support (dopamine >5 microgram/min/ NA/ Vasopressin) beyond the first 48 hours or new initiation of inotropes in absence of bleeding or diagnosed by a clinician.
80	septic_shock_reason	
81	complication_renal_failure	Measured by identifying a patient on dialysis or other renal replacement therapy
82	renal_failure_reason	
83	complication_Coagulopathy	Measured by transfusion of plasma /platelets Or deranged INR and low platelets
84	quality_of_life	EQ5D & EQ5D Y (for <18 years) questionnaire at discharge and at one month post discharge
85	number_of_hospitalizations_for_this_injury	Count of no of re-hospitalisations (other hospitals or same hospitals) after discharge at home (not transfer)
86	return_to_work	Return to any kind of work not necessarily same as pre injury
87	need_for_reexploration_or_resurgery	Need for resurgery for the same region for complication or missed injury
88	failure_of_conservative_management	Failure of conservative treatment which later needed intervention (radiological/vascular or surgical)
89	patient_satisfaction	This will be an ordinal scale recording of the overall satisfaction the patient had after discharge.
90	cost_of_treatment	Direct out-of-pocket costs for treatment to the patient including medicines, diagnostics, equipment, etc.

Table S4.1: Variable list. (continued)

Sr No	Variable	Description
91	selfambulatory_at_discharge	Was the patient able to ambulate on his own at discharge,
92	residents_confidence_in_managing_trauma_patients	

Supplemental Material 5

5.1 Interview: Patient Information Sheet

Patient Information Sheet

You are being invited to participate in a research study. Before you take part in this research study, the study must be explained to you and you must be given the chance to ask questions. Please read carefully the information provided here. If you agree to participate, please sign the informed consent form. You will be given a copy of this document to keep with you.

Protocol Title: Trauma Training Effectiveness Research Network

Principal Investigator: Name

PURPOSE OF THE RESEARCH STUDY

We are currently conducting research in this hospital to study the feasibility of assessing the effect of trauma life support training programs on care and outcome of patients with injury. Trauma means when a person has injuries that are serious. To select the outcomes most relevant to patients like you, we want to know what are the challenges you faced in returning back to normal life after the injury. We ask you to participate in this study because you presented to this hospital after having an injury.

STUDY PROCEDURES AND VISIT SCHEDULE

If you agree to participate, we will call you or a relative three months (90 days) after you arrived at this hospital to hear how you are. We will ask you for permission to visit you at your home. If you agree we will visit you at a time convenient to you to talk to you about health after discharge. You do not have to answer certain questions if you prefer not to. You can also end the interview whenever you want, even if all questions have not been answered. An audio recording of the interview may be taken.

WITHDRAWAL FROM STUDY

Participation in this study is completely voluntary. Even if you agree to participate now you are free to withdraw at any time without giving any reason for doing so. Withdrawing will not affect your ordinary treatment or the care given to you. To withdraw you contact any of the study contact persons on the numbers or emails listed below.

POSSIBLE RISKS, DISCOMFORTS AND INCONVENIENCES

We have not been able to identify any major risks associated with participating in this study. If you would at that point, or any other point of time, wish to withdraw from the study, you are free to do so.

POTENTIAL BENEFITS

Our research may help to study the most relevant outcomes for you to get back to normal life after the injury. Although this research will not affect the care you were given in this hospital at this time, its results might help you if you are injured again in the future, or

others that are injured. There is no assurance you will benefit from this study. However, your participation may contribute to the medical knowledge about the effect of the use of trauma life support training programs on the life of injured patients.

SUBJECT'S RIGHTS

Your participation in this study is entirely voluntary. Your questions will be answered clearly and to your satisfaction. In the event of any new information becoming available that may be relevant to your willingness to continue in this study, you will be informed in a timely manner by the Principal Investigator or his/her representative.

CONFIDENTIALITY OF STUDY AND MEDICAL RECORDS

The results of this research may be published as a scientific article; however, it will not be possible to identify you by reading any article that may result from this work. Further, data from this project will be shared with other researchers in India and abroad, but it will not be possible to identify you using only that data.

COSTS OF PARTICIPATION

If you take part in this study, there will be no charge levied on you. You will not receive any compensation for participating in this study.

RESEARCH RELATED INJURY AND COMPENSATION

The study being observational is not likely to cause any research related injury.

WHOM TO CONTACT IF YOU HAVE QUESTIONS

If you have questions about this research study and your rights or in the case of any injuries during the course of this study, you may contact:

Name

Department

Phone Number

Email

5.2 Interview: Patient Consent Form

Consent Form**Protocol Title: Trauma Training Effectiveness Research Network****Subject's Particulars**

Name:

Address:

Date of birth _____

Phone No

(dd/mm/yyyy)

I, _____ **agree / do not agree** to participate in the research study as described and on the terms set out in the Patient Information Sheet. The nature of my participation in the proposed research study has been explained to me by Dr/Mr/Ms _____. I have fully discussed and understood the purpose and procedures of this study. I have been given the Patient Information Sheet and the opportunity to ask questions about this study and have received satisfactory answers and information.

I understand that my participation is voluntary and that I am free to withdraw at any time, without giving any reasons and without my medical care being affected.

I also give permission for information in my medical records to be used for research. In any event of publication, I understand that this information will not bear my name or other identifiers and that due care will be taken to preserve the confidentiality of this information.

[Signature/Thumbprint (Right / Left) of patient]_____
(Date of signing)

Supplemental Material 6

6.1 Interview: Non-patient participant Information Sheet

Participant Information Sheet

You are being invited to participate in a research study. Before you take part in this research study, the study must be explained to you and you must be given the chance to ask questions. Please read carefully the information provided here. If you agree to participate, please sign the informed consent form. You will be given a copy of this document to keep with you.

Protocol Title: Trauma Training Effectiveness Research Network

Principal Investigator: Name

PURPOSE OF THE RESEARCH STUDY

We are currently conducting research in this hospital to study the feasibility of assessing the effect of trauma life support training programs on care and outcome of patients with injury. Trauma means when a person has injuries that are serious. To select the outcomes most relevant to patients we want to know what are the challenges patients face in returning back to normal life after the injury.

STUDY PROCEDURES

If you agree to participate, we will visit you at a time convenient to you to talk to you about challenges trauma patients face. You do not have to answer certain questions if you prefer not to. You can also end the interview whenever you want, even if all questions have not been answered. An audio recording of the interview may be taken.

WITHDRAWAL FROM STUDY

Participation in this study is completely voluntary. Even if you agree to participate now you are free to withdraw at any time without giving any reason for doing so. To withdraw you contact any of the study contact persons on the numbers or emails listed below.

POSSIBLE RISKS, DISCOMFORTS AND INCONVENIENCES

We have not been able to identify any major risks associated with participating in this study. If you would at that point, or any other point of time, wish to withdraw from the study, you are free to do so.

POTENTIAL BENEFITS

Our research may help to study the most relevant outcomes for patients to get back to normal life after the injury. There is no assurance you will benefit from this study. However, your participation may contribute to the medical knowledge about the effect of the use of trauma life support training programs on the life of injured patients.

PARTICIPANT'S RIGHTS

Your participation in this study is entirely voluntary. Your questions will be answered clearly and to your satisfaction. In the event of any new information becoming available that may be relevant to your willingness to continue in this study, you will be informed in a timely manner by the Principal Investigator or his/her representative.

CONFIDENTIALITY OF THE STUDY

The results of this research may be published as a scientific article; however, it will not be possible to identify you by reading any article that may result from this work. Further, data from this project will be shared with other researchers in India and abroad, but it will not be possible to identify you using only that data.

COSTS OF PARTICIPATION

If you take part in this study, there will be no charge levied on you. You will not receive any compensation for participating in this study.

RESEARCH RELATED INJURY AND COMPENSATION

The study being observational is not likely to cause any research related injury.

WHOM TO CONTACT IF YOU HAVE QUESTIONS

If you have questions about this research study and your rights during the course of this study, you may contact:

Name

Designation, Department

Phone Number

Email

6.2 Interview: Non-patient participant Consent Form

Consent Form

Protocol Title: Trauma Training Effectiveness Research Network

Subject's Particulars

Name:

Address:

Date of birth _____
(dd/mm/yyyy)

Phone No

I, _____ **agree / do not agree** to participate in the research study as described and on the terms set out in the Participant Information Sheet. The nature of my participation in the proposed research study has been explained to me by Dr/Mr/Ms _____ I have fully discussed and understood the purpose and procedures of this study. I have been given the Participant Information Sheet and the opportunity to ask questions about this study and have received satisfactory answers and information.

I understand that my participation is voluntary and that I am free to withdraw at any time, without giving any reasons.

In any event of publication, I understand that it will not bear my name or other identifiers and that due care will be taken to preserve the confidentiality of this information.

[Signature/Thumbprint (Right / Left) of participant]

(Date of signing)

Supplemental Material 7

Table S7.1: Shows which outcomes that will be assessed in which subgroups.

	All patients	Men	Women	Blunt multisystem	Penetrating	Shock	Severe traumatic brain injury	Elderly
All cause mortality within 30 days from the time of arrival to the emergency department	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
All cause mortality within 24 hours from the time of arrival to the emergency department	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Time to all cause mortality during follow up.	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Cause-specific in-hospital mortality.	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Adherence to the WHO trauma care checklist.	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Fluids for resuscitation in first one hour in patients.	Yes	No	No	No	No	Yes	No	No

Table S7.1: Shows which outcomes that will be assessed in which subgroups.
(continued)

	All patients	Men	Women	Blunt multisystem	Penetrating	Shock	Severe traumatic brain injury	Elderly
Massive transfusion, defined as four or more units of packed red blood cells, plasma or platelets transfused within the first 24 hours after arrival to the emergency department.	Yes	No	No	No	No	Yes	No	No
Time to first surgery.	Yes	No	No	No	No	Yes	No	No
Time to first intubation.	Yes	No	No	No	No	No	No	No
Time to CT scan.	Yes	No	No	No	No	No	No	No
Ventilator free days.	Yes	No	No	No	No	No	No	No
ICU free days.	Yes	No	No	No	No	No	No	No
Pulmonary complications.	Yes	No	No	No	No	No	No	No
Septic shock.	Yes	No	No	No	No	No	No	No
Renal failure.	Yes	No	No	No	No	No	No	No
Coagulopathy.	Yes	No	No	No	No	No	No	No
Length of stay.	Yes	No	No	No	No	No	No	No
Quality of life.	Yes	No	No	No	No	No	No	No
Number of hospitalizations after the index admission during the follow up period.	Yes	No	No	No	No	No	No	No
Return to work.	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No

Table S7.1: Shows which outcomes that will be assessed in which subgroups.
(continued)

	All patients	Men	Women	Blunt multisystem	Penetrating	Shock	Severe traumatic brain injury	Elderly
Need for unplanned re-exploration.	Yes	No	No	No	No	No	No	No
Failure of non-operative management.	Yes	No	No	No	No	No	No	No
Patient satisfaction.	Yes	Yes	Yes	Yes	No	No	No	No
Out-of-pocket expenditure.	Yes	No	No	No	No	No	No	No
Self-ambulatory.	Yes	No	No	No	No	No	No	No