

BMJ Open Acceptability of extending HPV-based cervical screening intervals from 3 to 5 years: an interview study with women in England

Martin Nemeč , Jo Waller , Jessica Barnes, Laura A V Marlow 

To cite: Nemeč M, Waller J, Barnes J, *et al.* Acceptability of extending HPV-based cervical screening intervals from 3 to 5 years: an interview study with women in England. *BMJ Open* 2022;**12**:e058635. doi:10.1136/bmjopen-2021-058635

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-058635>).

Received 01 November 2021
Accepted 08 April 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

Cancer Prevention Group, School of Cancer and Pharmaceutical Sciences, King's College London, London, UK

Correspondence to
Dr Laura A V Marlow;
l.marlow@kcl.ac.uk

ABSTRACT

Objectives The introduction of primary Human Papillomavirus (HPV) testing in the National Health Service (NHS) Cervical Screening Programme in England means the screening interval for 25–49 years can be extended from 3 to 5 years. We explored women's responses to the proposed interval extension.

Methods We conducted semi-structured phone/video interviews with 22 women aged 25–49 years. Participants were selected to vary in age, socioeconomic and screening history. We explored attitudes to the current 3-year interval, then acceptability of a 5-year interval. Interviews were transcribed verbatim and analysed using framework analysis.

Results Attitudes to the current 3-year interval varied; some wanted more frequent screening, believing cancer develops quickly. Some participants worried about the proposed change; others trusted it was evidence based. Frequent questions concerned the rationale and safety of longer intervals, speed of cancer development, the possibility of HPV being missed or cell changes occurring between screens. Many participants felt reassured when the interval change was explained alongside the move to HPV primary screening, of which most had previously been unaware.

Conclusions Communication of the interval change should be done in the context of broader information about HPV primary screening, emphasising that people who test negative for HPV are at lower risk of cell changes so can safely be screened every 5 years. The long time needed for HPV to develop into cervical cancer provides reassurance about safety, but it is important to be transparent that no screening test is perfect.

INTRODUCTION

By the end of 2019, Human Papillomavirus (HPV)-based screening had been introduced into the National Health Service (NHS) Cervical Screening Programme across England.¹ Tests for high-risk HPV have higher sensitivity than cytology-based screening, significantly reducing the likelihood of false negatives.² The superior negative predictive value of HPV-based screening (almost 100%)² means that 5-yearly screening is likely to be

Strengths and limitations of this study

- ⇒ This is the first study exploring acceptability of extended cervical screening intervals in England.
- ⇒ The findings suggest that explaining the introduction of HPV testing and the rationale for screening interval changes will improve acceptability.
- ⇒ The sample was self-selected, but included women with a range of screening experiences.
- ⇒ The study was carried out during the coronavirus pandemic which may have affected women's views.

safe. At present, the NHS Cervical Screening Programme recommends 3-yearly screening for women aged 25–49 years, moving to 5-yearly from age 50 to 64. A universal recommendation for 5-yearly screening would therefore represent an extension (from 3 to 5 years) for the younger cohort.¹ Other countries have recently implemented similar changes prolonging the interval between cervical screens to 5 years (eg, Australia)³ or 10 years (the Netherlands).⁴

Acceptability is an important consideration ahead of changes to health policy.⁵ Following a recent systematic review,⁵ it has been argued that 'All components of the screening programme should be clinically, socially and ethically acceptable to screening participants' (p. 427). However, previous research suggests that the transition to longer screening intervals may not be considered acceptable to all women.^{6–8} Following changes to the cervical screening programme in Australia (the 'renewal', where HPV-based testing, changes to age-based eligibility and extended intervals between screens were introduced together),⁹ research revealed public concerns. These included beliefs that women's health was being endangered and devalued as a result of budget cuts, and that changes would lead to missed or late diagnoses of cervical cancer.⁶ These concerns were related to limited

understanding of the rationale for the interval changes as well as limited knowledge of HPV regression and the slow progression from HPV infection to cervical cancer.¹⁰ More recently, a qualitative study suggested that Australian women wanted to be consulted about any further changes to the cervical screening programme prior to implementation.¹¹ Previous UK-based research suggests acceptability of extended screening intervals may be influenced by how the change is presented. A clearer presentation of the rationale for extended intervals could increase acceptability.¹²

While the term ‘acceptability’ has been used in health services research for many years,¹³ it has recently been defined and operationalised as a ‘multifaceted construct which reflects the extent to which people delivering or receiving a healthcare intervention consider it to be appropriate, based on anticipated or experienced cognitive and emotional responses.’ (Theoretical Framework of Acceptability, TFA).¹³ Sekhon *et al*¹⁴ argue that acceptability includes a number of constructs including affective attitude (how the person feels about the relevant healthcare intervention), burden (how difficult participating in the intervention seems), ethicality (how the intervention fits with one’s values), intervention coherence (how much the person understands the intervention), opportunity costs (what must be given up to engage in the intervention), perceived effectiveness (the perceived likelihood of the intervention achieving its purpose) and self-efficacy (perceived ability to perform the behaviour associated with the intervention).

This study aimed to explore acceptability of 5-yearly cervical screening intervals for women aged 25–49 years, drawing on the TFA as a broad framework.

METHODS

Design

This was an exploratory qualitative study, using one-to-one semistructured interviews.

Setting

The study was carried out in England, where a call-recall cervical screening programme invites women aged 25–64 for free cervical screening. Invitations and reminders are sent to everyone who is eligible and is registered with a primary care provider. The programme used liquid-based cytology with HPV triage until 2019. Following a large-scale pilot,¹⁵ primary HPV screening was rolled out in all areas over the course of 2019 but no changes have yet been made to screening intervals. As the roll-out was phased, there was no public announcement of the change, and women would only be made aware of it by reading their invitation letter and the accompanying information leaflet or by visiting the NHS website (<https://www.nhs.uk/conditions/cervical-screening/>).¹⁶

Participants

Participants were identified and contacted via email by a research recruitment agency with a large participant panel. The number of participants approached was not recorded. Those who were interested in taking part completed a short questionnaire providing information on demographic characteristics and cervical screening history. We used purposive sampling to ensure variation with respect to age (within the eligible age range: 25–49 years), socioeconomic background and cervical screening experience. Participants were emailed a participant information sheet and consent form ahead of the interview. Signed or initialled consent forms were returned by email or, where this was not possible, consent was collected verbally prior to interview. Twenty-two participants took part in a single interview and were paid £50 to thank them for their time. There were no withdrawals following consent.

Procedure

Interviews were conducted by phone (n=12) or video call (using Microsoft Teams; n=10) according to participant preference by either JB or LAVM, both female postdoctoral researchers with training and experience in semi-structured interviewing and no prior relationship with the participants. Interviewers introduced themselves as researchers from King’s College London, funded by PHE. Interviews lasted around 40 min (mean=39 min, range 14–60 min) and were audiorecorded. Field notes were not used.

A topic guide was used to ensure key topics of interest were covered. The topic guide (see online supplemental file 1) included three sections: (1) views about the current screening interval, (2) views about the possibility of extended intervals, (3) reactions to key messages about the interval change. Prompts were developed for the first two sections to explore the different ways in which participants may find screening intervals acceptable or unacceptable. These were designed to encourage discussion of different elements of acceptability as identified in the TFA.¹⁴ In section 2, participants were provided with additional information explaining the rationale for the 5-year interval in a stepped way before discussing views in further detail. This was provided verbally using a script (see Topic Guide, online supplemental file 1). In section 3, participants were presented with additional messages designed to reassure women about the interval change (online supplemental file 1). This part of the interview was designed to help us develop key messages for a future study,¹⁷ but also prompted further discussion so this part of the data was also analysed. The topic guide was not pilot tested.

Patient and public involvement

Neither patients nor the public were involved in the design, conduct, analysis or interpretation of this study.

Analysis

Interviews were transcribed verbatim by an external agency. Due to time-constraints, transcripts were not

returned to participants for comment or correction. Analysis began after completion of the first five interviews and continued alongside data collection. Whole transcripts were analysed as one entity, rather than as separate sections.

We used framework analysis,¹⁸ following the stages outlined by Gale *et al.*¹⁹ Inductive line by line coding was performed on the first three transcripts by MN (male research assistant with experience and training in qualitative analysis and no prior relationship with the participants) and LAVM after which, codes were agreed for subsequent analysis and were grouped into categories using MIRO, an online platform that allows creation and movement of virtual post-it-notes. This working analytical framework was then applied to the rest of the transcripts and refined throughout the process. Transcripts were coded in order and regular discussions on modifications of the framework were held (LAVM and MN). After the first nine transcripts had been coded, a final analytical framework was confirmed. This framework was then used to code the remaining transcripts (MN) before charting (entering summarised data) into a framework matrix (a code by case spreadsheet) in Excel. Data were then charted (by MN and checked by LAVM). MN and LAVM then interpreted the data with insights from JW (female postdoctoral researcher with training and experience in qualitative research and no prior relationship with the participants) and JB. Participant checking was not part of our analysis plan.

RESULTS

Sample characteristics

We carried out 22 interviews. We used a stopping criterion of three meaning that data collection stopped at the point where no new themes had emerged in the previous three interviews.²⁰

Demographic characteristics are shown in table 1. Four women were aged 25–29 years, 14 were aged 30–39 years and 4 were 40–49 years (age range: 25–49 years, mean age=34.9 years, SD=6.4 years). Participants were evenly distributed across occupational social grade and Any White versus ethnic minority groups. Seventeen participants reported having been screened in the last 3 years (up to date with screening) and five were overdue.

Summary of themes and processes that arose from the interviews

There were two broad primary topics within the interviews (1) women's acceptance of existing screening intervals and (2) their acceptance of the proposed 5-yearly screening interval (see figure 1). Within these topics specific themes related to beliefs about the existing and extended screening intervals. Throughout the interviews, as more information was provided, women's understanding of the rationale for the change improved. Two broader influencing factors ('personal risk' and 'trust and empowerment') were also identified. Each theme is described below, with example quotes (see online supplemental file 2, for additional participant quotes).

Table 1 Sociodemographic characteristics of the sample (n=22)

	N
Age	
25–29 years	4
30–39 years	14
40–49 years	4
Occupational social grade*	
AB (high)	6
C1	5
C2	6
DE (low)	5
Marital status	
Single	9
Married/civil partnership/cohabiting/engaged	10
Divorced	3
Working status	
Working (full-time or part-time)	16
Not working	4
Furloughed due to COVID-19	2
Ethnicity	
Any White background	11
Mixed ethnic background	3
Asian background	5
Black background	3
Sexual orientation	
Heterosexual/straight	16
Gay/Lesbian/Bisexual	5
Prefer not to say	1
Educational level	
None/GCSE or equivalent	9
A Level or equivalent	6
Postgraduate diploma	1
University degree or higher	6
Last cervical screen	
Within the last 3 years	17
More than 3 years ago	5
Had the HPV vaccine?	
Yes, I have	6
No/don't know	16
Heard of HPV before today?	
Yes, I have heard of it	18
No, I have not heard of it	4
Self-rated HPV knowledge	
Good/very good	6
Fair/poor/very poor	16

*Occupational social grade is based on the occupation of the household's chief income earner 'AB' (higher and intermediate managerial, administrative and professional), 'C1' (supervisory, clerical and junior managerial administrative and professional), 'C2' (skilled manual workers), and 'DE' (semi-skilled and unskilled manual workers, and state pensioners, casual and lowest workers, unemployed with state benefits only).
GCSE, General Certificate of Secondary Education; HPV, Human Papillomavirus.




Figure 1 Schematic representing the themes/processes from the interviews. HPV, Human Papillomavirus.

Acceptance of existing screening intervals

Acceptance of existing screening intervals was influenced by beliefs about the importance of prevention and early detection, the asymptomatic nature of cervical abnormalities and the speed of development of cervical cancer. These beliefs were driven by existing knowledge of cancer and how screening works and perceptions of existing public health messages about screening.

Regular cervical screening was felt to bring the benefits of prevention and early detection:

It may be treatable, but it might not be. And the treatability that might be linked in some way to an early detection. ... I think in a lot of people's mind, including mine, is that that you need to get in there quickly and you've got a better chance if things are treated early. [P13; 40–49 years; up-to-date with screening]

Cervical cancer was thought to be asymptomatic or to have symptoms only at a later stage so frequent screening was seen as way of improving outcomes. For some this meant screening more frequently than every 3 years would be considered an improvement to services:

...without the test you wouldn't know if you've got a problem with your cervix. It's not like you've got a symptom and you know you've got a symptom. [P16; 30–39 years; up-to-date]

Acceptance of screening intervals was also influenced by beliefs about the timeline of cervical cancer development. Understanding of the speed that cervical cancer develops varied:

I know that it's supposedly, like, if you do have cervical cancer then it's a very slow process from pre-cancerous to actually cancerous. [P18; 30–39 years; up-to-date]

...probably 36 months is a long time if, if after one smear test somebody has started developing a problem and then after 36 months compared to the last one, it might have gone if it, you know, it might go out of control... [P14; 30–39 years; overdue]

Acceptance of 5-yearly intervals

Initially, discussion about acceptance of 5-yearly screening reflected how women felt about the existing interval. Women who felt strongly that 3-yearly screening was important or not enough, were resistant to a 5-yearly interval, expressing shock and concern. Initial reactions centred around how a 5-yearly interval contradicted their existing beliefs about the importance of screening and early detection:

It's just- The whole—Every three years has always been for prevention, to make sure you're healthy, and

now suddenly, it's every five years. I don't know. [P7; 30–39 years; up-to-date]

After that [Jade Goody's death from cervical cancer] there was such a massive drive and a massive like— Please go and get tested. Please do this. Please do that [...] I really felt like it made a difference. But now, if the NHS is saying, you know, you can't even have one when people want it. That, to me, makes no sense. [P2; 25–29 years; up-to-date]

And I just feel like that gap between the screenings, I think personally, is much too long to catch something early enough... So it seems like quite a long time between them, yeah... I think it probably should be shorter... And then what do you do? Like how do you— Because you're not getting the screening, how do you know that it hasn't progressed to abnormal cells or to cervical cancer within that 5 year period if you're not being screened? [P6; 25–29 years; up-to-date]

For women who felt ambivalent about the current interval, less frequent screening was met with a more positive response and a recognition of the benefits. Perceived benefits included not needing as many screening appointments, the advantage of feeling reassured for longer and the potential to save the NHS money. For those who disliked the procedure, found it intrusive, worrying or inconvenient, the need for fewer screens was an obvious benefit of the interval change.

Especially as they're not the most comfortable things to go through either. So that'd be a positive in the fact that you wouldn't have to endure it every three years, [...] you don't have to go through it. The embarrassment. [P16; 30–39 years; up-to-date]

The new interval was also seen to provide reassurance for a longer period after a negative result. This view was more prominent among those without previous direct or indirect experience of abnormalities or cancer.

Um, and also, I guess that you know that you're like clear for five years. I mean, in theory, because that would mean that the NHS is giving you the confidence that because they think it can be so long between screenings, you've had your screening and then you can have, like, the reassurance that, you know, I'm kind of good to go for five years. [P6; 25–29 years; up-to-date]

So, it's kind of, it's probably a positive, you know, that you're not constantly being checked up on and made to feel guilty if it was meant to be more frequent and then you miss the frequency or, you know, things happen, if you go travelling, if you're away for some time. [P6; 25–29 years; up-to-date]

Those preferring shorter screening intervals described being more likely to attend as soon as they were due, when a longer interval is implemented:

I think I'd make it more of a point to get there as soon as my screening was due. Just in case... I'd be getting myself there as soon as. [P5; 30–39 years; up-to-date]

However, some participants wondered if the new interval could have an impact on screening attendance, because a longer interval may make it easier to forget to attend or may reduce perceptions of urgency.

Maybe there's even less of an impetus to respond to it and less of a sense of, it's important to respond to it. So, I would be concerned that it kind of, the lengthening of the time somehow undermines its importance. [P13; 40–49 years; up-to-date]

Finally, for some, the potential for a new test to save the NHS money was a positive consequence of the change, as long as it also resulted in effective screening.

...if they can save money, then great. The main thing is, just making sure it's effective. Yeah. [P10; 30–39 years; up-to-date]

The process of accepting a longer interval

During the interviews, many of the women seemed to go through a process of increasing confidence in the interval change, moving from their existing acceptance of 3-yearly screening to an acceptance of 5-yearly screening. This was driven by their existing awareness of HPV primary screening and information provided during the interview to address initial concerns about the risks associated with a longer interval.

Awareness of HPV primary screening

At the start of the interviews (section 1), not all women were aware of the move to HPV primary screening, leading some participants to question why there was an interval change happening now. In the absence of an explanation, a range of potential reasons for the interval change were put forward, including the need to cut costs and the availability of HPV vaccination.

If it's about like, capacity and saving money and I know it's probably a big strain, you know, to do all of that, but if that's the reason they want to make it a long period of time and this is the reasoning behind it, I think you're dicing, you're dicing with death a little bit there. [P5; 30–39 years; up-to-date]

Once HPV primary screening had been explained to them (in Section 2), some participants continued to question why a successful test such as the Pap smear would need to be changed at all.

I think there needs to be a valid reason as to why they're changing it now. Why didn't they start this from the beginning? [...] I don't know why they did it every three years, but if they did it every three years, there must have been a reason why they did that. [P7; 30–39 years; up-to-date]

Additional questions about the implications of HPV primary testing were raised, including if the patient experience during the examination itself would be different and what would happen for people with abnormal cells or HPV-positive results (ie, whether they would also be invited 5 yearly).

So, will, the process, like when you're actually in the nurse's office change, or will that still be exactly the same? [P6; 25–29 years; up-to-date]

The information that screening would be more frequent for HPV-positive individuals was reassuring:

Knowing that if there is a problem in that HPV then they can check it out every year. Hopefully it will sort itself out. I think it's actually better. [P22; 40–49 years; overdue]

Understanding the rationale for the changes to a longer interval

Across all sections, participants raised questions about the safety and effectiveness of an extended interval and needed additional information to help build their confidence. For some, even the potential for a small risk would be a concern. These safety questions were raised particularly by women in their 20s–30s.

But me personally, I would want to detect it as early as possible. Because you could be in that 1 or 2% like Jade Goody, you don't know, there's still a chance. [P8; 25–29 years; up-to-date]

Some women suggested that cervical cancer can appear between screens.

... you can get the HPV virus at any point in those five years. Right, so if you've had your test and you get it at year two, then you have to wait, you know, it's like three years before you have your next test. And in that time, it could develop into cervical cancer. So, I'm wondering would it then be too late? I don't know how quickly it develops and things like that. [P12; 30–39 years; up-to-date]

Concerns regarding the detection of HPV and cervical cancer also included the possibility of false negatives resulting from the use of the HPV test or the potential inability of the HPV test to identify abnormal cells without the presence of HPV.

So you—So you could, very small chance, have abnormal cells without HPV, and that wouldn't get picked up? [P4; 30–39 years; up-to-date]

These also included views that a longer interval might result in more adverse outcomes including more cancers, later diagnosis, more invasive treatment and more deaths:

Say there were abnormalities, but the test didn't detect that, probably because they took very small amount of sample or whatever. And the test didn't detect the abnormalities. And then there are five years now. So, in that—She has to wait five years before she

gets the next appointment, and then next appointment, they will discover that it has already happened, and the cells are already abnormal.... from the time the smear test was done when she was abnormal, but the test showed normal to five years, is it not going to be more risky for her life? [P14; 30–39 years; overdue]

Learning that the HPV test allows screening to pick up problems earlier in the cancer development process and that HPV infection takes a long time to develop into abnormal cells (more than 5 years), improved understanding of the interval change and was met with feelings of reassurance during sections 2 and 3 of the interviews.

I feel much comfortable knowing the information I've got now. Especially when, you know, the time it takes to develop. Yeah, it makes complete sense. And yeah, I'd be totally comfortable with that. [P12; 30–39 years; up-to-date]

That makes sense to me. I would rather start at the HPV stage and know you're being tested there rather than leaving it so it could possibly then develop onto the abnormal cells and then if it is every five years, from there I feel like it will probably be a lot better probably. [P15; 40–49 years; overdue]

Influencing factors

Two factors seemed to influence acceptance of screening intervals: perceptions of personal risk (based on age and screening history) and feelings of trust and empowerment. For some women these factors meant that 5-yearly screening was still not acceptable, even towards the end of the interview process when messages about the rationale and safety had been presented.

Perceptions of personal risk

Participants reflected on their personal risk of developing abnormal cells and cervical cancer when describing their views about the existing and proposed intervals. Individuals with prior experience of abnormal results tended to feel that the new interval would pose a risk, even if small. They referred to their past results, their own safety, and worried that they might be 'the rare case' to develop aggressive, fast-growing cancer. Past experiences with cervical and other types of cancer in the family were mentioned.

I wouldn't feel 100% comfortable. I really wouldn't, but that's because of my personal experiences. I think for anybody else, if it's all clear and it's always been clear then maybe, maybe it'd be alright. But I wouldn't like that. I'd want that option of three years.' [P19; 30–39 years; up-to-date]

...it then makes you worry even more because you think, hang on, does that increase my chances as well that it's in the family and should I be really worrying here? [P21; 30–39 years; overdue]

Age was also seen to impact personal risk. Younger participants expressed the need for a more individual

approach to screening (eg, based on age) and felt they may need screening more frequently:

I'm just saying that I think it is a worry for young women and I think they should look at an age bracket and maybe test them a bit more often. Does that make sense? [P16; 30–39 years; up-to-date]

The reasons women gave for this included beliefs that younger women were at more risk of cervical cancer, that implications for future fertility were more important in women who had not yet had children and that younger women had generally had fewer screens.

Five years would just be too long. Yeah, because when you first start going for your smear test at age 25, then to leave it again until 30, that's a big change. It's a long time to leave. So yeah, in my opinion, I wouldn't be happy with it being every five years. [P8; 25–29 years; up-to-date]

Older participants, by contrast, described themselves as feeling at lower risk:

No, I'll be honest, I feel like because I am—I never thought I'd say this, but because I am in the older bracket, I do feel a bit safer just knowing the causes of HPV and knowing that I've been fully checked out and things like that. So, I feel a lot more confident in terms of my having delayed my screening and not having gone in yet. [P15; 40–49 years; overdue]

Participants also assessed their risk based on their understanding of HPV transmission and their own relationship status, with those in longer term relationships and those who only had sex with women feeling at lower risk:

Yeah, I mean I suppose if you've not changed, you know, your partner or anything and you're still in a relationship and you've not got the worst strain of HPV, I'm presuming that you would be happy to continue to do it every five years [...] but, you know, if there is someone that, you know, has got more sexual partners and they are, you know, more frequently and stuff like that, they would really not want to wait five years. Because it's just like having a sexually transmitted infection, you might get treated for it and then go and pick it up again the next week. [P8; 25–29 years; up-to-date]

Perceptions of risk, especially in younger women aged 20–39 years, led some to raise the possibility of paying for private screening so they could be screened more often:

Could a woman, if they wanted to, sort of book in for a cervical smear more than the five year period? [P1; 30–39 years; up-to-date]

...you're forcing people to maybe go private for it and having to pay, and that's a shame, because I would probably end up going down that route. [P3; 30–39 years; up-to-date]

Concerns were voiced regarding the 'one size fits all' approach to screening. A more personalised approach to cervical screening, taking into account a woman's age and prior experiences (with HPV-positive results, abnormal cells, treatment, follow-up) was considered preferable.

Trust and empowerment

Trust and empowerment was also an overarching theme including perceptions of trust in healthcare professionals and decision-makers, and in science and research more generally. Some participants expected that any change to the screening programme would be a reasoned decision, based on consensus among experts. Trust in the government, the NHS, Public Health England and publicly funded bodies seemed to influence this:

...with the NHS within this country, I feel confident that they're doing the right thing, so I'd be comfortable going along with what they think is right...I have faith that the public health are doing the right thing. [P16; 30–39 years; up-to-date]

Participants also expressed their trust in science and research findings and wanted to see more information about research evidence supporting the interval change:

Like, the government would need to put things out and be like, 'This is what we're backing. We know the science, and this is it, and this is okay,' to reassure a lot of women.' [P3; 30–39 years; up-to-date]

For some participants, a change to extended intervals was seen as a cost-cutting exercise and sent a message about the value of women's health. There was a particular tendency to express this view among those with previous experience of abnormal results.

-and think, okay, well, maybe this, like, it could be based in science. It seems more likely to be, like, some kind of money-saving thing and then I would feel much less reassured. [P10; 30–39 years; up-to-date]

Well, these things are expensive, aren't they? And then they're trying to cut costs somewhere along the line and it's usually women's health, children's health, old people's health. [P22; 40–49 years; overdue]

Some described feeling powerless and frustrated about not being able to have more frequent screening, with a resignation that there was no choice but to accept the upcoming change. This was expressed exclusively by women with previous experience of abnormal results (own or of family/friends).

Um, well, if they change it for five years, then you have no choice. One voice is not going to change it. [P14; 30–39 years; overdue]

DISCUSSION

This is the first study in England to explore, in depth, women's attitudes to extending cervical screening intervals

in the context of the move to HPV primary screening. Acceptability of extended intervals varied between participants and for many this changed over the course of the interview as additional information was provided. More than 6 months after the full roll-out of HPV primary screening across England, many of our participants were unaware of this change. Information about HPV primary screening and the rationale for the interval change led to a clearer understanding and, for some, acceptance of a 5-year interval. Acceptability was influenced by views on the existing 3-year interval, which were underpinned by beliefs about the importance of prevention, the asymptomatic nature of cervical cell changes and perceptions of the timeline of cervical cancer development. It was also influenced by a range of factors including personal risk perceptions and trust in decision-makers.

Our study benefited from a sample that was diverse with respect to socioeconomic status, ethnic background, relationship status and screening experience. However, it was also self-selected and it is likely that the views of women who are not interested in screening or in participating in research were under-represented. Although there were no explicit assumptions brought to the research, all researchers involved were interested in cancer prevention and LAVM/JW brought potential bias of past experience interviewing women about cervical screening and publishing on this topic. JB and MN had no previous experience in qualitative cervical screening research. We were able to identify positive as well as negative attitudes towards a change to the screening interval. The use of semistructured interviews, during which women were provided with additional information, allowed us to explore how acceptability of extended screening might change as more information is provided.

Many of the themes raised towards the beginning of the interviews, prior to an explanation of the rationale for changing the interval, were similar to those highlighted in an Australian petition that followed implementation of extended intervals.¹⁰ For example, women expressed preferences for more frequent screening, perceptions that cancer develops fast and beliefs that longer intervals would put lives at risk. De-valuing of women's health and beliefs that the change was a cost-cutting exercise, were also expressed, in line with the Australian petition.⁶ The views expressed following the provision of information were strikingly similar to another Australian study¹¹ which found that despite initially low awareness of HPV primary screening and understanding of HPV, concerns about longer intervals could be alleviated for many women by presenting information explaining the rationale for the changes, including the introduction of HPV primary screening, HPV-to-cancer development and the length of time this takes. These findings are also consistent with a quantitative study that showed women who were presented with this information had a better understanding of the change.¹²

Many of our findings could be mapped onto the TFA.^{13 14} Of key importance was intervention coherence.

Without a good understanding of the role of HPV in cervical cancer development, HPV primary screening and the rationale for the longer interval, women struggled to find the change acceptable. Many raised issues with the perceived effectiveness of the new approach, questioning the safety of the longer interval, and expressing concerns about false negative results and more adverse outcomes. Affective attitude appeared to be important in acceptance of existing intervals and extended ones. For some participants, there were positive attitudes associated with having to undergo an uncomfortable procedure less often (once they were reassured about safety). Participants also expected to feel positive emotions following a 'normal' result with a longer interval, which would be reassuring for a longer period. Others expressed worry and other negative emotions at the prospect of a change which they felt powerless to challenge. Our theme of 'Trust and empowerment' could be related to the TFA construct of ethicality and suggests trust in decision-makers will likely affect the extent to which the change is seen as ethical and therefore acceptable. Opportunity costs, burden and self-efficacy were less obviously important in this context.

Perceived risk was discussed throughout the interviews and was influenced by age, previous experience of abnormal screening results and family history. These seemed to impact acceptability of a longer interval, consistent with other work.²¹ In a quantitative study exploring acceptability of interval changes following information exposure,¹² participants who were classed as irregular screening attenders showed more positive attitudes towards the interval extension. We also found that individuals who were overdue for their 3-yearly screen tended to be less negative about the prolonged interval.

Our study suggests that communication about the rationale for extending screening intervals will need to include a clear explanation of current HPV screening. Some participants wanted to see an explanation of the scientific evidence supporting the interval change, including information on the sensitivity of the HPV test, timeline of abnormal cells and cervical cancer development, and robustness of studies on which the decision to extend intervals was based. Conveying the messages that HPV testing has the potential to intervene earlier in the oncogenic process than cytology, and that the timeline from HPV exposure to cancer development is long, is likely to help increase acceptability. Multilevel communication, including public awareness campaigns, print and online information, as well as conversations between trusted health professionals and those attending for screening will be needed to ensure widespread understanding and acceptance.

Due to the COVID-19 pandemic we were unable to carry out face-to-face data collection and decided that interviews were more feasible and accessible as they could be done online or by phone. We acknowledge that focus groups may have resulted in deeper discussion about some of the topics discussed, but we do not expect that the themes would have been different. This study provided an

indication of the information that is important for women to understand and accept the proposed screening interval change. However, more research, with larger samples, is needed to examine the impact of specific messages and identify the best way of communicating information about HPV testing and the rationale for longer intervals to screening-eligible women. We developed a script to discuss the interval changes within the interviews, but there may be additional content that we did not discuss that would influence acceptability. For example, we did not explicitly discuss the potential for overdetection if HPV-based screening is used with a 3-yearly interval. Previous studies have suggested that diagrams can support understanding of the development of cervical cancer from HPV,¹² because some of our interviews were on the telephone we were not able to use a diagram. In addition, it may be useful to identify demographic subgroups who are less accepting of longer intervals so that targeted communication can be developed. Our findings suggested women who were regular screening attenders were more resistant to longer intervals but this needs further investigation. Although we reached a point where no new themes were being identified in the interviews (thematic saturation), the sample was relatively homogeneous with respect to screening history (mainly up to date), age, ethnic group, sexual orientation and HPV awareness. Had we been able to recruit a larger and more diverse sample, we may have identified additional themes and perspectives.

Conclusion

Women showed a range of cognitive and emotional responses to learning about a planned extension to cervical screening intervals. A number of factors influenced whether participants found the change to 5-yearly screening acceptable. These included attachment to the current 3-yearly interval, trust in decision-makers and understanding of the rationale for the change. When women had the opportunity to discuss the changes and were provided with more information they were, for the most part, reassured and more accepting of the interval change. This highlights the importance of good communication.

Twitter Jo Waller @Jo_WallerKCL

Contributors JW and LAVM conceived the study. JB, JW and LAVM designed the study materials. LAVM and JB conducted the interviews. MN and LAVM analysed the data and drafted the paper. All authors contributed to the final version of the manuscript. LAVM is guarantor for the work.

Funding This study was commissioned by Public Health England (PHE, no grant number was given). MN and LAVM were funded by PHE. JB and JW are funded by Cancer Research UK (grant numbers: C7492/A17219 and C8162/A25356).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by King's College London ethics committee (LRS-19/20-19298 and MOD-20/21-19298).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Data are available from the corresponding author on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Martin Nemeč <http://orcid.org/0000-0003-4228-661X>

Jo Waller <http://orcid.org/0000-0003-4025-9132>

Laura A V Marlow <http://orcid.org/0000-0003-1709-2397>

REFERENCES

- 1 Stubbs R. Significant landmark as primary HPV screening is offered across England, 2020. Available: <https://phescreening.blog.gov.uk/2020/01/23/significant-landmark-as-primary-hpv-screening-is-offered-across-england/> [Accessed 20 Oct 2021].
- 2 Koliopoulos G, Nyaga VN, Santesso N, *et al*. Cytology versus HPV testing for cervical cancer screening in the general population. *Cochrane Database Syst Rev* 2017;8:CD008587.
- 3 Australian Government Department of Health. Changes to the National cervical screening program, 2020. Available: <https://www.health.gov.au/initiatives-and-programs/national-cervical-screening-program/providing-cervical-screening/changes-to-the-national-cervical-screening-program> [Accessed 20 Oct 2021].
- 4 Polman NJ, Snijders PJF, Kenter GG, *et al*. HPV-based cervical screening: rationale, expectations and future perspectives of the new Dutch screening programme. *Prev Med* 2019;119:108–17.
- 5 World Health Organization. *WHO Handbook for Guideline development*. 2nd edn, 2014. <https://apps.who.int/iris/handle/10665/145714>
- 6 Obermair HM, Dodd RH, Bonner C, *et al*. 'It has saved thousands of lives, so why change it?' Content analysis of objections to cervical screening programme changes in Australia. *BMJ Open* 2018;8:e019171.
- 7 Oglivie GS, Smith LW, van Niekerk D, *et al*. Correlates of women's intentions to be screened for human papillomavirus for cervical cancer screening with an extended interval. *BMC Public Health* 2016;16:213.
- 8 Silver MI, Rositch AF, Burke AE, *et al*. Patient concerns about human papillomavirus testing and 5-year intervals in routine cervical cancer screening. *Obstet Gynecol* 2015;125:317–29.
- 9 Cancer Institute NSW. Renewal of the National cervical screening program. Available: <https://www.cancer.nsw.gov.au/prevention-and-screening/screening-and-early-detection/cervical-screening/information-for-health-professionals/renewal-of-the-national-cervical-screening-program> [Accessed 20 Oct 2021].
- 10 Dodd RH, Obermair HM, McCaffery KJ. A thematic analysis of attitudes toward changes to cervical screening in Australia. *JMIR Cancer* 2019;5:e12307.
- 11 Dodd RH, Nickel B, Smith MA, *et al*. Getting the timing right: women's views on the best time to announce changes to cancer screening policy recommendations. *Prev Med Rep* 2020;20:101268.
- 12 Hill E, Nemeč M, Marlow L, *et al*. Maximising the acceptability of extended time intervals between screens in the NHS cervical screening programme: an online experimental study. *J Med Screen* 2021;28:333–40.
- 13 Sekhon M, Cartwright M, Francis JJ. Acceptability of healthcare interventions: an overview of reviews and development of a theoretical framework. *BMC Health Serv Res* 2017;17:88.
- 14 Sekhon M, Cartwright M, Francis JJ. Acceptability of health care interventions: A theoretical framework and proposed research agenda. *Br J Health Psychol* 2018;23:519–31.



- 15 Rebolj M, Rimmer J, Denton K, *et al.* Primary cervical screening with high risk human papillomavirus testing: observational study. *BMJ* 2019;364:l240.
- 16 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349–57.
- 17 A V Marlow L, Nemeč M, Barnes J, *et al.* Testing key messages about extending cervical screening intervals. *Patient Educ Couns* 2022. doi:10.1016/j.pec.2022.04.006. [Epub ahead of print: 16 Apr 2022].
- 18 Ritchie J, Lewis J, Nicholls CM, *et al.* *Qualitative research practice: a guide for social science students and researchers*. Sage, 2013.
- 19 Gale NK, Heath G, Cameron E, *et al.* Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol* 2013;13:117.
- 20 Francis JJ, Johnston M, Robertson C, *et al.* What is an adequate sample size? Operationalising data saturation for theory-based interview studies. *Psychol Health* 2010;25:1229–45.
- 21 Obermair HM, McCaffery KJ, Dodd RH. “A Pap smear saved my life”: Personal experiences of cervical abnormalities shape attitudes to cervical screening renewal. *J Med Screen* 2020;27:223–6.

Appendix 1: Example quotes for each theme**Acceptance of existing screening intervals***General acceptance*

“Ah, I suppose it's alright. I wouldn't mind if it was a little bit longer, but I suppose it's alright.” [P11; 30-39 years; overdue]

“I think I'd rather go more often than not, if that makes sense? ... So, yeah, three years seems like a decent amount of time to leave it.” [P1; 30-39 years; up-to-date]

Prevention and early detection

“I think that prevention is really important, because if you can prevent or if you can catch something at the early stages, it's better than waiting.” [P7; 30-39 years; up-to-date]

“If you can prevent it, do you know what I mean, it's one of the only cancers that you are able to prevent if you catch it early. You don't have routine tests for any other type of cancer really” [P8; 25-29 years; up-to-date]

“I don't actually know how many-- What the odds are of getting cervical cancer is. I have no idea of like, if leaving it too late, what would happen. So, all of that is, yeah. I guess it's lack of information that doesn't make you-- It's not making me move any faster, because I actually don't know, if I don't check it tomorrow, what's the outcome of that? Or what could be-- Yeah. I don't know.” [P15; 40-49 years; overdue]

Cervical cancer symptoms

“If cancer of, you know, cervical cancer isn't caught early on when there's probably no symptoms, then it can be quite devastating. So, I think it's quite, it's very vital, really, that they have the tests, um, for people to try and pick up early. Yeah.” [P4; 30-39 years; up-to-date]

“It feels like it's something that you can't be proactive about until it gets to a point where there's symptoms and presumably when there's symptoms, then some things have fairly significantly developed.” [P13; 40-49 years, up-to-date]

Timeline of cervical cancer development

“When I think about cancer, I feel like it's something that's potentially done, it's quite quickly and can be life threatening...” [P13; 40-49 years, up-to-date]

Acceptance of 5-yearly intervals*General acceptance*

“Um, so for me, if I had to wait five years, I guess I'd have it in the back of my mind of, oh, I wonder if everything is fine and normal.” [P18; 30-39 years; up-to-date]

"Oh my God, that's terrible. Awful. I'm shocked. 5 years is really bad...I feel like, I would really be unhappy if I had to wait every five years. That would make me feel quite ill. I'd be upset with that.' [P3; 30-39 years; up-to-date]

"That makes me quite nervous. That seems like quite a, quite a long time. That would mean, you know, between my first one and my next one, that would be from 25 to 30. I'd be 30 the next time I had one and that, to me, makes me quite nervous.' [P6; 25-29 years; up-to-date]

"Because once you tell some people they're gonna be like, 'Hang on a minute. That's too long. Five years is far too long for me. I can't go without a screening.' And if they're not told anything on top of that, then you can go away really worried and quite sort of scared about the idea of it." [P20; 30-39 years; up-to-date]

Contradicting existing beliefs

"... if there's no symptoms until this extensive cell change. Then that feels like the five years I've got no recourse to any kind of medical support or help or intervention. Because I've got absolutely no idea what's going on." [P13; 40-49 years; up-to-date]

"...it's just not something I would ever want to leave that long... you could go a few years and be okay and then all of a sudden you've got that strain of the virus again, and then that's developing into abnormal cells. And still, you've got another two years before the next test. So that could rapidly spread or, you know, rapidly turn more severe." [P8; 25-29 years; up-to-date]

"Um, well, that sounds okay if you're testing earlier. But personally, I still think that five years is too long, even though you're testing earlier. It still could develop, couldn't it? So, I'd say five years is too long." [P17; 40-49 years; up-to-date]

'Yeah, but then, in between-- Say there, say there were abnormalities, but the test didn't detect that, probably because they took very small amount of sample or whatever. And the test didn't detect the abnormalities. And then there are five years now. So, in that-- She has to wait five years before she gets the next appointment, and then next appointment, they will discover that it has already happened, and the cells are already abnormal. So, in a case like this, what are they going to do?" [P14; 30-39 years; overdue]

Recognition of the benefits

"... if you go, and then you know what you're looking for, the HPV, and then they will say, 'Okay, no anomalies, you come back in five years,' you're gonna trust that you actually have no problems. So, it is actually nice to know that I don't have to keep on going ..." [P22; 40-49 years; overdue]

"The less you go, the better it is for an embarrassing-- From an embarrassment point of view..." [P5; 30-39 years; up-to-date]

"Right, so if they're normal [the cells] then it would be every five years instead of three. Okay. Yeah, I feel like it could, yeah, I guess it could make you feel quite reassured in the sense of, you know, your results are quite-- Well, they're normal, and so we're going every five years instead of three, could make some women feel as though that's like less pressure off them. And they're able to take that away from, you know, their health worries." [P9; 25-29 years; up-to-date]

Impact on screening attendance

"I think-- As far as I can remember it used to be five years [the screening interval] and it's quite easy to forget after that length of time to actually go to your appointment. I know they send you letters and stuff, but people move, don't they and what have you? So, yeah, it can be-- If it's more frequent then you're more likely to remember to go." [P1; 30-39 years; up-to-date]

"I think it [the interval increase] would just be, it would just increase my, not awareness of it, but increase my thinking, you know, I'd be focusing more on that. You know, every three years seems to be just nice at the moment. But any more than that and I'd be thinking, you know, I've got another year until I have my smear test and things like that." [P12; 30-39 years; up-to-date]

The process of accepting a longer interval

Awareness of HPV primary screening

"If the current plan is working, in terms of its reducing the incidents of cancer or deaths from cancer or whatever, why would we be changing it?" [P13; 40-49 years; up-to-date]

"Right, so it's switching to five years for the HPV. But will the smears still be getting done as well or not? ... Yeah, I mean, I think when you first said that I thought you meant they were going to be doing the HPV test alongside the smear." [P8; 25-29 years; up-to-date]

"Okay, um, another thing-- Is this because a majority of the female population have had their HPV vaccine, so is that why they want to, is that why they think that if they've had a vaccine they will, you know, have a positive result with HPV? Which means that if they're already vaccinated and they have their cervical screening, of course, there'll be less risk of them developing abnormal cells, that's why they want to move it to five years?" [P7; 30-39 years; up-to-date]

"Okay. But would the test be exactly, would the test be exactly the same like they do currently? I will want to know whether the test they are going to do, the fact that it's changing, would it be slightly different or would it be the same? Or will it take longer for them to give me a response or if it is abnormal, what would be their steps?" [P14; 30-39 years; overdue]

Understanding the rationale

"I probably-- I think I'd need information on how-- So if you get the HPV, if you get that virus, how long it takes for that to turn into abnormal cells or potentially, I'd need to know the time frames to figure out whether I think five years is a good time frame. Because obviously, I said before the three years I thought was too long because something could happen in the first few months after a smear test and then you've got to wait years. But it could actually take years for the virus to turn into abnormal cells. So, I don't know the time frames." [P4; 30-39 years; up-to-date]

"So probably like I've mentioned, how quickly the stages progress and if that time period is, like, safe to be able to detect something from, like, you know, incubation until, like, actually getting cancer." [P6; 25-29 years; up-to-date]

"The main thing is, just making sure it's effective. Yeah." [P10; 30-39 years; up-to-date]

"Yeah, so-- So people that have abnormal cells, do they all have HPV, I assume?" [P4; 30-39 years; up-to-date]

"I'd want to be sure. I'd want to know. I want to know a bit more about, like, why they think that it's sooner...I want to know the theories behind why they think that five years is acceptable with obviously the testing.' [P3; 30-39 years; up-to-date]

"Yes, I mean if they manage to catch it in the earlier stage, that would be really good, yeah.' [P11; 30-39 years; overdue]

"I feel much comfortable knowing the information I've got now. Especially when, you know, the time it takes to develop. Yeah, it makes complete sense. And yeah, I'd be totally comfortable with that." [P12; 30-39 years; up-to-date]

"Oh okay. I mean, when you-- When I have the kind of like time frame, I feel much better about it, you know? And I think that's probably like what's missing from a lot of, yeah, the way I'm thinking about it and also, it seems like from what you're saying that the type of cancer or the type of cells that you're kind of screening for in a smear, you know, it's very different from other types of cancer. And I guess, like, in my head, all types of cancer get kind of lumped into one type, and I guess that's where a lot of the misunderstandings happen. I think that's quite reassuring. I'd still look a little bit suspicious but, yeah." [P10; 30-39 years; up-to-date]

"...at the end of the day , if there's a way for them to directly test for the HPV and, you know, you've got the stage one , then I can understand why would be every five years , especially if your results come back as normal , then yes , that would make sense. Yeah." [P9; 25-29 years; up-to-date]

"That kind of explains quite a bit, actually. Which I didn't even know. So that kind of makes things a lot more clear around the whole process of it and obviously, there is a step before... Because if you're told about from the beginning and the abnormalities stage and then later down the line, if you do develop cervical cancer. So, to know about that as well, that's very useful. And obviously understanding the process now, why they do have longer in between screenings, is because they are testing for the infection first so that can eradicate it later down the line. So that's really useful to know." [P20; 30-39 years; up-to-date]

"And I suppose I can't get away from this idea that if you have a screening, it's-- And you get a clear result, it's clear at that point. What is it that then happens that makes cells start to change? And what happens-- I don't know. What happens then between that point and five years later, or four and a half years, or four years later? What happens to those cells?" [P13; 40-49 years; up-to-date]

Influencing factors

Perceptions of personal risk

"Um, I don't know, because I had these noncancerous cells which I had removed, if I was given that option of, "Okay, everything's clear, you're all alright, we could see you in the next four or five years," I'd still ask them, you know, "can I still be seen in three years?" ' [P19; 30-39 years; up-to-date]

"I think I'd feel a bit more-- I'd feel a bit nervous about that, just for the sake that, I mean, because I've had the experience of having the cell abnormalities sort of thing. You know, if I had been fine all throughout then it'd be fine. I wouldn't think twice about it. But yeah, I wouldn't want it to be any more than three years." [P12; 30-39 years; up-to-date]

“And um, you know, I'm married, so obviously, if I didn't have the HPV and then I went for my test and they say, “Well you still haven't got it”, well, the chances are, you know, unless I do something naughty that I'm probably not going to get it. Is that-- Am I thinking that right?” [P1; 30-39 years; up-to-date]

“Um, I guess it seems like a bit too long, but that opinion might be coloured by the fact that my mum did have to have a hysterectomy because she had a lot of pre-cancerous cells” [P18; 30-39 years; up-to-date]

Age as a risk factor

“That means if I want one, I have to go private, but I shouldn't have to go private for something so, you know, normal and so, you know, regular.” [P2; 25-29 years; up-to-date]

“Um, I don't know, would there be like a procedure in place where women would be able to request having it earlier? Or would that just be the standard every five years, maybe?” [P9; 25-29 years; up-to-date]

A personalised approach

“And I think, I don't-- What I'm saying is I don't think everyone should be bucketed into those three years. I think if you're over a certain age maybe, then get it more often. If you are, um, in a bucket, there's higher risk. Then you should be getting it more often. I don't think it's fair to say, ‘Everyone's either three years or every year.’ That just doesn't really work... So, it just, as if, you know, with an assumption that a 20-- A healthy 27, 28-year-old female. You know, it just, I don't know, it just seems silly to not, to wait five years.” [P2; 25-29 years; up-to-date]

Trust and empowerment

“Our health service is in such an almighty state that-- I mean, people tell me all sorts of absolute nonsense in the past, that I just kind of think, well, I'm not, I'm just not convinced by it.” [P13; 40-49 years; up-to-date]

“I know we have a very, very good health care system despite what gets said in the media sometimes. But I, I suppose I trust that we do it [screening] for three years for a reason.” [P21; 30-39 years; overdue]

“Um, yeah, more information sent to us would be ideal. I'd also like to see, um, in a few years obviously, statistics from 2019. Um, obviously they'd need, I don't know, 5, 10 years to see if it has actually changed anything.” [P5; 30-39 years; up-to-date]

The value of women's health

“I think in general and sort in medical science, um, it does feel like women are sort of getting like, well, maybe not so much anymore, but it sort of feels that women can get left behind a little bit.” [P1; 30-39 years; up-to-date]

“It doesn't sound-- It doesn't sound like they're taking into consideration the health of people; it sounds more like they're putting costs first.” [P7; 30-39 years; up-to-date]

Feeling powerless and frustrated

“I still-- I mean, if a decision was taken to every five years, then you'd have to accept it...” [P21; 30-39 years; overdue]

“I think three years is kind of too long, so if they made it even longer. Yeah. I don't think I'd be, you know, I'd probably have to deal with it...” [P4; 30-39 years; up-to-date]

Women's opinions on the length of time between cervical screenings (interview study)

TOPIC GUIDE

[START RECORDING]

1. INTRODUCTION (5 mins)

- Intro to researcher
- Acknowledgement of the current pandemic situation
- Study topic and funder, aims/objectives of study
- Explanation of confidentiality/anonymity
- Explanation of recording, duration of interview, nature of discussion, outputs/audiences for findings, data security and rights.
- Reminder of £50 payment, Saros will pay this within a week
- Go through consent process (reading out points from information sheet – should have been shared in advance so they had time to read and consider it)
- Ask whether they have any questions
- Ask whether they are happy to continue and collect consent

[START NEW RECORDING]

2. UNDERSTANDING OF AND FEELINGS REGARDING CERVICAL SCREENING (5 mins)

- Understanding of process/purpose of cervical screening
- Broad feelings/sentiments regarding cervical screening programme and smear tests

3. ACCEPTABILITY OF CURRENT SCREENING INTERVAL (10 mins)

- Awareness of current screening interval

Script: At the moment, the screening interval is 3 years for women in your age-group: from 25-49 years.

- Views re: current 3-year frequency in terms of:
 - How it makes them feel (affective attitude)
 - Amount of effort or how easy/difficult it is to be screened 3-yearly (burden)
 - How effective it is (perceived effectiveness)
Prompt: could it be shorter or longer?
 - Fit with personal values (ethicality)
 - Does it make sense; why they think the interval is 3 years (intervention coherence)
 - Any disadvantages to being screened 3-yearly (opportunity costs)
 - Confidence in ability to get screened every 3 years (self-efficacy)

3. ACCEPTABILITY OF NEW SCREENING INTERVAL (15 mins)

Script: We've talked about the 3 year screening interval for women aged 25-49 years. From next year, the screening programme is going to change so that women with normal screening results will be invited for their next screen in 5 years instead of 3.

- Initial reactions
 - Thoughts and feelings
 - Questions and concerns – what would they want to know about the change

Script: I am now going to give you a bit more background to help explain why the interval is changing:

There are three main steps in the development of cervical cancer. The first is infection with HPV, which is a virus. If the immune system doesn't clear the virus, it can cause the second step, abnormal cells. And thirdly, if the abnormal cells aren't treated, they can turn into cancer. Cervical screening

Women's opinions on the length of time between cervical screenings (interview study)

used to look for abnormal cells (the second step), but in 2019, cervical screening changed so samples are now tested for the HPV virus. Finding HPV picks up the first step in that process, at a really early stage. This means it is now considered safe to screen all women every 5 years.

[Check in] How this information influences their initial thoughts

IF NEEDED: Most people will get the virus at some point in their life. It is spread through close skin to skin contact during any type of sexual activity with a man or woman. HPV can stay in the body for many years. It can stay at very low or undetectable levels and not cause any problems.

- Now that you have listened to this information...

Views re: new 5-year frequency in terms of:

- How it makes them feel (affective attitude) Does information change their initial reactions
- Amount of effort or how easy/difficult it is to be screened 5-yearly (burden)
- How effective it is (perceived effectiveness)
Prompt: will 5-yearly screening find the women who need further tests and checks?
- Fit with personal values (ethicality)
- Does it make sense (intervention coherence)
- What do they have to give up to be able to get screened 5-yearly (opportunity costs)
Prompt: What would you feel you were missing out on if you had 5-yearly screening instead of 3-yearly?
- Confidence in ability to get screened every 5 years (self-efficacy)
- What else do you need to know?

4. RESPONSES TO KEY MESSAGES (15 mins)

Script: I am now going to read six messages that we think help to further explain the change in screening intervals. After each one I am going to ask what you think and whether you find the message helpful or not. You may want to rephrase the message in a way that makes sense to you, or say if something is confusing.

OK, ready (check), so the first message is...

1. If a woman has a negative HPV result, it's very rare for her to develop abnormal cells in less than 5 years [Prompt response]

2. If a test does not indicate (show) you have a HPV infection, it is safe to wait five years between tests. [Prompt response]

3. An HPV test is more accurate than the old smear test. [Prompt response]

4. Women aged 50-64 years already have cervical screening at 5-yearly intervals [Prompt response]

5. Screening more often than needed can lead to unnecessary treatment and worry [Prompt response]

6. Having screening 3 years after a negative HPV result is too soon for any problems to have developed. [Prompt response]

Thanks you, can you think of any other messages you would want to have?

5. CONCLUSION (5 mins)

- Thank participant for their time.
- Reiterate confidentiality, anonymity, right to withdraw from participation.
- Any further questions – contact research team by email. Signpost to support.