

Rapid Evidence Review: COVID-19 and Nurturing Care

Supplementary file 2: Summary of included studies

Responsive caregiving: Caregiver stress and mental health

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Abdelrahman, M. K., et al. Impact of social distancing on the mental health of parents and children in Qatar.	Sept-20	Qatar	N = 308 Parents of children <18 years	Cross-sectional	This online survey, conducted between June 11 and July 10, 2020, explores parents' mental health and coping strategies during COVID-19.	Results showed a significant positive correlation between social distancing practices and parents' activities with children and their coping strategies, suggesting that staying at home for an extended period and not going outside unless necessary motivate parents to develop activities for the children,
Adibelli, D., and A. Sumen. The Effect of the Coronavirus (Covid-19) Pandemic on Health-Related Quality of Life in Children.	Oct-20	Turkey	N = 597 Parents of children ages 7-13 years	Cross-sectional	The online survey, conducted between March 30 and April 20, 2020, explored the relationship between parents' mental health and children's quality of life during the pandemic.	45% of parent respondents reported that staying at home due to the pandemic negatively affected their mental health, and 17% reported a positive affect on their mental health. Quality of life scores (i.e., emotional well-being, family, friends) were lower among children of the parents who were fear/anxiety about coronavirus.
ARNEC. A survey on Perspectives on the impact of COVID-19 on young children and early childhood development in the Asia-Pacific region.	Jun-20	Multiple N=30	N = 684 ECD Stakeholders	Cross-sectional	The survey, conducted between 17 April 2020 and 16 May 2020, explored perspectives of the impact of COVID-19 on families and young children in Asia Pacific.	More than 50% of ECD respondents reported that families are under pandemic induced stress and facing challenges in coping under the heightened stress. Income losses, school/ECD centre closures and prolonged home stay are top reasons for family stress.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Asbury, K., et al. How is COVID-19 Affecting the Mental Health of Children with Special Educational Needs and Disabilities and Their Families?	Jul-20	UK	<i>N</i> = 241 Parents of children ages 5-18 with Special Educational Needs and Disabilities	Qualitative	In an online survey conducted between March 22-April 1, 2020, participants were asked to answer the following free response question: Please describe in your own words how the coronavirus outbreak is affecting your mental health and your child's mental health.	Both parents and children appear to be experiencing loss, worry and changes in mood and behaviour as a result of the rapid social changes that have occurred. Some parents reported feeling overwhelmed and described the impact of child understanding and awareness. Finally, a minority of parents reported that COVID-19 has had little impact on mental health in their family or has even led to improvements.
Brown, S. M., et al. Stress and parenting during the global COVID-19 pandemic.	Aug-20	USA	<i>N</i> = 183 Parents of children <18 years	Cross-sectional	An online survey, conducted from April 21 to May 9, 2020, to examine the relationships among COVID-19 risk factors, mental health risk factors, protective factors, parental perceived stress, and child abuse potential.	Greater COVID-19 related stressors and high anxiety and depressive symptoms are associated with higher parental perceived stress. Receipt of financial assistance and high anxiety and depressive symptoms are associated with higher child abuse potential. Conversely, greater parental support and perceived control during the pandemic are associated with lower perceived stress and child abuse potential. Results indicate racial and ethnic differences in COVID-19 related stressors. 10% reported a positive change: having more time with family.
Cameron, E. E., et al. Maternal Psychological Distress & Mental Health Service Use during the COVID-19 Pandemic.	Jul-20	Primarily Canada	<i>N</i> = 641 Mothers of children ≤8 years, including expectant mothers	Cross-sectional	An online survey, conducted between April 14 and April 28, 2020, described rates of maternal depressive and anxiety symptoms during the COVID-19 pandemic, and identified	In the full sample, 41.51% of mothers met the clinical cut-off scores of ≥ 13 for EPDS and ≥ 16 for CESD indicating depression. Over one-third of mothers (38.62%) indicated current financial strain.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
					risk and protective factors.	
Carroll, N., et al. The Impact of COVID-19 on Health Behavior, Stress, Financial and Food Security among Middle to High Income Canadian Families with Young Children.	Aug-20	Canada	N = 254 Parents of children ages 18 months to 5 years	Cross-sectional	An online survey administered between 20 April 2020 to 15 May 2020 to explore impact of COVID-19 on physical activity, screen time, and sleep among parents and children, as well as stress, financial and food insecurity of families.	Parents reported moderately high levels of stress. Several parents reported coping strategies they use to manage stress and life changes due to COVID-19 restrictions including being more mindful and focusing on the positive aspects of having more time together as a family, taking prescription medication or seeking therapy, and following their pre-COVID-19 routines
Ceulemans, M., et al. Mental Health Status of Pregnant and Breastfeeding Women during the COVID-19 Pandemic: A Call for Action.	Jul-20	Belgium	N = 5866 Pregnant and breastfeeding mothers	Cross-sectional	Online survey, limited details on sampling and recruitment, explored maternal mental health status using the Edinburgh Depression Scale and the Generalized Anxiety 7-item scale	The prevalence of self-reported major depressive symptoms (EDS ≥ 13) in pregnancy (25.3%) and postpartum (23.6%) was higher compared to estimates obtained prior to the pandemic. Generalized anxiety during the lockdown, with more than 40% of the women surveyed scoring ≥ 5 on the EDS subscale of anxiety. In addition, 14% met the criteria for high anxiety on the GAD-7 (≥ 10).
Chu, K., et al. Parenting Under Pressure	2020 <i>Preprint</i>	USA	N = 56 Parents of children ages 6-17 years	Mixed methods	Thematic analysis to identify themes in parents' and children's reflective writing about COVID-19, conducted between April to May 2020.	The most common themes identified from parent reports surrounded concerns about the impact of COVID-19 on their children, health concerns for others, and the stressful balancing act of parenting, schooling, and working from home. In addition, many parents reported gratitude for what they had, and reflected on the upsides of the pandemic for family relations.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Chung, G., et al. Associations Between Work-Family Balance, Parenting Stress, and Marital Conflicts During COVID-19 Pandemic in Singapore.	Jun-20 <i>Preprint</i>	Singapore	N = 258 Parents of children <12 years	Cross-sectional	An online survey disseminated from April 22, 2020, to May 5, 2020 explored work, family life and parenting during the pandemic.	Results indicated three profiles of Work-Family Balance (WFB) and social support: Profile 1 (Poor, n = 38, 19%); Profile 2 (Moderate, n = 76, 38%); and Profile 3 (Strong, n = 87, 43%). Parents' age, race, education, family income, any domestic helpers, and number of caregivers in the household did not predict membership in any of the profiles. Mothers were more likely than fathers to be in the Moderate and Poor profiles. Parents with older children were less likely to be in Profile 1 and 2. Parents who reported a greater impact of COVID-19 and/or with more children in the household were more likely to be in Profile 2 than in Profile 3.
Davenport, M. H., et al. Moms Are Not OK: COVID-19 and Maternal Mental Health.	Jun-20	Mostly USA and Canada	N = 900 Expectant and mothers of children ≤12 months	Cross-sectional	An online survey, conducted between April 14 and May 8, 2020, explored the COVID-19 pandemic on the mental health and physical activity of pregnant and postpartum women.	An EPDS score >13 was self-identified in 40.7% of participants. Moderate to high anxiety (STAI-state score >40) was identified in 29% of women before the pandemic vs. 72% of women currently.
Dhiman, S., et al. Impact of COVID-19 outbreak on mental health and perceived strain among caregivers tending children with special needs.	Oct-20	India	N = 264 caregivers of children with special needs	Cross-sectional	Caregivers completed an online survey from April 29, 2020 to May 22, 2020 that assessed demographics, use and perspective on tele-rehabilitation, homecare therapy, caregiver's strain and mental health	Prevalence of depression, anxiety and stress symptoms were 62.5%, 20.5% and 36.4%, respectively. A significant difference in caregiver strain was observed during the outbreak and pre-outbreak, as measured retrospectively. Caregivers not using tele-rehabilitation along with a perception of it being a poor medium for rehabilitation were at greater risks for poor mental health, while negative perception on homecare therapy were strongly associated with higher psychological symptoms and strain. Working from home was a risk factor for poor mental health. Caregivers of children

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
						older than 6 were less prone for depression, anxiety and stress symptoms.
Di Giorgio, E., et al. The Interplay between Mothers' and Children Behavioral and Psychological Factors during COVID-19: An Italian Study.	Aug-20	Italy	N = 245 Mothers of children ages 2 to 5 years	Cross-sectional	An online survey, conducted between April 1-9, 2020 to explore both mothers' habits, behaviors, and emotions and those of their children during the quarantine, and retrospectively, before the national lockdown (i.e., in late February)	Most respondents (66.5%) had to stop working or to start working from home. A general worsening of sleep quality and distortion of time experience in both mothers and children, as well as increasing emotional symptoms and self-regulation difficulties in children, was observed. Moreover, even when the interplay between the behavioral and psychological factors was investigated, the factor that seems to mostly impact both mothers' and children's psychological well-being was their sleep quality.
Dib, S., et al. Maternal Mental Health and Coping during the COVID-19 Lockdown in the UK: Data from the COVID-19 New Mum Study.	Sep-20	United Kingdom	N = 1329 Mothers of children ≤12 months	Cross-sectional	An online survey May 27, 2020, and will remain open until at least December 31, 2020, explores predictors of maternal mental health and coping during COVID-19.	Support with her own health (95% confidence interval [CI] 0.004–0.235), contacting infant support groups (95% CI –0.003 to 0.252), and higher gestational age of the infant (95% CI 0.000–0.063) predicted better mental health. Support with her own health and more equal division of household chores were associated with better coping.
Drouin, M., et al. How Parents and Their Children Used Social Media and Technology at the Beginning of the COVID-19 Pandemic and Associations with Anxiety.	Jul-20	USA	N = 260 Parents of children ≤18 years	Cross-sectional	Online survey data, collected from March 20-25, 2020, explored children's use of social media and technology during the COVID-19 pandemic and its association with anxiety.	92% of parents and 82% of children had increased their technology use to connect with others. Bivariate correlations revealed significant positive associations between parent anxiety and parent technology use, social media use, and checking messages/ news, but not connecting via video or taking time for self. Those with greater financial preparedness and older children were able to take time for themselves more frequently, and those with older children connected with others via video less frequently than those with younger children.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Evans, S., et al. From “It Has Stopped Our Lives” to “Spending More Time Together Has Strengthened Bonds”: The Varied Experiences of Australian Families During COVID-19.	Oct-20	Australia	N=2130 Parents of children 0-18 years	Qualitative	During an online survey conducted between April 8 and April 28, 2020, participants were asked to respond via an open-ended question about how COVID-19 had impacted their family.	Six themes were derived from the data, including “Boredom, depression and suicide: A spectrum of emotion,” “Families are missing the things that keep them healthy,” “Changing family relationships: The push pull of intimacy,” and others. Messages around loss and challenge were predominant, with many reporting mental health difficulties and strained family relationships. However, not all families were negatively impacted by the restrictions, with some reporting positive aspects, including opportunities for strengthening relationships, finding new hobbies, and developing appreciation, gratitude, and tolerance.
Farewell, C. V., et al. A Mixed-Methods Pilot Study of Perinatal Risk and Resilience During COVID-19.	Jul-20	USA	N = 31 Mothers of children ≤6 months and expectant mothers	Mixed methods	Mixed methods study conducted between March and April 2020 explored mental health and well-being during the perinatal period.	New moms shared that social isolation was significantly affecting their postpartum mental health. Positive impacts related to the COVID-19 pandemic, included being able to work from home during pregnancy, increased connection and bonding with their immediate family unit, partner supports in the home to share caregiving responsibilities; and increased access to remote postnatal and postpartum care.
Gassman-Pines, A., et al. COVID-19 and Parent-Child Psychological Well-being.	Oct-20	USA	N = 561 Parents/ caregivers of children ages 2 to 7 years	Longitudinal	Daily survey data were collected between February 20 and April 27, 2020, from hourly service workers with a young child in a large US city. A subsample of 561 completed a one-time survey about the effects of the crisis between March 23 and April 26.	Ordered probit models revealed that the frequency of parent-reported daily negative mood increased significantly since the start of the crisis. Many families have experienced hardships during the crisis, including job loss and income loss. Both parents’ and children’s well-being was strongly associated with the number of crisis-related hardships.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Günther-Bel, C., et al. A Mixed-method Study of Individual, Couple, and Parental Functioning During the State-regulated COVID-19 Lockdown in Spain.	Jul-20	Spain	N = 407 Adults including parents	Mixed methods	An online survey conducted between March 24 and April 7, 2020 explored individual and family well-being during the first 3 weeks of state-regulated lockdown.	An online survey conducted between March 24 and April 7, 2020, described perceived improvement as family connection, better communication, emotional expressiveness, teamwork spirit, and balance between individual and shared needs. The overall prevalence of improvement themes exceeded the prevalence of deterioration themes, mainly conflict atmosphere and unbalanced needs (most likely with children with children at home). Deterioration themes were associated with the presence of preschoolers.
Hamadani, J. D., et al. Immediate Impact of Stay-at-Home Orders to Control COVID-19 Transmission on Socioeconomic Conditions, Food Insecurity, Mental Health, and Intimate Partner Violence in Bangladeshi Women and Their Families: An Interrupted Time Series.	Aug-20	Bangladesh	N = 2417 Mothers of children ≤5 years	Quasi-experimental	An interrupted time series was used to compare data collected on income, food security, and mental health a median of 1 year and 2 years before the COVID-19 pandemic to data collected during the lockdown.	Mothers' depression and anxiety symptoms increased during the lockdown. Among women experiencing emotional or moderate physical violence, over half reported it had increased since the lockdown.
Hiraoka, D., and A. Tomoda. Relationship between Parenting Stress and School	Jun-20	Japan	N = 353 Parents of children <18 years	Cross-sectional	An online survey, conducted between 29 and 30 April 2020, administered the Parenting Stress Index –	Parental stress was significantly higher after school closures, compared to parental stress before school closures, based on recall ($t = 4.89$, $p < .01$, $d = 0.12$).

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Closures Due to the COVID-19 Pandemic.					Short Form (PSI-SF) to measure parenting stress during the pandemic.	
Joyce, Kayla M., et al. Changes in Maternal Substance Use During the COVID-19 Pandemic.	2020 <i>Preprint</i>	Canada	<i>N</i> = 508 Mothers of children ≤8 years	Cross-sectional	Online survey data collected April 14th to 28th 2020 to identify associations between mental health and substance use. The EPDS and GAD-7 were completed.	Of participating mothers, 54.9% did not change their substance use, 39.2% increased their use, and 5.9% decreased their use. Findings indicate that mothers with an anxiety disorder or clinically relevant anxiety symptomatology were more likely to report increased substance use during the COVID-19 pandemic.
Kalil, A., et al. Impact of the COVID-19 Crisis on Family Dynamics in Economically Vulnerable Households.	Oct-20	USA	<i>N</i> = 572 Parents of preschool-age child	Cross-sectional	A telephone survey, conducted from May 3 to July 20, 2020, to explore parental mental health and parent-child interaction during the pandemic.	Parental job and income losses are strongly associated with parents' depressive symptoms, stress, diminished sense of hope, and negative interactions with children. Parents who experience a job and income loss, as well as those with just income loss, are more likely to lose their temper or yell at their child. Parents who have job and income loss are more likely to be regular readers with their children.
Lee, S. J., et al. Parenting Activities and the Transition to Home-Based Education During the COVID-19 Pandemic.	Oct-20 <i>Preprint</i>	USA	<i>N</i> = 405 Parents with children <12 years	Mixed methods	An online survey, including open-ended questions, conducted from April 24-25, 2020, examined parental mental health and the transition to home-based learning.	78% of parents said they were educating their child at home due to COVID-19. Two out of every five parents met the PHQ-8 criteria for major depression or severe major depression (40.0%) and the GAD-7 criteria for moderate or severe anxiety (39.9%). Multivariate analyses indicated that, compared to non-depressed parents, parents who met criteria for probable major or severe depression and parenting stress were negatively associated with parents' perceived preparation to educate at home. Compared to parents with minimal or mild anxiety, parents with moderate or severe anxiety reported higher child anxiety. Content analyses of

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
						open-ended questions indicated that school closures were a significant disruption, followed by lack of physical activity, and social isolation.
Limbers, C. A., et al. Physical Activity Moderates the Association between Parenting Stress and Quality of Life in Working Mothers during the COVID-19 pandemic.	Oct-20	USA	<i>N</i> = 200 Full-time working mothers of children ≤5 years	Cross-sectional	Online survey of mothers completed during April 6th and April 13th, 2020, to examine physical activity, parenting stress and quality of life during the pandemic.	Conducted from results showed that greater parenting stress was associated with poorer maternal quality of life (physical health, psychological, social relationships, and environment quality of life), after controlling for socio-demographic variables. Multiple linear regression analysis suggest that moderate intensity physical activity may be one mechanism that attenuates the association between parenting stress and diminished social relationships quality of life during.
Marchetti, D. et al. Parenting-Related Exhaustion During the Italian COVID-19 Lockdown.	Oct-20	Italy	<i>N</i> = 1,226 Parents of children ≤13 years	Cross-sectional study	An online survey administered between April 3 and 14 2020 to explore emotional exhaustion, parental resilience, social connections, and psychological distress during the lockdown.	Results indicate that more than 80% of parents reported high levels of psychological distress, and nearly one in five (17%) reported significant parenting-related exhaustion, with mothers more severely affected. Multiple regression analyses showed that greater parenting-related exhaustion was predicted by psychological distress, lower parental resilience, fewer perceived social connections, and being single, as well as having a child with special needs, having a large number of children, and having younger children.
Mazza, C., et al. How Personality Relates to Distress in Parents during the Covid-19 Lockdown: The Mediating Role of Child's Emotional and Behavioral Difficulties and the Moderating	Aug-20	Italy	<i>N</i> = 833 Parents of children ages 3–13 years	Cross-sectional	An online survey was administered to 833 participants from 3 to 15 April 2020 to investigate psychological distress of parents during the lockdown and contributing factors.	For parents living only with child(ren), high levels of psychological distress depended exclusively on their levels of neuroticism. For parents living with at least one other person in addition to child(ren), distress levels were also mediated by child behavioral and emotional difficulties. Motherhood emerged as a significant factor contributing to greater distress. Furthermore, parent psychological distress decreased in line with increased child age.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Effect of Living with Other People.						
McRae, C. S., et al. Parents' Distress and Poor Parenting during COVID-19: The Buffering Effects of Partner Support and Cooperative Coparenting.	2020 <i>Preprint</i>	New Zealand	<i>N</i> = 362 Parents of children ages 4 to 5 years	Longitudinal	An online survey conducted from March 26 to April 28, 2020 assessed harsh parenting, warm/responsive parenting, and parent-child relationship quality among a cohort of parents who had already been surveyed pre-pandemic.	Parents who experienced greater distress during the lockdown reported harsher parenting, less warm/responsive parenting and lower parent-child relationship quality. Greater psychological distress predicted residual increases in harsh parenting when parents perceived low partner support, but the detrimental effects of parents' distress was buffered when parents perceived high levels of partner support.
Nolvi, S., et al. Parental Depressive and Anxiety Symptoms as a Response to the COVID-19 Pandemic: A Birth Cohort Follow-up Study.	2020 <i>Preprint</i>	Finland	<i>N</i> = 651 Parents of children ages 5-8 years	Longitudinal	Based on a sample from a longitudinal cohort study, the FinnBrain Birth Cohort Study, data collected from May 4 and June 7, 2020 using the EPDS, Symptom Checklist 90	Higher symptom level was associated with COVID-related stressors, being mother (vs. father), prior symptoms and prior negative life events. Importantly, there was a significant increase in both depressive and anxiety symptoms and in subjects scoring above the screening cut-off of depression from pre-pandemic to pandemic. The increase in symptoms more pronounced in mothers versus fathers.
Oskovi-Kaplan, Z. A., et al. The Effect of COVID-19 Pandemic and Social Restrictions on Depression Rates and Maternal Attachment in Immediate Postpartum	Sep-20	Turkey	<i>N</i> = 223 Mothers, 48 hrs postpartum	Cross-sectional study	The survey was conducted in June 2020. The low-risk term pregnant women who gave birth were given the surveys Edinburgh Postpartum Depression Scale (EPDS) and Maternal Attachment	Conducted in June 2020, the median score obtained from the EPDS was 7 and 15% of the women were determined to have a risk for postpartum depression. The authors suggest the postpartum depression risk during the COVID-19 pandemic was increased when compared with previous reports in Turkey, while it was still lower than the studies reported from other countries.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Women: A Preliminary Study					Inventory (MAI) within 48 h after birth.	
Patrick SW, et al. Well-being of Parents and Children During the COVID-19 Pandemic: A National Survey.	Oct-20	USA	<i>N</i> = 1011 Parents of children <18 years	Cross-sectional	As national online survey conducted from June 5-10, 2020 to determine how the pandemic and mitigation efforts affected the physical and emotional well-being of parents and children.	Since March 2020, 27% of parents reported worsening mental health for themselves, and 14% reported worsening behavioral health for their children. 48% reported loss of regular childcare, 16% reported change in insurance status, and 11% reported worsening food security.
Peltz, J. S., et al. Mediators of the Association between COVID-19-Related Stressors and Parents' Psychological Flexibility and Inflexibility: The Roles of Perceived Sleep Quality and Energy.	Jul-20	USA	<i>N</i> = 1003 Parents of children ages 5 to 18 years	Cross-sectional	Online survey conducted from March 27th to the end of April 2020 to explore COVID-19 related stressors, and	Path analyses suggested that higher levels of health-related stress were associated with lower levels of energy and lower perceived sleep quality, both of which were associated with higher levels of psychological inflexibility. In secondary analyses on the specific dimensions of flexibility and inflexibility, results suggested that lower energy levels indirectly linked health-related stress to lower levels of all 6 dimensions of flexibility and poorer sleep quality indirectly linked health-related stress to higher levels of self-as-content, fusion, and inaction.
Romero, E., et al. Testing the Effects of COVID-19 Confinement in Spanish Children: The Role of Parents' Distress, Emotional Problems and Specific Parenting.	Sep-20	Spain	<i>N</i> = 940 Parents of children ages 3 to 12 years	Cross-sectional	Online survey conducted in April 2020 to examine parenting distress and specific parenting practices (e.g., structured or avoidant parenting).	Data collected in April 2020 show that higher levels of parenting distress were significantly related with lower focused and structured parenting, and higher avoidant parenting. parenting distress was also affected by COVID-related stressors, including close contagion, keeping up with jobs and working at home.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Roos, L. E., et al. Supporting Families to Protect Child Health: Parenting Quality and Household Needs During the COVID-19 Pandemic.	Sep-20	Canada	<i>N</i> = 656 Parents or caregivers of children ages 1-8 years	Cross-sectional	Mixed methods were applied to data collected between April 14th - 28th, 2020 from the “Parenting During the Pandemic” survey. Data collection occurred through REDCap.	Caregiver depression, higher child parity, unmet childcare needs, and relationship distress predicted lower-quality parenting (i.e., more stress, fewer positive behaviors, and more negative behaviors). Caregiver depression was the most significant predictor across every parenting dimension, with analyses indicating medium effect sizes. Qualitative findings highlighted severe strains on parent capacities including managing psychological distress, limited social supports, and too much unstructured time.
Russell, B. S., et al. Initial Challenges of Caregiving During COVID-19: Caregiver Burden, Mental Health, and the Parent–Child Relationship.	Aug-20	USA	<i>N</i> = 420 Parents of children <18 years	Cross-sectional	Online survey conducted April 27–28, 2020s to explore family experiences and coping during the COVID-19 pandemic.	Caregiver burden was positively associated with generalized anxiety, depression. Male caregivers reported higher rates overall than female caregivers. Depression was significantly positively associated with child parent relationship conflict but not child parent relationship closeness. Caregiver burden was not significantly directly associated with child parent relationship conflict or child parent relationship closeness. Generalized anxiety did not significantly predict child parent relationship conflict or child parent relationship closeness.
Spinelli, M., et al. Parenting Stress During the COVID-19 Outbreak: Socioeconomic and Environmental Risk Factors and Implications for Children Emotion Regulation.	Sep-20	Italy	<i>N</i> = 810 Parents of children ages 2–14 years	Cross-sectional	Online survey from April 2nd to 7th, 2020 to examine level of household chaos, parenting stress, parent involvement during COVID-19.	Parents reporting higher levels of stress were less engaged with their children, they were less interested in children emotional well-being, they paid less attention to the child, and in general spent less time with the child. For SES at-risk families, the impact of parenting stress and involvement on children regulation strategies was stronger, with a protective role played by parental involvement on children’s negativity not evident for SES no-risk families. The majority of these families in our sample were characterized by a reduction of their

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
						income due to the loss of the job because of the pandemic.
Spinelli, M., et al. Parents' Stress and Children's Psychological Problems in Families Facing the COVID-19 Outbreak in Italy.	Jul-20	Italy	N = 854 Parents of children ages 2-14 years	Cross-sectional	Online survey conducted from April 2nd to 7th, 2020 to explore perceptions of individual and parent-child dyadic stress, and their children's emotional and behavioral problems.	Parents who reported more difficulties in dealing with quarantine show more stress. This, in turn, increases the children's problems. Living in a more at-risk area, the quality of the home environment, or the relation they have with the pandemic consequences, do not have an effect on families' well-being.
Westrupp, E., et al. Child, Parent, and Family Mental Health and Functioning in Australia During COVID-19: Comparison to Pre-pandemic Data.	Sep-20 <i>Preprint</i>	Australia	N = 2365 mothers and fathers of children 0-18 years	Repeat cross-sectional	Online survey assessing mental health, substance use, couple conflict, parenting, and family functioning during 'stage three' COVID-19 restrictions in April 2020 was compared to pre-pandemic data from four population-based surveys.	Compared to pre-pandemic estimates, during the pandemic period parents reported higher rates of parent mental health symptoms, higher parenting irritability, lower family positive, and higher alcohol consumption. In multivariable analyses, pre-existing financial deprivation and COVID-19 stressors were associated with greater severity in parent and child mental health symptoms, parent emotion dysregulation, parenting irritability, couple conflict and family positive/ negative expressiveness. Parents and children with pre-existing mental health conditions had elevated difficulties during the pandemic across most domains.
Willner, P., et al. Effect of the COVID-19 pandemic on the mental health of carers of people with intellectual disabilities.	Sep-20	UK	n = 107 carers of adults with intellectual disability; n = 100 carers of children <18 years with intellectual	Cross-sectional	An online survey conducted between 28 April and 5 June 2020 estimated whether the extent of mental health problems was greater than would have been seen under pre-pandemic	Relative to carers of children without intellectual disability, carers of both children and adults with intellectual disability had significantly greater levels of a wish fulfilment coping style, defeat/entrapment, anxiety, and depression. Differences were 2-3 times greater than reported in earlier pre-pandemic studies. Positive correlations were found between objective stress scores and all mental health outcomes. Despite

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
			disability; and <i>n</i> = 37 carers of children <18 years without intellectual disability		conditions, by comparing current data to earlier literature. The instrument assessed demographics, coping strategies, social support, anxiety, feelings of defeat/entrapment, and stress.	their greater mental health needs, carers of those with intellectual disability received less social support from a variety of sources.
Xu, Y., et al. Material Hardship and Parenting Stress among Grandparent Kinship Providers during the COVID-19 Pandemic: The Mediating Role of Grandparents' Mental Health.	Aug-20	USA	<i>N</i> = 362 Grandparent caregivers of children < 18 years	Cross-sectional	An online survey conducted in June 2020 to examine material hardship, parenting stress, and grandparents' mental health.	A total of 28.18 % (<i>n</i> = 102) of grandparents agreed that they experienced parenting stress during the COVID-19 pandemic. A comparison between grandparent kinship providers that experienced parenting stress and those that did not found that the latter had a significantly lower score on experiencing material hardship. Similarly, grandparents that did not report parenting stress had significantly better mental health status.
Yang, Y., et al. How did COVID-19 Impact the Lives and Perceived Well-being of Parents? Using the Case of Singapore to Investigate the Mechanisms.	Oct-20 Preprint	Singapore	<i>N</i> = 505 adults caring for children <18 years from 198 families	Cross-sectional	An online survey fielded in June 2020 collected retrospective self-reports on everyday activities, stressors, and well-being of parents and caregivers at three timepoints: before local transmission, after local transmission but before "circuit breaker", and during "circuit breaker".	Results showed significant changes in jobs and income, childcare arrangement, family dynamics, and parents' emotional well-being throughout the three timepoints. Both mothers and fathers reported to worry most about the health and safety of family members and self. Mothers' time spent on housework partially mediated the effect of lockdown on their emotional well-being, and parents' conflict with other adults in the household partially mediated the effects of both pandemic and lockdown on their emotional well-being. The effects of pandemic and lockdown were also moderated by parents' age, education level, and fathers' authoritarian values.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Zamarro, G., and M. Prados. Gender Differences in Couples' Division of Childcare, Work and Mental Health During COVID-19.	Jul-20	USA	<i>N</i> = 3980 adults, including parents of school-age children or children in daycare; 26,052 observations	Longitudinal	Uses data from eight waves of the online <i>Understanding Coronavirus in America Tracking Survey</i> , collected approximately every two weeks from March 10 to July 22, 2020 to understand gender differences within households on the impacts of COVID-19 on childcare provision, employment, working arrangements, and psychological distress levels.	Combining a measure of feelings of anxiety and depression, we find that psychological distress increased significantly early during the crisis, with 49% of mothers showing at least mild symptoms of psychological distress in early April, as compared to 41% of women without school-age children. (In the U.S., schools closed in all states between March 16-24) This new gap in psychological distress observed between mothers and women without school-age children appears to be driven by higher levels of psychological distress among mothers of elementary school-age and younger children.
Zanardo, V., et al. Psychological Impact of COVID-19 Quarantine Measures in Northeastern Italy on Mothers in the Immediate Postpartum Period.	Jun-20	Italy	<i>N</i> = 192 mothers post-partum, specifically mothers of singleton, healthy neonates delivered between March 8 and May 3	Case-control	Case-control study of mothers who gave birth during a COVID-19 quarantine period between March 8 and May 3, 2020, with a matched comparison group of matched postpartum women who delivered in the same period in 2019. Participants completed the Edinburgh Postnatal Depression Scale	A case-control study of mothers who gave birth during a COVID-19 quarantine period between March 8 and May 3, 2020 had significantly higher mean EPDS scores compared with the control group who delivered in the same period in 2019 (8.5 vs 6.34).

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
					(EPDS) on the second day postpartum.	

Responsive caregiving: Fathers' engagement in caregiving and gendered division of labour

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Andrew, Alison, et al. How Are Mothers and Fathers Balancing Work and Family under Lockdown?	May-20	United Kingdom	N = 3519 Parents of children ages 4-15 years	Online time-use diaries	Online survey conducted from 29 April and 15 May 2020 to examine paid work, exercise, personal care, childcare and sleep.	On average, fathers are now doing some childcare during 8 hours of the day, compared with 4 hours in 2014/15. Of parents who were in paid work prior to the lockdown, mothers are one-and-a-half times more likely than fathers to have either lost their job or quit since the lockdown began. In 2014/15, mothers were in paid work at 80% of the rate of fathers; now this is 70% of the fathers' rate. Compared with fathers, mothers are spending less time on paid work but more time on household responsibilities.
Carlson, Daniel L., et al. Changes in Parents' Domestic Labor During the COVID-19 Pandemic.	May-20	USA	N = 1,025 Parents of children ages 6-17 years	Cross-sectional	Survey conducted in late April 2020 to examine the division of domestic labor before and during the pandemic.	Results show an overall increase in domestic responsibilities for mothers, who were already doing most of the household labor, as well as an increase in fathers' contributions. Driven by increases in fathers' time spent on housework and childcare, we find that both mothers and fathers report a general shift toward more egalitarian divisions of household labor.
Craig, L., & Churchill, B. Dual-earner Parent Couples' Work and Care during COVID-19.	Jun-20	Australia	N = 1536 Parents of children <16 years	Cross-sectional	An online survey conducted from 7 May to 30 May 2020 examined changes in parental employment and childcare arrangements,	Before COVID-19, mothers averaged 3.41 daily hours in active care of children compared to 2.21 daily hours for fathers: the gender gap was about 54 per cent. This decreased to 40 per cent during COVID-19 because fathers' hours went up by 64 per cent (to 3.64 hours per day) whereas mothers went up by 50 per cent (to

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
					and their subjective feelings.	5.13 hours per day). Before COVID-19, almost 44 per cent of fathers and 41 per cent of mothers had been 'extremely' or 'somewhat' satisfied with how they divided their time between paid and unpaid work. During the restrictions, the proportions of fathers and mothers across these categories fell to 38 and 30 per cent, respectively.
Del Boca, Daniela, et al. Women's and Men's Work, Housework and Childcare, before and during COVID-19.	Aug-20	Italy	N = 800 Working women, of whom 67% have children of any age	Longitudinal	An online survey conducted in April and July 2019 and April 2020 to examine changes in employment status, hours of work, childcare, income and satisfaction regarding their work and family during the emergency.	Both men and women are spending more time on housework and childcare, but the distribution is unequal: with 68% of women spending more time in housework and only 40% of men. The percentages for childcare are 61% and 51%, respectively. Hence, while most of the burden has fallen on women, the additional childcare is more equally shared than housework. Yet this balance disappears when we consider the amount of time spent on housework: 64% of women and 58% of men increase the amount of housework they do. Another predictor of higher child-related workload is the age of the children: children younger than 10 years old require more time from both working mothers and fathers. Working women with children age 0–5 are those most likely to report it is difficult to balance work and family due to excessive domestic responsibilities.
Farré, Lúdia, et al. How the Covid-19 Lockdown Affected Gender Inequality in Paid and Unpaid Work in Spain.	Jul-20	Spain	N = 3894 adults from opposite-sex households with children (age range not specified)	Cross-sectional	An online survey, conducted between May 5-19, 2020 to explore household distribution of domestic and childcare tasks.	The lockdown increased the volume of childcare done by parents by 25% from 48 to 60 hours a week. Women were still doing most of the childcare during the lockdown, although men seem to have increased their participation. Before the lockdown, women were spending 15 more hours than men in childcare. During

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
						the lockdown, men increased their childcare time by almost 4 hours.
Heggeness, M. Estimating the Immediate Impact of the COVID-19 Shock on Parental Attachment to the Labor Market and the Double Bind of Mothers.	Oct-20	USA	175,969 observations from 62,702 unique parents of school age children	Quasi-experimental	Compares labor market attachment, non-work activity, hours worked, and earnings and wages by gender for those in areas with early school closures and stay-in-place orders to those in areas with delayed or no pandemic closures. Utilizes monthly panel data Current Population Survey for the months of January, February, March, April, and May of 2019 and 2020.	Mothers of school age children who maintained jobs in early closure states were 68.8% percent more likely than mothers in late closure states to not be working. In other words, mothers of school age children in early closure states were much more likely to take leave from work than women in general. There was no significant difference in leave time between fathers from early closure and late closure states.
Hjálmsdóttir, A., and Bjarnadóttir, V. S. 'I Have Turned into a Foreman Here at Home.' Families and Work-Life Balance in Times of Covid-19 in a Gender Equality Paradise.	Sep-20	Iceland	<i>N</i> = 37 Mothers of children (age range not specified)	Qualitative	The study draws on open-ended real-time diary entries, collected from March 26 to April 14, 2020 to explore impact of the COVID-19 pandemic on families and work-life balance.	In most of the cases, both parents primarily or solely worked from home during the time of the study, and most of them were working full-time the whole period, even though some worked reduced hours due to the pandemic. The diary entries of the mothers in the study demonstrate a gendered reality in which they experience burdens that seem to have escalated during the pandemic.
Lyttelton, T. et al. Gender Differences in Telecommuting and Implications for	Jul-20 <i>Preprint</i>	USA	<i>N</i> = 784 Employed parents of	Quasi-experimental	Uses nationally representative time diary data from the 2003-2018 American Time Use	Gender gaps in housework are larger for telecommuters, and, among telecommuters, larger on telecommuting days. Conversely, telecommuting may shrink the gender gap in childcare, particularly among

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Inequality at Home and Work.			children <18 years		Survey (ATUS) and the April and May 2020 COVID Impact Survey to describe the prevalence of parental telecommuting subjective wellbeing.	couples with two full time earners, although childcare more frequently impinges upon mothers' work time. Survey data collected following the March COVID19 stay-at-home orders show that telecommuting mothers more frequently report feelings of anxiety, loneliness and depression than telecommuting fathers.
Mangiavacchi, L. et al. Fathers Matter: Intra-Household Responsibilities and Children's Wellbeing During the Covid-19 Lockdown in Italy.	Jul-20	Italy	N = 3,352 Families with children < 16 years	Cross-sectional	Online survey data collected in April 2020 examines children's wellbeing, and parents' market and household work during the lockdown.	Labor market restrictions imposed during lockdown resulted in increases in fathers' involvement in childcare and homeschooling. Fathers' involvement had a positive association with child emotional wellbeing, but no impact on reading hours and educational progress.
Manzo, L. K. C., and Minello, A. Mothers, Childcare Duties, and Remote Working under COVID-19 Lockdown in Italy: Cultivating Communities of Care.	Jul-20	Italy	N = 20 Mothers of children age <5 years	Qualitative	Conducted virtual interviews from March to May 2020 to explore how the increase in remote working has created unequal parenting duties with respect to gender during the lockdown in Italy.	The results confirm that the reorganization of care during the COVID-19 pandemic has been unbalanced: women, including those working regularly, performed most of the household duties.
Qian, Y. and Fuller, S. COVID-19 and the Gender Employment Gap among Parents of Young Children.	Jul-20	Canada	N = 60,364 parents with a youngest child under 6 N=51,600 parents with a youngest	Quasi-experimental	pooled monthly data from the public use microdata files of Statistics Canada's Labour Force Survey (LFS) from February to May 2020 to examine the gender gap in	Among parents whose youngest child was under 6, there was a small gender employment gap of one percentage point in February; and this gap increased to 3.8 percentage points in March. Controlling for job and demographic characteristics has little impact on results, suggesting that, among parents of young children, the growing gap cannot be attributed simply to differences in the types of jobs mothers and fathers hold or their

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
			child aged 6–12 years		employment immediately before and also during the COVID-19 pandemic.	personal characteristics. The gender employment gap among parents with a child under 6 widened much more during the pandemic among the less educated.
Zamarro, G., and M. Prados. Gender Differences in Couples' Division of Childcare, Work and Mental Health During COVID-19.	Jul-20	USA	<i>N</i> = 3980 adults, including parents of school-age children or children in daycare; 26,052 observations	Longitudinal	Uses data from eight waves of an online survey collected approximately every two weeks from March 10 to July 22, 2020 to understand gender differences within households on the impacts of COVID-19 on childcare provision, employment, working arrangements, and psychological distress levels.	We find that increased need for childcare has put a strain on working parents of both genders, but overall, mothers have been carrying a heavier load on the provision of childcare. In households with children, 44% of women report being the sole provider of care for their children as compared to 14% of men. Even while currently working, women are 27 percentage points more likely to be the only providers of childcare than working men, and 41 percentage points less likely to report that their partner is the only provider of care. Among working parents who are married or living with a partner, women were 17 percentage points more likely than men to become the sole childcare provider during the pandemic.

Responsive caregiving: breastfeeding support

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Brown, A., and N. Shenker. Experiences of Breastfeeding during COVID-19: Lessons for Future Practical	Sep-20	United Kingdom	<i>N</i> = 1219 Mothers of children <2 years	Cross-sectional	Data were collected for 4 weeks during May–June 2020.	41.8% of mothers felt that breastfeeding was protected due to lockdown, but 27.0% of mothers struggled to get support and had numerous barriers stemming from lockdown with some stopped breastfeeding before they were ready. Mothers with a lower education, with more challenging living circumstances and from Black and minority ethnic backgrounds were more likely to find the impact of lockdown challenging and stop breastfeeding. Overall, 7.8% stated they were not supported to have skin to skin, 4.6%

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
and Emotional Support.						were not encouraged to breastfeed as soon as possible after birth, 24.6% were not given information on expressing milk, and 21.2% stated they received no breastfeeding support in hospitals. Overall, 68.7% of those still breastfeeding felt they had much more or a little more time to focus on breastfeeding.
Das, R. COVID-19, Perinatal Mental Health and the Digital Pivot.	Jun-20	United Kingdom	<i>N</i> = 14 Pregnant and new mothers	Qualitative	Participants interviewed online using a semi-structured guide to explore impacts of COVID-19 on mental health and wellbeing.	Infant feeding support, health visits and baby weigh-in sessions were the three services most missed in their in-person formats, and there was, across the board, a near unanimous assertion that these are sorely needed. A sense of feeling cheated of the joys of pregnancy and maternity, and intense self-imposed protective measures around the virus, and sometimes the feelings of not doing enough for an older child in the absence of schooling or childcare.
Saunders, B. and Hogg, S. Babies in Lockdown: Listening to Parent to Build Back Better.	Aug-20	United Kingdom	<i>N</i> = 5794 Currently pregnant and parents of children <2 years	Cross-sectional	Online survey from 29 April to 3 June 2020, to examine impact of COVID-19. The survey included a mix of close- and open-ended questions on experiences for expectant and new parents during lockdown.	For some respondents the lockdown brought advantages, such as increased time at home and the presence of fathers and other co-parents. However, for many the time was one of very considerable stress. Women cited feeling rushed out of hospital, poor follow-up care and poor access to information and support once at home as the main barriers to establishing breastfeeding with their babies. Black or Black British respondents were least likely to have had any contact with breastfeeding support organisations or charities. Younger respondents (<25 years of age) were most likely to cite the following as main concerns, all of which were higher than their older counterparts: their mental health, financial worries, relationship with others, eating a healthy diet, getting reliable pregnancy information and housing issues. Greater numbers of respondents whose income is below £16k reported to have concerns for their emotional and mental health, financial worries, and housing issues.
Spatz, D. L., and E. B. Froh. Birth and	Oct-20	USA	<i>N</i> = 3 New mothers	Qualitative	Virtual interviews conducted in March and April	All three mothers reported social isolation from not being able to go out and spend time with friends, families, or other new mothers. The mothers described how they had planned to spend time with support groups or with

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Breastfeeding in the Hospital Setting during the COVID-19 Pandemic.					2020 to understand the ways in which new families experience pregnancy and lactation.	other new mothers in their social circle but reported sadness of not being able to have those experiences due to social distancing. A great benefit to all the mothers was having their partners at home with them due to social distancing.
Vazquez-Vazquez, A., et al. The Impact of the Covid-19 Lockdown on the Experiences and Feeding Practices of New Mothers in the UK: Preliminary Data from the COVID-19 New Mum Study.	Oct-20	United Kingdom	N = 1365 Mothers of children ≤12 months	Cross-sectional	Online survey, conducted from May 27-June 3, 2020, with mothers to explore COVID-19 impact on birth experience and infant feeding practices	Delivery mode, skin-to-skin contact and breastfeeding initiation did not differ between groups. 'During lockdown' (DL) women had shorter hospital stays. 39% reported changes to their birth plan. Reflecting younger infant age, 59% of DL infants were exclusively breast-fed/ mixed fed versus 39% 'before lockdown' (BL) (p < 0.05). 13% reported a change in feeding; often related to lack of breastfeeding support, especially with practical problems. 45% of DL women reported insufficient feeding support. Among BL women, 57% and 69% reported decreased feeding support and childcare, respectively.

Opportunities for early learning

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Barnett, S., Jung, K., & Nores, M. Young children's home learning and	Aug-20	USA	N = 945 Parents of children ages 3-5 years	Cross-sectional	The survey was conducted between May 22 and June 5, 2020 about child's	The 2020 NIEER survey reported frequencies of engaging in reading, singing, teaching activities, and arts and crafts were lower than those in the 2019 NHES. The percentage reporting storytelling three or more times in the past week was higher

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
preschool participation experiences during the pandemic. NIEER 2020 Preschool Learning Activities Survey					daily activities during the pandemic.	than before the pandemic. Nearly three-quarters (74%) of children attending preschool programs had their programs closed as a result of the pandemic. More than 90% of public program classrooms closed compared to two-thirds of private programs.
Briggs, D. C. COVID-19: The Effect of Lockdown on Children's Remote Learning Experience – Parents' Perspective.	Oct-20	Nigeria	<i>N</i> = 260 parents of schoolchildren	Cross-sectional	An online survey distributed from May 9-June 8, 2020 explored parents' perspectives on education during stay-at-home periods	77.7% possess a computer device, 93.2% have internet services and 55.0% spend between N5,000 – N9,999 (13.1 – 26.2 USD) per month for internet services. Only 35.4% had their children participate in the ongoing radio/TV learning sessions, 46.1% educate their children with their computer devices at home, however, 40.0% preferred an online based method of homeschooling. Overall, 70.0% were dissatisfied with the level of homeschooling their children received. Respondents with higher education and income level were significantly more likely to possess a computer device, have internet access at home and prefer online classes when compared to low-income earning parents with secondary education and below.
Dayal, H. C., & Tiko, L. When are we going to have the real school? A case study of early childhood education and care teachers' experiences surrounding education during the	Oct-20	Small Pacific Island	<i>N</i> = 4 teachers/staff at early childhood education and care centres	Qualitative	Telephone interview were conducted to allow teachers to share their feelings and emotions about the pandemic and school closures; plans for delivering education during the pandemic; and proposed strategies	Teachers worried about students' health and safety and general well-being during complete lockdown. In one facility, each teacher created a Viber group with parents (a free app that has text, calling and video-calling features) to communicate about general well-being of children and to send learning materials. Parents who cannot access Viber are called by teachers on their phones. Teachers mentioned the need for more training in technical issues. Another facility is using Zoom for continuing education. Staff noted approximately 80% of students were able to continue education, including children who had returned to Australia and New Zealand.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
COVID-19 pandemic.					to overcome challenges.	Staff had a single laptop and modem, so teachers took turns delivering their virtual lessons at different time. Some parents did not have Zoom enabling resources at home or were unaware of how to use them. The latter was mainly for children who were looked after by grandparents, who may have lacked knowledge to use online technologies. Parents who faced such difficulties had 'print copy' teaching materials sent to them through email, or in a few cases, delivered to their homes by staff.
Dong, C., et al. Young Children's Online Learning during COVID-19 Pandemic: Chinese Parents' Beliefs and Attitudes.	Sep-20	China	<i>N</i> = 3275 Parents of children enrolled in ECE	Cross-sectional	This survey study was to explore parental beliefs and attitudes about online learning.	Most parents (92.7%) reported that their children had online learning experiences during the pandemic, and many (84.6%) spent less than a half-hour each time. The parents generally had negative beliefs about the values and benefits of online learning and preferred traditional learning in early childhood settings. They tended to resist and even reject online learning for three key reasons: shortcomings of online learning, young children's inadequate self-regulation, and their lack of time and professional knowledge in supporting children's online learning. Also, the hardship caused by the COVID-19 pandemic has made them suffer, thus more resistant to online learning at home.
Grover, S., et al. A Survey of Parents of Children Attending the Online Classes During the Ongoing COVID-19 Pandemic.	Oct-20	India	<i>N</i> = 289 guardians of children attending online classes	Cross-sectional	An online survey carried out between June 21 and July 17, 2020 assessed the impact of online classes on children and their parents or guardians.	A majority (93.8%) of children were studying in private schools and attending online live classes (83.7%). About three-fourth of the parents/guardians reported that online classes were less comfortable (81.7%) and less satisfactory (78.5%); guardians reported that children had poor attention and concentration (80.6%), lower learning of theoretical and practical aspects of the subject (82.4%), and lower ability to sit satisfactorily for the duration of the class (78.6%). The

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
						most common reported distraction was surfing internet (52.2%) or online games (52.2%).
Kim, J., et al. The Implications of COVID-19 on Early Learning Continuity in Ethiopia: Perspectives of Parents and Caregivers.	Oct-20	Ethiopia	N = 480 Parents of children <5	Cross-sectional	mobile phone surveys to understand how parents and caregivers with pre-primary aged children have	More than half of respondents reported they do not have children's books or learning materials at home, especially caregivers who are not literate. A large difference between urban and rural locations was also observed for families' access to electricity and technologies to support children's learning. Most have received little support from schools or local governments. Only 10% of caregivers reported they have been in contact with pre-primary teachers or school principals, with significant differences by household wealth and across regions. Only half report they have engaged in supporting learning activities for pre-primary children, which favours families living in urban areas. Mothers are most likely to be responsible for supporting children's learning at home; yet mothers are less likely to be literate. Around three quarters reported they play more often with their child since the COVID-19 crisis than before, including telling stories or singing songs. However, caregivers from poorer households or who are illiterate are less likely to engage in such activities. Only 12% said they had used radio lessons with their children.
Lee, S. J., et al. Parenting Activities and the Transition to Home-Based Education During the COVID-19 Pandemic.	Oct-20 <i>Preprint</i>	USA	N = 405 Parents with children <12 years	Mixed methods	An online survey, including open-ended questions, conducted from April 24-25, 2020, examined parental mental health and the transition to home-based learning.	78% of parents said they were educating their child at home due to COVID19. Most (77.1%) reported use of online tools for at-home education, including educational apps, social media, and school-provided electronic resources. Most parents were spending more time involved in daily caregiving of their children since COVID-19. Content analyses of open-ended questions indicated that school closures were a significant disruption, followed by lack of physical activity, and social isolation.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Mangiavacchi, L. et al. Fathers Matter: Intra-Household Responsibilities and Children's Wellbeing During the Covid-19 Lockdown in Italy.	Jul-20	Italy	N = 3,352 families with children < 16 years	Cross-sectional	Online survey data collected in April 2020 examines children's wellbeing, and parents' market and household work during the lockdown.	Only about 60% of primary school children had some live online classes and about 30% received only some learning material accompanied by homework to be delivered through an electronic registry. Only 18% of children aged 3-5 had some live on-line classes, and 37% received some material from their teachers. 45% of children aged 3-5 were never involved in any school activity at the date of interview. Changes in reading to preschool children, who listen to their parents reading, increased by almost half an hour more per day. During lockdown, all children watched about twice as much TV (about 3 hours for kindergarten and primary school children). Labor market restrictions imposed during lockdown resulted in increases in fathers' involvement in childcare and homeschooling. Fathers' involvement had a positive association with child emotional well-being, but no impact on reading hours and educational progress.
World Vision. Rapid Assessment of the Impact of COVID-19 on Child Wellbeing in Cambodia.	Jun-20	Cambodia	N = 222 household heads (survey); 42 key informants (local leaders, etc.); 65 Agricultural Cooperative leaders (survey)	Mixed methods	The rapid assessment was conducted from May 11-14, 2020 to assess food insecurity; education; and livelihoods among vulnerable families.	Confirmed that home learning is very challenging for children for a variety of reasons, including limited availability of learning materials at home and limited support received by children. Technological barriers (bad internet connectivity, costs of internet connection and limited availability of a device to connect) while significant were not the highest reported challenges. Overall, there was a consensus that school closures are going to lead to school dropout, which respondents link to the impact that COVID-19 is having on livelihoods and to the perceived increase in child labour.

Rapid Evidence Review: COVID-19 and Nurturing Care

Opportunities for early learning: Outdoor play and physical activity

Reference	Date released	Setting	Population	Study Design	Data Source	Summary of Findings
Carroll, N., et al. The Impact of COVID-19 on Health Behavior, Stress, Financial and Food Security among Middle to High Income Canadian Families with Young Children.	Aug-20	Canada	<i>N</i> = 254 Parents of children ages 18 months to 5 years	Cross-sectional	An online survey administered between 20 April 2020 to 15 May 2020 to explore impact of COVID-19 on physical activity, screen time, and sleep among parents and children, as well as stress, financial and food insecurity of families.	Many parents expressed concern about decreases in frequency and intensity of children's physical activity and increases in screen time. Parents reported physical activity is limited due to lack of space and variety in tools and toys. Parents also reported strategies used to increase physical activity, such as scheduling outdoor playtime. Specific resources that were frequently mentioned included resources on engaging children in physical activity or any other activities, e.g., crafts, cooking to decrease screen time, tips for grocery shopping during COVID-19, homeschooling, scheduling, and time management.
de Lannoy, L. et al. Regional Differences in Access to the Outdoors and Outdoor Play of Canadian Children and Youth during the COVID-19 Outbreak.	Oct-20	Canada	<i>N</i> = 1472 Parents of children ages 5–17 years	Cross-sectional	Used online survey data collected in April 2020. Parents were asked about their child's physical activity, sleep, sedentary time, time spent outdoors and in outdoor play in the last week and were asked to compare behaviours before and during the COVID-19 outbreak.	Change in child and youth outdoor behaviour ranged from 1 (much less play) to 5 (much more play) with 3 representing no change. Overall, we found that all regions exhibited a decrease in time spent outdoors and in outdoor play (i.e., below 3), with Ontario having experienced the greatest decline in both time spent outdoors and in outdoor play ($p < 0.001$).
Dodd, H., et al. Co-SPYCE Report One: Findings from 1728 parents/carers of 2-4 year olds on stress, child activities, child worries and need for support.	May-20	UK	<i>N</i> = 1728 caregivers of children aged 2-4 years	Cross-sectional	An online survey was completed between April 17 April and May 20, 2020 to explore parent/carer stress, how children are spending time, parent feelings about child play, and the need for parental support.	Over 80% of children are reported by participating parents/carers to be getting at least 30 minutes of exercise a day but only 22% are reported to be getting the recommended 3 hours. Almost half of the children (47%) are reported to be spending no time playing with another child in their household. The top 3 stressors for participating parents/carers were (i) work, (ii) child's screen time, (iii) their child's wellbeing. Participating parents/carers particularly want support around managing children's emotions, educational demands, behaviours and coming out of social isolation.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date released	Setting	Population	Study Design	Data Source	Summary of Findings
Drouin, M., et al. How Parents and Their Children Used Social Media and Technology at the Beginning of the COVID-19 Pandemic and Associations with Anxiety.	Jul-20	USA	N = 260 Parents of children ≤18 months	Cross-sectional	Online survey data, collected from March 20-25, 2020, explored children's use of social media and technology during the COVID-19 pandemic and its association with anxiety.	Mean differences revealed that children in the 0 to 5 year age range did not increase their technology use as much as older age groups, nor did they use social media/ social technology or video/phone calls as frequently.
Dunton, G., Do, B., & Wang, S. Early Effects of the COVID-19 Pandemic on Physical Activity and Sedentary Behavior in US Children.	Sep-20	USA	N = 211 Parents and guardians of children ages 5-13 years	Cross-sectional	online survey between April 25–May 16, 2020 assessed physical activity and sedentary behavior in previous day, as well as perceived changes in activities between pre-COVID-19 period and early-COVID-19 periods.	The most common physical activities during the early-COVID-19 period were free play/unstructured activity (e.g., running around, tag) (90% of children) and going for a walk (55% of children). Children engaged in about 90 min of school-related sitting and over 8 h of leisure-related sitting a day. Girls and older children generally spent more time in sedentary behaviors than boys and younger children. Children were more likely to perform physical activity at home indoors or on neighborhood streets during the early-vs. pre-COVID-19 periods.
Guerrero, M. D., et al. Canadian Children's and Youth's Adherence to the 24-h Movement Guidelines during the COVID-19 Pandemic: A Decision Tree Analysis.	Jun-20	Canada	N = 1472 Parents of children ages 5-17 years	Cross-sectional	Online survey fielded in April 2020 explored adherence and non-adherence to Canadian 24-hr movement guidelines during the COVID-19 outbreak. Survey assessed demographic, social, and environmental characteristics, as well as physical activity, screen time, and sleep.	Characteristics associated with adherence to recommendations included high parental perceived capability to restrict screen time, annual household income of >=\$100,000, increases in children's and youth's outdoor physical activity/sport since the outbreak began, being a boy, having parents younger than 43 years old, and small increases in children's and youth's sleep duration since the outbreak began. Characteristics associated with non-adherence to the recommendations included low parental perceived capability to restrict screen time, youth aged 12-17 years, decreases in children's and youth's outdoor physical activity/sport since the outbreak began, primary residences located in all provinces except Quebec, low parental perceived capability to support children's and

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date released	Setting	Population	Study Design	Data Source	Summary of Findings
						youth's sleep and physical activity, and annual household income of \leq \$99,999.
Mitra, R., et al. Healthy Movement Behaviours in Children and Youth During the COVID-19 Pandemic: Exploring the Role of the Neighbourhood Environment.	Aug-20	Canada	<i>N</i> = 1472 Parents of children ages 5-17 years	Cross-sectional	Secondary analysis of an online survey conducted in April 2020 explores patterns of physical activity, sedentary and sleep behaviours among children and youth during the COVID-19 pandemic. The survey included health behavior measures and built environment measures.	Cluster analysis identified two groups (those with increased outdoor activities, those with decreased outdoor activities). Compliance to 24-hour movement guidelines was low among both groups. For children (aged 5-11), living in houses (versus apartments) was correlated with increased outdoor activities; proximity to major roads was a barrier. Similarly, dwelling density was negatively associated with increased outdoor activity, suggesting that those living in high-density neighbourhoods may demonstrate less healthy movement behaviours during a pandemic. However, results also indicate that for those living in higher density areas, access to parks within 1 km increased the probability of being in cluster 1 (i.e. increased outdoor activities).
Moore, S. A., et al. Impact of the COVID-19 Virus Outbreak on Movement and Play Behaviours of Canadian Children and Youth: A National Survey.	Jul-20	Canada	<i>N</i> = 1472 Parents of children ages 5–17 years	Cross-sectional	Online survey assessed immediate changes in child movement and play behaviours during the COVID-19 outbreak. Behaviours included physical activity and play, sedentary behavior, and sleep.	Only 4.8% (2.8% girls, 6.5% boys) of children and 0.6% (0.8% girls, 0.5% boys) of youth were meeting combined movement behaviour guidelines during COVID-19 restrictions. Children and youth had lower PA levels, less outside time, higher SB (including leisure screen time), and more sleep during the outbreak. Parental encouragement and support, parental engagement in PA, and family dog ownership were positively associated with healthy movement behaviours. Although families spent less time in PA and more time in SB, several parents reported adopting new hobbies or accessing new resources.
Xiang, M., et al. Impact of COVID-19 Pandemic on Children and Adolescents' Lifestyle Behavior Larger than Expected.	Jul-20	China	<i>N</i> = 2426 children and adolescents (6–17 years)	Quasi-experimental	Longitudinal online surveys conducted from 3-21 January and 13-23 March 2020 to better understand physical activity and sedentary behavior among school children.	Overall, the median time spent in physical activity decreased drastically, from 540 min/week before the pandemic to 105 min/week during the pandemic. During the pandemic, prevalence of physically inactive students increased from 21.3% to 65.6%. Screen time considerably increased during the pandemic in total (+1730 min [or approximately 30 h] per week on average). Screen time during leisure was also prolonged, indicating that nearly a quarter of students engaged in long screen time for leisure.

Rapid Evidence Review: COVID-19 and Nurturing Care

Safety and Security: Child protection from violence and neglect

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Barboza, G. E., et al. A Spatiotemporal Analysis of the Impact of COVID-19 on Child Abuse and Neglect in the City of Los Angeles, California.	Sep-20	USA	Children <18 years	Quasi-experimental	Utilizes police reports from before and during the COVID-19 pandemic (July 24, 2019–January 20, 2020 vs. January 21–July 19, 2020) to explore association between implementation of social distancing protocols and reported child abuse and neglect. Neighborhood structural factors (e.g., school absenteeism, poverty, unemployment, housing insecurity and birth assets) are explored.	Results indicate a significant 8% decrease in the number of child abuse and neglect reports during the COVID-19 pandemic (January 21, 2020–July 19, 2020) compared to the time period immediately preceding it (July 24, 2019–January 20, 2020). Spatiotemporal analysis identified locations of emerging "hotspots" for reported child abuse and neglect during the pandemic. Neighbourhoods characterized by severe housing burden, the number of assets children have at birth, poverty, school absenteeism and labor force participation were significantly associated with new and intensifying hotspots of child abuse and neglect.
Baron, E. Jason, et al. Suffering in Silence: How COVID-19 School Closures Inhibit the Reporting of Child Maltreatment.	Aug-20	USA	School age children	Quasi-experimental	Uses publicly available, monthly county-level data for Florida from January 2004 to April 2020 to estimate the effect of COVID-19 school closures on the number of child maltreatment allegations. Leverage monthly data to predict the number of allegations that would be expected due to seasonal fluctuations and secular trends in March and April 2020.	Using counterfactual distribution of county-level data from Florida, results show the number of reported child maltreatment allegations was approximately 27% or 15,000 lower than expected for the months of March and April 2020, the first two months in which schools closed. Counties with higher numbers of educators trained to identify and report child maltreatment (e.g., school psychologists and school nurses) experience a disproportionately larger reduction in the number of child maltreatment allegations.
Bhopal, S., et al. Who Has Been Missed? Dramatic Decrease in Numbers of Children Seen for Child Protection Assessments during the Pandemic.	Jun-20	United Kingdom	Children ≤16 years	Quasi-experimental	Examined monthly referral data for children and young people attending child protection medical examination as part of a child safeguarding assessment from January to April 2020 in comparison with the same months in the last 2 years.	Results showed a one-third reduction in child protection assessments from January to April 2020, compared to previous years. The total number of assessments from January to April was 152 in 2018, 156 in 2019 and 99 in 2020.
Branstetter, S., et al. The Impact of the COVID-19 Pandemic	Oct-20 <i>Preprint</i>	Germany	N = 918 parents of	Cross-sectional	Online survey assessing the impact of the COVID-19 pandemic on families with young children from areas with	Many participants reported an impact of the COVID-19 pandemic on social and family life. About one third of the families (30.5%) were at

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
on Families in Germany.			children ages 18 months to 6 years		low and moderate COVID-19 prevalence were fielded at the end of the German lock-down period. Outcomes were parent-reported impact of the COVID-19 pandemic on family life, concerns and trust in political measures. Participants were part of two ongoing cohort studies.	least very much burdened by the situation, and the lack of social contacts was very/extremely bothersome for almost half of families. On the other hand, the majority coped well with the situation. About half had a clear daily schedule. Families indicated that they were enjoying the additional time they could spend together (60%) and that they had united as a family (56%). Nevertheless, 22.5% of the families reported relevantly more conflicts with children and 19% and 12% reported more conflicts between children or between the spouses, respectively.
Bullinger, L., et al. COVID-19 and Alleged Child Maltreatment.	Oct-20	USA	Children <18 years	Quasi-experimental	Compared weekly, county-level referral data from January 8, 2018 through May 30, 2020 for all 159 counties in Georgia to examine whether COVID-19 related policies affected child maltreatment referrals. Superimposed aggregate, anonymous cell phone tracking data to determine whether time at home is related to referrals.	Relative to 2018 and 2019 trends during the same time period, referral rates plummeted by approximately 55% following the emergency declaration. Maltreatment referrals begin to slowly return to pre-COVID-19 levels around 7 weeks following the emergency declaration, but never fully rebound during the study period. As time at home during the pandemic increased, the risk of supervisory neglect increased, with largest impacts in metropolitan counties and counties with more poverty.
Bullinger, L. R., et al COVID-19 and Crime: Effects of Stay-at-Home Orders on Domestic Violence.	Jul-20	USA	Children <18 years	Quasi-experimental	Uses cell phone block-level activity data and administrative 911 and crime data from 2019 and 2020 for Chicago to estimate the effects of Stay-at-Home Orders on calls for police service, crimes recorded by police, and arrests made relating to domestic violence.	Results show domestic-related police calls increased by 8% during the stay at home order. The effects are largest in areas with more households that are married with children, areas with relatively high rates of rental residences, and areas that experienced the largest changes in staying at home. Did not find significant effects for domestic crimes against children, as measured by physical child abuse. This result may indicate that although domestic crimes rose in March-April 2020, calls for child abuse fell, potentially due to lessened visibility of child victims and limitations of various reporting channels for child maltreatment crimes.

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Cabrera-Hernández, F., and Padilla-Romo, M. Hidden Violence: How COVID-19 School Closures Reduced the Reporting of Child Maltreatment.	Jul-20	Mexico	Children ages 3-15 years	Quasi-experimental	Leverage incident-level crime reports from Mexico City Attorney General's Office for January 2019-June 2020 to estimate the impact of school closures on the reporting of school-age child maltreatment.	The results show an average reduction of 21% and 30% in March 2020 for the reporting of child maltreatment among school-aged children, or a reduction of 7 to 10 cases from the average of 33 per week. These effects are higher for females, at a reduction of 66%, or 22 cases per week from an average of 33; and in poorer municipalities (reduction of 40% or 14 cases less a week, from an average of 34). These effects suggest that 220 incidents of child maltreatment for school-age females were not reported to the authorities in the 10 weeks after school closures.
Chung, G., et al. Mediating Effects of Parental Stress on Harsh Parenting and Parent-Child Relationship during Coronavirus (COVID-19) Pandemic in Singapore.	Jun-20	Singapore	N = 258 Parents of children <12 years	Cross-sectional	An online survey disseminated from April 22, 2020, to May 5, 2020 explored work, family life and parenting during the pandemic.	Parents who were more impacted by COVID-19 experienced more parenting stress than those who felt they were less impacted by COVID-19, and parents who experienced more parenting stress indicated that they had used more harsh parenting and felt less close with their children in the past weeks. There was no evidence that parents' perceived impact of COVID-19 influenced harsh parenting or parent-child relationships independent of its effect on parenting stress.
Dulieu, N. and Burgess, M. The Hidden Impact of COVID-19 on Child Rights.	Sep-20	Multiple countries (37 countries)	N = 17,565 Parents and caregivers of children <18 years	Cross-sectional	An online survey assessed implications of COVID-19 pandemic on Child Rights. Measured household-level characteristics, child rights, caregiver's well-being and family relationships, child education and well-being, among other topics. Sample included Save the Children program participants, specific population groups of interest to the program, and the general public.	Nearly one-third (32%) of the households reported that violence had occurred in the home, including children and/or adults being verbally or physically abused. Parents/caregivers in urban locations reported a higher rate of violence in the home (23%) compared to rural respondents (17%). A statistically significant positive correlation was found between reported income loss and reported levels of household violence.
Edwards, J. Protect a Generation: The	Sep-20	Multiple	N = 17,565 Parents	Mixed methods	An online survey conducted with Save the Children participant households assessed implications of	In response to the survey, more than 3 in 4 households reported an income loss since the beginning of the pandemic. Poorer households were

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Impact of COVID-19 on Children's Lives.		countries (16 countries)	and caregivers of children <18 years		COVID-19 pandemic on child rights, caregiver's well-being and family relationships, child education and well-being, among other topics. Qualitative data on adverse experiences was collected online from participant households in 46 countries via and a separate survey sent to populations of specific interest.	more likely to suffer income losses (82%) than those not classified as poor (70%). Urban households were also disproportionately affected by income loss, with respondents from urban areas almost twice as likely to say they have lost their job (61%) compared with those in rural areas (33%). 70% of respondents who suffered economic losses reported that they had not received government support. Nearly one third (32%) of households had a child, parent or caregiver who said that there had been physical or emotional violence in their home since the start of the pandemic. The number of households with a parent or caregiver respondent who said there was violence varied significantly across regions.
Garstang, J., et al. Effect of COVID-19 Lockdown on Child Protection Medical Assessments: A Retrospective Observational Study in Birmingham, UK.	Sep-20	United Kingdom	Children under the age of 18 years	Quasi-experimental	Compared data for all child protection medical examinations (CPME) at a large, urban healthcare network for 18-week periods in 2018, 2019 and 2020 to compare changes in non-accidental injury (NAI) or neglect during the COVID-19 school shutdowns. Assessed child demographic data, referral details, and whether there were physical findings to support NAI or neglect.	The results indicate a 39.7% reduction in referrals for CPME from March to June 2020, compared to the same period in 2018 and a 37.3% reduction from 2019 to 2020. There were significantly fewer girls referred in 2020. The children referred for CPME in 2020 had similar social care histories to other years with the majority being previously known to social care and approximately half being open cases at the time of referral.
Hansen, Alison L. Child Maltreatment Reporting Statistics During the Covid-19 Pandemic: A Cursory Analysis.	Aug-20	USA	Children served by state protection agencies	Quasi-experimental	The number of total allegations of child maltreatment between the months of March and June—a span of time representative of the COVID-19 pandemic thus far—were analyzed in five different states in the years 2017, 2018, 2019, and 2020.	Utilizing publicly reported child maltreatment data from five states (Florida, New Jersey, Arizona, Nevada and Indiana), results reveal a disproportionate decline in child maltreatment reports during the COVID-19 pandemic (March to June 2020), compared to previous years from 2017 to 2019, across all five states. Indiana and Nevada experienced the greatest percentage change in reports of alleged child maltreatment. Nevada experienced a 26.6% decrease in reports between

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
						2019 and 2020. Similarly, Indiana experienced a 24.3% decrease in reports of alleged child maltreatment in the same period.
Jentsch, B., and B. Schnock. Child Welfare in the Midst of the Coronavirus Pandemic-Emerging Evidence from Germany.	Sep-20	Germany	German- and English-language scientific articles, position papers and other gray literature	Review	A systematic search of German and English academic and gray literature using key word searches and the dates April 19, 2020 to August 27, 2020. Interim results of empirical studies conducted in Germany were included.	Results indicate that the number of reports on alleged child maltreatment had declined in March and April 2020. Out of 231 youth welfare agencies responding, 43% stated volume of reports had decreased in first four weeks following contact restrictions. A more comprehensive survey of agencies carried out in May 2020 similarly reported that alleged child maltreatment reports had not increased and, in some cases, decreased. A similar development was noted concerning number of children being taken into care.
Kovler, M. L., et al. Increased Proportion of Physical Child Abuse Injuries at a Level I Pediatric Trauma Center during the Covid-19 Pandemic.	Sep-20	USA	Children under the age of 15 years	Quasi-experimental	Retrospective review of patients at a pediatric trauma centre to identify injuries caused by physical child abuse in the month following the statewide closure of childcare facilities in Maryland. The proportion of PCA patients treated during the Covid-19 era were compared to the corresponding period in the preceding two years.	Results indicate an increase in the proportion of traumatic injuries caused by physical child abuse in March and April 2020, compared to the same period in 2019 and 2018. Eight patients with physical child abuse injuries were treated during the Covid-19 period (13% of total trauma patients), compared to four in 2019 (4%) and three in 2018 (3%). The median age of patients in the Covid-19 period was 11.5 months. Most patients were black (75%) with public health insurance (75%). All injuries were caused by blunt trauma, resulting in scalp/face contusions (63%), skull fractures (50%), intracranial hemorrhage (38%), and long bone fractures (25%).
Lawson, M., et al. Child Maltreatment during the COVID-19 Pandemic: Consequences of Parental Job Loss on Psychological and	Sep-20	USA	N = 342 parents who are in a committed romantic relations	Cross-sectional	Online survey data collected from mid-April to mid-May 2020, assessed experiences with COVID-19, the Parent-Child Conflict Tactics Scale, and the Family Crisis Oriented Personal Evaluation Scales.	Parental job loss was positively and significantly associated with psychological maltreatment and physical abuse towards children in the past week. Younger children were more likely to be physically abused and psychologically maltreated during the past week. Parents from families with more financial stability were less likely to physically abuse their children during the pandemic. Parents

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Physical Abuse towards Children.			hip and who have a child age 4 to 10 years			reporting more depressive symptoms were more likely to psychologically maltreat and physically abuse their children. Male children were more likely to be psychologically maltreated.
Lee, S. J., and Ward, K. P. Stress and Parenting During the Coronavirus Pandemic.	Mar-20	USA	N = 288 parents with a child age 0-12 that lives with them at least part of the time	Cross-sectional	Online survey, conducted March 24-25, 2020 to investigate mental health and wellbeing, parenting, and economic situation during the pandemic.	Parents are engaging in parent-child activities much more frequently since COVID-19. 81% of parents said that they praised or complimented their child(ren) very often, and 88% of parents said they and their children had shown love to each other very often in the past 2 weeks. However, most parents (52%) said financial concerns and social isolation (50%) were getting in the way of their parenting. Most parents (61%) shouted, yelled or screamed at their children at least once in the past 2 weeks; 20% spanked or slapped their child at least once in the past 2 weeks. Parents reported they have yelled or screamed at their children more often (19%) or increased discipline (15%) since the Coronavirus pandemic
Martins-Filho, P. R., et al. Decrease in Child Abuse Notifications during COVID-19 Outbreak: A Reason for Worry or Celebration?	Sep-20	Brazil	Children under the age of 18 years	Quasi-experimental	Compared data extracted from the official child maltreatment database (Coordination of Statistics and Criminal Analysis) for the first half of 2019 and 2020 in Sergipe state, Northeast Brazil.	The reported rate of child physical injury decreased in 2020 by 24.3% from 1 January to 30 June 2020, compared to the same time period in 2019. Reported rates were higher among girls (71.7%), children aged 12–17 years (69.8%) and Blacks/Browns (73.6%). These trends in child abuse notifications according to gender, age and race/ethnicity were not significantly different between 2020 and 2019.
Øverlien, C. (2020). The COVID-19 Pandemic and Its Impact on Children in Domestic Violence Refuges.	Aug-20	Norway	N = 46 Staff at domestic violence refuges	Cross-sectional	An online survey was distributed to all domestic violence refuges in Norway from April 8-23, 2020, with the aim of obtaining an overview of the impact of the COVID-19 crisis and the virus control measures on victims of domestic violence and	56% of respondents reported there had been a reduction in the number of requests from clients, many expressed concerns that it is 'too quiet out there.' Refuge staff reported being most concerned about children. Refuge staff were concerned that children exposed to violence who were living outside of the refuge would no longer receive the

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
					abuse. The survey contained 15 items with fixed answers, follow-up questions and an opportunity for the refugees to provide supplementary qualitative descriptions and examples.	help that they need from professionals; and, moreover, that the violence they experience would remain undisclosed, as contact between adults outside of the family and the abused child is dramatically reduced. Closed schools and daycare centres are of particular concern as school represents normality and a zone free from the abusive parent.
Rapoport, E., et al. Reporting of Child Maltreatment During the SARS-CoV-2 Pandemic in New York City from March to May 2020.	Sep-20	USA	Children under the age of 18 years	Quasi-experimental	Analyzed monthly data from New York City on the number of child maltreatment allegations using data from January 2015 to February 2020 to predict expected values for March, April, and May 2020.	The results show 29% fewer allegations of child maltreatment in March 2020 than forecasted. In April and May 2020, the number of reports was approximately half of what was expected. Significant decreases in child maltreatment reporting were also noted for all reporter subtypes including mandated reporters such as educational personnel, child care personnel and medical health personnel, and non-mandated reporters such as family members and neighbours.
Ritz, D., et al. The Hidden Impact of COVID-19 on Child Protection and Wellbeing.	Sep-20	Multiple countries	N = 17,565 Parents and caregivers of a child under age 18 years	Cross-sectional	An online survey assessed implications of COVID-19 pandemic on Child Rights. Measured household-level characteristics, child rights, caregiver's well-being and family relationships, child education and well-being, among other topics. Sample included Save the Children program participants, specific population groups of interest to the program, and the general public.	Over 1 in 5 parents/caregivers (22%) surveyed reported an increase in their use of negative or violent parenting methods. 32% of households had a child and/or parent/caregiver reporting that violence had occurred in the home, including children and/or adults being verbally or physically abused. The vast majority of parents/caregivers (89%) reported an increase in negative feelings due to the COVID-19 pandemic. While 83% of the parents/caregivers reported an increase in negative feelings when schools were closed between 1 to 4 weeks, the reports of negative feelings were higher for the vast majority of adults (95%) when schools had been closed for 17 to 19 weeks.
Sidpra, J., et al. Rise in the Incidence of Abusive Head Trauma	Jun-20	United	Children under the	Quasi-experimental	Reviewed the incidence of suspected acute head trauma at an urban hospital between 23 March and 23 April 2020—the month that the	Review of the incidence of suspected abusive head trauma indicated a 1493% increase per month compared to the same period over the previous 3 years (10 cases in comparison with a mean of 0.67

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
during the COVID-19 Pandemic.		Kingdom	age of 13 months		United Kingdom entered a period of national self-isolation—and compared this with the incidence in the previous 3 years.	cases per month). All families live in areas with a higher than average Index of Multiple Deprivation and 70% of parents had significant underlying vulnerabilities (e.g., criminal histories, mental health disorders, financial concerns).
Tener, D., et al. “How Does COVID-19 Impact Intrafamilial Child Sexual Abuse? Comparison Analysis of Reports by Practitioners in Israel and the US.	Oct-20	USA and Israel	N = 50 Child welfare and legal professionals	Qualitative	An open-ended, online questionnaire distributed during May 2020 compare professional perspectives and experiences working with intrafamilial child sexual abuse with respect to the context of the COVID-19 pandemic within the United States and Israel.	Professionals in both countries were extremely concerned by the lack of opportunity to report the abuse. A surge in reports of intrafamilial child sexual abuse (IFCSA) was predicted after the COVID-19 period ends. Findings reveal mostly negative changes in the dynamics of IFCSA families during COVID-19, including financial, environmental, and emotional hardships, as well as some positive changes in relationships among family members. Risks of transferring to internet based or telephonic therapeutic interventions included technical issues, limited internet access among ultra-Orthodox families, lack of proper separate space for therapy, and difficulties engaging younger children.
Toros, K., & Falch-Eriksen, A. A Child’s Right to Protection during the COVID-19 Crisis: An Exploratory Study of the Child Protective Services of Estonia.	Oct-20	Estonia	N = 81 Child protection workers	Qualitative	An online survey with three open-ended questions sent to child protection workers (CPW) in April and May 2020 collected case-worker narratives. Assessed the impact of COVID-19 on organisational design and, the workers’ ability to conduct a diagnosis of a child’s care context.	Results pointed towards challenges to facilitating children's protection during the pandemic. CPWs reported more difficulty identifying children and families in need; a decrease in referrals from the formal network (typically schools and kindergartens); perception of overload of the work by the networking partners; and a drop in provision of services for families in need, especially family therapy, psychology and child psychiatry. CPWs reported that child well-being and protection became a secondary consideration, and families' basic food needs and individual health protection from the virus became priorities.
United Nations Children’s Fund. Protecting Children	Aug-20	Multiple	N = 136 UNICEF	Cross-sectional	Online survey collected information from May 1-August 14, 2020 from 136 UNICEF country offices on	South Asia has the highest proportion of countries reporting service disruptions, followed by Eastern Europe and Central Asia. Case management and

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
from Violence in the Time of COVID-19: Disruptions in Prevention and Response Services.		countries	country offices		disruptions in service provision related to violence against children (VAC) as a result of the coronavirus.	home visits for children and women at risk of abuse are among the most commonly disrupted services. Around two thirds of countries with VAC-related service disruptions reported that at least one type of service had been severely affected. 70% of countries reported that mitigation measures had been put into place.
Whaling, K. et al. Reduced Child Maltreatment Prevention Service Case Openings during COVID-19.	May-20 <i>Preprint</i>	USA	Children under the age of 18 years	Quasi-experimental	Analyzed data from New York City's Administration for Children's Services for January, February, and March for the years 2013-2020 to investigate the frequency of child maltreatment prevention service case openings.	An examination of trend lines demonstrated that new prevention case openings have consistently peaked in the month of March for 2013-2019. New prevention case service openings in March 2020 do not peak, as they do in the preceding seven years. Results show new child maltreatment prevention case openings in March 2020 were 61% lower than mean new case openings over March 2013–2019.
World Vision. COVID-19 Rapid Impact Assessment Report.	May-20	Bangladesh	N = 12,713 people, including adults (53.2%) and children (46.8%)	Mixed methods	Used mixed method approach to understand impact of COVID-19 on most vulnerable children, families and communities across Bangladesh. Data were quantitative and qualitative-household (HH) surveys, focus group discussion and KIIs with community leaders, children groups, government representatives and faith leaders.	Loss of incomes among households has triggered stress among parents and caregivers who are now engaging in desperate negative behaviours towards their children with about 35% applying physical punishment or psychological aggression. More than half of the respondent (58%) indicating that they were to partially able, with one-third (33.7%) indicating capability while 7.2% respondent expressed that they cannot handle the situation.
World Vision. Rapid Assessment of the Impact of COVID-19 on Child Wellbeing in Cambodia.	Jun-20	Cambodia	N = 222 household heads (survey); 42 key informants (local leaders, etc.); 65 Agriculture	Mixed methods	The rapid assessment was conducted from May 11-14, 2020 to assess food insecurity; education; and livelihoods among vulnerable families.	57% of parents and caregivers reported that parenting has become more complicated or challenging since the beginning of the COVID-19 outbreak and, as a result, only 32% of them reported that they could adequately manage their children's behaviour. A significant number of local authorities, child club leaders, faith leaders and members of Commune Committees for Women and Children (CCWCs) interviewed mentioned that they are concerned by domestic violence in the

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
			ral Cooperative leaders (survey)			present context. However, the data from the household survey does not suggest a significant increase in the use of corporal punishments and additional research would be needed in this area to better assess the potential impact of the COVID-19 outbreak on domestic violence.
World Vision. Unmasking the Impact of COVID-19 on Asia's Most Vulnerable Children.	Aug-20	Bangladesh, Cambodia, India, Indonesia, Mongolia, Myanmar, Nepal, Philippines, and Sri Lanka	N = 19,065 parents/caregivers; and N = 14,669 children	Mixed methods	A household survey and child consultation were conducted in May 2020 via telephone and face to face to gain a better understanding on how the COVID-19 pandemic is impacting the lives of the most vulnerable children, their families and communities.	27% of parents/caregivers indicated that they had resorted to physical punishment or emotional abuse in the last month (39% for urban parents) while 25% of the children confirmed that their parents or caregivers used physical punishment and/or psychological aggression in the past month.

Safety and Security: Child Injury

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Bram J. T, et al. Where Have All the Fractures Gone?	Sep-20	USA	N = 1745 records of children	Quasi-experimental	Utilized hospital records from a level I pediatric trauma	Results indicate that pediatric fracture volume decreased 2.5-fold during the COVID-19 pandemic. There was a significant decrease in the incidence of fractures (22.5

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
The Epidemiology of Pediatric Fractures During the COVID-19 Pandemic.			under the age of 18 years		hospital from 15 March to 15 April 2020 to compare acute fractures during the COVID-19 pandemic with fractures during a pre-pandemic period at the same institution.	vs. 9.6 per day). The presenting age for all fractures decreased (7.5 vs. 9.4 years) because of decreased fracture burden among adolescents. There was also a decrease in the number of fractures requiring surgery (2.2 vs. 0.8 per day). During the pandemic, there was an increase in the proportion of injuries occurring at home (57.8% vs. 32.5%) or on bicycles (18.3% vs. 8.2%), but a decrease in those related to sports (7.2% vs. 26.0%) or playgrounds (5.2% vs. 9.0%).
Chaiyachati, B. H., et al. Trends in Pediatric Emergency Department Utilization after Institution of COVID-19 Mandatory Social Distancing.	Jul-20	USA	N = 231 records of children under age 21 years	Quasi-experimental	Compared emergency department (ED) visits to a tertiary urban children's hospital from January to April 2020 with the same date range during the 3 prior years (2017-2019).	Pediatric emergency healthcare use from January to April 2020 compared with the prior 3 years indicates a rapid decrease in emergency department visits (95 vs 286). Trauma, fever, and abdominal pain had higher proportions of high-acuity presentations in 2020. The average number of ingestions was higher in 2020 than in the comparison period (31 vs 25). The proportion of patients categorized as high acuity was higher in 2020 (59.4% vs 49.6%). Similarly, the proportions of patients admitted both overall (22.4% vs 18.5%) and to the intensive care unit (2.8% vs 1.7%) were higher in 2020.
Hamill J. K. and Sawyer M C. Reduction of childhood trauma during the COVID-19 Level 4 lockdown in New Zealand.	Jun-20	New Zealand	Children under the age of 18 years	Quasi-experimental	Compared patient data from New Zealand's only designated paediatric major trauma centre from period of lockdown from 26 March to 27 April 2020, to the same date range from the years 2016 to 2019	Results show total trauma admissions during the lockdown were at a 5-year low. There were 57 trauma admissions (four major trauma) during the lockdown compared to an average of 85 (six major trauma) during the same time period in previous years. No motor vehicle crash occupants were admitted. Nineteen percent of children were injured on bicycles compared to an average of 3.6% in previous years.
Li, H., et al. Changes in Children's Healthcare Visits During Coronavirus Disease-2019	May-20	China	Children under the age of 18 years	Quasi-experimental	Compared data from outpatient healthcare visits between January 1, 2020, and March 31, 2020, as well as historical data during the same period of 2019	Coincident with COVID-19 restrictions, the daily number of pediatric visits after January 25, 2020, decreased to about only one-quarter compared with the historical data in 2019. The number of injuries also rose significantly, especially indoor injuries such as finger injuries, foreign body entering eye or natural orifice, tibial fracture, eyelid laceration, and supracondylar

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Pandemic in Hangzhou, China.					for a tertiary Children's Hospital in a large Chinese City.	fracture of humerus. Some cases of epistaxis also were due to injuries. After January 25, 2020, there were 19 electrical injuries seen, compared with only 6 in the first 3 months in 2019.
Nabian M. H., et al. Epidemiological Pattern of Pediatric Trauma in COVID-19 Outbreak: Data from a Tertiary Trauma Center in Iran.	Sep-20	Iran	Children under the age of 18 years	Quasi-experimental	Compared patient data on pediatric fractures from a tertiary trauma center from March 1-April 15, 2020 with mean data from the past 2 years.	117 of 288 trauma patients during the 2020 period (40.62%) had a fractured bone (145 fractures). Patients were mostly boys, with a mean age of 9.87 years. The three most common fracture types in children involved arm fractures. Compared to non-COVID era, the number of pediatric trauma admissions dropped from 589 to 288. No significant change happened in the mean age, male/female ratio and percentage of motor vehicle accidents. Proportion of proximal humeral, proximal forearm, carpal, and hand fractures declined. The number of open fractures significantly dropped (from 12 to 2).
Raitio A, et al. Reduced Number of Pediatric Orthopedic Trauma Requiring Operative Treatment during COVID-19 Restrictions: A Nationwide Cohort Study.	Oct-20	Finland	Children under the age of 16 years	Quasi-experimental	Assessed pediatric emergency operations related to fractures, appendicitis and scrotal exploration for all five tertiary centers in Finland for the periods from March 1-May 31 for 2017-2020.	Results show a significant decrease (31%) in trauma operations compared to the same period during the previous three years. Daycare, school, and organized sports-related injuries decreased. The frequencies for scrotal exploration and surgery for acute appendicitis were similar during the pre-pandemic and pandemic era. There was no change in the age or sex of the children undergoing emergency surgery in four university hospitals; in one hospital, the age of the fracture patients was significantly lower during the pandemic months (median 12.2 versus 10.4 years).
Sugand K, et al. Impact of the COVID-19 Pandemic on Paediatric Orthopaedic Trauma Workload in Central London: A Multi-centre Longitudinal Observational	Aug-20	UK	Children under the age of 18 years	Quasi-experimental	Compared patient data from a large acute paediatric hospital service in urban UK during the first lockdown from 17 March to 28 April 2020 with the same period in 2019, including both acute paediatric orthopaedic trauma	There was a significant reduction in the odds of sporting-related mechanism of injuries (by 57%) in 2020 compared with 2019. There were only 97 referrals following the introduction of social distancing measures in 2020 compared with 302 in 2019. This represents a 68% reduction in paediatric injuries. The general demographic of those presenting with injuries changed between the 2 periods, with a significantly younger median age in 2020 and more girls. The pattern of injury also remained generally unchanged with upper limb injuries being the most common at 67% and 69%

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Study over the “Golden Weeks”					referrals and operative caseload.	respectively in 2019 and 2020. In 2020 there were 3% more safeguarding referrals for non-accidental injuries (not a statistically significant difference compared to 2019). There has also not been a large increase in delayed presentation of substantial injuries.
Turgut A, et al. Effect of COVID-19 Pandemic on the Fracture Demographics: Data from a Tertiary Care Hospital in Turkey.	Jul-20	Turkey	Children under the age of 17 years	Quasi-experimental	Assessed records of patients admitted to a tertiary care hospital with a new fracture during pandemic period of March 16 to May 22, 2020, compared to the same date range in 2018 and 2019. Compared pediatric (<=16 years) and non-pediatric (>16 years)	A total of 1794, 1747, and 670 fractures were observed in 2018, 2019, and 2020, respectively. Mean age of the patients in pediatric group was found to have decreased significantly in the pandemic period. The most common fracture sites in the pediatric population were the distal forearm and distal arm. The proportional increase in femoral and tibial shaft fractures in the pediatric group was found to be statistically significantly. In the pediatric group 6.8%, 7.7%, and 14.6% of the fractures were treated surgically in 2018, 2019, and 2020, respectively.
Verdoni F., et al. Effect of the COVID-19 Outbreak on Pediatric Patients’ Admissions to the Emergency Department in an Italian Orthopedic Trauma Hub.	Sep-20	Italy	Children under the age of 17 years	Quasi-experimental	The records of pediatric patients admitted to emergency room of an orthopedic trauma center during the pandemic period from 12 March to 5 May 2020 was compared to the same period in 2019.	Results show a reduction of 87% in admissions. A statistically significant difference between the two groups was found in the mean age of patients, that decreased from 11.4 prior to the pandemic to 8.6 years during the pandemic. a trend towards more severe codes was observed in the pandemic group and the diagnosis of fracture was significantly less frequent in non-pandemic group and injuries at home were more frequent (34.8% compared to 6.8% in the non-pandemic group).

Rapid Evidence Review: COVID-19 and Nurturing Care

Reference	Date Released	Setting	Population	Study Design	Data Source	Summary of Findings
Williams F. N., et al. COVID-19 related Admissions to a Regional Burn Center: The impact of Shelter-in-place Mandate.	Oct-20	USA	School-aged children from age 6 to 17 years	Quasi-experimental	Compared all burn center admissions from between 10 March and 22 May 2020 to a month-matched historical cohort from 2019 and summer-time admissions for the last five years.	Compared to the 2019 cohort, there was a 28% increase in admissions of school aged children in 2020 during the shelter-in-place order time frame. Comparing the percentage of school age admissions from 10 March to 22 May 2020 (11%) to the percentage of school age admissions during the summer breaks from 2015 to 2019, the percentages were essentially equivalent (11%, 11%, 12%, 10%, 10%, respectively). None of these comparisons were statistically significant. There were also no statistically significant differences in race, sex, or type of insurance. For school-age children, there was a 36% increase in those with free or low cost federal and state health insurance in 2020 compared to 2019, though this did not reach statistical significance.