Supplementary Table 1. Characteristics of participants based on phone call success for telephone interviews

| Characteristics | Phone call success | | |
|--------------------------|--------------------|-------------|--|
| | Yes (n=2894) | No (n=1401) | |
| Covide-19 confirmed case | 45% | 40% | |
| Age group | | | |
| < 18 years | 13% | 11% | |
| 18 – 30 years | 28% | 26% | |
| 31 – 40 years | 24% | 25% | |
| 41 – 50 years | 17% | 17% | |
| 51 – 60 years | 11% | 14% | |
| > 60 years | 7% | 8% | |
| Female sex | 46% | 48% | |
| Fever | 56% | 58% | |
| Cough | 48% | 48% | |
| Cold | 7% | 10% | |
| Shortness of Breath | 4% | 3% | |
| Body ache | 4% | 2% | |
| Headache | 0.5% | 0.5% | |
| Sore throat | 7% | 7% | |
| Weakness | 0.3% | 0.3% | |
| Anosmia | 1% | 1% | |
| Loose motion | 0.6% | 1% | |
| Runny nose | 0.5% | 0.3% | |

Supplementary Table 2. Baseline characteristics of non-hospitalized COVID-19 test positives and test negatives

| Characteristics | | COVID-19 | |
|--|------|------------|------------|
| | n | Negative | Positive |
| Age group, n (%) | 4149 | | |
| < 18 years | | 335 (14%) | 194 (10%) |
| 18 – 30 years | | 689 (29%) | 472 (27%) |
| 31 – 40 years | | 579 (24%) | 418 (23%) |
| 41 – 50 years | | 404 (17%) | 299 (17%) |
| 51 – 60 years | | 266 (11%) | 209 (12%) |
| > 60 years | | 125 (5%) | 159 (9 %) |
| Female sex, n (%) | 4159 | 1088 (45%) | 853 (49%) |
| BCG scar, n (%) | 2772 | 1285 (82%) | 1007 (83%) |
| ABO Blood group, n (%) | 2619 | | |
| A | | 355 (24%) | 262 (23%) |
| В | | 477 (32%) | 397 (35%) |
| AB | | 116 (8%) | 127 (11%) |
| О | | 520 (35%) | 365 (32%) |
| Pre-existing chronic disease | | | |
| COPD/Asthma/Respiratory illness, n (%) | 2815 | 164 (11%) | 115 (9%) |
| Hypertension, n (%) | 2816 | 259 (16%) | 262 (21%) |
| Ischemic heart disease (IHD), n (%) | 2814 | 57 (4%) | 57 (5%) |
| Chronic liver disease (CLD), n (%) | 2814 | 20 (1%) | 15 (1%) |
| Diabetes mellitus (DM), n (%) | 2816 | 186 (12%) | 167 (13%) |
| Hypothyroidism, n (%) | 2814 | 57 (4%) | 53 (4%) |
| Chronic kidney disease (CKD), n (%) | 2813 | 23 (1%) | 27 (2%) |
| Physical activity | 2772 | | |
| No | | 926 (60%) | 710 (58%) |
| Mild | | 515 (33%) | 428 (35%) |
| Moderate | | 62 (4%) | 59 (5%) |
| Vigorous | | 48 (3%) | 24 (2%) |
| Presenting symptoms | 4159 | | |
| Fever, n (%) | | 1125 (47%) | 1218 (70%) |

| Cough, n (%) | 1133 (47%) | 886 (51%) |
|----------------------------|------------|-----------|
| Cold, n (%) | 199 (8%) | 137 (8%) |
| Shortness of Breath, n (%) | 104 (4%) | 38 (2%) |
| Body ache, n (%) | 68 (3%) | 64 (4%) |
| Headache, n (%) | 11 (0.5%) | 9 (0.5%) |
| Sore throat, n (%) | 207 (9%) | 102 (6%) |
| Weakness, n (%) | 6 (0.3%) | 6 (0.3%) |
| Anosmia, n (%) | 16 (0.7%) | 33 (2%) |
| Loose motion, n (%) | 20 (0.8%) | 17 (1%) |
| Runny nose, n (%) | 14 (0.6%) | 5 (0.3%) |