

BMJ Open Causes of admissions and in-hospital mortality among patients admitted to critical care units in primary and secondary hospitals in Vietnam in 2018: a multicentre retrospective study

Vu Quoc Dat ,¹ Bui Thi Khanh Linh ,¹ Giang Bao Kim²

To cite: Dat VQ, Linh BTK, Kim GB. Causes of admissions and in-hospital mortality among patients admitted to critical care units in primary and secondary hospitals in Vietnam in 2018: a multicentre retrospective study. *BMJ Open* 2022;**12**:e061638. doi:10.1136/bmjopen-2022-061638

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-061638>).

Received 09 February 2022
Accepted 26 May 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

¹Department of Infectious Diseases, Hanoi Medical University, Hanoi, Viet Nam

²Institute for Preventive Medicine and Public Health, Hanoi Medical University, Hanoi, Viet Nam

Correspondence to

Dr Vu Quoc Dat;
quocdat181@yahoo.com

ABSTRACT

Objective The goal of this study was to describe the burden of disease and in-hospital mortality among patients admitted to the critical care units (CCUs) in Vietnam.

Design Retrospective study.

Setting The whole 1-year data of admissions to CCUs were collected from 34 hospitals from January to December 2018.

Participants A total of 44 013 episodes of admission to CCUs were analysed.

Primary outcome We used International Classification of Diseases-11 codes to assess the primary diagnosis associated with admissions and in-hospital mortality. Years of life lost (YLL) measure was further used to estimate the burden of disease.

Results The 0–5 years and ≥70 years age groups accounted for 14.8% (6508/44 013) and 26.1% (11 480/44 013) of all admissions, respectively. The most common diagnoses were diseases of the respiratory system (27.8% or 12 255/44 013), followed by unclassified symptoms, signs or clinical findings (13% or 5712/44 013), and diseases of the circulatory system (12.2% or 5380/44 013). Among 28 311 patients with available outcome data, 1681 individuals (5.9%) died during the hospitalisation. The in-hospital mortality rate increased with age, from 2.8% (86/3105) in under 5 years old age group to 23.1% (297/1288) in over 90-year age group. Diseases of the respiratory system was the leading causes of death in term of number of deaths (21.8% or 367/1681 of all deaths). Diagnosis of sepsis was associated with the highest in-hospital mortality (36.8%). The overall YLL under the age of 75 were 1287 per 1000 patients.

Conclusions CCUs in Vietnam faced wide differences in the burden of diseases. Sufficient infrastructure and adequate multidisciplinary training are essential to ensure the appropriate response to the current needs of population.

INTRODUCTION

Emergency care is an essential service of the hospitals which serves as a first point of contact to provide immediate treatment for individual with acute illness and life-threatening needs. Over 50% of mortality and over 40%

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ We conducted a large multicentre study on the pattern of admission and frequency of diagnosis in the critical care units in Vietnam, a lower-middle-income country.
- ⇒ We present mortality, cause of admission and the years of life lost due to premature death in critical care units in provincial and district hospitals in Vietnam.
- ⇒ The major limitation relates to the accuracy of the diagnosis and diagnostic coding in the study hospitals.

of all morbidities worldwide can be contributed by emergency medical conditions.¹ In May 2019, the 63rd World Health Assembly recognised that existing data do not provide adequate support for effective planning and resource allocation for emergency care and urged member states to identify the local acute illness burden for improving quality of emergency care.²

The International Statistical Classification of Diseases and Related Health Problems (often referred to as the International Classification of Diseases, ICD) has been used in more than 120 countries to monitor the incidence and prevalence of diseases.³ Vietnam has implemented the ICD since 1998 as a tool for reporting morbidity, cause of death and billing for health insurance in hospitals. The burden of disease in Vietnam has changed rapidly over the past two decades with a marked shift towards the predominance of non-communicable diseases. Between 2000 and 2017, the share of communicable disease has reduced a half (24%–14%), the share of non-communicable disease has increased almost 10% (63%–74%) while the share of injuries still remained stable at around 12%–13% (all measured in the disability-adjusted life-year

(DALY)).⁴ However, the research specifically measuring the burden of diseases in emergency departments (EDs) or intensive care capable EDs (referred to as critical care units or CCUs) in Vietnam has been limited. Recently, some studies on the pattern of hospitalisation, the causes of death or the hospital death certification have been performed,^{5–7} but none of these studies has provided a comprehensive description of the patterns of diagnosis or measured the premature death by the years of life lost (YLL) metric in patients with admission to CCUs in the country.

The lack of infrastructure and human resource brings significant challenges to the delivery of critical care in low-income and middle-income countries (LMICs).⁸ The emergence of COVID-19 has put further pressure on the critical care system in these countries that already struggling with the high burden of admission to the CCUs in the pre-COVID-19 era. An understanding of the conditions that involve an admission to CCU will allow policy-makers and planners to implement interventions that are effective in the national contexts to meet the care quality needs of the rapidly expanding ageing populations and to prevent the disruption of the critical care services in the pandemic. Our aim was to describe the frequency of diagnostic categories, in-hospital mortality, cause of death and the burden of premature mortality measuring by YLL in CCUs in Vietnam in 2018.

METHODS

Study design and participants

This was a retrospective study of all episodes of CCU admissions to selected provincial and district hospitals in Vietnam from 1 January 2018 to 31 December 2018. We purposely selected 5 out of 63 provinces to represent 5 geographical areas in Vietnam. In each province, we invited all CCUs in hospitals in the catchment area.

Data were collected from the electronic hospital management system, including patient's demographic information, diagnosis, dates of admission, length of stay and outcome at discharge. Diagnosis of individual patients in each episode of admission were originally recorded in the ICD, 10th revision (ICD-10) codes. Because the ICD, 11th revision (ICD-11) came into effect in January 2022, for the monitoring of any change in the pattern of diseases responsible for admission in the future, all ICD-10 codes were translated to the ICD-11 codes (version 05/2021) using WHO mapping tables with the correspondence between ICD-10 and ICD-11 codes and link the ICD-11 codes with their corresponding associated descriptive labels. Two independent investigators (VQD and BTKL) manually reviewed any non-unique code or discrepancies in mapping between ICD-10 and ICD-11 which were eventually resolved by the consensus of two reviewers with the reference to the majority of code assignments by treating doctors.

The outcomes from the hospitalisation were defined as death in patients who died in the hospital or was going

home to die imminently after hospital discharge (palliative discharge). Palliative discharge is a common practice in Vietnam when a patient is in a moribund state and expresses a wish to die at home. Deterioration was defined as any worsening of clinical conditions by the treating doctor's judgement at the time of discharge. The data on the outcomes at CCU discharge and interventions during CCU stay were not available for analysis.

Statistical analysis

For the analysis of pattern of admission, we included all episodes of admission. Consequently, one patient might have multiple episodes of admission to CCUs during a year. For each episode, the primary diagnosis which was defined as the main reason for CCU admission was counted. The primary diagnosis for that episode may be the same or different from the previous episode of admission. Individual patients were identified within the datasets by matching on full name, gender, date of birth and insurance numbers when they were available). The in-hospital mortality was calculated by dividing the number of deaths by the total number of individual patients.

YLL is a measure of premature mortality which is calculated from the number of deaths at each age multiplied by a standard life expectancy for the age at which death occurs.⁹ Vietnam's standard life expectancy at birth in 2019 for both sexes were 73.7 years old¹⁰ and to compare with other studies, we considered premature death were deaths under 75 years old. We also calculated the average YLL by dividing total of YLL by the number of premature death and the crude YLL rate by dividing total of YLL by the total number of patients aged under 75 years (expressed per 1000 persons).

We presented categorical variables as frequencies and percentages, while continuous variables as medians and IQR. We used the χ^2 test or Fisher's exact test to compare two binomial proportions. Statistical analyses were performed using SPSS V.27 (IBM). Significant differences were considered if two-sided p values were <0.05.

Patient and public involvement

Patients and the public were not involved in the design, recruitment or conduct of the study.

RESULTS

Between 1 January 2018 and 31 December 2018, there were 44013 admission episodes to CCUs from 34 hospitals in 5 provinces (namely Hanoi, Thai Nguyen, Ha Nam, Kon Tum and Can Tho), corresponding to 38534 unique patients. There were 3647 (9.5%) patients with two or more admission in the study year (figure 1). The median age was 53 years (IQR) 25–70) and included 53.4% male. The majority of CCU admissions in this study (31 569, 71.7%) were at the district hospitals. The median length of stay was 5 (IQR 2–8) days. A total of 1764/36 518 admissions (4.8%) had a length of stay of more than 14 days. The hospitalisation outcome was available for 28




Figure 1 Flow chart of CCU admission and outcome. CCU, critical care unit.

311/38 534 (73.5%) patients. Overall survival to hospital discharge was 94.1% (26 630/28 311).

Admission diagnosis in CCUs

Of 44013 episodes of admission to CCUs, the most common diagnosis were respiratory diseases (12255 episodes, 27.8%) in which, 38.7% was certain lower respiratory tract diseases (blockL1-CA2) and 34.4% was lung infections (blockL1-CA4), followed by unclassified symptoms, signs or clinical findings (5712 episodes or 13.0%) (51.8% was of digestive system or abdomen, blockL1-MD8), circulatory system diseases (5380 episodes,

12.2%) (51,2% was hypertensive diseases, blockL1-BA0), diseases of the nervous system (5019 episodes, 11.4%) (85.4% was cerebrovascular diseases, blockL1-8B0) and injury, poisoning or consequences of external causes (4186 episodes or 9.5%) (39.7% was harmful effect off substances, blockL1-NE6). The percentage of diagnosis in episodes of CCU admission varied by age groups (figure 2). The number of admissions peaked in children from 5 years old and younger and was lowest in the 5–10 years age group (figure 3). The respiratory disease was the leading cause of admission in the under 10 year age




Figure 2 Diagnosis on admission to CCU by age groups. CCU, critical care unit.

group (64% of admissions) and age groups of 61 years and older (29% of admissions) (figure 2).

The number of admissions in all study hospitals fluctuated across months but mostly driven by the admission by respiratory diseases (online supplemental figure 1). The proportions of admission to CCUs were significantly higher in provincial hospitals than in district hospital in regard to diseases of the nervous system (18.7% vs 8.5%, $p < 0.001$), certain infectious or parasitic diseases (8.9% vs 3.4%, $p < 0.001$), injury/poisoning (10.3% vs 9.2%, $p = 0.00$), endocrine, nutritional or metabolic diseases (3.6% vs 2.2%, $p < 0.001$) and neoplasms (2% vs 0.7%, $p < 0.001$) (figure 4).

Outcome of hospitalisation

The median length of stay for all admissions was 5 days (IQR 2–8). Among 28 311 patients with available outcome data and, 1681/28 311 (5.9%) died. The median age of death was 68 years (IQR 47–86) and 63.3% of death were in men. Men had significantly higher in-hospital mortality rate than women (3.8% vs 2.6%, $p < 0.001$). The median duration of hospital stay for the survivors and death cases was 4 (IQR: 2–8) and 3 (IQR: 2–7), respectively. More than three quarters of deaths (1290/1681 or 76.7%) occurred in provincial hospitals. The all-cause in-hospital mortality rates by age groups and mortality rate for selected groups of diagnosis are presented in figure 3 and online supplemental table 1. The most common causes of death by

ICD-11 block codes and chapter codes are shown in table 1 and online supplemental table 2, respectively.

The in-hospital mortality in patients over 75 years was 662/6,140 (10.8%). In total, 28 135 000 years were lost among patient admitted to the study CCUs. This yielded an overall YLL under the age of 75 years of 1287 per 1000 people. Table 1 presents the top 10 causes of death by ICD-11 block codes. Cerebrovascular diseases rank the first in the number of deaths (303/1681 or 18%) and the 11 when measuring in in-hospital mortality (8.7%) while sepsis ranked as the first specified cause of death in terms of in-hospital mortality (36.8%) and ranked fourth in term of death counts.

DISCUSSION

This study presents for the first time, data to quantify the burden of diseases in CCUs, for the most common reasons of admission and the in-hospital mortality of patients admitted to CCUs in Vietnam. Few studies on admission to the CCUs in LMICs have been reported earlier.

Burden of diseases in Vietnam has changed significantly in the past 20 years.⁴ The communicable diseases burden (measured in DALYs) decreased rapidly from 37% in 1990 to 14% in 2017. However, in the critical care settings, our study shows that the infectious diseases remain a major emergency medical condition and are the prime cause of




Figure 3 Number of admission and In-hospital mortality by age groups.

morbidity and mortality in terms of number of admission and in terms of YLL.

Our findings on the difference distribution of diseases in CCUs in provincial and district hospitals reflected the variation of healthcare services and capacity of clinical management. In 2019, we conducted a survey on the capacity and use of diagnostics and treatment for patients

with severe acute respiratory infections in 48 hospitals in Vietnam (including 32 hospitals in this study) at the same time as this study. Significant differences were observed in the district hospitals when compared with provincial hospitals in regard to availability of the microbiological diagnostics (9.4% vs 81.2%, $p < 0.001$), inflammatory markers of C reactive protein (31.2% vs 54.2%,




Figure 4 Cause of admission to CCUs by level of care. CCU, critical care unit.

Table 1 The top 20 causes of death in terms of years of life lost (YLL) and numbers of deaths by ICD-11 block level one codes

Diagnosis	No and proportion of death cases	In-hospital mortality (n,%)	Total YLL75	Proportion of YLLs75	YLL75 per 1000 patients	Length of stay to death (median, IQR)
Cerebrovascular diseases	303 (18.0%)	8.7%	3 211 000	12.3%	1380	3 (2–6)
Miscellaneous	259 (15.4%)	2.3%	5 686 000	38.2%	614	2 (1–6)
Lung infections	201 (12.0%)	7.2%	2 875 000	9.5%	1440	7 (3–15)
Sepsis	149 (8.9%)	36.8%	1 991 000	2.6%	8127	4 (2–11)
Harmful effects of substances	87 (5.2%)	6.1%	3 534 000	5.0%	2544	1 (1–2)
Respiratory failure	79 (4.7%)	18.1%	365 000	1.6%	1714	9 (5–15)
General symptoms, signs or clinical findings	67 (4.0%)	13.5%	1 507 000	2.2%	3712	2 (1–4.5)
Injuries to the head	66 (3.9%)	15.6%	1 140 000	1.6%	3167	2 (2–4)
Certain lower respiratory tract diseases	63 (3.7%)	2.1%	443 000	10.2%	217	4 (2–10)
Symptoms, signs or clinical findings of the digestive system or abdomen	61 (3.6%)	5.5%	994 000	4.0%	1062	3 (2–6)
Malignant neoplasms, except primary neoplasms of lymphoid, haematopoietic, central nervous system or related	50 (3.0%)	18.0%	587 000	1.0%	3073	6 (3–9)
Symptoms, signs or clinical findings of the circulatory system	50 (3.0%)	31.8%	1 060 000	0.7%	8217	2 (1–6)
Heart failure	39 (2.3%)	8.4%	280 000	1.6%	1466	3 (1–5)
Ischaemic heart diseases	37 (2.2%)	5.2%	78 000	2.5%	158	3 (1–5)
Diseases of liver	36 (2.1%)	11.2%	549 000	1.1%	1861	2.5 (1.5–7.5)
Injuries involving multiple body regions	29 (1.7%)	28.2%	439 000	0.4%	6362	2 (1–6)
Symptoms, signs or clinical findings of the respiratory system	28 (1.7%)	8.4%	706 000	1.1%	3017	2 (1–4)
Mental or behavioural symptoms, signs or clinical findings	22 (1.3%)	13.3%	363 000	0.8%	2771	2 (1–2)
Allergic or hypersensitivity conditions	20 (1.2%)	3.3%	890 000	2.2%	1537	2 (1–2)
Infection, unspecified	19 (1.1%)	31.7%	741 000	0.2%	14 529	2 (2–4)
Epilepsy or seizures	16 (1.0%)	5.0%	696 000	1.1%	2275	1.5 (1–2)
Total	1681 (100.0%)	5.9%	28 135 000	100.0%	1287	3 (2–7)

ICD, International Classification of Disease.

$p < 0.001$) and procalcitonin (3.1% vs 25%, $p < 0.001$) and mechanical ventilation (40.6% vs 60.4%, $p < 0.001$).¹¹ Provincial hospitals in Vietnam are the secondary hospitals which do not only treat referred patients from lower level of care (district hospitals or commune centres) but also admit patients from the community. Additionally, patient can directly self-refer to higher level of hospitals which resulting in under-utilisation of district hospitals and overcrowding in provincial hospitals, especially for given specialities which requires higher levels of skills and capacity.

Our findings are partly aligned with previous reports in all-cause mortality in both sexes and all ages showing that cardiovascular (ischaemic heart disease, stroke), respiratory diseases and injuries were the significant challenge. WHO's report in 2019 showed that ischaemic heart

disease, stroke, neonatal conditions, chronic obstructive pulmonary disease (COPD) and lower respiratory infections were the top five causes of death in term of death counts in LMICs.¹² At the global level in 2019, the top three diseases with highest burden (measuring in DALY) were neonatal disorder (32.9% of global DALYs), lower respiratory infections (11% of global DALYs) and diarrhoeal diseases (9.3% of global DALYs) in children younger than 10 years and were ischaemic heart diseases (16.2% of global DALY), stroke (13% of global DALY) and COPD (8.5% of global DALY) in patients from 75 years and older.¹³ In Vietnam, stroke (164.8 deaths per 100 000), ischaemic heart disease (95.3 deaths per 100 000), COPD (37.2 deaths per 100 000) and diabetes mellitus (35.1 deaths per 100 000) and road injuries (30.5 deaths per 100 000) were in the top-five rankings for total

number of death in 2019.¹⁴ Sepsis and harmful effects of substances were not listed in the top five causes of death in the general population, but we found that these were important conditions in emergency care settings.

The overall mortality in our study was higher than previous report. In a systematic review of data from 65 EDs in 59 LMICs countries between 2013 and 2014, the median mortality was 1.8% (IQR 0.5%–5.1%), highest in sub-Saharan Africa (median 3.4%, IQR: 0.5%–6.3%) and lowest in South Asia, East Asia and Pacific (0.3, IQR 0.2%–0.8%).¹⁵

In our study, patients aged 75 years or older presented 18.9% of all admission at CCUs despite the fact that they comprised 3.2% of general population.¹⁶ The percentage of population older than 75 years in Vietnam is expected to increase to 4.2% by 2030 and 6.6% by 2040.¹⁶ This will increase the demand for critical care resources and requires further evaluation to guide healthcare decision-making and changes in organisational models in CCU and delivery of critical care services during a pandemic.

The design of multicentre study, large population and data collection for the whole year of 2018 are the strengths of our study. The major limitations related to the diagnosis description and diagnostic coding by doctors in study hospitals. Data quality of diagnosis may vary between hospitals due to the accurate clinical diagnosis of disease depends on doctor's knowledge and skills, available laboratory investigations and diagnostic imaging procedures. However, there are differences in the capacity of clinical management and use of diagnostics between provincial and district hospitals in the country.¹¹ In Vietnam, treating doctors are responsible for assigning the ICD diagnostic code to each patient discharged from the hospital. The complex nature of ICD coding, patient's condition in critical care settings and the lack of coding experience pose further challenges for the accuracy and reliability of the diagnosis given across hospitals. Another limitation was the missing information on outcome for 26.5% (10 223/38 543) of patients because these were not actually entered into the electronic hospital management system. These missing data can cause bias in the estimation of in-hospital mortality.

Currently, there is limited cost-effectiveness analysis performed for critical care services in Vietnam. The results of this study in regard to the variety of diagnoses and outcomes of admission will direct the future studies on the establishment of CCU registry, the effectiveness of CCU treatments and estimates of the total cost to save a life-year. This information will be helpful to develop policies and strategies to make better-informed decisions in resource-constrained settings in the face of an ageing population in Vietnam.

Conclusion

Our study indicates a variation of needs of emergency care by age groups and levels of care among patients admitted to CCUs in provincial and district hospitals in Vietnam. These findings support the need to provide

multidisciplinary team training for clinicians in CCUs and the future studies should further assess the emergency and critical care capacity needs, ability of the healthcare setting to provide the emergency services and the quality of care (including minimal set of indicators) and gaps to mobilise the resources required for live saving during a pandemic.

Acknowledgements We are very grateful to the patients and staff at the participating hospitals in Ha Noi, Thai Nguyen, Ha Nam, Kon Tum and Can Tho. We would like to thank Dr Satoko Otsu and Vu Quang Hieu (WHO, Vietnam Country Office) for their tireless support and contribution to strengthening the capacity for clinicians in Vietnam and thank Dr Nguyen The Hung (National Hospital for Tropical Diseases) for his initial support in the implementation of this study.

Contributors VQD conceived and designed the study. VQD analysed the data and wrote the first draft of the report. VQD, BTKL and GBK involved to the acquisition and interpretation of data. All authors contributed to and approved the final report. VQD is guarantor.

Funding This work was supported by Viet Nam-WHO Country Office.

Disclaimer The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study was approved by the institutional review board (IRB) in the Hanoi Medical University (59/GCN-DDNCYSH-DHYHN).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Vu Quoc Dat <http://orcid.org/0000-0002-5904-5970>


Bui Thi Khanh Linh <http://orcid.org/0000-0002-7337-5925>

REFERENCES

- 1 Razzak J, Usmani MF, Bhutta ZA. Global, regional and national burden of emergency medical diseases using specific emergency disease indicators: analysis of the 2015 Global Burden of Disease Study. *BMJ Glob Health* 2019;4:e000733.
- 2 World Health Assembly. *Emergency care systems for universal health coverage: ensuring timely care for the acutely ill and injured*. Geneva: World Health Organization, 2019.
- 3 Executive Board. *International classification of diseases: international statistical classification of diseases and related health problems: update on the eleventh revision: report by the director-general*. Geneva: World Health Organization, 2018.
- 4 Glinskaya E, Feige ADK, Hoang VTL. *Vietnam - adapting to an aging society*. Washington, DC: World Bank Group, 2021.
- 5 Nguyen NTT, Dien TM, Schindler C, et al. Childhood hospitalisation and related deaths in Hanoi, Vietnam: a tertiary hospital database analysis from 2007 to 2014. *BMJ Open* 2017;7:e015260.



- 6 Hong TT, Phuong Hoa N, Walker SM, *et al.* Completeness and reliability of mortality data in Viet Nam: implications for the National routine health management information system. *PLoS One* 2018;13:e0190755.
- 7 Walton M, Harrison R, Chevalier A, *et al.* Improving hospital death certification in Vietnam: results of a pilot study of injury-related fatalities. *Inj Prev* 2018;24:324–31.
- 8 Murthy S, Leligdowicz A, Adhikari NKJ. Intensive care unit capacity in low-income countries: a systematic review. *PLoS One* 2015;10:e0116949.
- 9 World Health Organization. WHO methods and data sources for global burden of disease estimates 2000-2019. In: *Analytics DoDa, division of data AaDfi*. Geneva: World Health Organization, 2020.
- 10 World Health Organization. Life expectancy and healthy life expectancy. Data by country 2020, 2021. Available: <https://apps.who.int/gho/data/node.main.688> [Accessed 6 Jan 2022].
- 11 Dat VQ, Nguyen TH, Kim BG, *et al.* Capacity and use of diagnostics and treatment for patients with severe acute respiratory infections in the pre-COVID-19 era in district and provincial hospitals in Viet Nam. *Western Pac Surveill Response JWPSAR* 2021;12:28–36.
- 12 World Health Organization. The top 10 causes of death 2019, 2020. Available: <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death> [Accessed 11 Dec 2021].
- 13 GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the global burden of disease study 2019. *Lancet* 2020;396:1204–22.
- 14 World Health Organization. Global health estimates 2020: deaths by cause, age, sex, by country and by region, 2000-2019, 2020. Available: <https://www.who.int/data/gho/data/themes/mortality-and-global-health-estimates/ghe-leading-causes-of-death> [Accessed 13 Dec 2021].
- 15 Obermeyer Z, Abujaber S, Makar M, *et al.* Emergency care in 59 low- and middle-income countries: a systematic review. *Bull World Health Organ* 2015;93:577–86.
- 16 General Statistic Office, United Nations Population Fund. *Vietnam population projection for the period 2019-2069*. Hanoi: General Statistic Office, United Nations Population Fund, 2020.



Supplementary Figure 1. Pattern of admissions over months.

Supplementary Table 1. Mortality and years of life lost (YLL) grouped by ICD-11 chapter codes

Diagnosis	Number and proportion of death cases	In-hospital mortality (n,%)	Total YLL75	Proportion of YLLs75	YLL75 per 1000 patients	Length of stay to death (median, IQR)
Diseases of the respiratory system	367 (21.8%)	4.8%	4506000	25.9%	813	6 (3-13)
Diseases of the nervous system	324 (19.3%)	8.1%	3994000	14.0%	1432	2 (2-6)
Symptoms, signs or clinical findings, not elsewhere classified	241 (14.3%)	9.4%	5141000	9.8%	2476	2 (1-4)
Injury, poisoning or certain other consequences of external causes	210 (12.5%)	8.4%	5846000	8.8%	2557	2 (1-2)
Certain infectious or parasitic diseases	188 (11.2%)	14.0%	3218000	5.8%	3056	4 (2-10)
Diseases of the circulatory system	112 (6.7%)	2.8%	817000	13.4%	314	3 (1-5)
Diseases of the digestive system	83 (4.9%)	3.9%	1221000	7.1%	683	3 (2-8)
Neoplasms	59 (3.5%)	17.3%	809000	1.3%	3385	5 (2-9)
Miscellaneous	31 (1.8%)	3.5%	963000	3.1%	1259	2 (1-5)
Endocrine, nutritional or metabolic diseases	22 (1.3%)	2.4%	217000	3.3%	276	2 (1-3)
Diseases of the immune system	20 (1.2%)	3.3%	890000	2.2%	1532	2 (1-2)
Diseases of the genitourinary system	13 (0.8%)	3.0%	168000	1.5%	487	5 (2-11)
External causes of morbidity or mortality	8 (0.5%)	1.8%	223000	1.6%	556	3.5 (1.5-4.5)
Diseases of the ear or mastoid process	2 (0.1%)	0.4%	87000	1.8%	192	1 (1-1)
Pregnancy, childbirth or the puerperium	1 (0.1%)	0.7%	35000	0.5%	241	1 (1-1)
Total	1681 (100.0%)	5.9%	28135000	100.0%	1287	3 (2-7)

Supplementary Table 2. The number of deaths by ICD 4-character codes

	Causes of death	N (%)
CHAPTER 01 Certain infectious or parasitic diseases	Viral intestinal infections, unspecified	1 (0.06%)
	Gastroenteritis or colitis without specification of infectious agent	1 (0.06%)
	Tuberculosis of the respiratory system	2 (0.12%)
	Bacterial cellulitis, erysipelas and lymphangitis	1 (0.06%)
	Other specified pyogenic bacterial infection of skin and subcutaneous tissue	1 (0.06%)
	Tetanus	1 (0.06%)
	Bacterial infection of unspecified site	1 (0.06%)
	Viral encephalitis not elsewhere classified	2 (0.12%)
	Viral meningitis not elsewhere classified	1 (0.06%)
	Infectious abscess of the central nervous system	1 (0.06%)
	Dengue fever, unspecified	1 (0.06%)
	Acute viral hepatitis	4 (0.24%)
	Viral hepatitis, unspecified	2 (0.12%)
	Phaeohyphomycosis	1 (0.06%)
	Sepsis without septic shock	23 (1.37%)
	Sepsis with septic shock	126 (7.50%)
	Infection, unspecified	19 (1.13%)
CHAPTER 02 Neoplasms	Primary neoplasms of brain	6 (0.36%)
	Malignant haematopoietic neoplasms without further specification	1 (0.06%)
	Malignant neoplasms of other or ill-defined sites in the lip, oral cavity or pharynx	2 (0.12%)
	Malignant neoplasms of oesophagus	5 (0.30%)
	Malignant neoplasms of stomach	2 (0.12%)
	Malignant neoplasms of colon	5 (0.30%)
	Malignant neoplasm of pancreas	1 (0.06%)
	Malignant neoplasms of liver or intrahepatic bile ducts	11 (0.65%)
	Malignant neoplasms of gallbladder	1 (0.06%)
	Malignant neoplasms of bronchus or lung	10 (0.59%)
	Malignant neoplasms of thymus	1 (0.06%)
	Malignant neoplasms of heart, mediastinum or non-mesothelioma of pleura	2 (0.12%)
	Malignant neoplasms of breast, unspecified	3 (0.18%)
	Malignant neoplasms of prostate	1 (0.06%)
	Malignant neoplasms of bladder	1 (0.06%)
	Malignant neoplasms of thyroid gland	1 (0.06%)
	Malignant neoplasm metastasis in lung	4 (0.24%)
	Carcinoma in situ of bladder	1 (0.06%)
Benign osteogenic tumours	1 (0.06%)	
CHAPTER 03 Diseases of the blood or blood-forming organs	Anaemia due to chronic disease	1 (0.06%)
	Anaemias or other erythrocyte disorders, unspecified	8 (0.48%)
	Anaphylaxis	9 (0.54%)

	Causes of death	N (%)
CHAPTER 04 Diseases of the immune system	Allergic or hypersensitivity conditions of unspecified type	11 (0.65%)
CHAPTER 05 Endocrine, nutritional or metabolic diseases	Type 1 diabetes mellitus	3 (0.18%)
	Type 2 diabetes mellitus	3 (0.18%)
	Hypoglycaemia without associated diabetes	5 (0.30%)
	Increased secretion of glucagon	1 (0.06%)
	Protein deficiency	1 (0.06%)
	Hyperosmolality or hypernatraemia	1 (0.06%)
	Acidosis	1 (0.06%)
	Hyperkalaemia	1 (0.06%)
	Hypokalaemia	1 (0.06%)
CHAPTER 06 Mental, behavioural or neurodevelopmental disorders	Disorders of fluid, electrolyte or acid-base balance, unspecified	5 (0.30%)
	Developmental learning disorder	1 (0.06%)
	Dissociative disorders, unspecified	4 (0.24%)
	Disorders due to use of dissociative drugs including ketamine and phencyclidine [PCP]	1 (0.06%)
	Delirium	1 (0.06%)
CHAPTER 08 Diseases of the nervous system	Mental, behavioural or neurodevelopmental disorders, unspecified	2 (0.12%)
	Seizure due to acute causes	2 (0.12%)
	Types of seizures	11 (0.65%)
	Epilepsy or seizures, unspecified	3 (0.18%)
	Intracerebral haemorrhage	133 (7.91%)
	Subarachnoid haemorrhage	3 (0.18%)
	Intracranial haemorrhage, unspecified	32 (1.90%)
	Cerebral ischaemic stroke	72 (4.28%)
	Stroke not known if ischaemic or haemorrhagic	61 (3.63%)
	Late effects of cerebrovascular disease	2 (0.12%)
	Myelitis	1 (0.06%)
	Motor neuron disease	1 (0.06%)
	Hereditary sensory or autonomic neuropathy	1 (0.06%)
	Increased intracranial pressure	1 (0.06%)
Disorders of the meninges excluding infection	1 (0.06%)	
CHAPTER 10 Diseases of the ear or mastoid process	Disorders of vestibular function	2 (0.12%)
CHAPTER 11 Diseases of the circulatory system	Essential hypertension	8 (0.48%)
	Hypertensive heart disease	1 (0.06%)
	Hypotension, unspecified	5 (0.30%)
	Angina pectoris	1 (0.06%)
	Acute myocardial infarction	36 (2.14%)
	Pulmonary thromboembolism	1 (0.06%)
	Pulmonary hypertension	1 (0.06%)
	Pulmonary heart disease or diseases of pulmonary circulation, unspecified	5 (0.30%)
	Pericardial effusion	1 (0.06%)
	Mitral valve stenosis with insufficiency	1 (0.06%)

	Causes of death	N (%)
	Myocarditis	1 (0.06%)
	Conduction disorders	1 (0.06%)
	Ventricular tachyarrhythmia	2 (0.12%)
	Cardiac arrhythmia, unspecified	5 (0.30%)
	Congestive heart failure	3 (0.18%)
	Heart failure, unspecified	36 (2.14%)
	Acute arterial occlusion	1 (0.06%)
	Aortic aneurysm or dissection	1 (0.06%)
	Diseases of arteries or arterioles, unspecified	1 (0.06%)
	Deep vein thrombosis	1 (0.06%)
CHAPTER 12 Diseases of the respiratory system	Acute tonsillitis	1 (0.06%)
	Acute laryngitis or tracheitis	1 (0.06%)
	Acute upper respiratory infections of multiple and unspecified sites	1 (0.06%)
	Bronchitis	2 (0.12%)
	Chronic obstructive pulmonary disease	52 (3.09%)
	Asthma	8 (0.48%)
	Bronchiectasis	1 (0.06%)
	Pneumonia	198 (11.78%)
	Abscess of lung or mediastinum	2 (0.12%)
	Lung infections, unspecified	1 (0.06%)
	Pneumonitis due to solids and liquids	1 (0.06%)
	Respiratory conditions due to inhalation of chemicals, gases, fumes or vapours	3 (0.18%)
	Acute respiratory distress syndrome	2 (0.12%)
	Pulmonary oedema	11 (0.65%)
	Pneumothorax	2 (0.12%)
	Haemothorax	1 (0.06%)
Pleural effusion	1 (0.06%)	
Respiratory failure	79 (4.70%)	
CHAPTER 13 Diseases of the digestive system	Gastro-oesophageal reflux disease	1 (0.06%)
	Vascular disorders of the oesophagus	3 (0.18%)
	Gastritis	4 (0.24%)
	Gastric ulcer	5 (0.30%)
	Obstruction of small intestine	7 (0.42%)
	Appendicitis	3 (0.18%)
	Obstruction of large intestine	1 (0.06%)
	Polyp of large intestine	1 (0.06%)
	Infectious liver disease	1 (0.06%)
	Acute or subacute hepatic failure	2 (0.12%)
	Hepatic fibrosis or cirrhosis	15 (0.89%)
	Autoimmune liver disease	1 (0.06%)
	Certain specified diseases of liver	17 (1.01%)
	Cholelithiasis	2 (0.12%)
	Cholangitis	2 (0.12%)
	Acute pancreatitis	8 (0.48%)
	Peritonitis	6 (0.36%)
	Inguinal hernia	1 (0.06%)
Indeterminate colitis	1 (0.06%)	

	Causes of death	N (%)
	Diseases of the digestive system, unspecified	2 (0.12%)
CHAPTER 14 Diseases of the skin	Lichen planus	1 (0.06%)
	Skin disease of unspecified nature	1 (0.06%)
CHAPTER 16 Diseases of the genitourinary system	Pyonephrosis	1 (0.06%)
	Acute kidney failure	1 (0.06%)
	Chronic kidney disease	7 (0.42%)
	Kidney failure, unspecified	2 (0.12%)
	Urinary tract infection, site not specified	2 (0.12%)
CHAPTER 18 Pregnancy, childbirth or the puerperium	Complications following abortion, ectopic or molar pregnancy	1 (0.06%)
CHAPTER 19 Certain conditions originating in the perinatal period	Sepsis of foetus or newborn	1 (0.06%)
	Respiratory failure of newborn	1 (0.06%)
	Transitory disorders of carbohydrate metabolism specific to foetus or newborn	1 (0.06%)
	Certain specified transitory neonatal electrolyte or metabolic disturbances	1 (0.06%)
CHAPTER 20 Developmental anomalies	Congenital hydrocephalus	1 (0.06%)
	Congenital anomaly of a ventricle or the ventricular septum	1 (0.06%)
	Structural developmental anomalies of gallbladder, bile ducts or liver	1 (0.06%)
CHAPTER 21 Symptoms, signs or clinical findings, not elsewhere classified	Symptoms, signs or clinical findings involving consciousness	22 (1.31%)
	Abnormalities of heart beat	2 (0.12%)
	Cardiac arrest	48 (2.86%)
	Abnormalities of breathing	2 (0.12%)
	Haemorrhage from throat	1 (0.06%)
	Haemoptysis	2 (0.12%)
	Respiratory arrest	23 (1.37%)
	Abdominal or pelvic pain	4 (0.24%)
	Nausea or vomiting	1 (0.06%)
	Clinical manifestations of the digestive system	56 (3.33%)
	Cachexia	1 (0.06%)
	Fever of other or unknown origin	4 (0.24%)
	Haemorrhage, not elsewhere classified	1 (0.06%)
	Shock	60 (3.57%)
	Syncope and collapse	1 (0.06%)
	Other sudden death, cause unknown	1 (0.06%)
Other ill-defined and unspecified causes of mortality	9 (0.54%)	
Multi organ failure	3 (0.18%)	
CHAPTER 22 Injury, poisoning or certain other consequences of external causes	Injury of cranial nerves	1 (0.06%)
	Intracranial injury	65 (3.87%)
	Superficial injury of thorax	2 (0.12%)
	Injury of blood vessels at abdomen, lower back or pelvis level	1 (0.06%)
	Injury of intra-abdominal organs	5 (0.30%)
	Traumatic amputation of forearm	1 (0.06%)

	Causes of death	N (%)
	Fracture of femur	1 (0.06%)
	Superficial injury of ankle or foot	1 (0.06%)
	Crushing injuries involving multiple body regions	1 (0.06%)
	Unspecified multiple injuries	28 (1.67%)
	Burn of wrist or hand	1 (0.06%)
	Burns of external body surface, other specified site	1 (0.06%)
	Burns of multiple body regions	1 (0.06%)
	Harmful effects of drugs, medicaments or biological substances, not elsewhere classified	57 (3.39%)
	Harmful effects of or exposure to noxious substances, chiefly nonmedicinal as to source, not elsewhere classified	30 (1.78%)
	Effects of other specified external causes	3 (0.18%)
	Certain early complications of trauma, not elsewhere classified	11 (0.65%)
CHAPTER 23 External causes of morbidity or mortality	Unintentional exposure to or harmful effects of pesticides	1 (0.06%)
	Intentional self-harm by threat to breathing by suffocation from object covering mouth or nose	1 (0.06%)
	Intentional self-harm by and exposure to other and unspecified drug, medicament and biological substance	3 (0.18%)
	Intentional self-harm by exposure to or harmful effects of pesticides	1 (0.06%)
	Intentional self-harm by exposure to or harmful effects of other or unspecified substances chiefly nonmedicinal as to so	1 (0.06%)
	Exposure to or harmful effects of undetermined intent of other or unspecified drugs, medicaments or biological substance	1 (0.06%)
CHAPTER 24 Factors influencing health status or contact with health services	Surgical or postsurgical states, unspecified	3 (0.18%)
	Personal history of malignant neoplasm	1 (0.06%)