

Table S2 Area under the receiver operating characteristic curve of the physician's clinical diagnosis of influenza in each subgroup

	AUC (95% CI)
Age (years)	
<6 (n = 37)	0.69 (0.49 to 0.88)
≥6 (n = 298)	0.78 (0.73 to 0.83)
Type of doctor	
Staff physician (family physicians) (n = 171)	0.76 (0.69 to 0.84)
Senior resident (family medicine) (n = 88)	0.83 (0.74 to 0.91)
Other doctors (n = 76)	0.75 (0.64 to 0.86)
Past medical history of influenza	
Yes (n = 201)	0.80 (0.72 to 0.88)
No (n = 134)	0.76 (0.69 to 0.82)
Influenza vaccination	
Yes (n = 122)	0.80 (0.72 to 0.88)
No (n = 213)	0.76 (0.63 to 0.89)
Medication taken prior to medical visit	
Yes (n = 201)	0.77 (0.69 to 0.85)
No (n = 134)	0.78 (0.71 to 0.84)
Severity of current symptoms compared to those of a common cold	
Severe (n = 158)	0.77 (0.69 to 0.84)
Similar (n = 138)	0.76 (0.68 to 0.84)
Mild (n = 39)	0.73 (0.57 to 0.90)
Duration from symptom onset to RIDT (≤12 h)	
Yes (n = 35)	0.69 (0.51 to 0.88)
No (n = 300)	0.79 (0.74 to 0.84)
Duration from symptom onset to RIDT (≤24 h)	
Yes (n = 119)	0.77 (0.69 to 0.86)
No (n = 216)	0.78 (0.72 to 0.85)
Type of influenza	
A (n = 87)	0.78 (0.72 to 0.84)
B (n = 75)	0.76 (0.70 to 0.82)

AUC, area under the curve; CI, confidence interval; RIDT, rapid influenza diagnostic test