

Table S1. Characteristics of children who enrolled and were interviewed compared to individuals who could not be contacted or declined participation.

Characteristic	Individuals enrolled and interviewed, n (%) [†] (n=18)	Could not be contacted or declined participation, (n, %) [†] (n=36)	p value
Current age, years (median [IQR])	12.1 (9.7–14.5)	14.2 (11.4–16.0)	0.14 [‡]
Current age, ≥13 years	8 (44)	21 (58)	0.50 [§]
Years at diagnosis (median [IQR])	0.5 (0.2–4.0)	0.5 (0.0–4.8)	0.89 [‡]
Age at diagnosis, ≥4 years	5 (28)	10 (28)	1.00 [§]
Years since diagnosis (median [IQR])	9.8 (7.3–13.6)	11.2 (8.5–13.7)	0.72 [‡]
Gender, female	6 (33)	17 (47)	0.50 [§]
Laterality of glaucoma, bilateral	11 (61)	34 (94)	0.004 [¶]
Self-reported ancestry, European	16 (89)	21 (60)*	0.06 [§]
Glaucoma subtype, primary**	12 (67)	28 (78)	0.51 [¶]
BCVA better eye, impaired (<0.5)	3 (17)	6 (17)	1.00 [¶]
Molecular diagnosis identified	9 (50)	9 (25)	0.13 [§]
Family history, first degree	4 (22)	17 (49)*	0.12 [§]

[†]: n (%) presented unless otherwise specified

[‡]Mann-Whitney U test

[§]Chi-square test with Yates' correction for continuity

[¶]Fisher exact test

*Data missing for one individual (n=1)

**Primary glaucoma includes primary congenital glaucoma and juvenile open-angle glaucoma

IQR: interquartile range; BCVA: best-corrected visual acuity