

BMJ Open Prevalence, nature and trajectory of dysphagia postoesophageal cancer surgery: a prospective longitudinal study protocol

Michelle Hayes ^{1,2}, Anna Gillman,¹ Brona Wright,³ Sean Dorgan,³ Ian Brennan,⁴ Margaret Walshe,¹ Claire Donohoe,⁵ John V Reynolds,^{5,6} Julie Regan ¹

To cite: Hayes M, Gillman A, Wright B, *et al.* Prevalence, nature and trajectory of dysphagia postoesophageal cancer surgery: a prospective longitudinal study protocol. *BMJ Open* 2022;**12**:e058815. doi:10.1136/bmjopen-2021-058815

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-058815>).

Received 30 October 2021
Accepted 06 June 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Ms Michelle Hayes;
mhayes6@tcd.ie

ABSTRACT

Introduction Dysphagia is a common problem following oesophagectomy, and is associated with aspiration pneumonia, malnutrition, weight loss, prolonged enteral feeding tube dependence, in addition to an extended in-hospital stay and compromised quality of life (QOL). To date, the prevalence, nature and trajectory of post-oesophagectomy dysphagia has not been systematically studied in a prospective longitudinal design. The study aims (1) to evaluate the prevalence, nature and trajectory of dysphagia for participants undergoing an oesophagectomy as part of curative treatment, (2) to determine the risk factors for, and post-operative complications of dysphagia in this population and (3) to examine the impact of oropharyngeal dysphagia on health-related QOL across time points.

Methods and analysis A videofluoroscopy will be completed and analysed on both post-operative day (POD) 4 or 5 and at 6-months post-surgery. Other swallow evaluations will be completed preoperatively, POD 4 or 5, 1-month and 6-month time points will include a swallowing screening test, tongue pressure measurement, cough reflex testing and an oral hygiene evaluation. Nutritional measurements will include the Functional Oral Intake Scale to measure feeding tube reliance, Malnutrition Screening Tool and the Strength, Assistance With Walking, Rise From a Chair, Climb Stairs and Falls questionnaire. The Reflux Symptom Index will be administered to investigate aerodigestive symptoms commonly experienced by adults post-oesophagectomy. Swallowing-related QOL outcome measures will be determined using the European Organisation for Research and Treatment of Cancer QLQ-18, MD Anderson Dysphagia Inventory and the Swallowing Quality of Life Questionnaire.

Ethics and dissemination Ethical approval has been granted by the Tallaght University Hospital/St. James' Hospital Research Ethics Committee (JREC), Dublin, Ireland (Ref. No. 2021-Jul-310). The study results will be published in peer-reviewed journals and presented at national and international scientific conferences.

INTRODUCTION

The incidence of oesophageal cancer has increased markedly in the western world over the last 50 years, with the rates of the

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This is the first prospective longitudinal study to evaluate the presence, nature and trajectory of dysphagia following oesophagectomy and the impact of dysphagia on quality of life over a 6-month timeframe.
- ⇒ Public and patient representatives have contributed to the study design including the selection of meaningful outcome measures across time points.
- ⇒ This study will take place in a national specialist centre for oesophageal cancer, involving multidisciplinary experts in oesophageal cancer and two patient and public representatives as co-researchers.
- ⇒ This prospective longitudinal study will evaluate dysphagia post-oesophagectomy, across all open surgical approaches (including transthoracic and transhiatal).
- ⇒ Limitations include the single-centre study design, excludes patients who undergo a complex post-operative course (eg, post-operative pulmonary complications) who may present with an oropharyngeal dysphagia, and the long-term time point is limited to 6-months post surgery.

pathological subtype of adenocarcinoma linked to an increased prevalence of obesity, gastro-oesophageal reflux disease and Barrett's oesophagus.¹ The mainstay of curative treatment is surgery, often combined with preoperative combination chemoradiotherapy, or perioperative chemotherapy as per the MAGIC/FLOT or CROSS regimens.² Surgery for oesophageal cancer is major, with up to 5% risk of mortality and over 50% risk of morbidity, irrespective of whether surgery is via open, minimally invasive or robotic-assisted approaches.³⁻⁷ Post-operative complications include pulmonary dysfunction, atrial fibrillation and anastomotic leak.⁸ Post-operative pulmonary complications (PPCs) are among the most serious postoperative challenges occurring between 15% and 40%

of patients post-oesophagectomy, impacting length of stay in critical care units, increasing overall hospital stay with significant cost implications.^{9,10} Malnutrition, weight loss and sarcopaenia are common after surgery or combination therapies.^{11,12}

Due to centralisation of services and enhanced recovery programmes, operative mortality has decreased.¹³ Furthermore, the 5-year survival rates among survivors of oesophageal cancer have improved in high-income countries.¹⁴ This has led to a shift in focus to improving survivorship in adults who have undergone curative treatment for oesophageal cancer.¹⁵ The health-related quality of life (HR-QOL) among oesophageal cancer survivors varies considerably. Symptoms known to impact long-term survivors and HR-QOL include coughing, reflux and deterioration in swallowing function.¹⁶

Dysphagia is the most common presenting symptom for the majority of patients with oesophageal cancer.¹⁷ Following oesophageal resection, dysphagia continues to present post-operatively alongside other complications, which may be a result of, or further exacerbated by PPCs, recurrent laryngeal nerve (RLN) damage, neo-oesophageal strictures and gastrointestinal reflux.^{18–20} Interventions for oesophageal dysphagia include oesophageal dilation, stenting and thermal and chemical ablation therapy.²¹ Dysphagia may be associated with aspiration pneumonia, malnutrition, prolonged feeding tube dependence and an extended inpatient hospital stay.²²

Dysphagia is highly associated with compromised quality of life (QOL) among oesophageal cancer survivors.¹⁶ One-year post-oesophagectomy, almost half of survivors' report eating restrictions and other symptoms include dry mouth, taste problems, difficulty swallowing saliva and choking.¹⁵ At two years post-oesophagectomy, eating difficulties and reluctance to eat in front of others has been associated with psychological distress.²³ In a recent cross-sectional cohort study on patient-reported outcomes post-oesophagectomy, long-term symptom burden is common in this patient group, with swallowing/conduit problems being one of the six main problems reported.²⁴ Ten years post-operatively, swallowing difficulties persist for half of survivors.²⁵

Despite the prevalence of dysphagia post-oesophagectomy as well as its impact on QOL, the prevalence, nature and trajectory of dysphagia has been poorly studied in oesophageal cancer. Some small studies have identified impairment of oropharyngeal structures post-operatively during videofluoroscopy (VFS). Swallowing impairment following resection have been reported to include a reduction in tongue pressure, delayed initiation of the pharyngeal swallow, impaired biomechanics, RLN palsy and increased pharyngeal residue, which may increase the patients' risk of aspirating, silently aspirating and developing pneumonia.^{26–28} To date there has been no systematic research using a prospective longitudinal study design in this patient group. By determining the prevalence, nature and trajectory of dysphagia post-oesophagectomy the researchers anticipate that this

would inform future research and guidelines in prevention and management of dysphagia, including exercise-based dysphagia interventions, which may optimise clinical and QOL outcomes.

Study objectives

The primary objectives of this study are:

1. To establish the prevalence, nature, severity and trajectory of dysphagia post-oesophagectomy among adults who have undergone a transthoracic (2-stage or 3-stage) or a transhiatal oesophagectomy (THO).
2. To determine the impact of oropharyngeal dysphagia on HR-QOL in this population across short and long-term time points.

The secondary objectives of this study are:

1. To determine the risk factors for post-operative dysphagia among adults post-oesophagectomy.
2. To identify the post-operative complications of dysphagia within this clinical population.

METHODS AND ANALYSIS

Study design

This is a proposed prospective longitudinal study which will be reported according to the Strengthening the Reporting of Observational Studies in Epidemiology checklist (see online supplemental appendix A).²⁹

Study setting

The study will take place in a National Oesophageal Centre (NOC) where patients with a diagnosis of oesophageal cancer who are scheduled for oesophagectomy will be identified from the upper gastrointestinal (UGI) clinic. Recruitment will be completed in this clinic over a 2-year period using consecutive sampling. Adults with a diagnosis of oesophageal cancer who are due to have curative oesophageal cancer surgery within the study setting will be invited to participate in this study by an independent gatekeeper. This prospective research study will assess patients across four time points: (1) A preoperative assessment of swallow will be recorded using the Functional Oral Intake Scale (FOIS) at time of consent for this study prior to their surgery,³⁰ (2) on day 4 or 5 post-oesophageal resection, (3) 1-month post-surgery and (4) 6-months post-surgery. The research team will evaluate swallowing-related outcome measures across these four time points. Risk factors and post-operative complications have been selected based on research in this clinical population to date. This will create a large resource of original data, which will inform further studies targeting prevention, early detection and intervention in dysphagia.

Patient and public involvement (PPI)

PPI in research has evolved over the past decade demonstrating a positive impact on health-related research.³¹ Early collaboration is known to enhance the quality and relevance of research when setting research priorities important to both the researcher and PPI, while also guiding further research.³² This prospective longitudinal




Figure 1 Study flowchart. CRT, cough reflex testing; EORTC-20, EORTC Quality of Life Questionnaire-18; FOIS, Functional Oral Intake Scale; IOPI, Iowa Oral Performance Instrument; MDADI, MD Anderson Dysphagia Inventory; MST, malnutrition screening tool; OHAT, oral health assessment tool; PIL, patient information leaflet; SARC-F, Strength, Assistance with Walking, Rise from a Chair, Climb Stairs, and Falls; SWAL-QOL, The Quality of Life in Swallowing Disorders; UGI, upper gastrointestinal; VFS, videofluoroscopy.

study has two PPI representatives (one male (SD); one female (BW)) involved on the research team, both of whom have undergone curative oesophageal cancer treatment. The PPI representatives are participating throughout the research study from the research design to dissemination. The PPI committee initially reviewed resource materials including the consent forms, patient information leaflet and rated patient-reported outcome measures (PROMs) for their relevance and ease of use. PPI involvement will be recorded using the Guidance for Reporting Involvement of Patients and the Public 2 form,³³ ensuring quality and consistency throughout the research. The PPI will be an integral part of the Knowledge Exchange and Dissemination scheme plan.³⁴

Study participants

Eligibility criteria are listed below.

Inclusion criteria

- ▶ A diagnosis of oesophageal cancer as confirmed by biopsy.
- ▶ Treated with curative intent involving surgery, which may be either open or minimally invasive.
- ▶ +/- neoadjuvant/adjuvant therapy.

- ▶ Scheduled for either a transthoracic (2-stage or 3-stage) or THO.
- ▶ Adults (>18 years).
- ▶ Ability to provide informed consent as per ethical approval obtained.

Exclusion criteria

- ▶ Known metastatic disease.
- ▶ Unable to complete VFS due to post-operative complications on POD 4 or 5.
- ▶ Patients who experience prolonged intubation beyond enhanced recovery after surgery (ERAS) protocol (>2 days).
- ▶ Patients who have a tracheostomy inserted due to failed extubation, secondary to prolonged intubation or reintubated post-operatively.
- ▶ Premorbid conditions potentially causing oropharyngeal dysphagia such as an acute or progressive neurological disease, history of head and neck cancer.
- ▶ 2-stage oesophagectomy with a confirmed anastomotic leak based on failed water-soluble swallow study at POD 5.

Patient recruitment

The proposed study will recruit 60 adults with oesophageal cancer undergoing oesophagectomy for curative intent. Participants will be recruited from the NOC, at St. James's Hospital (SJH). Details on recruitment and data collection can be viewed in [figure 1](#).

ERAS protocol

All participants will be treated according to standardised ERAS care pathway (see online supplemental appendix B),³⁵ involving either multimodal therapy (pre-operative chemotherapy alone or combined with radiation therapy), as per the MAGIC/FLOT or CROSS regimens, respectively, or surgery only. Surgical resection is typically performed at least 6 weeks post neoadjuvant therapy. Date of expected discharge from hospital is POD 9 as per the local oesophagectomy integrated care pathway.

Study protocol

Videofluoroscopy

The prevalence, nature and trajectory of oropharyngeal dysphagia post-oesophageal resection will be examined using VFS, a reference standard instrumental evaluation of oropharyngeal dysphagia and the evaluation of aspiration risk. Two VFS examinations will be completed on patients undergoing transthoracic (2-stage or 3-stage) and THO across two time points, immediately post-operatively (POD 4 or 5) and at 6-months post-oesophagectomy. Where a water-soluble contrast swallow study is required for inpatients post 2-stage oesophagectomy, the VFS will be completed immediately after this study, once the radiologist has ruled out an anastomotic leak. If an anastomotic leak is determined in this test, the participant will be withdrawn from the research study and the UGI clinical team informed.




Figure 2 Videofluoroscopy protocol. DIGEST, Dynamic Imaging Grade of Swallowing Toxicity; IDDSI, International Dysphagia Diet Standardisation Initiative.

The VFS will be completed by one researcher (MH) using Siemen Axiom Luminos TF fluoroscopy in the study setting: (1) post-resection on POD 4 or 5 and (2) at 6 months postoesophagectomy. The VFS pulse and frame rates will be 25 frames per second as per international recommendations.³⁶ Maxibar (98.45% w/w powder for oral suspension) is the contrast medium that will be used for VFS studies. This will be mixed with food and fluids to be radiopaque, assisting in determining anatomical and physiological deficits, rating the severity of oropharyngeal dysphagia and identifying aspiration risk during the study. The VFS will take approximately 20 minutes to complete. A standardised VFS protocol will be completed with participants in a seated position in lateral view followed by an anterior–posterior (AP) view as depicted in figure 2. Standardised bolus volumes and consistencies will be administered as per the International Dysphagia Diet Standardisation Initiative (IDDSI).³⁷

Tongue pressure evaluation

To investigate the nature of oropharyngeal dysphagia within this population, tongue pressure will be measured at three study time points. The Iowa Oral Performance Instrument (IOPI) is a handheld device frequently employed in clinical dysphagia research which will be used to measure tongue pressure.³⁸ This evaluation will be conducted at the bedside (POD 4 or 5) and in outpatient clinics (1-month and 6-month time points) before the VFS examination, where relevant. Participants will be instructed to insert an air-filled bulb into the oral cavity and to use their tongue to press this bulb against their hard palate. Measures of anterior peak tongue pressure (kPa) and tongue endurance (s) will be obtained.

Cough reflex testing

To evaluate laryngeal sensation, a dose–response method of cough reflex testing (CRT) will be measured across two of the study time points (prior to VFS POD 4 or 5 and at the 6-month clinic). CRT involves inhalation of a single concentration of a tussive agent (citric acid) via a face-mask nebuliser for a fixed period (within 15s of starting the nebuliser).³⁹ For best sensitivity and specificity to detect silent aspiration risk and impaired laryngeal sensation, a dosage of 0.4–0.8mol/L of citric acid in 0.9% saline solution is recommended.⁴⁰ In this study, various increments of citric acid in conjunction with a placebo 0.9 normal saline will be administered. A cough response will be considered positive if two (C_2) or more consecutive strong coughs are triggered within the time period where citric acid is induced. A weak cough will be determined as a cough that does not appear strong to clear material from the airway and is deemed substantially weaker than their own volitional cough.⁴¹ Patients who do not cough may indicate a greater silent aspiration risk and will be documented as a negative result. The findings of the test will be marked as a pass or fail result.

Aerodigestive symptoms

Based on feedback from the patient representatives, aerodigestive symptoms including cough and reflux are commonly experienced post-oesophagectomy and, given their strong association with swallowing, will be captured alongside swallow status in this study. The Reflux Symptom Index⁴² has been selected to address this and will be administered to participants across three time points (POD 4 or 5, 1-month and at 6-months) within the research study.

Patient-reported outcome measures

Based on feedback from PPI representatives, the PROMs selected for this study include the MDADI,⁴³ SWAL-QOL⁴⁴ and the EORTC-18.⁴⁵ The MDADI and the SWAL-QOL are validated dysphagia-specific QOL measure which is commonly used in dysphagia research. The EORTC-18 is another PROMs developed specifically for oesophageal cancer which is frequently used to evaluate HR-QOL in oesophageal cancer research⁴⁶ (see table 1).

Primary outcome measures

VFS analysis

The primary researcher, an experienced SLT, will complete the VFS analysis. Modified Barium Swallow Impairment Profile (MBS-Imp) ratings will be used to rate the presence, severity and trajectory of any swallow pathophysiology.⁴⁷ Fifteen randomly selected VFS studies (25%) will be rerated by blinded researchers to minimise bias (AG and JR).

The following validated VFS analysis measures will be obtained:

1. MBS-Imp ratings to identify the presence, severity and nature of any swallow pathophysiology.⁴⁷ The VFS images will be analysed frame by frame and graded using

Table 1 Swallowing, nutritional and QOL measurements across all time points

Instrument	Time point 1: baseline function and consent	Time point 2: POD 4/5	Time point 3: 1 month	Time point 4: 6 months
1. Swallow screening tool (TOR-BSST) ⁶⁰		x	x	x
2. Cough reflex testing (CRT) ³⁹⁻⁴¹		x	x	x
3. Tongue pressure measurement (IOP) ³⁸		x	x	x
4. Aerodigestive symptoms: Symptom Reflux Index ⁴²		x	x	x
5. FOIS, IDDSI, SARC-F, MST, weight and BMI ^{30 37 50 51}		x	x	x
6. QOL measures:		x	x	x
▶ MDADI ⁴³				
▶ SWAL-QOL ⁴⁴				
▶ EORTC-18 ⁴⁵				

EORTC, Quality of Life Questionnaire-18; FOIS, Functional Oral Intake Scale; IDDSI, International Dysphagia Diet Standardisation Initiative; IDDSI, International Dysphagia Diet Standardisation Initiative; IOP, Iowa Oral Performance Instrument; MDADI, MD Anderson Dysphagia Inventory; MST, malnutrition screening tool; POD, post-operative day; QOL, quality of life; SARC-F, Strength, Assistance with Walking, Rise from a Chair, Climb Stairs, and Falls; SWAL-QOL, Swallowing Quality of Life questionnaire.

the standardised MBS-Imp to identify the presence, severity and nature of swallow pathophysiology across 17 components. The components closely examine physiological components including the oral, pharyngeal and oesophageal phases of swallowing via lateral and AP radiological positioning during VFS. Please see [table 2](#).

2. Penetration-Aspiration Scale (PAS) ratings to measure swallow safety and cough response to aspiration across all swallows.⁴⁸ The validated PAS will be used to evaluate aspiration and cough response to penetration

and aspiration. This is an 8-point ordinal scale, which characterises the depth and response to airway penetration/aspiration during a VFS study.

3. Dynamic Imaging Grade of Swallowing Toxicity (DIGEST) score⁴⁹ will be used to stratify participants into dysphagia and non-dysphagia subgroups.

Secondary outcomes

Risk factors

The data on risk factors and post-operative complications will be obtained from participants' medical charts and from a local research database. Potential predictor variables will include: (1) age, (2) gender, (3) pre-surgical chemo/radiation, (4) tumour staging, (5) tumour type (squamous cell carcinoma/adenocarcinoma), (6) surgery type, (7) surgery duration (measured in hours), (8) RLN damage, (9) presence/degree of sarcopaenia using the Strength, Assistance With Walking, Rise From a Chair, Climb Stairs, and Falls,⁵⁰ (10) malnutrition (weight, body mass index (BMI), >10% wt loss and malnutrition screening tool,⁵¹ (11) oral health assessment tool and (12) number of co-morbidities.

Post-operative complications

Post-operative complications data will be collected from medical records and post-oesophagectomy database at outpatient appointments (1-month and 6-month clinic). Data will be obtained on (1) length of stay in the Intensive Care Unit (ICU) (days >3 days); (2) time to oral intake (days >5 POD); (3) tube feeding duration (days >30 days post discharge); (4) presence of pneumonia as per American Thoracic Society (ATS) post-operative pneumonia score⁵² (as per local UGI database) (yes/no); (5) oesophageal strictures±dilatation/stenting; (6) mortality/survival rates; and (7) other complications as per the Esophageal Complications Consensus Group definitions.¹⁰

Table 2 Modified Barium Swallow Impairment Profile analysis components⁴⁷

Number	Physiological component
1	Lip closure
2	Tongue control during bolus hold
3	Bolus preparation/mastication
4	Bolus transport/lingual motion
5	Oral residue
6	Initiation of pharyngeal swallow
7	Soft palate elevation
8	Laryngeal elevation
9	Anterior hyoid excursion
10	Epiglottic movement
11	Laryngeal vestibular closure
12	Pharyngeal stripping wave
13	Pharyngeal contraction (AP view)
14	Pharyngo-oesophageal segment opening
15	Tongue base retraction
16	Pharyngeal residue
17	Oesophageal clearance (AP view)

AP, anterior-posterior.



Pneumonia

PPCs, primarily pneumonia, is a common post-operative complication, which may be infection associated or complicated by respiratory failure or acute respiratory distress syndrome (ARDS).⁵³ The risk is greater in patients with existing chronic obstructive pulmonary disease or in current smokers.⁵³ Other risk factors include age, gender, total number of lymph nodes resected and operation approach (transthoracic extended).⁵⁴ The ATS post-operative pneumonia score will be used to determine pneumonia in post-oesophagectomy patients. The ATS define hospital-acquired pneumonia as a pneumonia not incubating at the time of hospital admission, occurring >48 hours or more after admission whereas ventilated-acquired pneumonia is determined >48 hours post endotracheal intubation.⁵⁵ Pneumonia is suspected if the patient has radiographic infiltrates that is new or progressive in association with the following clinical findings suggestive of a pneumonia include: (1) new onset of fever, (2) purulent sputum, (3) leukocytosis and (4) a decline in oxygenation.⁵⁵ As pneumonia is the most prevalent complication post-oesophageal resection, research supports assessing patients for any swallowing dysfunction or predisposition to aspiration prior to commencing oral intake to reduce risk of post-operative complications and mortality.⁵⁶ As the prevalence, nature and trajectory of oropharyngeal dysphagia has not been determined in a prospective longitudinal study, its link and impact on pneumonia rates postoesophageal cancer surgery are relatively unknown.

Study size

This is an exploratory longitudinal study in an area with limited previous research or group comparisons and no reporting of effect size. Based on previous literature in this cancer cohort to estimate an effect size 0.5 at a significance level of 0.05 and a power of 0.8, a sample size of 60 is calculated for repeated measures. This sample estimate is consistent with other publications in this area.^{57 58}

Data analysis

SPSS V.22.0 will be used for statistical analyses.⁵⁹ Variables will be tested for normality using the Shapiro-Wilks test. Normally distributed variables will be summarised as mean and SD. Non-normally distributed data will be summarised as median and IQR. Categorical variables are presented as frequency (percentage). To establish changes in participant swallow outcomes across time points, repeated measures will be performed using repeated measures analysis of variance or Friedman tests.

To identify independent risk factors, multiple logistic regression will be performed. To identify complications of dysphagia, mean/median (depending on distribution of data) differences in length of hospital stay, pneumonia, sarcopaenia, tube-feeding reliance, mortality and QOL will be compared across dysphagia and non-dysphagia subgroups. The VFS protocol outlined includes a robust system of validated measures to detect oropharyngeal

dysphagia in this patient group. A strict data management plan includes data being stored securely, anonymously and processed in adherence to the general data protection regulator best practice guidelines in line with ethical approval.

Ethics and dissemination

Ethical approval has been obtained from the SJH-Tallaght University Hospital (TUH) Joint Research Ethics Committee (J-REC) (2021-Jul-310), alongside the SJH Research and Innovation (R&I) committee. The patient will be formally enrolled into the research study if meets the research criteria and informed consent has been obtained. The primary researcher (MH) involved will eliminate any potential risks to the participant. During the procedure, the patient may be at risk of aspiration if oropharyngeal dysphagia post-oesophagectomy is present. The researcher will inform the patient, refer to inpatient Speech and Language Therapy and Physiotherapy team and notify the UGI Surgeons. The Radiology department where the VFS will take place is located within SJH and is covered by the hospital response team. All adverse events will be documented, and any serious adverse incidences will be immediately reported to the patients' surgical team and to the research ethics committee.

Findings of the prospective longitudinal study will be disseminated via conference presentations including the World Dysphagia Summit, Dysphagia Research Society, The European Society of Swallowing Disorders and the International Society of Diseases of the Esophagus conference. The findings will be published in peer-reviewed academic journals. Study participants will be informed of study results.

DISCUSSION

Data collection and analysis will be completed at a NOC, where approximately 55–60 curative oesophageal resections are completed annually. This research study has not received any specific grants from funding agencies, and no known competing financial interests or personal conflict that could appear to influence the nature of this research has been declared.

As survival rates are improving among adults with oesophageal cancer, there has been a shift in research and clinical focus to optimise HR-QOL among survivors. Dysphagia is strongly associated with HR-QOL in this population, but relatively understudied in terms of prevalence and modifiable intervention target, and the studies proposed will provide comprehensive data on this cohort and inform further research and clinical advances in this context. Physiological changes impacting the oropharyngeal swallow across four separate time points will be determined using rigorous reference standard and validated swallowing assessment tools. A robust study design will be implemented, using a broad range of clinical swallowing outcomes. This will be examining the prevalence, nature

and trajectory of oropharyngeal dysphagia following oesophagectomy.

PPI will be a key strength in this research study. Patients' previous experience of the oesophageal cancer journey will provide invaluable insight and guidance across different time points in the study. Furthermore, the collaboration between the researcher and committee members will strengthen research priorities set out and aim to meet at different intervals throughout the research cycle within this study.

This study has some limitations that we acknowledge. Firstly, the risk of post-operative complications including ARDS, pneumothorax, risk of re-intubation, delirium, anastomotic leak who require medical interventions, will prevent recruitment into this study. Failure to collect data on patients with complex post-operative needs who may potentially present with an oropharyngeal dysphagia is recognised as a limitation. Patient retention may be challenging due to the increased risk of cancer recurrence in this population, ultimately impacting their ability to participate during the different time points. For this reason, it was decided to recruit patients up to 6-months post-resection rather than 1-year following oesophagectomy. The author acknowledges that the 6-month timeframe may not fully capture swallowing impairment and QOL measures following surgery, however this research group is also conducting another major study, examining the prevalence, nature and impact of dysphagia 1-year post-oesophagectomy and into survivorship.

This longitudinal study will create a large database encompassing detailed information about the presence, nature and trajectory of dysphagia in the post-oesophagectomy setting, its link to other complications and its impact on recovery of QOL. The database will inform the development of intervention programmes tailored to the unique needs of people with oesophageal cancer. The results will provide a large resource of original data and inform further studies targeting prevention and early intervention. Furthermore, the findings may target development of swallowing compensatory strategies and rehabilitation therapy to optimise swallow function and safety. The results may further inform current clinical practice and provide direction for future research.

Author affiliations

¹Department of Clinical Speech and Language Studies, Trinity College Dublin, Dublin, Ireland

²Senior Upper GI and ICU Speech and Language Therapist, St. James's Hospital, Dublin, Ireland

³Patient and Public Representative Group, Department of Clinical Speech and Language Studies, Trinity College Dublin, Dublin, Ireland

⁴Department of Radiology, St. James's Hospital, Dublin, Ireland

⁵Consultant Gastrointestinal Surgeon, Department of Surgery, St. James's Hospital, Dublin, Ireland

⁶Department of Surgery, Trinity College Dublin, Dublin, Ireland

Twitter Michelle Hayes @MichelleHayeSLT

Contributors MH and JR designed the study. MH wrote the protocol. AG, BW, SD, IB, MW, CD, JVR and JR reviewed the protocol paper.

Funding The authors have declared that they are not receiving any specific funding for this research.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Michelle Hayes <http://orcid.org/0000-0002-7996-8561>

Julie Regan <http://orcid.org/0000-0001-5816-4516>

REFERENCES

- Smyth EC, Lagergren J, Fitzgerald RC, *et al*. Oesophageal cancer. *Nat Rev Dis Primers* 2017;3:1–21.
- Triantafyllou T, Wijnhoven BPL. Current status of esophageal cancer treatment. *Chin J Cancer Res* 2020;32:271–86.
- Ashok A, Niyogi D, Ranganathan P, *et al*. The enhanced recovery after surgery (ERAS) protocol to promote recovery following esophageal cancer resection. *Surg Today* 2020;50:323–34.
- Scheepers JJG, van der Peet DL, Veenhof AAFA, *et al*. Thoracoscopic resection for esophageal cancer: a review of literature. *J Minim Access Surg* 2007;3:149.
- Gujjuri RR, Kamarajah SK, Markar SR. Effect of anastomotic leaks on long-term survival after oesophagectomy for oesophageal cancer: systematic review and meta-analysis. *Dis Esophagus* 2021;34:doaa085.
- Huang F-L, Yu S-J. Esophageal cancer: risk factors, genetic association, and treatment. *Asian J Surg* 2018;41:210–5.
- Ahmadinejad M, Soltanian A, Maghsoudi LH. Risk factors and therapeutic measures for postoperative complications associated with esophagectomy. *Ann Med Surg* 2020;55:167–73.
- Donlon NE, Ravi N, King S, *et al*. Modern oncological and operative outcomes in esophageal cancer: the St. James's Hospital experience. *Ir J Med Sci* 2021;190:297–305.
- Sheill G, Guinan E, O'Neill L, O'Neill L, *et al*. Preoperative exercise to improve fitness in patients undergoing complex surgery for cancer of the lung or oesophagus (PRE-HIIT): protocol for a randomized controlled trial. *BMC Cancer* 2020;20:1–11.
- Reynolds JV, Donlon N, Elliott JA, *et al*. Comparison of esophagectomy outcomes between a national center, a national audit collaborative, and an international database using the esophageal complications consensus group (ECCG) standardized definitions. *Dis Esophagus* 2021;34:doaa060.
- Elliott JA, Docherty NG, Eckhardt H-G, *et al*. Weight loss, satiety, and the postprandial gut hormone response after esophagectomy: a prospective study. *Ann Surg* 2017;266:82–90.
- Bolger JC, Loughney L, Tully R, *et al*. Perioperative prehabilitation and rehabilitation in esophagogastric malignancies: a systematic review. *Dis Esophagus* 2019;32:doz058.
- Reynolds JV, Donohoe CL, McGillycuddy E, *et al*. Evolving progress in oncologic and operative outcomes for esophageal and junctional cancer: lessons from the experience of a high-volume center. *J Thorac Cardiovasc Surg* 2012;143:1130–7.
- Arnold M, Rutherford MJ, Bardot A, *et al*. Progress in cancer survival, mortality, and incidence in seven high-income countries 1995–2014

- (ICBP SURVMARK-2): a population-based study. *Lancet Oncol* 2019;20:1493–505.
- 15 Schandl A, Johar A, Anandavadevelan P, *et al.* Patient-reported outcomes 1 year after oesophageal cancer surgery. *Acta Oncol* 2020;59:613–9.
 - 16 Donohoe CL, McGillicuddy E, Reynolds JV. Long-term health-related quality of life for disease-free esophageal cancer patients. *World J Surg* 2011;35:1853–60.
 - 17 Thrumurthy SG, Chaudry MA, Thrumurthy SSD, *et al.* Oesophageal cancer: risks, prevention, and diagnosis. *BMJ* 2019;366:l4373.
 - 18 Kato H, Miyazaki T, Sakai M, *et al.* Videofluoroscopic evaluation in oropharyngeal swallowing after radical esophagectomy with lymphadenectomy for esophageal cancer. *Anticancer Res* 2007;27:4249–54.
 - 19 Scholtemeijer MG, Seesing MFJ, Brenkman HJF, *et al.* Recurrent laryngeal nerve injury after esophagectomy for esophageal cancer: incidence, management, and impact on short- and long-term outcomes. *J Thorac Dis* 2017;9:S868–78.
 - 20 Ryan AM, Rowley SP, Healy LA, *et al.* Post-oesophagectomy early enteral nutrition via a needle catheter jejunostomy: 8-year experience at a specialist unit. *Clin Nutr* 2006;25:386–93.
 - 21 Dai Y, Li C, Xie Y. Interventions or dysphagia in oesophageal cancer. *Cochrane Database Syst Rev* 2014;2014:CD005048.
 - 22 Takatsu J, Higaki E, Hosoi T, *et al.* Clinical benefits of a swallowing intervention for esophageal cancer patients after esophagectomy. *Dis Esophagus* 2021;34:doaa094.
 - 23 Liu YJ, Schandl A, Markar S, *et al.* Psychological distress and health-related quality of life up to 2 years after oesophageal cancer surgery: nationwide population-based study. *BJS Open* 2021;5:zraa038.
 - 24 Markar SR, Sounderajah V, Johar A, *et al.* Patient-reported outcomes after oesophagectomy in the multicentre laser study. *Br J Surg* 2021;108:1090–6.
 - 25 Klevebro F, Johar A, Lagergren P. Impact of co-morbidities on health-related quality of life 10 years after surgical treatment of oesophageal cancer. *BJS Open* 2020;4:601–4.
 - 26 Yokoi A, Ekuni D, Yamanaka R, *et al.* Change in tongue pressure and the related factors after esophagectomy: a short-term, longitudinal study. *Esophagus* 2019;16:300–8.
 - 27 Tsubosa Y, Sato H, Nemoto M, *et al.* Experience of rehabilitation for swallowing disorders after esophagectomy for esophageal cancer. *Jpn J Gastroenterol Surg, Nihon Shokaki Geka Gakkai zasshi* 2005;38:571–6.
 - 28 Yuen MTY, Tsang RK, Wong IYH, *et al.* Long-term pharyngeal dysphagia after esophagectomy for esophageal cancer—an investigation using videofluoroscopic swallow studies. *Diseases of the Esophagus* 2019;32:doj068.
 - 29 von Elm E, Altman DG, Egger M, *et al.* The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Int J Surg* 2014;12:1495–9.
 - 30 Crary MA, Mann GDC, Groher ME. Initial psychometric assessment of a functional oral intake scale for dysphagia in stroke patients. *Arch Phys Med Rehabil* 2005;86:1516–20.
 - 31 Involve L. *Public involvement in research getting it right and making a difference: a unique forum for people who are interested in active public involvement in research*, 2008.
 - 32 Saini P, Hassan SM, Morasae EK, *et al.* The value of involving patients and public in health services research and evaluation: a qualitative study. *Res Involv Engagem* 2021;7:1–16.
 - 33 Staniszewska S, Brett J, Simer I, *et al.* GRIPP2 reporting checklists: tools to improve reporting of patient and public involvement in research. *BMJ* 2017;358:j3453.
 - 34 Health Research Board. Available: <https://www.hrb.ie/funding/funding-schemes/all-funding-schemes>
 - 35 Low DE, Allum W, De Manzoni G, *et al.* Guidelines for perioperative care in esophagectomy: enhanced recovery after surgery (ERAS[®]) society recommendations. *World J Surg* 2019;43:299–330.
 - 36 Martin-Harris B, Canon CL, Bonilha HS, *et al.* Best practices in modified barium swallow studies. *Am J Speech Lang Pathol* 2020;29:1078–93.
 - 37 Cichero JAY, Lam P, Steele CM, *et al.* Development of international terminology and definitions for texture-modified foods and thickened fluids used in dysphagia management: the IDDSI framework. *Dysphagia* 2017;32:293–314.
 - 38 Adams V, Mathisen B, Baines S, *et al.* A systematic review and meta-analysis of measurements of tongue and hand strength and endurance using the lowa oral performance instrument (IOP). *Dysphagia* 2013;28:350–69.
 - 39 Wallace E, Guiu Hernandez E, Ang A, *et al.* A systematic review of methods of citric acid cough reflex testing. *Pulm Pharmacol Ther* 2019;58:101827.
 - 40 Monroe MD, Manco K, Bennett R, *et al.* Citric acid cough reflex test: establishing normative data. *Speech, Language and Hearing* 2014;17:216–24.
 - 41 Morice AH, Fontana GA, Belvisi MG, *et al.* ERS guidelines on the assessment of cough. *Eur Respir J* 2007;29:1256–76.
 - 42 Belafsky PC, Postma GN, Koufman JA. Validity and reliability of the reflux symptom index (RSI). *J Voice* 2002;16:274–7.
 - 43 Chen AY, Frankowski R, Bishop-Leone J, *et al.* The development and validation of a dysphagia-specific quality-of-life questionnaire for patients with head and neck cancer: the M. D. Anderson dysphagia inventory. *Arch Otolaryngol Head Neck Surg* 2001;127:870–6.
 - 44 McHorney CA, Robbins J, Lomax K, *et al.* The SWAL-QOL and SWAL-CARE outcomes tool for oropharyngeal dysphagia in adults: III. documentation of reliability and validity. *Dysphagia* 2002;17:97–114.
 - 45 McKernan M. *The relationship between quality of life (EORTC QLQ C-20) and survival and treatment in patients with gastro-oesophageal cancer*, 2008.
 - 46 Churrua K, Pomare C, Ellis LA, *et al.* Patient-reported outcome measures (PROMs): a review of generic and condition-specific measures and a discussion of trends and issues. *Health Expectations* 2021;24:1015–24.
 - 47 Martin-Harris B, Humphries K, (Focht) Garand KL. The modified barium swallow impairment profile (MBSImP[™]) – innovation, dissemination and implementation. *Perspect ASHA Spec Interest Groups* 2017;2:129–38.
 - 48 Rosenbek JC, Robbins JA, Roecker EB, *et al.* A penetration-aspiration scale. *Dysphagia* 1996;11:93–8.
 - 49 Hutcheson KA, Barrow MP, Barringer DA, *et al.* Dynamic imaging grade of swallowing toxicity (digest): scale development and validation. *Cancer* 2017;123:62–70.
 - 50 Borges TC, Gomes TLN, Pimentel GD. Sarcopenia as a predictor of nutritional status and comorbidities in hospitalized patients with cancer: a cross-sectional study. *Nutrition* 2020;73:110703.
 - 51 Deftereos I, Djordjevic A, Carter VM, *et al.* Malnutrition screening tools in gastrointestinal cancer: a systematic review of concurrent validity. *Surg Oncol* 2021;38:101627.
 - 52 van der Sluis PC, Verhage RJJ, van der Horst S, *et al.* A new clinical scoring system to define pneumonia following esophagectomy for cancer. *Dig Surg* 2014;31:108–16.
 - 53 Rotstein C, Evans G, Born A, *et al.* Clinical practice guidelines for hospital-acquired pneumonia and ventilator-associated pneumonia in adults. *Can J Infect Dis Med Microbiol* 2008;19:19–53.
 - 54 Linden PA, Towe CW, Watson TJ, *et al.* Mortality after esophagectomy: analysis of individual complications and their association with mortality. *J Gastrointest Surg* 2020;24:1–7.
 - 55 Dela Cruz CS, Evans SE, Restrepo MI. Understanding the host in the management of pneumonia. *Ann Am Thorac Soc* 2021;18:1087–97.
 - 56 Atkins BZ, Shah AS, Hutcheson KA, *et al.* Reducing hospital morbidity and mortality following esophagectomy. *Ann Thorac Surg* 2004;78:56:1170–6.
 - 57 Citak E, Tulek Z, Uzel O. Nutritional status in patients with head and neck cancer undergoing radiotherapy: a longitudinal study. *Support Care Cancer* 2019;27:239–47.
 - 58 Guinan EM, Bennett AE, Doyle SL, *et al.* Measuring the impact of oesophagectomy on physical functioning and physical activity participation: a prospective study. *BMC Cancer* 2019;19:1–11.
 - 59 Leech NL, Barrett KC, Morgan GA. *IBM SPSS for intermediate statistics: use and interpretation*. Routledge, 2014.
 - 60 Martino R, Silver F, Teasell R, *et al.* The Toronto bedside swallowing screening test (TOR-BSST): development and validation of a dysphagia screening tool for patients with stroke. *Stroke* 2009;40:555–61.