

# BMJ Open Developing a core outcome set for physical activity interventions in primary schools: a modified-Delphi study

Bina Ram <sup>1</sup>, Kimberley A Foley <sup>1</sup>, Esther van Sluijs,<sup>2</sup> Dougal S Hargreaves,<sup>1,3</sup> Russell M Viner <sup>4</sup>, Sonia Saxena <sup>1</sup>

**To cite:** Ram B, Foley KA, van Sluijs E, *et al.* Developing a core outcome set for physical activity interventions in primary schools: a modified-Delphi study. *BMJ Open* 2022;**12**:e061335. doi:10.1136/bmjopen-2022-061335

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-061335>).

Received 24 January 2022  
Accepted 26 August 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Primary Care and Public Health, Imperial College London, London, UK

<sup>2</sup>MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge, UK

<sup>3</sup>Mohn Centre for Children's Health and Wellbeing, Imperial College London, London, UK

<sup>4</sup>Population, Policy and Practice Research Programme, UCL Institute of Child Health, London, UK

**Correspondence to**  
Dr Bina Ram;  
[b.ram@imperial.ac.uk](mailto:b.ram@imperial.ac.uk)

## ABSTRACT

**Objectives** To develop a core outcome set (COS) for physical activity interventions in primary schools.

**Design** Modified-Delphi study.

**Setting** The UK and international.

**Participants** 104 participants from four stakeholder groups (educators, public health professionals, health researchers, parents); 16 children (aged 8–9 years) from 1 London primary school.

**Interventions** Physical activity interventions.

**Methods** Four-stage process: (1) outcomes extracted from relevant studies identified from an umbrella review and a focus group; (2) list of outcomes produced and domains established; (3) stakeholders completed a two-round Delphi survey by rating (Round 1) and re-rating (Round 2) each outcome on a nine-point Likert Scale from 'not important' to 'critical': a >70% participant threshold identified the outcomes rated 'critical' to measure, and outcomes important to children were identified through a workshop; and (4) a stakeholder meeting to achieve consensus of the outcomes to include in the COS.

**Results** In total, 74 studies were extracted from 53 reviews. A list of 50 outcomes was produced and three domains were established: 'physical activity and health' (16 outcomes), 'social and emotional health' (22 outcomes) and 'educational performance' (12 outcomes). 104 participants completed survey Round 1; 65 participants completed both rounds. In total, 13 outcomes met the threshold; children identified 8 outcomes. Fourteen outcomes achieved consensus to produce the COS: five outcomes for physical activity and health (diet (varied and balanced), energy, fitness, intensity of physical activity, sleep (number of hours)); seven outcomes for social and emotional health (anxiety, depression, enjoyment, happiness, self-esteem, stress, well-being); and two outcomes for educational performance (concentration, focus).

**Conclusions** We have developed the first COS for physical activity interventions in primary schools in consultation with those interested in the development and application of an agreed standardised set of outcomes. Future studies including these outcomes will reduce heterogeneity across studies.

**Trial registration number** Core Outcome Measures in Effectiveness Trials Initiative registration number 1322; Results.


## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This is the first study to develop a core outcome set (COS) for physical activity interventions in primary schools.
- ⇒ The COS has been developed in consultation with participants from key stakeholder groups.
- ⇒ This study uses robust methodology as recommended by the Core Outcome Measures in Effectiveness in Trials Initiative.
- ⇒ There were an unbalanced number of participants in each stakeholder group.
- ⇒ The low representation of international participants may limit the use of this COS to UK schools only.

## INTRODUCTION

Increasing children's physical activity is a global health goal given the vast evidence showing benefits on physical, social, mental and cognitive health outcomes.<sup>1</sup> Health behaviours may become embedded in childhood; providing opportunities for children to engage in physical activities during the primary school years may lead to physically active lifestyles and improved health during adolescence and adulthood.<sup>2</sup> Many governments support the need for increased physical activity promotion in schools.<sup>3</sup> The WHO recommends that schools should organise and promote opportunities for children to regularly participate in physical activities.<sup>4</sup>

School settings are ideal as they have the potential to reach the majority of children across society<sup>5 6</sup> including those living in poverty. Socioeconomic inequalities have been associated with moderate and vigorous physical activities and may contribute to widening health inequalities.<sup>7</sup> Targeting schools therefore could help towards reducing the gap in physical activity among children.<sup>7 8</sup> As a result of governments and the WHO recommendations of physical activity promotion and engagement in schools, there are many physical activity interventions that are implemented. However, the interventions



**Figure 1** Process for developing a core outcome set for physical activity interventions in primary schools.

vary in design. Some interventions integrate additional physical education classes alongside compulsory physical education lessons,<sup>9</sup> while some may incorporate 10 min of physical activity into every school day.<sup>10</sup> There are also others which implement classroom movement breaks<sup>11</sup> or active mile interventions.<sup>12 13</sup>

There is considerable evidence showing the benefits of physical activity interventions in schools successfully increasing children's fitness<sup>14–17</sup> and reducing sedentary time<sup>18 19</sup>. There is also increasing evidence of improvements to children's social, emotional and cognitive outcomes.<sup>20–23</sup> However, due to the heterogeneity of the outcomes assessed across studies, definitive conclusions are challenging.<sup>20 22</sup> For example, to assess children's emotional health, one study may measure children's 'happiness', while another may measure 'depression'. Both these outcomes are conceptually different and difficult to compare. In 2013, a Cochrane review of 44 randomised control trials of physical activity interventions in schools for children aged 6–18 years found considerable variations in the outcomes measured, and the results could not be synthesised to establish intervention effects.<sup>24</sup> The review was updated in 2021; the authors concluded that due to the variability of results, heterogeneity and risk of bias across studies, the impacts of physical activity interventions in schools have shown small effects. These interventions may show small improvements to children's physical fitness but have little or no impact on other outcomes such as body mass index (BMI).<sup>25</sup>

Synthesising results from studies are likely to be of interest to a number of key groups including public health professionals, teachers, parents, healthcare researchers and policymakers. However, many of the outcomes measured in existing studies, although important to

measure, may vary in relevance to specific groups. For example, BMI is a frequently measured outcome from which important conclusions have been identified.<sup>26 27</sup> BMI may be considered highly important to healthcare practitioners but may not be considered as important to teachers who may instead place higher importance on cognitive outcomes. Lack of consultation with key groups when deciding which outcomes to measure in studies limits the relevance of findings to specific groups and may have possibly led to differences of outcomes measured across studies, thus preventing comparisons.

A core outcome set (COS) is an agreed set of standardised outcomes in a specific research area that is recommended to measure and report.<sup>28</sup> These sets should be developed in consultation with those who are interested in the development and application of an agreed set of outcomes.<sup>29</sup> The COS should be viewed as a minimum to measure and does not restrict additional outcomes of interest to be assessed. COSs were originally developed for clinical trials but are increasingly being used in other study designs, for example, in observational studies by practitioners and researchers to conduct their own assessments of interventions.<sup>28</sup> To our knowledge, there is not a COS for physical activity interventions in primary schools. Therefore, the development of a COS (the aim of this study) would contribute to this field of research by identifying the key outcomes to be studied, allowing for evidence synthesis to better understand the impact of physical activity interventions in schools on children's health.

## METHODS

### Design

The protocol for this work has been published in online supplemental file 1<sup>30</sup>; it was developed in accordance with the Core Outcome Measures in Effectiveness Trials (COMET) criteria<sup>29</sup> and prospectively registered accordingly.<sup>31</sup> We used a modified-Delphi method consisting of four stages to develop the COS (figure 1). First, we extracted outcomes and how they had been defined/described by the authors of relevant studies identified through an umbrella review and through a focus group with our steering committee (our steering committee includes health professionals, health researchers, academics and sports representatives from organisations such as Sport England and The Daily Mile Foundation). Second, after deduplication and combining similar outcomes, we created a long list and established domains determined by the outcomes. Third, we recruited participants from four key stakeholder groups (educators, health researchers, public health professionals and parents of children aged from 5 to 11 years) to complete a two-round Delphi survey. We also obtained children's views of what is important to them through a workshop. Fourth, we held a stakeholder meeting to achieve consensus on the outcomes to be included in the COS. We report the

study following the COS-STAndards for Reporting check-list (online supplemental file 2).<sup>32</sup>

### Stage 1: extraction of outcomes

For the umbrella review, we searched six databases (MEDLINE, EMBASE, PsycINFO, CINAHL, CENTRAL and the Cochrane Database of Systematic Reviews). Keywords used for the search were 'school', 'physical activity', 'exercise', 'physical education', 'fitness' and 'energy expenditure' and adapted to use database specific filters, that is, subject headings or medical subject headings. Reviews were limited to systematic reviews, meta-analyses or meta-syntheses and those published between 1990 and 2019. Relevant studies from these reviews were identified from which the outcomes extracted. We also held a focus group with our steering committee and used a nominal group technique to brainstorm outcomes and rate their importance to extract further outcomes that may not have been captured in our literature review. Descriptions of each outcome were guided by the published literature and discussions with our steering group.

### Stage 2: list of outcomes and establishing domains

We removed duplicate outcomes and merged those that were closely related, for example, outcomes of 'light physical activity', 'moderate physical activity' and 'vigorous physical activity' were combined into 'intensity of physical activity', to create a long list of outcomes. Descriptions were generated for each outcome based on those provided by authors of the relevant studies and discussions with our steering committee. Guided by the outcomes and descriptions, we established relevant domains by grouping similar outcomes that captured a broader concept.

### Stage 3: stakeholder recruitment, Delphi surveys and children's workshop

The purpose of the Delphi surveys was to identify which outcomes, from the long list we produced, were considered the most important to measure across key stakeholder groups.

#### Stakeholder recruitment

Through emails to our public health research and practitioner networks and through snowballing and social media, we recruited participants from four key stakeholder groups (educators (teachers, head teachers, school governors), health researchers, public health professionals and parents of primary school-aged children). Through discussions with our steering group, we identified the key stakeholder groups that would be the most interested in the development and implementation of an agreed set of outcomes to enhance this field of research. An information leaflet was made available to participants which included an electronic link to the Round 1 Delphi survey and study contact details. Through the Round 1 survey link, we obtained consent for participation, followed by participants registering their details (name and email address) and indicating which of the four stakeholder groups they identified with.

#### Delphi surveys

Using DelphiManager software,<sup>33</sup> we listed the outcomes with their descriptions by each domain in a Delphi survey conducted over two rounds (Round 1 took place during June 2020 and Round 2 in August 2020). Using the predefined Delphi survey guidelines,<sup>33</sup> we asked participants to rate the importance of each outcome using a nine-point Likert Scale ranging from 'not important to measure' to 'critical to measure' in Round 1. A rating of 10 could be indicated if participants felt they were unable to score an outcome. Ratings were grouped into three categories: 'not important to measure' (ratings of 1, 2 or 3); 'important but not critical to measure' (ratings of 4, 5 or 6); and 'critical to measure' (ratings of 7, 8 or 9). In addition, participants were asked to suggest any other outcomes that they felt were not captured. In line with our protocol, if more than two individual participants suggested the same additional outcome, this would be included in Round 2 for all participants to rate. For ratings in Round 2, participants were provided with feedback of Round 1 ratings categorised by stakeholder group and an option to rerate their initial ratings based on this feedback. Participants were sent three email reminders to complete Round 1; those who rated all outcomes in Round 1 were invited to complete Round 2. The criteria for outcomes considered most important to measure for each domain after Round 2 were defined a priori,  $\geq 70\%$  of all participants rating an outcome 'critical' and 15% or less rating it 'not important'.<sup>30</sup> None of the outcomes were removed between rounds.

#### Children's workshop

We recruited primary school children to take part in a workshop in December 2020 with consent obtained from parents via the school. Due to COVID-19, our access to schools was restricted. We partnered with one primary school in Greater London, UK. Guided by the list of outcomes, we engaged the children in a series of activities and discussions on physical activity and elicited the children's views on what they thought was important to measure.

#### Stage 4: stakeholder meeting

Participants who completed both survey rounds were invited to attend the stakeholder meeting in December 2020. Due to COVID-19 restrictions, the meeting was held virtually using the Zoom platform and we adapted the voting method (70%/15% threshold) as described in our protocol. Instead, to achieve consensus on the outcomes to be included in the COS, we led discussions around the ratings of outcomes in the Delphi surveys and children's views. We used the Zoom chat function for participants to indicate the most important outcomes and further discussion to agree the outcomes to be included in the COS.

#### Patient and public involvement

We have consulted with professional and public representatives within our steering committee and as part of



The Daily Mile Research Advisory Group. Both groups include public health professionals, health researchers, academic researchers and representatives from The Daily Mile Foundation, Sport England, London Marathon and London Sport. Our COS has been developed in consultation with educators, health researchers, public health professionals, parents and children through focus groups and workshops. We will widely advertise our COS through those involved in the development and also to child public health policymakers through our research networks.

## RESULTS

### Stage 1: extraction of outcomes

Our umbrella review identified 53 relevant papers from which 74 individual studies were extracted (online supplemental file 3); around 181 outcomes were identified from these studies. However, we identified variations across studies of how the outcomes were defined or described if at all. The steering committee focus group identified 34 outcomes. We created the description for each outcome guided by the literature and from discussions with our Steering Group.

### Stage 2: list of outcomes and establishing domains

The final list consisted of 50 outcomes (table 1) representing three domains: (1) physical activity and health (16 outcomes); (2) social and emotional health (22 outcomes); and (3) educational performance (12 outcomes). Two outcomes, 'sleep' and 'diet', were included in two domains as authors agreed that these outcomes in particular could be both a 'physical activity and health' and a 'social and emotional health' outcome. For example, sleep defined as number of hours slept as recommended for children was included in the physical activity and health domain, while sleep times/patterns/broken sleep was included in the social and emotional health domain. Similarly for the outcome of diet, eating well-balanced meals was included in the physical activity and health domain, while appetite was included in the social and emotional health domain (see table 1 for descriptions).

### Stage 3: stakeholder recruitment, Delphi surveys and children's workshop

#### Stakeholder recruitment

A total of 104 participants consented and registered their details. Ninety (87%) completed Round 1 in full of whom 65 (72%) also completed Round 2 in full. The 65 participants included 16 (25%) educators, 24 (37%) researchers, 13 (20%) public health professionals and 12 (18%) parents and represented 9 countries: the UK (80%), Brazil (6%) and Korea (5%), Australia, France, the Netherlands, Romania, Spain and Taiwan (2%).

#### Delphi surveys

In total, 13 outcomes met the >70% participant critical threshold: sleep (number of hours) and diet (varied

and balanced) in 'physical activity and health'; happiness, well-being, anxiety, self-esteem, depression, self-confidence, enjoyment and stress in 'social and emotional health'; and concentration, attention and focus in 'educational performance' (table 2). In Round 1, a further 29 outcomes were suggested, but after internal discussions, it was agreed that 16 of the suggestions overlapped with the outcomes that were listed in the survey, and the remaining 13 were proposed by only one participant and therefore not carried forward to Round 2. Mean Round 1 ratings of participants completing both Rounds were similar to those who completed Round 1 but did not complete Round 2 (6.33, SD 2.08 vs 6.48, SD 1.95, respectively) suggesting those who did not complete Round 2 would have scored similarly to those who did.

#### Children's workshop

In total, 16 children aged 8–9 years took part in the workshop, of which 50% were girls; 13% were Caucasian, 56% were Asian and 31% were black; 6% had special educational needs; and 75% had English as a second language. The children identified eight outcomes important to measure: five in 'physical activity and health' (energy, fitness, heart rate, muscle strength and weight) and three in 'social and emotional health' (happiness, mood and stress). Interestingly, children did not associate physical activity with any educational performance related outcomes.

#### Stage 4: stakeholder meeting

In total, 13 participants attended (2 educators, 2 parents and 9 researchers). Participants expressed that they had expected more outcomes under the domain of physical activity and health to be rated critical, that is, intensity of physical activity which had been rated critical by 63% (table 2). Through discussion, agreement was reached that this outcome is important to measure to be able to assess sustainability of physical activity interventions in schools. After review of the outcomes identified critical in the survey and the outcomes considered important to children, six outcomes were dropped and the additional outcome of intensity of physical activity was included (online supplemental file 4). Therefore, a total of 14 outcomes reached consensus for the COS: diet (varied and balanced), fitness, intensity of physical activity and sleep (number of hours) in the physical activity and health domain; anxiety, depression, enjoyment, happiness, self-esteem, stress and well-being in social and emotional health domain; and concentration and focus in the domain of educational performance (table 3). We sent the agreed set of outcomes for review to the stakeholders unable to attend the meeting. The wider group approved the COS.

## DISCUSSION

We have developed the first COS for physical activity interventions in primary schools. By using robust consensus

**Table 1** List of 50 outcomes and their descriptions by domain and the number of studies from which the outcomes were extracted

Domain	Outcomes measured	Description*	Studies†	
1: Physical activity and health	Active travel	To get to and from school, for example, walking, public transport, that is, train/tube/bus (do not include car, van, motorcycle), cycling and scooter	FG‡	
	Anthropometry§	Weight, height, body mass index body fat, body mass and waist circumference	34	
	Blood lipids	Fatty substances found in the blood (ie, cholesterol, triglycerides) which increase the risk of heart attack	2	
	Blood pressure	The force at which your heart pumps blood around your body and the resistance to the blood flow in the blood vessels	2	
	Diet	Varied and balanced diet including fruit and vegetables	FG‡	
	Energy levels/expenditure	The amount of energy needed to carry out physical functions such as breathing, exercising or digesting food	5	
	Fitness	Being fit and healthy for optimal health and overall well-being	16	
	Heart rate	Number of beats per minute to establish normal resting heart rate, high or low heart rate	5	
	Intensity of physical activity	Includes light activity (ie, taking a stroll); moderate activity (ie, cycling/swimming at regular pace, sweeping, washing windows); and vigorous activity (ie, aerobics, running, fast cycling, climbing stairs)	42	
	Leisure time activity	Time spent in activity for leisure during the day (ie, walking in the park, playing sports with friends/family)	FG‡	
	Motor skills	Skills that require using large muscles of the arms/legs/torso, that is, standing, walking, going up and down stairs, running, swimming, jumping, skipping, leaping and kicking	8	
	Musculoskeletal	Bone strength, bone mineral density and muscle¶	8	
	Peak oxygen intake	The maximal rate at which oxygen can be used by the body during maximal work	1	
	Sedentary time	Time spent sitting at desk, reading, sitting or lying down to watch television	7	
	Sleep	Between approximately 10–12 hours per night	FG‡	
	2: Social and emotional health	Step counts	Number of steps taken in a day	13
		Anxiety	Persistent feeling of worry, fear or nervousness	FG‡
Appetite		Eating well and regularly	FG*	
Body awareness		The ability to recognise one's body moves helping to understand how to relate to objects and people at home, at school and outdoors	1	
Body image		The perception one has of their physical self	1	
Depression		Feeling persistently sad for more than a few days	FG‡	
Empowerment		Feeling a sense of becoming stronger and more confident	FG‡	
Enjoyment		Taking pleasure in doing something	3	
Happiness		Feeling a sense of joy and contentment	FG‡	

Continued

Table 1 Continued

Domain	Outcomes measured	Description*	Studies†
	Mood	A state of mind or a feeling such as happy, sad, cheerful or angry	FG‡
	Peer support	Using one's own experiences to help others	1
	Resilience	The ability to recover quickly from difficulties	FG‡
	Satisfaction	A sense of fulfilling a need, desire or appetite	FG‡
	Self-confidence	A feeling of trust in one's abilities, qualities and judgement	FG‡
	Self-efficacy	A person's belief of their capacity to perform behaviours necessary to produce specific performance attainments	2
	Self-esteem	A factor that influences people's choices and decisions which results in them either taking or not taking care of themselves and explore their full potential	1
	Self-expression	The communication of one's personality, feelings or opinions	FG‡
	Self-perception	Attitudes towards own preferences and behaviour	1
	Sickness	Feeling unwell, nauseous and dizzy	FG‡
	Sleep patterns	Sleep patterns/achieving less than recommended (10–12 hours)/broken sleep	FG‡
	Social interaction	An exchange between two or more people	FG‡
	Stress	Feeling under pressure or threatened	FG‡
	Well-being	Feeling well, happy, healthy and ability to manage stress	FG‡
3: Educational performance	Academic performance	Measurement of a child's achievement over a range of academic subjects	20
	Attention	Taking notice of someone or something	6
	Classroom behaviour	How children are acting in the classroom in response to what is going on or present around them	15
	Cognitive development/function	How children think, explore and figure things out	6
	Concentration	Ability to focus on task	16
	Engagement	The degree of attention, curiosity, interest, optimism and passion that children show when they are learning or being taught	5
	Executive functioning	A set of mental skills including working memory, flexible thinking and self-control to apply to everyday learning, work, and daily life	4
	Focus	Ability to concentrate and not easily distracted	16
	Maths	The study of numbers, shapes and patterns	16
	Reading	A cognitive process that involves decoding symbols to arrive at meaning, the primary purpose of which is to understand the text	8
	Working memory/inhibition	A cognitive system with a limited capacity that can hold information temporarily and is important for reasoning, decision-making and behaviour	6
	Writing	A form of communication to express language using symbols; being able to understand grammar, punctuation, spelling and vocabulary	5

\*Descriptions were guided by the published literature and our steering group.

†From the 74 studies identified from the 53 relevant reviews.

‡FG=outcome identified by our focus group (steering group).

§Anthropometry was presented as 'bioimpedance' to participants, changed to 'anthropometry' based on reviewer suggestions.

¶Muscle was not included in the original description presented to participants. This was added based on reviewer suggestions.

**Table 2** Outcomes rated ‘not important’ and ‘critical’ to measure after Delphi survey Round 2 (n=60)

Domain	Outcome	Participants rating outcomes ‘not important’ %	Participants rating outcomes ‘critical’ %	
1. Physical activity and health	Active travel	3	51	
	Anthropometry*	15	26	
	Blood lipids	32	14	
	Blood pressure	28	14	
	Diet (varied and balanced)†	3	71†	
	Energy	8	26	
	Fitness	0	60	
	Heart rate	20	17	
	Intensity of physical activity	3	63	
	Leisure time activity	3	62	
	Motor skills	8	46	
	Musculoskeletal	12	20	
	Oxygen peak intake	29	9	
	Sedentary time	3	63	
	Sleep (number of hours)†	3	85†	
	Step counts	12	23	
	2. Social and emotional health	Anxiety†	0	78†
		Appetite	8	42
		Body awareness	2	46
Body image		2	66	
Depression		3	74	
Empowerment		2	42	
Enjoyment†		0	74†	
Happiness†		0	85†	
Mood		0	51	
Peer support		0	46	
Resilience		3	55	
Satisfaction		2	46	
Self-confidence†		0	74†	
Self-efficacy		2	68	
Self-esteem†		0	75†	
Self-expression		8	34	
Self-perception		2	51	
Sickness		12	40	
Sleep patterns		3	69	
Social interaction	0	65		
Stress	0	72†		
Well-being†	0	85†		

Continued

**Table 2** Continued

Domain	Outcome	Participants rating outcomes ‘not important’ %	Participants rating outcomes ‘critical’ %
3. Educational performance	Academic performance	2	57
	Attention†	0†	74†
	Classroom behaviour	2	68
	Cognition	2	54
	Concentration†	0	75†
	Engagement	0	69
	Executive functioning	2	46
	Focus†	3	72†
	Maths	8	55
	Memory	2	48
	Reading	8	51
	Writing	8	48

\*Anthropometry was presented as ‘bioimpedance’ to the participants. This was changed based on reviewer comments.  
 †Ratings that met the threshold ( $\leq 15\%$  agreement of the outcome rated ‘not important’ and  $>70\%$  agreement of the outcome rated ‘critical’ to measure.

methods and multidisciplinary stakeholder groups, we have achieved consensus on the outcomes considered important to measure. Implementation of this COS in future studies will reduce heterogeneity between studies allowing for evidence synthesis and will also be relevant to wider audiences.

During the consensus meeting, it was noted that the survey identified only two outcomes (sleep and diet) in

**Table 3** Core outcome set for physical activity interventions in primary schools

Domain	Outcome
Physical activity and health	Diet (varied and balanced)
	Energy
	Fitness
	Intensity of physical activity
	Sleep (number of hours)
Social and emotional health	Anxiety
	Depression
	Enjoyment
	Happiness
	Self-esteem
Educational performance	Stress
	Well-being
Educational performance	Concentration
	Focus



the domain of physical activity and health as critical to measure, while the outcomes 'physical activity intensity' and 'fitness' did not meet the threshold. Outcomes that may fit under this domain include moderate physical activity, vigorous physical activity, moderate-to-vigorous physical activity and heart rate, which are more commonly studied but these did not meet the critical threshold in our survey. This potentially reflects the heterogeneity across studies of the outcomes that should be measured under broader concepts. As discussed in our consensus meeting, the under-representation of outcomes rated critically important in the physical activity domain may have been due to the specificity of outcomes listed. For example, researchers agree that physical activity should be measured but do not agree on which specific outcome to measure it. This would explain the wide variation of physical activity outcomes that were identified from the published literature. Physical activity can have many benefits beyond measuring its impact on particular health or clinical outcomes. Therefore, our participants agreed that measuring physical activity is important and should be included.

In the published literature, we found only 10 studies which measured outcomes that related to mental health, yet all our stakeholders placed critical importance on many of the outcomes under the domain of social and emotional health. These findings may be explained by the growing awareness of poor mental health in children and the growing evidence base of associations between increased physical and better mental health. The importance placed on mental health perhaps indicates a shift in focus from measuring physiological outcomes and towards measuring mental health when assessing physical activity interventions in primary schools. This may allow health professionals/researchers/teachers/parents to be able tackle better mental health in childhood which may lead to better mental health in adolescence and adulthood. These findings further support the need for a COS in this field. Our study has provided a better understanding that to achieve better overall health and well-being in children, both physical and mental health are important to measure.

Functional precursors of performance-related outcomes (concentration, attention and focus) met the critical threshold than actual educational attainment outcomes of reading, writing and maths which are more commonly assessed in previous studies and by schools. A possible explanation for this is that to improve educational attainment, physical activity interventions need to help to improve cognition (ie, concentration, focus). These interventions may therefore have an indirect effect on improving reading, writing and maths by improving cognition. Schools provide children with learning a range of subjects. However, if increased physical activity in schools enhances children's learning by improving their physical and mental health, this will likely increase the acceptability of physical activity interventions in schools. This may therefore generate

a greater interest from schools to implement these interventions.

Although we are not aware of another COS that specifically evaluates interventions aimed at increasing children's physical activity in primary schools or other settings such as in the community, there are several existing frameworks for assessing these interventions. A systematic review by Cassar *et al*<sup>34</sup> identified 14 frameworks applied across 27 papers<sup>34</sup> which included reach, effectiveness, adoption, implementation and maintenance framework,<sup>35</sup> ecological framework for understanding effective implementation,<sup>36</sup> multilevel implementation quality framework<sup>37</sup> and a conceptual framework for implementation.<sup>38</sup> The review found that the frameworks were primarily used for interpreting results and analyses rather than being used as a planning tool for outcomes to be measured or for understanding results.<sup>34</sup> Another review by Damschroder *et al*<sup>39</sup> also found little evidence that frameworks for school-based physical activity interventions were used to guide the data collection.<sup>39</sup> Findings from these reviews imply that the frameworks to assess these interventions provide little emphasis on the planning of what should be measured and perhaps explain the heterogeneity of outcomes measured to date. A study by McKay *et al*<sup>40</sup> prioritised a list of frameworks to improve the quality and consistency of implementing interventions to ensure that interventions are effectively delivered to achieve population level benefits.<sup>40</sup> COSs should be used to inform the choice of outcomes<sup>41</sup> and our COS contributes to an important gap in these frameworks and can add to them by providing a guide on the minimum set of outcomes to measure in future studies of physical activity interventions in primary schools. It is important to note however that the existing research from physical activity intervention studies has enabled important findings of outcomes that are more commonly measured such as BMI<sup>42</sup> and physical activity<sup>43</sup> and have allowed for a better understanding of the impacts of these interventions on these outcomes. But any COSs currently being developed are mainly centred around childhood obesity<sup>44-46</sup> which is complex; tackling childhood obesity requires comprehensive, multicomponent strategies. Developing COSs require the need to consider the aims and scale of the intervention, the population groups being targeted and the needs of the stakeholders. Our COS, focused on physical activity interventions in primary schools and developed in consultation with those who would benefit the most to better understand intervention effects, should be considered as part of a set of tools for wider improvement of health in primary schools.

Our study's strengths include: we have developed the first COS for physical activity interventions in primary schools, to our knowledge, and used robust methodology as recommended by the COMET to capture a wide range of outcomes to reach consensus. Our inclusion of participants from four key stakeholder groups representing nine countries, as well as incorporating views of children, ensures the relevance of outcomes



to measure for the target population. We also ensured that the domains were not predetermined. We instead established the domains led by the list of outcomes and their descriptions, thus avoiding any researcher bias. However, there are limitations to our study. The descriptions of each outcome were guided by the published literature. We had found variations in how the outcomes were described across studies. This resulted in our descriptions for each outcome either being a definition, suggestion, implying a positively directed relationship or a combination of these. Further research is needed to identify neutral descriptions of outcomes. The low attendance of participants in our consensus meeting which did not include a representation for the educators stakeholder group, may have possibly limited further discussions of the outcomes that should be included in the COS. However, the final list of outcomes was circulated to all the participants who completed both rounds of the Delphi survey and an opportunity to comment further was provided before the final outcome set was agreed. As we recruited participants through several methods including advertising on our research network websites and through snowballing, we are not aware of how many potential participants were targeted for our research and did not participate. Although our participants represented nine countries, most were UK based. The educators and health researcher stakeholder groups included participants from five countries, while participants from two countries represented the public health professional and parent groups. All stakeholder groups had a UK participant representation between 71% and 95%. The outcomes identified from our umbrella review were not limited to UK-based studies, but the lower proportion of participants representing other countries and in each stakeholder group may have prevented the identification of other outcomes that may be more relevant. Other countries and cultures may differ in the importance placed on physical activity in schools and may focus on other aspects such as educational attainment. This may bias our COS towards outcomes relevant to UK audiences. COVID-19 restrictions limited our reach to primary schools and year groups to target for our workshops; children from different year groups may have considered additional or fewer outcomes important. In addition, our representation of children with English as a second language was much higher (75%) than the average number of children with English as a second language in London primary schools (48%).<sup>47</sup> The development of our COS during the COVID-19 pandemic may have influenced our findings. It has been widely reported that school closures and restrictions have reduced opportunities for children to be physically active and has increased poorer mental health.<sup>48 49</sup> This may perhaps explain the higher number of outcomes in the domain of social and emotional health that met the threshold in our surveys. Finally, it may be challenging for future studies to include all 14 outcomes identified in our COS. However, as our outcomes have been grouped into three

main domains, researchers may choose to include the outcomes within the domain of interest.

The development of our COS is timely; several interventions that have been implemented in schools in recent year may have stopped due to COVID-19. These interventions are likely to resume and may be more important to assess now due to the negative impacts the pandemic has had on children's physical activity and mental health. Our COS would be relevant to future studies assessing the impact of physical activity interventions in primary schools such as The Daily Mile, a popular active mile intervention reaching one in five state-funded primary schools in England,<sup>50</sup> and recommended by England's National Obesity Plan.<sup>51</sup> Despite its reach, the evidence of its impact remains limited or inconsistent.<sup>52-55</sup>

Our COS would benefit from identifying the best assessment tools to measure the outcomes that are readily available to those implementing physical activity interventions in schools. COMET suggests that a COS use should first aim to establish which outcomes are important to measure, and then aim to identify which assessment tools would be the most accessible for end users.<sup>56</sup> There is a low uptake of COSs in randomised control trials due to lack of recommendations of valid measures, lack of involvement of key stakeholders and those implementing or assessing interventions not being aware of a COS in their field of research.<sup>56</sup> Our next step is to identify assessment tools that are readily available to measure the outcomes in our COS. Recommendations of assessment tools would further enhance the quality and consistency of results in studies using our COS.

Prevention and public health approaches in early life to reduce health inequalities and improve health of the whole population may be a better investment than treating disease in the population that generally arises later in life.<sup>57 58</sup> The robust processes that we have applied in this study could be repeated to inform an adolescent (young people aged 12–17 years) focused COS. Physical activity is low among the secondary school population<sup>59</sup> and poorer mental health is also increasing among this age group.<sup>60</sup> We recommend that our COS is included as part of a wider set of tools and frameworks that should be developed to standardise the outcomes to measure other areas of children and young people's health such as weight and nutrition.<sup>61</sup> This would allow for improved health to continue during adolescence and adulthood.

## Conclusion

Our COS identifies the outcomes that are most important to measure for studies of physical activity interventions in primary schools. Next, we aim to identify the assessment tools to measure these outcomes. Wide use of our COS in future studies will reduce heterogeneity allowing for evidence synthesis to better understand intervention effects on children's health and cognition during the primary school years.

Twitter Bina Ram @DrBinaRam and Sonia Saxena @SoniaKSaxena

**Acknowledgements** The authors are grateful for support from the National Institute for Health Research (NIHR) School for Public Health Research (SPHR) and the NIHR Applied Research Collaboration Northwest London (ARC NWL). This paper is independent research supported by the NIHR ARC NWL. The authors thank their steering committee for their input and guidance and Esta Orchard for facilitating the children's workshop and stakeholder meeting. The authors thank all the participants who took part in the study including Molly Adkin, Teatske Altenburg, Thomas Beaney, CÅ-ntia Botton, Anna Chalkley, Paul Chapman, Lotte de Vries, Stuart Fairclough, Lawrence Foweather, Chris Gale, Charan Gill, Muhamet Halilaj, Deirdre Harrington, Frances Hillier-Brown, Josie Hopkins, Daniel Katon, Michelle Kelly-Irving, Amy King, Edward Maile, Natalie McConnon, Catherine McKay, Colin Moran, Francesca Neale, Dasha Nicholls, Nikita Punjabi, Anelise Reis Gaya, Mairead Ryan, Tajvir Singh, Jorien Slot-Heijs, Rodrigo Sudatti Delevatti, Cathryn Taylor, Shannon Tessier, Daniel Umpierre, Esther van Sluijs, Tishya Venkatraman, Julian Winn, Kathryn Woods-Townsend, Callum Wright, Maddie Yang. We would also like to thank the children, the PE specialist teacher (Mr Alex Wilke) and the head teacher (Mr Raphael Moss) at Elsley Primary School. The authors thank The Daily Mile Foundation for funding and support.

**Contributors** KAF, SS and BR conceived and designed the study. BR and KAF designed study materials. BR was responsible for managing all components of the study including recruitment, data collection and analyses. SS, EvS, DSH and RMV advised on the project. BR wrote the first draft of the manuscript. All authors contributed to re-drafts and approved the final manuscript. The guarantor (SS) accepts full responsibility for the finished work and/or the conduct of the study, had access to the data, and controlled the decision to publish.

**Funding** This work was funded by The Daily Mile Foundation supported by INEOS and was supported by the Children and Families Programme, National Institute for Health Research School for Public Health Research (NIHR SPHR) (Grant Reference Number PD-SPH-2015 (PROG-CYP-WP2)) and the NIHR Applied Research Collaboration Northwest London (NIHR ARC NWL). BR was funded by The Daily Mile Foundation. SS was funded by The Daily Mile Foundation, NIHR SPHR and the NIHR ARC NWL. KAF was supported by NIHR ARC NWL. DSH was supported by the NIHR SPHR and NIHR ARC NWL. EvS was supported by the Medical Research Council (Grant MC\_UU\_12015/7). The NIHR School for Public Health Research is a partnership between the Universities of: Sheffield; Bristol; Cambridge; Imperial; and University College London; The London School for Hygiene and Tropical Medicine; LiLaC—a collaboration between the Universities of Liverpool and Lancaster; and Fuse—The Centre for Translational Research in Public Health a collaboration between Newcastle, Durham, Northumbria, Sunderland and Teesside Universities. The Department of Primary Care and Public Health is grateful for support from the NIHR Biomedical Research Centre funding scheme, NIHR SPHR and NIHR ARC NWL.

**Disclaimer** The funders had no role in the analyses or results of this study. The views expressed in this publication are those of the authors and not necessarily those of NIHR, the Department of Health and Social Care, or The Daily Mile Foundation.

**Competing interests** SS and BR received funding from The Daily Mile Foundation. SS, BR and KAF are members of The Daily Mile Research Advisory Group. All other authors declare no competing interests.

**Patient and public involvement** Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was approved by Imperial College London research ethics committee: Reference Number 19IC5428. Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. All data relevant to the study are included in the article or uploaded as supplementary information.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Bina Ram <http://orcid.org/0000-0003-0023-1573>  
 Kimberley A Foley <http://orcid.org/0000-0003-3664-8100>  
 Russell M Viner <http://orcid.org/0000-0003-3047-2247>  
 Sonia Saxena <http://orcid.org/0000-0003-3787-2083>

#### REFERENCES

- 1 Chaput J-P, Willumsen J, Bull F, *et al.* 2020 WHO guidelines on physical activity and sedentary behaviour for children and adolescents aged 5-17 years: summary of the evidence. *Int J Behav Nutr Phys Act* 2020;17:141.
- 2 NCMP. *National child measurement programme (NCMP), England 2018/19 school year*. United Kingdom: National Statistics, NHS Digital, 2019.
- 3 OECD. *Future of education 2030: making physical education dynamic and inclusive for 2030. international curriculum analysis*. Paris, France: Organisation for Economic Co-operation and Development (OECD), 2019.
- 4 WHO. *Report of the Commission on ending childhood obesity*. Geneva, Switzerland: World Health Organisation, 2016.
- 5 Wang Y, Cai L, Wu Y, *et al.* What childhood obesity prevention programmes work? A systematic review and meta-analysis. *Obes Rev* 2015;16:547-65.
- 6 Waters E, de Silva-Sanigorski A, Hall BJ. Interventions for preventing obesity in children. *Cochrane Database Syst Rev* 2011;12:CD001871.
- 7 Love R, Adams J, Atkin A, *et al.* Socioeconomic and ethnic differences in children's vigorous intensity physical activity: a cross-sectional analysis of the UK millennium cohort study. *BMJ Open* 2019;9:e027627.
- 8 Marmot M, Allen J, Boyce T. *Health equity in England: the Marmot review 10 years on*. London: Institute of Health Equity, 2020.
- 9 Kriemler S, Zahner L, Schindler C, *et al.* Effect of school based physical activity programme (kiss) on fitness and adiposity in primary schoolchildren: cluster randomised controlled trial. *BMJ* 2010;340:c785.
- 10 Liu A, Hu X, Ma G, *et al.* Evaluation of a classroom-based physical activity promoting programme. *Obes Rev* 2008;9 Suppl 1:130-4.
- 11 Daly-Smith AJ, Zvolinsky S, McKenna J. Systematic review of acute physically active learning and classroom movement breaks on children's physical activity, cognition, academic performance and classroom behaviour: understanding critical design features. *Bmj Open Sport Exerc* 2018;4.
- 12 The Daily Mile Foundation. The daily mile. Available: <https://thedailymile.co.uk/>
- 13 Marathon Kids UK, 2010. Available: <https://www.marathonkids.co.uk/>
- 14 Kriemler S, Meyer U, Martin E, *et al.* Effect of school-based interventions on physical activity and fitness in children and adolescents: a review of reviews and systematic update. *Br J Sports Med* 2011;45:923-30.
- 15 Hartwig TB, Sanders T, Vasconcellos D, *et al.* School-Based interventions modestly increase physical activity and cardiorespiratory fitness but are least effective for youth who need them most: an individual participant pooled analysis of 20 controlled trials. *Br J Sports Med* 2021;55:721-9.
- 16 Pozuelo-Carrascosa DP, García-Hermoso A, Álvarez-Bueno C, *et al.* Effectiveness of school-based physical activity programmes on cardiorespiratory fitness in children: a meta-analysis of randomised controlled trials. *Br J Sports Med* 2018;52:1234-40.
- 17 Minatto G, Barbosa Filho VC, Berria J, *et al.* School-Based interventions to improve cardiorespiratory fitness in adolescents: systematic review with meta-analysis. *Sports Med* 2016;46:1273-92.
- 18 van Grieken A, Ezendam NPM, Paulis WD, *et al.* Primary prevention of overweight in children and adolescents: a meta-analysis of the effectiveness of interventions aiming to decrease sedentary behaviour. *Int J Behav Nutr Phys Act* 2012;9:61.
- 19 Hegarty LM, Mair JL, Kirby K, *et al.* School-Based interventions to reduce sedentary behaviour in children: a systematic review. *AIMS Public Health* 2016;3:520-41.
- 20 Andermo S, Hallgren M, Nguyen T-T-D, *et al.* School-related physical activity interventions and mental health among children: a systematic review and meta-analysis. *Sports Med Open* 2020;6:25.

- 21 Álvarez-Bueno C, Pesce C, Cavero-Redondo I, *et al.* The Effect of Physical Activity Interventions on Children's Cognition and Metacognition: A Systematic Review and Meta-Analysis. *J Am Acad Child Adolesc Psychiatry* 2017;56:729–38.
- 22 Watson A, Timperio A, Brown H, *et al.* Effect of classroom-based physical activity interventions on academic and physical activity outcomes: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act* 2017;14:114.
- 23 Carter T, Pascoe M, Bastounis A, *et al.* The effect of physical activity on anxiety in children and young people: a systematic review and meta-analysis. *J Affect Disord* 2021;285:10–21.
- 24 Dobbins M, Husson H, DeCorby K, *et al.* School-Based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6 to 18. *Cochrane Database Syst Rev* 2013;2:CD007651.
- 25 Neil-Sztramko SE, Caldwell H, Dobbins M. School-Based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6 to 18. *Cochrane Database Syst Rev* 2021;9:CD007651.
- 26 Hollis JL, Sutherland R, Campbell L, *et al.* Effects of a 'school-based' physical activity intervention on adiposity in adolescents from economically disadvantaged communities: secondary outcomes of the 'Physical Activity 4 Everyone' RCT. *Int J Obes* 2016;40:1486–93.
- 27 Mei H, Xiong Y, Xie S, *et al.* The impact of long-term school-based physical activity interventions on body mass index of primary school children - a meta-analysis of randomized controlled trials. *BMC Public Health* 2016;16:205.
- 28 Williamson PR, Altman DG, Blazey JM, *et al.* Developing core outcome sets for clinical trials: issues to consider. *Trials* 2012;13:132.
- 29 Initiative C. Core outcome measures in effectiveness trials, 2010. Available: <https://www.comet-initiative.org/>
- 30 Foley KA, Venkatraman T, Ram B, *et al.* Protocol for developing a core outcome set for evaluating school-based physical activity interventions in primary schools. *BMJ Open* 2019;9:e031868.
- 31 Saxena S, Foley K, Venkatraman T. Registration to comet initiative: developing a core outcomes set for evaluation of physical activity interventions in primary schools UK, 2019. Available: <https://www.comet-initiative.org/Studies/Details/1322>
- 32 Kirkham JJ, Gorst S, Altman DG, *et al.* Core outcome Set-Standards for reporting: the COS-STAR statement. *PLoS Med* 2016;13:e1002148.
- 33 Initiative C. Delphi manager UK. Available: <https://www.comet-initiative.org/delphimanager/>
- 34 Cassar S, Salmon J, Timperio A, *et al.* Adoption, implementation and sustainability of school-based physical activity and sedentary behaviour interventions in real-world settings: a systematic review. *Int J Behav Nutr Phys Act* 2019;16:120.
- 35 Glasgow RE, Vogt TM, Boles SM. Evaluating the public health impact of health promotion interventions: the RE-AIM framework. *Am J Public Health* 1999;89:1322–7.
- 36 Durlak JA, DuPre EP. Implementation matters: a review of research on the influence of implementation on program outcomes and the factors affecting implementation. *Am J Community Psychol* 2008;41:327–50.
- 37 Domitrovich CE, Bradshaw CP, Poduska JM, *et al.* Maximizing the implementation quality of evidence-based preventive interventions in schools: a conceptual framework. *Adv Sch Ment Health Promot* 2008;1:6–28.
- 38 Fixsen DL, Naoom SF, Blase KA. Implementation research: a synthesis of the literature. Tampa, Florida: the National implementation research network, University of South Florida, Louis de la parte Florida mental health Institute. *FMHl Publication* 2005:231.
- 39 Damschroder LJ, Aron DC, Keith RE, *et al.* Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci* 2009;4:50.
- 40 McKay H, Naylor P-J, Lau E, *et al.* Implementation and scale-up of physical activity and behavioural nutrition interventions: an evaluation roadmap. *Int J Behav Nutr Phys Act* 2019;16:102.
- 41 Williamson PR, de Ávila Oliveira R, Clarke M, *et al.* Assessing the relevance and uptake of core outcome sets (an agreed minimum collection of outcomes to measure in research studies) in Cochrane systematic reviews: a review. *BMJ Open* 2020;10:e036562.
- 42 Liu Z, Xu H-M, Wen L-M, *et al.* A systematic review and meta-analysis of the overall effects of school-based obesity prevention interventions and effect differences by intervention components. *Int J Behav Nutr Phys Act* 2019;16:95.
- 43 Love R, Adams J, van Sluijs EMF. Are school-based physical activity interventions effective and equitable? A meta-analysis of cluster randomized controlled trials with accelerometer-assessed activity. *Obes Rev* 2019;20:859–70.
- 44 Wirix A, Altenburg T, van de Beek C. Defining core outcomes for school-based intervention studies on preventing childhood overweight and obesity (Ref: 971): comet initiative; 2017, -2019. Available: <https://www.comet-initiative.org/studies/details/971> [Accessed July 2021].
- 45 Botton C, De-Nardi A, Pfeifer L. Core outcomes in children and adolescents with overweight/obesity exposed to physical activity interventions (Ref: 1356) comet Initiative2019-2020. Available: <https://www.comet-initiative.org/studies/details/1356> [Accessed July 2021].
- 46 Benson AC, Torode ME, Fiatarone Singh MA. Effects of resistance training on metabolic fitness in children and adolescents: a systematic review. *Obesity Reviews* 2007;9:070522225912002-???.
- 47 Department for Education. Percentage of pupils by first language, borough. UK, 2018. Available: <https://data.london.gov.uk/dataset/percentage-pupils-first-language-borough> [Accessed June 2022].
- 48 Youth Sports Trust. *Evidence paper: the impact of Covid-19 restrictions on children and young people.* Leicestershire, UK, 2020.
- 49 Viner R, Russell S, Saull R. Impacts of school closures on physical and mental health of children and young people: a systematic review. *medRxiv* 2021.
- 50 Venkatraman T, Honeyford K, Costelloe CE. Sociodemographic profiles, educational attainment and physical activity associated with the daily Mile™ registration in primary schools in England: a national cross-sectional linkage study. *J Epidemiol Community Health* 2020;20:jech-2020-214203–8.
- 51 HM Government. a plan for action Chapter 2. In: *Childhood obesity.* United Kingdom, 2018.
- 52 Chesham RA, Booth JN, Sweeney EL, *et al.* The daily mile makes primary school children more active, less sedentary and improves their fitness and body composition: a quasi-experimental pilot study. *BMC Med* 2018;16:64.
- 53 Morris JL, Daly-Smith A, Archbold VJS, *et al.* The daily Mile™ initiative: exploring physical activity and the acute effects on executive function and academic performance in primary school children. *Psychol Sport Exerc* 2019;45:101583.
- 54 Chalkley AE, Routen AC, Harris JP, *et al.* "I Just Like the Feeling of It, Outside Being Active": Pupils' Experiences of a School-Based Running Program, a Qualitative Study. *J Sport Exerc Psychol* 2020;42:48–58.
- 55 Marchant E, Todd C, Stratton G, *et al.* The daily mile: Whole-school recommendations for implementation and sustainability. A mixed-methods study. *PLoS One* 2020;15:e0228149.
- 56 Hughes KL, Clarke M, Williamson PR. A systematic review finds core outcome set uptake varies widely across different areas of health. *J Clin Epidemiol* 2021;129:114–23.
- 57 The Academy of medical sciences. *Improving the health of the public by 2040: optimising the research environment for a healthier, fairer future.* London, UK, 2016.
- 58 WHO. *Global action plan on physical activity 2018–2030: more active people for a healthier world Geneva.* World Health Organization, 2018.
- 59 van Sluijs EMF, Ekelund U, Crochemore-Silva I, *et al.* Physical activity behaviours in adolescence: current evidence and opportunities for intervention. *Lancet* 2021;398:429–42.
- 60 WHO. Adolescent mental health: the world health organisation, 2020. Available: <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health> [Accessed July 2021].
- 61 Hargreaves D, Mates E, Menon P, *et al.* Strategies and interventions for healthy adolescent growth, nutrition, and development. *Lancet* 2022;399:198–210.