

Supplementary Table 1. Baseline characteristics of study patients overall and by PASE questionnaire respondent

	PASE questionnaire respondent		
	Overall	Next of kin	Patient
N	625	61 (10%)	564 (90%)
PASE score quartile, n (%)			
1st	157 (25%)	15 (25%)	142 (25%)
2nd	156 (25%)	18 (30%)	138 (24%)
3rd	156 (25%)	14 (23%)	142 (25%)
4th	156 (25%)	14 (23%)	142 (25%)
Study treatment, n (%)			
Citalopram	309 (49%)	31 (51%)	278 (49%)
Placebo	316 (51%)	30 (49%)	286 (51%)
Female, n (%)	215 (34%)	27 (44%)	188 (33%)
Age, median (IQR)	69 (60, 77)	70 (63, 77)	69 (59, 77)
Cohabitation, n (%)	409 (65%)	38 (62%)	371 (66%)
History of smoking, n (%)	195 (32%)	24 (41%)	171 (31%)
Missing	17	2	15
Diabetes, n (%)	71 (11%)	6 (9.8%)	65 (12%)
Missing	7	0	7
Hypertension, n (%)	325 (52%)	32 (52%)	293 (52%)
Missing	5	0	5
Atrial Fibrillation, n (%)	102 (17%)	15 (25%)	87 (16%)
Missing	7	1	6
Prior MI, n (%)	52 (8.4%)	7 (12%)	45 (8.1%)
Missing	9	1	8
Prior TIA, n (%)	16 (2.6%)	2 (3.3%)	14 (2.5%)
Missing	9	0	9
PAD, n (%)	25 (4.1%)	4 (6.6%)	21 (3.8%)
Missing	14	0	14
Baseline NIHSS, median (IQR)	3 (2, 6)	11 (4, 18.5)	3 (2, 5)
Missing	12	2	10

Reperfusion, n(%)	237 (38%)	33 (54%)	204 (36%)
Follow-up time, days, median (IQR)	180 (161, 185)	175 (40, 183)	180 (167, 185)

PASE: Physical Activity Scale for the Elderly, MI: myocardial infarction, PAD: peripheral arterial disease, NIHSS: National Institute of Health Stroke Scale, IQR: interquartile range