


BMJ Open Effect of environmental enrichment on relapse rates in patients with severe alcohol use disorder: protocol for a randomised controlled trial

Lila Barillot ^{1,2}, Claudia Chauvet,^{2,3} Marc Besnier,^{1,2} Nematollah Jaafari,^{1,2} Marcello Solinas,^{2,4} Armand Chatard^{1,2}

To cite: Barillot L, Chauvet C, Besnier M, *et al.* Effect of environmental enrichment on relapse rates in patients with severe alcohol use disorder: protocol for a randomised controlled trial. *BMJ Open* 2023;**13**:e069249. doi:10.1136/bmjopen-2022-069249

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-069249>).

Received 14 October 2022
Accepted 06 April 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Université de Poitiers, Centre de Recherches sur la Cognition et l'Apprentissage, Poitiers, Nouvelle-Aquitaine, France

²Unité de Recherche Clinique Pierre Deniker, Centre Hospitalier Henri Laborit, Poitiers, Poitou-Charentes, France

³Service Hospitalo-Universitaire de Psychiatrie et de Psychologie Médicale, Centre Hospitalier Henri Laborit, Poitiers, France

⁴Université de Poitiers, Laboratoire de Neurosciences Expérimentales et Cliniques U1084, Poitiers, France

Correspondence to

Lila Barillot;
lila.barillot@univ-poitiers.fr

ABSTRACT

Introduction Alcohol use disorder (AUD) ranks among the most prevalent psychiatric disorders worldwide. Despite current treatments, more than half of patients relapse within weeks after treatment. In animal models, exposure to environmental enrichment (EE) has been shown to be a promising approach to reduce relapse. However, controlled, multimodal EE is difficult to transpose to humans. To address this gap, this study aims at assessing the effectiveness of exposure to a newly designed EE protocol during AUD treatment in reducing relapse to alcohol use. Our EE will allow an enhancement of the standard intervention, and will combine several promising enrichment factors identified in the literature—physical activity, cognitive stimulation, mindfulness and virtual reality (VR).

Methods and analysis A randomised controlled trial involving 135 participants receiving treatment for severe AUD will be conducted. Patients will be randomised to an intervention enhancement group or a control group. The enhanced intervention will consist of six 40-min sessions of EE spread over 9 days. During the first 20 min of these sessions, patients will practise mindfulness in multisensory VR, in virtual environments designed to practise mindfulness and use it to regulate craving induced by virtual cues or stress. Then, participants will practise indoor cycling combined with cognitive training exercises. The control group will undergo standard management for AUD. The primary outcome is relapse assessed at 2 weeks after treatment, using a questionnaire and biological indicators. Relapse will be defined as drinking at least five drinks per occasion or drinking at least five times a week. It is predicted that the group receiving the EE intervention will have a lower relapse rate than the control group. The secondary outcomes are relapse at 1 month and 3 months after treatment, craving and drug-seeking behaviour, mindfulness skills acquisition and the effect of the intervention enhancement on the perceived richness of the daily environment, assessed by questionnaires and neuropsychological tasks.

Ethics and dissemination All participants have to give written informed consent to the investigator. This study is approved by the Ethics Committee Nord Ouest IV of Lille (reference number 2022-A01156-37). Results will be disseminated through presentations, peer-reviewed journals and seminar conferences. All information on

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This study is the first to attempt to transpose the complex environmental enrichment (EE) used in animal models to humans for the treatment of alcohol use disorder.
- ⇒ Our EE protocol is designed to target the main triggers of relapse, which are stress and cue reactivity.
- ⇒ Repeated assessments and follow-up during the period of vulnerability for alcohol-dependent patients allow a robust evaluation of the outcomes.
- ⇒ The EE used for this intervention does not include specific forms of social stimulation.
- ⇒ Neither the participants nor the therapist can be blinded to the treatment, as this trial includes complex behavioural interventions.

ethical considerations and open science practices can be accessed at <https://osf.io/b57uj/>

Trial registration number NCT05577741.

INTRODUCTION

Alcohol use disorder (AUD) is one of the most prevalent psychiatric disorders, affecting 107 million people worldwide.^{1,2} It is associated with high morbidity and mortality, causing 3 million deaths per year.² Relapse is the most significant obstacle to rehabilitation. Despite the existence of several treatments, 85% of patients treated for AUD relapse, even after long periods of abstinence.³ In particular, the first 3 months of abstinence constitute the period of greatest vulnerability,⁴ and more than half of patients will consume their first drink (lapse) within 2 weeks.⁵ Craving is one of the main predictors of relapse^{6–9} and is triggered by drug use, drug-associated environmental cues and stress.^{10–12} It is therefore necessary to manage craving, through therapy allowing patients to better handle daily stress and cue exposure.

Environmental enrichment

Environmental enrichment (EE) is a promising experimental paradigm to reduce craving and the risk of relapse. The role of the environment in the development and continuation of addictive behaviour is well demonstrated.^{13–21} In addition, environmental conditions during periods of abstinence represent an interesting opportunity for intervention.²¹ Preclinical studies have found that EE combining different types of stimulation can prevent the development and maintenance of addictive behaviours. EE combines complex social, cognitive and physical stimulation (a large cage, new toys, a racing wheel) that aim to improve sensory, cognitive and motor functions. In animal models of addiction (and for different types of drugs), housing addicted individuals in an EE during a period of abstinence led to extinction of the addictive behaviour and a reduced risk of relapse compared with control animals housed in standard cages.^{18 22 23} This non-pharmacological strategy would prevent relapse by altering the stress response and drug-seeking behaviour, resulting in decreased brain reactivity to cues.^{14 17–19} Behavioural and neurobiological evidence suggest that modulation of the reactivity to stress plays a major role in the effects of EE.^{24 25}

Human studies have examined the effect of different aspects of EE, such as physical activity, cognitive or social stimulation, separately on addiction.¹⁸ However, no study has integrated these different components into a model of EE to provide multimodal stimulation. Yet, preclinical studies suggest that cognitive and physical stimulation can induce additive or complementary action leading to greater neurogenesis when combined, which can produce better therapeutic results.²⁶

Environmental components in the treatment of addiction

Different components of EE have already shown promising results in humans. Physical exercise has positive effects on methamphetamine,²⁷ alcohol²⁸ and nicotine addiction.^{29–32} In particular, several studies have shown positive effects of exercise on stress, mood and craving in alcohol-dependent patients, suggesting that physical activity could be effective in treating addiction.³³ Regarding cognitive stimulation, studies that integrated cognitive training of several cognitive functions (attention, memory, executive functions) have revealed positive effects on different types of addiction, improving cognition, well-being and the compulsive aspect of craving.^{34–37} Interestingly, there exist commercial solutions such as the cognitive bike (vélo-cognitif) that allow performing at the same time physical and cognitive exercise in comfortable, easy and safe conditions that are adapted to hospital settings.

A recent review highlights the potential of combining cognitive exercises with alternative interventions such as mindfulness that can impact both non-cognitive and cognitive processes, particularly executive functions known to be strongly impaired in alcohol-dependent patients.²⁶ Mindfulness practice allows training

attentional reorientation, metacognition, inhibitory control, emotion regulation and interoception. Thus, mindfulness serves as cognitive-behavioural training that promotes well-being while targeting emotional regulation and addiction mechanisms.^{38 39} Many studies have shown positive effects of mindfulness practice on stress levels.^{40 41} In particular, the Mindfulness Based Stress Reduction programme is widely used to improve stress regulation by teaching people to practice mindfulness in stressful situations.⁴² Furthermore, a number of studies suggest that mindfulness interventions allow the reduction of craving, drug consumption and the relapse rate in tobacco and alcohol addiction.^{43 44} In addition, some authors have found that among the methods used in training executive functions, mindfulness and physical activity are particularly promising, facilitating a general improvement in tasks other than those used for training.⁴⁵ An intervention combining these different techniques could therefore be more effective in training the deregulated cognitive and affective processes involved in addiction.²⁶

Virtual reality (VR) is increasingly used in medical protocols to enrich the environment of patients suffering from various cognitive disorders.^{46 47} VR allows exposing patients to different levels of enrichment and stimulation in secure and controlled environments. Moreover, VR helps promote patients' well-being and stimulates them at the cognitive level.⁴⁸ Another benefit is that VR can simulate proximal and contextual cues of risky situations for patients (being in a bar or at a party with people drinking and offering alcohol).^{49 50} Several studies have shown that exposure to cues in VR is particularly effective in inducing craving.^{51–53} Therefore, VR is increasingly used for addiction treatment, mainly in cue exposure protocols, to try to extinguish the stimulus (cue) response (drug consumption) association.^{54 55} In addition, an interesting feature of VR is that it can be used to induce stress,^{56 57} a known trigger of craving and relapse.¹² VR could therefore allow patients to practise regulating their cravings induced by cues or stress in a secure setting, preparing them for a return to everyday life.

VR could facilitate mindfulness practice. Indeed, practicing mindfulness can be complicated for beginners, who may have difficulty staying focused.⁵⁸ In particular, many alcohol-dependent patients have comorbidities, such as depression^{59 60}; these can lead to a loss of motivation that can make active participation on the part of the patient more difficult. One study showed that VR, as a very immersive technology, could compensate for these difficulties by facilitating the allocation of attentional resources to the virtual environment (VE), thus reducing distracting thoughts.⁶¹ The combination of VR and mindfulness can therefore be an interesting tool to provide EE in humans. Practising mindfulness in VEs that induce craving through cues or stress could be particularly useful in training patients to learn how to cope with these situations in their real life.

Aims of the study

The main aim of this study is to assess the effectiveness of exposure to EE combining physical activity, cognitive activity using cognitive bikes and mindfulness in VR to prevent AUD relapse. Our randomised control trial will allocate half of the patients to a control group that will receive only the standard treatment for AUD and half to an intervention enhancement group that will receive several sessions of EE in addition to the standard treatment. We hypothesise that the relapse rate in the group receiving the EE intervention will be lower than in the control group at 2 weeks, 1 month and 3 months after the intervention or the 10th day of inclusion. We also expect the EE intervention to induce a greater decrease in patients' craving and drug-seeking behaviour than standard treatment. It is predicted that the EE intervention will improve patients' mindfulness skills. Finally, we believe that the intervention, by providing alternative rewarding stimuli to drug taking, should encourage a change in behaviour²⁴ and thus in lifestyle (modification of one's environment), and should therefore increase the perceived richness of the daily environment.

METHODS AND ANALYSIS

Trial design

This study is a randomised, controlled, non-blinded trial with two parallel arms comparing an EE intervention group to a control group following standard care. Participants will be randomised at a 1/1 ratio to one of two groups. Both groups will complete a battery of tests and questionnaires on the 1st day of their inclusion and on the 10th day to evaluate craving and mindfulness skills before and after the EE intervention. Follow-ups will be conducted at 2 weeks, 1 month and 3 months after treatment to assess relapse. In addition to these measurement sessions, the intervention enhancement group will carry out six EE sessions.

Study setting

This study will take place in the laboratories of the Pierre Deniker Intersectoral Clinical Research Unit in Psychiatry at the Henri Laborit University Hospital Centre (CHL) in Poitiers, France. The study will end as soon as the number $n=135$ participants is reached or at the end of the 2-year inclusion period.

Participants

One hundred and thirty-five patients undergoing alcohol treatment will be recruited from the Calliope Addiction Unit at the CHL, or from the University Hospital of Poitiers. The base rate of relapse at this site is estimated to be about 50% at 1 month and 60% at 3 months.⁶² Calculation of the sample size is reported in the sample size section.

The inclusion criteria are as follows: patients aged 18–65 treated for alcohol addiction at the CHL in an open ward or at the University Hospital of Poitiers for

at least 48 hours with severe AUD according to the 5th edition of the Diagnostic and Statistical Manual of Mental Disorders classification; benefitting from social security personally or through a third party in accordance with French law on research involving humans; and having signed the informed consent form after having received written information.

The exclusion criteria are as follows: disabling cognitive impairment; susceptibility to cybersickness; cardiological pathologies that could compromise the participation of the patient (detected by an ECG); advanced pulmonary or renal diseases or any unstable and serious medical conditions that could compromise the participation of the patient (subject to the judgement of a doctor); hypertension; ataxia; uncompensated or unstable psychiatric pathology; pregnancy; breast feeding; simultaneous participation in another trial; any other current addiction except addiction to tobacco, Tetrahydrocannabinol and benzodiazepines; being an employee of the investigator or of the clinical study site; being a patient protected by law; not covered by state health insurance; and being unable to complete the questionnaire based on the opinion of the investigator.

Randomisation

Patients are randomised 1:1 to either the control or intervention EE enhancement group. A block randomisation is used with a block size of 4, using an allocation sequence generated from RStudio by a biostatistician.

Intervention

The enhanced intervention consists of six sessions of exposure to EE spread over 9 days (depending on the inclusion date). These sessions take place in addition to the standard intervention and are planned with the patient so as not to interfere with other activities (workshops or therapeutic meetings). The EE is produced using two innovative tools combining different types of stimulation:

The practice of mindfulness in multisensory VR

A total of six virtual environments (VEs) have been developed in collaboration with Sensiks (Amsterdam, Netherlands) using Unity software. An Oculus headset and two joysticks are used to interact with the VEs. To mimic real life and guarantee the participant's immersion, the following interactions are possible in the VEs using the joysticks: moving around the environment by teleportation, catching and throwing virtual objects and ordering a virtual drink using a menu. Teleportation consists of pointing the joystick at predetermined points in the environments, represented by white circles on the ground. When the user points to one of the circles, he/she is immediately teleported to that location, thus limiting the risk of cybersickness due tovection. For each VE, mindfulness instructions are broadcast to enable guided mindfulness while exploring the environments. The mindfulness instructions have been prerecorded by



Figure 1 Virtual environments presented: forest (VE 1), beach (VE 2), bar (VE 3), party (VE 4), parachute jump (VE 5), plane (VE 6).

a therapist specialising in therapeutic relaxation and can be found in online supplemental file 1. These instructions take into account the VE presented and guide the participant through the scenario.

Description of the VEs (figure 1):

Environments 1 and 2 represent relaxing natural places: a virtual forest for the former and a sandy beach for the latter. In these VEs, the participants can catch and throw natural objects (flowers, mushrooms, shells).

Environments 3 and 4 feature places with cues associated with alcohol consumption: bottles of alcohol and avatars drinking in an appropriate context. Mindfulness instructions guide the participant through the following scenario. For VE 3, after a walk down a virtual street, the participant has to buy a bottle of water in a store and then order a coffee in a bar. The participant can grab objects (bottle of alcohol, cigarettes, coins) and buy or order a drink using a virtual menu presenting several choices of alcohol or soft drinks. The scenario for VE 4 is a virtual party in a house. Some avatars dance, smoke and drink alcohol in the living room, and others sit and chat or play cards. The session consists of the participant sitting with them and then going to the kitchen to get a bottle of water. Interactions with objects are the same as in environment 3 (possibility to grab bottles of alcohol, cigarettes or decorative objects), and a menu allows selecting a drink from the fridge, including a glass of water, wine, beer or fruit juice.

Environments 5 and 6 present stressful contexts. The scenario for VE 5 consists of a virtual parachute jump. The participant is immersed in a virtual aeroplane environment with avatars showing signs of stress (frequently looking right, left or out the window; shaky hands; leaning forward), and they have to jump after them.

The participant cannot move freely or grab objects in this environment. The scenario unfolds gradually for 20 min. After the jump, the participant falls into a void, their parachute opens and they gradually descend. At the end of the session, the participant is at ground level. For VE 6, the scene takes place in a virtual aeroplane environment in which there is turbulence. The participant can explore the plane by teleporting. An announcement warning of turbulence is broadcast, and the participant is teleported into a seat and can no longer move. The turbulence consists of shaking of the plane, and there is a thunderstorm, falling luggage and avatars expressing fear (looking right and left, holding their heads in their hands, screaming and sobbing). Once the turbulence has subsided, the participant can grab a book or a bottle of water using the joystick.

The VR sessions take place in a multisensory cabin that allows enriching the experience by potentiating the immersion and the experience of mindfulness. This cabin enables a more embodied and realistic experience by generating sounds, smells, air and heat (figure 2). This device (the Sensory reality pod) was designed by Sensiks as a modular framework that includes electronic modules and programmable actuators to generate the defined stimulations at the desired times. The modules and actuators are linked through a central device. The VR cabin measures 119×119×224 cm. This tool provides a multisensory experience that adapts to the VE presented. The appropriate smells (notably forest, beach, alcohol, tobacco, coffee, gasoline), sounds, airflow and heat are programmed for each VE and evolve according to the exploration of the environment or as the session progresses (eg, when moving in the sun in the VE, heaters are switched on).

For each of the six sessions, patients complete 20 min of guided mindfulness while exploring a VE. The first two sessions aim to teach participants the practice of mindfulness through immersion in relaxing environments (VEs 1 and 2). Next, to train the patients to better control craving induced by cues, they are gradually exposed to VEs containing cues meant to arouse the desire to consume alcohol (VEs 3 and 4). The aim is to get used to being confronted with cues without it precipitating consumption by learning to regulate cue-induced craving through mindfulness. Finally, to train the patients to regulate stress, a powerful inducer of craving and relapse,¹² they are gradually exposed to environments that can induce stress (VEs 5 and 6). Mindfulness instructions guide the patient to regulate stress and stress-induced craving. The goal of this stress induction is for patients to learn to better regulate their stress in daily life, thus reducing their risk of relapse. During VR mindfulness sessions, patients' cardiac and respiratory activity is monitored using a breathing belt and a heartbeat ear clip.

Performing cognitive exercises while cycling

The second tool used for this intervention is the cognitive bike (Vélo-cognitif, figure 3), designed by RevLim




Figure 2 Sensory reality pod: Device created by Sensiks containing programmable actuators to generate different types of stimulation using heaters, an odour diffusion module connected to bottles containing fragrances, an audio system and fans. The authors (including the person pictured in the figure) declare that this photograph was illustrated by one of the coauthors, and grant permission and give their consent to BMJ open for the use of this photograph for publication, including print or web-based publications. The authors understand that with their authorisation the photograph can never be revoked.

to practise a cognitive and physical dual task. This tool combines an exercise bike and a touch pad offering cognitive training games. The bike has different levels of resistance. The cognitive exercises are designed by HappyNeuron, a network of scientific experts specialising in cognitive training. These exercises include training in inhibition, attention, memory and visuo-spatial skills through playful games of various difficulty levels. By simultaneously stimulating motor skills and cognition, this tool offers a playful activity in which the participants are able to see themselves progressing and which could reduce



Figure 3 Vélo-cognitif (cognitive bike) consisting of an exercise bike to which a digital touch pad is attached to allow cognitive exercises to be performed while pedalling. The authors (including the person pictured in the figure) declare that this photograph was illustrated by one of the coauthors, and grant permission and give their consent to BMJ open for the use of this photograph for publication, including print or web-based publications. The authors understand that with their authorisation the photograph can never be revoked.

stress through physical activity.⁶³ In a pilot study using the cognitive bike with patients suffering from AUD, most patients reported having enjoyed the activity.⁶⁴ This tool could therefore help improve well-being and quality of life by providing a rewarding physical activity while exercising executive functions.

Control group

Patients in the control group are treated according to the standard protocol used at the Laborit Psychiatric Hospital. This consists of a clinical and biological examination performed by a health professional following the patient's admission to hospital. A treatment for withdrawal symptoms (benzodiazepines) is then administered under medical supervision. The patient benefits from individual medical and psychological monitoring. Workshops and therapeutic meetings are offered.

Measures

Primary outcome

The primary outcome is the proportion of individuals who relapsed 2 weeks after treatment. Relapse has been defined as drinking at least five drinks per occasion or drinking at least five times a week.^{65 66} Maintenance of abstinence and relapse are usually assessed using

the Alcohol Timeline Followback (TLFB) assessment method.⁶⁷ However, this method relies on verbal reports and may thus be sensitive to under-reporting. Relapse is also assessed using biological indicators—a breathalyser, and a blood test for the measurements of carbohydrate deficient transferrin (CDT) and gamma-glutamyl transpeptidase (GGT). CDT is a glycoprotein synthesised by the liver and is a specific marker indicating the presence of alcohol in the blood. Elevated CDT levels suggest recent high alcohol consumption, particularly if other liver-associated enzymes (such as GGT) are elevated.⁶⁸ These measures will be used as combined indicators of relapse in order to corroborate the patient's report and to objectify the relapse, to avoid a possible reporting bias. Indeed, underestimation of consumption is frequent, and may be related to low insight or social desirability bias,^{62 69 70} hence it is of interest to combine subjective with objective indicators of relapse. In this clinical trial, we will consider that a relapse occurred if at least one of the three indicators points to a relapse: (1) if in the TLFB the patient indicates consumption at least five times a week or at least five drinks per occasion; (2) if there is a significant increase in CDT and GGT since the day 10 blood test; or (3) if the breathalyser is positive. We will consider patients as non-relapsers if none of these indicators (TLFB, CDT and GGT, breathalyser) are positive.

Secondary outcomes

1. Mid-term relapse

Relapse is assessed using the same measures at 1 month and 3 months.

2. Craving

- Explicit craving, defined as the conscious desire to consume alcohol, is assessed in two ways:
 - o Subjective craving during the past week is assessed using the Obsessive Compulsive Drinking Scale questionnaire,⁷¹ which measures an individual's alcohol consumption and attempts to control it over the past week.
 - o Craving induced by cues is assessed using the craving induction protocol of Fox *et al.*⁷² This protocol induces craving through personalised cues, which consist of a short text written by the patient. This text should describe a memory of a situation of strong craving that led to consumption and should detail the context, the physical sensations and the state of mind during that scene. This half-page script is recorded by the experimenter, and then the audio is played to the patient during a second session. The patient indicates his level of craving before and after listening to the script on a Visual Analogue Scale ranging from 0 to 10, where 0 represents no desire to consume and 10 represents an extremely strong craving.

- To obtain a behavioural assessment in relation to the automatic component of craving and drug-seeking behaviour, we also use three implicit measures (that will be combined into a single score of implicit craving):
 - o Identification with the drug is assessed using a standard Implicit Association Test.⁷³ This test measures whether alcohol is more strongly associated with the self or others using images referring to alcohol or neutral images as target categories and words referring to the self or others as attribute categories (eg: 'me', 'I', 'myself', 'mine' or 'they', 'them', 'their', 'others').⁷³
 - o A test of the seeking for alcohol-related stimuli based on the probabilistic image choice task of Moeller *et al.*⁷⁴ and adapted to alcohol is used. This test allows evaluating the preference for the drug among other reinforcers via four categories of images (drug, pleasant, unpleasant and neutral). There is a choice of four decks of cards, face down, each containing a majority of one of the image categories. A large image of the selected deck is presented on the screen for 2000 milliseconds, and then the subjects can select one of the decks again. A pseudo-randomisation described by Moeller *et al.*⁷⁵ aims at reducing awareness of the identity of the deck while allowing a preference to be established. We adapted this task to alcohol addiction using images related to alcohol consumption (a glass of beer, a glass of wine, people drinking in a bar, etc).
 - o Attentional bias towards alcohol is assessed using the task of Soleymani *et al.*⁷⁶ a visual research test for alcohol-related stimuli (eye tracking). This task consists of freely viewing several sets of 16 images of alcoholic and non-alcoholic drinks, while an eye-tracker records the location of the first fixation and the total fixation time for each image. The strength of the attentional bias is determined by these two parameters.
- 3. Mindfulness
 - Mindfulness skills acquisition is measured using the Five Facets Mindfulness questionnaire.^{77 78} This questionnaire assesses the tendency to be in a state of mindfulness in daily life based on five facets of mindfulness: observation (being attentive to one's internal/external states), description (being able to verbalise one's internal experiences), action with awareness (not automatic), non-judgment of internal and external experience (posture of acceptance, non-evaluative) and non-reactivity (detachment, non-response to internal states).^{79 80} We use the 15-item version of the questionnaire.
 - The development of mindfulness skills is also assessed during mindfulness sessions using heart rate, respiratory rate and salivary cortisol measurements. These measures allow us to monitor the patient's attention to instructions, the stress induced by the en-

Table 1 Study schedule of enrolment and assessments by time points (Tn)

	Enrolment	Enhanced intervention	Post-intervention	Follow-ups		
Time point	T0	T1	T2	T3	T4	T5
Eligibility screening	X					
Informed consent	X					
Group allocation	X					
Primary outcome						
TLFB				X	X	X
Breathalyser				X	X	X
Blood test			X	X	X	X
Secondary outcomes						
OCDS	X			X	X	X
Craving induced by cues	X		X			
IAT	X		X			
Alcohol seeking	X		X			
Alcohol attentional bias	X		X			
MPSEQ	X			X	X	X
FFMQ	X	X	X			
Salivary cortisol	X	X				

FFMQ, Five Facets Mindfulness Questionnaire; IAT, Implicit Association Test; MPSEQ, Measurement of the Perception of a Stimulating Environment Questionnaire; OCDS, Obsessive Compulsive Drinking Scale; TLFB, Alcohol Timeline Followback .

vironments and whether mindfulness can effectively regulate this stress. These measurements also allow the calculation of heart rate variability associated with self-regulation skills, which is commonly used in research on mindfulness-based interventions.^{81–83}

4. Richness of daily environment

We evaluate the effect of the intervention on the perceived richness of the daily environment. The perceived richness of the daily environment is assessed using the Measurement of the Perception of a Stimulating Environment Questionnaire (MPSEQ) (Chatard A, Barillot L, Besnier M, *et al.* Measurement of the Perception of a Stimulating Environment Questionnaire, unpublished, online supplemental file 2). Composed of 13 items, this questionnaire evaluates to what extent the individual perceives their environment as stimulating. Each item is a declarative statement referring to the stimulations, activities or satisfaction and entertainment that the person perceives or realises in their life (eg: ‘My immediate environment is rich in sensations and stimulation of all kinds’). The individual expresses the extent to which they agree or disagree with each statement using a scale ranging from 1 to 7, with 1 corresponding to completely disagree and 7 to completely agree.

Study schedule

Summary [table 1](#) (online supplemental file).

Potential participants are invited to a 15 min informative meeting. Eligibility screening is conducted, and patients are invited to participate after being provided a description of the study. A pretest in the VR cabin is

offered to the patients to allow them to get acquainted with VR and to check that they do not show signs of cybersickness.

Patients are reconvened (T0), and if they wish to participate in the study and if they meet all the eligibility criteria they sign the consent form and are randomly assigned to one of two groups. They complete baseline measures of their craving and salivary cortisol and complete questionnaires about their initial mindfulness skills and their perception of the richness of their daily environment (online supplemental table 1).

Intermediate measures take place for participants undergoing the intervention enhancement (T1). The acquisition of mindfulness skills is assessed after the second session of intervention enhancement. Salivary cortisol is measured after session 5 and session 6, which are expected to induce stress in the patient.

On the 10th day of inclusion or after completing the enhanced intervention (depending on the group) (T2), participants are seen again for a measurement session during which craving and mindfulness skills are assessed. A blood test is performed to obtain a baseline measurement of CDT and GGT. Relapse is assessed in both groups according to the method described in the measures section at 2 weeks (T3), 1 month (T4) and 3 months (T5) after T2.

Sample size

The required sample size was computed to have 80% statistical power to detect a significant reduction of relapse rate of at least 25% on our primary outcome (with

$p < 0.05$, two-tailed). We considered a 25% reduction in relapse rate as the smallest effect size of interest in our study. This effect size seems realistic to us and appears to be the smallest effect size of clinical interest given the investment of material and human time in this intervention. The smallest effect size of interest is the smallest effect that (1) researchers personally care about, (2) is theoretically interesting or (3) has practical relevance.⁸⁴ The relapse rate on our site is about 50% on average after 1 month.⁶² This is consistent with other studies in which nearly half of the patients relapsed in the weeks following hospital discharge for alcohol withdrawal.^{85 86} With a relapse rate of 50% after 1 month, there is significant potential to reduce the relapse rate by 25%. It is quite possible that the intervention will have a much smaller effect size. However, if the relapse rate is not reduced by at least 25%, the intervention would be considered to have no efficacy or too low to be of clinical relevance.

The a priori power analysis for sample size indicates that the required sample size is 116 for our primary outcome. However, we will include 135 patients to compensate for possible missing data. For the secondary outcomes (craving, mindfulness skills, perceived richness of the daily environment) this sample size ($N=135$) provides adequate power ($1-\beta=0.82$) to detect a medium effect size or a larger effect size (Cohen's $d > 0.50$, with $p < 0.05$ (two-sided)).

Data analysis

Data analysts will be blind to the condition: the condition will be letter coded rather than explicitly 'control' or 'intervention'. All analyses will be conducted using both RStudio and SPSS V.23.0 software. A descriptive analysis of the study population will be performed. Qualitative variables will be expressed as a proportion with 95% CI. Quantitative variables will be expressed as mean and SD or as median and IQR. A value of $p < 0.05$ (two-tailed) will be considered statistically significant. In intention-to-treat analyses, missing data will be simulated using a multiple-imputation technique with interim values.

The initial comparability resulting from the randomisation will be checked using tests appropriate to the distribution (parametric or non-parametric) and type (quantitative/qualitative) of the variables studied.

The primary outcome (relapse at 2 weeks) will be analysed using an independent sample proportion test (χ^2) comparing the differences according to the groups with a two-tailed alpha risk of 5%. The difference between the two groups for relapse at 1 month and 3 months (secondary outcomes) will be analysed in the same way. An independent sample t-test (Student's t-test) will be used on the other secondary outcomes (craving, mindfulness skills and perceived richness of the daily environment) to compare the differences before and after the enhanced intervention, and before and during the enhanced intervention for the intervention enhancement group (within-participant), with a two-tailed alpha risk of 5%.

ETHICS AND DISSEMINATION

All participants have to give written informed consent to the investigator. This study is approved by the Ethics Committee Nord Ouest IV of Lille (reference number 2022-A01156-37). Results will be disseminated through presentations, peer-reviewed journals and seminar conferences. All information on ethical considerations and open science practices can be accessed at <https://osf.io/b57uj/>.

PATIENT AND PUBLIC INVOLVEMENT

Patients or public were not involved in the design or conduct of the study.

DISCUSSION

This study aims at evaluating the effectiveness of exposure to EE sessions in reducing relapse in patients receiving treatment for AUD. This study is the first attempt to transpose the EE approach described in preclinical studies²⁴ combining different types of stimulation to humans. Our study should inform about the potential of this strategy to treat addiction by promoting long-term abstinence and reducing the incidence of relapse. Our EE provides rewarding stimulations as alternatives to drug taking while training the cognitive and emotional processes deregulated in AUD,⁶⁴ and provide skills that can be reused in daily life in stressful or cue-exposure situations to resist craving and avoid relapse.¹⁸

In terms of study limitations, it is possible that the acceptability of VR and mindfulness⁸⁷⁻⁹⁰ can play a role in the effectiveness of the intervention enhancement. We have chosen to exclude people who show a cybersickness type of discomfort in the VR. Regarding mindfulness, repeated measures of mindfulness skills should help to control for this acceptability bias. Another limitation is that we chose to induce stress through VR during the last two EE sessions, even though stress is supposed to be a sort of functional opposite of EE and can therefore interfere with the enrichment procedure.^{24 25} However, the sessions may be considered to be part of EE because they provide cognitive training to learn to regulate stress through mindfulness. Another possible weakness of the EE in this study is that it does not include additional social stimulation compared with standard treatment, while several animal studies have shown that social stimulation is an important component of EE,²⁴ and social support in humans is an important part of addiction treatment.¹⁸ Therefore, further studies are needed to investigate whether a protocol of EE that includes more specific forms of social stimulation would be more effective than the present one. Finally, it will be important in the future to perform parametric studies to investigate whether more or fewer EE sessions can increase or decrease the benefits of this EE protocol.

Contributors AC, NJ and MS were responsible for identifying the research question and designing the study protocol. MB, LVB and CC contributed to the

development of the protocol and study design. LVB wrote the first draft of this manuscript, which was revised and modified by all authors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Lila Barillot <http://orcid.org/0000-0002-4327-1308>

REFERENCES

- Ritchie H, Roser M. Alcohol consumption [Our World Data]. 2018. Available: <https://ourworldindata.org/alcohol-consumption>
- World Health Organization. Global status report on alcohol and health 2018. 2019.
- Batra A, Müller CA, Mann K, et al. Alcohol dependence and harmful use of alcohol. *Dtsch Arztebl Int* 2016;113:arztebl.2016.0301:301–10..
- Hunt WA, Barnett LW, Branch LG. Relapse rates in addiction programs. *J Clin Psychol* 1971;27:455–6. 10.1002/1097-4679(197110)27:4<455::aid-jclp2270270412>3.0.co;2-r Available: [http://dx.doi.org/10.1002/1097-4679\(197110\)27:4%3C455::AID-JCLP2270270412%3E3.0.CO;2-R](http://dx.doi.org/10.1002/1097-4679(197110)27:4%3C455::AID-JCLP2270270412%3E3.0.CO;2-R)
- Manning V, Staiger PK, Hall K, et al. Cognitive bias modification training during inpatient alcohol detoxification reduces early relapse: a randomized controlled trial. *Alcohol Clin Exp Res* 2016;40:2011–9. 10.1111/acer.13163 Available: <http://dx.doi.org/10.1111/acer.13163>
- Kharb R, Shekhawat LS, Beniwal RP, et al. Relationship between craving and early relapse in alcohol dependence: a short-term follow-up study. *Indian J Psychol Med* 2018;40:315–21. 10.4103/IJPSYM.IJPSYM_558_17 Available: http://dx.doi.org/10.4103/IJPSYM.IJPSYM_558_17
- Stohs ME, Schneekloth TD, Geske JR, et al. Alcohol craving predicts relapse after residential addiction treatment. *Alcohol and Alcoholism* 2019;54:167–72. 10.1093/alcalc/agy093 Available: <http://dx.doi.org/10.1093/alcalc/agy093>
- Sliedrecht W, Seesink HJ, Vrijmoeth C, et al. Alcohol use disorder relapse factors: an exploratory investigation of craving, alcohol dependence severity, and meaning in life. *Addiction Research & Theory* 2022;30:351–9. 10.1080/16066359.2022.2040488 Available: <http://dx.doi.org/10.1080/16066359.2022.2040488>
- Vafaie N, Kober H. Association of drug cues and craving with drug use and relapse: a systematic review and meta-analysis. *JAMA Psychiatry* 2022;79:641–50. 10.1001/jamapsychiatry.2022.1240 Available: <http://dx.doi.org/10.1001/jamapsychiatry.2022.1240>
- Pohorecky LA. Stress and alcohol interaction: an update of human research. *Alcohol Clin Exp Res* 1991;15:438–59. 10.1111/j.1530-0277.1991.tb00543.x Available: <http://dx.doi.org/10.1111/j.1530-0277.1991.tb00543.x>
- Self DW. Neural substrates of drug craving and relapse in drug addiction. *Ann Med* 1998;30:379–89. 10.3109/07853899809029938 Available: <http://dx.doi.org/10.3109/07853899809029938>
- Sinha R, Li CSR. Imaging stress- and cue-induced drug and alcohol craving: association with relapse and clinical implications. *Drug Alcohol Rev* 2007;26:25–31. 10.1080/09595230601036960 Available: <http://dx.doi.org/10.1080/09595230601036960>
- Sinha R. How does stress increase risk of drug abuse and relapse? *Psychopharmacology (Berl)* 2001;158:343–59. 10.1007/s002130100917 Available: <http://dx.doi.org/10.1007/s002130100917>
- Solinas M, Chauvet C, Thiriet N, et al. Reversal of cocaine addiction by environmental enrichment. *Proc Natl Acad Sci USA* 2008;105:17145–50. 10.1073/pnas.0806889105 Available: <http://dx.doi.org/10.1073/pnas.0806889105>
- Solinas M, Thiriet N, Rawas RE, et al. Environmental enrichment during early stages of life reduces the behavioral, neurochemical, and molecular effects of cocaine. *Neuropsychopharmacol* 2009;34:1102–11. 10.1038/npp.2008.51 Available: <http://dx.doi.org/10.1038/npp.2008.51>
- Stairs DJ, Bardo MT. Neurobehavioral effects of environmental enrichment and drug abuse vulnerability. *Pharmacol Biochem Behav* 2009;92:377–82. 10.1016/j.pbb.2009.01.016 Available: <http://dx.doi.org/10.1016/j.pbb.2009.01.016>
- Chauvet C, Goldberg SR, Jaber M, et al. Effects of environmental enrichment on the incubation of cocaine craving. *Neuropharmacology* 2012;63:635–41. 10.1016/j.neuropharm.2012.05.014 Available: <http://dx.doi.org/10.1016/j.neuropharm.2012.05.014>
- Galaj E, Barrera ED, Ranaldi R. Therapeutic efficacy of environmental enrichment for substance use disorders. *Pharmacol Biochem Behav* 2020;188:172829.
- Solinas M, Chauvet C, Lafay-Chebassier C, et al. Environmental enrichment-inspired pharmacological tools for the treatment of addiction. *Curr Opin Pharmacol* 2021;56:22–8.
- Enoch MA. Genetic and environmental influences on the development of alcoholism: resilience vs. risk. *Ann N Y Acad Sci* 2006;1094:193–201. 10.1196/annals.1376.019 Available: <http://dx.doi.org/10.1196/annals.1376.019>
- Ewald DR, Strack RW, Orsini MM. Rethinking addiction. *Glob Pediatr Health* 2019;6:2333794X18821943. 10.1177/2333794X18821943 Available: <http://dx.doi.org/10.1177/2333794X18821943>
- Sikora M, Nicolas C, Istini M, et al. Generalization of effects of environmental enrichment on seeking for different classes of drugs of abuse. *Behav Brain Res* 2018;341:109–13.
- Li X, Meng L, Huang K, et al. Environmental enrichment blocks reinstatement of ethanol-induced conditioned place preference in mice. *Neuroscience Letters* 2015;599:92–6. 10.1016/j.neulet.2015.05.035 Available: <http://dx.doi.org/10.1016/j.neulet.2015.05.035>
- Solinas M, Thiriet N, Chauvet C, et al. Prevention and treatment of drug addiction by environmental enrichment. *Prog Neurobiol* 2010;92:572–92.
- Crofton EJ, Zhang Y, Green TA. Inoculation stress hypothesis of environmental enrichment. *Neuroscience & Biobehavioral Reviews* 2015;49:19–31.
- Sampedro-Piquero P, Ladrón de Guevara-Miranda D, Pavón FJ, et al. Neuroplastic and cognitive impairment in substance use disorders: a therapeutic potential of cognitive stimulation. *Neuroscience & Biobehavioral Reviews* 2019;106:23–48. 10.1016/j.neubiorev.2018.11.015 Available: <http://dx.doi.org/10.1016/j.neubiorev.2018.11.015>
- Rawson RA, Chudzynski J, Mooney L, et al. Impact of an exercise intervention on methamphetamine use outcomes post-residential treatment care. *Drug Alcohol Depend* 2015;156:21–8.
- Georgakouli K, Manthou E, Georgoulas P, et al. Exercise training reduces alcohol consumption but does not affect HPA-axis activity in heavy drinkers. *Physiology & Behavior* 2017;179:276–83. 10.1016/j.physbeh.2017.07.003 Available: <http://dx.doi.org/10.1016/j.physbeh.2017.07.003>
- Elibero A, Janse Van Rensburg K, Drobos DJ. Acute effects of aerobic exercise and hatha yoga on craving to smoke. *Nicotine & Tobacco Research* 2011;13:1140–8. 10.1093/ntr/ntr163 Available: <http://dx.doi.org/10.1093/ntr/ntr163>
- Williams DM, Dunsiger S, Whiteley JA, et al. Acute effects of moderate intensity aerobic exercise on affective withdrawal symptoms and cravings among women smokers. *Addict Behav* 2011;36:894–7. 10.1016/j.addbeh.2011.04.001 Available: <http://dx.doi.org/10.1016/j.addbeh.2011.04.001>
- Kurti AN, Dallery J. Effects of exercise on craving and cigarette smoking in the human laboratory. *Addict Behav* 2014;39:1131–7.
- Prapavessis H, De Jesus S, Fitzgeorge L, et al. Exercise to enhance smoking cessation: the getting physical on cigarette randomized control trial. *Ann Behav Med* 2016;50:358–69. 10.1007/s12160-015-9761-9 Available: <http://dx.doi.org/10.1007/s12160-015-9761-9>
- Giménez-Meseguer J, Tortosa-Martínez J, Cortell-Tormo JM. The benefits of physical exercise on mental disorders and quality of life in substance use disorders patients. systematic review and meta-

- analysis. *Int J Environ Res Public Health* 2020;17:3680. 10.3390/ijerph17103680 Available: <https://doi.org/10.3390/ijerph17103680>
- 34 Czuchry M, Dansereau DF. Cognitive skills training: impact on drug abuse counseling and readiness for treatment. *Am J Drug Alcohol Abuse* 2003;29:1–18. 10.1081/ada-120018837 Available: <http://dx.doi.org/10.1081/ADA-120018837>
- 35 Rupp CI, Kemmler G, Kurz M, et al. Cognitive remediation therapy during treatment for alcohol dependence. *J Stud Alcohol Drugs* 2012;73:625–34. 10.15288/jsad.2012.73.625 Available: <http://dx.doi.org/10.15288/jsad.2012.73.625>
- 36 Rass O, Schacht RL, Buckheit K, et al. A randomized controlled trial of the effects of working memory training in methadone maintenance patients. *Drug Alcohol Depend* 2015;156:38–46.
- 37 Rezapour T, Hatami J, Farhoudian A, et al. Cognitive rehabilitation for individuals with opioid use disorder: a randomized controlled trial *Neuropsychol Rehabil* 2019;29:1273–89. 10.1080/09602011.2017.1391103 Available: <http://dx.doi.org/10.1080/09602011.2017.1391103>
- 38 Garland EL, Froeliger B, Howard MO. Mindfulness training targets neurocognitive mechanisms of addiction at the attention-appraisal-emotion interface. *Front Psychiatry* 2014;4:173.
- 39 Garland EL, Gaylord SA, Boettiger CA, et al. Mindfulness training modifies cognitive, affective, and physiological mechanisms implicated in alcohol dependence: results of a randomized controlled pilot trial. *J Psychoactive Drugs* 2010;42:177–92. 10.1080/02791072.2010.10400690 Available: <http://dx.doi.org/10.1080/02791072.2010.10400690>
- 40 Morton ML, Helminen EC, Felver JC. A systematic review of mindfulness interventions on psychophysiological responses to acute stress. *Mindfulness* 2020;11:2039–54.
- 41 Mohan A, Sharma R, Bijlani RL. Effect of meditation on stress-induced changes in cognitive functions. *J Altern Complement Med* 2011;17:207–12.
- 42 Bishop SR. What do we really know about mindfulness-based stress reduction? *Psychosom Med* 2002;64:71–83.
- 43 Creswell JD. Mindfulness interventions. *Annu Rev Psychol* 2017;68:491–516. 10.1146/annurev-psych-042716-051139 Available: <http://dx.doi.org/10.1146/annurev-psych-042716-051139>
- 44 Hsu SH, Grow J, Marlatt GA. Mindfulness and addiction. *Recent Dev Alcohol* 2008;18:229–50. 10.1007/978-0-387-77725-2_13 Available: http://dx.doi.org/10.1007/978-0-387-77725-2_13
- 45 Audiffren M, André N, Baumeister RF. Training willpower: reducing costs and valuing effort. *Front Neurosci* 2022;16:699817.
- 46 Lei C, Sunzi K, Dai F, et al. Effects of virtual reality rehabilitation training on gait and balance in patients with Parkinson's disease: a systematic review. *PLoS One* 2019;14:e0224819.
- 47 Riaz W, Khan ZY, Jawaid A, et al. Virtual reality (VR) -based environmental enrichment in older adults with mild cognitive impairment (MCI) and mild dementia. *Brain Sci* 2021;11:1103. 10.3390/brainsci11081103 Available: <http://dx.doi.org/10.3390/brainsci11081103>
- 48 Riva G, Serino S. Virtual reality in the assessment, understanding and treatment of mental health disorders. *JCM* 2020;9:3434.
- 49 Giovancarli C, Malbos E, Baumstarck K, et al. Virtual reality cue exposure for the relapse prevention of tobacco consumption: a study protocol for a randomized controlled trial. *Trials* 2016;17:96. 10.1186/s13063-016-1224-5 Available: <http://dx.doi.org/10.1186/s13063-016-1224-5>
- 50 Lebiecka Z, Skoneczny T, Tyburski E, et al. Is virtual reality cue exposure a promising adjunctive treatment for alcohol use disorder? *JCM* 2021;10:2972. 10.3390/jcm10132972 Available: <http://dx.doi.org/10.3390/jcm10132972>
- 51 Bordnick PS, Traylor A, Copp HL, et al. Assessing reactivity to virtual reality alcohol based cues. *Addict Behav* 2008;33:743–56.
- 52 Ghiță A, Hernández-Serrano O, Fernández-Ruiz Y, et al. Cue-elicited anxiety and alcohol craving as indicators of the validity of ALCO-VR software: a virtual reality study. *J Clin Med* 2019;8:1153.
- 53 Ryan JJ, Kreiner DS, Chapman MD, et al. Virtual reality cues for binge drinking in college students. *CyberPsychology & Behavior* 2009;091110182143098.
- 54 Segawa T, Baudry T, Bourla A, et al. Virtual reality (VR) in assessment and treatment of addictive disorders: a systematic review. *Front Neurosci* 2019;13:1409. 10.3389/fnins.2019.01409 Available: <http://dx.doi.org/10.3389/fnins.2019.01409>
- 55 Mazza M, Kammler-Sücker K, Leménager T, et al. Virtual reality: a powerful technology to provide novel insight into treatment mechanisms of addiction. *Transl Psychiatry* 2021;11:817.
- 56 Zimmer P, Buttlar B, Halbeisen G, et al. Virtually stressed? A refined virtual reality adaptation of the trier social stress test (tsst) induces robust endocrine responses. *Psychoneuroendocrinology* 2019;101:186–92.
- 57 Martens MA, Antley A, Freeman D, et al. It feels real: physiological responses to a stressful virtual reality environment and its impact on working memory. *J Psychopharmacol* 2019;33:1264–73. 10.1177/0269881119860156 Available: <http://dx.doi.org/10.1177/0269881119860156>
- 58 Linehan M. *DBT Skills Training Manual*. New York: Guilford Publications, 2014.
- 59 Grant BF, Stinson FS, Dawson DA, et al. Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders. *Arch Gen Psychiatry* 2004;61:807.
- 60 Boden JM, Fergusson DM. Alcohol and depression. *Addiction* 2011;106:906–14.
- 61 Navarro-Haro MV, Modrego-Alarcón M, Hoffman HG, et al. Evaluation of a mindfulness-based intervention with and without virtual reality dialectical behavior therapy® mindfulness skills training for the treatment of generalized anxiety disorder in primary care: a pilot study. *Front Psychol* 2019;10:55. 10.3389/fpsyg.2019.00055 Available: <http://dx.doi.org/10.3389/fpsyg.2019.00055>
- 62 Dandaba M, Serra W, Harika-Germaneau G, et al. Predicting relapse in patients with severe alcohol use disorder: the role of alcohol insight and implicit alcohol associations. *Addict Behav* 2020;107:106433.
- 63 Edenfield TM, Blumenthal JA. Exercise and stress reduction. In: Contrada R, Baum A, eds. *The Handbook of Stress Science: Biology, Psychology, and Health*. New York: Springer Publishing Company, 2011: 301–19.
- 64 Chauvet C. Stratégies environnementales et pharmacologiques Du traitement de la rechute à la cocaïne: Du comportement à la neurobiologie. 2011.
- 65 O'Malley SS, Jaffe AJ, Chang G, et al. Naltrexone and coping skills therapy for alcohol dependence. A controlled study. *Arch Gen Psychiatry* 1992;49:881–7. 10.1001/archpsyc.1992.01820110045007 Available: <http://dx.doi.org/10.1001/archpsyc.1992.01820110045007>
- 66 Volpicelli JR, Alterman AI, Hayashida M, et al. Naltrexone in the treatment of alcohol dependence. *Arch Gen Psychiatry* 1992;49:876–80. 10.1001/archpsyc.1992.01820110040006 Available: <http://dx.doi.org/10.1001/archpsyc.1992.01820110040006>
- 67 Litten RZ, Allen JP. Measuring alcohol consumption. Totowa, NJ, 1992.
- 68 Allen JP, Anton R. Biomarkers as aids to identification of relapse in alcoholic patients. In: *In Recent Developments in Alcoholism 2002*. Boston, MA: Springer 25–38.
- 69 Goldstein RZ, Craig ADB, Bechara A, et al. The neurocircuitry of impaired insight in drug addiction. *Trends Cogn Sci* 2009;13:372–80.
- 70 Kim JS, Park BK, Kim GJ, et al. The role of alcoholics' insight in abstinence from alcohol in male Korean alcohol dependents. *J Korean Med Sci* 2007;22:132–7.
- 71 Anton RF, Moak DH, Latham P. The obsessive compulsive drinking scale: a self-rated instrument for the quantification of thoughts about alcohol and drinking behavior. *Alcoholism Clin Exp Res* 1995;19:92–9. 10.1111/j.1530-0277.1995.tb01475.x Available: <http://www.blackwell-synergy.com/toc/acer/19/1>
- 72 Fox HC, Bergquist KL, Hong K-I, et al. Stress-Induced and alcohol cue-induced craving in recently abstinent alcohol-dependent individuals. *Alcohol Clin Exp Res* 2007;31:395–403. 10.1111/j.1530-0277.2006.00320.x Available: <https://doi.org/10.1111/j.1530-0277.2006.00320.x>
- 73 Lindgren KP, Neighbors C, Teachman BA, et al. Implicit alcohol associations, especially drinking identity, predict drinking over time. *Health Psychol* 2016;35:908–18. 10.1037/hea0000396 Available: <https://doi.org/10.1037/hea0000396>
- 74 Moeller SJ, Beebe-Wang N, Woicik PA, et al. Choice to view cocaine images predicts concurrent and prospective drug use in cocaine addiction. *Drug Alcohol Depend* 2013;130:178–85.
- 75 Moeller SJ, Maloney T, Parvaz MA, et al. Enhanced choice for viewing cocaine pictures in cocaine addiction. *Biol Psychiatry* 2009;66:169–76. 10.1016/j.biopsych.2009.02.015 Available: <http://dx.doi.org/10.1016/j.biopsych.2009.02.015>
- 76 Soleymani A, Ivanov Y, Mathot S, et al. Free-viewing multi-stimulus eye tracking task to index attention bias for alcohol versus soda cues: satisfactory reliability and criterion validity. *Addict Behav* 2020;100:106117.
- 77 Baer RA, Carmody J, Hunsinger M. Weekly change in mindfulness and perceived stress in a mindfulness-based stress reduction program. *J Clin Psychol* 2012;68:755–65. 10.1002/jclp.21865 Available: <http://dx.doi.org/10.1002/jclp.21865>
- 78 Baer RA, Smith GT, Hopkins J, et al. Using self-report assessment methods to explore facets of mindfulness. *Assessment* 2006;13:27–45. 10.1177/1073191105283504 Available: <https://doi.org/10.1177/1073191105283504>

- 79 Baer RA, Smith GT, Lykins E, *et al.* FFMQ-15: 15-item five-facet mindfulness questionnaire. *Assessment* 2008;15:329–42.
- 80 Heeren A, Douilliez C, Peschard V, *et al.* Cross-Cultural validity of the five facets mindfulness questionnaire: adaptation and validation in a French-speaking sample. *European Review of Applied Psychology* 2011;61:147–51. 10.1016/j.erap.2011.02.001 Available: <https://doi.org/10.1016/j.erap.2011.02.001>
- 81 Christodoulou G, Salami N, Black DS. The utility of heart rate variability in mindfulness research. *Mindfulness* 2020;11:554–70. 10.1007/s12671-019-01296-3 Available: <http://dx.doi.org/10.1007/s12671-019-01296-3>
- 82 Kirk U, Axelsen JL. Heart rate variability is enhanced during mindfulness practice: a randomized controlled trial involving a 10-day online-based mindfulness intervention. *PLoS One* 2020;15:e0243488.
- 83 Shearer A, Hunt M, Chowdhury M, *et al.* Effects of a brief mindfulness meditation intervention on student stress and heart rate variability. *International Journal of Stress Management* 2016;23:232–54. 10.1037/a0039814 Available: <http://dx.doi.org/10.1037/a0039814>
- 84 Anvari F, Lakens D. Using anchor-based methods to determine the smallest effect size of interest. *Journal of Experimental Social Psychology* 2021;96:104159.
- 85 Boothby LA, Doering PL. Acamprosate for the treatment of alcohol dependence. *Clin Ther* 2005;27:695–714. 10.1016/j.clinthera.2005.06.015 Available: <https://doi.org/10.1016/j.clinthera.2005.06.015>
- 86 Manning V, Garfield JBB, Staiger PK, *et al.* Effect of cognitive bias modification on early relapse among adults undergoing inpatient alcohol withdrawal treatment: a randomized clinical trial. *JAMA Psychiatry* 2021;78:133–40. 10.1001/jamapsychiatry.2020.3446 Available: <https://doi.org/10.1001/jamapsychiatry.2020.3446>
- 87 Bautista T. Acceptability of mindfulness-based intervention among women with substance use disorder. 2020.
- 88 Kiken LG, Garland EL, Bluth K, *et al.* From a state to a trait: trajectories of state mindfulness in meditation during intervention predict changes in trait mindfulness. *Pers Individ Dif* 2015;81:41–6.
- 89 Amaro H, Spear S, Vallejo Z, *et al.* Feasibility, acceptability, and preliminary outcomes of a mindfulness-based relapse prevention intervention for culturally-diverse, low-income women in substance use disorder treatment. *Substance Use & Misuse* 2014;49:547–59.
- 90 Warland A, Paraskevopoulos I, Tsekleves E, *et al.* The feasibility, acceptability and preliminary efficacy of a low-cost, virtual-reality based, upper-limb stroke rehabilitation device: a mixed methods study. *Disabil Rehabil* 2019;41:2119–34.

Mindfulness instructions for mindfulness practice in virtual environments (VEs)

- **VE1: Immersion in a forest**

First of all, go and discover this environment.

Your senses are a precious help. Observe the scenery of this forest, the luminosity, the smells, the possible sounds, in a global way

To immerse yourself in this place, stop for a few moments. Close your eyes. Pay attention only to the sounds that are present. From the most distant (perhaps the wind blowing in the distance) to the closest (your breathing). Observe, without judging...

Observe if this causes things to happen to you: for example, do you have any thoughts? Observe them passing and let them go as they came. Sensations (e.g. tension in the body? tingling? a feeling of lightness...)? Welcome them, without judging. Let them come and go as they came. An emotion (fear, anger, sadness, surprise, disgust, joy...)? Welcome it, without judgement. Take them into account as they are, in the present moment, without judgment. Then let them go as they came, that is, without trying to hold them back, and without trying to make them go: their departure is gradual, passive.

To finish, I suggest that you focus your attention on a specific object, for example a leaf, a mushroom, a flower... whatever feels good to you... Choose what makes you happy.

Approach this object and give it your full attention.

Without judgement, observe its colours, its shape. Does it have a scent? Does it make sounds when you bring it close to your ear? Take a few moments to observe...

When you take this time, observe if you have thoughts, sensations, emotions... Without judgement. Taste the present moment... The "here and now".

Observe if these sensations are pleasant to you, if they leave an imprint on you, a pleasant taste.

We are coming to the end of this experience, thank you for participating and see you soon for the next discovery together.

- **VE2: Walking along a beach**

First of all, discover the environment.

Your five senses are of great help. Observe the beach, the colours, the possible smells, the sounds

To immerse yourself in this place, stop for a few moments. Close your eyes. Pay attention only to the sounds that are present. From the most distant (perhaps the wind blowing in the distance, the sound of the waves) to the closest (your breathing). Observe, without judging...

Observe if this causes things to happen to you: for example, do you have thoughts? Observe them passing and let them go as they came. Sensations (e.g. tension in the body? tingling? a feeling of lightness...)? Welcome them, without judging. Let them come and go as they came. An emotion (fear, anger, sadness, surprise, disgust, joy...)? Welcome it, without judgement. Take them into account as they are, in the present moment, without judgment. Then let them go as they came, that is to say, without trying to hold them back, and without trying to make them go away: their departure is done gradually, passively.

Finally, I suggest that you focus your attention on a particular object, for example a shell, something that seems good to you... Choose what you like. Approach it and give it your full attention.

Without judgment, observe its colours, its shape. Does it have a scent? Does it make sounds when you bring it close to your ear? Take a few moments to observe...

When you take this time, observe if you have thoughts, sensations, emotions... Without judgement. Taste the present moment... The "here and now".

Observe if these sensations are pleasant to you, if they leave an imprint on you, a pleasant taste.

We are coming to the end of this experience, thank you for participating and see you soon for the next discovery together.

- **VE3: Walking down the street, buying something in a shop, then ordering a coffee in a bar**

First of all, make this new environment your own.

Your five senses are a great help. Observe your surroundings. Observe the street you are on, the landscape, the smells of the city, the sounds.

You can go into the shop near you to buy a bottle of water. Focus on the elements around you. The sounds, from the most distant (perhaps the music in the shop, conversations) to the closest (your breathing). Observe, without judging...

You are going to walk towards the exit. Once outside, you see a café in front of you. Walk slowly towards it and enter it.

Continue to focus on your sensations, the wind caressing your face, the sounds of the street. As you enter the café, observe whether this causes anything to happen to you: for example, do you have any sensations (e.g. tension in your body? tingling)?

Welcome them, without judging. Let them come and go as they came. An emotion (fear, anger, sadness, surprise, disgust, joy...)? Welcome it, without judgement. Take them into account as they are, in the present moment, without judgment. Then let them go as they came. Let them go..., without trying to hold them back, and without trying to make them go away: their departure happens gradually, passively. Let it happen...

You are going to sit at the counter and order a coffee. Focus on the smells and sounds of conversation around you. Observe if these sensations are pleasant to you, if they leave an imprint on you, a pleasant taste or not. Once again, without judging, welcoming things as they are. To finish, I suggest that you focus your attention on a particular object, for example a cup of coffee, which seems good to you... Choose what pleases you. Approach it and give it your full attention.

Without judgment, observe its colours, its shape. Does it smell? Does it make sounds when you bring it close to your ear? Take a few moments to observe...

When you take this time, observe if you have thoughts, sensations, emotions... Without judgement. Taste the present moment... The "here and now".

Observe if these sensations are pleasant to you, if they leave an imprint on you, a pleasant taste or not. Once again, without judging, welcoming things as they are.

We are coming to the end of this experience, thank you for participating and see you soon for the next discovery together.

- **VE4: Party in a house with other people who smoke and drink alcohol**

First of all, get used to this new environment.

Your five senses are of great help. Observe your surroundings. Observe the house you are in, the rooms, the smells in the house, the sounds.

Walk around the house and focus on the elements around you. The sounds, from the most distant (perhaps the music in the house, conversations) to the closest (your breathing). Observe, without judging...

Let any sensations, thoughts or emotions within you quietly and passively subside. You let it happen... Without trying to change your breathing, without trying to hold anything back... You let the sensations that may be present happen, until they disappear by themselves. You let it happen.

Go to the table and sit down with the other people. Observe if this causes things to happen in you: for example, do you have any sensations (e.g. tension in the body? tingling?)? Welcome them, without judging. Let them come and go as they came. An emotion (fear, anger, sadness, surprise, disgust, joy...)? Welcome it, without judgement. Take them into account as they are, in the present moment, without judgment. Then let them go as they came. Let them go..., without trying to hold them

back, and without trying to make them go away: their departure happens gradually, passively. Let it happen...

You get up and go to the kitchen. Look in the fridge for a bottle of water. Focus on the smells and sounds of conversation around you. Observe if these sensations are pleasant to you, if they leave an imprint on you, a pleasant taste.

Go back to your seat and continue to focus on the sounds and smells. Observe whether this causes you to do something: for example, do you have any sensations (e.g. tension in the body? tingling?)? Welcome them, without judging. Let them come and go as they came. An emotion (fear, anger, sadness, surprise, disgust, joy...)? Accept it, without judgement. Take them into account as they are, in the present moment, without judgment. Then let them go as they came. Let them go..., without trying to hold them back, and without trying to make them go away: their departure happens gradually, passively. Let it happen...

To finish, I suggest that you focus your attention on a particular object, for example a plate, a flower, something that seems good to you... Choose what you like. Approach it and focus your attention on this object.

Without judgment, observe its colours, its shape. Does it have a scent? Does it make sounds when you bring it close to your ear? Take a few moments to observe... When you take this time, observe if you have thoughts, sensations, emotions... Without judgement. Taste the present moment... The "here and now".

Observe if these sensations are pleasant to you, if they leave an imprint on you, a pleasant taste or not. Once again, without judging, welcoming things as they are.

We are coming to the end of this experience, thank you for participating and see you soon for the next discovery together.

- **VE5: A parachute jump**

First of all, get used to this new environment.

Your five senses are a great help. Observe your environment, around you. Observe the cabin you are in, the landscape, the sounds.

Before approaching the edge, you can focus on yourself. Close your eyes. Pay attention only to the sounds that are present. From the most distant (perhaps the wind blowing in the distance) to the closest (your breathing). Observe, without judging...

The jumping will now begin. Try to keep the calmness you have achieved. Let the sensations come to you without trying to anticipate them.

Continue to focus on your sensations, the wind caressing your face... Observe if this causes things to happen to you: for example, do you have any sensations (e.g. tension in your body? tingling?)? Welcome them, without judging. Let them come and go as

they came. An emotion (fear, anger, sadness, surprise, disgust, joy...)? Welcome it, without judgement. Take them into account as they are, in the present moment, without judgment. Then let them go as they came. Let them go..., without trying to hold them back, and without trying to make them go away: their departure happens gradually, passively. Let it happen...

Perhaps you notice an increase in your heart rate, your heart beating faster..., or tensions in your body, muscles that stiffen...

You let it happen, without judgment. Let it happen, without trying to control anything. Let these sensations pass, the emotions that come, as they have come. Let it happen, passively, so that they leave quietly as they came. You let it happen..., without trying to modify your breathing. You let it happen, in confidence, quietly.

Continue to observe and feel your environment. Try to perceive again the elements around you, the noises, perhaps the wind

Let all sensations, thoughts and emotions in you quietly and passively subside. You let it happen... Without trying to change your breathing, without trying to hold anything inside you... You let the sensations that may be present happen, until they disappear by themselves. You let it happen. The parachute will now open. In the same way, observe if these sensations are pleasant for you, if they leave an imprint in you, a pleasant taste or not. Once again, without judging, welcoming things as they are.

We are coming to the end of this experience, thank you for participating and see you soon for the next discovery together.

- **VE6: Flying with turbulence**

First of all, get used to this new environment.

Your five senses are of great help. Observe your surroundings. Observe the cabin you are in, the landscape through the window, the smells, the sounds. Take time to explore the cabin. As the captain announces, turbulence will start in the plane. You can return to your seat. Close your eyes and let the sensations come to you without trying to anticipate them.

Continue to focus on your sensations, the ventilation of the cabin on your face, the sounds of your surroundings.

The belt signals come on and you hear thunder in the distance. Observe if this causes things to happen to you: for example, do you have any sensations (e.g. tension in your body? tingling?)? Welcome them, without judging. Let them come and go as they came. An emotion (fear, anger, sadness, surprise, disgust, joy...)? Welcome it, without judgement. Take them into account as they are, in the present moment, without judgment. Then let them go as they came. Let them go..., without trying to hold them back, and without trying to make them go away: their departure happens gradually, passively. Let it happen...

Perhaps you notice an increase in your heart rate, your heart beating faster..., or tensions in your body, muscles that tighten...

You let it happen, without judgment. Let it happen, without trying to control anything. Let these sensations pass, the emotions that come, as they have come. Let it happen, passively, so that they go away quietly as they came. You let it happen..., without trying to modify your breathing. You let it happen, in confidence, quietly.

Around you, you observe the luggages falling and the elements falling over. Focus on the way it makes you feel. Accept it and let it go without chasing it or holding it back. Perhaps you notice an increase in your heart rate, your heart beating faster..., or tension in your body, muscles tightening...

You let it happen, without judgment. Let it happen, without trying to control anything. Let these sensations pass, the emotions that come, as they have come. Let it happen, passively, so that they go away quietly as they came. You let it happen..., without trying to modify your breathing. You let it happen, in confidence, quietly.

The plane calms down again and the turbulence goes away. Continue to observe and feel your surroundings during the turbulence. Observe if these sensations are pleasant for you, if they leave an imprint on you, a pleasant sensation or not. Again, without judging, welcoming things as they are.

Let all sensations, thoughts or emotions in you quietly and passively subside. You let it happen... Without trying to change your breathing, without trying to hold anything inside you... You let the sensations that may be present happen, until they disappear by themselves. You let it happen. To finish, I suggest that you focus your attention on a particular object, for example a book, something that seems good to you... Choose what you like.

Approach it and give it your full attention. Without judgment, observe its colours, its shape. Does it smell? Does it make sounds when you bring it close to your ear? Take a few moments to observe... When you take this time, observe if you have thoughts, sensations, emotions... Without judgement. Taste the present moment... The "here and now".

Observe if these sensations are pleasant to you, if they leave an imprint on you, a pleasant taste or not. Once again, without judging, welcoming things as they are.

We are coming to the end of this experience, thank you for participating and see you soon for the next discovery together.

Measurement of the Perception of a Stimulating Environment Questionnaire (MPSEQ)

(Chatard A, Barillot L, Besnier M, et al. Measurement of the Perception of a Stimulating Environment Questionnaire, unpublished)

Please indicate your degree of agreement with the statements below using the following response scale: 1 = Strongly disagree; 2 = Disagree; 3 = Somewhat disagree; 4 = Neither agree nor disagree; 5 = Somewhat agree; 6 = Agree; 7 = Strongly agree

1. My immediate environment is rich in sensations and stimuli of all kinds ____
2. Often, I tell myself that my life is not very exciting ____
3. I practice many physical and/or sports activities ____
4. My physical and/or sports activities are a source of personal development ____
5. I have a rich inner life ____
6. I play and/or listen to a lot of music ____
7. I watch a lot of exiting and stimulating films and documentaries ____
8. I have a lot of friends and social connections ____
9. My social relationships are rich and stimulating ____
10. I do many diverse and varied activities ____
11. I have a passion in life (ex: a hobby) ____
12. I live in a stimulating environment ____
13. I have the impression that my daily environment is boring, that it is always the same ____

Table 1

Study schedule of enrolment and assessments by time points (Tn).

S1, S2, S3, S4, S5 and S6 refer to the 6 different EE sessions of the enhanced intervention.

Time point	Enrolment	Enhanced intervention						Post-intervention	Follow-ups		
	T0	T1						T2	T3	T4	T5
		S1	S2	S3	S4	S5	S6				
Eligibility screening	X										
Informed consent	X										
Group allocation	X										
Primary outcome											
TLFB									X	X	X
Breathalyser									X	X	X
Blood test								X	X	X	X
Secondary outcomes											
OCDS	X								X	X	X
Craving induced by cues	X							X			
IAT	X							X			
Alcohol seeking	X							X			
Alcohol attentional bias	X							X			
MPSEQ	X								X	X	X
FFMQ	X		X					X			
Salivary cortisol	X					X	X				