




BMJ Open Oral-only antibiotics for bone and joint infections in children: study protocol for a nationwide randomised open-label non-inferiority trial

Allan Bybeck Nielsen ¹, Luise Borch,^{2,3} Mads Damkjaer ^{4,5}, Jonathan Peter Glenthøj,⁶ Ulla Hartling,⁷ Thomas Ulrik Hoffmann,⁸ Mette Holm,⁹ Annett Helleskov Rasmussen,⁴ Lisbeth Samsø Schmidt,¹⁰ Kjeld Schmiegelow,¹ Lone Graff Stensballe ¹, Ulrikka Nygaard,¹ Local Investigators

To cite: Bybeck Nielsen A, Borch L, Damkjaer M, *et al.* Oral-only antibiotics for bone and joint infections in children: study protocol for a nationwide randomised open-label non-inferiority trial. *BMJ Open* 2023;**13**:e072622. doi:10.1136/bmjopen-2023-072622

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2023-072622>).

Received 15 February 2023
Accepted 14 May 2023

ABSTRACT

Introduction Children with bone and joint infections are traditionally treated with intravenous antibiotics for 3–10 days, followed by oral antibiotics. Oral-only treatment has not been tested in randomised trials.

Methods and analysis Children (3 months to 18 years) will be randomised 1:1 with the experimental group receiving high-dose oral antibiotics and the control group receiving intravenous antibiotics with a shift in both groups to standard oral antibiotics after clinical and paraclinical improvement. Children in need of acute surgery or systemic features requiring intravenous therapy, including septic shock, are excluded. The primary outcome is defined as a normal blinded standardised clinical assessment 6 months after end of treatment. Secondary outcomes are non-acute treatment failure and recurrent infection. Outcomes will be compared by a non-inferiority assumption with an inferiority margin of 5%.

Ethics and dissemination The trial has the potential to reduce unnecessary hospitalisation and use of intravenous antibiotics in children with bone or joint infections. Due to the close follow-up, exclusion of severely ill children and predefined criteria for discontinuation of the allocated therapy, we expect the risk of treatment failure to be minimal.

Trial registration number NCT04563325.

INTRODUCTION

In recent years, the necessity of intravenous antibiotic treatment of children and adults with severe bacterial infections has been questioned as the evidence hereof is lacking.¹ Oral-only treatment has been demonstrated to be safe and non-inferior to intravenous treatment for children with, for example, severe pneumonia² and febrile urinary tract infections³ including urinary tract infections with bacteraemia.⁴ In large prospective randomised trials, early shift from intravenous to oral antibiotic treatment and even oral-only antibiotic treatment has been demonstrated to be safe and non-inferior to longer

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The prospective randomised design with blinded evaluation of the primary outcome will reduce potential bias.
- ⇒ The primary outcome, sequelae at 6 months, is of clear clinical relevance.
- ⇒ The pragmatic design with full integration into daily clinical practice ensures a high transferability of the generated data.
- ⇒ The 12-month follow-up reduces the risk of missing long-term sequelae.
- ⇒ The pragmatic criteria of inclusion involve a risk of including children with non-bacterial bone or joint disease.

courses of intravenous treatment for adults with infectious diseases such as endocarditis^{5,6} and bone and joint infection (BJI).^{1,7} Several oral antibiotics have demonstrated good bone and joint penetration profiles compared with their respective plasma concentrations.⁸

Prospective studies on children with BJI have demonstrated that individualised antibiotic therapy with significantly reduced treatment duration is non-inferior to the previous long-term treatment^{9–12} and that only 2–4 days of intravenous therapy followed by 1–3 weeks of oral therapy is sufficient in most cases. The recommendations of individualised and shorter treatment have been adopted in Denmark with good results.¹³ These significant changes in treatment strategies raise the question of whether most children with BJI could be treated with oral antibiotics only. One small case series¹⁴ and one retrospective case-control study¹⁵ found oral-only therapy as a valid alternative to the current standard treatment, but the issue has not been investigated in randomised controlled trials (RCTs).



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Allan Bybeck Nielsen;
allan.bybeck.nielsen@regionh.dk

**Box 1** Criteria of inclusion and exclusion**Inclusion criteria**

The participant must meet both of the following criteria:

1. Age: 3 months to 18 years at study entry.
2. Antibiotic treatment of uncomplicated bone or joint infection.

Exclusion criteria

Any of the following criteria exclude the participant from the study:

1. Severe disease at admission, for example, septic shock, or any concomitant invasive infection, for example, necrotising fasciitis, requiring intravenous antibiotics in the opinion of the treating clinician.
2. Complicated bone or joint infection, for example, prosthetic material, infection secondary to or complicated by trauma, severe pyomyositis or other substantial soft tissue infections.
3. Expected need of major surgery within the first 24 hours of treatment, for example, drilling, debridement, fenestration, surgical drainage, synovectomy. Minor surgery as diagnostic surgical bone biopsy or diagnostic joint fluid aspiration including lavage is not a criterion for exclusion.
4. Significant comorbidities that might influence the choice of treatment or the course of the infection, for example, immunodeficiency or sickle cell anaemia.
5. Previous bone or joint infection.
6. Antibiotic therapy for more than 24 hours before inclusion.
7. Documented pathogen with limited treatment options that do not permit randomisation, for example, the pathogen is only sensitive to intravenous antibiotics.
8. Prior enrolment in the trial.

METHODS AND ANALYSIS**Study hypothesis and design**

We hypothesise that oral-only antibiotic therapy is non-inferior (non-inferior margin 5%) to initial intravenous therapy, followed by oral therapy in children and adolescents with BJI and test that hypothesis in a nationwide, multicentre, randomised, controlled, open-label, non-inferiority trial.

Included children will be randomised 1:1 with the experimental group receiving high-dose oral antibiotic therapy and the control group receiving intravenous antibiotic therapy with a shift in both groups to standard oral therapy after clinical and paraclinical improvement.

Trial participants

Children and adolescents are considered for inclusion when they present with suspicion of BJI. Recruitment will be nationwide from all 18 paediatric departments in Denmark. Request a complete list of study sites and primary investigators at allan.bybeck.nielsen@regionh.dk.

Criteria of inclusion are listed in [box 1](#).

Delayed inclusion

Most patients will be included in the study before the start of treatment. If inclusion is not performed before administration of antibiotic therapy (eg, the treating physician is not aware of the project), up to 24 hours of antibiotic treatment will be accepted before inclusion in the study to achieve adequate participant enrolment.

Early termination of falsely included patients

To reflect the daily clinical setting and facilitate timely entry to the study, the patients are eligible for entry based on the available clinical information, often before the final diagnosis is made. In most cases, the clinical suspicion of infection is sustained (eg, by positive microbiology or imaging), and the study treatment is completed with or without further evidence of infection. If the suspicion of infection is abandoned during the study period, it will lead to early termination from the study according to the following criteria:

1. The suspicion of infection is abandoned, and antibiotic therapy is stopped prematurely.
2. The course of the disease is explained by another diagnosis (not BJI) that is made during treatment or follow-up. Examples:
 1. Imaging reveals another diagnosis, for example, bone tumour, Legg-Calve-Perthes disease, or soft tissue infection with no involvement of bone or joint structures.
 2. Recurrent symptoms with no confirmation of infection leads to another final diagnosis, such as juvenile idiopathic arthritis, chronic recurrent multifocal osteomyelitis or other non-infectious diseases.
 3. Lyme arthritis is suspected, and the antibiotic treatment is changed targeting *Borrelia burgdorferi*.

Early terminated patients without BJI will not continue follow-up in the trial. They will be included in the analysis of the safety outcomes, but not in the analysis of any other outcomes. For transparency, details on all early terminated patients will be reported.

Minimising bias

Staff and participants are not blinded to the treatment allocation, since we consider intravenous placebo treatment to be unethical in this group of children and adolescents. To minimise bias, the evaluation of the primary outcome is blinded.

Diagnostic procedures and procedures during antibiotic treatment

According to randomisation the child will receive (1) oral-only antibiotic therapy or (2) intravenous antibiotic therapy with shift to oral therapy after clinical and paraclinical improvement. All other aspects of diagnosing and treating the infection will follow current guidelines on paediatric BJI. This includes recommendations for mobilisation, physiotherapy, analgesic drugs, etc, which are all permitted during the trial. A throat swab for *Kingella kingae* will be added to the routine diagnostic procedures. All children will be evaluated with routine blood samples and clinical examination including grading of symptoms and pain score. See online supplemental appendix 1 for collection of extra material. The parents and/or the child will complete a daily registration of temperature and pain score, which will be reported weekly in an electronic questionnaire directly linked to the electronic case report form (eCRF). In these questionnaires, the parents and/

	STUDY PERIOD									
	Enrolment Allocation		Intervention (high dose IV vs. high dose Oral)				Standard follow-up treatment		Follow-up	
	D ₋₁	D ₀	D ₁	D ₂	D ₃	...	D _†	D _{†+7/21/28}	M ₆	M ₁₂
Enrollment:										
Eligibility screen	(x)	x								
Informed consent		x								
Allocation		x								
Intervention:										
High dose IV (standard group)		x	x	x	x	x				
High dose oral (experimental group)		x	x	x	x	x				
Standard dose oral (both groups)							x			
End of therapy (both groups)								x		
Assessments:										
Clinical assessment	(x)	x	(x)	(x)	(x)	(x)	x	x		x
Blood samples / imaging	(x)	x	(x)	(x)	(x)	(x)	x	x	(x)	x
Parental registration (temperature, pain, side effects)		x	x	x	x	x	x	x		
Blinded clinical assessment (primary endpoint)									X	
Assessment of non-primary endpoints			x	x	x	x	x	x	x	x
24/7 hotline to hospital staff		x	x	x	x	x	x	x	x	x

x = performed

(x) = performed if clinically relevant

D1 = Day 1, D2 = Day 2 etc.

D_† = Day of clinical/paraclinical improvement

D_{†+7/21/28} = 7 days (septic arthritis), 21 days (osteomyelitis) or 28 days (spondylodiscitis) after D_†

M6 = Month 6, M12 = Month 12

Figure 1 Participant timeline.

or the child will also report adherence (number of missed doses) and drug side effects. The participant timeline is illustrated in [figure 1](#).

Choice of antibiotic treatment and predefined treatment strategies

The antibiotic treatment is administered as ‘initial treatment’ (intravenous vs high-dose oral) with a shift to ‘follow-up treatment’ (oral) after clinical and paraclinical improvement.

Initial treatment (randomised)

Empiric intravenous treatment (standard group):

- ▶ Ceftriaxone 100 mg/kg/day (maximum 4000 mg/day).

Empiric oral treatment (experimental group):

- ▶ Age <5 years:
 - Amoxicillin+clavulanate 1:8, 100 mg/kg/day divided every 8 hours (maximum 3000 mg amoxicillin/day).
- ▶ Age >5 years:
 - Dicloxacillin 200 mg/kg/day divided every 6 hours (maximum 8000 mg dicloxacillin/day).

Follow-up treatment (not randomised)

Empiric oral treatment (both groups):

- ▶ Age <5 years:
 - Amoxicillin+clavulanate 1:4, 50 mg/kg/day divided every 8 hours (maximum 1500 mg amoxicillin/day).
- ▶ Age >5 years:
 - Dicloxacillin, 100 mg/kg/day divided every 6 hours (maximum 4000 mg dicloxacillin/day).

If infection with *Staphylococcus aureus* is suspected or confirmed, oral rifampicin 20 mg/kg/day divided every 8 hours (maximum 900 mg/day) can be added to the empiric therapy in both groups for optimal penetration and staphylococcal coverage.

According to daily clinical practice, the empiric antibiotic can be adjusted within the same route of administration (oral to oral or intravenous to intravenous) due to susceptibility testing, allergic reactions, adverse events or patient preferences to increase adherence (eg, taste) in both groups.

Treatment duration

In both groups, the treatment duration is individualised according to the rate of improvement. The minimum duration of the ‘initial treatment’ is 3 days with a shift to ‘follow-up treatment’ after clinical improvement (pain and mobility) and paraclinical improvement (decreasing

C reactive protein (CRP)). The follow-up treatment is given for 4 weeks (spondylodiscitis), 3 weeks (other bone infections) or 1 week (joint infection only). The follow-up treatment will only be terminated if the child has no symptoms and a normal clinical examination, otherwise, the antibiotic therapy can be prolonged until these criteria are fulfilled.

Follow-up

All children will receive a clinical follow-up after 6 and 12 months. Any complaints or positive findings during follow-up will be evaluated and treated according to local guidelines. All patients and parents will be instructed to contact the department in the case of any symptoms indicating BJI during follow-up.

Registration of information from patient records

During the treatment and follow-up, the local primary investigator will maintain contact with the clinical team to identify potential outcomes. The primary investigator will register information in the patient's eCRF in a predesigned database (software version: REDCap 9.1.0). Registered data will include age, sex, medical history, symptoms, clinical findings, blood samples, microbiological findings, imaging, treatment details and absence/presence of predefined outcomes. Missing data will be retrieved by a telephone interview if possible.

Primary outcome

Proportion of children with sequelae 6 months after initiation of treatment defined as abnormal mobility or function of the affected joint/bone. Evaluated by blinded clinical examination by a qualified paediatrician and/or paediatric orthopaedic surgeon (6 months).

Blinding

The assessor (qualified paediatrician or paediatric surgeon) of the primary outcome will be blinded for the intervention as well as any other details on the course of disease. The assessor will be informed about the age of the child as well as the approximate location of the infection and will perform a predefined systematic clinical examination of the relevant area (categories: (1) lower extremities, (2) upper extremities including clavulae and scapulae and (3) column, costae and sternum). The exact anatomical location including side (left or right) will not be revealed. The child and parents will be followed by a study nurse who will secure that the blinding is respected. The primary outcome is met if there are any positive findings related to the previously infected bone or joint.

If indicated, any clinical findings will be further evaluated according to standard clinical practice and if this evaluation fails in confirming the sequelae (eg, suspicion of clinical scoliosis followed by normal scoliosis imaging), the primary outcome is not met.

Secondary outcomes

1. Non-acute treatment failure. Proportion of children with change of antibiotic therapy due to non-acute treatment failure (28 days).

This will be evaluated by two paediatric specialists and is suggested by the following parameters:

1. Temperature above 38.5°C after more than 72 hours of antibiotic therapy.
 2. Increasing CRP after more than 96 hours of antibiotic therapy.
 3. No improvement in mobility or pain after 120 hours of antibiotic therapy.
2. Recurrent infection. Proportion of children with recurrence of symptoms and signs (same anatomical location) after completion of antibiotic treatment requiring further antibiotic administration (6 months).

Other predefined outcome measures

Safety outcomes and other predefined outcome measures are listed in [table 1](#).

Discontinuation/withdrawal of participants from study treatment

If the allocated treatment is no longer compatible with good clinical care, the randomised strategy will be discontinued. The participant will continue follow-up in the trial and will be included in the primary analyses (intention to treat, ITT) but not in secondary analyses (per-protocol). Routine clinical care consistent with the new information will be recommended. Reasons for discontinuation/withdrawal of the randomised strategy include:

1. Development of severe disease or complications (safety outcome no. 1).
2. Non-acute treatment failure (secondary outcome no. 1). Non-acute treatment failure will be evaluated by two paediatric specialists, the local paediatrician responsible for the treatment, and one of the chief investigators. The decision to change therapy will be based on a complete evaluation of the clinical condition and the treating paediatrician will make the final decision.
3. Unable to receive oral treatment, for example, vomiting with no improvement on antinausea medications. If the patient is only temporarily unable to receive oral treatment, for example, intercurrent gastroenteritis or vomiting primarily due to pain or fever, intravenous treatment for maximum 24 hours will be accepted without discontinuation of the allocated oral strategy.
4. Unable to maintain intravenous access. If the participant is temporarily without intravenous access, for example, waiting for anaesthetic assistance, oral treatment for maximum 24 hours will be accepted without discontinuation of the allocated intravenous strategy.
5. No suitable medication exists within the allocated strategy (intravenous vs oral) due to susceptibility testing, contraindications or adverse reactions.
6. Non-adherence to the allocated therapy.
7. Withdrawal of consent from participant/parents.

Table 1 Safety outcomes and other predefined outcome measures

No	Category	Specification	Time frame
1	Safety	Proportion of children with severe complications during antibiotic treatment, for example, need for intensive care, septic shock, organ failure, pyomyositis, endocarditis, deep venous thrombosis.	28 days
2	Safety	Proportion of children with need for surgical intervention during antibiotic treatment. Diagnostic surgical intervention (diagnostic joint aspiration or diagnostic bone biopsy) excluded	28 days
3	Safety	Proportion of children with treatment-related adverse events for example, complications of intravenous access (infection, need for replacement, extravasation) and drug side effects reported by medical staff or by parents (electronic questionnaire)	3 months
4	Exploratory	Time to apyrexia from initiation of antibiotic treatment	28 days
5	Exploratory	Level of mobility and pain assessed by daily grading of symptoms by medical staff and daily standardised pain scores from participants and/or parents. Score systems: Visual Analogue Scale or Face Legs Activity Cry Consolability scale, both with scores from 0 (no pain) to 10 (worst pain).	14 days
6	Exploratory	Total duration of antibiotic therapy	3 months
7	Exploratory	Proportion of children with sequelae, for example, abnormal mobility and growth abnormalities, assessed by clinical examination by a qualified paediatrician 12 months after the initiation of treatment, accepted range 11–14 months	14 months
8	Exploratory	Proportion of children with radiological abnormalities assessed by a qualified radiologist 12 months after initiation of treatment, accepted range 11–14 months	14 months
9	Exploratory	Secondary infection with antimicrobial-resistant organisms or <i>Clostridioides difficile</i>	3 months

Statistical methods and analyses

The plan for statistical analyses is based on the estimand framework.^{16 17}

Sample size and power calculation

Outcomes will be compared by a non-inferiority assumption with an inferiority margin of 5% and an expected treatment success of 99% in both groups. With a one-sided significance level (alpha) of 2.5% and accounting for a 10% rate of drop-out, a sample size of 180 children in the principal stratum, 90 children in each group, will provide a power of 90% to detect non-inferiority.

Randomisation

Web-based computer randomisation (TrialPartner by DEFACTUM) will be used to allocate patients to the treatment groups at a 1:1 ratio stratified by CRP (<35 vs ≥35). The randomisation will be in randomly permuted blocks with varying block sizes of 4 and 6.

Target population/the principal stratum

The principal stratum will include all randomised children but exclude children with the intercurrent event of early termination, that is, children with no BJI. All randomised children will be included in the analysis of the three safety outcomes. For the remaining outcomes, including the primary and secondary outcomes, only data from children in the principal stratum will be analysed.

Statistical analyses for the primary outcome

Based on the principal stratum, the population-level summary measure will be the risk difference, calculated as the proportion with sequelae after 6 months in the

experimental group (oral only) minus the proportion with sequelae after 6 months in the control group (intravenous+oral), with one-sided 97.5% confidence limits. If the absolute upper one-sided 97.5% CI is less than 5%, then the criteria of non-inferiority will be met.

Based on ITT principle, the primary analysis of the primary outcome will include all participants in the principal stratum.

A secondary per-protocol analysis (PPP) will also be performed excluding all participants experiencing one or more of the predefined intercurrent events leading to a discontinuation of the allocated treatment strategy.

Missing data

The primary outcome will be evaluated in all participants who attend the follow-up after 6 months. We only expect missing data if the participant is lost to follow-up or withdraw their consent. Due to the close follow-up at each local department, we expect missing data to be less than 5% and equal in both groups. If the missing data are more than 5%, we will impute data based on available knowledge for the patient and the observations of the other patients in the same randomised group. Sensitivity analyses will include tipping point analyses for the imputation.

Statistical analyses for the secondary outcomes

Based on the principal stratum, the primary analysis (ITT) described in 6.4.2 will be repeated for the secondary outcomes: ‘non-acute treatment failure’ and ‘recurrent infection’. The secondary analysis (PPP) described above will be repeated for ‘recurrent infection’ but not for

'non-acute treatment failure' while this outcome is one of the intercurrent events excluding the participant from the secondary analysis (PPP).

Missing data

If a participant experiences one of the secondary outcomes, it will be registered in the medical record. Due to the close follow-up and the national design including complete surveillance of all paediatric departments in Denmark with access to all electronic medical records, only cases where participants withdraw from all follow-up or relocate to another country are at risk for missing data on secondary outcomes. We expect this to be less than 5% and no imputation is planned for the secondary outcomes.

Statistical analyses for the safety outcomes

For safety outcomes 1 and 2, we will report the total number of participants in each treatment group meeting these outcomes.

For safety outcome 3, the outcome will be the fraction of days the child is affected by one or more of these non-serious adverse events (SAEs). The two randomised groups will be compared using the Wilcoxon-Mann-Whitney test. The population-level summary measure will be the difference between the means of the fractions with two-sided 95% CIs based on the normal approximation for the estimated mean difference. The mean of the individual fractions in each randomised group will also be presented with two-sided 95% CIs based on the normal approximation for the estimated mean.

In addition, we will also for each randomised group present the total number of non-SAEs, the number of children who had one or more non-SAEs, the number of non-SAEs per person-year at risk and the total number of days with one or more non-SAE.

Risks and safety monitoring

All antibiotics used in the trial are licensed agents with well described safety profiles approved for treating BJIs in children. The trial is not a clinical trial of an investigational medical product. The following, therefore, describes our own procedures for safety reporting.

Since our hypothesis is founded on the existing literature^{14 15} as well as the general development in antibiotic treatment for children in recent years²⁻⁴ and since we are excluding children with severe disease, we consider the risk of insufficient treatment to be minimal. Due to the close follow-up, we expect to promptly recognise insufficient treatment and evaluate whether a change in treatment is needed. In our recent retrospective study on Danish paediatric BJI infections,¹³ approximately 5% of children receiving the current standard treatment needed a change in antibiotic therapy due to insufficient effect (data not published). In the current study, we expect a similar need for change in therapy in both groups. The infection and the antibiotic treatment may result in discomfort (pain, fever), side effects (loose stool,

abdominal pain, rash) or rarely other complications, but the risk of these events is not increased due to participation in the study. Unexpected adverse effects may occur, but the close follow-up including a 24/7 hotline to the hospital staff, enables us to react promptly to any unexpected incidents. Children and adolescents in the experimental group can avoid the establishment of intravenous access which in most children is a painful procedure. On the other hand, children in the experimental group need to take oral antibiotics for a longer period.

SAEs are not expected in this study due to the exclusion of patients with septic appearance and the fact that paediatric BJI is a condition not associated with mortality or severe morbidity. Only 1 of 82 children in our recent retrospective study of paediatric BJI developed sepsis or any other SAE.¹³

Safety and data monitoring committee

For monitoring the study, an independent safety and data monitoring committee will be established. The committee will consist of two independent paediatricians and a biostatistician with experience in paediatric infectious diseases and in the monitoring of RCTs will be consulted if needed. The committee will function independently of all other individuals conducting the study and all potential and SAEs will be reported to the committee. The committee will review all safety data by treatment arm every 6–12 months. The committee will be empowered to advise suspending the trial.

Roles, funding and patient compensation

The project is initiated by chief investigators Allan Bybeck Nielsen and Ulrikka Nygaard (sponsor). Each site has a primary investigator. The protocol is finalised by the listed authors which includes the chief investigators, main collaborators (Professor Lone Graff Stensballe and Professor Kjeld Schmiegelow) as well as primary investigators from sites with an expected inclusion of more than 10 participants. Funding is received from The Research Foundation of Copenhagen University Hospital, Copenhagen Health Science Partners and Innovation Fund Denmark. None of the investigators or main collaborators are financially attached to private enterprises, foundations, etc that have interests in the research project. Neither the patients enrolled in the project, nor their families, will receive any kind of payment for participating in the project.

ETHICS AND DISSEMINATION

The treating paediatrician will obtain informed consent according to the ethical approval, see information for parents in online supplemental appendix 2. As described in 'risks and safety monitoring', we consider the risk of harm to the participants to be minimal. The hypothesis has never been tested before and the radical change in treatment cannot be made in a scientific ethical sense without being based on a study at the current level of

evidence. The hypothesis has not been tested in adults and it is not possible to perform the study on adults and transmit the data directly to children, as there are large differences in the course of BJIs between children and adults. If the experimental therapy is non-inferior to the standard therapy, the study is expected to lead to a radical change in the treatment of BJI in children and adolescents. The results of the project (whether positive, negative or inconclusive) will be submitted for publication in scientific peer-reviewed international journals. All investigators and main collaborators will be given the opportunity of authorship of the main publication(s) according to the International Committee of Medical Journal Editors (ICJME) criteria.

The study will be performed under the approval of The Ethics Committee of the Capital Region of Denmark (Ref.: H-20009117) and conducted according to the guidelines and recommendations of Good Clinical Practice and the Declaration of Helsinki.

Patient and public involvement

Parents of children in different age groups were involved in designing the participant information. No patient, parent or the public was involved in designing, writing or editing the protocol.

Trial registration, trial status and protocol amendments

The study is registered at www.clinicaltrials.gov (NCT04563325), trial registration dataset in online supplemental appendix 3. Recruitment was started on 15 September 2020. All sites are active, and 148 patients have been recruited in the principal stratum by 1 February 2023.

In version 1.0 of the protocol and at the initial registration on www.clinicaltrials.gov, all outcomes, except the primary outcome, were categorised as secondary outcomes. After finishing the statistical analysis plan, the categories of the outcomes were changed so that only two secondary outcomes are now registered. The remaining nine are now registered as 'Other Outcome Measures' including three safety outcomes corresponding to the current protocol version 2.0. In this revision of the outcome categories after the initiation of the study, four exploratory outcomes were dismissed:

1. Level of CRP. Comment: Will be reported but not considered as an outcome.
2. Disability days. Comment: This registration was missed in the electronic questionnaire and data are not registered.
3. Duration of intravenous antibiotics. Comment: Will be reported but not considered as an outcome.
4. Adherence (parental report of missed doses). Comment: Will be reported but not considered as an outcome.

DISCUSSION

With the current protocol, we aim to investigate if oral-only antibiotic treatment is non-inferior to the current

standard therapy of initially intravenous antibiotic treatment followed by oral treatment in children and adolescents with BJI.

The strengths of our study are the randomised controlled design with a blinded primary outcome evaluation and the national inclusion of all 18 paediatric departments in Denmark. The trial is fully embedded into daily clinical practice and reflects standard practice in all aspects other than randomisation and collection of data. The main limitation is the pragmatic criteria of inclusion without strict diagnostic criteria introducing a potential risk of including children with non-bacterial arthritis or osteomyelitis. We aim to reduce this risk by early termination of children if a non-infectious condition is established during treatment or follow-up. The pragmatic design ensures timely inclusion of patients and reflects daily clinical practice where antibiotic therapy is often initiated before strict diagnostic confirmation of bacterial infection.

The choice of a non-inferiority design was based on the combination of (1) the high treatment success of the current intravenous treatment and the overall favourable outcomes of these types of infection, (2) the practical aspects of realistic recruitment and (3) the potential large benefits of the experimental oral treatment. This choice is in accordance with current recommendations on non-inferiority clinical trials.¹⁸ We initially decided to use a 10% non-inferiority margin, which was based on consensus among a wide range of infectious disease specialists, paediatric infectious disease specialists and paediatric orthopaedic surgeons balancing the potential risks and benefits of oral treatment. This margin was further supported by published guidelines suggesting similar non-inferiority margins when evaluating treatment of other bacterial infections.¹⁹ After 6-month follow-up of the first 100 children, no sequelae were registered in the trial, equaling a treatment success of 100% in both groups. Due to this very high overall treatment success, we reduced the non-inferiority margin to 5% and adjusted the sample size estimation to 184 (initially 180) with expected treatment success of 99%.

Based on extrapolated Danish data from 2012 to 2017,¹³ more than 100 children are diagnosed with BJI in Denmark each year. If the experimental therapy is successful, these children will benefit significantly from the new treatment in terms of shorter admissions, avoidance of intravenous access, earlier return to everyday activities and better quality of life during the treatment. The results are expected to influence the treatment strategies of children with BJIs worldwide. Especially in countries with similar resistance spectrum where the results can be directly applied, but also in countries with a more challenging resistance spectrum, where the general demand (often non-evidence based) of intravenous antibiotics for severe infections will be challenged.

Study sites

Department of Pediatrics and Adolescent Medicine, Lillebaelt Hospital, Kolding

Department of Pediatrics and Adolescent Medicine, Regional Hospital, Viborg

Department of Pediatrics and Adolescent Medicine, Sydvestjysk Hospital, Esbjerg

Department of Pediatrics and Adolescent Medicine, Aalborg University Hospital

Department of Pediatrics and Adolescent Medicine, Nordsjælland Hospital, Hillerød

Department of Pediatrics and Adolescent Medicine, Slagelse Hospital

Department of Pediatrics and Adolescent Medicine Pediatrics, Herlev Hospital

Department of Pediatrics and Adolescent Medicine, Zealand University Hospital, Roskilde

Department of Pediatrics and Adolescent Medicine, Regional Hospital Vestjylland, Herning

Department of Pediatrics and Adolescent Medicine, Holbæk Hospital

Department of Pediatrics and Adolescent Medicine, Aarhus University Hospital

Department of Pediatrics and Adolescent Medicine, Regional Hospital Randers

Department of Pediatrics and Adolescent Medicine, Hjørring Hospital

Department of Pediatrics and Adolescent Medicine, Sønderjylland Hospital, Aabenraa

Department of Pediatrics and Adolescent Medicine, Nykøbing Falster Hospital

Department of Pediatrics and Adolescent Medicine, Hvidovre Hospital

Department of Pediatrics and Adolescent Medicine, Odense University Hospital

Department of Pediatrics and Adolescent Medicine, Rigshospitalet, Copenhagen University Hospital

Author affiliations

¹Department of Paediatrics and Adolescent Medicine, Copenhagen University Hospital, Kobenhavn, Denmark

²Department of Paediatrics and Adolescent Medicine, Gødstrup Hospital, Herning, Denmark

³NIDO - Centre for Research and Education, Gødstrup Hospital, Herning, Denmark

⁴Department of Paediatrics and Adolescent Medicine, Lillebaelt Hospital - University Hospital of Southern Denmark, Kolding, Denmark

⁵Regional Health Research, University of Southern Denmark, Odense, Denmark

⁶Department of Paediatrics and Adolescent Medicine, Copenhagen University Hospital, Hillerød, Denmark

⁷Department of Paediatrics and Adolescent Medicine, Odense University Hospital, Odense, Denmark

⁸Department of Paediatrics and Adolescent Medicine, Copenhagen University Hospital, Hvidovre, Denmark

⁹Department of Paediatrics and Adolescent Medicine, Aarhus University Hospital, Aarhus, Denmark

¹⁰Department of Paediatrics and Adolescent Medicine, Copenhagen University Hospital, Herlev, Denmark

Collaborators The authors thank all the local investigators (Grethe Lemvik Mikkelsen, Regional Hospital Viborg; Jens Jakob Herrche Petersen, Sydvestjysk Hospital, Esbjerg; Jesper Thaarup, Aalborg University Hospital; Kim Kristensen; Lise

Heilmann Jensen, Zealand University Hospital, Roskilde; Marie Cecilie Lawaetz, Holbæk Hospital; Morten Søndergaard Lindhard, Regional Hospital Randers; Pawel Andrej Marcinski, Hjørring Hospital; Lotte Høeg Hansen, Sønderjylland Hospital, Aabenraa, Tanja Hübertz Horsager, Regional Hospital Viborg; Maren Rytter, Slagelse Hospital; Tatjana Zaharov, Nykøbing Falster Hospital) for their big effort in preparing and informing the individual sites about the trial before initiation. We also thank Jakob Hjort, data manager, Department of Clinical Medicine, Aarhus University, for establishing the web-based randomisation module as well as Ulrik Justesen, Odense University Hospital for microbiological advice.

Contributors ABN and UN initiated and planned the study, and they contributed equally to the initial conception and design of the study protocol. ABN and UN drafted the first and subsequent versions of the study protocol. LB, MD, JPG, UH, TUH, MH, AHR, LSS, KS and LGS made substantial contributions to the design of the study, and they critically revised the study protocol. All authors read and approved the final manuscript and agree to be accountable for all aspects of the work.

Funding Funding is received from The Research Foundation of Copenhagen University Hospital, Copenhagen Health Science Partners and Innovation Fund Denmark. The funding covers salaries for study doctors, statisticians, and additional expenses (eg, software and hardware). The funding is administered through research accounts at the Copenhagen University Hospital Rigshospitalet.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Allan Bybeck Nielsen <http://orcid.org/0000-0002-9035-830X>

Mads Damkjaer <http://orcid.org/0000-0001-7410-8573>

Lone Graff Stensballe <http://orcid.org/0000-0003-1569-153X>

REFERENCES

- 1 Spellberg B, Aggrey G, Brennan MB, *et al*. Use of novel strategies to develop guidelines for management of Pyogenic Osteomyelitis in adults: A Wikiguideelines group consensus statement. *JAMA Netw Open* 2022;5:e2211321.
- 2 Lodha R, Kabra SK, Pandey RM, *et al*. Antibiotics for community-acquired pneumonia in children. *Cochrane Database of Systematic Reviews* 2013.
- 3 Strohmeier Y, Hodson EM, Willis NS, *et al*. Antibiotics for acute Pyelonephritis in children. *Cochrane Database Syst Rev* 2014:CD003772.
- 4 Hoberman A, Wald ER, Hickey RW, *et al*. Oral versus initial intravenous therapy for urinary tract infections in young febrile children. *Pediatrics* 1999;104:79–86.
- 5 Iversen K, Ihlemann N, Gill SU, *et al*. Partial oral versus intravenous antibiotic treatment of Endocarditis. *N Engl J Med* 2019;380:415–24.
- 6 Heldman AW, Hartert TV, Ray SC, *et al*. Oral antibiotic treatment of right-sided Staphylococcal Endocarditis in injection drug users: prospective randomized comparison with parenteral therapy. *Am J Med* 1996;101:68–76.
- 7 Wald-Dickler N, Holtom PD, Phillips MC, *et al*. Oral is the new IV. challenging decades of blood and bone infection dogma: A systematic review. *Am J Med* 2022;135:369–79.

- 8 Thabit AK, Fatani DF, Bamakhrama MS, *et al.* Antibiotic penetration into bone and joints: an updated review. *Int J Infect Dis* 2019;81:128–36.
- 9 Valenzuela MC, Viviani ST. Short duration of initial intravenous treatment in 70 pediatric patients with Osteoarticular infections. *Rev Chil Infectologia Organo Of Soc Chil Infectologia* 2008;25:30–6.
- 10 Jagodzinski NA, Kanwar R, Graham K, *et al.* Prospective evaluation of a shortened regimen of treatment for acute Osteomyelitis and septic arthritis in children. *J Pediatr Orthop* 2009;29:518–25.
- 11 Peltola H, Pääkkönen M, Kallio P, *et al.* Versus long-term antimicrobial treatment for acute Hematogenous Osteomyelitis of childhood: prospective, randomized trial on 131 culture-positive cases. *Pediatr Infect Dis J* 2010;29:1123–8.
- 12 Peltola H, Pääkkönen M, Kallio P, *et al.* Prospective, randomized trial of 10 days versus 30 days of antimicrobial treatment, including a short-term course of parenteral therapy, for childhood septic arthritis. *CLIN INFECT DIS* 2009;48:1201–10. 10.1086/597582 Available: <http://www.journals.uchicago.edu/toc/cid/48/9>
- 13 Nielsen AB, Nygaard U, Hoffmann T, *et al.* Short Individualised treatment of bone and joint infections in Danish children. *Arch Dis Child* 2019;104:205–6.
- 14 Alcobendas R, Remesal A, Murias S, *et al.* Outpatients with acute Osteoarticular infections had favourable outcomes when they received just oral antibiotics without intravenous antibiotics. *Acta Paediatr* 2018;107:1792–7.
- 15 Roul-Levy A, Looten V, Bachy M, *et al.* Oral ambulatory treatment of acute Osteomyelitis in children: A case-control study. *Pediatr Emerg Care* 2016;32:154–6.
- 16 International Council for Harmonisation. Addendum on Estimands and sensitivity analysis in clinical trials to the guideline OM statistical principles for clinical trials E9(R). n.d. Available: https://database.ich.org/sites/default/files/E9-R1_Step4_Guideline_2019_1203.pdf
- 17 Clark TP, Kahan BC, Phillips A, *et al.* Estimands: bringing clarity and focus to research questions in clinical trials *BMJ open* 2022;12:E052953. *BMJ Open* 2022;12:e052953.
- 18 FDA guidance on non-inferiority clinical trials. n.d. Available: <https://www.fda.gov/media/78504/download>
- 19 EMA guideline on the evaluation of medicinal products indicated for treatment of bacterial infections. EMA/844951/2018Rev. n.d. Available: https://www.ema.europa.eu/en/documents/scientific-guideline/draft-guideline-evaluation-medicinal-products-indicated-treatment-bacterial-infections-revision-3_en.pdf

Appendix 1: Biological specimens

Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in this trial/future use:

When routine samples (blood, joint fluid, bone biopsy, throat swabs or any other material) are collected, extra material will be collected to test and improve diagnostic tools in children with infections. Samples will only be obtained in addition to routine samples, i.e. blood will be obtained at the same time as routine blood samples without extra cannulations and joint fluid or bone material will only be obtained if joint puncture or bone biopsy is performed according to the national guideline for pediatric bone and joint infections. The maximum volume of blood taken for analysis is 2 ml/kg/day, the maximum volume of joint fluid is 10 ml (in addition to routine analysis) and the maximum number of bone biopsies is 2 (in addition to routine biopsies). These samples will be stored and analyzed according to the "CHILD" protocol approved by the Danish National Ethics Committee (protocol no. H-18065635). According to the "CHILD" protocol, these samples will be analyzed for pathogen-targeted PCR, host transcriptomics and other -omics, inflammatory parameters (cytokines and their receptors, immunomodulatory agents), and antibiotic concentrations. No analysis on the human genome of the child or adolescent will be performed. The samples will be saved in a research biobank until 31.12.2029. The remaining material will be transferred to a biobank and might be used for future research projects, but only after additional approval from the Ethics Committee and the Danish data protection agency.

APPENDIX 2: Parent information and consent form

Information for parents

Bone and joint infections in children and adolescents - a research project

Original title: Oral antibiotika til børn og unge med led- og knogleinfektioner. Et nationalt, randomiseret, kontrolleret forsøg.

Summary:

- Children and young people with bone and joint infections are treated with intravenous antibiotics for about 3-5 days followed by oral antibiotics for 1-4 weeks.
- We want to investigate whether oral antibiotics from day one is as effective as the current treatment. If this is confirmed, children with bone and joint infections can be treated at home and avoid unnecessary needlesticks.

Dear parents

We hereby ask if you would allow your child to participate in a research project on the treatment of children and adolescents with bone or joint infections. Before you decide, please read this information carefully and we will then talk to you about the project. Participation is voluntary, and you can withdraw your consent at any time and without explanation. We expect 200 children and young people from all over Denmark to participate in the project.

Purpose:

We want to investigate whether oral antibiotics (tablets or oral solutions) from day one (new treatment) are as effective as the current treatment, where antibiotics are given intravenously for the first days followed by oral treatment.

Background:

Today, we treat many infections in children and adolescents with oral antibiotics. However, joint and bone infections are still treated with intravenous antibiotics for the first 3-5 days followed by oral antibiotics for 1-4 weeks. The duration of intravenous antibiotic has been greatly reduced in recent years and some studies have shown that oral antibiotics throughout the full treatment period may be sufficient. However, so far, no studies have compared the two treatments.

Project plan:

Your child will randomly be selected to receive one of two treatments:

1. **New treatment:** Oral antibiotics from day one. The first dose is taken at the hospital, but then the treatment can take place either at home or in hospital depending on your child's condition.
2. **Current treatment:** Intravenous antibiotics from day one until there is improvement in the condition (typically 3-5 days). Then switch to oral antibiotics. The intravenous treatment will be performed in the hospital.

All children and adolescents will be followed closely by doctors and nurses for the effect of treatment (development in pain, redness, swelling, temperature). During treatment at home, all families will be

contacted by telephone twice a week by a project nurse as an additional follow-up. In addition, the Children and Adolescents Department can be contacted 24 hours a day throughout the treatment period. The effect of the treatment will be assessed with blood tests, among other things. The total duration of treatment follows common guidelines in both groups (usually 2-4 weeks) depending on the condition of the child.

There will be a follow up appointment after 6 and 12 months, during which time your child will be examined by a doctor, and an X-ray may be taken.

If your child does not participate in the project, he/she will receive the current treatment.

Sampling and storage of samples:

Children involved in the project will have blood tests taken according to the standard practice of treating bone and joint infections. When these standard blood samples are drawn, a small amount of blood will also be drawn for additional analyses. If other samples are taken due to standard practice, such as fluid from a joint, extra material will be stored if possible. In some cases, you will be asked for an additional sample, such as saliva, urine, feces or throat swab. The collected material will be stored in a research biobank until December 31st, 2029. It will be used to measure antibiotic concentrations and to investigate new methods for diagnosing infectious conditions in children and adolescents. The material will be encoded for anonymity and will not be directly traceable back to your child. After 2029, excess material will be transferred to a biobank for future research and will only be applicable after new approval from the Scientific Ethics Committee. You can always have the material from your child destroyed.

Side effects, risks, complications and disadvantages:

We ask you and your child to participate in this project because your child is among the children where we do not know if the best treatment is oral or intravenous antibiotics. There is a small risk that the new treatment is insufficient, which is why we will follow your child closely. If the treatment is insufficient, it will immediately be reassessed and, if necessary, changed. A small number of children are insufficiently treated with the current treatment and need a change of treatment along the way. We expect this to be the case with the new treatment as well.

Antibiotics can cause side effects such as abdominal pain, diarrhea and skin rash. We expect the side effects to be the same in the two groups. Extra blood samples will be taken at the same time as the regular blood samples, and your child will not be subjected to more needlesticks than usual. The total amount of blood taken each time will be a maximum of 1-2% of the child's blood volume and this will have no effect on the health of the child.

There may be risks to the project that we do not yet know, but we do not expect increased side effects or complications either in the short or long term.

Managing personal data:

The research team will obtain relevant health information in the electronic patient record relating to the disease episode, including the condition of the child, blood tests, bacterial examinations and scans. If we need more information, we will contact you after hospitalization with questions about the course of the disease. Before the analysis, the samples collected and data from medical records data will be assigned a

code so that the child's civil registration number (CPR number) does not appear directly. Data will be stored in a database created for the research project. The Danish Data Protection Act and Data Protection Regulation will be respected.

Benefits of the project

The expected benefits of the project include that most children and adolescents with bone or joint infections in the future will be able to receive oral antibiotics from day one. Hereby time at the hospital as well as insertion of intravenous catheters are reduced. This reduces insecurity in the child and increases the quality of life. There is no direct benefit for your child, but children who are randomized to the new treatment will avoid the insertion of an intravenous catheter.

Exclusion from the project

Your child can be excluded from the project if he/she appears seriously ill, if an intravenous catheter cannot be inserted, oral treatment cannot be performed, major surgery is needed, or another diagnosis is made along the way.

Financial support

The project has received funding from the Danish National Hospital's Research Foundation (DKK 3.25 million) and Copenhagen Health Science Partners (DKK 0.5 million) for salaries for researchers, analyses, statisticians and dissemination. None of the associated researchers have economic or commercial interests in the results of the study or its beneficiaries. There are no financially affiliated companies.

The project has been approved by the Scientific Ethics Committee with protocol number (j.nr. H-20009117). On the next page you will find information about your rights as a participant in a research project.

We hope that this information has helped you decide whether your child can take part in the project. You are welcome to contact us if you would like to know more about the project. It is possible to obtain information about the results when these are published on the project's website: childathome.dk.

Sincerely,

National primary investigator:

Allan Bybeck Nielsen, MD
BørneUngeKlinikken
Rigshospitalet
Tlf +45 3545 3545

Local primary investigator:

Participants rights in a health science research project

Information from the Scientific Ethics Committee, translated from "Forsøgspersoners rettigheder I et sundhedsvidenskabeligt forskningsprojekt"

As a parent to a participant in a biomedical research project, please note that:

- Your child's participation in the research project is entirely voluntary and he/she may only participate after you have been informed verbally and in writing about the research project and signed the consent form
- You may withdraw your consent verbally, in writing or by any other clear indication at any time and leave the project. Withdrawing your consent does not affect your right to current or future treatment or any other rights you or your child may have
- You are entitled to bring a family member, friend or acquaintance to the introductory consultation
- You are entitled to take time to think things over before you sign the consent form
- Information about your child's health as well as other personal and confidential information about your child that comes to light in connection with the research project are covered by a duty of confidentiality
- Storage of personal information on your child, including information contained in your child's blood samples and tissue, will be in accordance with regulations in the Danish Data Protecting Act and the Danish Health Act
- In accordance with the Danish Act on Free Access to Public Records, you have legal right of access to experimental protocols. This means that you can both access all documents concerning your child's participation in the study, except for the sections that contain trade secrets or confidential information about other people.
- In accordance with the Danish Act Governing the Right to Complain and to Obtain Compensation in the Danish Healthcare System, you may lodge a complaint and seek compensation. If you are injured during the trial, please contact the Patient Compensation Association; for more information visit www.patienterstatningen.dk.

October 2018

APPENDIX 3: Trial registration data set

DATA CATEGORY	INFORMATION
Primary registry and trial identifying number	ClinicalTrials.gov: NCT04563325
Date of registration in primary registry	18th September 2020
Secondary identifying numbers	H-20009117 (The Ethics Committee of the Capital Region of Denmark)
Source(s) of monetary or material support	Copenhagen University Hospital Rigshospitalet
Primary sponsor	Copenhagen University Hospital Rigshospitalet, Ulrikka Nygaard
Secondary sponsor(s)	N/A
Contact for public queries	Allan Bybeck Nielsen allan.bybeck.nielsen@regionh.dk
Contact for scientific queries	Allan Bybeck Nielsen allan.bybeck.nielsen@regionh.dk
Public title	ORAL-ONLY ANTIBIOTICS FOR BONE AND JOINT INFECTIONS IN CHILDREN
Scientific title	ORAL-ONLY ANTIBIOTICS FOR BONE AND JOINT INFECTIONS IN CHILDREN - A nationwide randomized open-label non-inferiority trial.
Countries of recruitment	Denmark
Health condition(s) or problem(s) studied	Antibiotic treatment of bone and joint infections in children and adolescents
Intervention(s)	Standard: IV + oral antibiotic treatment vs. Experimental: Oral only antibiotic treatment.
Key inclusion and exclusion criteria	Inclusion: - Age 3 months to 18 years - Antibiotic therapy for bone or joint infection. Exclusion: - Severe disease at admission - Complicated bone or joint infection - Expected need of major surgery within the first 24 hours of treatment - Significant co-morbidities - Previous bone or joint infection. - Antibiotic therapy for more than 24 hours before inclusion.
Study type	Open label, randomized 1:1, non-inferiority trial. Investigator of the primary endpoint is blinded to treatment allocation.
Date of first enrolment	Sep 2020
Target sample size	180
Recruitment status	Recruiting
Primary outcome	Proportion of children with sequelae 6 months after initiation of treatment defined as abnormal mobility or function of the affected joint/bone. Evaluated by blinded clinical examination by a qualified pediatrician and/or pediatric orthopedic surgeon.
Secondary outcomes	- Non acute treatment failure - Recurrent infection