




# BMJ Open Personal strategies to reduce the effects of landscape fire smoke on asthma-related outcomes: a protocol for systematic review and meta-analysis

Tesfalidet Beyene <sup>1,2</sup>, Peter G Gibson,<sup>1,2,3</sup> VE Murphy <sup>1,2</sup>, Megan E Jensen,<sup>1,2</sup> Vanessa M McDonald <sup>2,3,4</sup>

**To cite:** Beyene T, Gibson PG, Murphy VE, *et al*. Personal strategies to reduce the effects of landscape fire smoke on asthma-related outcomes: a protocol for systematic review and meta-analysis. *BMJ Open* 2024;**14**:e069516. doi:10.1136/bmjopen-2022-069516

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2022-069516>).

Received 24 October 2022  
Accepted 18 December 2023



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>School of Medicine and Public Health, University of Newcastle, Newcastle, NSW, Australia

<sup>2</sup>Asthma and Breathing Research Program, Hunter Medical Research Institute, Newcastle, NSW, Australia

<sup>3</sup>Department of Respiratory and Sleep Medicine, John Hunter Hospital, Newcastle, NSW, Australia

<sup>4</sup>School of Nursing and Midwifery, University of Newcastle, Newcastle, NSW, Australia

## Correspondence to

Prof Vanessa M McDonald;  
Vanessa.McDonald@newcastle.edu.au

## ABSTRACT

**Introduction** Landscape fire smoke (LFS) contains several hazardous air pollutants that are known to be detrimental to human health. People with asthma are more vulnerable to the health impact of LFS than general populations. The aim of this review is to investigate the effectiveness of personal strategies to reduce the effect of LFS on asthma-related outcomes.

**Methods and analysis** We will electronically search databases such as Medline, Embase, CINAHL and Cochrane Clinical Trials Register to identify eligible articles for the review. Screening of search results and data extraction from included studies will be completed by two independent reviewers. The risk of bias (RoB 2) will be assessed using the Risk of Bias Assessment Tool for Non-Randomised Studies for observational studies, the Cochrane Collaboration tool for assessing the RoB 2 for randomised controlled trials (RCTs) and the Risk Of Bias In Nonrandomized Studies of Interventions tool for non-RCTs. A random-effect meta-analysis will be performed to determine the pooled summary of findings of the included studies. If meta-analysis is not possible, we will conduct a narrative synthesis. Findings will be reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement.

**Ethics and dissemination** This study will synthesise the available evidence obtained from published studies and as such, no ethical approval is required. The review will be disseminated through peer-reviewed publications and conference presentations.

**PROSPERO registration number** CRD42022341120.

## INTRODUCTION

In recent years, unprecedented landscape fire (including bushfires, prescribed fires and uncontrolled wildfires, tropical deforestation fires, peat fires, agricultural burning and grass fires) events have been occurring more frequently around the world and with increased intensity and duration in many fire-prone regions.<sup>1 2</sup> Landscape fire smoke (LFS) contains several air pollutants such as particulate matter (PM), carbon monoxide, nitrogen dioxide, nitric oxide and volatile

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This study will include observational, randomised control trials and before and after studies from a comprehensive list of bibliographic databases, to summarise personal strategies to reduce the effects of landscape fire smoke (LFS) on asthma-related outcomes.
- ⇒ The review will provide updated information on the effectiveness of personal strategies to reduce the effects of LFS on asthma-related outcomes.
- ⇒ The methods will adhere to Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.
- ⇒ Language restriction to English may exclude additional studies published in other languages.

organic compounds that are known to cause adverse health effects.<sup>3 4</sup> A study of global mortality attributable to smoke from landscape fires estimated that 339 000 deaths annually are attributable to LFS.<sup>5</sup> In addition, some people, particularly those with asthma, are at higher risk of adverse health outcomes following exposure to fire smoke.<sup>6 7</sup> Asthma is a common illness worldwide with a global prevalence of approximately 262 million people.<sup>8</sup>

Epidemiological studies have found an association between exposure to LFS and increased respiratory morbidity, mortality and healthcare utilisation.<sup>9–11</sup> A systematic review and meta-analysis of 20 studies that investigated the relationship between landscape fire-related PM<sub>2.5</sub> and asthma-related outcomes reported that fire-related PM<sub>2.5</sub> increases hospital admission and emergency department visits for asthma.<sup>12</sup> A cross-sectional survey examining the impact of the 2019/2020 catastrophic bushfires on people with asthma in Australia found that people with asthma were more likely to report respiratory symptoms and healthcare service

utilisation than those without asthma.<sup>6</sup> Taken together, these findings show the significant health impact of LFS on people with asthma.

A study on the impact of prolonged bushfire smoke exposure in people with severe asthma reported that most people with severe asthma reported respiratory and non-respiratory symptoms during the bushfire period. This was despite most people with severe asthma implementing actions (such as staying indoors and avoiding going outdoors) to minimise or avoid exposure to bushfire smoke.<sup>10</sup> Similarly, a study on the impact of prolonged landscape fires on women with asthma found that even though most women with asthma took action to minimise exposure to landscape fire smoke, they experienced respiratory and non-respiratory symptoms during landscape fire period.<sup>13</sup> A study of the impact of bushfires on people with and without asthma revealed that those with asthma were more likely to take measures to minimise exposure to bushfire smoke than those without asthma.<sup>6</sup>

Governmental organisations have developed guidance about smoke mitigation strategies for the public to improve health outcomes.<sup>14</sup> However, there is a dearth of data regarding the effectiveness of personal strategies to reduce the effect of LFS on health outcomes, particularly for vulnerable populations.<sup>15 16</sup> A workshop convened by the Centre for Air pollution, energy and health Research underscored the need for assessing the efficacy of smoke mitigation strategies such as the use of air cleaners and making houses more airtight to reduce the adverse health outcomes.<sup>17</sup> Furthermore, a report by the European Respiratory Society highlighted the need for healthcare professionals to have information to advise patients about air pollution including fire smoke since they are facing daily concerns from patients, particularly vulnerable populations, about the effect that air pollution can have on their health.<sup>18</sup> Given that landscape fires are increasing in frequency and duration, it is imperative to examine personal strategies for people with asthma that are designed to reduce the effect of LFS on asthma-related outcomes. Furthermore, Vardoulakis *et al* suggested further research to understand if health advice during bushfire smoke can effectively lower the risk of adverse health outcomes.<sup>19</sup> A position statement about asthma and LFS highlighted the importance of improving health communication during fire period to lower health risks and preparation for future events.<sup>20</sup> To the best of our knowledge, no published systematic review has investigated the effectiveness of personal smoke mitigation strategies in reducing asthma related outcomes. Thus, the aim of this systematic review and meta-analysis is to synthesise evidence on personal strategies for people with asthma who were exposed to LFS to reduce the negative effect of LFS on asthma-related outcomes.

## Study question

We aim to answer the following question: for people with asthma exposed to LFS do personal smoke mitigation strategies improve asthma-related outcomes?

## METHODS AND ANALYSIS

### Reporting of the review findings

The protocol was developed in accordance with the Preferred Reporting Items for Systematic review and Meta-analysis Protocols (PRISMA-P) statement.<sup>21</sup> The protocol for this review has been registered with PROSPERO (CRD42022341120). The PRISMA (2020) statement will be used to report the findings.<sup>22</sup> This review will commence on the 15 October 2023 and end on the 25 January 2024.

### Data sources and search strategy

We developed the search strategy with the support of a librarian and other researchers and pilot tested it before the final search. A search will be conducted of the electronic Medline, Embase, CINAHL and Cochrane Clinical Trials Register databases to identify eligible articles for the review. We will use Medical Subject Headings (Mesh), keywords and free text search terms. The keywords include: (“Asthma” or “wheeze\*”) and (“bushfire” or “wildfire” or “wildland fire” or “landscape fire”) and (“mitigat\*” or “reduc\*” or “interven\*” or “avoid\*” or “personal protect\*”) and (“hospital\*” or “admission\*”, “emergenc\*”, “physicia\*”, “doctor”, “dispensatio\*”, “visit\*”, “medication”, “attendanc\*”) which will be used to identify all potential studies. The complete list of keywords is included in online supplemental table S1). The search strategy for Medline is supplemented with this protocol (online supplemental table S2). Reference lists of selected articles will be assessed to identify other possible studies of interest.

### Inclusion criteria

1. Studies involving people 7 years old or above with diagnosis of asthma or self-reported asthma.
2. People who were exposed to LFS.
3. Cross-sectional, cohort, case-control, randomised controlled trials (RCTs), before and after studies in which landscape fire, asthma-related outcomes and personal strategies to reduce LFS exposure were studied. For this review, landscape fires include vegetation fires (bushfires/wildfire, prescribed and cultural burns, forest, savanna, grassland and shrubland fires and agricultural burning), open coal mine and peat fires. To determine the reliability of LFS exposure assessment in each study, we will review the methodologies, measurement techniques and definitions employed.

### Exclusion criteria

1. Studies that recruited participants based on other respiratory conditions.
2. Studies published in languages other than English.

3. Other articles such as case reports, case series, letters, conference abstracts, letters and thesis.
4. Studies that examined other type of air pollution and asthma related outcomes.

## POPULATION, INTERVENTION, COMPARATOR AND OUTCOMES

### Population

People with asthma who were exposed to LFS. Asthma will be defined as doctor-diagnosed asthma fulfilling American Thoracic Society criteria or self-reported asthma.

### Interventions

Individual strategies aimed at minimising or avoiding personal exposure to LFS. The strategies could be physical (eg, wearing face masks), behavioural (eg, staying at home or avoiding outdoor physical activity, temporary relocation and nutrition), pharmacological (eg, use of medications and appropriate inhaler) and technological (eg, use of mobile technology, use of air conditioning or air purifiers). The intervention could involve a single strategy or multiple strategies.

### Comparator

People with asthma who were exposed to LFS and did not implement strategies to avoid/minimise personal exposure to LFS.

### Outcomes

Asthma-related outcomes include asthma symptoms, exacerbation (including hospitalisation, emergency department visits, unscheduled doctor visits and oral corticosteroid use), quality of life, asthma control, lung function (forced vital capacity and forced expiratory volume) and asthma medication use.

### Selection of studies

Studies obtained from databases will be exported to EndNote V.X9.1 citation manager and will then be exported to Covidence,<sup>23</sup> a software designed for conducting systematic reviews. Two independent reviewers (TB and PGG) will assess the studies, based on inclusion criteria. The title of the studies and abstracts will be screened to identify for full-text review. Then, full texts of potentially eligible studies will be assessed against eligibility criteria. Only studies that are approved by both reviewers will be included in the review. Any disagreements will be resolved by discussion or by consultation with a third reviewer (VMcD). Studies excluded from the review will be recorded along with the reason for their exclusion at the full-text screening stage. A final list of studies will be prepared for data extraction.

### Data extraction

Two independent reviewers (TB) will complete the data extraction form, using Covidence data extraction template with clear inclusion and exclusion criteria. Periodic meetings will be held to ensure consistency

in data extraction. A third reviewer (VMcD) will act as mediator in the event of disagreement. For each study, data recorded will include the first author's last name, year of publication, study setting (including country), study design, study period, sample size, response rate, population, personal smoke mitigation strategies, comparison groups, exposure and outcome definitions, and effect estimates. The corresponding authors of the studies included will be contacted to obtain missing information.

### Risk of bias assessment

Two review authors (TB) will assess the risk of bias independently for each study using the Risk of Bias Assessment Tool for Non-Randomized Studies (RoBANS)<sup>24</sup> for observational studies, Cochrane Collaboration tool for assessing the risk of bias<sup>25</sup> for RCTs and the Risk Of Bias In Nonrandomized Studies of Interventions tool for non-RCTs.<sup>26</sup> RoBANS consists of six domains: selection bias, confounding bias, performance bias, detection bias, attrition bias and reporting bias. Each domain will be allocated one of three possible categories for each study: 'low risk', 'high risk' and 'unclear'. The ROB classes for the RCTs were 'low risk of bias', 'some concerns,' or 'high risk of bias'. Any discrepancies arising during assessment will be resolved through discussion. Certainty of evidence will be assessed using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) framework adapted to the environment and occupational health.<sup>27</sup> The GRADE framework uses a four-level scale (ie, high, moderate, low and very low certainty) to illustrate the strength of the evidence for each outcome.

### Data synthesis and analysis

We will perform a narrative description of the studies included, including the study population, the personal strategies and the outcomes. We will use tables and figures to summarise the selected studies and results. R V.4.0.3 (Vienna, Austria) will be used for data entry and statistical analysis.<sup>28</sup> A random-effect meta-analysis will be performed to determine the pooled summary of findings of the included studies, where possible. The impact of individual strategies on the outcome of interest will be evaluated to compare the various personal strategies. This includes analysing data to assess how each strategy influences the outcomes' occurrence, allowing for a direct comparison of their effectiveness. To assess heterogeneity among studies, we will calculate the  $I^2$  and  $\tau^2$  statistic,<sup>29</sup> which describes the percentage of total variation among studies due to heterogeneity rather than chance. Sensitivity analysis will be conducted to assess the stability of the results and to test whether any individual study influences the meta-analysis. Where possible, subgroup analysis will be conducted based on severity of asthma, type and

duration of strategies (long-term vs short-term strategies), duration of exposures to LFS, country and study design. Potential publication bias will be examined by visual inspection of funnel plot asymmetry and Egger's test. If meta-analysis is not possible, we will conduct a narrative synthesis.

### Patient and public involvement

A consumer reference group was conducted and people with asthma identified the need to effective strategies to reduce the effects of LFS.

### DISCUSSION

To our knowledge, this systematic review and meta-analysis will be the first to synthesise literature on personal strategies for people with asthma to reduce the effects of LFS on asthma-related outcomes. This review will provide strong evidence on the effectiveness of personal strategies for people with asthma to minimise asthma-related outcomes. Furthermore, the findings will also provide information for healthcare providers to advise people with asthma on personal strategies, which have potential to improve asthma-related outcomes.

### ETHICS AND DISSEMINATION PLAN

Ethical approval is not required for the proposed review. Findings of this systematic review and meta-analysis will be published in a peer-reviewed journal and presented at conferences.

**Twitter** Tesfalidet Beyene @Tesfalidetlove <sup>ORCID</sup>, Megan E Jensen @DrMeganEJensen and Vanessa M McDonald @nessmcd

**Contributors** All authors were involved in the conception of the study question. TB developed the search strategy, wrote and prepared the protocol. PGG, VMur, MEJ and VMcD revised the protocol. All authors read and revised the protocol and consented to the publication of the article.

**Funding** This work is supported by the Australian Government Department of Health and Aged Care Medical Research Futures Fund

**Competing interests** None declared.

**Patient and public involvement** A consumer reference group was conducted and people with asthma identified the need to effective strategies to reduce the effects of LFS.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iDs

Tesfalidet Beyene <http://orcid.org/0000-0003-0774-7708>

VE Murphy <http://orcid.org/0000-0003-3282-1324>

Vanessa M McDonald <http://orcid.org/0000-0001-9890-3408>

### REFERENCES

- Xu R, Yu P, Abramson MJ, et al. Wildfires, Global Climate Change, and Human Health. *N Engl J Med* 2020;383:2173–81.
- Johnston FH, Borchers-Arriagada N, Morgan GG, et al. Unprecedented health costs of smoke-related PM<sub>2.5</sub> from the 2019–20 Australian megafires. *Nat Sustain* 2021;4:42–7.
- Urbanski SP, Hao WM, Baker S. Chemical composition of wildland fire emissions. *Dev Environ Sci* 2009;8:79–108.
- Reid CE, Brauer M, Johnston FH, et al. Critical Review of Health Impacts of Wildfire Smoke Exposure. *Environ Health Perspect* 2016;124:1334–43.
- Johnston FH, Henderson SB, Chen Y, et al. Estimated global mortality attributable to smoke from landscape fires. *Environ Health Perspect* 2012;120:695–701.
- Bui D, Davis S, Flynn A, et al. Impact of recent catastrophic bushfires on people with asthma in Australia: Health, social and financial burdens. *Respirology* 2021;26:296–7.
- MacIntyre CR, Nguyen P-Y, Trent M, et al. Adverse Health Effects in People with and without Preexisting Respiratory Conditions during Bushfire Smoke Exposure in the 2019/2020 Australian Summer. *Am J Respir Crit Care Med* 2021;204:368–71.
- Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet* 2020;396:1204–22.
- Chen G, Guo Y, Yue X, et al. Mortality risk attributable to wildfire-related PM<sub>2.5</sub> pollution: a global time series study in 749 locations. *Lancet Planet Health* 2021;5:e579–87.
- Beyene T, Harvey ES, Van Buskirk J, et al. 'Breathing Fire': impact of prolonged bushfire smoke exposure in people with severe asthma. *Int J Environ Res Public Health* 2022;19:7419.
- Reid CE, Jerrett M, Tager IB, et al. Differential respiratory health effects from the 2008 northern California wildfires: a spatiotemporal approach. *Environ Res* 2016;150:227–35.
- Borchers Arriagada N, Horsley JA, Palmer AJ, et al. Association between fire smoke fine particulate matter and asthma-related outcomes: systematic review and meta-analysis. *Environ Res* 2019;179:108777.
- Beyene T, Murphy VE, Gibson PG, et al. The impact of prolonged landscape fire smoke exposure on women with asthma in Australia. *BMC Pregnancy Childbirth* 2022;22:919.
- US Environmental Protection Agency. Wildfire smoke: a guide for public health officials. 2019. Available: <https://www.epa.gov/> [Accessed 3 Aug 2022].
- Fish JA, Peters MDJ, Ramsey I, et al. Effectiveness of public health messaging and communication channels during smoke events: a rapid systematic review. *J Environ Manage* 2017;193:247–56.
- Van Deventer D, Marecaux J, Doubleday A, et al. Wildfire smoke risk communication efficacy: a content analysis of Washington state's 2018 statewide smoke event public health messaging. *J Public Health Manag Pract* 2021;27:607–14.
- Cowie CT, Wheeler AJ, Tripovich JS, et al. Policy implications for protecting health from the hazards of fire smoke. a panel discussion report from the workshop landscape fire smoke: protecting health in an Era of escalating fire risk. *Int J Environ Res Public Health* 2021;18:5702:11..
- Powell P, Brunekreef B, Grigg J. How do you explain the risk of air pollution to your patients? *Breathe* 2016;12:201–3.
- Vardoulakis S, Jalaludin BB, Morgan GG, et al. Bushfire smoke: urgent need for a national health protection strategy. *Med J Aust* 2020;212:349–353.
- McDonald VM, Archbold G, Beyene T, et al. Asthma and landscape fire smoke: a thoracic society of Australia and New Zealand position statement. *Respirology* 2023;28:1023–35.
- Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
- Covidence systematic review software, Veritas health innovation. Melbourne, Australia. n.d. Available: [www.covidence.org](http://www.covidence.org)

- 24 Kim SY, Park JE, Lee YJ, *et al.* Testing a tool for assessing the risk of bias for nonrandomized studies showed moderate reliability and promising validity. *J Clin Epidemiol* 2013;66:408–14.
- 25 Sterne JAC, Savović J, Page MJ, *et al.* RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366:l4898.
- 26 Sterne JA, Hernán MA, Reeves BC, *et al.* ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355:i4919.
- 27 Morgan RL, Thayer KA, Bero L, *et al.* GRADE: assessing the quality of evidence in environmental and occupational health. *Environ Int* 2016;92–93:611–6.
- 28 R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing, 2020.
- 29 Borenstein M, Higgins JP, Hedges LV, *et al.* Basics of meta-analysis:  $I^2$  is not an absolute measure of heterogeneity. *Res Synth Methods* 2017;8:5–18.

**Table S1 Search term combination through Boolean operators**

<b>Population</b>		<b>Exposure</b>		<b>Intervention</b>		<b>Outcomes</b>
“Asthma” or “wheez*”	<b>AND</b>	“bushfire” or “wildfire” or “wildland fire” or “landscape fire” or “open fire” or “forest fire” or “coal mine fire” or “peat fire” or “crop burn*” or “stubble burn*” or “habitat fire” or “agricultur* fire” or “grass fire” or “savanna* fire” or “prescribed burn*” or “prescribed fire” or “vegetation fire” or “hazard reduction burn*” or “planned burn*” or “fuel reduction burn*” or “controlled burn*” or “cool burn*”	<b>AND</b>	“mitigat*” or “minimi*” or “reduc*” or “interven*” or “avoid*” or “lower*” or “reduction behavior” or “behavioral change” or “personal protect*” or “Staying indoors” or “indoor*” or “exercise” or “physical activity*” or “masks” or “facemask*” or “respirator*” or “monitor* or “smartphone* or “phone* or “app” or “air purifier” or “air filter” or “portable” or “nasal filter” or “air cleaner” or “ventilation” or “air conditioner*” or “forecast” or “air quality index” or “diet*” or “vitamin*” or “Antioxidant*” or “supplement*” or “aspirin” or “statin*”	<b>AND</b>	symptom*” or “quality of Life” or “asthma control”, or “exacerbation” or “lung function” or “medication” or “hospital*” or “admission*”, “emergenc*”, “physicia*” or “doctor”, “dispensation”, or “visit*” or “attendance*” or “asthma-related outcomes”

**Table S2 OVID Medline search strategy**

#	Searches	Results
1	Asthma/ or Asthma.mp.	191661
2	wheez*.mp.	15194
3	bushfire.mp.	266
4	wildfire.mp. or Wildfires/	2494
5	wildland fire.mp. or Wildfires/	1085
6	landscape fire.mp.	49
7	open fire.mp.	156
8	forest fire.mp. or Wildfires/	1465
9	coal mine fire.mp.	39
10	peat fire.mp.	31
11	crop burn*.mp.	24
12	stubble burn.mp. or Wildfires/	938
13	habitat fire.mp.	4
14	agricultur* fire.mp.	13
15	grass fire.mp.	15
16	savanna* fire.mp.	27
17	prescribed burn*.mp.	356
18	prescribed fire.mp.	290
19	vegetation fire.mp.	50
20	hazard reduction burn*.mp.	5
21	planned burn.mp.	8
22	fuel reduction burn*.mp.	8
23	controlled burn*.mp.	107
24	cool burn*.mp.	5
25	mitigat*.mp.	127977
26	minimi*.mp.	235519
27	reduc*.mp.	4071848
28	interven*.mp.	1318322
29	avoid*.mp.	460695
30	lower*.mp.	2319407
31	reduction behavior.mp.	15059
32	behavioral change.mp.	4305
33	Staying indoors.mp.	40

34	indoor*.mp.	42530
35	Exercise/ or exercise.mp.	406296
36	physical activity*.mp.	136933
37	Masks/ or masks.mp.	23357
38	facemask*.mp.	2000
39	respirator*.mp. or N95 Respirators/	670633
40	monitor*.mp.	1121827
41	Smartphone/ or Mobile Applications/ or smartphone*.mp. or Cell Phone/	35135
42	phone*.mp.	63548
43	app.mp.	35425
44	air purifier.mp. or Air Filters/	656
45	air filter.mp. or Air Filters/	992
46	portable.mp.	36806
47	nasal filter.mp.	21
48	air cleaner.mp.	154
49	ventilation.mp. or Ventilation/	159522
50	air conditioner*.mp. or Air Conditioning/	3234
51	forecast.mp. or Forecasting/	100100
52	air quality index.mp.	652
53	Diet/ or diet*.mp.	847724
54	vitamin*.mp.	295675
55	Antioxidants/ or Antioxidant*.mp.	287034
56	Dietary Supplements/ or supplement*.mp.	407359
57	aspirin.mp. or Aspirin/	72031
58	statin*.mp.	49907
59	symptom*.mp.	1357888
60	quality of Life.mp. or "Quality of Life"/	418624
61	asthma control.mp.	7578
62	exacerbation.mp.	42714
63	lung function.mp.	38975
64	medication.mp.	286961
65	Hospitalization/ or hospital*.mp.	384049
66	hospital.mp. or Hospitals/	1468633
67	admission*.mp.	279646
68	emergenc*.mp.	529135



69	Physicians/ or physicia*.mp.	626158
70	doctor.mp.	58354
71	dispensation.mp.	1155
72	Emergency Service, Hospital/ or visit*.mp. or Office Visits/	353667
73	attendance*.mp.	32712
74	asthma-related outcomes.mp.	174
75	1 or 2	197314
76	3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24	4073
77	25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 or 35 or 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 or 52 or 53 or 54 or 55 or 56 or 57 or 58	9741843
78	59 or 60 or 61 or 62 or 63 or 64 or 65 or 66 or 67 or 68 or 69 or 70 or 71 or 72 or 73 or 74	4382933
79	75 and 76 and 77 and 78	82