

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

The global contribution of suicide to maternal mortality: a systematic review protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2024-087669
Article Type:	Protocol
Date Submitted by the Author:	16-Apr-2024
Complete List of Authors:	Simmons, Emma; University of Oxford, Medical Sciences Division Gong, Jenny; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health Quigley, Maria; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health Alderdice, Fiona; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health; Queen's University Belfast, School of Nursing and Midwifery Harrison, Sian; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health Fellmeth, Gracia; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health
Keywords:	Systematic Review, Suicide & self-harm < PSYCHIATRY, Maternal medicine < OBSTETRICS

SCHOLARONE™
Manuscripts

Title: The global contribution of suicide to maternal mortality: a systematic review protocol

Authors

Emma Simmons¹, Jenny Gong², Maria Quigley², Fiona Alderdice^{2,3}, Siân Harrison², Gracia Fellmeth²

Affiliations

1. Medical Sciences Division, University of Oxford, Oxford, UK
2. National Perinatal Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK
3. School of Nursing and Midwifery, Queen's University Belfast, Belfast, UK

Contact emails

emma.simmons@pmb.ox.ac.uk

jenny.gong@ndph.ox.ac.uk

maria.quigley@npeu.ox.ac.uk

fiona.alderdice@npeu.ox.ac.uk

sian.harrison@npeu.ox.ac.uk

gracia.fellmeth@ndph.ox.ac.uk

Corresponding author

Gracia Fellmeth

National Perinatal Epidemiology Unit (NPEU)

Nuffield Department of Population Health

University of Oxford

Old Road Campus

Roosevelt Drive

Oxford OX3 7LF

United Kingdom

Tel. 01865 289745

Email: gracia.fellmeth@ndph.ox.ac.uk

Keywords

Suicide; maternal; perinatal; systematic review; global

Word count

2592

1
2
3 **TITLE:** The global contribution of suicide to maternal mortality: a systematic review protocol
4
5

6 **ABSTRACT**

7
8 **Introduction:** Maternal suicide is a significant contributor to maternal mortality with
9 devastating consequences for women, families and society. Maternal mortality reporting
10 systems differ across countries and there is no up-to-date overview of maternal suicide
11 deaths globally. This systematic review aims to synthesise the evidence on maternal suicide.
12 The primary objective is to determine the contribution of suicide towards maternal
13 mortality globally and explore differences between geographical regions. The secondary
14 objectives are to summarise the availability and quality of data globally and to describe how
15 suicide deaths are classified across different countries.
16
17

18
19 **Methods and analysis:** This protocol follows the Preferred Reporting Items for Systematic
20 Reviews and Meta-analyses for systematic review protocols (PRISMA-P) guidelines. Medline,
21 Embase, PsychINFO, Global Health and CINAHL databases and the grey literature were
22 searched with no date or language restrictions. Observational studies, national surveys and
23 reports that present data on maternal deaths due to suicide occurring during pregnancy,
24 intrapartum and in the post-partum period will be included. Screening, data extraction and
25 quality assessment will be conducted independently by two reviewers. Results will be
26 summarised narratively. If sufficient outcome data is available, random-effects meta-
27 analyses will be conducted to determine global pooled estimates of suicide-related
28 maternal mortality rates and the proportion of maternal deaths attributable to suicide.
29
30
31

32 **Ethics and dissemination:** Ethical approval is not required for this systematic review. Results
33 will be written up for publication in a peer-reviewed journal and findings will be shared at
34 national and international conferences.
35
36

37 **Prospero registration number:** CRD42023429072
38
39

40 **ARTICLE SUMMARY**

41 **Strengths and limitations of this study**

- 42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- This will be the first systematic review of the contribution of suicide to maternal deaths globally.
 - We will follow rigorous methods and report our methodology and results in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.
 - The search strategy is highly comprehensive and was developed with an experienced university librarian.
 - We will conduct the search without language or date limitations, allowing us to include non-English publications which are often omitted in systematic reviews.
 - We will conduct an extensive grey literature search to identify non-indexed and non-academic publications including maternal mortality databases and surveillance reports.
 - A limitation of our review is that we will exclude maternal deaths due to injury or accidents.

INTRODUCTION

Rationale

Maternal suicide, defined as death by suicide occurring during pregnancy or within one year of giving birth, is a significant contributor to maternal mortality with devastating consequences for women, families and society. Across a number of high-income countries including the United Kingdom and the USA, suicide is one of the leading causes of maternal death.^{1,2} In the UK, the most recent data suggests that between 2019-2021 almost 40% of maternal deaths between six weeks and twelve months after childbirth were due to suicide.¹ In lower-resource settings, where rates of all-cause maternal mortality are often higher overall, the contribution of suicide to maternal deaths is less clear. A previous review reported a maternal suicide prevalence of 1% across low- and middle-income countries (LMIC) with a wide range in prevalence between regions.³

Understanding the magnitude of suicide-related maternal mortality is important to improve assessment, identification and reporting of maternal suicide deaths and inform prevention strategies. However, there are a number of challenges in accurately estimating the burden. Firstly, there are a number of different definitions of maternal death (**Figure 1**). The World Health Organization (WHO) uses the terms *maternal death* and *late maternal death*, which are defined in the International Classification of Diseases (11th Revision; ICD-11) and which differ from each other only in terms of the time periods they cover.^{4,5} Maternal deaths and late maternal deaths are combined in the ICD-11 under a new grouping of *comprehensive maternal deaths*.⁴ Others, notably the US Centres for Disease Control and Prevention (CDC), refer to *pregnancy-associated* and *pregnancy-related deaths*, which both span the same time period but differ according to the designated cause of death.⁶ Confusingly, the WHO also refers to *pregnancy-related death*, but defines this term differently from the CDC both in terms of the causes included and the time period covered.⁵ In reports and publications presenting maternal death statistics, there is often a lack of clarity around which definition was applied.

Figure 1. Definitions of maternal death
[INSERT Figure 1 here]

Secondly, there are differences in the denominator used to calculate rates of maternal mortality. The commonly reported maternal mortality ratio (MMR) reports the number of maternal deaths during a given time period per 100,000 live births during the same time period.⁵ The use of live births as a denominator means that deaths occurring during pregnancy and stillbirths are not captured by the MMR. To address this, some countries such as the UK also report the number of maternal deaths per 100,000 maternities, defined as the number of women giving birth, including stillbirth, at or beyond 24 weeks' gestation.¹ When maternities are used as the denominator, the statistic is often referred to as a maternal mortality *rate* (rather than ratio), but the terminology is inconsistent and in practice the distinction between MMR and maternal mortality rates is often unclear. These differences create challenges when comparing statistics between countries or across different reporting systems.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Thirdly, there is variation in the availability and quality across different regions of the world. High quality data on maternal suicide requires an accurate identification of all maternal deaths, as well as an accurate designation of cause of death. Some countries have systems in place to identify and investigate maternal deaths and their causes at a regional or national level. Such systems include systematic maternal mortality surveillance systems, confidential enquiries and Reproductive Age Mortality Surveys (RAMOS). The assignment of cause of death can include the use of formal death registration certificates and verbal autopsies in which family members are asked to provide information on circumstances around the death. In countries with less robust death reporting systems it is difficult to determine how accurate maternal death reporting is.

A final challenge is the misclassification of deaths by suicide to other causes such as injury or accidents. This is thought to be common given the stigma associated with suicide and its criminalisation in some societies.³ A review of maternal suicide in LMIC found that reclassifying the leading suicide methods from injuries to suicide increased the pooled prevalence of pregnancy-related suicide deaths from 1.00 to 1.68.³ Misclassification is compounded by the fact that until the most recent revision of the ICD, maternal suicide did not have a specific code. There has also been a change in the recommended classification of maternal suicides. The ICD-10 categorised maternal deaths due to injuries or mental disorders as indirect maternal deaths.⁷ The WHO subsequently published a derivation of the ICD-10: this ICD-Maternal Mortality (ICD-MM) calls for maternal suicides to be classified as direct maternal deaths.⁸ This discrepancy has led to variations in the reporting of maternal suicide as direct or indirect causes of death, with some statistics reporting maternal suicide as an incidental cause of death and other statistics excluding suicides from their maternal mortality data altogether.⁹

There are several existing systematic reviews of maternal suicide but these are limited to specific perinatal populations or geographical regions, and there is no up-to-date synthesis of the global evidence for all perinatal women. Lindahl *et al.* (2005) reviewed prevalence of suicide ideation, attempts and deaths in pregnancy and the post-partum period globally; however this included studies published up to 2002 and their findings need updating.¹⁰ Fuhr *et al.* (2014) assessed the contribution of suicide and injuries to perinatal mortality in LMIC only.³ Amiri *et al.* (2021) assessed rates of suicide ideation, attempts and mortality globally among post-partum women only.¹¹ Chin *et al.* (2022) reviewed prevalence, risk factors, outcomes and interventions for perinatal deaths by suicide but focused on a limited number of studies that were not included in an earlier review by Mangla *et al.* (2019).² Chin *et al.* (2022) also restricted their search to studies published in English, and most of the identified studies were conducted in the USA.²

Our systematic review expands upon previous reviews by including suicides occurring during pregnancy and the post-partum period in all countries globally with no language or date restrictions. Results will enable an estimation of the contribution of suicide to maternal deaths globally and a comparison of the burden across geographical regions. Identifying countries reporting either very high or very low rates will inform further work to determine whether differences in rates reflect true differences or reporting- or coding-related factors. Our review will also show whether maternal suicide deaths are reported as direct, indirect or incidental maternal deaths, informing a possible need for further clarification of the ICD-

1
2
3 11 and ICD-MM classifications. Finally, our review will identify countries for which no data
4 on maternal suicide is available, an important requirement in meeting the United Nations'
5 Sustainable Development Goal (SDG) 3.¹²
6
7

8 **Objectives**

9

10 This systematic review aims to identify and synthesise evidence on maternal suicide deaths
11 globally. The primary objective is to determine the contribution of suicide towards maternal
12 mortality and explore differences between geographical regions. The secondary objectives
13 are to summarise the availability and quality of data globally and to describe how suicide
14 deaths are classified across different countries.
15
16

17 **METHODS AND ANALYSIS**

18

19 **Registration and protocol adherence**

20

21 This systematic review protocol is reported in adherence with the Preferred Reporting Items
22 for Systematic Reviews and Meta-Analyses for systematic review protocols (PRISMA-P)
23 guidelines (**Supplementary Information S1**).¹³ The review was registered on PROSPERO
24 (CRD42023429072) on 25 May 2023. If important amendments to the protocol are required
25 during the course of the review, these will be recorded as amendments on PROSPERO and
26 reported in the final publication as a deviation from the protocol.
27
28
29

30 **Definitions**

31

32 For the purposes of this review, we define maternal suicide as the death of a woman by
33 suicide at any time during pregnancy, intrapartum or within one year of giving birth. The
34 perinatal period is the period of pregnancy and up to one year after giving birth.
35
36
37

38 **Eligibility criteria**

39

40 *Population, intervention, comparison and outcome framework*

41 The population of interest is women at any stage of pregnancy or within one year of giving
42 birth. The outcome of interest is death by suicide during the perinatal period. There is no
43 intervention or control for this systematic review of global maternal suicide rates.
44
45

46 *Inclusion and exclusion criteria*

47 We will include any study that reports data on the number of maternal deaths due to
48 suicide along with appropriate denominator data (either total live births or total maternities
49 during the same time period as maternal deaths). Studies reporting the proportion of all
50 maternal deaths that were due to suicide will also be included. We will include cohort,
51 cross-sectional and case control study designs as well as data from national surveys,
52 databases and reports such as mortality statistics and confidential enquiries. Qualitative
53 studies will be excluded as they are unlikely to contain numerator and denominator data on
54 maternal suicide and maternal death. There will be no minimum threshold applied to the
55 sample size or numbers of suicides reported. Studies that do not report data on the
56 denominator to allow calculation of the MMR attributable to suicide or the proportion of
57
58
59
60

deaths that are due to suicide will be excluded. Studies in which suicide deaths cannot be differentiated from deaths from injury or other causes will also be excluded. We will include studies published in any language.

Patient and public involvement

There was no patient or public involvement in the design of this systematic review.

Search strategy and data sources

Medline, Embase, PsychINFO, Global Health and CINAHL databases were searched with no date or language restrictions on 5 May 2023. A search strategy using terms relevant to perinatal status, mortality and suicide was developed in collaboration with a university research librarian and adapted for each database. The search strategy used for Medline is shown in **Table 1**. In order to capture relevant evidence not published in the databases above, we conducted an extensive grey literature search. This involved searches of *Google* and *Google Scholar* using the relevant terms listed above to capture non-academic articles such as reports and statistical databases. We also searched the websites of the World Health Organization (WHO), the United Nations Children's Fund (UNICEF), the United Nations Population Fund (UNFPA), UN Women, the World Bank and the Global Burden of Disease Study to identify relevant reports. All records identified by the database search and grey literature searches were combined and duplicates removed using *Endnote*.¹⁴ De-duplicated references were imported into *Covidence* for screening, data extraction and quality assessment.¹⁵

Table 1. Search strategy for Medline

1	Pregnancy/ or Pregnant Women/
2	Postnatal Care/ or Postpartum Period/
3	1 or 2
4	Mortality/
5	3 and 4
6	maternal mortality/ or maternal death/
7	((pregnan* or maternal or postnatal or post-natal or postpart* or post-part*) and (death? or mortality or fatalit*)).ti.
8	((pregnan* or maternal or postnatal or post-natal or postpart* or post-part*) adj3 (death? or mortality or fatalit*)).ti,ab,kf.
9	5 or 6 or 7 or 8
10	"cause of death"/
11	Death Certificates/
12	exp suicide/
13	"cause of death".ti,ab,kf.
14	(suicid* or overdos* or self harm or self injury or violen*).ti,ab,kf.
15	10 or 11 or 12 or 13 or 14
16	9 and 15

Data extraction

1
2
3 Screening of titles and abstracts was performed independently by two reviewers (ES and JG)
4 to exclude studies which did not meet inclusion criteria. Any disagreements were discussed
5 and resolved with a third reviewer (SH and GF). Full-texts were retrieved. Any missing full
6 texts were searched for using a university library database, online journal archives, *Google*
7 and by contacting study authors. Any full-texts that could not be retrieved after this search
8 process were excluded. Full-texts of remaining studies were screened independently by two
9 reviewers (ES, JG, SH, GF) and any disagreements were resolved through discussion.
10 Reasons for excluding full-texts were recorded. Two reviewers (ES, JG, SH, GF) will
11 independently extract data from included studies using a standardised and piloted data
12 extraction form consisting of the following fields:
13
14
15

- 16 • **Publication details:** lead author, publication year, publication language
- 17 • **Setting:** country, region, type of setting (e.g. hospital or community deaths), dates
18 covered, whether data was collected at national or regional level
- 19 • **Participants:** perinatal status (pregnant or post-partum), definition of maternal
20 death used, age, any sociodemographic characteristics (e.g. socioeconomic
21 background, ethnicity, income, parity)
- 22 • **Methods of assessment:** method of identifying maternal deaths, method of
23 ascertaining cause of death, classification of suicide (as indirect, direct or incidental
24 cause of death)
- 25 • **Outcomes:** total maternal deaths, total live births, total maternities, number of
26 suicide deaths, proportion of deaths due to suicide, accuracy measures (e.g.
27 standard deviation, p-values or 95% confidence intervals), mode of suicide
28
29
30
31
32

33 If reported data is unclear or insufficient, authors of the study will be contacted to request
34 the additional data required for inclusion in the review.
35

36 **Quality assessment and risk of bias**

37
38 Two reviewers (ES, JG, SH, GF) will independently assess the quality of each included study.
39 We will use criteria adapted from a study by Grollman and Ronsmans¹² to assess quality and
40 risk of bias. This will include assessment of three domains: quality of the method of death
41 ascertainment; completeness of the cause of death assignation; and quality of the method
42 of assigning cause of death. Each study will be assigned a rating of low, medium or high
43 quality in each of these categories using the attributes outlined by Grollman and
44 Ronsmans.¹⁶ Fuhr *et al.* (2014) used an adapted version of these criteria, with only two
45 categories: quality of the method of death ascertainment and completeness of the cause of
46 death assignation.³ We have chosen this method as there is a precedent for its use in a
47 similar review, and other risk of bias assessment tools have limited application to studies
48 looking at causes of death. Any disagreements will be discussed with the other review
49 authors (FA, MQ) until a consensus is reached.
50
51
52
53

54 **Data synthesis**

55
56 A narrative synthesis of included studies will be conducted and a table of study
57 characteristics will be compiled to provide an overview of all included studies. Countries for
58 which data on maternal suicide deaths is available will be mapped and colour-coded
59
60

1
2
3 (red/amber/green) according to the quality of evidence. The proportion of countries
4 classifying maternal suicide deaths as direct, indirect or incidental deaths will also be
5 reported. If sufficient data is available, random-effects meta-analyses will be conducted to
6 generate pooled estimates of suicide-related maternal mortality rates. Depending on the
7 data identified and comparability between studies, we will use either live births or
8 maternities as the denominator or conduct a separate pooled estimate for each. Forest
9 plots will be used to present data visually. Statistical heterogeneity between studies will be
10 calculated using the I^2 statistic. If data allows and sufficient studies are identified, we will
11 carry out subgroup analyses to explore differences in suicide death rates according to
12 perinatal status (pregnancy vs. post-partum); country income classification (high-income vs.
13 low- and middle-income countries); geographical region (using WHO geographical regions);
14 and study quality. The presence of publication bias will be assessed using funnel plots.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ETHICS AND DISSEMINATION

No primary data is being collected for this study therefore ethical approval is not required. Results of this study will be published in a peer-reviewed publication and presented at conferences. Raw data will be published as supplementary material in the final publication.

DISCUSSION

This systematic review and meta-analysis will produce a synthesis of the available evidence on maternal suicide globally. This will allow comparisons to be made between countries on suicide-related maternal mortality rates and the overall contribution of suicide toward maternal mortality. Summarising the availability and quality of data available in different regions of the world will identify priorities for future research. Identifying countries with high or low maternal suicide rates will also inform future studies of risk factors and preventive interventions in different populations.

There are a number of limitations to this study. It is possible that some of the data retrieved will be of poor quality, limiting its reliability and making comparison between areas challenging. The wide variety of methods and definitions used in studies identifying maternal suicide may also make it difficult to draw direct comparisons, and it may not be possible to calculate pooled estimates. By excluding maternal deaths classified as accidental or injury-related we may be under-estimating the true burden of suicide-related maternal mortality. The lack of data in some regions could also limit the power of analyses. There has also been a difficulty in acquiring the full texts of several of the included abstracts. Every effort was made to identify these texts, however a significant number of texts had to be excluded. This could impact whether we have captured the full breadth of evidence on this topic.

Acknowledgements We would like to thank Ms. Nia Roberts, Senior Outreach Librarian at the University of Oxford Bodleian Health Care Libraries, for her guidance and support in the development of the search strategy for this review.

Data statement No data was gathered for this protocol.

1
2
3 **Author contributions** ES, SH and GF developed the protocol. ES developed the search
4 strategy with input from SH and GF. ES, JG, SH and GF contributed to screening. ES, JG, SH
5 and GF will conduct data extraction and quality assessment. MQ and FA contributed to the
6 analysis plan. All authors contributed to the writing of this manuscript. SH and GF act as
7 guarantors for the review.
8
9

10 **Funding statement:** GF is funded by a Clinical Research Fellowship from the Nuffield
11 Department of Population Health, University of Oxford. The funder had no role in the design
12 or development of this protocol.
13
14

15 **Competing interests statement:** All authors declare no competing interests.
16
17

18 REFERENCES

- 19
20
21 1. Knight M, Bunch K, Felker A, Patel R, Kotnis R, Kenyon S, Kurinczuk JJ (Eds.) on behalf of
22 MBRRACE-UK. Saving Lives, Improving Mothers' Care Core Report - Lessons learned to
23 inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths
24 and Morbidity 2019-21. Oxford: National Perinatal Epidemiology Unit, University of Oxford
25 2023.
26
27 2. Chin K, Wendt A, Bennett IM, Bhat A. Suicide and Maternal Mortality. *Curr Psychiatry Rep*
28 2022;24:239–275.
29
30 3. Fuhr DC, Calvert C, Ronsmans C, Chandra PS, Sikander S, De Silva MJ, Patel V.
31 Contribution of suicide and injuries to pregnancy-related mortality in low-income and
32 middle-income countries: a systematic review and meta-analysis. *Lancet Psychiatry*. 2014
33 Aug;1(3):213-25.
34
35 4. World Health Organization. International Statistical Classification of Diseases and Related
36 Health Problems (11th edition; ICD-11). 2021. Geneva: World Health Organization.
37
38 5. World Health Organization. Trends in maternal mortality 2000 to 2020: estimates by
39 WHO, UNICEF, UNFPA, World Bank Group and UNDESA/Population Division. 2023. Geneva:
40 World Health Organization.
41
42 6. Centers for Disease Control and Prevention. Pregnancy Mortality Surveillance System.
43 2024. Available at [www.cdc.gov/reproductivehealth/maternal-mortality/pregnancy-](http://www.cdc.gov/reproductivehealth/maternal-mortality/pregnancy-mortality-surveillance-system.htm)
44 [mortality-surveillance-system.htm](http://www.cdc.gov/reproductivehealth/maternal-mortality/pregnancy-mortality-surveillance-system.htm) (accessed 11 April 2024).
45
46 7. World Health Organization. International Statistical Classification of Diseases and Related
47 Health Problems (10th edition; ICD-10). 2016. Geneva: World Health Organization.
48
49 8. World Health Organization. The WHO Application of ICD-10 to Deaths during Pregnancy,
50 Childbirth and Puerperium: ICD-MM. 2012. Geneva: World Health Organization.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 9. Hasegawa J, Katsuragi S, Tanaka H, Kubo T, Sekizawa A, Ishiwata I, Ikeda T. How should
4 maternal death due to suicide be classified? Discrepancy between ICD-10 and ICD-MM.
5 BJOG. 2020 May;127(6):665-667.
6
7
- 8 10. Lindahl V, Pearson JL, Colpe L. Prevalence of suicidality during pregnancy and the
9 postpartum. Arch Womens Ment Health. 2005 Jun;8(2):77-87.
10
11
- 12 11. Amiri S, Behnezhad S. The global prevalence of postpartum suicidal ideation, suicide
13 attempts, and suicide mortality: A systematic review and meta-analysis. International
14 Journal of Mental Health. 2021;50:311–336.
15
16
- 17 12. United Nations. Transforming Our World: The 2030 Agenda for Sustainable
18 Development. 2015. Available at: <https://sdgs.un.org/2030agenda> (accessed 08/04/2024).
19
20
- 21 13. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart
22 LA. Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-
23 P) 2015 statement. Syst Rev. 2015;4(1):1.
24
25
- 26 14. The EndNote Team. EndNote (Version X9). 2013. Philadelphia, PA: Clarivate.
27
28
- 29 15. Covidence Systematic Review software. 2023. Melbourne: Veritas Health Innovation.
30
31
- 32 16. Grollman C, Ronsmans C. Systematic review of the proportion of pregnancy-related
33 deaths attributed to HIV in population-based studies in sub-Saharan Africa. Trop Med Int
34 Health. 2014 Jan;19(1):83-97.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Definitions of maternal death

- **Maternal death** is defined by the World Health Organization (WHO) as the death of a woman from direct or indirect obstetric causes while pregnant or within 42 days after the end of pregnancy from causes associated with, or exacerbated by, pregnancy or its management, irrespective of the duration or outcome of the pregnancy.⁵
- **Late maternal death** is defined by the WHO as a death of a woman from direct or indirect obstetric causes more than 42 days but less than one year after termination of pregnancy.⁵
- **Comprehensive maternal death** is defined by the International Classification of Diseases (11th Revision) WHO as maternal deaths and late maternal deaths combined.⁴
- **Pregnancy-related death** is defined by the WHO as a death of a woman while pregnant or within 42 days of termination of pregnancy, irrespective of the cause of death. This definition includes obstetric, non-obstetric, unintentional (accidental) and incidental causes of death.⁵
- **Pregnancy-related death** is defined by the US Centers for Disease Control and Prevention (CDC) as a death during or within one year of pregnancy from a pregnancy complication, a chain of events initiated by pregnancy or the aggravation of an unrelated condition by the physiological effects of pregnancy.⁶
- **Pregnancy-associated death** is defined by the CDC as a death during or within one year of pregnancy, regardless of the cause of death.⁶

PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol*

Section and topic	Item No	Checklist item	Page Reported
ADMINISTRATIVE INFORMATION			
Title:			
Identification	1a	Identify the report as a protocol of a systematic review	2
Update	1b	If the protocol is for an update of a previous systematic review, identify as such	N/A
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number	2, 3
Authors:			
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author	1
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review	9
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments	5
Support:			
Sources	5a	Indicate sources of financial or other support for the review	9
Sponsor	5b	Provide name for the review funder and/or sponsor	9
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol	9
INTRODUCTION			
Rationale	6	Describe the rationale for the review in the context of what is already known	3-5
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO)	5, 6
METHODS			
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review	6
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage	6
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated	Table 1

Study records:			
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review	6
Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)	7
Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators	7
Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications	7
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale	7
Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis	8
Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised	8
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I^2 , Kendall's τ)	8
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression)	8
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned	8
Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)	8
Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)	8

*** It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.

BMJ Open

The global contribution of suicide to maternal mortality: a systematic review protocol

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2024-087669.R1
Article Type:	Protocol
Date Submitted by the Author:	31-Jul-2024
Complete List of Authors:	Simmons, Emma; University of Oxford, Medical Sciences Division Gong, Jenny; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health Daskalopoulou, Zoe; University of Oxford, Nuffield Department of Population Health Quigley, Maria; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health Alderdice, Fiona; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health; Queen's University Belfast, School of Nursing and Midwifery Harrison, Sian; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health Fellmeth, Gracia; University of Oxford, National Perinatal Epidemiology Unit, Nuffield Department of Population Health
Primary Subject Heading:	Mental health
Secondary Subject Heading:	Obstetrics and gynaecology, Public health, Global health
Keywords:	Systematic Review, Maternal medicine < OBSTETRICS, Suicide & self-harm < PSYCHIATRY

SCHOLARONE™
Manuscripts

Title: The global contribution of suicide to maternal mortality: a systematic review protocol

Authors

Emma Simmons¹, Jenny Gong², Zoe Daskalopoulou², Maria Quigley², Fiona Alderdice^{2,3}, Siân Harrison², Gracia Fellmeth²

Affiliations

1. Medical Sciences Division, University of Oxford, Oxford, UK
2. National Perinatal Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK
3. School of Nursing and Midwifery, Queen's University Belfast, Belfast, UK

Contact emails

emma.simmons@pmb.ox.ac.uk

jenny.gong@ndph.ox.ac.uk

zoe.daskalopoulou@npeu.ox.ac.uk

maria.quigley@npeu.ox.ac.uk

fiona.alderdice@npeu.ox.ac.uk

sian.harrison@npeu.ox.ac.uk

gracia.fellmeth@ndph.ox.ac.uk

Corresponding author

Gracia Fellmeth

National Perinatal Epidemiology Unit (NPEU)

Nuffield Department of Population Health

University of Oxford

Old Road Campus

Roosevelt Drive

Oxford OX3 7LF

United Kingdom

Tel. 01865 289745

Email: gracia.fellmeth@ndph.ox.ac.uk

Keywords

Suicide; maternal; perinatal; systematic review; global

Word count

2877

1
2
3 **TITLE:** The global contribution of suicide to maternal mortality: a systematic review protocol
4
5

6 **ABSTRACT**

7
8 **Introduction:** Maternal suicide is a significant contributor to maternal mortality with
9 devastating consequences for women, families and society. Maternal mortality reporting
10 systems differ across countries and there is no up-to-date overview of maternal suicide
11 deaths globally. This systematic review aims to synthesise the evidence on maternal suicide.
12 The primary objective is to determine the contribution of suicide towards maternal
13 mortality globally and explore differences between geographical regions. The secondary
14 objectives are to summarise the availability and quality of data globally and to describe how
15 suicide deaths are classified across different countries.
16
17

18
19 **Methods and analysis:** This protocol follows the Preferred Reporting Items for Systematic
20 Reviews and Meta-analyses for systematic review protocols (PRISMA-P) guidelines. Medline,
21 Embase, PsychINFO, Global Health and CINAHL databases and the grey literature were
22 searched with no date or language restrictions. Observational studies, national surveys and
23 reports that present data on maternal deaths due to suicide occurring during pregnancy,
24 intrapartum and in the post-partum period will be included. Screening, data extraction and
25 quality assessment will be conducted independently by two reviewers. Results will be
26 summarised narratively. If sufficient outcome data is available, random-effects meta-
27 analyses will be conducted to determine global pooled estimates of suicide-related
28 maternal mortality rates and the proportion of maternal deaths attributable to suicide.
29
30
31

32 **Ethics and dissemination:** Ethical approval is not required for this systematic review. Results
33 will be written up for publication in a peer-reviewed journal and findings will be shared at
34 national and international conferences.
35
36

37 **Prospero registration number:** CRD42023429072
38
39

40 **ARTICLE SUMMARY**

41 **Strengths and limitations of this study**

- 42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- We will follow rigorous methods and report our methodology and results in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.
 - The search strategy is highly comprehensive and was developed with an experienced university librarian.
 - We will conduct the search without language or date limitations, allowing us to include non-English publications which are often omitted in systematic reviews.
 - We will conduct an extensive grey literature search to identify non-indexed and non-academic publications including maternal mortality databases and surveillance reports.
 - A limitation of our review is that we will exclude maternal deaths due to injury, accidents or substance use.

INTRODUCTION

Rationale

Maternal suicide, defined as death by suicide occurring during pregnancy or within one year of giving birth, is a significant contributor to maternal mortality with devastating consequences for women, families and society. Across a number of high-income countries including the United Kingdom and the USA, suicide is one of the leading causes of maternal death.^{1,2} In the UK, the most recent data suggests that between 2019-2021 almost 40% of maternal deaths between six weeks and twelve months after childbirth were due to suicide.¹ In lower-resource settings, where rates of all-cause maternal mortality are often higher overall, the contribution of suicide to maternal deaths is less clear. A previous review reported a maternal suicide prevalence of 1% across low- and middle-income countries (LMIC) with a wide range in prevalence between regions.³

Understanding the magnitude of suicide-related maternal mortality is important to improve assessment, identification and reporting of maternal suicide deaths and inform prevention strategies. However, there are a number of challenges in accurately estimating the burden. Firstly, there are a number of different definitions of maternal death (**Figure 1**). The World Health Organization (WHO) uses the terms *maternal death* and *late maternal death*, which are defined in the International Classification of Diseases (11th Revision; ICD-11) and which differ from each other only in terms of the time periods they cover.^{4,5} Maternal deaths and late maternal deaths are combined in the ICD-11 under a new grouping of *comprehensive maternal deaths*.⁴ Others, notably the US Centres for Disease Control and Prevention (CDC), refer to *pregnancy-associated* and *pregnancy-related deaths*, which both span the same time period but differ according to the designated cause of death.⁶ Confusingly, the WHO also refers to *pregnancy-related death*, but defines this term differently from the CDC both in terms of the causes included and the time period covered.⁵ In reports and publications presenting maternal death statistics, there is often a lack of clarity around which definition was applied.

Figure 1. Definitions of maternal death
[INSERT Figure 1 here]

Secondly, there are differences in the denominator used to calculate rates of maternal mortality. The commonly reported maternal mortality ratio (MMR) reports the number of maternal deaths during a given time period per 100,000 live births during the same time period.⁵ The use of live births as a denominator means that deaths occurring during pregnancy and stillbirths are not captured by the MMR. To address this, some countries such as the UK also report the number of maternal deaths per 100,000 maternities, defined as the number of women giving birth, including stillbirth, at or beyond 24 weeks' gestation.¹ When maternities are used as the denominator, the statistic is often referred to as a maternal mortality *rate* (rather than ratio), but the terminology is inconsistent and in practice the distinction between MMR and maternal mortality rates is often unclear. These differences create challenges when comparing statistics between countries or across different reporting systems.

1
2
3 Thirdly, there is variation in the availability and quality across different regions of the world.
4 High quality data on maternal suicide requires an accurate identification of all maternal
5 deaths, as well as an accurate designation of cause of death. Some countries have systems
6 in place to identify and investigate maternal deaths and their causes at a regional or
7 national level. Such systems include systematic maternal mortality surveillance systems,
8 confidential enquiries and Reproductive Age Mortality Surveys (RAMOS). The assignment of
9 cause of death can include the use of formal death registration certificates and verbal
10 autopsies in which family members are asked to provide information on circumstances
11 around the death. In countries with less robust death reporting systems it is difficult to
12 determine how accurate maternal death reporting is.
13
14
15

16 A final challenge is the misclassification of deaths by suicide to other causes such as injury or
17 accidents. This is thought to be common given the stigma associated with suicide and its
18 criminalisation in some societies.³ A review of maternal suicide in LMIC found that
19 reclassifying the leading suicide methods from injuries to suicide increased the pooled
20 prevalence of pregnancy-related suicide deaths from 1.00 to 1.68.³ Misclassification is
21 compounded by the fact that until the most recent revision of the ICD, maternal suicide did
22 not have a specific code. There has also been a change in the recommended classification of
23 maternal suicides. The ICD-10 categorised maternal deaths due to injuries or mental
24 disorders as indirect maternal deaths.⁷ The WHO subsequently published a derivation of the
25 ICD-10: this ICD-Maternal Mortality (ICD-MM) calls for maternal suicides to be classified as
26 direct maternal deaths.⁸ This discrepancy has led to variations in the reporting of maternal
27 suicide as direct or indirect causes of death, with some statistics reporting maternal suicide
28 as an incidental cause of death and other statistics excluding suicides from their maternal
29 mortality data altogether.⁹
30
31
32
33
34

35 There are several existing systematic reviews of maternal suicide but these are limited to
36 specific perinatal populations or geographical regions, and there is no up-to-date synthesis
37 of the global evidence for all perinatal women. Lindahl *et al.* (2005) reviewed prevalence of
38 suicide ideation, attempts and deaths in pregnancy and the post-partum period globally;
39 however this included studies published up to 2002 and their findings need updating.¹⁰ Fuhr
40 *et al.* (2014) assessed the contribution of suicide and injuries to perinatal mortality in LMIC
41 only.³ Amiri *et al.* (2021) assessed rates of suicide ideation, attempts and mortality globally
42 among post-partum women only.¹¹ Chin *et al.* (2022) reviewed prevalence, risk factors,
43 outcomes and interventions for perinatal deaths by suicide but focused on a limited number
44 of studies that were not included in an earlier review by Mangla *et al.* (2019).² Chin *et al.*
45 (2022) also restricted their search to studies published in English, and most of the identified
46 studies were conducted in the USA.²
47
48
49
50

51 Our systematic review expands upon previous reviews by including suicides occurring during
52 pregnancy and the post-partum period in all countries globally with no language or date
53 restrictions. Results will enable an estimation of the contribution of suicide to maternal
54 deaths globally and a comparison of the burden across geographical regions. Identifying
55 countries reporting either very high or very low rates will inform further work to determine
56 whether differences in rates reflect true differences or reporting- or coding-related factors.
57 Our review will also show whether maternal suicide deaths are reported as direct, indirect
58 or incidental maternal deaths, informing a possible need for further clarification of the ICD-
59
60

1
2
3 11 and ICD-MM classifications. Finally, our review will identify countries for which no data
4 on maternal suicide is available, an important requirement in meeting the United Nations'
5 Sustainable Development Goal (SDG) 3.¹²
6
7

8 **Objectives**

9

10 This systematic review aims to identify and synthesise evidence on maternal suicide deaths
11 globally. The primary objective is to determine the contribution of suicide towards maternal
12 mortality and explore differences between geographical regions. The secondary objectives
13 are to summarise the availability and quality of data globally and to describe how suicide
14 deaths are classified across different countries.
15
16

17 **METHODS AND ANALYSIS**

18

19 **Registration and protocol adherence**

20

21 This systematic review protocol is reported in adherence with the Preferred Reporting Items
22 for Systematic Reviews and Meta-Analyses for systematic review protocols (PRISMA-P)
23 guidelines (**Supplementary Information S1**).¹³ The review was registered on PROSPERO
24 (CRD42023429072) on 25 May 2023. If important amendments to the protocol are required
25 during the course of the review, these will be recorded as amendments on PROSPERO and
26 reported in the final publication as a deviation from the protocol. The start date for the
27 review was January 2023 and the planned end date is July 2025.
28
29
30
31

32 **Definitions**

33

34 For the purposes of this review, we define maternal suicide as the death of a woman by
35 suicide at any time during pregnancy, intrapartum or within one year of giving birth. The
36 perinatal period is the period of pregnancy and up to one year after giving birth.
37
38
39

40 **Eligibility criteria**

41

42 *Population, intervention, comparison and outcome framework*

43 The population of interest is women at any stage of pregnancy or within one year of giving
44 birth. The outcome of interest is death by suicide during the perinatal period. There is no
45 intervention or control for this systematic review of global maternal suicide rates.
46
47

48 *Inclusion and exclusion criteria*

49 We will include any study that reports data on the number of maternal deaths due to
50 suicide along with appropriate denominator data (either total live births or total maternities
51 during the same time period as maternal deaths). Studies reporting the proportion of all
52 maternal deaths that were due to suicide will also be included. We will include cohort,
53 cross-sectional and case control study designs as well as data from national surveys,
54 databases and reports such as mortality statistics and confidential enquiries. Qualitative
55 studies will be excluded as they are unlikely to contain numerator and denominator data on
56 maternal suicide and maternal death. There will be no minimum threshold applied to the
57 sample size or numbers of suicides reported. Studies that do not report data on the
58
59
60

denominator to allow calculation of the MMR attributable to suicide or the proportion of deaths that are due to suicide will be excluded. Studies in which suicide deaths cannot be differentiated from deaths from injury or other causes will also be excluded. We will include studies published in any language.

Patient and public involvement

There was no patient or public involvement in the design of this systematic review.

Search strategy and data sources

Medline, Embase, PsychINFO, Global Health and CINAHL databases were searched with no date or language restrictions on 5 May 2023. A search strategy using terms relevant to perinatal status, mortality and suicide was developed in collaboration with a university research librarian and adapted for each database. The search strategy used for Medline is shown in **Table 1**. In order to capture relevant evidence not published in the databases above, we conducted an extensive grey literature search. This involved searches of *Google* and *Google Scholar* using the relevant terms listed above to capture non-academic articles such as reports and statistical databases. We also searched the websites of the World Health Organization (WHO), the United Nations Children's Fund (UNICEF), the United Nations Population Fund (UNFPA), UN Women, the World Bank and the Global Burden of Disease Study to identify relevant reports. All records identified by the database search and grey literature searches were combined and duplicates removed using *Endnote*.¹⁴ De-duplicated references were imported into *Covidence* for screening, data extraction and quality assessment.¹⁵ Prior to submitting the review for publication, the search strategy will be re-run to identify any studies published since the original search.

Table 1. Search strategy for Medline

1	Pregnancy/ or Pregnant Women/
2	Postnatal Care/ or Postpartum Period/
3	1 or 2
4	Mortality/
5	3 and 4
6	maternal mortality/ or maternal death/
7	((pregnan* or maternal or postnatal or post-natal or postpart* or post-part*) and (death? or mortality or fatalit*)).ti.
8	((pregnan* or maternal or postnatal or post-natal or postpart* or post-part*) adj3 (death? or mortality or fatalit*)).ti,ab,kf.
9	5 or 6 or 7 or 8
10	"cause of death"/
11	Death Certificates/
12	exp suicide/
13	"cause of death".ti,ab,kf.
14	(suicid* or overdos* or self harm or self injury or violen*).ti,ab,kf.
15	10 or 11 or 12 or 13 or 14
16	9 and 15

Data extraction

Screening of titles and abstracts was performed independently by two reviewers (ES and JG) to exclude studies which did not meet inclusion criteria. Any disagreements were discussed and resolved with a third reviewer (SH and GF). Full-texts were retrieved. Any missing full texts were searched for using a university library database, online journal archives, *Google* and by contacting study authors. Any full-texts that could not be retrieved after this search process were excluded. Full-texts of remaining studies were screened independently by two reviewers (ES, JG, SH, GF) and any disagreements were resolved through discussion. Reasons for excluding full-texts were recorded. Two reviewers (ES, JG, SH, GF) will independently extract data from included studies using a standardised and piloted data extraction form consisting of the following fields:

- **Publication details:** lead author, publication year, publication language
- **Setting:** country, region, type of setting (e.g. hospital or community deaths), dates covered, whether data was collected at national or regional level
- **Participants:** perinatal status (pregnant or post-partum), definition of maternal death used, age, any sociodemographic characteristics (e.g. socioeconomic background, ethnicity, income, parity)
- **Methods of assessment:** method of identifying maternal deaths, method of ascertaining cause of death, classification of suicide (as indirect, direct or incidental cause of death)
- **Outcomes:** total maternal deaths, total live births, total maternities, number of suicide deaths, proportion of deaths due to suicide, accuracy measures (e.g. standard deviation, p-values or 95% confidence intervals), mode of suicide

If reported data is unclear or insufficient, authors of the study will be contacted to request the additional data required for inclusion in the review. Publications in English, French, German, Greek, Italian, Spanish and Portuguese will be assessed by the study authors. For all other languages, translation support will be sought from wider colleagues.

Quality assessment and risk of bias

Two reviewers (ES, JG, SH, GF) will independently assess the quality of each included study. We will use criteria adapted from a study by Grollman and Ronsmans¹² to assess quality and risk of bias. This will include assessment of three domains: quality of the method of death ascertainment; completeness of the cause of death assignment; and quality of the method of assigning cause of death. Each study will be assigned a rating of low, medium or high quality in each of these categories using the attributes outlined by Grollman and Ronsmans.¹⁶ We have chosen this method as there is a precedent for its use in a similar review, and other risk of bias assessment tools have limited application to studies looking at causes of death. Any disagreements will be discussed with the other review authors (FA, MQ) until a consensus is reached.

Data synthesis

1
2
3 A narrative synthesis of included studies will be conducted and a table of study
4 characteristics will be compiled to provide an overview of all included studies. Countries for
5 which data on maternal suicide deaths is available will be mapped and colour-coded
6 (red/amber/green) according to the quality of evidence. The proportion of countries
7 classifying maternal suicide deaths as direct, indirect or incidental deaths will also be
8 reported.
9
10

11
12 Random-effects meta-analyses will be conducted if numerator and denominator data for
13 the same standardised outcome is reported by at least three studies. The primary outcome
14 will be pooled estimates of suicide-related maternal mortality rates. Depending on the data
15 identified and comparability between studies, we will use either live births or maternities as
16 the denominator or conduct a separate pooled estimate for each. Forest plots will be used
17 to present data visually. Statistical heterogeneity between studies will be calculated using
18 the I^2 statistic. If data allows and sufficient studies are identified, we will carry out subgroup
19 analyses to explore differences in suicide death rates according to perinatal status
20 (pregnancy vs. post-partum); country income classification (high-income vs. low- and
21 middle-income countries); geographical region (using WHO geographical regions); and study
22 quality. The secondary outcome will be a pooled estimate of the proportion of all maternal
23 deaths caused by suicide, using number of suicide deaths as the numerator and total
24 number of maternal deaths as the denominator. If meta-analysis is not possible due to a
25 lack of standardised outcome reporting or due to fewer than three studies presenting the
26 same outcome. The presence of publication bias will be assessed using funnel plots. Meta-
27 analyses will be conducted using STATA.
28
29
30
31

32 **ETHICS AND DISSEMINATION**

33
34
35 No primary data is being collected for this study therefore ethical approval is not required.
36 Results of this study will be published in a peer-reviewed publication and presented at
37 conferences. Raw data will be published as supplementary material in the final publication.
38
39

40 **DISCUSSION**

41
42 This systematic review and meta-analysis will produce a synthesis of the available evidence
43 on maternal suicide globally. This will allow comparisons to be made between countries on
44 suicide-related maternal mortality rates and the overall contribution of suicide toward
45 maternal mortality. Summarising the availability and quality of data available in different
46 regions of the world will identify priorities for future research. Identifying countries with
47 high or low maternal suicide rates will also inform future studies of risk factors and
48 preventive interventions in different populations.
49
50

51
52 There are a number of limitations to this study. It is possible that some of the data retrieved
53 will be of poor quality, limiting its reliability and making comparison between areas
54 challenging. The wide variety of methods and definitions used in studies identifying
55 maternal suicide may also make it difficult to draw direct comparisons, and it may not be
56 possible to calculate pooled estimates. By excluding maternal deaths attributed to
57 accidents, injury or substance use, we may be under-estimating the true burden of suicide-
58 related maternal mortality. However, given that it is not possible to establish whether these
59
60

1
2
3 deaths were intentional or non-intentional, we felt it was a safer and more robust approach
4 to exclude these deaths. The lack of data in some regions could also limit the power of
5 analyses. There has also been a difficulty in acquiring the full texts of several of the included
6 abstracts. Every effort was made to identify these texts, however a significant number of
7 texts had to be excluded. This could impact whether we have captured the full breadth of
8 evidence on this topic.
9
10

11
12 **Acknowledgements** We would like to thank Ms. Nia Roberts, Senior Outreach Librarian at
13 the University of Oxford Bodleian Health Care Libraries, for her guidance and support in the
14 development of the search strategy for this review.
15

16
17 **Data statement** No data was gathered for this protocol.
18

19
20 **Author contributions** ES, SH and GF developed the protocol. ES developed the search
21 strategy with input from SH and GF. ES, JG, ZD, SH and GF contributed to screening. ES, JG,
22 ZD, SH and GF will conduct data extraction and quality assessment. SH, GF, MQ and FA
23 contributed to the analysis plan. All authors contributed to the writing of this manuscript.
24 SH and GF act as guarantors for the review.
25

26
27 **Funding statement:** GF is funded by a Clinical Research Fellowship from the Nuffield
28 Department of Population Health, University of Oxford. The funder had no role in the design
29 or development of this protocol.
30

31
32 **Competing interests statement:** All authors declare no competing interests.
33

34 REFERENCES

- 35
36 1. Knight M, Bunch K, Felker A, Patel R, Kotnis R, Kenyon S, Kurinczuk JJ (Eds.) on behalf of
37 MBRRACE-UK. Saving Lives, Improving Mothers' Care Core Report - Lessons learned to
38 inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths
39 and Morbidity 2019-21. Oxford: National Perinatal Epidemiology Unit, University of Oxford
40 2023.
41
42
43 2. Chin K, Wendt A, Bennett IM, Bhat A. Suicide and Maternal Mortality. *Curr Psychiatry Rep*
44 2022;24:239–275.
45
46
47 3. Fuhr DC, Calvert C, Ronsmans C, Chandra PS, Sikander S, De Silva MJ, Patel V.
48 Contribution of suicide and injuries to pregnancy-related mortality in low-income and
49 middle-income countries: a systematic review and meta-analysis. *Lancet Psychiatry*. 2014
50 Aug;1(3):213-25.
51
52
53 4. World Health Organization. International Statistical Classification of Diseases and Related
54 Health Problems (11th edition; ICD-11). 2021. Geneva: World Health Organization.
55
56
57 5. World Health Organization. Trends in maternal mortality 2000 to 2020: estimates by
58 WHO, UNICEF, UNFPA, World Bank Group and UNDESA/Population Division. 2023. Geneva:
59 World Health Organization.
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
6. Centers for Disease Control and Prevention. Pregnancy Mortality Surveillance System. 2024. Available at www.cdc.gov/reproductivehealth/maternal-mortality/pregnancy-mortality-surveillance-system.htm (accessed 11 April 2024).
7. World Health Organization. International Statistical Classification of Diseases and Related Health Problems (10th edition; ICD-10). 2016. Geneva: World Health Organization.
8. World Health Organization. The WHO Application of ICD-10 to Deaths during Pregnancy, Childbirth and Puerperium: ICD-MM. 2012. Geneva: World Health Organization.
9. Hasegawa J, Katsuragi S, Tanaka H, Kubo T, Sekizawa A, Ishiwata I, Ikeda T. How should maternal death due to suicide be classified? Discrepancy between ICD-10 and ICD-MM. *BJOG*. 2020 May;127(6):665-667.
10. Lindahl V, Pearson JL, Colpe L. Prevalence of suicidality during pregnancy and the postpartum. *Arch Womens Ment Health*. 2005 Jun;8(2):77-87.
11. Amiri S, Behnezhad S. The global prevalence of postpartum suicidal ideation, suicide attempts, and suicide mortality: A systematic review and meta-analysis. *International Journal of Mental Health*. 2021;50:311–336.
12. United Nations. Transforming Our World: The 2030 Agenda for Sustainable Development. 2015. Available at: <https://sdgs.un.org/2030agenda> (accessed 08/04/2024).
13. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA. Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015;4(1):1.
14. The EndNote Team. EndNote (Version X9). 2013. Philadelphia, PA: Clarivate.
15. Covidence Systematic Review software. 2023. Melbourne: Veritas Health Innovation.
16. Grollman C, Ronsmans C. Systematic review of the proportion of pregnancy-related deaths attributed to HIV in population-based studies in sub-Saharan Africa. *Trop Med Int Health*. 2014 Jan;19(1):83-97.

Figure 1. Definitions of maternal death

- **Maternal death** is defined by the World Health Organization (WHO) as the death of a woman from direct or indirect obstetric causes while pregnant or within 42 days after the end of pregnancy from causes associated with, or exacerbated by, pregnancy or its management, irrespective of the duration or outcome of the pregnancy.⁵
- **Late maternal death** is defined by the WHO as a death of a woman from direct or indirect obstetric causes more than 42 days but less than one year after termination of pregnancy.⁵
- **Comprehensive maternal death** is defined by the International Classification of Diseases (11th Revision) WHO as maternal deaths and late maternal deaths combined.⁴
- **Pregnancy-related death** is defined by the WHO as a death of a woman while pregnant or within 42 days of termination of pregnancy, irrespective of the cause of death. This definition includes obstetric, non-obstetric, unintentional (accidental) and incidental causes of death.⁵
- **Pregnancy-related death** is defined by the US Centers for Disease Control and Prevention (CDC) as a death during or within one year of pregnancy from a pregnancy complication, a chain of events initiated by pregnancy or the aggravation of an unrelated condition by the physiological effects of pregnancy.⁶
- **Pregnancy-associated death** is defined by the CDC as a death during or within one year of pregnancy, regardless of the cause of death.⁶

PRISMA-P (Preferred Reporting Items for Systematic review and Meta-Analysis Protocols) 2015 checklist: recommended items to address in a systematic review protocol*

Section and topic	Item No	Checklist item	Page Reported
ADMINISTRATIVE INFORMATION			
Title:			
Identification	1a	Identify the report as a protocol of a systematic review	1, 2
Update	1b	If the protocol is for an update of a previous systematic review, identify as such	N/A
Registration	2	If registered, provide the name of the registry (such as PROSPERO) and registration number	2, 5
Authors:			
Contact	3a	Provide name, institutional affiliation, e-mail address of all protocol authors; provide physical mailing address of corresponding author	1
Contributions	3b	Describe contributions of protocol authors and identify the guarantor of the review	9
Amendments	4	If the protocol represents an amendment of a previously completed or published protocol, identify as such and list changes; otherwise, state plan for documenting important protocol amendments	5
Support:			
Sources	5a	Indicate sources of financial or other support for the review	9
Sponsor	5b	Provide name for the review funder and/or sponsor	9
Role of sponsor or funder	5c	Describe roles of funder(s), sponsor(s), and/or institution(s), if any, in developing the protocol	9
INTRODUCTION			
Rationale	6	Describe the rationale for the review in the context of what is already known	3-5
Objectives	7	Provide an explicit statement of the question(s) the review will address with reference to participants, interventions, comparators, and outcomes (PICO)	5, 6
METHODS			
Eligibility criteria	8	Specify the study characteristics (such as PICO, study design, setting, time frame) and report characteristics (such as years considered, language, publication status) to be used as criteria for eligibility for the review	5, 6
Information sources	9	Describe all intended information sources (such as electronic databases, contact with study authors, trial registers or other grey literature sources) with planned dates of coverage	6
Search strategy	10	Present draft of search strategy to be used for at least one electronic database, including planned limits, such that it could be repeated	Table 1

Study records:			
Data management	11a	Describe the mechanism(s) that will be used to manage records and data throughout the review	6
Selection process	11b	State the process that will be used for selecting studies (such as two independent reviewers) through each phase of the review (that is, screening, eligibility and inclusion in meta-analysis)	7
Data collection process	11c	Describe planned method of extracting data from reports (such as piloting forms, done independently, in duplicate), any processes for obtaining and confirming data from investigators	7
Data items	12	List and define all variables for which data will be sought (such as PICO items, funding sources), any pre-planned data assumptions and simplifications	7
Outcomes and prioritization	13	List and define all outcomes for which data will be sought, including prioritization of main and additional outcomes, with rationale	8
Risk of bias in individual studies	14	Describe anticipated methods for assessing risk of bias of individual studies, including whether this will be done at the outcome or study level, or both; state how this information will be used in data synthesis	7
Data synthesis	15a	Describe criteria under which study data will be quantitatively synthesised	8
	15b	If data are appropriate for quantitative synthesis, describe planned summary measures, methods of handling data and methods of combining data from studies, including any planned exploration of consistency (such as I^2 , Kendall's τ)	8
	15c	Describe any proposed additional analyses (such as sensitivity or subgroup analyses, meta-regression)	8
	15d	If quantitative synthesis is not appropriate, describe the type of summary planned	8
Meta-bias(es)	16	Specify any planned assessment of meta-bias(es) (such as publication bias across studies, selective reporting within studies)	8
Confidence in cumulative evidence	17	Describe how the strength of the body of evidence will be assessed (such as GRADE)	8

*** It is strongly recommended that this checklist be read in conjunction with the PRISMA-P Explanation and Elaboration (cite when available) for important clarification on the items. Amendments to a review protocol should be tracked and dated. The copyright for PRISMA-P (including checklist) is held by the PRISMA-P Group and is distributed under a Creative Commons Attribution Licence 4.0.**

From: Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart L, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;349(jan02 1):g7647.