



Rationing of total knee replacement: A cost-effectiveness analysis on a large trial dataset

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2011-000332
Article Type:	Research
Date Submitted by the Author:	26-Aug-2011
Complete List of Authors:	Dakin, Helen; University of Oxford, Health Economics Research Centre, Department of Public Health Gray, Alastair; University of Oxford, Health Economics Research Centre, Department of Public Health Fitzpatrick, Ray; University of Oxford, Department of Public Health Maclennan, Graeme; University of Aberdeen, Health Services Research Unit Murray, David; University of Oxford, Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences
Primary Subject Heading:	Health services research
Keywords:	Economic evaluation, total knee replacement, heterogeneity, cost-utility analysis, randomised controlled trial

SCHOLARONE™
Manuscripts

Only

Rationing of total knee replacement: A cost-effectiveness analysis on a large trial dataset

Running header:

Rationing of total knee replacement

Key words:

Economic evaluation; total knee replacement; heterogeneity; subgroup analysis; randomised controlled trial.

Authors:

Helen Dakin,¹ Alastair Gray,¹ Ray Fitzpatrick,² Graeme MacLennan,³ and David Murray⁴

Authors' affiliation:

¹ Health Economics Research Centre, Department of Public Health, University of Oxford, Old Road Campus, Oxford, UK

² Department of Public Health, University of Oxford, Old Road Campus, Oxford, UK

³ Health Services Research Unit, University of Aberdeen, Aberdeen, UK

⁴ Nuffield Department of Orthopaedic Surgery, University of Oxford, UK

Corresponding author:

Helen Dakin

Researcher

Health Economics Research Centre

Department of Public Health, University of Oxford

Old Road Campus

Headington

Oxford

OX3 7LF

Email: helen.dakin@dph.ox.ac.uk

Tel: 01865 289422

Fax: 01865 289271

Word count: 3,398 (main text of introduction, methods, results and discussion)

Number of figures: 3

Number of tables: 2

Ethics: The KAT trial was approved by the Multi Centre Research Ethics Committee for Scotland in November 1998 (research protocol MREC/98/0/100) and was approved by the Local Research Ethics Committees in each study centre recruiting trial participants. Further details are available on request.

Data sharing statement: No additional data available.

Protocol: Available at <http://www.hta.ac.uk/protocols/199500100001.pdf>.

Abstract

Objectives: Many UK primary care trusts (PCTs) have recently introduced eligibility criteria restricting total knee replacement (TKR) to patients with low pre-operative Oxford Knee Scores (OKS) to cut expenditure. We evaluate these criteria by assessing the cost-effectiveness of TKR compared with no knee replacement for patients with different baseline characteristics from an NHS perspective.

Design: The cost-effectiveness of TKR in different patient subgroups was assessed using regression analyses of patient-level data from the Knee Arthroplasty Trial (KAT), a large pragmatic randomised controlled trial comparing different knee prostheses. Costs and quality-adjusted life-years (QALYs) observed in the KAT trial over a five-year time horizon following TKR were compared with conservative assumptions about the costs and outcomes that would have been accrued had TKR not been performed.

Results: On average, primary TKR and five years of subsequent care cost £7,458 per patient (standard deviation, SD: £4,058), and patients gained an average of 1.33 (SD: 1.43) QALYs. As result, TKR cost £5,623 per QALY gained. Although costs and health outcomes varied with age and sex, TKR cost less than £20,000 per QALY gained for patients of ASA grade 1 or 2 who had baseline OKS of 39 or below, and for ASA grade 3 patients with OKS of 34 or below, even with highly conservative assumptions about costs and outcomes without TKR. Body mass index (BMI) had no significant effect on costs or outcomes. Restricting TKR to patients with pre-operative OKS of 26 or less would inappropriately deny a highly cost-effective treatment to more than 10,000 patients annually.

Conclusions: TKR is highly cost-effective for most current patients if the NHS is willing to pay £20,000-£30,000 per QALY gained. At least 97% of TKR patients in England have more severe symptoms than the thresholds we have identified, suggesting further rationing by OKS is probably unjustified.

Article summary

Article focus

- We assess the cost-effectiveness of total knee replacement (TKR) compared with no knee replacement for patients with different baseline characteristics from a National Health Service (NHS) perspective.
- In particular, we assess the appropriateness of eligibility criteria recently introduced by many UK primary care trusts (PCTs), which restrict TKR to patients with low pre-operative Oxford Knee Scores (OKS) to cut expenditure.

Key messages

- We find TKR to be highly cost-effective, costing £5,623 per quality-adjusted life-year (QALY) gained for the average patient.
- TKR costs less than £20,000 per QALY gained for healthy patients with OKS of ≤ 39 , or ≤ 34 for patients who have other conditions restricting their daily activities.
- We find no evidence to support the criteria for restricting access to TKR that have been proposed by some PCTs and calculate that restricting TKR to those patients with pre-operative OKS of 26 or less would deny a highly cost-effective treatment to more than 10,000 patients per year.

Strengths and limitations of this study

- This is the first study assessing how the cost-effectiveness of TKR varies with OKS and the first assessing the clinical/economic implications of the newly-introduced rationing criteria.
- Analyses are based on patient-level data from a large pragmatic trial with detailed, prospective collection of utilities, baseline characteristics and all major knee-related NHS resource use.
- Our study makes several highly conservative assumptions: in particular, assuming that patients would have accrued no knee-related costs and remained at baseline utility without TKR. Furthermore, the KAT sample included only 37 patients with pre-operative OKS > 35 . As result, TKR may be also cost-effective for some patients with OKS above 39.

Introduction

Total knee replacement (TKR) is a highly effective treatment for patients with bone-on-bone osteoarthritis and significant knee symptoms, producing substantial reductions in symptoms (particularly pain)[1] and lasting for at least 15 years in 83-94% of cases.[2] However, about 18% of patients consider the outcomes of their surgery to be only fair or poor,[1] and a small proportion experience complications.[3-5] Previous economic evaluations have found TKR to be highly cost-effective, with a cost-effectiveness ratio between €1,276 and \$18,300 per quality-adjusted life-year (QALY) gained for the average patient[6-10]: well below the £20,000-£30,000 per QALY range that the National Institute for Health and Clinical Excellence (NICE) considers cost-effective.[11] However, studies suggest that the costs and benefits of TKR vary between patient subgroups, with TKR being more cost-effective in younger patients,[9] those attending high-volume centres[7] and those deemed low-risk based on age, comorbidities, and poverty criteria.[7] There is also evidence that older patients have higher medical costs,[12] although other studies suggest that knee replacement is also cost-effective in nonagenarians[13] and draw conflicting conclusions on how benefits vary with body mass index (BMI).[14-16]

In the current economic environment there is great pressure to reduce NHS expenditure and numerous Primary Care Trusts (PCTs) have proposed cutting costs by limiting access to joint replacement[17-24] or classifying these procedures as being of limited value.[23-25] Since around 60,000 primary TKR procedures are conducted in England and Wales each year,[26] savings could be made with little loss of patient welfare *if* it were possible to reliably predict which patients obtain little/no benefit from TKR based on explicit, evidence-based criteria. Although the widely-used Oxford Knee Score (OKS) was developed to assess outcomes of knee replacement in trial populations,[27] several PCTs have set maximum OKS thresholds (ranging from 18 to 32), above which patients are ineligible for TKR.[17-23] However, we are not aware of any clinical or cost-effectiveness justification for these thresholds. Furthermore, many patients with higher OKS have significant symptoms or limitations to daily activities that could be improved by surgery. If rationing policies are to be adopted, cost-effectiveness analysis offers the only fair basis for designing them, since it ensures that NHS resources are allocated to maximise the health gained from available resources. As treatments are generally considered cost-effective if they cost less than £20,000-£30,000 per quality-adjusted life-year (QALY) gained,[11] restricting TKR to those patients for whom the procedure

1
2
3 costs less than £20,000-£30,000 per QALY gained would be a reasonable approach,
4 potentially realising savings that could be invested in other treatments giving greater
5 health gains. However, we are aware of no published evidence assessing how the
6 cost-effectiveness of TKR varies with OKS.
7
8
9

10
11 We therefore use data from the Knee Arthroplasty Trial (KAT)[28, 29] to assess the
12 appropriateness of rationing knee replacement by OKS, age, BMI, and clinical
13 characteristics, by calculating the cost-effectiveness of TKR in different patient
14 subgroups. In contrast to previous studies evaluating the cost-effectiveness of TKR
15 that relied upon data from small cohorts with short follow-up,[6-10, 13] the KAT trial
16 dataset provides detailed, prospective individual patient data on baseline
17 characteristics, treatments, complications, costs, and quality of life, and now has
18 more than five years' follow-up on 2352 patients.
19
20
21
22
23
24
25
26
27

28 **Methods**

29
30 We assessed the cost-effectiveness of primary TKR compared with no knee
31 replacement for patients with different baseline characteristics. Our analysis took a
32 National Health Service (NHS) perspective and excluded personal and social
33 services from the analysis, although TKR is likely to delay admission to residential
34 care,[13] reduce personal care costs and help recipients to continue paid
35 employment. A cost-utility analysis was conducted, which assessed cost-
36 effectiveness in terms of cost per quality-adjusted life-year (QALY) gained to capture
37 the quality of life gains associated with TKR.
38
39
40
41
42
43

44
45 Data on costs and quality of life following TKR were taken from the KAT trial
46 (ISRCTN 45837371): a pragmatic, partial-factorial, unblinded randomised controlled
47 trial (RCT) in which 2352 participants attending 34 UK centres were randomly
48 allocated to undergo TKR with/without a metal-backed tibial component, with/without
49 patellar resurfacing, and/or with/without a mobile bearing.[28, 29] Five-year trial
50 results demonstrated that patients experience substantial improvements in quality of
51 life and functional status following knee replacement that are maintained out to five
52 years[29]; further follow-up is ongoing. Our analysis took a five-year time horizon in
53 line with the results published to date[29]; costs and health benefits occurring more
54 than five years after primary TKR were conservatively excluded.
55
56
57
58
59
60

1
2
3 KAT participants completed the Oxford Knee Score (OKS) and EQ-5D
4 questionnaires immediately before and three months after TKR and annually
5 thereafter. The OKS is a validated 12-item questionnaire to assess health status and
6 outcomes in relation to knee replacement,[27] while the EQ-5D is a generic health
7 state preference measure.[30] Health state preference values (“utilities”) for EQ-5D
8 profiles were based on time-trade-off valuations by members of the UK general
9 public.[30] The number of QALYs that each patient accrued following TKR was
10 calculated as the area under the utility curve, with linear interpolation between utility
11 measurements; those patients who died were assumed to remain at the last
12 observed utility until death.
13
14
15
16
17
18
19
20

21 Healthcare resource use data (including knee-related outpatient, general practice
22 and physiotherapy consultations, and theatre time, hospital days, complications and
23 knee components used during the primary admission for TKR and during any knee-
24 related readmissions) were collected prospectively for all KAT participants and
25 valued at 2007-08 prices, as described previously.[29] Costs incurred after Year 1
26 were discounted at 3.5% per year based on current UK recommendations.[31]
27
28
29
30
31

32 The number of QALYs gained from TKR was calculated as the number of QALYs
33 observed minus the number of QALYs expected without TKR. Since all KAT
34 participants underwent TKR, assumptions were needed to estimate the costs and
35 QALYs that would have been accrued if no TKR procedures had occurred in the five-
36 year time horizon. We assumed that:
37
38
39

- 40 • Without TKR, patients would have remained at their baseline EQ-5D utility for five
41 years or until death (if sooner). In practice, patients’ symptoms and quality of life
42 are likely to deteriorate due to worsening arthritis[16, 32] and increasing age.[33]
43
- 44 • The seven KAT participants who died before hospital discharge would have
45 survived for five years at baseline utility without TKR but that TKR had no effect
46 on the date of death for other patients. This assumption is highly conservative
47 since those patients who die soon after joint replacement tend to be older[5, 34]
48 and have comorbidities that could have led to death without TKR.[5]
49
- 50 • Patients would not have used any healthcare resources due to their knee
51 problems if they had not undergone TKR. This assumption is highly conservative
52 since osteoarthritis patients will receive medical management without TKR and a
53 recent Canadian case-control study suggested that TKR recipients accrued lower
54 costs 6-18 months after the procedure than matched controls who had arthritis
55 but no TKR.[32]
56
57
58
59
60

1
2
3
4
5 Since these assumptions are likely to underestimate the benefits and overestimate
6 the additional costs of TKR, we evaluated the effect of relaxing these assumptions in
7 sensitivity analyses.
8
9

10
11 The 7% of data on baseline characteristics, resource use and quality of life that were
12 missing were imputed by multiple imputation using the ice command[35-37] (version
13 1.3.0) within Stata Version 11.0 (College Station, TX), thereby avoiding the bias and
14 inefficiency associated with complete case analysis.[37-39] Treatment indicators and
15 the covariates included in subsequent regression analyses were included in the
16 imputation model to avoid bias.[37] Fifty imputed datasets were generated and
17 results combined.
18
19
20
21
22

23
24 Regression models were used to estimate how the costs and QALYs gained from
25 TKR vary with baseline characteristics and to predict outcomes for different patient
26 groups without subdividing the trial population into small patient subgroups. Since
27 costs and QALYs were highly skewed, generalised linear models (GLM) with gamma
28 family distributions were used to predict total costs and the QALYs accrued with or
29 without TKR; prior to regression, the number of QALYs accrued was subtracted from
30 the maximum number of QALYs that could have been accrued over the five-year trial
31 period to ensure that all QALY data were positive. We evaluated the effect of six
32 baseline characteristics on costs and QALYs with/without TKR:
33
34
35
36
37
38

- 39 • Female sex;
- 40 • Age at time of operation;
- 41 • Baseline OKS (using the new, 0-48, scoring system[40]);
- 42 • American Society of Anesthesiologists (ASA) grade 3 (symptomatic illness
43 with minimal restriction on daily life) rather than grades 1 (fit & healthy) or 2
44 (asymptomatic illness with no restriction on daily life);
- 45 • BMI (kg/m²);
- 46 • Presence of arthritis in one knee rather than in both knees or generally.
47
48
49
50
51
52
53

54 Since the introduction of biologic therapies has changed management of rheumatoid
55 arthritis and as rheumatoid arthritis patients and those with severe comorbidities are
56 likely have very different costs and outcomes from most patients, 108 rheumatoid
57 patients and 13 patients of ASA grade 4 were excluded from the analysis.
58
59
60

1
2
3 All regression analyses were conducted in Stata Version 11. Bootstrapping was used
4 to capture uncertainty around cost-effectiveness ratios and correlations between
5 costs and QALYs with/without TKR; the regression models for costs and outcomes
6 were run for each of 125 bootstrap samples drawn from each imputed KAT dataset
7 and results combined using Rubin's rule.[38] Predicted costs and the predicted
8 number of QALYs accrued with and without TKR in different patient groups were
9 calculated from regression coefficients and used to estimate incremental cost-
10 effectiveness ratios (ICERs). Bootstrap results were used to calculate cost-
11 effectiveness acceptability curves (CEACs),[41] which plot the probability of
12 treatment being cost-effective against the ceiling ratio (i.e. the maximum that society
13 is willing or able to pay per QALY gained).

14
15
16
17
18
19
20
21
22
23 We also calculated number of patients in England and Wales eligible for TKR under
24 different rationing policies based on 61,651 patients undergoing primary TKR each
25 year.[26] The proportion of patients eligible for TKR under different rationing criteria
26 was based on analysis of an extract of patient-level Patient Reported Outcome
27 Measures[1] (PROMs) data on admissions for knee replacement up to 31st
28 December 2010, supplemented where necessary by the baseline characteristics from
29 the KAT sample.

30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Results

Of the 2131 KAT participants who had osteoarthritis, were ASA grades 1-3 and
underwent TKR as part of the trial, the mean baseline OKS was 18 (standard
deviation, SD: 7.5), mean age 71 (SD: 8.0), mean BMI 30 (SD: 5.5), and 45%
(956/2131) were female (Figure 1); further baseline characteristics of the trial
population have been reported previously.[28]

Mean EQ-5D utility rose substantially within three months of TKR from 0.39 at
baseline to 0.71 at one year, and declined gradually thereafter (Figure 1, Table 1).
However, baseline utility and the quality of life gains associated with TKR varied
between patient subgroups. In particular, women and patients with low OKS (i.e. poor
knee function), higher ASA grade or morbid obesity at baseline tended to have lower
quality of life at all timepoints (Figure 1), although those in the lowest OKS quintile
experienced significantly greater health gains from TKR than the top quintile
($p < 0.001$; calculated using linear regression combining imputed datasets using `mim`
command).

Table 1: EQ-5D utility, costs and estimated QALYs for subgroups with different baseline OKS

OKS decile		1 (worst)	2	3	4	5	6	7	8	9	10 (best)	All patients
Baseline OKS range		<9	9-11	12-13	14-15	16-17	18-19	20-21	22-24	25-27	>27	0-48
No. KAT participants		204	213	186	232	193	224	195	256	194	234	2131
EQ-5D utility (SE)	Baseline	0.54 (0.007)	0.58 (0.006)	0.64 (0.006)	0.66 (0.005)	0.71 (0.005)	0.71 (0.005)	0.70 (0.004)	0.74 (0.004)	0.61 (0.004)	0.67 (0.003)	0.39 (0.002)
	3 mths	0.51 (0.007)	0.60 (0.007)	0.68 (0.006)	0.70 (0.006)	0.71 (0.006)	0.75 (0.005)	0.69 (0.006)	0.78 (0.004)	0.75 (0.005)	0.77 (0.004)	0.68 (0.002)
	1 year	0.49 (0.008)	0.54 (0.007)	0.66 (0.007)	0.67 (0.006)	0.68 (0.007)	0.73 (0.005)	0.66 (0.007)	0.76 (0.005)	0.79 (0.005)	0.84 (0.004)	0.71 (0.002)
	2 year	0.47 (0.008)	0.53 (0.007)	0.64 (0.007)	0.63 (0.006)	0.67 (0.007)	0.71 (0.006)	0.68 (0.007)	0.72 (0.005)	0.78 (0.005)	0.81 (0.005)	0.68 (0.002)
	3 year	0.43 (0.008)	0.49 (0.008)	0.61 (0.007)	0.62 (0.007)	0.65 (0.007)	0.68 (0.007)	0.64 (0.007)	0.70 (0.006)	0.76 (0.006)	0.79 (0.006)	0.66 (0.002)
	4 year	0.43 (0.008)	0.50 (0.008)	0.57 (0.008)	0.58 (0.007)	0.63 (0.007)	0.67 (0.007)	0.59 (0.007)	0.68 (0.006)	0.71 (0.007)	0.75 (0.006)	0.63 (0.002)
	5 year	0.47 (0.006)	0.58 (0.005)	0.65 (0.005)	0.65 (0.005)	0.67 (0.008)	0.73 (0.007)	0.47 (0.006)	0.58 (0.005)	0.65 (0.005)	0.65 (0.005)	0.61 (0.002)
Total cost (SE)*		£8657 (£157)	£7715 (£81)	£7495 (£73)	£7081 (£63)	£7185 (£69)	£7567 (£82)	£7619 (£102)	£7128 (£61)	£7399 (£94)	£6917 (£67)	£7458 (£28)
QALYs with TKR (SE)*		2.17 (0.029)	2.48 (0.027)	2.94 (0.024)	2.99 (0.022)	3.13 (0.025)	3.30 (0.022)	3.09 (0.025)	3.40 (0.019)	3.50 (0.020)	3.67 (0.020)	3.08 (0.025)
QALYs without TKR (SE)*		0.15 (0.02)	0.62 (0.022)	0.94 (0.026)	1.14 (0.026)	1.63 (0.029)	2.04 (0.026)	2.39 (0.026)	2.55 (0.021)	2.78 (0.02)	3.02 (0.02)	1.75 (0.031)
QALY gain from TKR (SE)*		2.02 (0.032)	1.86 (0.031)	2.00 (0.033)	1.85 (0.031)	1.51 (0.034)	1.26 (0.030)	0.69 (0.027)	0.85 (0.023)	0.73 (0.02)	0.65 (0.02)	1.33 (0.01)
ICER*		£4,295	£4,153	£3,747	£3,836	£4,770	£6,007	£10,971	£8,391	£10,167	£10,697	£5,623

* Discounted at 3.5% per annum

Abbreviations: ICER, incremental cost-effectiveness ratio; OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])[40]; QALY, quality-adjusted life-year; SE, standard error around the mean; TKR, total knee replacement.

On average, each admission for primary TKR cost £6,363 (SD: £1,702). Readmissions, revision procedures and GP, outpatient, and physiotherapy consultations related to the study knee over the following five years cost a further £1,095 (SD: £3,579) per patient, giving a total cost of £7,458 (SD: £4,058) per patient. TKR was more costly in patients with low baseline OKS, but the greater QALY gains meant that TKR was better value for money in this patient group than in patients with moderate symptoms (Table 1). Subgroup analyses suggested that TKR cost less than £11,000/QALY gained across all OKS deciles: well below the £20,000-£30,000/QALY gained range that NICE currently consider good value for money.[11]

However, cost-effectiveness also varied substantially between patients within each OKS decile. We therefore used regression models to estimate the costs, QALY gains and cost-effectiveness of TKR in a wide range of patient subgroups while controlling for multiple baseline characteristics. Our results indicated that four factors have a significant effect on cost-effectiveness: age, sex, baseline OKS and ASA grade (Table 2). BMI and the presence of arthritis in other joints besides the study knee had no significant effect on either the costs or QALY gains associated with TKR ($p>0.05$).

Table 2: Results of the regression analysis

Baseline characteristic	Coefficient (SE)		
	QALY loss with TKR†	QALY loss without TKR†	Total costs with TKR
Female sex	0.056 (0.065)	-0.001 (0.054)	593 (181)*
Age at operation (years)	0.007 (0.003)*	0.002 (0.004)	-8 (12)
Pre-operative OKS	-0.040 (0.004)*	-0.087 (0.004)*	-52 (12)*
ASA grade 3	0.414 (0.067)*	0.227 (0.087)*	492 (208)*
Constant	1.712 (0.201)*	4.274 (0.325)*	8573 (881)*

* $p<0.05$

† QALY loss indicates the number of discounted QALYs that would have been accrued if patients had experienced an EQ-5D utility of 1 for five years (4.67 QALYs) minus the number of discounted QALYs that the patient actually experienced.

The incremental cost-effectiveness ratio (expressed as cost/QALY) for any given patient equals: $(\text{female} \times 593 - \text{age} \times 8 - \text{OKS} \times 52 + \text{ASAgrade3or4} \times 492 + 8573) / ((\text{female} \times -0.001 + \text{age} \times 0.002 - \text{OKS} \times 0.087 + \text{ASAgrade3} \times 0.227 + 4.274) - (\text{female} \times 0.056 + \text{age} \times 0.007 - \text{OKS} \times 0.040 + \text{ASAgrade3} \times 0.414 + 1.712))$. Costs and QALYs were discounted at 3.5% per annum.

Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy, 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life); OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])[40]; QALY, quality-adjusted life-year; TKR, total knee replacement.

Figure 2 shows the cost-effectiveness of TKR for each multivariate group, using charts similar in design to widely-used cardiovascular risk tables.[42] The charts demonstrate that TKR is clearly cost-effective across a wide range of patient groups. For example, TKR costs less than £20,000/QALY gained for 71-year-old men with no symptomatic illness unless baseline OKS is above 40 (equivalent to having very

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

minor problems on up to eight of the 12 activities mentioned in OKS or moderate-severe pain only on exertion). The maximum OKS at which TKR becomes cost-ineffective was lower for older patients, women, and those with symptomatic disease, but TKR remained cost-effective for all patients with OKS below 32 regardless of age, sex, or ASA grade. A simplified analysis controlling only for OKS and ASA grade suggested that TKR is cost-effective for all ASA grade 1 or 2 patients with baseline OKS of 39 or less, and all ASA grade 3 patients with OKS ≤ 34 .

There was substantial uncertainty around the cost-effectiveness ratios for patients with very high baseline OKS, since only 37 trial participants had OKS above 35 at baseline. However, we estimate that there is a 99% probability that TKR is cost-effective (at a £20,000/QALY ceiling ratio) for the average ASA grade 1-2 patient with a baseline OKS of 35, and a 53% probability for ASA grade 3 (Figure 3).

Since each additional baseline characteristic increases the complexity of the eligibility criteria and may also raise ethical or equity concerns, we estimated the number of patients who would receive TKR under different eligibility criteria. This analysis suggested that restricting TKR to those patients for whom the operation was predicted to cost less than £20,000/QALY gained based on the simplified model (allowing only for ASA grade and OKS) would lead to 2000 fewer TKR procedures being conducted in England and Wales each year (3% fewer than the 61,651 procedures conducted in 2009/10[26]). If the decision rules were simplified further to provide TKR to all patients with baseline OKS ≤ 38 , all but 1700 of the 61,651 patients currently receiving TKR in England and Wales in 2009/10[26] would be eligible. Allowing for other baseline characteristics had minimal effect on patient numbers or predicted costs or benefits, suggesting that decisions about suitability for TKR can be based on OKS alone.

Sensitivity analyses were conducted to assess the effect of extending the time horizon of our analysis up to 10 years, and of assuming that EQ-5D utility would have fallen by 0.5% per year without TKR,[33] that all knee-related ambulatory consultations would have occurred without TKR, and that the mean length of stay has fallen by 41% (5.9 days[26] vs 10.0) since the KAT operations took place in 1999-2003. Changing any of these assumptions produced lower ICERs than the base case analysis, confirming that the current analysis is highly conservative and suggesting that TKR may be cost-effective for ASA grade 1-2 patients with baseline

1
2
3 OKS as high as 43. Excluding patients with incomplete data had no effect on
4 conclusions.
5
6
7

8 **Discussion**

9
10 We found TKR to cost £5,623 per QALY gained for the average patient (similar to
11 estimates from previous studies[6-10]) and therefore represent very good value for
12 money compared with the £20,000-£30,000/QALY threshold typically used in NHS
13 decision-making.[11] Our study also confirms previous findings[7, 9, 16] that costs
14 and benefits vary with age and comorbidity. We found pre-operative OKS to be the
15 best predictor of post-operative costs, outcomes and cost-effectiveness.
16
17
18
19

20
21 Our analyses were based on patient-level data from a large pragmatic trial with
22 detailed, prospective collection of costs, utilities and baseline characteristics that
23 permit extensive subgroup analyses. Furthermore, unlike some previous economic
24 evaluations, our study took full account of uncertainty and included all knee-related
25 NHS costs accrued in the first five years after TKR. The main limitation of our
26 analysis was the small number of patients with high OKS. Since KAT recruited only
27 37 patients with OKS >35, further studies are required to evaluate TKR in this patient
28 subgroup. Analyses were also based on *post hoc* comparisons between outcomes
29 before and after TKR, rather than between randomly-assigned treatment groups. As
30 a result, our study does not provide unbiased estimates of the QALY gains from TKR,
31 and regression towards the mean may have caused our study to overestimate the
32 quality of life improvement attributable to TKR and/or the effect of pre-operative OKS
33 on QALY gains.
34
35
36
37
38
39
40
41
42

43
44 We made several highly conservative assumptions: in particular, assuming that
45 patients would have accrued no knee-related costs and would have remained at
46 baseline utility without TKR, whereas previous studies have found that patients
47 accrue substantial costs and show clinical worsening without TKR.[32] We also
48 assumed that those patients who died before hospital discharge would have survived
49 for five years if they had not had TKR; in reality, those patients dying soon after TKR
50 tend to be older[5, 34] and frailer[5] pre-operatively and some studies suggest that
51 patients undergoing hip replacement have higher long-term survival than the general
52 population[43, 44] (although this may be due to patient selection[34, 43]). When
53 these assumptions were relaxed, TKR was cost-effective for ASA grade 1-2 patients
54 with baseline OKS \leq 43. The costs and benefits of TKR are likely to vary between
55
56
57
58
59
60

1
2
3 centres and over time due to variations in patient demographics, component prices,
4 length of stay, and operation time; sensitivity analyses demonstrated that TKR may
5 have become more cost-effective since the KAT procedures were conducted in 1999-
6 2003 due to reductions in length of stay.
7
8
9

10
11 Our analyses show that if TKR were to be rationed based on cost-effectiveness, OKS
12 would be a reasonable tool to use to set the threshold, although there may, of
13 course, be practical difficulties with using a patient-reported measure to determine
14 treatment eligibility: particularly if patients know how their questionnaire responses
15 will be used or if test-retest reliability is low in this setting. Based on regression and
16 subgroup analyses, we can be confident that TKR costs less than £20,000/QALY for
17 all ASA grade 1-2 patients with baseline OKS ≤ 39 and for ASA grade 3 patients with
18 OKS ≤ 34 . Using these thresholds to determine who received treatment would avoid
19 around 2000 TKR procedures per year in England and Wales, saving around £11.8
20 million per year. However, the higher thresholds suggested by sensitivity analyses
21 using less conservative assumptions suggest that all TKR is cost-effective for all but
22 100 patients per year, which is unlikely to achieve sufficient savings to warrant
23 rationing of TKR. Although the cost-effectiveness of knee replacement was also
24 influenced by gender and age, taking account of these factors as well as OKS and
25 ASA grade would increase the complexity of eligibility criteria, reduce equality of
26 access to healthcare and have minimal effect on NHS budgets.
27
28
29
30
31
32
33
34
35
36
37
38

39 Our analysis shows that the thresholds proposed by some PCTs[17-23] are
40 inappropriate and would deny a highly cost-effective treatment to thousands of
41 patients with severe arthritis. The current situation also introduces a postcode lottery
42 with different PCTs using different eligibility criteria. We estimate that limiting knee
43 replacement to patients with OKS ≤ 26 would prevent more than 10,000 people from
44 receiving TKR. This would be inequitable as TKR costs £10,697 per QALY gained for
45 this group, whereas NICE routinely recommend treatments with ICERs in the region
46 of £20,000-£30,000/QALY gained[11, 45, 46] and are willing to pay substantially
47 more for end of life care.[47] Furthermore, the finding that TKR and subsequent
48 monitoring is more costly for patients with low OKS suggests that delaying TKR until
49 symptoms have deteriorated may be a false economy. Although some PCTs have
50 considered restricting TKR further for obese patients,[17, 20, 24, 48] we found BMI to
51 have no significant effect on costs ($p=0.442$), QALY gains ($p=0.098$) or the incidence
52 of peri- or post-operative complications ($p>0.26$): even when ASA grade was omitted
53 from regression analyses.
54
55
56
57
58
59
60

1
2
3
4
5 Eligibility criteria for rheumatoid arthritis patients and patients with ASA grade 4
6 remain unclear, although ASA grade 4 patients would normally be considered unfit
7 for surgery, and the decision to operate on such patients would be based on a case-
8 by-case consideration of whether the benefits outweigh the risks. While our study
9 focused on the costs and benefits of TKR, patients with high baseline OKS often
10 undergo unicompartmental knee replacement, which is cheaper than TKR due to
11 lower component prices, fewer complications and shorter inpatient stay.[49-51] The
12 OKS threshold for unicompartmental replacement may therefore be even higher than
13 for TKR, since it produces marked quality of life improvements in patients with high
14 baseline scores.[49] This remains an area for future research.
15
16
17
18
19
20
21
22

23 **Acknowledgements**

24
25 The authors would like to thank all those involved in the KAT trial for their role in
26 running the trial and collecting the data used in this study, particularly other members
27 of the KAT Project Management Group: Suzanne Breeman; Marion Campbell; Jackie
28 Ellington; Nick Fiddian; Adrian Grant; Linda Johnston; and Richard Morris.
29
30
31
32

33 The Knee Arthroplasty Trial is funded by the NIHR Health Technology Assessment
34 Programme (project number 95/10/01) and will be published in full as a Health
35 Technology Assessment report. Visit the HTA programme website for further project
36 information. Additional funding for research support in clinical centres was provided
37 by Howmedica Osteonics; Zimmer; DePuy, a Johnson and Johnson company; Corin
38 Medical; Smith and Nephew Healthcare; Biomet Merck; and Wright Cremascoli. The
39 Health Economics Research Centre obtains support from the National Institute of
40 Health Research. The Health Services Research Unit is core funded by the Chief
41 Scientist Office of the Scottish Government Health Directorates. PROMs data were
42 reused with the permission of The Health and Social Care Information Centre,
43 copyright 2011, with all rights reserved. The views and opinions expressed therein
44 are those of the authors and do not necessarily reflect those of the HTA programme,
45 NIHR, NHS, Information Centre or the Department of Health.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Competing interests

All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare no support from any organisation for the submitted work with the exception of those declared in the funding statement. AG, RF and GM declare that they have no financial relationships with any organisations that might have an interest in the submitted work in the previous three years and no other relationships or activities that could appear to have influenced the submitted work. DM receives royalties paid to him by Biomet and HD has undertaken consultancy work for Pfizer on drugs for rheumatoid arthritis but both declare that they have no other financial relationships with any organisations that might have an interest in the submitted work in the previous three years and no other relationships or activities that could appear to have influenced the submitted work.

Funding

The Knee Arthroplasty Trial is funded by the NIHR Health Technology Assessment Programme (project number 95/10/01). Additional funding for research support in clinical centres was provided by Howmedica Osteonics; Zimmer; DePuy, a Johnson and Johnson company; Corin Medical; Smith and Nephew Healthcare; Biomet Merck; and Wright Cremascoli.

The Health Economics Research Centre obtains support from the National Institute of Health Research. The Health Services Research Unit is core funded by the Chief Scientist Office of the Scottish Government Health Directorates.

The authors conducted the research independently of all funding organisations. The funders had no role in the collection, analysis or interpretation of data, writing of the manuscript or the decision to publish. The authors had full access to all of the data (including statistical reports and tables) in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Contributorship statement

HD, AG and DM conceived and developed the principles and methods underpinning the current analysis. GM was involved in primary data collection and database management, while HD conducted the analysis under AG's supervision and guidance from other authors. HD drafted the manuscript with input from all other authors; all

1
2
3 authors have approved the final manuscript. The KAT trial was designed and run by
4 the KAT trial group, which comprised Suzanne Breeman, Marion Campbell, HD,
5 Jackie Ellington, Nick Fiddian, RF, Adrian Grant, AG, Linda Johnston, GM, Richard
6 Morris and DM.
7
8
9

10 11 12 **References**

- 13
14 1 Hospital Episode Statistics. Provisional Monthly Patient Reported Outcome
15 Measures (PROMs) in England. April 2009 – April 2010: Pre- and postoperative data:
16 Experimental statistics. 2010 [updated 2010; cited 6th April 2011]; Available from:
17 <http://www.hqip.org.uk/assets/NCAPOP-Library/PROMsreport09101.pdf>.
18
19 2 Labek G, Thaler M, Janda W, et al. Revision rates after total joint replacement:
20 cumulative results from worldwide joint register datasets. *J Bone Joint Surg Br*
21 2011;**93**:293-7.
22
23 3 The National Collaborating Centre for Chronic Conditions. Osteoarthritis: National
24 clinical guideline for care and management in adults. London: Royal College of
25 Physicians; 2008 [updated 2008; cited 29th June 2011]; Available from:
26 <http://www.nice.org.uk/nicemedia/live/11926/39720/39720.pdf>.
27
28 4 Huddleston JI, Maloney WJ, Wang Y, et al. Adverse events after total knee
29 arthroplasty: a national Medicare study. *J Arthroplasty* 2009;**24**:95-100.
30
31 5 Memtsoudis SG, Della Valle AG, Besculides MC, et al. Risk factors for
32 perioperative mortality after lower extremity arthroplasty: a population-based study of
33 6,901,324 patient discharges. *J Arthroplasty* 2010;**25**:19-26.
34
35 6 Navarro Espigares JL, Hernandez Torres E. Cost-outcome analysis of joint
36 replacement: evidence from a Spanish public hospital. *Gac Sanit* 2008;**22**:337-43.
37
38 7 Losina E, Walensky RP, Kessler CL, et al. Cost-effectiveness of total knee
39 arthroplasty in the United States: patient risk and hospital volume. *Arch Intern Med*
40 2009;**169**:1113-21; discussion 21-2.
41
42 8 Rasanen P, Paavolainen P, Sintonen H, et al. Effectiveness of hip or knee
43 replacement surgery in terms of quality-adjusted life years and costs. *Acta Orthop*
44 2007;**78**:108-15.
45
46 9 Rissanen P, Aro S, Sintonen H, et al. Costs and cost-effectiveness in hip and knee
47 replacements. A prospective study. *Int J Technol Assess Health Care* 1997;**13**:575-
48 88.
49
50 10 Lavernia CJ, Guzman JF, Gachupin-Garcia A. Cost effectiveness and quality of
51 life in knee arthroplasty. *Clin Orthop Relat Res* 1997:134-9.
52
53
54
55
56
57
58
59
60

1
2
3 11 National Institute for Health and Clinical Excellence. Social value judgements:
4 Principles for the development of NICE guidance. Second Edition. 2008 [updated
5 2008; cited 17th November 2010]; Available from:

6
7
8 <http://www.nice.org.uk/media/C18/30/SVJ2PUBLICATION2008.pdf>.

9
10 12 Brander VA, Malhotra S, Jet J, et al. Outcome of hip and knee arthroplasty in
11 persons aged 80 years and older. Clin Orthop Relat Res 1997;67-78.

12
13 13 Karuppiah SV, Banaszkiwicz PA, Ledingham WM. The mortality, morbidity and
14 cost benefits of elective total knee arthroplasty in the nonagenarian population. Int
15 Orthop 2008;**32**:339-43.

16
17 14 McQueen DA, Long MJ, Algotar AM, et al. The effect of obesity on quality-of-life
18 improvement after total knee arthroplasty. Am J Orthop (Belle Mead NJ)
19 2007;**36**:E117-20, E27.

20
21 15 de Leeuw JM, Villar RN. Obesity and quality of life after primary total knee
22 replacement Knee 1998;**5**:119-23.

23
24 16 Cushnaghan J, Bennett J, Reading I, et al. Long-term outcome following total
25 knee arthroplasty: a controlled longitudinal study. Ann Rheum Dis 2009;**68**:642-7.

26
27 17 NHS Oxfordshire. INTERIM Treatment Threshold Statement: Knee arthroplasty
28 (Reference PS 188). 2010 [updated 2010; cited 5th April 2011]; Available from:

29
30
31
32 [http://www.oxfordshirepct.nhs.uk/professional-resources/priority-setting/lavender-](http://www.oxfordshirepct.nhs.uk/professional-resources/priority-setting/lavender-statements/documents/PS188Kneereplacement.pdf)
33 [statements/documents/PS188Kneereplacement.pdf](http://www.oxfordshirepct.nhs.uk/professional-resources/priority-setting/lavender-statements/documents/PS188Kneereplacement.pdf).

34
35 18 NHS Gloucestershire. Guidelines for Management of Orthopaedic Conditions.
36 2010 [updated 2010; cited 5th April 2011]; Available from:

37
38
39 [http://www.glospct.nhs.uk/pdf/services/orthopaedic_referral_guidelines_%20march%](http://www.glospct.nhs.uk/pdf/services/orthopaedic_referral_guidelines_%20march%2010v3.pdf)
40 [2010v3.pdf](http://www.glospct.nhs.uk/pdf/services/orthopaedic_referral_guidelines_%20march%2010v3.pdf).

41
42 19 NHS Worcestershire. Commissioning policy for musculoskeletal surgical
43 interventions. 2010 [updated 2010; cited 5th April 2011]; Available from:

44
45
46 [http://www.worcestershire.nhs.uk/file_download.aspx?id=86bae3e2-4d88-4416-](http://www.worcestershire.nhs.uk/file_download.aspx?id=86bae3e2-4d88-4416-92e6-9eeda22e8208)
47 [92e6-9eeda22e8208](http://www.worcestershire.nhs.uk/file_download.aspx?id=86bae3e2-4d88-4416-92e6-9eeda22e8208).

48
49 20 Milton Keynes O, Berkshire East, Berkshire West & Buckinghamshire (MOBBB)
50 Priorities Committee,. Minutes of meeting held on Wednesday 15 December 2010.

51
52 2010 [updated 2010; cited 5th April 2011]; Available from: [http://www.sph.nhs.uk/sph-](http://www.sph.nhs.uk/sph-psu/sph-files/mobbb-minutes/mobbb-minutes-15-december-2010)
53 [psu/sph-files/mobbb-minutes/mobbb-minutes-15-december-2010](http://www.sph.nhs.uk/sph-psu/sph-files/mobbb-minutes/mobbb-minutes-15-december-2010).

54
55 21 South West London Public Health Network. 2010/11 South West London Effective
56 Commissioning Initiative. 2010 [updated 2010; cited 5th April 2011]; Available from:

57
58
59 [http://www.swlphnetwork.org.uk/ uploads/documents/effective-commissioning-](http://www.swlphnetwork.org.uk/uploads/documents/effective-commissioning-document-master-version-2010-11-final.doc)
60 [document-master-version-2010-11-final.doc](http://www.swlphnetwork.org.uk/uploads/documents/effective-commissioning-document-master-version-2010-11-final.doc).

1
2
3 22 NHS Manchester. Topics for inclusion or change in the pct non-commissioned
4 procedures list. 2010 [updated 2010; cited 5th April 2011]; Available from:
5
6 http://www.manchester.nhs.uk/document_uploads/19%20January%202011/G2turnar
7 [oundAppendix%20%20TR.doc.](http://www.manchester.nhs.uk/document_uploads/19%20January%202011/G2turnar)
8

9
10 23 NHS Derbyshire County. NHS Derbyshire County Prior Approval: Procedures of
11 Limited Clinical Value Summary. 2010 [updated 2010; cited 29th June 2011];
12 Available from:
13
14 <http://www.phcn.nhs.uk/Derbyshire%20County%20Procedures%20of%20Limited%2>
15 [0Clinical%20Value%20100525%20draft.pdf.](http://www.phcn.nhs.uk/Derbyshire%20County%20Procedures%20of%20Limited%2)
16
17

18 24 Jeffreys B. Surgeons raise alarm over waiting. Journal [serial on the Internet].
19 2011 Date: Available from: <http://www.bbc.co.uk/news/health-12964360>.

20
21 25 NHS Warwickshire. Report to the PCT board 10th November 2010. 2010 [updated
22 2010 6th April 2011; cited 29th June 2011]; Available from:
23
24 <http://www.warwickshire.nhs.uk/trustboardenclosures/November2010/Enc%208%20>
25 [%20In%20Year%20Demand%20Management%20of%20Electives%205-11-](http://www.warwickshire.nhs.uk/trustboardenclosures/November2010/Enc%208%20)
26 [2010.pdf.](http://www.warwickshire.nhs.uk/trustboardenclosures/November2010/Enc%208%20)
27
28

29 26 Hospital Episode Statistics. Main procedures and interventions: 4 character
30 inpatient data 2009-10. 2010 [updated 2010; cited 10th February 2011]; Available
31 from:
32
33 <http://www.hesonline.nhs.uk/Ease/servlet/ContentServer?siteID=1937&categoryID=2>
34 [15.](http://www.hesonline.nhs.uk/Ease/servlet/ContentServer?siteID=1937&categoryID=2)
35
36

37 27 Dawson J, Fitzpatrick R, Murray D, et al. Questionnaire on the perceptions of
38 patients about total knee replacement. J Bone Joint Surg Br 1998;**80**:63-9.

39 28 Johnston L, MacLennan G, McCormack K, et al. The Knee Arthroplasty Trial
40 (KAT) design features, baseline characteristics, and two-year functional outcomes
41 after alternative approaches to knee replacement. J Bone Joint Surg Am
42 2009;**91**:134-41.
43
44

45 29 Breeman S, Campbell M, Dakin H, et al. Patellar Resurfacing in Total Knee
46 Replacement: Five-Year Clinical and Economic Results of a Large Randomized
47 Controlled Trial. J Bone Joint Surg Am 2011;**93**:1473-81.
48
49

50 30 Dolan P. Modeling valuations for EuroQol health states. Med Care 1997;**35**:1095-
51 108.
52
53

54 31 HM Treasury. The Green Book: Appraisal and Evaluation in Central Government.
55 London: TSO; 2003.
56
57

58 32 Hawker GA, Badley EM, Croxford R, et al. A population-based nested case-
59 control study of the costs of hip and knee replacement surgery. Med Care
60 2009;**47**:732-41.

- 1
2
3 33 Kind P, Dolan P, Gudex C, et al. Variations in population health status: results
4 from a United Kingdom national questionnaire survey. *Bmj* 1998;**316**:736-41.
5
6 34 Lie SA, Pratt N, Ryan P, et al. Duration of the increase in early postoperative
7 mortality after elective hip and knee replacement. *J Bone Joint Surg Am* 2010;**92**:58-
8 63.
9
10 35 Royston P. Multiple imputation of missing values. *Stata Journal* 2004;**4**:227-41.
11
12 36 Royston P. Multiple imputation of missing values: further update of ice, with an
13 emphasis on interval censoring. *Stata Journal* 2007;**7**:445-64.
14
15 37 White IR, Royston P, Wood AM. Multiple imputation using chained equations:
16 Issues and guidance for practice. *Stat Med* 2011;**30**:377-99.
17
18 38 Briggs A, Clark T, Wolstenholme J, et al. Missing... presumed at random: cost-
19 analysis of incomplete data. *Health Econ* 2003;**12**:377-92.
20
21 39 Yu LM, Burton A, Rivero-Arias O. Evaluation of software for multiple imputation of
22 semi-continuous data. *Stat Methods Med Res* 2007;**16**:243-58.
23
24 40 Murray DW, Fitzpatrick R, Rogers K, et al. The use of the Oxford hip and knee
25 scores. *J Bone Joint Surg Br* 2007;**89**:1010-4.
26
27 41 Fenwick E, Claxton K, Sculpher M. Representing uncertainty: the role of cost-
28 effectiveness acceptability curves. *Health Econ* 2001;**10**:779-87.
29
30 42 British Medical Association. British National Formulary 61: Cardiovascular risk
31 prediction charts for primary prevention. March 2011 [updated March 2011; cited
32 12th April 2011]; Available from: <http://bnf.org/bnf/bnf/current/204016.htm>.
33
34 43 Barrett J, Losina E, Baron JA, et al. Survival following total hip replacement. *J*
35 *Bone Joint Surg Am* 2005;**87**:1965-71.
36
37 44 Lie SA, Engesaeter LB, Havelin LI, et al. Mortality after total hip replacement: 0-
38 10-year follow-up of 39,543 patients in the Norwegian Arthroplasty Register. *Acta*
39 *Orthop Scand* 2000;**71**:19-27.
40
41 45 Dakin HA, Devlin NJ, Odeyemi IA. "Yes", "No" or "Yes, but"? Multinomial
42 modelling of NICE decision-making. *Health Policy* 2006;**77**:352-67.
43
44 46 Devlin N, Parkin D. Does NICE have a cost-effectiveness threshold and what
45 other factors influence its decisions? A binary choice analysis. *Health Econ*
46 2004;**13**:437-52.
47
48 47 National Institute for Health and Clinical Excellence. Supplementary guidance on
49 appraising life-extending, end of life treatments. July 2009 [updated July 2009; cited
50 23rd February 2011]; Available from:
51 <http://www.nice.org.uk/media/E4A/79/SupplementaryAdviceTACEoL.pdf>.
52
53 48 Coombes R. Rationing of joint replacements raises fears of further cuts. *BMJ*
54 2005;**331**:1290.
55
56
57
58
59
60

1
2
3 49 Xie F, Lo NN, Tarride JE, et al. Total or partial knee replacement? Cost-utility
4 analysis in patients with knee osteoarthritis based on a 2-year observational study.
5 Eur J Health Econ 2010;**11**:27-34.
6
7

8 50 Robertsson O, Borgquist L, Knutson K, et al. Use of unicompartmental instead of
9 tricompartmental prostheses for unicompartmental arthrosis in the knee is a cost-
10 effective alternative. 15,437 primary tricompartmental prostheses were compared
11 with 10,624 primary medial or lateral unicompartmental prostheses. Acta Orthop
12 Scand 1999;**70**:170-5.
13
14
15

16 51 Slover J, Espehaug B, Havelin LI, et al. Cost-effectiveness of unicompartmental
17 and total knee arthroplasty in elderly low-demand patients. A Markov decision
18 analysis. J Bone Joint Surg Am 2006;**88**:2348-55.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Figure 1: EQ-5D utility by subgroup

For peer review only

Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy, 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life, 4=symptomatic illness with severe restriction on life); OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])[40]; TKR, total knee replacement. Error bars show 95% confidence intervals.

1
2
3 **Figure 2:** Cost-effectiveness prediction charts estimated based on predictions of
4 regression models.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32


33 Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy,
34 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life);
35 OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to
36 48 [no problems])[40]; QALY, quality-adjusted life-year; TKR, total knee replacement.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 3: Cost-effectiveness acceptability curves for different patient subgroups at different baseline Oxford Knee Scores (OKS). A: American Society of Anesthesiologists (ASA) grade 1 and 2 patients. B: ASA grade 3 patients.

For peer review only



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49



Simplified model: OKS and ASA only



Full model



Legend

- Clearly cost-effective: <£20,000/QALY gained
- Cost-ineffective: >£30,000/QALY gained
- Possibly cost-effective: £20,000-£30,000/QALY gained
- TKR dominated (no clinical benefit)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49



Health economic checklist for “Rationing of total knee replacement: A cost-effectiveness analysis on a large trial dataset”

Study design	Page
(1) The research question is stated	5, paragraphs 2 & 3
(2) The economic importance of the research question is stated	4-5
(3) The viewpoint(s) of the analysis are clearly stated and justified	5, paragraph 3
(4) The rationale for choosing the alternative programmes or interventions compared is stated	4-5
(5) The alternatives being compared are clearly described	5, paragraph 3
(6) The form of economic evaluation used is stated	5, paragraph 3
(7) The choice of form of economic evaluation is justified in relation to the questions addressed	5, paragraph 3
Data collection	
(8) The source(s) of effectiveness estimates used are stated	5, paragraph 4
(9) Details of the design and results of effectiveness study are given (if based on a single study)	5, paragraph 4
(10) Details of the method of synthesis or meta-analysis of estimates are given (if based on an overview of a number of effectiveness studies)	N/A
(11) The primary outcome measure(s) for the economic evaluation are clearly stated	5, paragraph 3
(12) Methods to value health states and other benefits are stated	6, paragraph 1
(13) Details of the subjects from whom valuations were obtained are given	6, paragraph 1
(14) Productivity changes (if included) are reported separately	N/A
(15) The relevance of productivity changes to the study question is discussed	5, paragraph 3
(16) Quantities of resources are reported separately from their unit costs	Given in reference 29
(17) Methods for the estimation of quantities and unit costs are described	Given in reference 29
(18) Currency and price data are recorded	N/A
(19) Details of currency of price adjustments for inflation or currency conversion are given	Given in reference 29
(20) Details of any model used are given	N/A
(21) The choice of model used and the key parameters on which it is based are justified	N/A
Analysis and interpretation of results	
(22) Time horizon of costs and benefits is stated	5, paragraph 4
(23) The discount rate(s) is stated	6, paragraph 2
(24) The choice of rate(s) is justified	6, paragraph 2
(25) An explanation is given if costs or benefits are not discounted	N/A
(26) Details of statistical tests and confidence intervals are given for stochastic data	8, paragraph 1
(27) The approach to sensitivity analysis is given	11, paragraph 4
(28) The choice of variables for sensitivity analysis is justified	11, paragraph 4
(29) The ranges over which the variables are varied are stated	N/A
(30) Relevant alternatives are compared	5, paragraph 3
(31) Incremental analysis is reported	Tables 1 & 2
(32) Major outcomes are presented in a disaggregated as well as aggregated form	Tables 1 & 2
(33) The answer to the study question is given	12, paragraph 2 & page 13 paragraphs 2-3
(34) Conclusions follow from the data reported	5, 12-13
(35) Conclusions are accompanied by the appropriate caveats	12-13

Authors may enter N/A if an item on the checklist is not appropriate, but this is only acceptable for items 9,10, 12-15, 20, 21, 23-29, and 31.



Rationing of total knee replacement: A cost-effectiveness analysis on a large trial dataset

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2011-000332.R1
Article Type:	Research
Date Submitted by the Author:	20-Dec-2011
Complete List of Authors:	Dakin, Helen; University of Oxford, Health Economics Research Centre, Department of Public Health Gray, Alastair; University of Oxford, Health Economics Research Centre, Department of Public Health Fitzpatrick, Ray; University of Oxford, Department of Public Health Maclennan, Graeme; University of Aberdeen, Health Services Research Unit Murray, David; University of Oxford, Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences
Primary Subject Heading:	Health services research
Secondary Subject Heading:	Rheumatology, Health policy, Health economics, Surgery
Keywords:	Economic evaluation, total knee replacement, heterogeneity, cost-utility analysis, randomised controlled trial

SCHOLARONE™
Manuscripts

Rationing of total knee replacement: A cost-effectiveness analysis on a large trial dataset

Running header:

Rationing of total knee replacement

Key words:

Economic evaluation; total knee replacement; heterogeneity; subgroup analysis; randomised controlled trial.

Authors:

Helen Dakin,¹ Alastair Gray,¹ Ray Fitzpatrick,² Graeme MacLennan,³ and David Murray⁴

Collaborator: The KAT Trial Group

Authors' affiliation:

¹ Health Economics Research Centre, Department of Public Health, University of Oxford, Old Road Campus, Oxford, UK

² Department of Public Health, University of Oxford, Old Road Campus, Oxford, UK

³ Health Services Research Unit, University of Aberdeen, Aberdeen, UK

⁴ Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, UK

Corresponding author:

Helen Dakin

Researcher

Health Economics Research Centre

Department of Public Health, University of Oxford

Old Road Campus

Headington

Oxford

OX3 7LF

Email: helen.dakin@dph.ox.ac.uk

Tel: 01865 289422

Fax: 01865 289271

Word count: 3,691 (main text of introduction, methods, results and discussion)

Number of figures: 3

Number of tables: 2

Ethics: The KAT trial was approved by the Multi Centre Research Ethics Committee for Scotland in November 1998 (research protocol MREC/98/0/100) and was approved by the Local Research Ethics Committees in each study centre recruiting trial participants. Further details are available on request.

Data sharing statement: No additional data available.

Protocol: Available at <http://www.hta.ac.uk/protocols/199500100001.pdf>.

Abstract

Objectives: Many UK primary care trusts (PCTs) have recently introduced eligibility criteria restricting total knee replacement (TKR) to patients with low pre-operative Oxford Knee Scores (OKS) to cut expenditure. We evaluate these criteria by assessing the cost-effectiveness of TKR compared with no knee replacement for patients with different baseline characteristics from an NHS perspective.

Design: The cost-effectiveness of TKR in different patient subgroups was assessed using regression analyses of patient-level data from the Knee Arthroplasty Trial (KAT), a large, pragmatic, randomised trial comparing knee prostheses.

Setting: 34 UK hospitals.

Participants: 2131 osteoarthritis patients undergoing TKR.

Interventions and outcome measures: Costs and quality-adjusted life-years (QALYs) observed in the KAT trial within five years of TKR were compared with conservative assumptions about the costs and outcomes that would have been accrued had TKR not been performed.

Results: On average, primary TKR and five years of subsequent care cost £7,458 per patient (standard deviation, SD: £4,058), and patients gained an average of 1.33 (SD: 1.43) QALYs. As result, TKR cost £5,623 per QALY gained. Although costs and health outcomes varied with age and sex, TKR cost less than £20,000 per QALY gained for patients with ASA grade 1-2 who had baseline OKS <40, and for ASA grade 3 patients with OKS <35, even with highly conservative assumptions about costs and outcomes without TKR. Body mass index had no significant effect on costs or outcomes. Restricting TKR to patients with pre-operative OKS <27 would inappropriately deny a highly cost-effective treatment to >10,000 patients annually.

Conclusions: TKR is highly cost-effective for most current patients if the NHS is willing to pay £20,000-£30,000 per QALY gained. At least 97% of TKR patients in England have more severe symptoms than the thresholds we have identified, suggesting further rationing by OKS is probably unjustified.

Trial registration: ISRCTN 45837371

Article summary

Article focus

- We assess the cost-effectiveness of total knee replacement (TKR) compared with no knee replacement for patients with different baseline characteristics from a National Health Service (NHS) perspective.
- In particular, we assess the appropriateness of eligibility criteria recently introduced by many UK primary care trusts (PCTs), which restrict TKR to patients with low (i.e. poor) pre-operative Oxford Knee Scores (OKS) to cut expenditure.

Key messages

- We find TKR to be highly cost-effective, costing £5,623 per quality-adjusted life-year (QALY) gained for the average patient.
- TKR costs less than £20,000 per QALY gained for healthy patients with OKS of <40, or <35 for patients who have other conditions restricting their daily activities.
- We find no evidence to support the criteria for restricting access to TKR that have been proposed by some PCTs and calculate that restricting TKR to those patients with pre-operative OKS of 26 or less would deny a highly cost-effective treatment to more than 10,000 patients per year.

Strengths and limitations of this study

- This is the first study assessing how the cost-effectiveness of TKR varies with OKS and the first assessing the clinical/economic implications of the newly-introduced rationing criteria.
- Analyses are based on patient-level data from a large pragmatic trial with detailed, prospective collection of utilities, baseline characteristics and all major knee-related NHS resource use, including revisions and ambulatory care.
- Our study makes several highly conservative assumptions: in particular, assuming that patients would have accrued no knee-related costs and remained at baseline utility without TKR. Furthermore, the KAT sample included only 37 patients with pre-operative OKS >35. As result, TKR may be also cost-effective for some patients with OKS above 39.

Introduction

Total knee replacement (TKR) is a highly effective treatment for patients with bone-on-bone osteoarthritis and significant knee symptoms, producing substantial reductions in symptoms (particularly pain)[1] and lasting for at least 15 years in 83-94% of cases.[2] However, about 18% of patients consider the outcomes of their surgery to be only fair or poor,[1] and a small proportion experience complications.[3-5] Previous economic evaluations have found TKR to be highly cost-effective, with a cost-effectiveness ratio between €1,276 and \$18,300 per quality-adjusted life-year (QALY) gained for the average patient[6-10]: well below the £20,000-£30,000 per QALY range that the National Institute for Health and Clinical Excellence (NICE) considers cost-effective.[11] However, studies suggest that the costs and benefits of TKR vary between patient subgroups, with TKR being more cost-effective in younger patients,[9] those attending high-volume centres[7] and those deemed low-risk based on age, comorbidities, and poverty criteria.[7] There is also evidence that older[12] and obese[13] patients have higher medical costs, although other studies suggest that knee replacement is also cost-effective in nonagenarians[14] and draw conflicting conclusions on how benefits vary with body mass index (BMI).[15-17]

In the current economic environment there is great pressure to reduce NHS expenditure and numerous Primary Care Trusts (PCTs) have proposed cutting costs by limiting access to joint replacement[18-25] or classifying these procedures as being of limited value.[24-26] Since around 60,000 primary TKR procedures are conducted in England and Wales each year,[27] savings could be made with little loss of patient welfare *if* it were possible to reliably predict which patients obtain little/no benefit from TKR based on explicit, evidence-based criteria. Although the widely-used Oxford Knee Score (OKS) was developed to assess outcomes of knee replacement in trial populations,[28] several PCTs have set maximum OKS thresholds (ranging from 18 to 32), above which patients are ineligible for TKR.[18-24] However, we are not aware of any clinical or cost-effectiveness justification for these thresholds. Furthermore, many patients with higher OKS have significant symptoms or limitations to daily activities that could be improved by surgery. If rationing policies are to be adopted, cost-effectiveness analysis offers the only fair basis for designing them, since it ensures that NHS resources are allocated to maximise the health gained from available resources. As treatments are generally considered cost-effective if they cost less than £20,000-£30,000 per quality-adjusted life-year (QALY) gained,[11] restricting TKR to those patients for whom the procedure

1
2
3 costs less than £20,000-£30,000 per QALY gained would be a reasonable approach,
4 potentially realising savings that could be invested in other treatments giving greater
5 health gains. However, we are aware of no published evidence assessing how the
6 cost-effectiveness of TKR varies with OKS.
7
8
9

10 We therefore use data from the Knee Arthroplasty Trial (KAT)[29, 30] to assess the
11 appropriateness of rationing knee replacement by OKS, age, BMI, and clinical
12 characteristics, by calculating the cost-effectiveness of TKR in different patient
13 subgroups. In contrast to previous studies evaluating the cost-effectiveness of TKR
14 that relied upon data from small cohorts with short follow-up,[6-10, 14] the KAT trial
15 dataset provides detailed, prospective individual patient data on baseline
16 characteristics, treatments, complications, costs, and quality of life, and now has
17 more than five years' follow-up on 2352 patients.
18
19
20
21
22
23
24
25

26 **Methods**

27 We assessed the cost-effectiveness of primary TKR compared with no knee
28 replacement for patients with different baseline characteristics. Our analysis took a
29 National Health Service (NHS) perspective and excluded personal and social
30 services from the analysis, although TKR is likely to delay admission to residential
31 care,[14] reduce personal care costs and help recipients to continue paid
32 employment. A cost-utility analysis was conducted, which assessed cost-
33 effectiveness in terms of cost per quality-adjusted life-year (QALY) gained to capture
34 the quality of life gains associated with TKR.
35
36
37
38
39
40

41 Data on costs and quality of life following TKR were taken from the KAT trial
42 (ISRCTN 45837371): a pragmatic, partial-factorial, unblinded randomised controlled
43 trial (RCT) in which 2352 participants attending 34 UK centres were randomly
44 allocated to undergo TKR with/without a metal-backed tibial component, with/without
45 patellar resurfacing, and/or with/without a mobile bearing.[29, 30] Five-year trial
46 results demonstrated that patients experience substantial improvements in quality of
47 life and functional status following knee replacement that are maintained out to five
48 years[30]; further follow-up is ongoing. Our analysis took a five-year time horizon in
49 line with the results published to date[30]; costs and health benefits occurring more
50 than five years after primary TKR were conservatively excluded.
51
52
53
54
55
56
57
58
59
60

1
2
3 KAT participants completed the Oxford Knee Score (OKS) and EQ-5D
4 questionnaires immediately before and three months after TKR and annually
5 thereafter. The OKS is a validated 12-item questionnaire for assessing health status
6 and outcomes in relation to knee replacement that gives an unweighted total score
7 ranging from 0 (severe problems on all items) to 48 (no problems on any item).[28]
8 The EQ-5D is a generic health state preference measure[31]; health state preference
9 values (“utilities”) for EQ-5D profiles were based on time-trade-off valuations by
10 members of the UK general public.[31] EQ-5D utilities range from -0.594 (extreme
11 problems on all five domains) to 1 (perfect health) and indicate the value placed on
12 different levels of health. The number of QALYs that each patient accrued following
13 TKR was calculated as the area under the utility curve, with linear interpolation
14 between utility measurements; those patients who died were assumed to remain at
15 the last observed utility until death.
16
17
18
19
20
21
22
23

24 Healthcare resource use data (including knee-related outpatient, general practice
25 and physiotherapy consultations, and theatre time, hospital days, complications and
26 knee components used during the primary admission for TKR and during any knee-
27 related readmissions or revisions) were collected prospectively for all KAT
28 participants and valued at 2007-08 prices, as described previously.[30] Costs
29 incurred after Year 1 were discounted at 3.5% per year based on current UK
30 recommendations.[32]
31
32
33
34
35

36 The number of QALYs gained from TKR was calculated as the number of QALYs
37 observed minus the number of QALYs expected without TKR. Since all KAT
38 participants underwent TKR, assumptions were needed to estimate the costs and
39 QALYs that would have been accrued if no TKR procedures had occurred in the five-
40 year time horizon. We assumed that:
41
42
43

- 44 • Without TKR, patients would have remained at their baseline EQ-5D utility for five
45 years or until death (if sooner). In practice, patients’ symptoms and quality of life
46 are likely to deteriorate due to worsening arthritis[17, 33] and increasing age.[34]
- 47 • The seven KAT participants who died before hospital discharge would have
48 survived for five years at baseline utility without TKR but that TKR had no effect
49 on the date of death for other patients. This assumption is highly conservative
50 since those patients who die soon after joint replacement tend to be older[5, 35]
51 and have comorbidities that could have led to death without TKR.[5]
- 52 • Patients would not have used any healthcare resources due to their knee
53 problems if they had not undergone TKR. This assumption is highly conservative
54
55
56
57
58
59
60

1
2
3 since osteoarthritis patients will receive medical management without TKR and a
4 recent Canadian case-control study suggested that TKR recipients accrued lower
5 costs 6-18 months after the procedure than matched controls who had arthritis
6 but no TKR.[33]
7
8
9

10 Since these assumptions are likely to underestimate the benefits and overestimate
11 the additional costs of TKR, we evaluated the effect of relaxing these assumptions in
12 sensitivity analyses.
13
14

15 The 7% of data on baseline characteristics, resource use and quality of life that were
16 missing were imputed by multiple imputation using the ice command[36-38] (version
17 1.3.0) within Stata Version 11.0 (College Station, TX), thereby avoiding the bias and
18 inefficiency associated with complete case analysis.[38-40] Treatment indicators and
19 the covariates included in subsequent regression analyses were included in the
20 imputation model to avoid bias.[38] Fifty imputed datasets were generated and
21 results combined.
22
23
24
25
26
27
28

29 Regression models were used to estimate how the costs and QALYs gained from
30 TKR vary with baseline characteristics and to predict outcomes for different patient
31 groups without subdividing the trial population into small patient subgroups. Since
32 costs and QALYs were highly skewed, generalised linear models (GLM) with gamma
33 family distributions were used to predict total costs and the QALYs accrued with or
34 without TKR; prior to regression, the number of QALYs accrued was subtracted from
35 the maximum number of QALYs that could have been accrued over the five-year trial
36 period to ensure that all QALY data were positive. We evaluated the effect of six
37 baseline characteristics on costs and QALYs with/without TKR:
38
39
40
41
42

- 43 • Male sex;
- 44 • Age at time of operation;
- 45 • Baseline OKS (using the new, 0-48, scoring system[41]);
- 46 • American Society of Anesthesiologists (ASA) grade 3 (symptomatic illness
47 with minimal restriction on daily life) rather than grades 1 (fit & healthy) or 2
48 (asymptomatic illness with no restriction on daily life);
- 49 • BMI (kg/m²);
- 50 • Presence of arthritis in one knee rather than in both knees or generally.
51
52
53
54
55
56
57
58
59
60

1
2
3 Since the introduction of biologic therapies has changed management of rheumatoid
4 arthritis and as rheumatoid arthritis patients and those with severe comorbidities are
5 likely have very different costs and outcomes from most patients, 108 rheumatoid
6 arthritis patients and 13 patients with ASA grade 4 were excluded from the analysis.
7
8 One hundred 100 patients who were randomised to total vs unicompartmental knee
9 replacement or died or withdrew from the trial before surgery were also excluded
10 from the analysis, giving a final sample size of 2131.
11
12

13
14
15 All regression analyses were conducted in Stata Version 11. Bootstrapping was used
16 to capture uncertainty around cost-effectiveness ratios and correlations between
17 costs and QALYs with/without TKR; the regression models for costs and outcomes
18 were run for each of 125 bootstrap samples drawn from each imputed KAT dataset
19 and results combined using Rubin's rule.[39] Predicted costs and the predicted
20 number of QALYs accrued with and without TKR in different patient groups were
21 calculated from regression coefficients and used to estimate incremental cost-
22 effectiveness ratios (ICERs). Bootstrap results were used to calculate cost-
23 effectiveness acceptability curves (CEACs),[42] which plot the probability of
24 treatment being cost-effective against the ceiling ratio (i.e. the maximum that society
25 is willing or able to pay per QALY gained).
26
27
28
29
30
31
32

33 We also calculated the number of patients in England and Wales eligible for TKR
34 under different rationing policies based on 61,651 patients undergoing primary TKR
35 each year.[27] The proportion of patients eligible for TKR under different rationing
36 criteria was based on analysis of an extract of patient-level Patient Reported
37 Outcome Measures[1] (PROMs) data on admissions for knee replacement up to 31st
38 December 2010, supplemented where necessary by the baseline characteristics from
39 the KAT sample.
40
41
42
43
44

45 46 **Results**

47 Of the 2131 KAT participants who had osteoarthritis, were ASA grades 1-3 and
48 underwent TKR as part of the trial, the mean baseline OKS was 18 (standard
49 deviation, SD: 7.5), mean age 71 (SD: 8.0), mean BMI 30 (SD: 5.5), and 45%
50 (956/2131) were male (Figure 1); further baseline characteristics of the trial
51 population have been reported previously.[29]
52
53
54
55
56
57
58
59
60

1
2
3 Mean EQ-5D utility rose substantially within three months of TKR from 0.39 at
4 baseline to 0.71 at one year, and declined gradually thereafter (Figure 1, Table 1).
5
6 However, baseline utility and the quality of life gains associated with TKR varied
7
8 between patient subgroups. In particular, women and patients with low OKS (i.e. poor
9
10 knee function), higher ASA grade or obesity at baseline tended to have lower quality
11
12 of life at all timepoints (Figure 1), although those in the lowest OKS quintile
13
14 experienced significantly greater health gains from TKR than the top quintile
15
16 ($p < 0.001$; calculated using linear regression, combining imputed datasets using the
17
18 mim command).

18
19 On average, each admission for primary TKR cost £6,363 (SD: £1,702).
20
21 Readmissions, revision procedures and GP, outpatient, and physiotherapy
22
23 consultations related to the study knee over the following five years cost a further
24
25 £1,095 (SD: £3,579) per patient, giving a total cost of £7,458 (SD: £4,058) per
26
27 patient. TKR was more costly in patients with low baseline OKS, but the greater
28
29 QALY gains meant that TKR was better value for money in this patient group than in
30
31 patients with moderate symptoms (Table 1). Subgroup analyses suggested that TKR
32
33 cost less than £11,000/QALY gained across all OKS deciles: well below the £20,000-
34
35 £30,000/QALY gained range that NICE currently consider good value for money.[11]

36
37 However, cost-effectiveness also varied substantially between patients within each
38
39 OKS decile. We therefore used regression models to estimate the costs, QALY gains
40
41 and cost-effectiveness of TKR in a wide range of patient subgroups while controlling
42
43 for multiple baseline characteristics. Our results indicated that four factors have a
44
45 significant effect on cost-effectiveness: age, sex, baseline OKS and ASA grade
46
47 (Table 2). BMI and the presence of arthritis in other joints besides the study knee had
48
49 no significant effect on either the costs or QALY gains associated with TKR ($p > 0.05$).
50
51
52
53
54
55
56
57
58
59
60

Table 1: EQ-5D utility, costs and estimated QALYs for subgroups with different baseline OKS

OKS decile		1 (worst)	2	3	4	5	6	7	8	9	10 (best)	All patients
Baseline OKS range		<9	9-11	12-13	14-15	16-17	18-19	20-21	22-24	25-27	>27	0-48
No. KAT participants		204	213	186	232	193	224	195	256	194	234	2131
EQ-5D utility (SE)†	Baseline	0.03 (0.004)	0.14 (0.005)	0.21 (0.006)	0.25 (0.006)	0.36 (0.006)	0.46 (0.006)	0.55 (0.005)	0.57 (0.004)	0.61 (0.004)	0.67 (0.003)	0.39 (0.002)
	3 mths	0.54 (0.007)	0.58 (0.006)	0.64 (0.006)	0.66 (0.005)	0.71 (0.005)	0.71 (0.005)	0.70 (0.004)	0.74 (0.004)	0.75 (0.005)	0.77 (0.004)	0.68 (0.002)
	1 year	0.51 (0.007)	0.60 (0.007)	0.68 (0.006)	0.70 (0.006)	0.71 (0.006)	0.75 (0.005)	0.69 (0.006)	0.78 (0.004)	0.79 (0.005)	0.84 (0.004)	0.71 (0.002)
	2 year	0.49 (0.008)	0.54 (0.007)	0.66 (0.007)	0.67 (0.006)	0.68 (0.007)	0.73 (0.005)	0.66 (0.007)	0.76 (0.005)	0.78 (0.005)	0.81 (0.005)	0.68 (0.002)
	3 year	0.47 (0.008)	0.53 (0.007)	0.64 (0.007)	0.63 (0.006)	0.67 (0.007)	0.71 (0.006)	0.68 (0.007)	0.72 (0.005)	0.76 (0.006)	0.79 (0.006)	0.66 (0.002)
	4 year	0.43 (0.008)	0.49 (0.008)	0.61 (0.007)	0.62 (0.007)	0.65 (0.007)	0.68 (0.007)	0.64 (0.007)	0.70 (0.006)	0.71 (0.007)	0.75 (0.006)	0.63 (0.002)
	5 year	0.43 (0.008)	0.50 (0.008)	0.57 (0.008)	0.58 (0.007)	0.63 (0.007)	0.67 (0.007)	0.59 (0.007)	0.68 (0.006)	0.67 (0.008)	0.73 (0.007)	0.61 (0.002)
Total cost (SE)*		£8657 (£157)	£7715 (£81)	£7495 (£73)	£7081 (£63)	£7185 (£69)	£7567 (£82)	£7619 (£102)	£7128 (£61)	£7399 (£94)	£6917 (£67)	£7458 (£28)
QALYs with TKR (SE)*		2.17 (0.029)	2.48 (0.027)	2.94 (0.024)	2.99 (0.022)	3.13 (0.025)	3.30 (0.022)	3.09 (0.025)	3.40 (0.019)	3.50 (0.020)	3.67 (0.020)	3.08 (0.025)
QALYs without TKR (SE)*		0.15 (0.02)	0.62 (0.022)	0.94 (0.026)	1.14 (0.026)	1.63 (0.029)	2.04 (0.026)	2.39 (0.026)	2.55 (0.021)	2.78 (0.02)	3.02 (0.02)	1.75 (0.031)
QALY gain from TKR (SE)*		2.02 (0.032)	1.86 (0.031)	2.00 (0.033)	1.85 (0.031)	1.51 (0.034)	1.26 (0.030)	0.69 (0.027)	0.85 (0.023)	0.73 (0.02)	0.65 (0.02)	1.33 (0.01)
ICER*		£4,295	£4,153	£3,747	£3,836	£4,770	£6,007	£10,971	£8,391	£10,167	£10,697	£5,623

* Discounted at 3.5% per annum

† EQ-5D utility is measured on a scale from 1 (perfect health) to -0.594, where 0 is equivalent to death.

Abbreviations: ICER, incremental cost-effectiveness ratio; OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])[41]; QALY, quality-adjusted life-year; SE, standard error around the mean; TKR, total knee replacement.

Table 2: Results of the regression analysis

Baseline characteristic	Coefficient (SE)		
	QALY loss with TKR†	QALY loss without TKR†	Total costs with TKR (£)
Male sex	0.056 (0.065)	-0.001 (0.054)	593 (181)*
Age at operation (years)	0.007 (0.003)*	0.002 (0.004)	-8 (12)
Pre-operative OKS	-0.040 (0.004)*	-0.087 (0.004)*	-52 (12)*
ASA grade 3	0.414 (0.067)*	0.227 (0.087)*	492 (208)*
Constant	1.712 (0.201)*	4.274 (0.325)*	8573 (881)*

* p<0.05

† QALY loss indicates the number of discounted QALYs that would have been accrued if patients had experienced an EQ-5D utility of 1 for five years (4.67 QALYs) minus the number of discounted QALYs that the patient actually experienced.

The incremental cost-effectiveness ratio (expressed as cost/QALY) for any given patient equals: (male x 593 - age x 8 - OKS x 52 + ASAgrade3 x 492 + 8573)/((male x -0.001 + age x 0.002 - OKS x 0.087 + ASAgrade3 x 0.227 + 4.274) - (male x 0.056 + age x 0.007 - OKS x 0.040 + ASAgrade3 x 0.414 + 1.712). Costs and QALYs were discounted at 3.5% per annum.

Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy, 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life); OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])[41]; QALY, quality-adjusted life-year; TKR, total knee replacement.

Figure 2 shows the cost-effectiveness of TKR for each multivariate group, using charts similar in design to widely-used cardiovascular risk tables.[43] The charts demonstrate that TKR is clearly cost-effective across a wide range of patient groups. For example, TKR costs less than £20,000/QALY gained for 71-year-old women with no symptomatic illness unless baseline OKS is above 40 (equivalent to having very minor problems on up to eight of the 12 activities mentioned in OKS or moderate-severe pain only on exertion). The threshold OKS at which TKR becomes cost-ineffective was lower for older patients, men, and those with symptomatic disease, but TKR remained cost-effective for all patients with OKS below 32 regardless of age, sex, or ASA grade. A simplified analysis controlling only for OKS and ASA grade suggested that TKR is cost-effective for all ASA grade 1 or 2 patients with baseline OKS <40, and all ASA grade 3 patients with OKS <35.

There was substantial uncertainty around the cost-effectiveness ratios for patients with very high baseline OKS, since only 37 trial participants had OKS above 35 at baseline. However, we estimate that there is a 99% probability that TKR is cost-effective (at a £20,000/QALY ceiling ratio) for the average ASA grade 1-2 patient with a baseline OKS of 35, and a 53% probability for ASA grade 3 (Figure 3).

Since each additional baseline characteristic increases the complexity of the eligibility criteria and may also raise ethical or equity concerns, we estimated the number of patients who would receive TKR under different eligibility criteria. This analysis suggested that restricting TKR to those patients for whom the operation was

1
2
3 predicted to cost less than £20,000/QALY gained based on the simplified model
4 (allowing only for ASA grade and OKS) would lead to 2000 fewer TKR procedures
5 being conducted in England and Wales each year (3% fewer than the 61,651
6 procedures conducted in 2009/10[27]). If the decision rules were simplified further to
7 provide TKR to all patients with baseline OKS <39, all but 1700 of the 61,651 patients
8 currently receiving TKR in England and Wales in 2009/10[27] would be eligible.
9 Allowing for other baseline characteristics had minimal effect on patient numbers or
10 predicted costs or benefits, suggesting that decisions about suitability for TKR can be
11 based on OKS alone.
12

13
14
15
16
17
18 Sensitivity analyses were conducted to assess the effect of extending the time
19 horizon of our analysis up to 10 years, and of assuming that EQ-5D utility would have
20 fallen by 0.5% per year without TKR,[34] that all knee-related ambulatory
21 consultations would have occurred without TKR, and that the mean length of stay
22 has fallen by 41% (5.9 days[27] vs 10.0) since the KAT operations took place in
23 1999-2003. Changing any of these assumptions produced lower ICERs than the
24 base case analysis, confirming that the current analysis is highly conservative and
25 suggesting that TKR may be cost-effective for ASA grade 1-2 patients with baseline
26 OKS as high as 43. Excluding patients with incomplete data had no effect on
27 conclusions. Although statistically significant, adding an age-squared term to the
28 regression model had no effect on the conclusions, while a sensitivity analysis adding
29 an OKS-squared term suggested that TKR may be cost-effective for all patients of
30 ASA grade 1 or 2. Recoding BMI as a dummy variable indicating whether or not
31 patients were obese (BMI ≥30) did not change the conclusions and obesity had no
32 significant effect on costs or QALYs.
33
34
35
36
37
38
39
40
41
42
43
44

45 Discussion

46 We found TKR to cost £5,623 per QALY gained for the average patient (similar to
47 estimates from previous studies[6-10]) and therefore represent very good value for
48 money compared with the £20,000-£30,000/QALY threshold typically used in NHS
49 decision-making.[11] Our study also confirms previous findings[7, 9, 17] that costs
50 and benefits vary with age and comorbidity, but evaluates such characteristics, and
51 pre-operative knee function, in multivariate analyses, which has not (to our
52 knowledge) been done previously. We found pre-operative OKS to be the best
53 predictor of post-operative costs, outcomes and cost-effectiveness.
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Our analyses were based on patient-level data from a large trial with detailed, prospective collection of costs, utilities and baseline characteristics that permit extensive subgroup analyses. KAT used a pragmatic design, with broad inclusion criteria and no restrictions on peri- or post-operative care other than randomised aspects of component design[29, 30]; as result, the trial cohort is similar to the national PROMs cohort and costs and benefits are likely to be comparable to those in routine clinical practice. Furthermore, unlike some previous economic evaluations, our study took full account of uncertainty and included all knee-related NHS costs accrued in the first five years after TKR. The main limitation of our analysis was the small number of patients with high OKS. Since KAT recruited only 37 patients with OKS >35, further studies are required to evaluate TKR in this patient subgroup. Analyses were also based on *post hoc* comparisons between outcomes before and after TKR, rather than between randomly-assigned treatment groups. As a result, our study does not provide unbiased estimates of the QALY gains from TKR, and regression towards the mean may have caused our study to overestimate the quality of life improvement attributable to TKR and/or the effect of pre-operative OKS on QALY gains.

We made several highly conservative assumptions: in particular, assuming that patients would have accrued no knee-related costs and would have remained at baseline utility without TKR, whereas previous studies have found that patients accrue substantial costs and show clinical worsening without TKR.[33] We also assumed that those patients who died before hospital discharge would have survived for five years if they had not had TKR; in reality, those patients dying soon after TKR tend to be older[5, 35] and frailer[5] pre-operatively and some studies suggest that patients undergoing hip replacement have higher long-term survival than the general population[44, 45] (although this may be due to patient selection[35, 44]). When these assumptions were relaxed, TKR was cost-effective for ASA grade 1-2 patients with baseline OKS <44. The costs and benefits of TKR are likely to vary between centres and over time due to variations in patient demographics, component prices, length of stay, and operation time; sensitivity analyses demonstrated that TKR may have become more cost-effective since the KAT procedures were conducted in 1999-2003 due to reductions in length of stay.

Our analyses show that if TKR were to be rationed based on cost-effectiveness, OKS would be a reasonable tool to use to set the threshold, although there may, of

1
2
3 course, be practical difficulties with using a patient-reported measure to determine
4 treatment eligibility: particularly if patients know how their questionnaire responses
5 will be used or if test-retest reliability is low in this setting. The decision to operate
6 must also take into consideration other factors, such as radiographic findings and
7 patient choice. Based on regression and subgroup analyses, we can be confident
8 that TKR costs less than £20,000/QALY for all ASA grade 1-2 patients with baseline
9 OKS <40 and for ASA grade 3 patients with OKS <35. Using these thresholds to
10 determine who received treatment would avoid around 2000 TKR procedures per
11 year in England and Wales, saving around £11.8 million per year. However, the
12 higher thresholds suggested by sensitivity analyses using less conservative
13 assumptions suggest that TKR is cost-effective for all but 100 patients per year,
14 which is unlikely to achieve sufficient savings to warrant rationing of TKR. Although
15 the cost-effectiveness of knee replacement was also influenced by gender and age,
16 taking account of these factors as well as OKS and ASA grade would increase the
17 complexity of eligibility criteria, reduce equality of access to healthcare and have
18 minimal effect on NHS budgets.
19
20
21
22
23
24
25
26
27
28

29 The results also demonstrate that the thresholds proposed by some PCTs[18-24] are
30 inappropriate and would deny a highly cost-effective treatment to thousands of
31 patients with severe arthritis. The current situation also introduces a postcode lottery
32 with different PCTs using different eligibility criteria. We estimate that limiting knee
33 replacement to patients with OKS ≤ 26 would prevent more than 10,000 people from
34 receiving TKR. This would be inequitable as TKR costs £10,697 per QALY gained for
35 this group, whereas NICE routinely recommend treatments with ICERs in the region
36 of £20,000-£30,000/QALY gained[11, 46, 47] and are willing to pay substantially
37 more for end of life care.[48] Furthermore, the finding that TKR and subsequent
38 monitoring is more costly for patients with low OKS suggests that delaying TKR until
39 symptoms have deteriorated may be a false economy. Although some PCTs have
40 considered restricting TKR further for obese patients,[18, 21, 25, 49] we found BMI to
41 have no significant effect on costs ($p=0.442$), QALY gains ($p=0.098$) or the incidence
42 of peri- or post-operative complications ($p>0.26$). Although BMI remained non-
43 significant even when ASA grade was omitted from regression analyses, it is difficult
44 to isolate the effect of obesity from other correlated variables (e.g. comorbidity); as
45 result, further research on obesity may be warranted: particularly since a previous
46 study observed a significant correlation between BMI and costs.[13]
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Eligibility criteria for rheumatoid arthritis patients and patients with ASA grade 4
4 remain unclear, although ASA grade 4 patients would normally be considered unfit
5 for surgery, and the decision to operate on such patients would be based on a case-
6 by-case consideration of whether the benefits outweigh the risks. While our study
7 focused on the costs and benefits of TKR, patients with high baseline OKS often
8 undergo unicompartmental knee replacement; further research is needed to evaluate
9 how the cost-effectiveness of unicompartmental replacement varies with OKS.
10
11
12
13

14
15 In conclusion, TKR is highly cost-effective for the vast majority of patients who
16 currently undergo this procedure in the UK. Although costs and health benefits vary
17 with baseline OKS, the rationing thresholds proposed by some PCTs[18-24] are not
18 supported by evidence on health outcomes or cost-effectiveness.
19
20
21
22

23 **Acknowledgements**

24 The authors would like to thank all those involved in the KAT trial for their role in
25 running the trial and collecting the data used in this study, particularly other members
26 of the KAT Project Management Group: Suzanne Breeman; Marion Campbell; Jackie
27 Ellington; Nick Fiddian; Adrian Grant; Linda Johnston; and Richard Morris.
28
29
30
31

32 The Knee Arthroplasty Trial is funded by the NIHR Health Technology Assessment
33 Programme (project number 95/10/01) and will be published in full as a Health
34 Technology Assessment report. Visit the HTA programme website for further project
35 information. Additional funding for research support in clinical centres was provided
36 by Howmedica Osteonics; Zimmer; DePuy, a Johnson and Johnson company; Corin
37 Medical; Smith and Nephew Healthcare; Biomet Merck; and Wright Cremascoli. The
38 Health Economics Research Centre obtains support from the National Institute of
39 Health Research. The Health Services Research Unit is core funded by the Chief
40 Scientist Office of the Scottish Government Health Directorates. PROMs data were
41 reused with the permission of The Health and Social Care Information Centre,
42 copyright 2011, with all rights reserved. The views and opinions expressed therein
43 are those of the authors and do not necessarily reflect those of the HTA programme,
44 NIHR, NHS, Information Centre or the Department of Health.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Competing interests

All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare no support from any organisation for the submitted work with the exception of those declared in the funding statement. AG, RF and GM declare that they have no financial relationships with any organisations that might have an interest in the submitted work in the previous three years and no other relationships or activities that could appear to have influenced the submitted work. DM receives royalties paid to him by Biomet and HD has undertaken consultancy work for Pfizer on drugs for rheumatoid arthritis but both declare that they have no other financial relationships with any organisations that might have an interest in the submitted work in the previous three years and no other relationships or activities that could appear to have influenced the submitted work.

Funding

The Knee Arthroplasty Trial is funded by the NIHR Health Technology Assessment Programme (project number 95/10/01). Additional funding for research support in clinical centres was provided by Howmedica Osteonics; Zimmer; DePuy, a Johnson and Johnson company; Corin Medical; Smith and Nephew Healthcare; Biomet Merck; and Wright Cremascoli.

The Health Economics Research Centre obtains support from the National Institute of Health Research. The Health Services Research Unit is core funded by the Chief Scientist Office of the Scottish Government Health Directorates. The Biomedical Research Unit at Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences is funded by the National Institute of Health Research.

The authors conducted the research independently of all funding organisations. The funders had no role in the collection, analysis or interpretation of data, writing of the manuscript or the decision to publish. The authors had full access to all of the data (including statistical reports and tables) in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Contributorship statement

HD, AG and DM conceived and developed the principles and methods underpinning the current analysis. GM was involved in primary data collection and database management, while HD conducted the analysis under AG's supervision and guidance from DM, GM and RF. HD drafted the manuscript with input from all other authors; all authors have approved the final manuscript and were involved in the interpretation of results. The KAT trial was conceived, designed and run by the KAT trial group, which comprised Suzanne Breeman, Marion Campbell, HD, Jackie Ellington, Nick Fiddian, RF, Adrian Grant, AG, Linda Johnston, GM, Richard Morris and DM.

References

- 1 Hospital Episode Statistics. Provisional Monthly Patient Reported Outcome Measures (PROMs) in England. April 2009 – April 2010: Pre- and postoperative data: Experimental statistics. 2010 [updated 2010; cited 6th April 2011]; Available from: <http://www.hqip.org.uk/assets/NCAPOP-Library/PROMsreport09101.pdf>.
- 2 Labek G, Thaler M, Janda W, et al. Revision rates after total joint replacement: cumulative results from worldwide joint register datasets. *J Bone Joint Surg Br* 2011;**93**:293-7. doi: 93-B/3/293 [pii] 10.1302/0301-620X.93B3.25467
- 3 The National Collaborating Centre for Chronic Conditions. Osteoarthritis: National clinical guideline for care and management in adults. London: Royal College of Physicians; 2008 [updated 2008; cited 29th June 2011]; Available from: <http://www.nice.org.uk/nicemedia/live/11926/39720/39720.pdf>.
- 4 Huddleston JI, Maloney WJ, Wang Y, et al. Adverse events after total knee arthroplasty: a national Medicare study. *J Arthroplasty* 2009;**24**:95-100. doi: S0883-5403(09)00168-5 [pii] 10.1016/j.arth.2009.05.001
- 5 Memtsoudis SG, Della Valle AG, Besculides MC, et al. Risk factors for perioperative mortality after lower extremity arthroplasty: a population-based study of 6,901,324 patient discharges. *J Arthroplasty* 2010;**25**:19-26. doi: S0883-5403(08)00855-3 [pii] 10.1016/j.arth.2008.11.010
- 6 Navarro Espigares JL, Hernandez Torres E. Cost-outcome analysis of joint replacement: evidence from a Spanish public hospital. *Gac Sanit* 2008;**22**:337-43. doi: 13125355 [pii]
- 7 Losina E, Walensky RP, Kessler CL, et al. Cost-effectiveness of total knee arthroplasty in the United States: patient risk and hospital volume. *Arch Intern Med* 2009;**169**:1113-21; discussion 21-2. doi: 169/12/1113 [pii] 10.1001/archinternmed.2009.136

- 1
2
3 8 Rasanen P, Paavolainen P, Sintonen H, et al. Effectiveness of hip or knee
4 replacement surgery in terms of quality-adjusted life years and costs. *Acta Orthop*
5 2007;**78**:108-15. doi: 776344017 [pii] 10.1080/17453670610013501
6
7 9 Rissanen P, Aro S, Sintonen H, et al. Costs and cost-effectiveness in hip and knee
8 replacements. A prospective study. *Int J Technol Assess Health Care* 1997;**13**:575-
9 88.
10
11 10 Lavernia CJ, Guzman JF, Gachupin-Garcia A. Cost effectiveness and quality of
12 life in knee arthroplasty. *Clin Orthop Relat Res* 1997:134-9.
13
14 11 National Institute for Health and Clinical Excellence. Social value judgements:
15 Principles for the development of NICE guidance. Second Edition. 2008 [updated
16 2008; cited 17th November 2010]; Available from:
17 <http://www.nice.org.uk/media/C18/30/SVJ2PUBLICATION2008.pdf>.
18
19 12 Brander VA, Malhotra S, Jet J, et al. Outcome of hip and knee arthroplasty in
20 persons aged 80 years and older. *Clin Orthop Relat Res* 1997:67-78.
21
22 13 Dowsey MM, Liew D, Choong PF. Economic burden of obesity in primary total
23 knee arthroplasty. *Arthritis Care Res (Hoboken)* 2011;**63**:1375-81. doi:
24 10.1002/acr.20563
25
26 14 Karuppiah SV, Banaszkiwicz PA, Ledingham WM. The mortality, morbidity and
27 cost benefits of elective total knee arthroplasty in the nonagenarian population. *Int*
28 *Orthop* 2008;**32**:339-43. doi: 10.1007/s00264-007-0324-y
29
30 15 McQueen DA, Long MJ, Algotar AM, et al. The effect of obesity on quality-of-life
31 improvement after total knee arthroplasty. *Am J Orthop (Belle Mead NJ)*
32 2007;**36**:E117-20, E27.
33
34 16 de Leeuw JM, Villar RN. Obesity and quality of life after primary total knee
35 replacement *Knee* 1998;**5**:119-23.
36
37 17 Cushnaghan J, Bennett J, Reading I, et al. Long-term outcome following total
38 knee arthroplasty: a controlled longitudinal study. *Ann Rheum Dis* 2009;**68**:642-7.
39 doi: ard.2008.093229 [pii] 10.1136/ard.2008.093229
40
41 18 NHS Oxfordshire. INTERIM Treatment Threshold Statement: Knee arthroplasty
42 (Reference PS 188). 2010 [updated 2010; cited 5th April 2011]; Available from:
43 [http://www.oxfordshirepct.nhs.uk/professional-resources/priority-setting/lavender-](http://www.oxfordshirepct.nhs.uk/professional-resources/priority-setting/lavender-statements/documents/PS188Kneereplacement.pdf)
44 [statements/documents/PS188Kneereplacement.pdf](http://www.oxfordshirepct.nhs.uk/professional-resources/priority-setting/lavender-statements/documents/PS188Kneereplacement.pdf).
45
46 19 NHS Gloucestershire. Guidelines for Management of Orthopaedic Conditions.
47 2010 [updated 2010; cited 5th April 2011]; Available from:
48 [http://www.glospct.nhs.uk/pdf/services/orthopaedic_referral_guidelines_%20march%](http://www.glospct.nhs.uk/pdf/services/orthopaedic_referral_guidelines_%20march%2010v3.pdf)
49 [2010v3.pdf](http://www.glospct.nhs.uk/pdf/services/orthopaedic_referral_guidelines_%20march%2010v3.pdf).
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 20 NHS Worcestershire. Commissioning policy for musculoskeletal surgical
4 interventions. 2010 [updated 2010; cited 5th April 2011]; Available from:
5
6 [http://www.worcestershire.nhs.uk/file_download.aspx?id=86bae3e2-4d88-4416-](http://www.worcestershire.nhs.uk/file_download.aspx?id=86bae3e2-4d88-4416-92e6-9eeda22e8208)
7
8 [92e6-9eeda22e8208](http://www.worcestershire.nhs.uk/file_download.aspx?id=86bae3e2-4d88-4416-92e6-9eeda22e8208).
- 9 21 Milton Keynes O, Berkshire East, Berkshire West & Buckinghamshire (MOBBB)
10 Priorities Committee,. Minutes of meeting held on Wednesday 15 December 2010.
11 2010 [updated 2010; cited 5th April 2011]; Available from: [http://www.sph.nhs.uk/sph-](http://www.sph.nhs.uk/sph-psu/sph-files/mobbb-minutes/mobbb-minutes-15-december-2010)
12
13 [psu/sph-files/mobbb-minutes/mobbb-minutes-15-december-2010](http://www.sph.nhs.uk/sph-psu/sph-files/mobbb-minutes/mobbb-minutes-15-december-2010).
- 14 22 South West London Public Health Network. 2010/11 South West London Effective
15 Commissioning Initiative. 2010 [updated 2010; cited 5th April 2011]; Available from:
16
17 [http://www.swlphnetwork.org.uk/ uploads/documents/effective-commissioning-](http://www.swlphnetwork.org.uk/uploads/documents/effective-commissioning-document-master-version-2010-11-final.doc)
18
19 [document-master-version-2010-11-final.doc](http://www.swlphnetwork.org.uk/uploads/documents/effective-commissioning-document-master-version-2010-11-final.doc).
- 20 23 NHS Manchester. Topics for inclusion or change in the pct non-commissioned
21 procedures list. 2010 [updated 2010; cited 5th April 2011]; Available from:
22
23 [http://www.manchester.nhs.uk/document_uploads/19%20January%202011/C2turnar](http://www.manchester.nhs.uk/document_uploads/19%20January%202011/C2turnaroundAppendix%20%20TR.doc)
24
25 [oundAppendix%20%20TR.doc](http://www.manchester.nhs.uk/document_uploads/19%20January%202011/C2turnaroundAppendix%20%20TR.doc).
- 26 24 NHS Derbyshire County. NHS Derbyshire County Prior Approval: Procedures of
27 Limited Clinical Value Summary. 2010 [updated 2010; cited 29th June 2011];
28 Available from:
29
30 [http://www.phcn.nhs.uk/Derbyshire%20County%20Procedures%20of%20Limited%2](http://www.phcn.nhs.uk/Derbyshire%20County%20Procedures%20of%20Limited%20Clinical%20Value%20100525%20draft.pdf)
31
32 [0Clinical%20Value%20100525%20draft.pdf](http://www.phcn.nhs.uk/Derbyshire%20County%20Procedures%20of%20Limited%20Clinical%20Value%20100525%20draft.pdf).
- 33 25 Jeffreys B. Surgeons raise alarm over waiting. Journal [serial on the Internet].
34 2011 Date: Available from: <http://www.bbc.co.uk/news/health-12964360>.
- 35 26 NHS Warwickshire. Report to the PCT board 10th November 2010. 2010 [updated
36 2010 6th April 2011; cited 29th June 2011]; Available from:
37
38 [http://www.warwickshire.nhs.uk/trustboardenclosures/November2010/Enc%208%20](http://www.warwickshire.nhs.uk/trustboardenclosures/November2010/Enc%208%20%20In%20Year%20Demand%20Management%20of%20Electives%205-11-2010.pdf)
39
40 [%20In%20Year%20Demand%20Management%20of%20Electives%205-11-](http://www.warwickshire.nhs.uk/trustboardenclosures/November2010/Enc%208%20%20In%20Year%20Demand%20Management%20of%20Electives%205-11-2010.pdf)
41
42 [2010.pdf](http://www.warwickshire.nhs.uk/trustboardenclosures/November2010/Enc%208%20%20In%20Year%20Demand%20Management%20of%20Electives%205-11-2010.pdf).
- 43 27 Hospital Episode Statistics. Main procedures and interventions: 4 character
44 inpatient data 2009-10. 2010 [updated 2010; cited 10th February 2011]; Available
45 from:
46
47 [http://www.hesonline.nhs.uk/Ease/servlet/ContentServer?siteID=1937&categoryID=2](http://www.hesonline.nhs.uk/Ease/servlet/ContentServer?siteID=1937&categoryID=215)
48
49 [15](http://www.hesonline.nhs.uk/Ease/servlet/ContentServer?siteID=1937&categoryID=215).
- 50 28 Dawson J, Fitzpatrick R, Murray D, et al. Questionnaire on the perceptions of
51 patients about total knee replacement. J Bone Joint Surg Br 1998;**80**:63-9. doi:
52
53 29 Johnston L, MacLennan G, McCormack K, et al. The Knee Arthroplasty Trial
54 (KAT) design features, baseline characteristics, and two-year functional outcomes
55
56
57
58
59
60

- 1
2
3 after alternative approaches to knee replacement. *J Bone Joint Surg Am*
4 2009;**91**:134-41. doi: 91/1/134 [pii] 10.2106/JBJS.G.01074
5
6 30 Breeman S, Campbell M, Dakin H, et al. Patellar Resurfacing in Total Knee
7 Replacement: Five-Year Clinical and Economic Results of a Large Randomized
8 Controlled Trial. *J Bone Joint Surg Am* 2011;**93**:1473-81.
9
10 31 Dolan P. Modeling valuations for EuroQol health states. *Med Care* 1997;**35**:1095-
11 108.
12
13 32 HM Treasury. *The Green Book: Appraisal and Evaluation in Central Government*.
14 London: TSO; 2003.
15
16 33 Hawker GA, Badley EM, Croxford R, et al. A population-based nested case-
17 control study of the costs of hip and knee replacement surgery. *Med Care*
18 2009;**47**:732-41. doi: 10.1097/MLR.0b013e3181934553
19
20 34 Kind P, Dolan P, Gudex C, et al. Variations in population health status: results
21 from a United Kingdom national questionnaire survey. *Bmj* 1998;**316**:736-41. doi:
22
23 35 Lie SA, Pratt N, Ryan P, et al. Duration of the increase in early postoperative
24 mortality after elective hip and knee replacement. *J Bone Joint Surg Am* 2010;**92**:58-
25 63. doi: 92/1/58 [pii] 10.2106/JBJS.H.01882
26
27 36 Royston P. Multiple imputation of missing values. *Stata Journal* 2004;**4**:227-41.
28
29 37 Royston P. Multiple imputation of missing values: further update of ice, with an
30 emphasis on interval censoring. *Stata Journal* 2007;**7**:445-64.
31
32 38 White IR, Royston P, Wood AM. Multiple imputation using chained equations:
33 Issues and guidance for practice. *Stat Med* 2011;**30**:377-99. doi: 10.1002/sim.4067
34
35 39 Briggs A, Clark T, Wolstenholme J, et al. Missing... presumed at random: cost-
36 analysis of incomplete data. *Health Econ* 2003;**12**:377-92.
37
38 40 Yu LM, Burton A, Rivero-Arias O. Evaluation of software for multiple imputation of
39 semi-continuous data. *Stat Methods Med Res* 2007;**16**:243-58. doi: 16/3/243 [pii]
40 10.1177/0962280206074464
41
42 41 Murray DW, Fitzpatrick R, Rogers K, et al. The use of the Oxford hip and knee
43 scores. *J Bone Joint Surg Br* 2007;**89**:1010-4. doi: 89-B/8/1010 [pii]
44 10.1302/0301-620X.89B8.19424
45
46 42 Fenwick E, Claxton K, Sculpher M. Representing uncertainty: the role of cost-
47 effectiveness acceptability curves. *Health Econ* 2001;**10**:779-87.
48
49 43 British Medical Association. *British National Formulary 61: Cardiovascular risk*
50 prediction charts for primary prevention. March 2011 [updated March 2011; cited
51 12th April 2011]; Available from: <http://bnf.org/bnf/bnf/current/204016.htm>.
52
53 44 Barrett J, Losina E, Baron JA, et al. Survival following total hip replacement. *J*
54 *Bone Joint Surg Am* 2005;**87**:1965-71. doi: 87/9/1965 [pii] 10.2106/JBJS.D.02440
55
56
57
58
59
60

1
2
3 45 Lie SA, Engesaeter LB, Havelin LI, et al. Mortality after total hip replacement: 0-
4 10-year follow-up of 39,543 patients in the Norwegian Arthroplasty Register. Acta
5 Orthop Scand 2000;**71**:19-27. doi: 10.1080/00016470052943838

6
7 46 Dakin HA, Devlin NJ, Odeyemi IA. "Yes", "No" or "Yes, but"? Multinomial
8 modelling of NICE decision-making. Health Policy 2006;**77**:352-67.

9
10 47 Devlin N, Parkin D. Does NICE have a cost-effectiveness threshold and what
11 other factors influence its decisions? A binary choice analysis. Health Econ
12 2004;**13**:437-52.

13
14 48 National Institute for Health and Clinical Excellence. Supplementary guidance on
15 appraising life-extending, end of life treatments. July 2009 [updated July 2009; cited
16 23rd February 2011]; Available from:

17
18 <http://www.nice.org.uk/media/E4A/79/SupplementaryAdviceTACEoL.pdf>.

19
20 49 Coombes R. Rationing of joint replacements raises fears of further cuts. BMJ
21 2005;**331**:1290. doi: 331/7528/1290 [pii] 10.1136/bmj.331.7528.1290
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 **Figure 1: EQ-5D utility by subgroup**
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

For peer review only

Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy, 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life); OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])[41]; TKR, total knee replacement. Error bars show 95% confidence intervals. EQ-5D utility is measured on a scale from 1 (perfect health) to -0.594, where 0 is equivalent to death.

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

1
2
3 **Figure 2:** Cost-effectiveness prediction charts estimated based on predictions of
4 regression models.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30







31 Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy,
32 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life);
33 OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to
34 48 [no problems])[41]; QALY, quality-adjusted life-year; TKR, total knee replacement.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure 3:** Cost-effectiveness acceptability curves for different patient subgroups at
4 different baseline Oxford Knee Scores (OKS). A: American Society of
5 Anesthesiologists (ASA) grade 1 and 2 patients. B: ASA grade 3 patients.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

-000332 on 30 January 2012. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2021 by guest. Protected by copyright.



1
2
3
4
5
6
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43



Full model

Simplified model: OKS and ASA only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43





Legend

- Clearly cost-effective: <£20,000/QALY gained
- Possibly cost-effective: £20,000-£30,000/QALY gained
- Cost-ineffective: >£30,000/QALY gained
- TKR dominated (no clinical benefit)

-000332 on 30 January 2012. Added from <http://bmjopen.bmj.com/> on May 13, 2021 by guest. Protected by copyright.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43



1-000332 on 30 January 2012. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2015 by guest. Protected by copyright.

Rationing of total knee replacement: A cost-effectiveness analysis on a large trial dataset

Running header:

Rationing of total knee replacement

Key words:

Economic evaluation; total knee replacement; heterogeneity; subgroup analysis; randomised controlled trial.

Authors:

Helen Dakin,¹ Alastair Gray,¹ Ray Fitzpatrick,² Graeme MacLennan,³ and David Murray⁴

Collaborator: [The KAT Trial Group](#)

Formatted: Font: Not Bold

Authors' affiliation:

¹ Health Economics Research Centre, Department of Public Health, University of Oxford, Old Road Campus, Oxford, UK

² Department of Public Health, University of Oxford, Old Road Campus, Oxford, UK

³ Health Services Research Unit, University of Aberdeen, Aberdeen, UK

⁴ Nuffield Department of Orthopaedics, [Rheumatology and Musculoskeletal Sciences](#), University of Oxford, UK

Deleted: Surgery

Corresponding author:

Helen Dakin
Researcher
Health Economics Research Centre
Department of Public Health, University of Oxford
Old Road Campus
Headington
Oxford
OX3 7LF
Email: helen.dakin@dph.ox.ac.uk
Tel: 01865 289422
Fax: 01865 289271

Word count: 3,691, (main text of introduction, methods, results and discussion)

Formatted: Not Highlight

Number of figures: 3

Deleted: 398

Number of tables: 2

Ethics: The KAT trial was approved by the Multi Centre Research Ethics Committee for Scotland in November 1998 (research protocol MREC/98/0/100) and was approved by the Local Research Ethics Committees in each study centre recruiting trial participants. Further details are available on request.

Data sharing statement: No additional data available.

Protocol: Available at <http://www.hta.ac.uk/protocols/199500100001.pdf>.

Formatted: Right: 0.25"

Abstract

Objectives: Many UK primary care trusts (PCTs) have recently introduced eligibility criteria restricting total knee replacement (TKR) to patients with low pre-operative Oxford Knee Scores (OKS) to cut expenditure. We evaluate these criteria by assessing the cost-effectiveness of TKR compared with no knee replacement for patients with different baseline characteristics from an NHS perspective.

Design: The cost-effectiveness of TKR in different patient subgroups was assessed using regression analyses of patient-level data from the Knee Arthroplasty Trial (KAT), a large, pragmatic, randomised trial comparing knee prostheses.

Setting: 34 UK hospitals.

Participants: 2131 osteoarthritis patients undergoing TKR.

Interventions and outcome measures: Costs and quality-adjusted life-years (QALYs) observed in the KAT trial within five years of TKR were compared with conservative assumptions about the costs and outcomes that would have been accrued had TKR not been performed.

Results: On average, primary TKR and five years of subsequent care cost £7,458 per patient (standard deviation, SD: £4,058), and patients gained an average of 1.33 (SD: 1.43) QALYs. As result, TKR cost £5,623 per QALY gained. Although costs and health outcomes varied with age and sex, TKR cost less than £20,000 per QALY gained for patients with ASA grade 1-2 who had baseline OKS <40, and for ASA grade 3 patients with OKS <35, even with highly conservative assumptions about costs and outcomes without TKR. Body mass index had no significant effect on costs or outcomes. Restricting TKR to patients with pre-operative OKS <27, would inappropriately deny a highly cost-effective treatment to >10,000 patients annually.

Conclusions: TKR is highly cost-effective for most current patients if the NHS is willing to pay £20,000-£30,000 per QALY gained. At least 97% of TKR patients in England have more severe symptoms than the thresholds we have identified, suggesting further rationing by OKS is probably unjustified.

Trial registration: ISRCTN 45837371

Deleted: controlled

Deleted: different

Deleted: over

Deleted: a

Deleted: -

Deleted: time horizon

Deleted: following

Deleted: ¶

Deleted: of

Deleted: or

Deleted: of 39 or below

Deleted: of 34 or below

Deleted: (BMI)

Deleted: of 26 or less

Deleted: more than

Formatted: Right: 0.25"

Article summary

Article focus

- We assess the cost-effectiveness of total knee replacement (TKR) compared with no knee replacement for patients with different baseline characteristics from a National Health Service (NHS) perspective.
- In particular, we assess the appropriateness of eligibility criteria recently introduced by many UK primary care trusts (PCTs), which restrict TKR to patients with low (*i.e.* [poor](#)) pre-operative Oxford Knee Scores (OKS) to cut expenditure.

Key messages

- We find TKR to be highly cost-effective, costing £5,623 per quality-adjusted life-year (QALY) gained for the average patient.
- TKR costs less than £20,000 per QALY gained for healthy patients with OKS of [<40](#), or [<35](#) for patients who have other conditions restricting their daily activities.
- We find no evidence to support the criteria for restricting access to TKR that have been proposed by some PCTs and calculate that restricting TKR to those patients with pre-operative OKS of 26 or less would deny a highly cost-effective treatment to more than 10,000 patients per year.

Strengths and limitations of this study

- This is the first study assessing how the cost-effectiveness of TKR varies with OKS and the first assessing the clinical/economic implications of the newly-introduced rationing criteria.
- Analyses are based on patient-level data from a large pragmatic trial with detailed, prospective collection of utilities, baseline characteristics and all major knee-related NHS resource use, [including revisions and ambulatory care](#).
- Our study makes several highly conservative assumptions: in particular, assuming that patients would have accrued no knee-related costs and remained at baseline utility without TKR. Furthermore, the KAT sample included only 37 patients with pre-operative OKS >35. As result, TKR may be also cost-effective for some patients with OKS above 39.

Deleted: ≤39

Deleted: ≤34

Formatted: Right: 0.25"

Introduction

Total knee replacement (TKR) is a highly effective treatment for patients with bone-on-bone osteoarthritis and significant knee symptoms, producing substantial reductions in symptoms (particularly pain)[1] and lasting for at least 15 years in 83-94% of cases.[2] However, about 18% of patients consider the outcomes of their surgery to be only fair or poor,[1] and a small proportion experience complications.[3-5] Previous economic evaluations have found TKR to be highly cost-effective, with a cost-effectiveness ratio between €1,276 and \$18,300 per quality-adjusted life-year (QALY) gained for the average patient[6-10]: well below the £20,000-£30,000 per QALY range that the National Institute for Health and Clinical Excellence (NICE) considers cost-effective.[11] However, studies suggest that the costs and benefits of TKR vary between patient subgroups, with TKR being more cost-effective in younger patients,[9] those attending high-volume centres[7] and those deemed low-risk based on age, comorbidities, and poverty criteria.[7] There is also evidence that older[12] and obese[13] patients have higher medical costs, although other studies suggest that knee replacement is also cost-effective in nonagenarians[14] and draw conflicting conclusions on how benefits vary with body mass index (BMI).[15-17]

In the current economic environment there is great pressure to reduce NHS expenditure and numerous Primary Care Trusts (PCTs) have proposed cutting costs by limiting access to joint replacement[18-25] or classifying these procedures as being of limited value.[24-26] Since around 60,000 primary TKR procedures are conducted in England and Wales each year,[27] savings could be made with little loss of patient welfare if it were possible to reliably predict which patients obtain little/no benefit from TKR based on explicit, evidence-based criteria. Although the widely-used Oxford Knee Score (OKS) was developed to assess outcomes of knee replacement in trial populations,[28] several PCTs have set maximum OKS thresholds (ranging from 18 to 32), above which patients are ineligible for TKR.[18-24]. However, we are not aware of any clinical or cost-effectiveness justification for these thresholds. Furthermore, many patients with higher OKS have significant symptoms or limitations to daily activities that could be improved by surgery. If rationing policies are to be adopted, cost-effectiveness analysis offers the only fair basis for designing them, since it ensures that NHS resources are allocated to maximise the health gained from available resources. As treatments are generally considered cost-effective if they cost less than £20,000-£30,000 per quality-adjusted life-year (QALY) gained,[11] restricting TKR to those patients for whom the procedure

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Deleted: [12]

Field Code Changed

Deleted: [13]

Field Code Changed

Deleted: [14-16]

Field Code Changed

Deleted: [17-24]

Field Code Changed

Deleted: [23-25]

Field Code Changed

Deleted: [26]

Field Code Changed

Deleted: [27]

Field Code Changed

Deleted: [17-23]

Field Code Changed

Formatted: Right: 0.25"

costs less than £20,000-£30,000 per QALY gained would be a reasonable approach, potentially realising savings that could be invested in other treatments giving greater health gains. However, we are aware of no published evidence assessing how the cost-effectiveness of TKR varies with OKS.

We therefore use data from the Knee Arthroplasty Trial (KAT)^[29, 30] to assess the appropriateness of rationing knee replacement by OKS, age, BMI, and clinical characteristics, by calculating the cost-effectiveness of TKR in different patient subgroups. In contrast to previous studies evaluating the cost-effectiveness of TKR that relied upon data from small cohorts with short follow-up,^[6-10, 14] the KAT trial dataset provides detailed, prospective individual patient data on baseline characteristics, treatments, complications, costs, and quality of life, and now has more than five years' follow-up on 2352 patients.

Methods

We assessed the cost-effectiveness of primary TKR compared with no knee replacement for patients with different baseline characteristics. Our analysis took a National Health Service (NHS) perspective and excluded personal and social services from the analysis, although TKR is likely to delay admission to residential care,^[14] reduce personal care costs and help recipients to continue paid employment. A cost-utility analysis was conducted, which assessed cost-effectiveness in terms of cost per quality-adjusted life-year (QALY) gained to capture the quality of life gains associated with TKR.

Data on costs and quality of life following TKR were taken from the KAT trial (ISRCTN 45837371): a pragmatic, partial-factorial, unblinded randomised controlled trial (RCT) in which 2352 participants attending 34 UK centres were randomly allocated to undergo TKR with/without a metal-backed tibial component, with/without patellar resurfacing, and/or with/without a mobile bearing.^[29, 30] Five-year trial results demonstrated that patients experience substantial improvements in quality of life and functional status following knee replacement that are maintained out to five years^[30]; further follow-up is ongoing. Our analysis took a five-year time horizon in line with the results published to date^[30]; costs and health benefits occurring more than five years after primary TKR were conservatively excluded.

Field Code Changed
Deleted: [28, 29]

Field Code Changed
Deleted: [6-10, 13]

Formatted: English (U.S.)

Deleted: [13]
Field Code Changed

Field Code Changed
Deleted: [28, 29]

Field Code Changed
Deleted: [29]

Field Code Changed
Deleted: [29]

Formatted: Right: 0.25"

BMJ Open: first published as 10.1136/bmjopen-2011-006332 on 30 January 2012. Downloaded from http://bmjopen.bmj.com/ on May 13, 2021 by guest. Protected by copyright.

KAT participants completed the Oxford Knee Score (OKS) and EQ-5D questionnaires immediately before and three months after TKR and annually thereafter. The OKS is a validated 12-item questionnaire [for assessing](#) health status and outcomes in relation to knee replacement [that gives an unweighted total score ranging from 0 \(severe problems on all items\) to 48 \(no problems on any item\).](#)^[28] The EQ-5D is a generic health state preference measure^[31]; health state preference values (“utilities”) for EQ-5D profiles were based on time-trade-off valuations by members of the UK general public.^[31] EQ-5D utilities range from -0.594 (extreme problems on all five domains) to 1 (perfect health) and indicate the value placed on different levels of health. The number of QALYs that each patient accrued following TKR was calculated as the area under the utility curve, with linear interpolation between utility measurements; those patients who died were assumed to remain at the last observed utility until death.

Healthcare resource use data (including knee-related outpatient, general practice and physiotherapy consultations, and theatre time, hospital days, complications and knee components used during the primary admission for TKR and during any knee-related readmissions [or revisions](#)) were collected prospectively for all KAT participants and valued at 2007-08 prices, as described previously.^[30] Costs incurred after Year 1 were discounted at 3.5% per year based on current UK recommendations.^[32]

The number of QALYs gained from TKR was calculated as the number of QALYs observed minus the number of QALYs expected without TKR. Since all KAT participants underwent TKR, assumptions were needed to estimate the costs and QALYs that would have been accrued if no TKR procedures had occurred in the five-year time horizon. We assumed that:

- Without TKR, patients would have remained at their baseline EQ-5D utility for five years or until death (if sooner). In practice, patients’ symptoms and quality of life are likely to deteriorate due to worsening arthritis^[17, 33] and increasing age.^[34]
- The seven KAT participants who died before hospital discharge would have survived for five years at baseline utility without TKR but that TKR had no effect on the date of death for other patients. This assumption is highly conservative since those patients who die soon after joint replacement tend to be older^[5, 35] and have comorbidities that could have led to death without TKR.^[5]
- Patients would not have used any healthcare resources due to their knee problems if they had not undergone TKR. This assumption is highly conservative

Deleted: to

Deleted: ,[27]

Deleted: while t

Deleted: .

Field Code Changed

Deleted: [30]

Deleted: H

Field Code Changed

Deleted: [30]

Field Code Changed

Deleted: [29]

Field Code Changed

Deleted: [31]

Deleted: [16, 32]

Field Code Changed

Deleted: [33]

Field Code Changed

Field Code Changed

Deleted: [5, 34]

Field Code Changed

Formatted: Right: 0.25"

since osteoarthritis patients will receive medical management without TKR and a recent Canadian case-control study suggested that TKR recipients accrued lower costs 6-18 months after the procedure than matched controls who had arthritis but no TKR.^[33]

Since these assumptions are likely to underestimate the benefits and overestimate the additional costs of TKR, we evaluated the effect of relaxing these assumptions in sensitivity analyses.

The 7% of data on baseline characteristics, resource use and quality of life that were missing were imputed by multiple imputation using the ice command^[36-38] (version 1.3.0) within Stata Version 11.0 (College Station, TX), thereby avoiding the bias and inefficiency associated with complete case analysis.^[38-40] Treatment indicators and the covariates included in subsequent regression analyses were included in the imputation model to avoid bias.^[38] Fifty imputed datasets were generated and results combined.

Regression models were used to estimate how the costs and QALYs gained from TKR vary with baseline characteristics and to predict outcomes for different patient groups without subdividing the trial population into small patient subgroups. Since costs and QALYs were highly skewed, generalised linear models (GLM) with gamma family distributions were used to predict total costs and the QALYs accrued with or without TKR; prior to regression, the number of QALYs accrued was subtracted from the maximum number of QALYs that could have been accrued over the five-year trial period to ensure that all QALY data were positive. We evaluated the effect of six baseline characteristics on costs and QALYs with/without TKR:

- Male sex;
- Age at time of operation;
- Baseline OKS (using the new, 0-48, scoring system^[41]);
- American Society of Anesthesiologists (ASA) grade 3 (symptomatic illness with minimal restriction on daily life) rather than grades 1 (fit & healthy) or 2 (asymptomatic illness with no restriction on daily life);
- BMI (kg/m²);
- Presence of arthritis in one knee rather than in both knees or generally.

Deleted: [32]

Field Code Changed

Field Code Changed

Deleted: [35-37]

Field Code Changed

Deleted: [37-39]

Field Code Changed

Deleted: [37]

Deleted: Fem

Field Code Changed

Deleted: [40]

Formatted: Right: 0.25"

Since the introduction of biologic therapies has changed management of rheumatoid arthritis and as rheumatoid arthritis patients and those with severe comorbidities are likely have very different costs and outcomes from most patients, 108 rheumatoid arthritis patients and 13 patients with ASA grade 4 were excluded from the analysis. One hundred 100 patients who were randomised to total vs unicompartmental knee replacement or died or withdrew from the trial before surgery were also excluded from the analysis, giving a final sample size of 2131.

All regression analyses were conducted in Stata Version 11. Bootstrapping was used to capture uncertainty around cost-effectiveness ratios and correlations between costs and QALYs with/without TKR; the regression models for costs and outcomes were run for each of 125 bootstrap samples drawn from each imputed KAT dataset and results combined using Rubin's rule.^[39] Predicted costs and the predicted number of QALYs accrued with and without TKR in different patient groups were calculated from regression coefficients and used to estimate incremental cost-effectiveness ratios (ICERs). Bootstrap results were used to calculate cost-effectiveness acceptability curves (CEACs)^[42] which plot the probability of treatment being cost-effective against the ceiling ratio (i.e. the maximum that society is willing or able to pay per QALY gained).

We also calculated the number of patients in England and Wales eligible for TKR under different rationing policies based on 61,651 patients undergoing primary TKR each year.^[27] The proportion of patients eligible for TKR under different rationing criteria was based on analysis of an extract of patient-level Patient Reported Outcome Measures^[1] (PROMs) data on admissions for knee replacement up to 31st December 2010, supplemented where necessary by the baseline characteristics from the KAT sample.

Results

Of the 2131 KAT participants who had osteoarthritis, were ASA grades 1-3 and underwent TKR as part of the trial, the mean baseline OKS was 18 (standard deviation, SD: 7.5), mean age 71 (SD: 8.0), mean BMI 30 (SD: 5.5), and 45% (956/2131) were male (Figure 1); further baseline characteristics of the trial population have been reported previously.^[29]

Deleted: of

Field Code Changed

Deleted: [38]

Field Code Changed

Deleted: [41]

Deleted: [26]

Field Code Changed

Field Code Changed

Deleted: fe

Field Code Changed

Deleted: [28]

Formatted: Right: 0.25"

1
2
3
4
5
6 Mean EQ-5D utility rose substantially within three months of TKR from 0.39 at
7 baseline to 0.71 at one year, and declined gradually thereafter (Figure 1, Table 1).
8 However, baseline utility and the quality of life gains associated with TKR varied
9 between patient subgroups. In particular, women and patients with low OKS (i.e. poor
10 knee function), higher ASA grade or obesity at baseline tended to have lower quality
11 of life at all timepoints (Figure 1), although those in the lowest OKS quintile
12 experienced significantly greater health gains from TKR than the top quintile
13 ($p < 0.001$; calculated using linear regression, combining imputed datasets using [the](#)
14 [mim](#) command).
15
16
17

18
19
20 On average, each admission for primary TKR cost £6,363 (SD: £1,702).
21 Readmissions, revision procedures and GP, outpatient, and physiotherapy
22 consultations related to the study knee over the following five years cost a further
23 £1,095 (SD: £3,579) per patient, giving a total cost of £7,458 (SD: £4,058) per
24 patient. TKR was more costly in patients with low baseline OKS, but the greater
25 QALY gains meant that TKR was better value for money in this patient group than in
26 patients with moderate symptoms (Table 1). Subgroup analyses suggested that TKR
27 cost less than £11,000/QALY gained across all OKS deciles: well below the £20,000-
28 £30,000/QALY gained range that NICE currently consider good value for money.^[11]
29
30
31

32
33 However, cost-effectiveness also varied substantially between patients within each
34 OKS decile. We therefore used regression models to estimate the costs, QALY gains
35 and cost-effectiveness of TKR in a wide range of patient subgroups while controlling
36 for multiple baseline characteristics. Our results indicated that four factors have a
37 significant effect on cost-effectiveness: age, sex, baseline OKS and ASA grade
38 (Table 2). BMI and the presence of arthritis in other joints besides the study knee had
39 no significant effect on either the costs or QALY gains associated with TKR ($p > 0.05$).
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Deleted: morbid

Field Code Changed

Formatted: Right: 0.25"

Table 1: EQ-5D utility, costs and estimated QALYs for subgroups with different baseline OKS

OKS decile	1 (worst)	2	3	4	5	6	7	8	9	10 (best)	All patients	
Baseline OKS range	<9	9-11	12-13	14-15	16-17	18-19	20-21	22-24	25-27	>27	0-48	
No. KAT participants	204	213	186	232	193	224	195	256	194	234	2131	
EQ-5D utility (SE)†	Baseline	0.03 (0.004)	0.14 (0.005)	0.21 (0.006)	0.25 (0.006)	0.36 (0.006)	0.46 (0.006)	0.55 (0.005)	0.57 (0.004)	0.61 (0.004)	0.67 (0.003)	0.39 (0.002)
	3 mths	0.54 (0.007)	0.58 (0.006)	0.64 (0.006)	0.66 (0.005)	0.71 (0.005)	0.71 (0.005)	0.70 (0.004)	0.74 (0.004)	0.75 (0.005)	0.77 (0.004)	0.68 (0.002)
	1 year	0.51 (0.007)	0.60 (0.007)	0.68 (0.006)	0.70 (0.006)	0.71 (0.006)	0.75 (0.005)	0.69 (0.006)	0.78 (0.004)	0.79 (0.005)	0.84 (0.004)	0.71 (0.002)
	2 year	0.49 (0.008)	0.54 (0.007)	0.66 (0.007)	0.67 (0.006)	0.68 (0.007)	0.73 (0.005)	0.66 (0.007)	0.76 (0.005)	0.78 (0.005)	0.81 (0.005)	0.68 (0.002)
	3 year	0.47 (0.008)	0.53 (0.007)	0.64 (0.007)	0.63 (0.006)	0.67 (0.007)	0.71 (0.006)	0.68 (0.007)	0.72 (0.005)	0.76 (0.006)	0.79 (0.006)	0.66 (0.002)
	4 year	0.43 (0.008)	0.49 (0.008)	0.61 (0.007)	0.62 (0.007)	0.65 (0.007)	0.68 (0.007)	0.64 (0.007)	0.70 (0.006)	0.71 (0.007)	0.75 (0.006)	0.63 (0.002)
	5 year	0.43 (0.008)	0.50 (0.008)	0.57 (0.008)	0.58 (0.007)	0.63 (0.007)	0.67 (0.007)	0.59 (0.007)	0.68 (0.006)	0.67 (0.008)	0.73 (0.007)	0.61 (0.002)
Total cost (SE)*	£8657 (£157)	£7715 (£81)	£7495 (£73)	£7081 (£63)	£7185 (£69)	£7567 (£82)	£7619 (£102)	£7128 (£61)	£7399 (£94)	£6917 (£67)	£7458 (£28)	
QALYs with TKR (SE)*	2.17 (0.029)	2.48 (0.027)	2.94 (0.024)	2.99 (0.022)	3.13 (0.025)	3.30 (0.022)	3.09 (0.025)	3.40 (0.019)	3.50 (0.020)	3.67 (0.020)	3.08 (0.025)	
QALYs without TKR (SE)*	0.15 (0.02)	0.62 (0.022)	0.94 (0.026)	1.14 (0.026)	1.63 (0.029)	2.04 (0.026)	2.39 (0.026)	2.55 (0.021)	2.78 (0.02)	3.02 (0.02)	1.75 (0.031)	
QALY gain from TKR (SE)*	2.02 (0.032)	1.86 (0.031)	2.00 (0.033)	1.85 (0.031)	1.51 (0.034)	1.26 (0.030)	0.69 (0.027)	0.85 (0.023)	0.73 (0.02)	0.65 (0.02)	1.33 (0.01)	
ICER*	£4,295	£4,153	£3,747	£3,836	£4,770	£6,007	£10,971	£8,391	£10,167	£10,697	£5,623	

* Discounted at 3.5% per annum

† EQ-5D utility is measured on a scale from 1 (perfect health) to -0.594, where 0 is equivalent to death.

Abbreviations: ICER, incremental cost-effectiveness ratio; OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems]); QALY, quality-adjusted life-year; SE, standard error around the mean; TKR, total knee replacement.

Formatted ... [3]

Deleted: 0.54 (0.007)

Deleted: 0.58 (0.006)

Deleted: 0.64 (0.006)

Deleted: 0.66 (0.005)

Deleted: 0.71 (0.005)

Deleted: 0.71 (0.005)

Deleted: 0.70 (0.004)

Deleted: 0.74 (0.004)

Deleted: 0.61 (0.004)

Deleted: 0.67 (0.003)

Deleted: 0.39 (0.002)

Formatted ... [2]

Formatted ... [4]

Deleted: 0.51 (0.007)

Deleted: 0.60 (0.007)

Deleted: 0.68 (0.006)

Deleted: 0.70 (0.006)

Deleted: 0.71 (0.006)

Deleted: 0.75 (0.005)

Deleted: 0.69 (0.006)

Deleted: 0.78 (0.004)

Deleted: 0.75 (0.005)

Deleted: 0.77 (0.004)

Deleted: 0.68 (0.002)

Formatted ... [5]

Deleted: 0.49 (0.008)

Deleted: 0.54 (0.007)

Deleted: 0.66 (0.007)

Deleted: 0.67 (0.006)

Deleted: 0.68 (0.007)

Deleted: 0.73 (0.005)

Deleted: 0.66 (0.007)

Deleted: 0.76 (0.005)

Deleted: 0.79 (0.005)

Deleted: 0.84 (0.004)

Deleted: 0.71 (0.002)

Formatted ... [6]

Deleted: 0.47 (0.008)

Deleted: 0.53 (0.007)

Deleted: 0.64 (0.007)

Deleted: 0.63 (0.006)

Deleted: 0.67 (0.007)

Deleted: 0.68 (0.007)

Table 2: Results of the regression analysis

Baseline characteristic	Coefficient (SE)		
	QALY loss with TKR†	QALY loss without TKR†	Total costs with TKR (£)
Male sex	0.056 (0.065)	-0.001 (0.054)	593 (181)*
Age at operation (years)	0.007 (0.003)*	0.002 (0.004)	-8 (12)
Pre-operative OKS	-0.040 (0.004)*	-0.087 (0.004)*	-52 (12)*
ASA grade 3	0.414 (0.067)*	0.227 (0.087)*	492 (208)*
Constant	1.712 (0.201)*	4.274 (0.325)*	8573 (881)*

* p<0.05

† QALY loss indicates the number of discounted QALYs that would have been accrued if patients had experienced an EQ-5D utility of 1 for five years (4.67 QALYs) minus the number of discounted QALYs that the patient actually experienced.

The incremental cost-effectiveness ratio (expressed as cost/QALY) for any given patient equals: $(\text{male} \times 593 - \text{age} \times 8 - \text{OKS} \times 52 + \text{ASAgrade3} \times 492 + 8573) / ((\text{male} \times -0.001 + \text{age} \times 0.002 - \text{OKS} \times 0.087 + \text{ASAgrade3} \times 0.227 + 4.274) - (\text{male} \times 0.056 + \text{age} \times 0.007 - \text{OKS} \times 0.040 + \text{ASAgrade3} \times 0.414 + 1.712))$. Costs and QALYs were discounted at 3.5% per annum.

Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy, 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life); OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])^[41]; QALY, quality-adjusted life-year; TKR, total knee replacement.

Figure 2 shows the cost-effectiveness of TKR for each multivariate group, using charts similar in design to widely-used cardiovascular risk tables.^[43] The charts demonstrate that TKR is clearly cost-effective across a wide range of patient groups. For example, TKR costs less than £20,000/QALY gained for 71-year-old women with no symptomatic illness unless baseline OKS is above 40 (equivalent to having very minor problems on up to eight of the 12 activities mentioned in OKS or moderate-severe pain only on exertion). The threshold OKS at which TKR becomes cost-ineffective was lower for older patients, men, and those with symptomatic disease, but TKR remained cost-effective for all patients with OKS below 32 regardless of age, sex, or ASA grade. A simplified analysis controlling only for OKS and ASA grade suggested that TKR is cost-effective for all ASA grade 1 or 2 patients with baseline OKS <40, and all ASA grade 3 patients with OKS <35.

There was substantial uncertainty around the cost-effectiveness ratios for patients with very high baseline OKS, since only 37 trial participants had OKS above 35 at baseline. However, we estimate that there is a 99% probability that TKR is cost-effective (at a £20,000/QALY ceiling ratio) for the average ASA grade 1-2 patient with a baseline OKS of 35, and a 53% probability for ASA grade 3 (Figure 3).

Since each additional baseline characteristic increases the complexity of the eligibility criteria and may also raise ethical or equity concerns, we estimated the number of patients who would receive TKR under different eligibility criteria. This analysis suggested that restricting TKR to those patients for whom the operation was

Deleted: Fem
Formatted

Deleted: fe
Deleted: or4
Deleted: fe
Deleted: fe

Field Code Changed
Deleted: [40]

Field Code Changed
Deleted: [42]

Deleted: maximum
Deleted: wo

Formatted: Not Highlight
Deleted: of 39 or less
Formatted: Not Highlight
Deleted: ≤34

Formatted: Right: 0.25"

1
2
3
4
5
6 predicted to cost less than £20,000/QALY gained based on the simplified model
7 (allowing only for ASA grade and OKS) would lead to 2000 fewer TKR procedures
8 being conducted in England and Wales each year (3% fewer than the 61,651
9 procedures conducted in 2009/10^[27]). If the decision rules were simplified further to
10 provide TKR to all patients with baseline OKS <39, all but 1700 of the 61,651 patients
11 currently receiving TKR in England and Wales in 2009/10^[27] would be eligible.
12
13 Allowing for other baseline characteristics had minimal effect on patient numbers or
14 predicted costs or benefits, suggesting that decisions about suitability for TKR can be
15 based on OKS alone.
16
17

18
19
20 Sensitivity analyses were conducted to assess the effect of extending the time
21 horizon of our analysis up to 10 years, and of assuming that EQ-5D utility would have
22 fallen by 0.5% per year without TKR,^[34] that all knee-related ambulatory
23 consultations would have occurred without TKR, and that the mean length of stay
24 has fallen by 41% (5.9 days^[27] vs 10.0) since the KAT operations took place in
25 1999-2003. Changing any of these assumptions produced lower ICERs than the
26 base case analysis, confirming that the current analysis is highly conservative and
27 suggesting that TKR may be cost-effective for ASA grade 1-2 patients with baseline
28 OKS as high as 43. Excluding patients with incomplete data had no effect on
29 conclusions. Although statistically significant, adding an age-squared term to the
30 regression model had no effect on the conclusions, while a sensitivity analysis adding
31 an OKS-squared term suggested that TKR may be cost-effective for all patients of
32 ASA grade 1 or 2. Recoding BMI as a dummy variable indicating whether or not
33 patients were obese (BMI ≥30) did not change the conclusions and obesity had no
34 significant effect on costs or QALYs.
35
36
37
38
39

40 Discussion

41
42
43 We found TKR to cost £5,623 per QALY gained for the average patient (similar to
44 estimates from previous studies^[6-10]) and therefore represent very good value for
45 money compared with the £20,000-£30,000/QALY threshold typically used in NHS
46 decision-making.^[11] Our study also confirms previous findings^[7, 9, 17] that costs
47 and benefits vary with age and comorbidity, but evaluates such characteristics, and
48 pre-operative knee function, in multivariate analyses, which has not (to our
49 knowledge) been done previously. We found pre-operative OKS to be the best
50 predictor of post-operative costs, outcomes and cost-effectiveness.
51
52
53
54
55
56
57
58
59
60

Field Code Changed

Deleted: [26]

Deleted: ≤38

Field Code Changed

Deleted: [26]

Deleted: [33]

Field Code Changed

Field Code Changed

Deleted: [26]

Formatted: English (U.K.)

Field Code Changed

Field Code Changed

Field Code Changed

Deleted: [7, 9, 16]

Formatted: Right: 0.25"

Our analyses were based on patient-level data from a large trial with detailed, prospective collection of costs, utilities and baseline characteristics that permit extensive subgroup analyses. [KAT used a pragmatic design, with broad inclusion criteria and no restrictions on peri- or post-operative care other than randomised aspects of component design](#)[29, 30]; as result, the trial cohort is similar to the [national PROMs cohort and costs and benefits are likely to be comparable to those in routine clinical practice](#). Furthermore, unlike some previous economic evaluations, our study took full account of uncertainty and included all knee-related NHS costs accrued in the first five years after TKR. The main limitation of our analysis was the small number of patients with high OKS. Since KAT recruited only 37 patients with OKS >35, further studies are required to evaluate TKR in this patient subgroup. Analyses were also based on *post hoc* comparisons between outcomes before and after TKR, rather than between randomly-assigned treatment groups. As a result, our study does not provide unbiased estimates of the QALY gains from TKR, and regression towards the mean may have caused our study to overestimate the quality of life improvement attributable to TKR and/or the effect of pre-operative OKS on QALY gains.

We made several highly conservative assumptions: in particular, assuming that patients would have accrued no knee-related costs and would have remained at baseline utility without TKR, whereas previous studies have found that patients accrue substantial costs and show clinical worsening without TKR.[33] We also assumed that those patients who died before hospital discharge would have survived for five years if they had not had TKR; in reality, those patients dying soon after TKR tend to be older[5, 35] and frailer[5] pre-operatively and some studies suggest that patients undergoing hip replacement have higher long-term survival than the general population[44, 45], (although this may be due to patient selection[35, 44]). When these assumptions were relaxed, TKR was cost-effective for ASA grade 1-2 patients with baseline OKS <44. The costs and benefits of TKR are likely to vary between centres and over time due to variations in patient demographics, component prices, length of stay, and operation time; sensitivity analyses demonstrated that TKR may have become more cost-effective since the KAT procedures were conducted in 1999-2003 due to reductions in length of stay.

Our analyses show that if TKR were to be rationed based on cost-effectiveness, OKS would be a reasonable tool to use to set the threshold, although there may, of

Deleted: pragmatic

Deleted: [32]

Field Code Changed

Field Code Changed

Deleted: [5, 34]

Field Code Changed

Field Code Changed

Deleted: [43, 44]

Field Code Changed

Deleted: [34, 43]

Deleted: ≤43

Formatted: Right: 0.25"

course, be practical difficulties with using a patient-reported measure to determine treatment eligibility: particularly if patients know how their questionnaire responses will be used or if test-retest reliability is low in this setting. [The decision to operate must also take into consideration other factors, such as radiographic findings and patient choice.](#) Based on regression and subgroup analyses, we can be confident that TKR costs less than £20,000/QALY for all ASA grade 1-2 patients with baseline OKS [<40](#) and for ASA grade 3 patients with OKS [<35](#). Using these thresholds to determine who received treatment would avoid around 2000 TKR procedures per year in England and Wales, saving around £11.8 million per year. However, the higher thresholds suggested by sensitivity analyses using less conservative assumptions suggest that TKR is cost-effective for all but 100 patients per year, which is unlikely to achieve sufficient savings to warrant rationing of TKR. Although the cost-effectiveness of knee replacement was also influenced by gender and age, taking account of these factors as well as OKS and ASA grade would increase the complexity of eligibility criteria, reduce equality of access to healthcare and have minimal effect on NHS budgets.

[The results also demonstrate](#) that the thresholds proposed by some PCTs[\[18-24\]](#) are inappropriate and would deny a highly cost-effective treatment to thousands of patients with severe arthritis. The current situation also introduces a postcode lottery with different PCTs using different eligibility criteria. We estimate that limiting knee replacement to patients with OKS ≤ 26 would prevent more than 10,000 people from receiving TKR. This would be inequitable as TKR costs £10,697 per QALY gained for this group, whereas NICE routinely recommend treatments with ICERs in the region of £20,000-£30,000/QALY gained[\[11, 46, 47\]](#) and are willing to pay substantially more for end of life care.[\[48\]](#) Furthermore, the finding that TKR and subsequent monitoring is more costly for patients with low OKS suggests that delaying TKR until symptoms have deteriorated may be a false economy. Although some PCTs have considered restricting TKR further for obese patients,[\[18, 21, 25, 49\]](#) we found BMI to have no significant effect on costs ($p=0.442$), QALY gains ($p=0.098$) or the incidence of peri- or post-operative complications ($p>0.26$). [Although BMI remained non-significant](#) even when ASA grade was omitted from regression analyses, [it is difficult to isolate the effect of obesity from other correlated variables \(e.g. comorbidity\): as result, further research on obesity may be warranted: particularly since a previous study observed a significant correlation between BMI and costs.](#)[\[13\]](#)

Deleted: ≤ 39 Deleted: ≤ 34

Deleted: all

Deleted: Our analysis

Deleted: shows

Deleted: [17-23]

Field Code Changed

Field Code Changed

Deleted: [11, 45, 46]

Field Code Changed

Deleted: [47]

Field Code Changed

Deleted: [17, 20, 24, 48]

Deleted: :

Deleted: .

Formatted: Right: 0.25"

Eligibility criteria for rheumatoid arthritis patients and patients with ASA grade 4 remain unclear, although ASA grade 4 patients would normally be considered unfit for surgery, and the decision to operate on such patients would be based on a case-by-case consideration of whether the benefits outweigh the risks. While our study focused on the costs and benefits of TKR, patients with high baseline OKS often undergo unicompartmental knee replacement; [further research is needed to evaluate how the cost-effectiveness of unicompartmental replacement varies with OKS.](#)

[In conclusion, TKR is highly cost-effective for the vast majority of patients who currently undergo this procedure in the UK. Although costs and health benefits vary with baseline OKS, the rationing thresholds proposed by some PCTs\[18-24\] are not supported by evidence on health outcomes or cost-effectiveness.](#)

Acknowledgements

The authors would like to thank all those involved in the KAT trial for their role in running the trial and collecting the data used in this study, particularly other members of the KAT Project Management Group: Suzanne Breeman; Marion Campbell; Jackie Ellington; Nick Fiddian; Adrian Grant; Linda Johnston; and Richard Morris.

The Knee Arthroplasty Trial is funded by the NIHR Health Technology Assessment Programme (project number 95/10/01) and will be published in full as a Health Technology Assessment report. Visit the HTA programme website for further project information. Additional funding for research support in clinical centres was provided by Howmedica Osteonics; Zimmer; DePuy, a Johnson and Johnson company; Corin Medical; Smith and Nephew Healthcare; Biomet Merck; and Wright Cremascoli. The Health Economics Research Centre obtains support from the National Institute of Health Research. The Health Services Research Unit is core funded by the Chief Scientist Office of the Scottish Government Health Directorates. PROMs data were reused with the permission of The Health and Social Care Information Centre, copyright 2011, with all rights reserved. The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the HTA programme, NIHR, NHS, Information Centre or the Department of Health.

Deleted: , which is cheaper than TKR due to lower component prices, fewer complications and shorter inpatient stay.[49-51] The OKS threshold for unicompartmental replacement may therefore be even higher than for TKR, since it produces marked quality of life improvements in patients with high baseline scores.[49]

Deleted: This remains an area for future research

Formatted: Right: 0.25"

Competing interests

All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare no support from any organisation for the submitted work with the exception of those declared in the funding statement. AG, RF and GM declare that they have no financial relationships with any organisations that might have an interest in the submitted work in the previous three years and no other relationships or activities that could appear to have influenced the submitted work. DM receives royalties paid to him by Biomet and HD has undertaken consultancy work for Pfizer on drugs for rheumatoid arthritis but both declare that they have no other financial relationships with any organisations that might have an interest in the submitted work in the previous three years and no other relationships or activities that could appear to have influenced the submitted work.

Funding

The Knee Arthroplasty Trial is funded by the NIHR Health Technology Assessment Programme (project number 95/10/01). Additional funding for research support in clinical centres was provided by Howmedica Osteonics; Zimmer; DePuy, a Johnson and Johnson company; Corin Medical; Smith and Nephew Healthcare; Biomet Merck; and Wright Cremascoli.

The Health Economics Research Centre obtains support from the National Institute of Health Research. The Health Services Research Unit is core funded by the Chief Scientist Office of the Scottish Government Health Directorates. [The Biomedical Research Unit at Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences is funded by the National Institute of Health Research.](#)

The authors conducted the research independently of all funding organisations. The funders had no role in the collection, analysis or interpretation of data, writing of the manuscript or the decision to publish. The authors had full access to all of the data (including statistical reports and tables) in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Formatted: Right: 0.25"

Contributorship statement

HD, AG and DM conceived and developed the principles and methods underpinning the current analysis. GM was involved in primary data collection and database management, while HD conducted the analysis under AG's supervision and guidance from [DM, GM and RF](#). HD drafted the manuscript with input from all other authors; all authors have approved the final manuscript [and were involved in the interpretation of results](#). The KAT trial was [conceived](#), designed and run by the KAT trial group, which comprised Suzanne Breeman, Marion Campbell, HD, Jackie Ellington, Nick Fiddian, RF, Adrian Grant, AG, Linda Johnston, GM, Richard Morris and DM.

References

- 1 [Hospital Episode Statistics. Provisional Monthly Patient Reported Outcome Measures \(PROMs\) in England. April 2009 – April 2010: Pre- and postoperative data: Experimental statistics. 2010 \[updated 2010; cited 6th April 2011\]; Available from: <http://www.hqip.org.uk/assets/NCAPOP-Library/PROMsreport09101.pdf>.](#)
- 2 [Labek G, Thaler M, Janda W, et al. Revision rates after total joint replacement: cumulative results from worldwide joint register datasets. J Bone Joint Surg Br 2011;**93**:293-7. doi: 93-B/3/293 \[pii\] 10.1302/0301-620X.93B3.25467](#)
- 3 [The National Collaborating Centre for Chronic Conditions. Osteoarthritis: National clinical guideline for care and management in adults. London: Royal College of Physicians; 2008 \[updated 2008; cited 29th June 2011\]; Available from: <http://www.nice.org.uk/nicemedia/live/11926/39720/39720.pdf>.](#)
- 4 [Huddleston JI, Maloney WJ, Wang Y, et al. Adverse events after total knee arthroplasty: a national Medicare study. J Arthroplasty 2009;**24**:95-100. doi: S0883-5403\(09\)00168-5 \[pii\] 10.1016/j.arth.2009.05.001](#)
- 5 [Memsoudis SG, Della Valle AG, Besculides MC, et al. Risk factors for perioperative mortality after lower extremity arthroplasty: a population-based study of 6,901,324 patient discharges. J Arthroplasty 2010;**25**:19-26. doi: S0883-5403\(08\)00855-3 \[pii\] 10.1016/j.arth.2008.11.010](#)
- 6 [Navarro Espigares JL, Hernandez Torres E. Cost-outcome analysis of joint replacement: evidence from a Spanish public hospital. Gac Sanit 2008;**22**:337-43. doi: 13125355 \[pii\]](#)
- 7 [Losina E, Walensky RP, Kessler CL, et al. Cost-effectiveness of total knee arthroplasty in the United States: patient risk and hospital volume. Arch Intern Med 2009;**169**:1113-21; discussion 21-2. doi: 169/12/1113 \[pii\] 10.1001/archinternmed.2009.136](#)

Deleted: other authors

Formatted: Indent: Left: 0", First line: 0", Line spacing: 1.5 lines

Field Code Changed

Formatted: Font: (Default) Arial, 11 pt

Formatted: French (France)

Formatted: French (France)

Formatted: Font: Bold, French (France)

Formatted: French (France)

Formatted: Font: (Default) Arial, 11 pt

Formatted: French (France)

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Right: 0.25"

- 8 Rasanen P, Paavolainen P, Sintonen H, et al. Effectiveness of hip or knee replacement surgery in terms of quality-adjusted life years and costs. *Acta Orthop* 2007;**78**:108-15. doi: 776344017 [pii].10.1080/17453670610013501
- 9 Rissanen P, Aro S, Sintonen H, et al. Costs and cost-effectiveness in hip and knee replacements. A prospective study. *Int J Technol Assess Health Care* 1997;**13**:575-88.
- 10 Lavernia CJ, Guzman JF, Gachupin-Garcia A. Cost effectiveness and quality of life in knee arthroplasty. *Clin Orthop Relat Res* 1997:134-9.
- 11 National Institute for Health and Clinical Excellence. Social value judgements: Principles for the development of NICE guidance. Second Edition. 2008 [updated 2008; cited 17th November 2010]; Available from: <http://www.nice.org.uk/media/C18/30/SVJ2PUBLICATION2008.pdf>.
- 12 Brander VA, Malhotra S, Jet J, et al. Outcome of hip and knee arthroplasty in persons aged 80 years and older. *Clin Orthop Relat Res* 1997:67-78.
- 13 Dowsey MM, Liew D, Choong PF. Economic burden of obesity in primary total knee arthroplasty. *Arthritis Care Res (Hoboken)* 2011;**63**:1375-81. doi: 10.1002/acr.20563
- 14 Karuppiyah SV, Banaszkiwicz PA, Ledingham WM. The mortality, morbidity and cost benefits of elective total knee arthroplasty in the nonagenarian population. *Int Orthop* 2008;**32**:339-43. doi: 10.1007/s00264-007-0324-y
- 15 McQueen DA, Long MJ, Algotar AM, et al. The effect of obesity on quality-of-life improvement after total knee arthroplasty. *Am J Orthop (Belle Mead NJ)* 2007;**36**:E117-20, E27.
- 16 de Leeuw JM, Villar RN. Obesity and quality of life after primary total knee replacement *Knee* 1998;**5**:119-23.
- 17 Cushnaghan J, Bennett J, Reading I, et al. Long-term outcome following total knee arthroplasty: a controlled longitudinal study. *Ann Rheum Dis* 2009;**68**:642-7. doi: ard.2008.093229 [pii].10.1136/ard.2008.093229
- 18 NHS Oxfordshire. INTERIM Treatment Threshold Statement: Knee arthroplasty (Reference PS 188). 2010 [updated 2010; cited 5th April 2011]; Available from: <http://www.oxfordshirepct.nhs.uk/professional-resources/priority-setting/lavender-statements/documents/PS188Kneereplacement.pdf>.
- 19 NHS Gloucestershire. Guidelines for Management of Orthopaedic Conditions. 2010 [updated 2010; cited 5th April 2011]; Available from: http://www.glospct.nhs.uk/pdf/services/orthopaedic_referral_guidelines_%20march%2010v3.pdf.

Formatted: Portuguese (Portugal)

Formatted: Font: Bold, Portuguese (Portugal)

Formatted: Portuguese (Portugal)

Formatted: Font: (Default) Arial, 11 pt

Formatted: Portuguese (Portugal)

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: (Default) Arial, 11 pt

Formatted: Right: 0.25"

20 NHS Worcestershire. [Commissioning policy for musculoskeletal surgical interventions](#). 2010 [updated 2010; cited 5th April 2011]; Available from: http://www.worcestershire.nhs.uk/file_download.aspx?id=86bae3e2-4d88-4416-92e6-9eeda22e8208.

Formatted: Font: (Default) Arial, 11 pt

21 Milton Keynes O, Berkshire East, Berkshire West & Buckinghamshire (MOBBB) Priorities Committee. Minutes of meeting held on Wednesday 15 December 2010. 2010 [updated 2010; cited 5th April 2011]; Available from: <http://www.sph.nhs.uk/sph-psu/sph-files/mobbb-minutes/mobbb-minutes-15-december-2010>.

Formatted: Font: (Default) Arial, 11 pt

22 South West London Public Health Network. 2010/11 South West London Effective Commissioning Initiative. 2010 [updated 2010; cited 5th April 2011]; Available from: <http://www.swlphnetwork.org.uk/uploads/documents/effective-commissioning-document-master-version-2010-11-final.doc>.

Formatted: Font: (Default) Arial, 11 pt

23 NHS Manchester. Topics for inclusion or change in the pct non-commissioned procedures list. 2010 [updated 2010; cited 5th April 2011]; Available from: http://www.manchester.nhs.uk/document_uploads/19%20January%202011/C2turnaroundAppendix%20%20TR.doc.

Formatted: Font: (Default) Arial, 11 pt

24 NHS Derbyshire County. NHS Derbyshire County Prior Approval: Procedures of Limited Clinical Value Summary. 2010 [updated 2010; cited 29th June 2011]; Available from: <http://www.phcn.nhs.uk/Derbyshire%20County%20Procedures%20of%20Limited%20Clinical%20Value%20100525%20draft.pdf>.

Formatted: Font: (Default) Arial, 11 pt

25 Jeffreys B. Surgeons raise alarm over waiting. Journal [serial on the Internet]. 2011 Date: Available from: <http://www.bbc.co.uk/news/health-12964360>.

Formatted: Font: (Default) Arial, 11 pt

26 NHS Warwickshire. Report to the PCT board 10th November 2010. 2010 [updated 2010 6th April 2011; cited 29th June 2011]; Available from: <http://www.warwickshire.nhs.uk/trustboardenclosures/November2010/Enc%208%20%20In%20Year%20Demand%20Management%20of%20Electives%205-11-2010.pdf>.

Formatted: Font: (Default) Arial, 11 pt

27 Hospital Episode Statistics. Main procedures and interventions: 4 character inpatient data 2009-10. 2010 [updated 2010; cited 10th February 2011]; Available from:

<http://www.hesonline.nhs.uk/Ease/servlet/ContentServer?siteID=1937&categoryID=215>.

Formatted: Font: (Default) Arial, 11 pt

28 Dawson J, Fitzpatrick R, Murray D, et al. Questionnaire on the perceptions of patients about total knee replacement. *J Bone Joint Surg Br* 1998;**80**:63-9. doi:

Formatted: Font: Bold

29 Johnston L, MacLennan G, McCormack K, et al. The Knee Arthroplasty Trial (KAT) design features, baseline characteristics, and two-year functional outcomes

Formatted: Right: 0.25"

after alternative approaches to knee replacement. *J Bone Joint Surg Am* 2009;**91**:134-41. doi: 91/1/134 [pii] 10.2106/JBJS.G.01074

30 Breeman S, Campbell M, Dakin H, et al. Patellar Resurfacing in Total Knee Replacement: Five-Year Clinical and Economic Results of a Large Randomized Controlled Trial. *J Bone Joint Surg Am* 2011;**93**:1473-81.

31 Dolan P. Modeling valuations for EuroQol health states. *Med Care* 1997;**35**:1095-108.

32 HM Treasury. *The Green Book: Appraisal and Evaluation in Central Government*. London: TSO: 2003.

33 Hawker GA, Badley EM, Croxford R, et al. A population-based nested case-control study of the costs of hip and knee replacement surgery. *Med Care* 2009;**47**:732-41. doi: 10.1097/MLR.0b013e3181934553

34 Kind P, Dolan P, Gudex C, et al. Variations in population health status: results from a United Kingdom national questionnaire survey. *Bmj* 1998;**316**:736-41. doi: 10.1136/bmj.316.7157.736

35 Lie SA, Pratt N, Ryan P, et al. Duration of the increase in early postoperative mortality after elective hip and knee replacement. *J Bone Joint Surg Am* 2010;**92**:58-63. doi: 92/1/58 [pii] 10.2106/JBJS.H.01882

36 Royston P. Multiple imputation of missing values. *Stata Journal* 2004;**4**:227-41.

37 Royston P. Multiple imputation of missing values: further update of ice, with an emphasis on interval censoring. *Stata Journal* 2007;**7**:445-64.

38 White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011;**30**:377-99. doi: 10.1002/sim.4067

39 Briggs A, Clark T, Wolstenholme J, et al. Missing... presumed at random: cost-analysis of incomplete data. *Health Econ* 2003;**12**:377-92.

40 Yu LM, Burton A, Rivero-Arias O. Evaluation of software for multiple imputation of semi-continuous data. *Stat Methods Med Res* 2007;**16**:243-58. doi: 16/3/243 [pii] 10.1177/0962280206074464

41 Murray DW, Fitzpatrick R, Rogers K, et al. The use of the Oxford hip and knee scores. *J Bone Joint Surg Br* 2007;**89**:1010-4. doi: 89-B/8/1010 [pii] 10.1302/0301-620X.89B8.19424

42 Fenwick E, Claxton K, Sculpher M. Representing uncertainty: the role of cost-effectiveness acceptability curves. *Health Econ* 2001;**10**:779-87.

43 British Medical Association. *British National Formulary 61: Cardiovascular risk prediction charts for primary prevention*. March 2011 [updated March 2011; cited 12th April 2011]; Available from: <http://bnf.org/bnf/bnf/current/204016.htm>.

44 Barrett J, Losina E, Baron JA, et al. Survival following total hip replacement. *J Bone Joint Surg Am* 2005;**87**:1965-71. doi: 87/9/1965 [pii] 10.2106/JBJS.D.02440

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: French (France)

Formatted: Font: Bold, French (France)

Formatted: French (France)

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: Bold

Formatted: French (France)

Formatted: Font: Bold, French (France)

Formatted: French (France)

Formatted: Font: Bold

Formatted: Font: (Default) Arial, 11 pt

Formatted: Font: (Default) Arial, 11 pt

Formatted: French (France)

Formatted: Font: Bold, French (France)

Formatted: French (France)

Formatted: Font: (Default) Arial, 11 pt

Formatted: French (France)

Formatted: Right: 0.25"

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[45 Lie SA, Engesaeter LB, Havelin LI, et al. Mortality after total hip replacement: 0-10-year follow-up of 39,543 patients in the Norwegian Arthroplasty Register. Acta Orthop Scand 2000;71:19-27. doi: 10.1080/00016470052943838](#)

[46 Dakin HA, Devlin NJ, Odeyemi IA. "Yes", "No" or "Yes, but"? Multinomial modelling of NICE decision-making. Health Policy 2006;77:352-67.](#)

[47 Devlin N, Parkin D. Does NICE have a cost-effectiveness threshold and what other factors influence its decisions? A binary choice analysis. Health Econ 2004;13:437-52.](#)

[48 National Institute for Health and Clinical Excellence. Supplementary guidance on appraising life-extending, end of life treatments. July 2009 \[updated July 2009; cited 23rd February 2011\]; Available from: <http://www.nice.org.uk/media/E4A/79/SupplementaryAdviceTACEoL.pdf>.](#)

[49 Coombes R. Rationing of joint replacements raises fears of further cuts. BMJ 2005;331:1290. doi: 331/7528/1290 \[pii\].10.1136/bmj.331.7528.1290](#)

Formatted: Font: Bold

Formatted ... [11]

Formatted: Font: (Default) Arial, 11 pt

Formatted ... [12]

Formatted: Indent: Left: 0", Hanging: 0.5"

Deleted: 1 Hospital Episode Statistics. Provisional Monthly Patient Reported Outcome Measures (PROMs) in England. April 2009 – April 2010: Pre- and postoperative data: Experimental statistics. 2010 [updated 2010; cited 6th April 2011]; Available from: <http://www.hqip.org.uk/assets/NCAPOP-Library/PROMsreport09101.pdf>.

2 Labek G, Thaler M, Janda W, et al. Revision rates after total joint replacement: cumulative results from worldwide joint register datasets. J Bone Joint Surg Br 2011;93:293-7.

3 The National Collaborating Centre for Chronic Conditions. Osteoarthritis: National clinical guideline for care and management in adults. London: Royal College of Physicians; 2008 [updated 2008; cited 29th June 2011]; Available from: <http://www.nice.org.uk/nicemedia/live/11926/39720/39720.pdf>.

4 Huddleston JI, Maloney WJ, Wang Y, et al. Adverse events after total knee arthroplasty: a national Medicare study. J Arthroplasty 2009;24:95-100.

5 Memtsoudis SG, Della Valle AG, Besculides MC, et al. Risk factors for perioperative mortality after lower extremity arthroplasty: a population-based study of 6,901,324 patient discharges. J Arthroplasty 2010;25:19-26.

6 Navarro Espigares JL, Hernandez Torres E. Cost-outcome analysis of joint replacement: evidence from a Spanish public hospital. Gac Sanit 2008;22:337-43.

7 Losina E, Walensky RP, Kessler CL, et al. Cost-effectiveness of total knee arthroplasty in the United States: patient risk and hospital volume. Arch Intern Med 2009;169:1113-21; discussion 21-2.

8 Rasanen P, Paavolainen P, Sintonen H, et al. Effectiveness of hip or knee replacement surgery in terms of quality-adjusted life years and costs. Acta Orthop 2007;78:108-15.

9 Rissanen P, Aro S, Sintonen H, et al. Costs and cost-effectiveness in hip and knee replacements. A prospective study. Int J Technol Assess Health Care 1997;13:575-88.

10 Lavernia CJ, Guzman JF, Gachupin-Garcia A. Cost effectiveness and quality of life in knee arthroplasty. Clin Orthop Relat Res 1997:134-9.

11 National Institute for Health and Clinical Excellence. Social value judgements: Principles for the development of NICE guidance. Second Edition. 2008 [updated 2008; cited 17th November 2010]; Available from: <http://www.nice.org.uk/media/C18/30/SVJ2PUBLICATION2008.pdf>.

12 Brander VA, Malhotra S, Jet J, et al. Outcome of hip and knee arthroplasty in persons aged 80 years and older. Clin Orthop Relat Res 1997:67-78.

13 Karupiah SV, Banaszkiwicz PA, Ledingham WM. The mortality, morbidity and cost benefits of elective total knee arthroplasty. J Bone Joint Surg Br 2004;86:108-12.

Formatted ... [13]

Formatted ... [14]

Formatted: Right: 0.25"

BMJ Open: first published as 10.1136/bmjopen-2011-000332 on 30 January 2012. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2021 by guest. Protected by copyright.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Figure 1: EQ-5D utility by subgroup

For peer review only

Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy, 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life); OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])^[41]; TKR, total knee replacement.
Error bars show 95% confidence intervals. EQ-5D utility is measured on a scale from 1 (perfect health) to -0.594, where 0 is equivalent to death.

Formatted: English (U.S.)

Deleted: , 4=symptomatic illness with severe restriction on life

Field Code Changed

Deleted: [40]

Formatted: Right: 0.25"

Figure 2: Cost-effectiveness prediction charts estimated based on predictions of regression models.

Abbreviations: ASA, American Society of Anesthesiologists classification (1=completely fit and healthy, 2=some illness but no effect on daily activity, 3=symptomatic illness with minimal restriction on life); OKS, Oxford Knee Score (new scoring system running from zero [severe problems on all functions] to 48 [no problems])^[41]; QALY, quality-adjusted life-year; TKR, total knee replacement.

For peer review only

Deleted: [40]

Field Code Changed

Formatted: Right: 0.25"

BMJ Open: first published as 10.1136/bmjopen-2011-000332 on 30 January 2012. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2021 by guest. Protected by copyright.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 3: Cost-effectiveness acceptability curves for different patient subgroups at different baseline Oxford Knee Scores (OKS). A: American Society of Anesthesiologists (ASA) grade 1 and 2 patients. B: ASA grade 3 patients.

For peer review only

Formatted: Right: 0.25"

BMJ Open: first published as 10.1136/bmjopen-2011-000332 on 30 January 2012. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2021 by guest. Protected by copyright.

1 12/20/2011 9:58:00 AM

4 12/20/2011 9:58:00 AM

7 12/20/2011 9:58:00 AM

10 12/20/2011 9:58:00 AM

13 12/15/2011 3:33:00 PM

ported Outcome Measures
operative data: Experimental
e from:
[rt06101.pdf](#).

For peer review only

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

BMJ Open: first published as 10.1136/bmjopen-2011-000332 on 30 January 2012. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2021 by guest. Protected by copyright.

1 effectiveness and quality of life in knee

2
3 value judgements: Principles for
4
5 updated 2008; cited 17th November
6
7 [SVJ2PUBLICATION2008.pdf](#).

8 knee arthroplasty in persons aged

9
10
11 mortality, morbidity and cost

12
13 population. Int Orthop

14
15
16 Obesity on quality-of-life

17
18 (Mead NJ) 2007;36:E117-20, E27.

19 primary total knee replacement Knee

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

BMJ Open: first published as 10.1136/bmjopen-2011-000332 on 30 January 2012. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2021 by guest. Protected by copyright.

2011]; Available from:

[ve-commissioning-document-](#)

1
2
3 Non-commissioned procedures
4
5
6
7 [nary%202011/C2turnaroundAppe](#)

9 approval: Procedures of Limited

10 2011]; Available from:

11 [es%20of%20Limited%20Clinical%2](#)

16 on the Internet]. 2011 Date:

19 ber 2010. 2010 [updated 2010 6th

For peer review only

BMJ Open: first published as 10.1136/bmjopen-2011-000332 on 30 January 2012. Downloaded from <http://bmjopen.bmj.com/> on May 13, 2021 by guest. Protected by copyright.

early postoperative mortality after

10;92:58-63.

Journal 2004;4:227-41.

update of ice, with an emphasis on

g chained equations: Issues and

med at random: cost-analysis of

for multiple imputation of semi-

Oxford hip and knee scores. J

For peer review only

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

onal study. Eur J Health Econ

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ompartmental instead of
in the knee is a cost-effective
e compared with 10,624 primary
e and 1999;**70**:170-5.

s of unicompartmental and total
decision analysis. J Bone Joint Surg

12/20/2011 9:58:00 AM

12/20/2011 9:58:00 AM

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Health economic checklist for “Rationing of total knee replacement: A cost-effectiveness analysis on a large trial dataset”

Study design	Page
(1) The research question is stated	5, paragraphs 2 & 3
(2) The economic importance of the research question is stated	4-5
(3) The viewpoint(s) of the analysis are clearly stated and justified	5, paragraph 3
(4) The rationale for choosing the alternative programmes or interventions compared is stated	4-5
(5) The alternatives being compared are clearly described	5, paragraph 3
(6) The form of economic evaluation used is stated	5, paragraph 3
(7) The choice of form of economic evaluation is justified in relation to the questions addressed	5, paragraph 3
Data collection	
(8) The source(s) of effectiveness estimates used are stated	5, paragraph 4
(9) Details of the design and results of effectiveness study are given (if based on a single study)	5, paragraph 4
(10) Details of the method of synthesis or meta-analysis of estimates are given (if based on an overview of a number of effectiveness studies)	N/A
(11) The primary outcome measure(s) for the economic evaluation are clearly stated	5, paragraph 3
(12) Methods to value health states and other benefits are stated	6, paragraph 1
(13) Details of the subjects from whom valuations were obtained are given	6, paragraph 1
(14) Productivity changes (if included) are reported separately	N/A
(15) The relevance of productivity changes to the study question is discussed	5, paragraph 3
(16) Quantities of resources are reported separately from their unit costs	Given in reference 29
(17) Methods for the estimation of quantities and unit costs are described	Given in reference 29
(18) Currency and price data are recorded	N/A
(19) Details of currency of price adjustments for inflation or currency conversion are given	Given in reference 29
(20) Details of any model used are given	N/A
(21) The choice of model used and the key parameters on which it is based are justified	N/A
Analysis and interpretation of results	
(22) Time horizon of costs and benefits is stated	5, paragraph 4
(23) The discount rate(s) is stated	6, paragraph 2
(24) The choice of rate(s) is justified	6, paragraph 2
(25) An explanation is given if costs or benefits are not discounted	N/A
(26) Details of statistical tests and confidence intervals are given for stochastic data	8, paragraph 1
(27) The approach to sensitivity analysis is given	11, paragraph 4
(28) The choice of variables for sensitivity analysis is justified	11, paragraph 4
(29) The ranges over which the variables are varied are stated	N/A
(30) Relevant alternatives are compared	5, paragraph 3
(31) Incremental analysis is reported	Tables 1 & 2
(32) Major outcomes are presented in a disaggregated as well as aggregated form	Tables 1 & 2
(33) The answer to the study question is given	12, paragraph 2 & page 13 paragraphs 2-3
(34) Conclusions follow from the data reported	5, 12-13
(35) Conclusions are accompanied by the appropriate caveats	12-13

Authors may enter N/A if an item on the checklist is not appropriate, but this is only acceptable for items 9,10, 12-15, 20, 21, 23-29, and 31.