



Sedentary behavior and life expectancy in the United States: A cause-deleted life table analysis

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2012-000828
Article Type:	Research
Date Submitted by the Author:	09-Jan-2012
Complete List of Authors:	Katzmarzyk, Peter; Pennington Biomedical Research Center, Lee, I-Min; Harvard Medical School
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology
Keywords:	PREVENTIVE MEDICINE, physical activity, lifestyle

SCHOLARONE™
Manuscripts

Peer Review Only

1
2
3
4 **Sedentary behavior and life expectancy in the United States: A cause-deleted life table**
5 **analysis**
6

7 Peter T. Katzmarzyk, I-Min Lee
8
9

10
11
12 Pennington Biomedical Research Center, Louisiana State University System, Baton Rouge, LA,
13 70808 Peter T. Katzmarzyk professor
14

15 Division of Preventive Medicine, Brigham & Women's Hospital and Harvard Medical School,
16 Boston, MA, 02215 I-Min Lee associate professor
17
18

19 Correspondence to: peter.katzmarzyk@pbrc.edu
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives: To determine the impact of sitting and television viewing on life expectancy in the United States.

Design: Prevalence-based cause-deleted life table analysis.

Setting: Summary relative risks of all-cause mortality associated with sitting and television viewing were obtained from a meta-analysis of available prospective cohort studies.

Prevalences of sitting and television viewing were obtained from the U.S. National Health and Nutrition Examination Survey.

Primary Outcome Measure: Life expectancy at birth.

Results: The estimated gains in life expectancy in the US population were 1.91 years for reducing excessive sitting to less than three hours per day and a gain of 1.38 years from reducing excessive television viewing to less than 2 hours per day. The lower and upper limits from a sensitivity analysis which involved simultaneously varying the estimates of RR (using the upper and lower bounds of the 95% CI) and the prevalence of television viewing ($\pm 20\%$) were 0.74 and 3.54 years for sitting and 0.48 and 2.50 years for television viewing, respectively.

Conclusions: Reducing sedentary behaviors such as sitting and television viewing has the potential to increase life expectancy in the United States.

Subject Headings: public health; epidemiology

Keywords:

Article Summary

Article focus

- This paper presents the results of an analysis aimed at determining the effects of sedentary behavior on life expectancy in the United States

Key messages

- The analyses indicate that population life expectancy in the United States would be 1.91 years higher if adults reduced their time spent sitting to less than three hours per day, and 1.38 years higher if they reduced television viewing to less than 2 hours per day.

Strengths and limitations

- The use of the well-accepted prevalence-based methodology to estimate the public health burden of sedentary behavior is a marked strength, which allows for comparability with the effects of other established risk factors.
- This study relied on self-reported engagement in sedentary behaviors rather than an objective measurement, which is a limitation.

1
2
3 Over 60 years of research on physical activity and health culminated in the release of the
4 *2008 Physical Activity Guidelines for Americans*, which recommend that adults should
5 accumulate at least 150 minutes of moderate-to-vigorous physical activity per week to attain the
6 health benefits associated with physical activity.¹ Recently, there has been considerable interest
7 in understanding the role of sedentary behaviors on health, independent of overall physical
8 activity levels,²⁻⁴ since one can be both sedentary and physically active (e.g., an office worker
9 who sits most of his work hours, but who also jogs regularly).

10
11 Sedentary behaviors which involve sitting for extended periods are ubiquitous in modern
12 society. Based on self-reports, a recent survey of 20 countries documented a median of 300
13 minutes per day spent sitting, ranging from ≤ 180 minutes per day in Portugal, Brazil and
14 Colombia to ≥ 360 minutes per day in Taiwan, Norway, Hong Kong, Saudi Arabia and Japan.⁵
15 Several studies have demonstrated positive associations between sedentary behaviors
16 including sitting and television viewing and health outcomes such as type 2 diabetes,
17 cardiovascular disease mortality and all-cause mortality.⁶ Thus, excessive time spent in
18 sedentary behavior is undoubtedly having an impact on public health. A recent study from
19 Australia estimated that television viewing reduced life expectancy at birth by 1.8 years in men
20 and 1.5 years in women.⁷ The purpose of this study was to determine the impact of sedentary
21 behaviors on life expectancy in the United States.

22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Methods

A prevalence-based approach was used to estimate the impact of sedentary behavior on life expectancy. The population-attributable fraction (PAF) was computed from the prevalences of sedentary behaviors (defined here as sitting and television viewing) and the relative risks (RR) of all-cause mortality associated with these behaviors. The PAF equation used was $\sum P_i(RR_i - 1/RR_i)$, where P is the prevalence of the risk factor among cases in stratum i (i=1 to 3 in this study; details are provided below). This equation produces internally valid estimates when confounding exists and adjusted RRs must be used.⁸

Relative Risks Associated with Sedentary Behaviors

Sedentary behavior can be captured globally using questionnaires that address total daily sitting time, or time spent in specific sedentary behaviors like television viewing, reading, or computer use. Published studies on the associations between sitting or television viewing and all-cause mortality were identified through a literature search of MEDLINE. We pooled RR estimates from each study separately for sitting or television viewing and all cause-mortality using a random effects meta-analysis. Pooled RR estimates were obtained for two levels of sitting and two levels of television viewing relative to a referent group in each case. In order to maintain consistency across studies, the age- and sex-adjusted RR estimates were used from each study. In cases where the authors presented only multivariable-adjusted RR estimates, or used different exposure categories, we contacted them and asked them to provide this information. All authors contacted complied with our requests. MIX 2.0 software was used to conduct the meta-analysis.⁹

Prevalence of Sedentary Behavior

The prevalences of time spent sitting and television viewing were obtained from the National Health and Nutrition Examination Survey (NHANES). NHANES uses a complex, multistage, probability sampling design to select participants who are representative of the civilian, noninstitutionalized U.S. population.¹⁰ The prevalence of sitting (<3 h, 3-5.9 h, and ≥6 h/day) among non-pregnant adults 18 years of age and older was determined from responses to the following question in the 2007-08 NHANES: "How much time do you usually spend sitting or reclining on a typical day?" The prevalence of television viewing (<2 h, 2-3.9 h, and ≥4 h/day) among non-pregnant adults 18 years of age and older was determined from responses to the following question in the 2005-06 NHANES: "Over the past 30 days, on average how many hours per day did you sit and watch TV or videos?" Data analysis followed the guidelines of the National Center for Health Statistics for analysis of NHANES data due to the complex sampling design and methods.¹¹

1
2
3 The PAF equation used in this study requires the prevalence of sedentary behavior among
4 cases (i.e. decedents) rather than from the source population (i.e. NHANES prevalence).
5
6 Therefore, each population prevalences obtained from NHANES was adjusted by using the
7
8 average case:source prevalence ratio (i.e. the prevalences among cases divided by the
9
10 prevalences among the baseline source population from the prospective cohort studies).
11
12

13 Gains in Life Expectancy

14
15 The PAFs for all-cause mortality associated with sitting and television viewing were
16
17 computed using the summary RR estimates obtained from the meta-analyses and the adjusted
18
19 prevalences from NHANES, as described above. Potential gains in life expectancy attributable
20
21 to reducing sedentary behaviors were estimated using a cause-deleted life-table analysis, which
22
23 estimates years of life gained at birth if a specific cause of death is eliminated. The most current
24
25 abridged life table for the United States (2009) was downloaded from the World Health
26
27 Organization website.¹² The PAFs computed for sedentary behavior were used to reduce the
28
29 mortality rates in the life table for adults aged 18 years and older, and the life expectancy at
30
31 birth was recalculated from the new mortality rates. The difference between the current life
32
33 expectancy and the cause-deleted life expectancy represents the estimated gain in life
34
35 expectancy from reducing the prevalence of sedentary behavior.
36
37
38
39

40 Sensitivity Analysis

41
42 A sensitivity analysis was used to estimate the effects of simultaneously varying the
43
44 estimates of RR (using the upper and lower bounds of the 95% CI) and the prevalences of
45
46 sitting and television viewing ($\pm 20\%$).
47
48

49 **Results**

50
51 The characteristics and main results of the prospective cohort studies included in the meta-
52
53 analyses of sedentary behavior and mortality are presented in Table 1. Two studies were
54
55 included that investigated the association between sitting and all-cause mortality.¹³⁻¹⁴ The
56
57 exposure levels of sitting in the first study were (1) none/ $\frac{1}{4}$ of the time, (2) $\frac{1}{2}$ the time, and (3) $\frac{3}{4}$
58
59
60

1
2
3 of the time/all of the time based on activities that participants did most days of the week,¹³
4
5 whereas the levels in the second study were (1) <3 h, (2) 3-5 h, and (3) ≥6 h of leisure-time
6
7 sitting.¹⁴ Three studies were included that studied the association between television viewing
8
9 and all-cause mortality.¹⁵⁻¹⁷ The levels of television viewing in two of the studies were (1) <2 h,
10
11 (2) 2-3.9 h, and (3) ≥4 h,¹⁵⁻¹⁶ whereas the levels in one study were (1) <2.5 h, (2) 2.5-3.6 h, and
12
13 (3) >3.6 h.¹⁷
14
15

16
17 Figure 1 presents the results of the meta-analysis for sitting and all-cause mortality. The
18
19 pooled relative risks were 1.18 (1.14 - 1.21) and 1.45 (1.39-1.51) for levels 2 and 3 versus level
20
21 1, respectively. Figure 2 presents the results for the meta-analysis for television viewing and all-
22
23 cause mortality. The pooled relative risks were 1.17 (1.04 - 1.32) and 1.49 (1.22-1.82) for levels
24
25 2 and 3 versus level 1, respectively.
26

27
28 The prevalences of sitting and television viewing in cases (decedents) and in the source
29
30 population at baseline in the prospective studies are presented in Table 2, along with the
31
32 average case:source prevalence ratio. These ratios were applied to the population prevalences
33
34 obtained from NHANES in order to estimate the prevalences among cases in the population,
35
36 which are required for inclusion in the PAF calculations. Figure 3 presents the weighted
37
38 population prevalences of sitting and television viewing in NHANES, along with the prevalences
39
40 after adjustment for the average case:source prevalence ratios from Table 2.
41

42
43 The results of the life table analyses indicate a gain in life expectancy from reducing the
44
45 prevalence of sedentary behavior from levels 2 or 3 to level 1 results in a gain of 1.91 years for
46
47 reducing sitting prevalence and a gain of 1.38 years from reducing television viewing
48
49 prevalence. The lower and upper limits from the sensitivity analyses were 0.74 and 3.54 years
50
51 for sitting and 0.48 and 2.50 years for television viewing, respectively.
52

53 Discussion

54
55 The results of this study indicate that limiting sitting to less than three hours per day and
56
57 limiting television viewing to less than two hours per day may increase life expectancy at birth in
58
59
60

1
2
3 the United States by approximately 1.9 and 1.4 years, respectively, assuming a causal
4 relationship. The PAF provides a *theoretical* estimate of the effects of a risk factor on an
5 outcome at the population level, in this case, all-cause mortality. The results indicate that
6 sedentary behaviors are accounting for between 1.4 and 1.9 years of life expectancy at birth.
7
8 This should not be interpreted to mean that people who are more sedentary can expect to live
9 1.4 or 1.9 years less than someone who does not engage in these behaviors as much. Life
10 expectancy is a population statistic and it does not apply to individuals. A recent meta-analysis
11 of television viewing and all-cause mortality estimated that the relative risk of all-cause mortality
12 was 1.13 (95% CI: 1.07-1.18) per 2 hours of daily television viewing, which corresponded with
13 104 deaths per 100,000 people in the United States.⁴

14
15
16 This study has several strengths and limitations that warrant discussion. The use of the well-
17 accepted prevalence-based PAF methodology to estimate the public health burden of sedentary
18 behavior is a marked strength, which allows for comparability with the effects of other
19 established risk factors. However, the PAF provides a *theoretical* estimate of the effects of risk
20 factor on a health outcome, and further research is required using an incidence-based
21 approach. Our analysis assumes that there is a causal relationship between sedentary behavior
22 and mortality. While studies using randomized designs are not possible, further observational
23 studies, which control for confounding, will add to the evidence for causation. Another major
24 strength of this study is the use of representative population data from NHANES to quantify the
25 exposure of the population to sedentary behaviors. This study relied on self-reported
26 engagement in sedentary behaviors. Future studies should attempt to better quantify sedentary
27 behavior using objective activity monitors; however, the estimates of sitting reported in NHANES
28 are similar to those obtained for the U.S. in a study of 20 countries⁵ which indicates some face
29 validity to the results. For television viewing, the categories of exposure reported in the cohort
30 studies (< 2 h, 2-3.9 h, ≥4 h) matched the categories reported in NHANES; however, the studies
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 on sitting used different exposure categories (i.e. total sitting time versus leisure-time sitting
4 only), so assumptions had to be made when estimating the exposure levels in NHANES.
5
6

7
8 The results of several recent studies have suggested that the effects of sedentary behavior
9 on health may be independent of the effects of physical activity *per se*.^{2,18} It has been estimated
10 that a lack of leisure-time physical activity accounts for approximately 0.9 years of life
11 expectancy at birth in Canada.¹⁹ Given that the studies used to derive the summary RR
12 estimates for the current study in many cases included leisure-time physical activity as a
13 covariate in multivariable-adjusted models and this did not appreciably change the estimates of
14 RR for sedentary behavior, the estimates of the effects on life expectancy may also be
15 considered independent. Current life expectancy in the United States (2009) is 78.5 years.¹²
16
17 The effects of sedentary behavior on life expectancy reported in this study are on a similar order
18 of magnitude as other chronic disease risk factors. For example, it has been estimated that
19 obesity accounts for between 0.30 and 1.08 years of population life expectancy at birth in the
20 United States, depending on gender, ethnicity, and severity of obesity.²⁰ A more recent study
21 has estimated that the current distribution of body mass index, compared to an optimal
22 distribution (a mean of 21 kg/m²), accounts for 1.3 years of current life expectancy at birth in
23 both males and females in the United States.²¹ Results from the same study indicate that
24 smoking is also associated with 2.5 years and 1.8 years of life expectancy at birth in males and
25 females, respectively.²¹
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 Sitting time is a global measure of sedentary behavior, whereas television viewing is
45 somewhat more specific. A recent review found that domain-specific behaviors such as
46 television viewing are recalled with more reliability than global measures of sitting and sedentary
47 behavior.²² The degree to which differences in reliability between the measures used in this
48 study may have affected the estimates is not known. There is some evidence to suggest that
49 using a single global question to measure sitting produces lower estimates than more detailed,
50 domain-specific questions.²²⁻²³ Thus, it is likely that the prevalences of higher levels of sitting
51
52
53
54
55
56
57
58
59
60

1
2
3 reported in this study from NHANES are likely conservative. Using objective monitoring
4
5 (accelerometry) in the 2003-2003 NHANES, Matthews et al. reported that U.S. adults spend
6
7 approximately 7.7 hours per day engaged in sedentary behavior.²⁴ There are several potential
8
9 mechanisms that could explain the association between sedentary behavior and all-cause
10
11 mortality rates. Sedentary behavior is associated with an increased risk of the development of
12
13 chronic conditions such type 2 diabetes and cardiovascular disease.⁶ Further, human and
14
15 animal studies indicate that sedentary behavior is associated with elevated cardiometabolic
16
17 biomarkers and a poor risk factor profile.²⁵⁻²⁶ For example, hind limb suspension (unloading) in
18
19 rats results in marked immediate decreases in lipoprotein lipase activity, triglyceride uptake into
20
21 red skeletal muscle, and reductions in the concentration of HDL cholesterol.²⁷ Future
22
23 intervention research is required to determine the causal pathways between sedentary behavior
24
25 and health outcomes that have the potential to impact mortality rates.
26
27

28
29 In conclusion, the results of this study indicate that extended sitting time and television
30
31 viewing are having a meaningful impact on life expectancy in the United States. Further
32
33 research using intervention designs is required to determine the effects of reducing sedentary
34
35 behavior on health outcomes and to make recommendations regarding the safe levels of
36
37 sedentary behavior for the population.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends

Figure 1. Risk of all-cause Mortality Associated with Sitting. Level 3 corresponds to $\frac{3}{4}$ of all of the time for Katzmarzyk et al. 2009 and ≥ 6 h for Patel et al. 2010. Level 2 corresponds to $\frac{1}{2}$ of the time for Katzmarzyk et al. 2009 and 3-5 h for Patel et al. 2010. Level 1 corresponds to $\frac{1}{4}$ of the time for Katzmarzyk et al. 2009 and < 3 h for Patel et al. 2010.

Figure 2. Risk of all-cause Mortality Associated with Television Viewing. Level 3 corresponds to ≥ 4 h for Dunstan et al. 2010 and Stamatakis et al. 2011 and ≥ 3.6 h for Wijndaele et al. 2011. Level 2 corresponds to 2-3.9 h for Dunstan et al. 2010 and Stamatakis et al. 2011 and 2.5-3.6 h for Wijndaele et al. 2011. Level 1 corresponds to < 2 h for Dunstan et al. 2010 and Stamatakis et al. 2011 and < 2.4 h for Wijndaele et al. 2011.

Figure 3. Prevalences of A) sitting and B) television viewing in the U.S. National Health and Nutrition Examination Survey. *Adjusted prevalences of sitting and television viewing using the average case/source prevalence ratio obtained from cohort studies of sedentary behavior and all-cause mortality.

1
2
3 **Funding:** This research received no specific grant from any funding agency in the public,
4 commercial or non-profit sectors.
5
6

7 **Competing Interests:** All authors have completed the Unified Competing Interest form at
8 www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and
9
10 declare that both authors have no financial or non-financial interests that may be relevant to the
11
12 submitted work.
13
14

15 **Data Sharing Statement:** There are no additional data available.
16
17

18 **Author Contributions:** Both authors were responsible study conception and design. PTK was
19 responsible for data analysis. Both authors drafted and critically revised the manuscript, and
20
21 approved the final version to be published. PTK is the guarantor for the study.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

er review only

References

1. US Department of Health and Human Services. *2008 Physical Activity Guidelines for Americans*: US Department of Health and Human Services, Centers for Disease Control and Prevention, 2008.
2. Katzmarzyk PT. Physical activity, sedentary behavior, and health: paradigm paralysis or paradigm shift? *Diabetes* 2010;59(11):2717-25.
3. Thorp AA, Owen N, Neuhaus M, et al. Sedentary behaviors and subsequent health outcomes in adults a systematic review of longitudinal studies, 1996-2011. *Am J Prev Med* 2011;41(2):207-15.
4. Grontved A, Hu FB. Television viewing and risk of type 2 diabetes, cardiovascular disease, and all-cause mortality: a meta-analysis. *JAMA* 2011;305(23):2448-55.
5. Bauman A, Ainsworth BE, Sallis JF, et al. The descriptive epidemiology of sitting: A 20-country comparison using the International Physical Activity Questionnaire (IPAQ). *Am J Prev Med* 2011;41(2):228-35.
6. Proper KI, Singh AS, van Mechelen W, et al. Sedentary behaviors and health outcomes among adults: a systematic review of prospective studies. *Am J Prev Med* 2011;40(2):174-82.
7. Veerman JL, Healy GN, Cobiac LJ, et al. Television viewing time and reduced life expectancy: a life table analysis. *Br J Sports Med* 2011.
8. Rockhill B, Newman B, Weinberg C. Use and misuse of population attributable fractions. *Am J Public Health* 1998;88(1):15-9.
9. Bax L. Professional software for meta-analysis in Excel. Version 2.01.4. BostatXL; <http://www.meta-analysis-made-easy.com> 2011.
10. Centers for Disease Control and Prevention (CDC). National Health and Nutrition Examination Survey. Available at: <http://www.cdc.gov/nchs/nhanes.htm>. Access verified October 7, 2011. 2011.

- 1
2
3 11. Centers for Disease Control and Prevention (CDC), National Center for Health Statistics.
4
5 The National Health and Nutrition Examination Survey Analytic and Reporting Guidelines.
6
7 Available at: http://www.cdc.gov/nchs/nhanes/nhanes2003-2004/analytical_guidelines.htm
8
9 (Accessed October 7, 2011). 2006.
- 10
11 12. World Health Organization. Global Health Observatory Data Repository. Available at:
12
13 <http://apps.who.int/ghodata/> (Accessed October 7, 2011). 2011.
- 14
15 16. Katzmarzyk PT, Church TS, Craig CL, et al. Sitting time and mortality from all causes,
17
18 cardiovascular disease, and cancer. *Med Sci Sports Exerc* 2009;41:998-1005.
- 19
20 21. Patel AV, Bernstein L, Deka A, et al. Leisure time spent sitting in relation to total mortality in
22
23 a prospective cohort of US adults. *Am J Epidemiol* 2010;172(4):419-29.
- 24
25 26. Dunstan DW, Barr EL, Healy GN, et al. Television viewing time and mortality: the Australian
27
28 Diabetes, Obesity and Lifestyle Study (AusDiab). *Circulation* 2010;121(3):384-91.
- 29
30 31. Stamatakis E, Hamer M, Dunstan DW. Screen-based entertainment time, all-cause
32
33 mortality, and cardiovascular events population-based study with ongoing mortality and
34
35 hospital events follow-up. *J Am Coll Cardiol* 2011;57(3):292-9.
- 36
37 38. Wijndaele K, Brage S, Besson H, et al. Television viewing time independently predicts all-
38
39 cause and cardiovascular mortality: The EPIC Norfolk Study *Int J Epidemiol* 2011;40:150-59.
- 40
41 42. Tremblay MS, Colley RC, Saunders TJ, et al. Physiological and health implications of a
42
43 sedentary lifestyle. *Appl Physiol Nutr Metab* 2010;35(6):725-40.
- 44
45 46. Katzmarzyk PT. Physical inactivity and life expectancy in Canada. *J Phys Act Health*
46
47 2006;3:381-89.
- 48
49 50. Olshansky SJ, Passaro DJ, Hershow RC, et al. A potential decline in life expectancy in the
50
51 United States in the 21st century. *N Engl J Med* 2005;352:1138-45.
- 52
53 54. Danaei G, Rimm EB, Oza S, et al. The promise of prevention: the effects of four preventable
54
55 risk factors on national life expectancy and life expectancy disparities by race and county in
56
57 the United States. *PLoS medicine* 2010;7(3):e1000248.
- 58
59
60

- 1
2
3 22. Healy GN, Clark BK, Winkler EA, et al. Measurement of adults' sedentary time in population-
4 based studies. *Am J Prev Med* 2011;41(2):216-27.
5
6
7 23. Clemes SA, David BM, Zhao Y, et al. Validity of two self-report measures of sitting time. *J*
8 *Phys Act Health* 2011;In Press.
9
10 24. Matthews CE, Chen KY, Freedson PS, et al. Amount of time spent in sedentary behaviors in
11 the United States, 2003-2004. *Am J Epidemiol* 2008;167:875-81.
12
13 25. Healy GN, Matthews CE, Dunstan DW, et al. Sedentary time and cardio-metabolic
14 biomarkers in US adults: NHANES 2003-06. *Eur Heart J* 2011;32(5):590-7.
15
16 26. Hamilton MT, Hamilton DG, Zderic TW. Exercise physiology versus inactivity physiology: an
17 essential concept for understanding lipoprotein lipase regulation. *Exerc Sport Sci Rev*
18 2004;32(4):161-6.
19
20 27. Bey L, Hamilton MT. Suppression of skeletal muscle lipoprotein lipase activity during
21 physical inactivity: a molecular reason to maintain daily low-intensity activity. *The Journal of*
22 *physiology* 2003;551(Pt 2):673-82.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Characteristics and main results of the prospective cohort studies included in the meta-analysis.

Study	Participants	Age at Baseline (years)	Follow-up (years)	Person-years	Deaths (n)	Results			
Sitting Time									
Katzmarzyk et al. 2009 Canada Fitness Survey	N = 17,013 men and women	18 - 90	12	204,732	1,832	Deaths	<u>None/¼ Time</u>	<u>½ Time</u>	<u>¾ Time/All</u>
						Person-yr	826	542	464
						Crude RR*	117,965	52,346	34,421
						Age-Sex-HR**	1.00	1.48	1.93
							1.00	1.14	1.51
Patel et al. 2010 Cancer Prevention Study II Nutrition Cohort	N = 123,216 men and women	50 - 74	14	1,610,728	19,230	Deaths	<u><3 h</u>	<u>3-5 h</u>	<u>≥6 h</u>
						Person-yr	7,068	9,194	2,968
						Crude RR*	755,214	688,709	166,805
						Age-Sex-HR**	1.00	1.43	1.90
							1.00	1.18	1.44
TV Viewing									
Dunstan et al. 2010 Australian Diabetes, Obesity and Lifestyle Study (AusDiab)	N = 8,800 men and women	≥ 25	6.6	58,087	284	Deaths	<u><2 h</u>	<u>2-3.9</u>	<u>≥4 h</u>
						Person-yr	105	125	54
						Crude RR*	33,024	20,737	4326
						Age-Sex-HR**	1.00	1.89	3.93
							1.00	1.20	1.67
Stamatakis et al. 2011 Scottish Health Survey	N = 4,512 men and women	≥ 35	4.3	19,364	325	Deaths	<u><2 h</u>	<u>2-3.9 h</u>	<u>≥4 h</u>
						Person-yr	42	138	146
						Crude RR*	3,328	10,548	5,488
						Age-Sex-HR**	1.00	1.04	2.11
							1.00	1.13	1.77
Wijndaele et al. 2011 EPIC - Norfolk Study	N = 13,197 men and women	45 - 79	9.5	124,902	1,270	Deaths	<u><2.5 h</u>	<u>2.5-3.6 h</u>	<u>≥3.6 h</u>
						Person-yr	291	414	565
						Crude RR*	41,936	42,288	40,680
						Age-Sex-HR**	1.00	1.41	2.17
							1.00	1.17	1.32

*relative risk ratio; **age- and sex-adjusted hazard ratio.

Table 2. Prevalences of sitting and television viewing at baseline in the prospective cohort studies included in the meta-analysis.

Study	Prevalences			
Sitting Time				
Katzmarzyk et al. 2009		<u>None/¼ Time</u>	<u>½ Time</u>	<u>¾ Time/All</u>
Canada Fitness Survey	Source Prevalence (%)*	56.9	25.7	17.4
	Case Prevalence (%)**	45.1	29.6	25.3
	Case/Source Ratio	0.79	1.15	1.45
Patel et al. 2010		<u><3 h</u>	<u>3-5 h</u>	<u>≥6 h</u>
Cancer Prevention Study II	Source Prevalence (%)*	46.2	43.1	10.7
Nutrition Cohort	Case Prevalence (%)**	36.8	47.8	15.4
	Case/Source Ratio	0.80	1.11	1.44
Average Case/Source Ratio***		0.79	1.13	1.45
TV Viewing				
Dunstan et al. 2010		<u><2 h</u>	<u>2-3.9 h</u>	<u>≥4 h</u>
Australian Diabetes, Obesity and Lifestyle Study (AusDiab)	Source Prevalence (%)*	56.5	35.9	7.6
	Case Prevalence (%)**	37.0	44.0	19.0
	Case/Source Ratio	0.65	1.23	2.50
Stamatakis et al. 2011		<u><2 h</u>	<u>2-3.9 h</u>	<u>≥4 h</u>
Scottish Health Survey	Source Prevalence (%)*	17.1	54.1	28.8
	Case Prevalence (%)**	13.4	45.2	41.4
	Case/Source Ratio	0.78	0.84	1.44
Wijndaele et al. 2011		<u><2.5 h</u>	<u>2.5-3.6 h</u>	<u>≥3.6 h</u>
EPIC - Norfolk Study	Source Prevalence (%)*	33.3	33.8	32.9
	Case Prevalence (%)**	22.9	32.6	44.5
	Case/Source Ratio	0.69	0.96	1.35
Average Case/Source Ratio***		0.71	1.01	1.76

*prevalences of sitting or TV viewing in the source population at baseline; **prevalences of sitting or TV viewing in cases (decedents) at baseline;

***average case/source prevalence ratio from the prospective cohort studies.

Figure 1.




Figure 2.





Figure 3.





Sedentary behavior and life expectancy in the United States: A cause-deleted life table analysis

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2012-000828.R1
Article Type:	Research
Date Submitted by the Author:	07-Feb-2012
Complete List of Authors:	Katzmarzyk, Peter; Pennington Biomedical Research Center, Lee, I-Min; Harvard Medical School
Primary Subject Heading:	Public health
Secondary Subject Heading:	Epidemiology
Keywords:	PREVENTIVE MEDICINE, physical activity, lifestyle

SCHOLARONE™
Manuscripts

Peer Review Only

1
2
3
4 **Sedentary behavior and life expectancy in the United States: A cause-deleted life table**
5 **analysis**
6

7 Peter T. Katzmarzyk, I-Min Lee
8
9

10
11
12 Pennington Biomedical Research Center, Louisiana State University System, Baton Rouge, LA,
13 70808 Peter T. Katzmarzyk professor
14

15 Division of Preventive Medicine, Brigham & Women's Hospital and Harvard Medical School,
16 Boston, MA, 02215 I-Min Lee associate professor
17
18

19 Correspondence to: peter.katzmarzyk@pbrc.edu
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Abstract

Objectives: To determine the impact of sitting and television viewing on life expectancy in the United States.

Design: Prevalence-based cause-deleted life table analysis.

Setting: Summary relative risks of all-cause mortality associated with sitting and television viewing were obtained from a meta-analysis of available prospective cohort studies.

Prevalences of sitting and television viewing were obtained from the U.S. National Health and Nutrition Examination Survey.

Primary Outcome Measure: Life expectancy at birth.

Results: The estimated gains in life expectancy in the US population were 2.00 years for reducing excessive sitting to less than three hours per day and a gain of 1.38 years from reducing excessive television viewing to less than two hours per day. The lower and upper limits from a sensitivity analysis which involved simultaneously varying the estimates of RR (using the upper and lower bounds of the 95% CI) and the prevalence of television viewing ($\pm 20\%$) were 1.39 and 2.69 years for sitting and 0.48 and 2.51 years for television viewing, respectively.

Conclusions: Reducing sedentary behaviors such as sitting and television viewing has the potential to increase life expectancy in the United States.

Subject Headings: public health; epidemiology

Keywords: Preventive medicine; physical activity; lifestyle

Article Summary

Article focus

- This paper presents the results of an analysis aimed at determining the effects of sedentary behavior on life expectancy in the United States

Key messages

- The analyses indicate that population life expectancy in the United States would be 2.00 years higher if adults reduced their time spent sitting to less than three hours per day, and 1.38 years higher if they reduced television viewing to less than two hours per day.

Strengths and limitations

- The use of the well-accepted prevalence-based methodology to estimate the public health burden of sedentary behavior is a marked strength, which allows for comparability with the effects of other established risk factors.
- This study relied on self-reported engagement in sedentary behaviors rather than an objective measurement, which is a limitation.

1
2
3 Over 60 years of research on physical activity and health culminated in the release of the
4 *2008 Physical Activity Guidelines for Americans*, which recommend that adults should
5 accumulate at least 150 minutes of moderate-to-vigorous physical activity per week to attain the
6 health benefits associated with physical activity.¹ Recently, there has been considerable interest
7 in understanding the role of sedentary behaviors on health, independent of overall physical
8 activity levels,²⁻⁴ since one can be both sedentary and physically active (e.g., an office worker
9 who sits most of his work hours, but who also jogs regularly).

10
11 Sedentary behaviors which involve sitting for extended periods are ubiquitous in modern
12 society. Based on self-reports, a recent survey of 20 countries documented a median of 300
13 minutes per day spent sitting, ranging from ≤ 180 minutes per day in Portugal, Brazil and
14 Colombia to ≥ 360 minutes per day in Taiwan, Norway, Hong Kong, Saudi Arabia and Japan.⁵
15 Several studies have demonstrated positive associations between sedentary behaviors
16 including sitting and television viewing and health outcomes such as type 2 diabetes,
17 cardiovascular disease mortality and all-cause mortality.⁶ Thus, excessive time spent in
18 sedentary behavior is undoubtedly having an impact on public health. A recent study from
19 Australia estimated that television viewing reduced life expectancy at birth by 1.8 years in men
20 and 1.5 years in women.⁷ The purpose of this study was to determine the impact of sedentary
21 behaviors on life expectancy in the United States.

22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Methods

A prevalence-based approach was used to estimate the impact of sedentary behavior on life expectancy. The population-attributable fraction (PAF) was computed from the prevalences of sedentary behaviors (defined here as sitting and television viewing) and the relative risks (RR) of all-cause mortality associated with these behaviors. The PAF equation used was $\sum P_i(RR_i - 1/RR_i)$, where P is the prevalence of the risk factor among cases in stratum i (i=1 to 3 in this study; details are provided below). This equation produces internally valid estimates when confounding exists and adjusted RRs must be used.⁸

Relative Risks Associated with Sedentary Behaviors

Sedentary behavior can be captured globally using questionnaires that address total daily sitting time, or time spent in specific sedentary behaviors like television viewing, reading, or computer use. Published studies on the associations between sitting or television viewing and all-cause mortality were identified through a literature search of MEDLINE using search terms related to the purpose of this study (“physical inactivity”, “sedentary”, “cohort study”, “mortality” and “death”). Studies which reported on the relationship between sedentary and outcomes other than all-cause mortality were excluded from consideration. We pooled RR estimates from each study separately for sitting or television viewing and all cause-mortality using a random effects meta-analysis. Pooled RR estimates were obtained for two levels of sitting and two levels of television viewing relative to a referent group in each case. In order to maintain consistency across studies, the age- and sex-adjusted RR estimates were used from each study. In cases where the authors presented only multivariable-adjusted RR estimates, or used different exposure categories, we contacted them and asked them to provide this information. MIX 2.0 software was used to conduct the meta-analysis.⁹

Prevalence of Sedentary Behavior

The prevalences of time spent sitting and television viewing were obtained from the National Health and Nutrition Examination Survey (NHANES). NHANES uses a complex, multistage, probability sampling design to select participants who are representative of the civilian, noninstitutionalized U.S. population.¹⁰ The prevalences were obtained from the most recent NHANES data available for adults (2009-2010 for sitting; 2005-2006 for television viewing). The prevalence of sitting (<3 h, 3-5.9 h, and ≥6 h/day) among non-pregnant adults 18 years of age and older was determined from responses to the following question in the 2009-10 NHANES: “How much time do you usually spend sitting on a typical day?” The prevalence of television viewing (<2 h, 2-3.9 h, and ≥4 h/day) among non-pregnant adults 18 years of age and older was determined from responses to the following question in the 2005-06 NHANES: “Over the past

1
2
3 30 days, on average how many hours per day did you sit and watch TV or videos?" Data
4
5 analysis followed the guidelines of the National Center for Health Statistics for analysis of
6
7 NHANES data due to the complex sampling design and methods.¹¹
8
9

10 The PAF equation used in this study requires the prevalence of sedentary behavior among
11 cases (i.e. decedents) rather than from the source population (i.e. NHANES prevalence).
12
13 Therefore, each population prevalences obtained from NHANES was adjusted by using the
14
15 **weighted** average case:source prevalence ratio (i.e. the prevalences among cases divided by
16
17 the prevalences among the baseline source population from the prospective cohort studies).
18
19

20 Gains in Life Expectancy

21
22 The PAFs for all-cause mortality associated with sitting and television viewing were
23
24 computed using the summary RR estimates obtained from the meta-analyses and the adjusted
25
26 prevalences from NHANES, as described above. Potential gains in life expectancy attributable
27
28 to reducing sedentary behaviors were estimated using a cause-deleted life table analysis, which
29
30 estimates years of life gained at birth **if deaths from a specific cause are eliminated from the**
31
32 **current death rates.**^{12 13} The most current abridged life table for the United States (2009) was
33
34 downloaded from the World Health Organization website.¹⁴ The PAFs computed for sedentary
35
36 behavior were used to reduce the mortality rates in the life table for adults aged 18 years and
37
38 older, and the life expectancy at birth was recalculated from the new mortality rates. The
39
40 difference between the current life expectancy and the cause-deleted life expectancy represents
41
42 the estimated gain in life expectancy from reducing the prevalence of sedentary behavior.
43
44
45

46 Sensitivity Analysis

47
48 A sensitivity analysis was used to estimate the effects of simultaneously varying the
49
50 estimates of RR (using the upper and lower bounds of the 95% CI) and the prevalences of
51
52 sitting and television viewing ($\pm 20\%$).
53
54
55
56
57
58
59
60

Results

The characteristics and main results of the prospective cohort studies included in the meta-analyses of sedentary behavior and mortality are presented in Table 1. Two studies were included that investigated the association between sitting and all-cause mortality.^{15 16} The exposure levels of sitting in the first study were (1) none/¼ of the time, (2) ½ the time, and (3) ¾ of the time/all of the time based on activities that participants did most days of the week,¹⁵ whereas the levels in the second study were (1) <3 h, (2) 3-5 h, and (3) ≥6 h of leisure-time sitting.¹⁶ Thus, three categories of exposure were used in each cohort study, and the prevalence of sitting categories from NHANES were obtained for three groups (<3 h, 3-5.9 h, and ≥6 h/day). However, given that the exposure categories from the Canadian cohort study were not quantifiable in terms of absolute hours/day, some misclassification may have occurred when combining the results.

Three studies were included that studied the association between television viewing and all-cause mortality.¹⁷⁻¹⁹ The levels of television viewing in two of the studies were (1) <2 h, (2) 2-3.9 h, and (3) ≥4 h,^{17 18} whereas the levels in one study were (1) <2.5 h, (2) 2.5-3.6 h, and (3) >3.6 h.¹⁹ The prevalence of television viewing obtained from NHANES (<2 h, 2-3.9 h, and ≥4 h/day) match quite well with the exposure categories from the cohort studies.

Figure 1 presents the results of the meta-analysis for sitting and all-cause mortality. The pooled relative risks were 1.18 (1.14 - 1.21) and 1.45 (1.39-1.51) for levels 2 and 3 versus level 1, respectively. Figure 2 presents the results for the meta-analysis for television viewing and all-cause mortality. The pooled relative risks were 1.17 (1.04 - 1.32) and 1.49 (1.22-1.82) for levels 2 and 3 versus level 1, respectively.

The prevalences of sitting and television viewing in cases (decedents) and in the source population at baseline in the prospective studies are presented in Table 2, along with the average case:source prevalence ratio. These ratios were applied to the population prevalences obtained from NHANES in order to estimate the prevalences among cases in the population,

1
2
3 which are required for inclusion in the PAF calculations. Figure 3 presents the weighted
4
5 population prevalences of sitting and television viewing in NHANES, along with the prevalences
6
7 after adjustment for the average case:source prevalence ratios from Table 2.
8
9

10 The PAFs for all-cause mortality associated with sitting and television viewing were 27% and
11
12 19%, respectively. The results of the life table analyses indicate a gain in life expectancy from
13
14 reducing the prevalence of sedentary behavior from levels 2 or 3 to level 1 results in a gain of
15
16 2.00 years for reducing sitting prevalence and a gain of 1.38 years from reducing television
17
18 viewing prevalence. The lower and upper limits from the sensitivity analyses were 1.39 and 2.69
19
20 years for sitting and 0.48 and 2.51 years for television viewing, respectively.
21
22

23 Discussion

24
25 The results of this study indicate that limiting sitting to less than three hours per day and
26
27 limiting television viewing to less than two hours per day may increase life expectancy at birth in
28
29 the United States by approximately 2.0 and 1.4 years, respectively, assuming a causal
30
31 relationship. The PAF provides a *theoretical* estimate of the effects of a risk factor on an
32
33 outcome at the population level, in this case, all-cause mortality. The results indicate that
34
35 sedentary behaviors are accounting for between 1.4 and 2.0 years of life expectancy at birth.
36
37 This should not be interpreted to mean that people who are more sedentary can expect to live
38
39 1.4 or 2.0 years less than someone who does not engage in these behaviors as much. Life
40
41 expectancy is a population statistic and it does not apply to individuals. A recent meta-analysis
42
43 of television viewing and all-cause mortality estimated that the relative risk of all-cause mortality
44
45 was 1.13 (95% CI: 1.07-1.18) per 2 hours of daily television viewing, which corresponded with
46
47 104 deaths per 100,000 people in the United States.⁴
48
49

50
51 This study has several strengths and limitations that warrant discussion. The use of the well-
52
53 accepted prevalence-based PAF methodology to estimate the public health burden of sedentary
54
55 behavior is a marked strength, which allows for comparability with the effects of other
56
57 established risk factors. However, the PAF provides a *theoretical* estimate of the effects of risk
58
59
60

1
2
3 factor on a health outcome, and further research is required using an incidence-based
4
5 approach. Our analysis assumes that there is a causal relationship between sedentary behavior
6
7 and mortality. While studies using randomized designs are not possible, further observational
8
9 studies, which control for confounding, will add to the evidence for causation. Another major
10
11 strength of this study is the use of representative population data from NHANES to quantify the
12
13 exposure of the population to sedentary behaviors. However, our analysis estimated the overall
14
15 gains in life expectancy at the population level, and assumes that the effects of sedentary time
16
17 on all-cause mortality are consistent across age and demographic sub-groups of the population.
18
19 Each of the cohort studies provided multivariable-adjusted RR estimates for sedentary behavior
20
21 and mortality using different combinations of covariates, and we chose to use summary RR
22
23 estimates based on RR adjusted for age and sex in order to maintain consistency across
24
25 studies. The degree to which this approach has yielded an overestimation of the independent
26
27 effect of sedentary behavior on life expectancy is not known.
28
29
30

31 This study relied on self-reported engagement in sedentary behaviors which introduces the
32
33 possibility for error and recall bias. Future cohort studies should attempt to better quantify
34
35 sedentary behavior using objective activity monitors; however, the self-reported estimates of
36
37 sitting reported in NHANES are similar to those obtained for the U.S. in a study of 20 countries⁵
38
39 which indicates some face validity to the results. For television viewing, the categories of
40
41 exposure reported in two of the cohort studies (< 2 h, 2-3.9 h, ≥ 4 h) matched the categories
42
43 reported in NHANES; however, Wijndaele et al.¹⁹ used a lower threshold for the upper category
44
45 (> 3.6 h) which may have resulted in an underestimate of the effects of television viewing on life
46
47 expectancy. The studies on sitting used different exposure categories (i.e. total sitting time
48
49 versus leisure-time sitting only), so assumptions had to be made when estimating the exposure
50
51 levels in NHANES. Inaccuracies associated with the assessment of sedentary behavior using
52
53 self-report methods in the cohort studies would have led to regression dilution bias, and resulted
54
55 in under-estimates of the association with all-cause mortality.
56
57
58
59
60

1
2
3 The results of several recent studies have suggested that the effects of sedentary behavior
4 on health may be independent of the effects of physical activity *per se*.²⁰ It has been estimated
5 that a lack of leisure-time physical activity accounts for approximately 0.9 years of life
6 expectancy at birth in Canada.²¹ Given that the studies used to derive the summary RR
7 estimates for the current study in many cases included leisure-time physical activity as a
8 covariate in multivariable-adjusted models and this did not appreciably change the estimates of
9 RR for sedentary behavior, the estimates of the effects on life expectancy may also be
10 considered independent. Current life expectancy in the United States (2009) is 78.5 years.¹⁴
11 The effects of sedentary behavior on life expectancy reported in this study are on a similar order
12 of magnitude as other chronic disease risk factors. For example, it has been estimated that
13 obesity accounts for between 0.30 and 1.08 years of population life expectancy at birth in the
14 United States, depending on gender, ethnicity, and severity of obesity.²² A more recent study
15 has estimated that the current distribution of body mass index, compared to an optimal
16 distribution (a mean of 21 kg/m²), accounts for 1.3 years of current life expectancy at birth in
17 both males and females in the United States.²³ Results from the same study indicate that
18 smoking is also associated with 2.5 years and 1.8 years of life expectancy at birth in males and
19 females, respectively.²³

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Sitting time is a global measure of sedentary behavior, whereas television viewing is
somewhat more specific. A recent review found that domain-specific behaviors such as
television viewing are recalled with more reliability than global measures of sitting and sedentary
behavior.²⁴ The degree to which differences in reliability between the measures used in this
study may have affected the estimates is not known. There is some evidence to suggest that
using a single global question to measure sitting produces lower estimates than more detailed,
domain-specific questions.^{24 25} Thus, it is likely that the prevalences of higher levels of sitting
reported in this study from NHANES are likely conservative. Using objective monitoring
(accelerometry) in the 2003-2003 NHANES, Matthews et al. reported that U.S. adults spend

1
2
3 approximately 7.7 hours per day engaged in sedentary behavior.²⁶ There are several potential
4 mechanisms that could explain the association between sedentary behavior and all-cause
5 mortality rates. Sedentary behavior is associated with an increased risk of the development of
6 chronic conditions such type 2 diabetes and cardiovascular disease.⁶ Further, human and
7 animal studies indicate that sedentary behavior is associated with elevated cardiometabolic
8 biomarkers and a poor risk factor profile.^{27 28} For example, hind limb suspension (unloading) in
9 rats results in marked immediate decreases in lipoprotein lipase activity, triglyceride uptake into
10 red skeletal muscle, and reductions in the concentration of HDL cholesterol.²⁹ Future
11 intervention research is required to determine the causal pathways between sedentary behavior
12 and health outcomes that have the potential to impact mortality rates.
13
14
15
16
17
18
19
20
21
22
23
24

25 In conclusion, the results of this study indicate that extended sitting time and television
26 viewing are having a meaningful impact on life expectancy in the United States. **Given that the**
27 **results from objective monitoring of sedentary time in NHANES has indicated that adults spend**
28 **an average of 55% of their day engaged in sedentary pursuits²⁶, a significant shift in behavior**
29 **change at the population level is required to make demonstrable improvements in life**
30 **expectancy.** Further research using intervention designs is required to determine the effects of
31 reducing sedentary behavior on health outcomes and to make recommendations regarding the
32 safe levels of sedentary behavior for the population.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends

Figure 1. Risk of all-cause Mortality Associated with Sitting. Level 3 corresponds to $\frac{3}{4}$ of all of the time for Katzmarzyk et al. 2009 and ≥ 6 h for Patel et al. 2010. Level 2 corresponds to $\frac{1}{2}$ of the time for Katzmarzyk et al. 2009 and 3-5 h for Patel et al. 2010. Level 1 corresponds to $\frac{1}{4}$ of the time for Katzmarzyk et al. 2009 and < 3 h for Patel et al. 2010.

Figure 2. Risk of all-cause Mortality Associated with Television Viewing. Level 3 corresponds to ≥ 4 h for Dunstan et al. 2010 and Stamatakis et al. 2011 and ≥ 3.6 h for Wijndaele et al. 2011. Level 2 corresponds to 2-3.9 h for Dunstan et al. 2010 and Stamatakis et al. 2011 and 2.5-3.6 h for Wijndaele et al. 2011. Level 1 corresponds to < 2 h for Dunstan et al. 2010 and Stamatakis et al. 2011 and < 2.4 h for Wijndaele et al. 2011.

Figure 3. Prevalences of A) sitting and B) television viewing in the U.S. National Health and Nutrition Examination Survey. *Adjusted prevalences of sitting and television viewing using the weighted average case/source prevalence ratio obtained from cohort studies of sedentary behavior and all-cause mortality.

1
2
3 **Funding:** This research received no specific grant from any funding agency in the public,
4 commercial or non-profit sectors.
5
6

7 **Competing Interests:** All authors have completed the Unified Competing Interest form at
8 www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and
9
10 declare that both authors have no financial or non-financial interests that may be relevant to the
11 submitted work.
12
13

14 **Data Sharing Statement:** There are no additional data available.
15
16

17 **Author Contributions:** Both authors were responsible study conception and design. PTK was
18 responsible for data analysis. Both authors drafted and critically revised the manuscript, and
19 approved the final version to be published. PTK is the guarantor for the study.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

er review only

References

1. US Department of Health and Human Services. *2008 Physical Activity Guidelines for Americans*: US Department of Health and Human Services, Centers for Disease Control and Prevention, 2008.
2. Katzmarzyk PT. Physical activity, sedentary behavior, and health: paradigm paralysis or paradigm shift? *Diabetes* 2010;59(11):2717-25.
3. Thorp AA, Owen N, Neuhaus M, Dunstan DW. Sedentary behaviors and subsequent health outcomes in adults a systematic review of longitudinal studies, 1996-2011. *Am J Prev Med* 2011;41(2):207-15.
4. Grontved A, Hu FB. Television viewing and risk of type 2 diabetes, cardiovascular disease, and all-cause mortality: a meta-analysis. *JAMA* 2011;305(23):2448-55.
5. Bauman A, Ainsworth BE, Sallis JF, et al. The descriptive epidemiology of sitting: A 20-country comparison using the International Physical Activity Questionnaire (IPAQ). *Am J Prev Med* 2011;41(2):228-35.
6. Proper KI, Singh AS, van Mechelen W, Chinapaw MJ. Sedentary behaviors and health outcomes among adults: a systematic review of prospective studies. *Am J Prev Med* 2011;40(2):174-82.
7. Veerman JL, Healy GN, Cobiac LJ, et al. Television viewing time and reduced life expectancy: a life table analysis. *Br J Sports Med* 2011.
8. Rockhill B, Newman B, Weinberg C. Use and misuse of population attributable fractions. *Am J Public Health* 1998;88(1):15-9.
9. Bax L. Professional software for meta-analysis in Excel. Version 2.01.4. BostatXL; <http://www.meta-analysis-made-easy.com> 2011.
10. Centers for Disease Control and Prevention (CDC). National Health and Nutrition Examination Survey. Available at: <http://www.cdc.gov/nchs/nhanes.htm>. Access verified October 7, 2011. 2011.

- 1
2
3 11. Centers for Disease Control and Prevention (CDC), National Center for Health Statistics.
4
5 The National Health and Nutrition Examination Survey Analytic and Reporting Guidelines.
6
7 Available at: http://www.cdc.gov/nchs/nhanes/nhanes2003-2004/analytical_guidelines.htm
8
9 (Accessed October 7, 2011). 2006.
- 10
11 12. Manuel DG, Luo W, Ugnat A-M, Mao Y. Cause-deleted health-adjusted life expectancy of
12
13 Canadians with selected chronic conditions. *Chronic diseases in Canada* 2003;24:108-15.
- 14
15 13. Newman SC. Formulae for cause-deleted life tables. *Statistics in medicine* 1987;6(4):527-8.
- 16
17 14. World Health Organization. Global Health Observatory Data Repository. Available at:
18
19 <http://apps.who.int/ghodata/> (Accessed October 7, 2011). 2011.
- 20
21 22
23 15. Katzmarzyk PT, Church TS, Craig CL, Bouchard C. Sitting time and mortality from all
24
25 causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc* 2009;41:998-1005.
- 26
27 16. Patel AV, Bernstein L, Deka A, et al. Leisure time spent sitting in relation to total mortality in
28
29 a prospective cohort of US adults. *Am J Epidemiol* 2010;172(4):419-29.
- 30
31 17. Dunstan DW, Barr EL, Healy GN, et al. Television viewing time and mortality: the Australian
32
33 Diabetes, Obesity and Lifestyle Study (AusDiab). *Circulation* 2010;121(3):384-91.
- 34
35 18. Stamatakis E, Hamer M, Dunstan DW. Screen-based entertainment time, all-cause
36
37 mortality, and cardiovascular events population-based study with ongoing mortality and
38
39 hospital events follow-up. *J Am Coll Cardiol* 2011;57(3):292-9.
- 40
41 19. Wijndaele K, Brage S, Besson H, et al. Television viewing time independently predicts all-
42
43 cause and cardiovascular mortality: The EPIC Norfolk Study *Int J Epidemiol* 2011;40:150-59.
- 44
45 20. Tremblay MS, Colley RC, Saunders TJ, Healy GN, Owen N. Physiological and health
46
47 implications of a sedentary lifestyle. *Appl Physiol Nutr Metab* 2010;35(6):725-40.
- 48
49 21. Katzmarzyk PT. Physical inactivity and life expectancy in Canada. *J Phys Act Health*
50
51 2006;3:381-89.
- 52
53 22. Olshansky SJ, Passaro DJ, Hershow RC, et al. A potential decline in life expectancy in the
54
55 United States in the 21st century. *N Engl J Med* 2005;352:1138-45.
- 56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
23. Danaei G, Rimm EB, Oza S, et al. The promise of prevention: the effects of four preventable risk factors on national life expectancy and life expectancy disparities by race and county in the United States. *PLoS medicine* 2010;7(3):e1000248.
24. Healy GN, Clark BK, Winkler EA, et al. Measurement of adults' sedentary time in population-based studies. *Am J Prev Med* 2011;41(2):216-27.
25. Clemes SA, David BM, Zhao Y, Han X, Brown WJ. Validity of two self-report measures of sitting time. *J Phys Act Health* 2011;In Press.
26. Matthews CE, Chen KY, Freedson PS, et al. Amount of time spent in sedentary behaviors in the United States, 2003-2004. *Am J Epidemiol* 2008;167:875-81.
27. Healy GN, Matthews CE, Dunstan DW, Winkler EA, Owen N. Sedentary time and cardio-metabolic biomarkers in US adults: NHANES 2003-06. *Eur Heart J* 2011;32(5):590-7.
28. Hamilton MT, Hamilton DG, Zderic TW. Exercise physiology versus inactivity physiology: an essential concept for understanding lipoprotein lipase regulation. *Exerc Sport Sci Rev* 2004;32(4):161-6.
29. Bey L, Hamilton MT. Suppression of skeletal muscle lipoprotein lipase activity during physical inactivity: a molecular reason to maintain daily low-intensity activity. *The Journal of physiology* 2003;551(Pt 2):673-82.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 1. Characteristics and main results of the prospective cohort studies included in the meta-analysis.

Study	Participants	Age at Baseline (years)	Follow-up (years)	Person-years	Deaths (n)	Results			
Sitting Time									
Katzmarzyk et al. 2009 Canada Fitness Survey	N = 17,013 men and women	18 - 90	12	204,732	1,832	Deaths	<u>None/¼ Time</u>	<u>½ Time</u>	<u>¾ Time/All</u>
						Person-yr	826	542	464
						Crude RR*	117,965	52,346	34,421
						Age-Sex-HR**	1.00	1.48	1.93
							1.00	1.14	1.51
Patel et al. 2010 Cancer Prevention Study II Nutrition Cohort	N = 123,216 men and women	50 - 74	14	1,610,728	19,230	Deaths	<u><3 h</u>	<u>3-5 h</u>	<u>≥6 h</u>
						Person-yr	7,068	9,194	2,968
						Crude RR*	755,214	688,709	166,805
						Age-Sex-HR**	1.00	1.43	1.90
							1.00	1.18	1.44
TV Viewing									
Dunstan et al. 2010 Australian Diabetes, Obesity and Lifestyle Study (AusDiab)	N = 8,800 men and women	≥ 25	6.6	58,087	284	Deaths	<u><2 h</u>	<u>2-3.9</u>	<u>≥4 h</u>
						Person-yr	105	125	54
						Crude RR*	33,024	20,737	4326
						Age-Sex-HR**	1.00	1.89	3.93
							1.00	1.20	1.67
Stamatakis et al. 2011 Scottish Health Survey	N = 4,512 men and women	≥ 35	4.3	19,364	325	Deaths	<u><2 h</u>	<u>2-3.9 h</u>	<u>≥4 h</u>
						Person-yr	42	138	146
						Crude RR*	3,328	10,548	5,488
						Age-Sex-HR**	1.00	1.04	2.11
							1.00	1.13	1.77
Wijndaele et al. 2011 EPIC - Norfolk Study	N = 13,197 men and women	45 - 79	9.5	124,902	1,270	Deaths	<u><2.5 h</u>	<u>2.5-3.6 h</u>	<u>≥3.6 h</u>
						Person-yr	291	414	565
						Crude RR*	41,936	42,288	40,680
						Age-Sex-HR**	1.00	1.41	2.17
							1.00	1.17	1.32

*relative risk ratio; **age- and sex-adjusted hazard ratio.

Table 2. Prevalences of sitting and television viewing at baseline in the prospective cohort studies included in the meta-analysis.

Study	Prevalences			
Sitting Time				
Katzmarzyk et al. 2009		<u>None/¼ Time</u>	<u>½ Time</u>	<u>¾ Time/All</u>
Canada Fitness Survey	Source Prevalence (%)*	56.9	25.7	17.4
	Case Prevalence (%)**	45.1	29.6	25.3
	Case/Source Ratio	0.79	1.15	1.45
Patel et al. 2010		<u><3 h</u>	<u>3-5 h</u>	<u>≥6 h</u>
Cancer Prevention Study II	Source Prevalence (%)*	46.2	43.1	10.7
Nutrition Cohort	Case Prevalence (%)**	36.8	47.8	15.4
	Case/Source Ratio	0.80	1.11	1.44
Average Case/Source Ratio***		0.80	1.11	1.44
TV Viewing				
Dunstan et al. 2010		<u><2 h</u>	<u>2-3.9 h</u>	<u>≥4 h</u>
Australian Diabetes, Obesity and Lifestyle Study (AusDiab)	Source Prevalence (%)*	56.5	35.9	7.6
	Case Prevalence (%)**	37.0	44.0	19.0
	Case/Source Ratio	0.65	1.23	2.50
Stamatakis et al. 2011		<u><2 h</u>	<u>2-3.9 h</u>	<u>≥4 h</u>
Scottish Health Survey	Source Prevalence (%)*	17.1	54.1	28.8
	Case Prevalence (%)**	13.4	45.2	41.4
	Case/Source Ratio	0.78	0.84	1.44
Wijndaele et al. 2011		<u><2.5 h</u>	<u>2.5-3.6 h</u>	<u>≥3.6 h</u>
EPIC - Norfolk Study	Source Prevalence (%)*	33.3	33.8	32.9
	Case Prevalence (%)**	22.9	32.6	44.5
	Case/Source Ratio	0.69	0.96	1.35
Average Case/Source Ratio***		0.69	1.03	1.75

*prevalences of sitting or TV viewing in the source population at baseline; **prevalences of sitting or TV viewing in cases (decedents) at baseline;

***weighted-average case/source prevalence ratio from the prospective cohort studies.

Figure 1.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

Weight **RR (95% CI)**

Level 2 versus Level 1

Patel et al. 2010 93.2% 1.18 (1.15 – 1.22)

Katzmarzyk et al;. 2009 6.8 % 1.14 (1.02 – 1.27)

Total 100% 1.18 (1.14 – 1.21)

Level 3 versus Level 1

Patel et al. 2010 87.9% 1.44 (1.38 – 1.50)

Katzmarzyk et al;. 2009 12.1% 1.51 (1.35 – 1.69)

Total 100% 1.45 (1.39 – 1.51)

0 0.5 1 1.5 2

RR (95% CI)

Figure 2.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58





Figure 3.

A)



B)

