



Obstructive Lung Disease and Edentulism in the Atherosclerosis Risk in Communities(ARIC) Study

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2012-001615
Article Type:	Research
Date Submitted by the Author:	12-Jul-2012
Complete List of Authors:	Offenbacher, Steven; Univ of NC School of Dentistry/NC Oral Health Institute/Koury Oral Health Sciences, Periodontology Beck, James; Univ of NC School of Dentistry/NC Oral Health Institute/Koury Oral Health Sciences, Dental Ecology; Univ of NC School of Dentistry/NC Oral Health Institute/Koury Oral Health Sciences, Dental Ecology Barros, Silvana; Univ of NC School of Dentistry/NC Oral Health Institute/Koury Oral Health Sciences, Periodontology Suruki, Robert; Worldwide Epidemiology, GlaxoSmithKline R&D Loewy, Zvi; Touro College of Pharmacy, Pharmaceutical and Biomedical Sciences
Primary Subject Heading:	Dentistry and oral medicine
Secondary Subject Heading:	Dentistry and oral medicine
Keywords:	Pulmonary disease, Edentulism, Chronic Obstructive disease, bronchitis

SCHOLARONE™
Manuscripts

only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Guideline
Not applicable

For peer review only

1
2
3 Word Count: 2763
4

5 **OBSTRUCTIVE LUNG DISEASE AND EDENTULISM IN THE ATHEROSCLEROSIS RISK IN**
6
7 **COMMUNITIES (ARIC) STUDY**
8
9

10
11 Running Title: COPD and Edentulism
12

13
14 **Keywords:** pulmonary disease, edentulism, chronic obstructive pulmonary disease, bronchitis
15
16

17
18 Steven Offenbacher, DDS¹, James D Beck, PhD², Silvana P Barros, DDS¹, Robert Suruki ,ScD³ and Zvi
19
20 G Loewy, PhD⁴
21
22

23
24
25 ¹Department of Periodontology, School of Dentistry, University of North Carolina , Chapel Hill, NC
26
27 27599
28

29
30 ²Department of Dental Ecology, School of Dentistry, University of North Carolina, Chapel Hill, NC
31
32 27599
33

34
35 ³Worldwide Epidemiology, GlaxoSmithKline R&D, Research Triangle Park, NC 27709-3398
36

37
38 ⁴Department of Pharmaceutical and Biomedical Sciences, Touro College of Pharmacy, New York, NY
39
40 10027 and Department of Biochemistry & Molecular Biology, New York Medical College, Valhalla, NY,
41
42 10595
43

44
45 *Address correspondence to

46 Steve Offenbacher, DDS, PhD, MMSc

47 Department of Periodontology, UNC School of Dentistry

48 North Carolina Oral Health Institute

49 PO Box 14290

50 Durham, NC 27709

51 Phone: (919)425-3597 Fax: (919)425-3530

52 Email: steve_offenbacher@dentistry.unc.edu
53
54
55
56
57
58
59
60

Abstract:

Objectives: We examined the potential association between prior COPD and edentulism, and whether the association varied by COPD severity using data from the Dental ARIC Study.

Design: Case-control

Setting: Community dwelling subjects from four US communities.

Participants and Measurements: Cases were identified as edentulous (without teeth) and subjects with one or more natural teeth were identified as dentate. COPD cases were defined by spirometry measurements that showed the ratio of forced expiratory volume (one second) to vital capacity to be less than 0.7. The severity of COPD cases was also scored using a modified Global Initiative for Chronic Obstructive Lung Disease classification criteria (GOLD scoring I-IV). Multiple-logistic regression was used to examine the association between COPD and edentulism, while adjusting for age, gender, center/race, ethnicity, education level, income, diabetes, hypertension, coronary heart disease and congestive heart failure, body mass index, smoking, smokeless tobacco use and alcohol consumption.

Results: 13,465 participants were included in this analysis (2,087 edentulous; 11,378 dentate). Approximately 28.3% of edentulous participants had prior COPD compared to 19.6% among dentate participants ($p < 0.0001$). After adjustment for potential confounders, we observed a 1.3(1.08-1.62) and 2.5(1.68-3.63) fold increased risk of edentulism among GOLD II and GOLD III/IV COPD, respectively, as compared to the non-COPD/dentate referent. Given the short period of time between the measurements of COPD (Visit2) and dentate status (Visit4) relative to the natural history of both diseases, neither temporality nor insight as to the directionality of the association can be ascertained.

Conclusion: We found a significant association between prior COPD and edentulism, with evidence of a positive incremental effect seen with increasing GOLD classification.

INTRODUCTION

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Chronic obstructive pulmonary disease (COPD) is currently the fifth leading cause of death world-wide, and is expected to rise to fourth place in 2030.¹ Although COPD has a heterogeneous clinical presentation primarily characterized by airflow limitation, generally associated with an abnormal inflammatory response of the lungs², it is also often associated with a significant systemic inflammatory response³ which has been correlated with adverse clinical effects^{3,4}.

Over the past decade there has been an increased interest in the link between respiratory diseases and oral infection. Higher levels of tooth-associated microorganisms (plaque) and periodontal disease have been associated with COPD^{5,6,7}. Recently, in a very comprehensive oral microbiology study, high levels of diverse microbes- including several implicated in COPD and denture stomatitis- were identified on previously worn dentures⁸. Thus both the teeth and the dentures are non-desquamating oral surfaces that allow the oral biofilm to emerge eliciting periodontal disease or denture stomatitis, respectively. This raises the question as to whether denture-associated microbes may contribute to COPD in a manner analogous to that similar to periodontal organisms. Because oral infections and denture-based sources of sepsis are both treatable and preventable, the potential association of high microbial burden of oral origin to COPD may represent modifiable risk factors that may potentially improve respiratory-related morbidity. It has been suggested by Garcia and colleagues⁷ that the observed association between oral health and COPD may be due to aspiration of oral pathogens, or a common underlying host susceptibility trait that places individuals at risk for both conditions such as an exaggerated inflammatory response that is triggered by oral microbes.

Although numerous studies have examined the association between oral health and respiratory diseases in dentate individuals, the published literature are scant with respect to studies that have examined the association between edentulism and COPD status/severity. We undertook the current study to examine the potential association between COPD and edentulism using data from the Atherosclerosis Risk in

1
2
3 Communities (ARIC) Study. These data include not only dentate status and full-mouth periodontal
4
5 examination measurements, but also includes results from direct pulmonary function tests and
6
7 symptomology assessments that enabled us to classify the levels of COPD present.
8
9

14 **METHODS**

15
16 We sought to test the hypothesis that the prevalence of edentulism was greater among individuals with a
17
18 reported history of lung disease, (i.e. COPD) and that the association with edentulism varied by COPD
19
20 severity as determined by a modified Global Initiative for Chronic Obstructive Lung disease (GOLD)
21
22 criteria. Our second hypothesis was that among dentate subjects there was a difference in the prevalence
23
24 of periodontal disease according to the severity of COPD as measured by GOLD criteria.
25
26

27 **Study Population and Study Design**

28
29 The entire cohort from the ARIC study was 15,792 and the available dental cohort Dental Atherosclerosis
30
31 Risk in Communities (D-ARIC) data were used to study participants with both a periodontal examination
32
33 and completed spirometry data. The cohort selection process and reasons for exclusion appear in Figure 1.
34
35 Descriptions of the cohort and the periodontal examination procedure have been described previously⁹.
36
37 Eligible participants were all aged 45-64 at study entry (i.e., Visit 1) and had spirometry measurements at
38
39 Visit 2 (3 years later) and periodontal examinations at D-ARIC Visit 4 (6 years from visit 2).
40
41
42
43
44

45 **Case Definitions**

46
47 Edentulism case definitions were defined as study participants without any natural teeth or implants.
48
49 Participants with one or more natural teeth (comprising 11,378 subjects) were selected as the dentate
50
51 subjects. The absence, presence or severity of periodontal disease among dentate subjects was determined
52
53 using the joint American Association of Periodontology/Center for Disease Control (AAP/CDC)
54
55 consensus definitions of Health, Early and Advanced disease.¹⁰
56
57
58
59
60

This investigation represents the first report that uses actual pulmonary function data to assess lung function as an exposure related to edentulism or periodontal disease. The measured exposure of pulmonary function was classified both as a dichotomous (e.g., COPD vs. no COPD) and ordinal categorical variable (e.g., no lung disease, restricted disease, COPD GOLD Stage I, Stage II, Stage III, and Stage IV), using the following criteria.

Lung Obstruction

Dichotomous variable (COPD/no COPD)—Prior lung obstruction (or COPD) was defined based on spirometry measurements collected during Visit 2. Participants with a ratio of forced expiratory volume in one second (FEV_1) and forced vital capacity (FVC) less than 0.7 were determined to have lung obstruction consistent with COPD.

Ordinal categorical variable (COPD GOLD Stage)—Severity of COPD was also examined in this study among COPD participants using a modification of the spirometry-based criteria for severity outlined in the GOLD guidelines.^{11,12}

GOLD Stage	Spirometry Criteria
No lung disease	$FEV_1/FVC \geq 0.7$ and $FEV_1 \geq 80\%$ predicted
Restricted disease	$FEV_1/FVC \geq 0.7$ and $FVC < 80\%$ predicted
Stage 0	Presence of respiratory symptoms in the absence of any lung function abnormality; and no lung disease
Stage I	$FEV_1/FVC < 0.7$ and $FEV_1 \geq 80\%$ predicted
Stage II	$FEV_1/FVC < 0.7$ and $50\% \leq FEV_1 < 80\%$ predicted
Stage III	$FEV_1/FVC < 0.7$ and $30\% \leq FEV_1 < 50\%$ predicted
Stage IV	$FEV_1/FVC < 0.7$ and $FEV_1 < 30\%$ predicted

Variables of Interest

Smoking data (both for cigarettes and cigars) collected during Visit 4 were used to adjust for the confounding effect of smoking on the relationship between lung obstruction (COPD) and being edentulous. Smoking was treated as a five-level categorical variable, as described in detail previously⁷. Cardiovascular disease_(CVD) comorbidities that may be associated with both history of lung obstruction and being edentulous were also examined using data from Visit 4. Other potential confounders examined included race and examination center, ethnicity, education level, income, frequency of dental visits, diabetes, body mass index, alcohol use, and medication utilization. These data were extracted from Visit 4.

Data Analysis

Multivariate logistic regression was used to examine the association between COPD and edentulous status. This was followed by a logistic regression analysis of the dentate subjects comparing the COPD prevalence and severity of dentate patients without periodontal disease to that of dentate patients with periodontal disease. The exposure variable, which is lung obstruction, was examined as both a dichotomous variable (e.g., positive prior COPD vs. no COPD) and by logistic models using nominal variables (e.g., no lung disease, restricted disease, GOLD 0, Stage I, Stage II, Stage III, and Stage IV) based upon spirometry data collected as part of the ARIC examination. For the no COPD or GOLD reference group we included: subjects with and without chronic respiratory symptoms with no lung function abnormality, and subjects with GOLD 0 and restricted disease. Multiple regression models were used to control for confounding. When adjusting for potential confounding variables we included in certain models variables that are known to modify either periodontal disease status or COPD, including Body mass index and smokeless tobacco. We developed minimally and fully-adjusted logistical models incorporating significant confounders and effect modifiers to compute odds ratios (OR) and 95% confidence intervals (CI). SAS Version 9.2 (Cary, NC). We also included traditional risk factors for edentulism, periodontal disease and COPD- such as age, even if not statistically significant in this cohort. A p-value of <0.05 was considered statistically significant for this study.

Sample Size and Power Calculations

We knew *a priori* that the ARIC cohort who had spirometry data from Visit 2 included 2,087 edentulous subjects, as determined by questionnaire data at Visit 4. We also had 11,378 dentate subjects with spirometry data at Visit 2, of which 5,155 subjects had a full-mouth periodontal exam completed. Thus, if approximately 9% of the 13,465 subjects (2,087 edentulous and 11,378 dentate) were determined to have COPD, we estimated that we had >99% power to detect a significant difference of 1% at $p=0.05$ in the prevalence of COPD when comparing edentulous vs. dentate subjects.

RESULTS

ARIC subject participation was based upon the inception cohort of 15,792 subjects. The Visit 4 screenings were broadly divided into 11,378 dentate and 2,087 edentulous subjects who also had spirometry data. These subjects are used for analyses in Tables 1-6 on the relationship between dentate status and COPD. Subjects were excluded ($n=2,327$) from the original inception participants for incomplete data, dental implants only, and missing Visit 4 screening. Table 7 is based upon the 8,274 subjects eligible for the oral exam, of which 6967 (84.1%) were actually examined. Of those examined, the periodontal portion of the exam was not completed on 174 subjects, resulting in 6,793 subjects with full periodontal data.

The distribution of study subjects according to dentate status and GOLD classification is shown in Table 1. Among the edentulous subjects (2087), there were 591 or 28.3% with COPD (Gold Stages I-IV). With the exception of GOLD I where the prevalence of edentulism was 14.7 percent, the prevalence of edentulous individuals increased from 16.4 percent for GOLD 0 to 35.6% for GOLD III/IV. . The crude odds ratio (not shown in Table 1) for edentulism being associated with prior COPD vs no COPD was 1.62 (95% CI, 1.46-1.80). Approximately 13 percent of individuals with no chronic respiratory symptoms and

1
2
3 no lung function abnormality (i.e. no lung disease) were edentulous. There were 2232 subjects with a
4
5 prior COPD (19.6%) among 11,378 dentate subjects.
6

7 Table 2 presents the prevalence of various subject characteristics according to GOLD classification. In
8
9 this table, the GOLD reference group for statistical testing combines the following three categories of
10
11 subjects: no lung disease, GOLD 0, and subjects with chronic respiratory symptoms but no lung function
12
13 abnormality. The general patterns reflected in Table 2 show that compared to the reference group those
14
15 subjects in the GOLD III-IV category were much more likely to be edentulous, to smoke, to use
16
17 smokeless tobacco, to be African American or North Carolina Caucasians, male, and somewhat more
18
19 likely to have lower levels of income and education, be older and more likely to have chronic heart failure
20
21 and heart disease.
22
23
24
25
26

27 Table 3 presents multivariate logistic models for the relationship between edentulous /dentate status and
28
29 GOLD stages using the GOLD reference group described above. In an unadjusted model individuals with
30
31 GOLD II and GOLD III/IV scores were at significantly higher odds for edentulism that increased with
32
33 increasing GOLD stage severity (OR=2.06; 95% CI 1.79-2.38 and OR=3.38; 95% CI 2.61-4.39,
34
35 respectively). Minimally adjusted models (race/center, sex and age) showed similar significant effects for
36
37 GOLD II and GOLD III/IV with little attenuation. The increased risk of being edentulous for GOLD II
38
39 and GOLD III/IV categories remain significant, but have slightly lower odds ratios. However, the fully
40
41 adjusted model showed more attenuation with the odds for being edentulous being 1.3 (1.08-1.62) and 2.5
42
43 (1.68-3.63) times greater for GOLD II disease and the most severe category GOLD III/IV, respectively.
44
45
46
47
48

49 Table 4 shows the association between periodontal status and GOLD classifications for the dentate
50
51 subjects. Because of small cell sizes in the GOLD III and IV groups, we combined them with GOLD II.
52
53 Using the CDC/AAP classification of periodontal disease, which is based upon clusters of clinical signs,
54
55 there is a significant positive association between higher GOLD stage and severe periodontal disease
56
57 (p<0.0001). Subjects with GOLD I or GOLD II-IV were more likely to have severe periodontitis than the
58
59
60

1
2
3 reference group. With respect to the individual periodontal variables measured as means or extent scores,
4
5 individuals with a GOLD classification of I were more likely to have a greater extent of periodontal
6
7 pocket depths of 4mm or more, greater extent of attachment loss of 3mm. or more, a higher mean
8
9 attachment loss and a greater extent of plaque in their mouths than the reference group. Those with
10
11 GOLD scores of II-IV were significantly higher on all periodontal measures and had fewer teeth
12
13 compared to the reference group.
14
15

16
17
18 Logistic models for the association between GOLD status and periodontal status using the AAP/CDC
19
20 definitions appear in Table 5. As in Table 4, small cell size resulted in grouping the more severe GOLD
21
22 stages together as GOLD II/III/IV. In general, the odds ratios for severe periodontal disease increased
23
24 with more severe GOLD stage in the crude, minimally adjusted and fully adjusted models. There is also
25
26 an increase in odds as the severity of periodontal disease increases for the two GOLD categories shown
27
28 using the periodontally healthy group as the reference group within each GOLD category. Although
29
30 adjusting for relevant confounders generally decreased the odds, there was still a statistically significant
31
32 association between higher GOLD stage and severe periodontal disease in the fully adjusted model
33
34 (OR=1.41; 95% CI 1.07-1.87).
35
36

37
38
39 The relationship between spirometry based GOLD classification and self-reported physician-based
40
41 diagnosis of bronchitis or emphysema is shown in Table 6. Participants classified as GOLD III/IV were
42
43 significantly more likely to have self-reported bronchitis or emphysema. There is a trend for a positive
44
45 association except for the GOLD zero category. Interestingly, 65.9% of subjects with GOLD III/IV
46
47 report no diagnosis of COPD suggesting that self-report of bronchitis or emphysema is not a strong
48
49 predictor of GOLD status. In fact, of all subjects reporting a positive diagnosis of bronchitis or
50
51 emphysema 53% demonstrated no evidence of chronic respiratory symptoms and no lung function
52
53 abnormality.
54
55
56
57
58
59
60

DISCUSSION

We found a significant association between prior COPD and edentulism, with evidence of an increasing prevalence of edentulism with increasing GOLD classification. Although there was some reduction in the magnitude of effect in the adjusted models, the association between dentate status and GOLD classification remained significant even in the fully adjusted model, with a 1.3 fold increased odds of being edentulous in the GOLD II category and a 2.5 fold increased odds of being edentulous in the most severe GOLD III/IV category in the fully adjusted regression model. This represents the first report of an association between edentulism and COPD in a population with GOLD measurements.

In the subgroup of dentate subjects, there was a significant association between the GOLD II - IV group vs. the GOLD reference group and prevalent periodontal disease with a significant increase in the odds of entry level and severe periodontal disease in the crude and minimally adjusted models. In the fully adjusted model, only the association between GOLD stage II/III/IV and severe periodontal disease [OR 1.41 (1.07-1.87)] remained significant. This is the first population report to include both full-mouth periodontal examination data as well as Spirometry assessments, thereby providing direct confirmation of the association between severe periodontal disease and COPD, as well as a dose-response relationship between these conditions.

While there is no previous published information available about the association between edentulism status and COPD, there is evidence in the literature of an association between periodontal disease and COPD, consistent with the findings of the current study^{5,6,7}. The results of this investigation were based upon standardized pulmonary function tests and suggest that studies that are based upon self-reported diagnoses can potentially result in significant misclassification error. Our data shown in Table 6 suggest that of the 12,297 subjects reporting no diagnosis, 2,390 subjects or 19.4% were classified as GOLD I-IV.

1
2
3 Among those 433 subjects with GOLD I-IV the positive predictive rate was only 15.3%. These findings
4 suggest that an important strength of the current study was the classification of disease based on an
5 objective biological measure (e.g., spirometry) rather than a self-reported measure.
6
7
8
9

10
11 Additional strengths of the current study include the large sample size, the inclusion of African
12 Americans providing some racial diversity of the sample, the use of a full-mouth examination protocol
13 and the collection of information about key confounding variables, including but not limited to: tobacco
14 use, hypertension, diabetes, and income. Furthermore, the outcomes of interest, edentulism and
15 periodontal disease, were evaluated using a comprehensive oral examination at Visit 4.
16
17
18
19
20
21
22
23

24
25 The current study must also be considered in light of certain limitations. First, the design of the current
26 study does not allow us to determine the directionality or temporality of any association found. It has been
27 reported previously that the dental ARIC sub-sample is healthier than the greater ARIC sample and that
28 the edentulous subjects within the ARIC dataset have overall poorer health,⁹ as compared to the ARIC
29 dental sample. Thus, it is possible that the observed relationship between both edentulism and periodontal
30 disease with prior COPD may be due in part to shared risk factors. There is also the potential bias due to
31 differential loss to follow-up prior to Visit 4. The current study included only those subjects with dental
32 history information at Visit 4. Finally, it is possible that participants with COPD may have been more
33 likely to be lost to follow-up (compared to participants without COPD) or that participants with more
34 severe COPD may have had a lower probability of surviving to Visit 4 than participants with mild COPD
35 or normal lung function. There are no data in this report to mechanistically link edentulism with
36 increased risk for COPD. Previous studies of periodontal disease associations with COPD have discussed
37 the potential role of periodontal infection serving as a chronic repository of pathogenic oral organisms
38 that could be aspirated into the airway to challenge the lungs⁶. Periodontal infections are also associated
39 with increases in systemic markers of inflammation including biomarkers such as C-reactive protein
40 (CRP)¹³. By analogy, edentulism is commonly associated with denture infections (denture mucositis)
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 which induce mucosal inflammation and unclean dentures can also pose as a chronic reservoir of oral
4 pathogens. However, the potential role of denture-based infection as a potential risk factor for COPD
5 remains to be elucidated.
6
7
8
9

10 11 12 13 14 **CONCLUSIONS**

15
16
17
18 Edentulism was associated with prior COPD, with an incremental effect observed with increasing GOLD
19 classification. Additionally, among dentate individuals, severe periodontal disease was associated with
20 GOLD classification. This is one of the first analyses of COPD and dentate status, and expands upon the
21 available literature regarding periodontal disease and COPD. Further research is warranted to explore the
22 relationship of edentulism and periodontal disease with COPD.
23
24
25
26
27
28

29 30 31 **Conflict of Interest**

32
33 There are no potential conflicts of interest that exist with any companies/organizations whose products or
34 services may be discussed in this article.
35
36
37

38 39 **Source of Funding**

40 This work was supported by a grant from GlaxoSmithKline and we declare that the sponsor had no role in
41 the study design, collection, analysis and interpretation of data; in the writing of the manuscript; and in
42 the decision to submit the manuscript for publication.
43
44

45 46 **Responsibility for Data**

47 This is to verify that we have full access to all the data in this study and take full responsibility for the
48 integrity of the data and accuracy of the data analysis, including and especially any adverse effects.
49
50
51
52
53
54

55 56 **Abbreviations:**

57 AAP/CDC – American Association of Periodontology/Center for Disease Control
58
59
60

1
2
3 ARIC – Atherosclerosis Risk in Communities
4 CHF – Congestive Heart Failure
5 CI – Confidence Intervals
6 COPD – Chronic Obstructive Pulmonary Disease
7 CVD – Cardiovascular Disease
8 D-ARIC – Dental Atherosclerosis Risk in Communities
9 FEV – Forced Expiratory Volume
10 FVC – Forced Vital Capacity
11 GOLD – Global Initiative for Chronic Obstructive Lung Disease
12 MAL – Mean Attachment Loss
13 NHANES – National Health and Nutrition Examination Survey
14 OR – Odds Ratio
15 SES – Social Economic Status
16 WHO – World Health Organization
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Wouters EFM. Economic analysis of the confronting COPD survey: an overview of results. *Respiratory Medicine* 2003;**97**(C): S3-S14.
2. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006;**3**(11):e442.
3. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med*. 2007;**176**(6):532-55.
4. van Eeden, SF, Sin DD. Chronic Obstructive Pulmonary Disease: A Chronic Systemic Inflammatory Disease. *Respiration* 2008;**75**(2):224-38.
5. Scannapieco FA, Ho AW. Potential associations between chronic respiratory disease and periodontal disease: analysis of National Health and Nutrition Examination Survey III. *J Periodontol* 2001;**72**(1):50-6.
6. Scannapieco FA, Papandonatos GD, Dunford RG. Associations between oral conditions and respiratory disease in a national sample survey population. *Ann Periodontol* 1998;**3**(1):251-6.
7. Garcia RI, Nunn ME, Vokonas PS. Epidemiologic associations between periodontal disease and chronic obstructive pulmonary disease. *Ann Periodontol* 2001;**6**(1):71-7.
8. Glass RT, Conrad RS, Bullard JW, Goodson LB, Mehta N, Lech SJ et al. Evaluation of microbial flora found in previously worn prostheses from the Northeast and Southwest regions of United States. *Journal of Prosthetic Dentistry* 2010;**103**(6):384-9.
9. Elter JR, Offenbacher S, Toole JF, Beck JD. Relationship of periodontal disease and edentulism to stroke/TIA. *J Dent Res* 2003;**82**(12):998-1001.
10. Page RC, Eke PI. Case definitions for use in population-based surveillance of periodontitis. *J Periodontol* 2007;**78**(7 Suppl): 1387-99.
11. Pauwels RA, Buist AS, Calverley PMA, Jenkins CR, Hurd SS. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop Summary. *American Journal of Respiratory and Critical Care Medicine* 2001;**163**(5): 1256-76.
12. Mannino DM, Doherty DE, Buist SA. Global Initiative on Obstructive Lung Disease (GOLD) classification of lung disease and mortality: findings from the Atherosclerosis Risk in Communities (ARIC) study. *Respiratory Med* 2006;**100**(1):115-22.
13. Slade GD, Ghezzi EM, Heiss G, Beck JD, Riche E, Offenbacher S. Relationship between Periodontal Disease and C-Reactive Protein among Adults in the Atherosclerosis Risk in Communities Study. *Arch Intern Med*. 2003;**163**(10):1172-9.

Table 1. Dentate Status by GOLD Stage

	No Chronic Respiratory Symptoms & No Lung Function Abnormality	Restricted Disease (FEV ₁ /FVC≥0.7 and FVC<80% predicted)	GOLD 0 (Chronic Respiratory Symptoms & No Lung Function Abnormality)	GOLD I	GOLD II	GOLD III/IV	p-value*
Edentulous	1294 (13.3%)	151 (24.6%)	51 (16.4%)	208 (14.7%)	290 (25.2%)	93 (35.6%)	
Dentate	8422 (86.7%)	464 (75.5%)	260 (83.6%)	1204 (85.3%)	860 (74.8%)	168 (64.4%)	<0.0001

*Chisq

Table 2. Demographics by GOLD Stage

	GOLD Reference*	GOLD I	GOLD II	GOLD III/IV	p-value**
Edentulous	1496 (14.1%)	208 (14.7%)	290 (25.2%)	93 (35.6%)	
Dentate	9146 (85.9%)	1204 (85.3%)	860 (74.8%)	168 (64.4%)	<0.0001
Race/Center					
MS African Americans	2654 (25.1%)	191 (13.6%)	196 (17.1%)	73 (28.4%)	
NC African Americans	285 (2.7%)	35 (2.5%)	27 (2.4%)	11 (4.3%)	
NC White	2211 (20.9%)	401 (28.5%)	344 (30.0%)	84 (32.7%)	
Washington Co, MD	2680 (25.3%)	350 (24.9%)	337 (29.4%)	56 (21.8%)	
Surburban Minneapolis	2747 (26.0%)	430 (30.6%)	243 (21.2%)	33 (12.8%)	<0.0001
Female	6344 (59.6%)	684 (48.4%)	457 (39.7%)	125 (47.9%)	
Male	4298 (40.4%)	728 (51.6%)	693 (60.3%)	136 (52.1%)	<0.0001
Mean Age (StdDev)	62.4 (5.59)	64.2 (5.69)	64.9 (5.57)	64.2 (5.70)	<0.0001
Mean BMI (StdDev)	29.2 (5.68)	27.1 (4.81)	27.6 (5.34)	27.3 (6.16)	<0.0001
Diabetes (Yes)	1573 (17.4%)	132 (10.8%)	179 (19.7%)	36 (20.7%)	
No	7486 (82.6%)	1090 (89.2%)	732 (80.4%)	138 (79.3%)	<0.0001
Hypertension (Yes)	3591 (39.4%)	388 (31.9%)	374 (40.7%)	74 (41.8%)	
No	5535 (60.7%)	829 (68.1%)	545 (59.3%)	103 (58.2%)	<0.0001
Income (Low)	2718 (31.1%)	365 (31.2%)	321 (37.2%)	72 (41.6%)	
Medium	3071 (35.2%)	440 (37.6%)	316 (36.7%)	59 (34.1%)	
High	2945 (33.7%)	366 (31.3%)	225 (26.1%)	42 (24.3%)	<0.0001
Education (Basic)	2226 (21.0%)	291 (20.6%)	332 (28.9%)	93 (35.8%)	
Intermediate	4360 (41.0%)	564 (40.0%)	480 (41.8%)	101 (38.9%)	
Advanced	4038 (38.0%)	556 (39.4%)	336 (29.3%)	66 (25.4%)	<0.0001
Alcohol Use (Yes)	4404 (48.4%)	657 (53.9%)	463 (51.1%)	78 (44.3%)	
No	4695 (51.6%)	561 (46.1%)	444 (49.0%)	98 (55.7%)	0.001
Smoke (Current, Heavy)	748 (8.6%)	191 (16.4%)	256 (29.8%)	33 (20.6%)	
Current, Light	245 (2.8%)	34 (2.9%)	15 (1.7%)	1 (0.6%)	
Former, Heavy	1304 (14.9%)	309 (26.5%)	339 (39.4%)	76 (47.5%)	
Former, Light	1981 (22.7%)	228 (19.6%)	100 (11.6%)	17 (10.6%)	
Never	4448 (51.0%)	403 (34.6%)	150 (17.4%)	33 (20.6%)	<0.0001
Smokeless Tobacco (Yes)	699 (7.7%)	131 (10.8%)	138 (15.2%)	34 (19.3%)	
No	8401 (92.3%)	1088 (89.3%)	769 (84.8%)	142 (80.7%)	<0.0001
Coronary Heart Disease (Yes)	718 (6.9%)	106 (7.6%)	159 (14.2%)	31 (13.0%)	
No	9727 (93.1%)	1281 (92.4%)	963 (85.8%)	208 (87.0%)	<0.0001
Congestive Heart Failure (Yes)	191 (2.1%)	25 (2.1%)	41 (4.6%)	9 (5.1%)	
No	8878 (97.9%)	1194 (98.0%)	861 (95.5%)	166 (94.9%)	<0.0001

*GOLD Reference: No Chronic Respiratory Symptoms & No Lung Function Abnormality + GOLD Zero (Chronic Respiratory Symptoms & No Lung Function Abnormality) + Restrictive (FEV1/FVC \geq 0.70 and FVC < 80% predicted)

**Chisq or t-test

Table 3: Generalized Logits Models: Association between edentulism and GOLD Stages as compared to dentate subjects

	GOLD Reference*	GOLD I	GOLD II	GOLD III/IV
Model 1 Crude				
Edentulous	Reference	1.06 (0.90-1.24)	2.06 (1.79-2.38)	3.38 (2.61-4.24)
Model 2 Minimally Adjusted**				
Edentulous	Reference	1.12 (0.93-1.34)	1.92 (1.61-2.28)	3.17 (2.28-4.38)
Model 3 Fully Adjusted ****				
Edentulous	Reference	1.05 (0.86-1.28)	1.32 (1.08-1.62)	2.47 (1.68-3.64)

*GOLD Reference: No Chronic Respiratory Symptoms & No Lung Function Abnormality + GOLD Zero (Chronic Respiratory Symptoms & No Lung Function Abnormality) + Restrictive (FEV1/FVC \geq 0.70 and FVC < 80% predicted)

**Model 2 is minimally adjusted for race/center (5-level), male, age (years)

***Model 3 is fully adjusted for race/center (5-level), male, age (years), diabetes, hypertension, smoking (5-levels), smokeless tobacco, alcohol, income (3-level), education (3-level), CFH and CHD

Table 4. Periodontal measures among the 5,155 dentate subjects by GOLD stages

	GOLD Reference*	GOLD I	GOLD II/III/IV	Overall p-value
CDC/AAP Definition				
Healthy N (SD)	2378 (43.0%)	281 (38.1%)	151 (28.5%)	
Entry N (SD)	2271 (41.1%)	295 (40.0%)	233 (44.1%)	
Severe N (SD)	878 (15.9%)	161 (21.9%)	145 (27.4%)	<0.0001
Mean (StdDev)				
Extent PD >=4mm	7.05 (11.3)	8.12 (12.3) ^a	9.66 (14.8) ^d	<0.0001
Mean PD	1.89 (0.56)	1.94 (0.61)	2.03 (0.74) ^d	<0.0001
Extent AL >=3mm	21.7 (21.8)	27.6 (26.0) ^d	34.3 (28.7) ^d	<0.0001
Mean AL	1.72 (0.96)	1.94 (1.15) ^d	2.29 (1.43) ^d	<0.0001
Extent BOP	25.2 (23.4)	25.1 (24.3)	27.7 (26.4) ^a	0.07
Extent PQ >=1	42.5 (38.3)	39.3 (37.7) ^a	49.9 (39.3) ^d	<0.0001
Number of Teeth	22.0 (6.99)	21.5 (7.26)	18.8 (8.09) ^d	<0.0001

^a = p<0.05, b=p<0.01, c=p<0.001, d=p<0.0001,

All superscripted p values are compared to reference group by Chi square or GLM

Table 5: Generalized Logits Models: Periodontal status (CDC/AAP Definition) by GOLD stages

	GOLD I	GOLD II/III/IV
Model 1 Crude		
Healthy	Reference	Reference
Entry	1.10 (0.92-1.31)	1.62 (1.31-2.00)
Severe	1.55 (1.26-1.91)	2.60 (2.05-3.31)
Model 2 Minimally Adjusted*		
Healthy	Reference	Reference
Entry	0.99 (0.83-1.19)	1.42 (1.14-1.77)
Severe	1.46 (1.17-1.82)	2.17 (1.67-2.80)
Model 3 Fully Adjusted **		
Healthy	Reference	Reference
Entry	0.95 (0.79-1.14)	1.10 (0.87-1.39)
Severe	1.23 (0.97-1.55)	1.41 (1.07-1.87)

*Model 2 is minimally adjusted for race/center (5-level), male, age (years)


**Model 3 is fully adjusted for race/center (5-level), male, age (years), diabetes, hypertension, smoking (5-levels), smokeless tobacco, alcohol, income (3-level), education (3-level)

Table 6: Self reported physician diagnosis of bronchitis or emphysema according to GOLD stages .

	No Chronic Respiratory Symptoms & No Lung Function Abnormality	GOLD Zero (Chronic Respiratory Symptoms & No Lung Function Abnormality)	Restricted Disease (FEV₁/FVC≥0.7 and FVC<80% predicted)	GOLD I	GOLD II	GOLD III/IV	p-value^a
Self Report (Yes)	633 (6.5%)	65 (20.9%)	64 (10.4%)	143 (10.1%)	201 (17.5%)	89 (34.1%)	
No	9083 (93.5%)	246 (79.1%)	551 (89.6%)	1269 (89.7%)	949 (82.5%)	172 (65.9%)	<0.0001

a) Chi square

Participant Flow Diagram



001615 on 18 December 2012. Downloaded from <http://bmjopen.bmj.com/> on April 18, 2024 by guest. Protected by copyright.



Obstructive Lung Disease and Edentulism in the Atherosclerosis Risk in Communities(ARIC) Study

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2012-001615.R1
Article Type:	Research
Date Submitted by the Author:	17-Oct-2012
Complete List of Authors:	Offenbacher, Steven; Univ of NC School of Dentistry/NC Oral Health Institute/Koury Oral Health Sciences, Periodontology Beck, James; Univ of NC School of Dentistry/NC Oral Health Institute/Koury Oral Health Sciences, Dental Ecology; Univ of NC School of Dentistry/NC Oral Health Institute/Koury Oral Health Sciences, Dental Ecology Barros, Silvana; Univ of NC School of Dentistry/NC Oral Health Institute/Koury Oral Health Sciences, Periodontology Suruki, Robert; Worldwide Epidemiology, GlaxoSmithKline R&D Loewy, Zvi; Touro College of Pharmacy, Pharmaceutical and Biomedical Sciences
Primary Subject Heading:	Dentistry and oral medicine
Secondary Subject Heading:	Respiratory medicine, Cardiovascular medicine
Keywords:	Pulmonary disease, Edentulism, Chronic Obstructive disease, bronchitis

SCHOLARONE™
Manuscripts

only

1
2
3 Word Count: 2763
4

5 **OBSTRUCTIVE AIRWAY DISEASE AND EDENTULISM IN THE ATHEROSCLEROSIS RISK**
6
7 **IN COMMUNITIES (ARIC) STUDY**
8
9

10
11 Running Title: COPD and Edentulism
12

13
14 **Keywords:** pulmonary disease, edentulism, chronic obstructive pulmonary disease, bronchitis
15
16

17
18 Steven Offenbacher, DDS¹, James D Beck, PhD², Silvana P Barros, DDS¹, Robert Suruki ,ScD³ and Zvi
19
20 G Loewy, PhD⁴
21
22

23
24
25 ¹Department of Periodontology, School of Dentistry, University of North Carolina , Chapel Hill, NC
26
27 27599
28

29 ²Department of Dental Ecology, School of Dentistry, University of North Carolina, Chapel Hill, NC
30
31 27599
32

33 ³Worldwide Epidemiology, GlaxoSmithKline R&D, Research Triangle Park, NC 27709-3398
34

35
36 ⁴Department of Pharmaceutical and Biomedical Sciences, Touro College of Pharmacy, New York, NY
37
38 10027 and Department of Biochemistry & Molecular Biology, New York Medical College, Valhalla, NY,
39
40 10595
41

42 *Address correspondence to
43

44 Steve Offenbacher, DDS, PhD, MMSc
45

46 Department of Periodontology, UNC School of Dentistry
47

48 North Carolina Oral Health Institute
49

50 PO Box 14290
51

52 Durham, NC 27709
53

54 Phone: (919)425-3597 Fax: (919)425-3530
55

56 Email: steve_offenbacher@dentistry.unc.edu
57
58
59
60

Abstract:

Objectives: We examined the potential association between prior COPD and edentulism, and whether the association varied by COPD severity using data from the Dental ARIC Study.

Design: Cross-sectional **Setting:** Community dwelling subjects from four US communities.

Participants and Measurements: Cases were identified as edentulous (without teeth) and subjects with one or more natural teeth were identified as dentate. COPD cases were defined by spirometry measurements that showed the ratio of forced expiratory volume (one second) to vital capacity to be less than 0.7. The severity of COPD cases was also scored using a modified Global Initiative for Chronic Obstructive Lung Disease classification criteria (GOLD scoring I-IV). Multiple-logistic regression was used to examine the association between COPD and edentulism, while adjusting for age, gender, center/race, ethnicity, education level, income, diabetes, hypertension, coronary heart disease and congestive heart failure, body mass index, smoking, smokeless tobacco use and alcohol consumption.

Results: 13,465 participants were included in this analysis (2,087 edentulous; 11,378 dentate). Approximately 28.3% of edentulous participants had prior COPD compared to 19.6% among dentate participants ($p < 0.0001$). After adjustment for potential confounders, we observed a 1.3(1.08-1.62) and 2.5(1.68-3.63) fold increased risk of edentulism among GOLD II and GOLD III/IV COPD, respectively, as compared to the non-COPD/dentate referent. Given the short period of time between the measurements of COPD (Visit2) and dentate status (Visit4) relative to the natural history of both diseases, neither temporality nor insight as to the directionality of the association can be ascertained.

Conclusion: We found a statistically significant association between prior COPD and edentulism, with evidence of a positive incremental effect seen with increasing GOLD classification.

INTRODUCTION

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Chronic obstructive pulmonary disease (COPD) is currently the fifth leading cause of death world-wide, and is expected to rise to fourth place in 2030.¹ Although COPD has a heterogeneous clinical presentation primarily characterized by airflow limitation, generally associated with an abnormal inflammatory response of the lungs², it is also often associated with a significant systemic inflammatory response³ which has been correlated with adverse clinical effects^{3,4}.

Over the past decade there has been an increased interest in the link between respiratory diseases and oral infection. Higher levels of tooth-associated microorganisms (plaque) and periodontal disease have been associated with COPD^{5,6,7}. Recently, in a very comprehensive oral microbiology study, high levels of diverse microbes- including several implicated in COPD and denture stomatitis- were identified on previously worn dentures⁸. Thus both the teeth and the dentures are non-desquamating oral surfaces that allow the oral biofilm to emerge eliciting periodontal disease or denture stomatitis, respectively. This raises the question as to whether denture-associated microbes may contribute to COPD in a manner analogous to that similar to periodontal organisms. Because oral infections and denture-based sources of sepsis are both treatable and preventable, the potential association of high microbial burden of oral origin to COPD may represent modifiable risk factors that may potentially improve respiratory-related morbidity. It has been suggested by Garcia and colleagues⁷ that the observed association between oral health and COPD may be due to aspiration of oral pathogens, or a common underlying host susceptibility trait that places individuals at risk for both conditions such as an exaggerated inflammatory response that is triggered by oral microbes.

Although numerous studies have examined the association between oral health and respiratory diseases in dentate individuals, the published literature are scant with respect to studies that have examined the association between edentulism and COPD status/severity. We undertook the current study to examine the potential association between COPD and edentulism using data from the Atherosclerosis Risk in

1
2
3 Communities (ARIC) Study. We hypothesized that edentulism would be associated with a diagnosis of
4
5 COPD and display a higher prevalence with more spirometry categories of COPD. These data include not
6
7 only dentate status and full-mouth periodontal examination measurements, but also includes results from
8
9 direct pulmonary function tests and symptomology assessments that enabled us to classify the levels of
10
11 COPD present.
12

13 14 15 16 17 18 **METHODS** 19

20 We sought to test the hypothesis that the prevalence of edentulism was greater among individuals with a
21
22 reported history of lung disease, (i.e. COPD) and that the association with edentulism varied by COPD
23
24 severity as determined by a modified Global Initiative for Chronic Obstructive Lung disease (GOLD)
25
26 criteria. Our second hypothesis was that among dentate subjects there was a difference in the prevalence
27
28 of periodontal disease according to the severity of COPD as measured by GOLD criteria.
29

30 31 **Study Population and Study Design** 32

33 The entire cohort from the ARIC study was 15,792 and the available dental cohort Dental Atherosclerosis
34
35 Risk in Communities (D-ARIC) data were used to study participants with both a periodontal examination
36
37 and completed spirometry data. The cohort selection process and reasons for exclusion appear in Figure 1.
38
39 Descriptions of the cohort and the periodontal examination procedure have been described previously⁹.
40
41 Eligible participants were all aged 45-64 at study entry (i.e., Visit 1) and had spirometry measurements at
42
43 Visit 2 (3 years later) and periodontal examinations at D-ARIC Visit 4 (6 years from visit 2).
44
45
46
47
48

49 **Case Definitions** 50

51 Edentulism case definitions were defined as study participants without any natural teeth or implants.
52
53 Participants with one or more natural teeth (comprising 11,378 subjects) were selected as the dentate
54
55 subjects. The absence, presence or severity of periodontal disease among dentate subjects was determined
56
57
58
59
60

using the joint American Association of Periodontology/Center for Disease Control (AAP/CDC) consensus definitions of Health, Early and Advanced disease.¹⁰

This investigation represents the first report that uses actual pulmonary function data to assess lung function as an exposure related to edentulism or periodontal disease. The measured exposure of pulmonary function was classified both as a dichotomous (e.g., COPD vs. no COPD) and ordinal categorical variable (e.g., no lung disease, restrictive disease, COPD GOLD Stage I, Stage II, Stage III, and Stage IV), using the following criteria.

Airway obstruction

Dichotomous variable (COPD/no COPD)—Prior airway obstruction (or COPD) was defined based on spirometry measurements collected during Visit 2. Participants with a ratio of forced expiratory volume in one second (FEV₁) and forced vital capacity (FVC) less than 0.7 were determined to have airway obstruction consistent with COPD.

Ordinal categorical variable (COPD GOLD Stage)—Severity of COPD was also examined in this study among COPD participants using a modification of the spirometry-based criteria for severity outlined in the GOLD guidelines.^{11,12}

GOLD Stage	Spirometry Criteria
No lung disease	FEV ₁ /FVC ≥ 0.7 and FEV ₁ ≥ 80% predicted
Restrictive disease	FEV ₁ /FVC ≥ 0.7 and FVC < 80% predicted
Stage 0	Presence of respiratory symptoms in the absence of any lung function abnormality; and no lung disease
Stage I	FEV ₁ /FVC < 0.7 and FEV ₁ ≥ 80% predicted
Stage II	FEV ₁ /FVC < 0.7 and 50% ≤ FEV ₁ < 80% predicted
Stage III	FEV ₁ /FVC < 0.7 and 30% ≤ FEV ₁ < 50% predicted
Stage IV	FEV ₁ /FVC < 0.7 and FEV ₁ < 30% predicted

Variables of Interest

Smoking data (both for cigarettes and cigars) collected during Visit 4 were used to adjust for the confounding effect of smoking on the relationship between airway obstruction (COPD) and being edentulous. Smoking was treated as a five-level categorical variable, as described in detail previously⁷.

Cardiovascular disease_(CVD) comorbidities that may be associated with both history of airway obstruction and being edentulous were also examined using data from Visit 4. Other potential confounders examined included race and examination center, ethnicity, education level, income, frequency of dental visits, diabetes, body mass index, alcohol use, and medication utilization. These data were extracted from Visit 4.

Data Analysis

Multivariate logistic regression was used to examine the association between COPD and edentulous status. This was followed by a logistic regression analysis of the dentate subjects comparing the COPD prevalence and severity of dentate patients without periodontal disease to that of dentate patients with periodontal disease. The exposure variable, which is airway obstruction, was examined as both a dichotomous variable (e.g., positive prior COPD vs. no COPD) and by logistic models using nominal variables (e.g., no lung disease, restrictive disease, GOLD 0, Stage I, Stage II, Stage III, and Stage IV) based upon spirometry data collected as part of the ARIC examination. For the no COPD or GOLD reference group we included: subjects with and without chronic respiratory symptoms with no lung function abnormality, and subjects with GOLD 0 and restrictive disease. Multiple regression models were used to control for confounding. When adjusting for potential confounding variables we included in certain models variables that are known to modify either periodontal disease status or COPD, including Body mass index and smokeless tobacco. We developed minimally and fully-adjusted logistical models incorporating significant confounders and effect modifiers to compute odds ratios (OR) and 95% confidence intervals (CI). SAS Version 9.2 (Cary, NC). We also included traditional risk factors for

1
2
3 edentulism, periodontal disease and COPD- such as age, even if not statistically significant in this cohort.
4
5 A p-value of <0.05 was considered statistically significant for this study.
6
7
8
9

10 **Sample Size and Power Calculations**

11 We knew *a priori* that the ARIC cohort who had spirometry data from Visit 2 included 2,087 edentulous
12 subjects, as determined by questionnaire data at Visit 4. We also had 11,378 dentate subjects with
13 spirometry data at Visit 2, of which 5,155 subjects had a full-mouth periodontal exam completed. Thus, if
14 approximately 9% of the 13,465 subjects (2,087 edentulous and 11,378 dentate) were determined to have
15 COPD, we estimated that we had >99% power to detect a significant difference of 1% at p=0.05 in the
16 prevalence of COPD when comparing edentulous vs. dentate subjects.
17
18
19
20
21
22
23
24
25
26

27 **RESULTS**

28 ARIC subject participation was based upon the inception cohort of 15,792 subjects. The Visit 4
29 screenings were broadly divided into 11,378 dentate and 2,087 edentulous subjects who also had
30 spirometry data. These subjects are used for analyses in Tables 1-6 on the relationship between dentate
31 status and COPD. Subjects were excluded (n=2,327) from the original inception participants for
32 incomplete data, dental implants only, and missing Visit 4 screening. Table 7 is based upon the 8,274
33 subjects eligible for the oral exam, of which 6967 (84.1%) were actually examined. Of those examined,
34 the periodontal portion of the exam was not completed on 174 subjects, resulting in 6,793 subjects with
35 full periodontal data.
36
37
38
39
40
41
42
43
44
45
46
47
48

49 The distribution of study subjects according to dentate status and GOLD classification is shown in Table
50 1. Among the edentulous subjects (2087), there were 591 or 28.3% with COPD (Gold Stages I-IV). With
51 the exception of GOLD I where the prevalence of edentulism was 14.7 percent, the prevalence of
52 edentulous individuals increased from 16.4 percent for GOLD 0 to 35.6% for GOLD III/IV. . The crude
53 odds ratio (not shown in Table 1) for edentulism being associated with prior COPD vs no COPD was 1.62
54
55
56
57
58
59
60

1
2
3 (95% CI, 1.46-1.80). Approximately 13 percent of individuals with no chronic respiratory symptoms and
4 no lung function abnormality (i.e. no lung disease) were edentulous. There were 2232 subjects with a
5 prior COPD (19.6%) among 11,378 dentate subjects.
6
7

8
9
10 Table 2 presents the prevalence of various subject characteristics according to GOLD classification. In
11 this table, the GOLD reference group for statistical testing combines the following three categories of
12 subjects: no lung disease, GOLD 0, and subjects with chronic respiratory symptoms but no lung function
13 abnormality. The general patterns reflected in Table 2 show that compared to the reference group those
14 subjects in the GOLD III-IV category were much more likely to be edentulous, to smoke, to use
15 smokeless tobacco, to be African American or North Carolina Caucasians, male, and somewhat more
16 likely to have lower levels of income and education, be older and more likely to have chronic heart failure
17 and heart disease.
18
19
20
21
22
23
24
25
26
27
28

29
30 Table 3 presents multivariate logistic models for the relationship between edentulous /dentate status and
31 GOLD stages using the GOLD reference group described above. In an unadjusted model individuals with
32 GOLD II and GOLD III/IV scores were at significantly higher odds for edentulism that increased with
33 increasing GOLD stage severity (OR=2.06; 95% CI 1.79-2.38 and OR=3.38; 95% CI 2.61-4.39,
34 respectively). Minimally adjusted models (race/center, sex and age) showed similar significant effects for
35 GOLD II and GOLD III/IV with little attenuation. The increased risk of being edentulous for GOLD II
36 and GOLD III/IV categories remain significant, but have slightly lower odds ratios. However, the fully
37 adjusted model showed more attenuation with the odds for being edentulous being 1.3 (1.08-1.62) and 2.5
38 (1.68-3.63) times greater for GOLD II disease and the most severe category GOLD III/IV, respectively.
39
40
41
42
43
44
45
46
47
48
49
50

51 Table 4 shows the association between periodontal status and GOLD classifications for the dentate
52 subjects. Because of small cell sizes in the GOLD III and IV groups, we combined them with GOLD II.
53 Using the CDC/AAP classification of periodontal disease, which is based upon clusters of clinical signs,
54 there is a significant positive association between higher GOLD stage and severe periodontal disease
55
56
57
58
59
60

1
2
3 (p<0.0001). Subjects with GOLD I or GOLD II-IV were more likely to have severe periodontitis than the
4 reference group. With respect to the individual periodontal variables measured as means or extent scores,
5 individuals with a GOLD classification of I were more likely to have a greater extent of periodontal
6 pocket depths of 4mm or more, greater extent of attachment loss of 3mm. or more, a higher mean
7 attachment loss and a greater extent of plaque in their mouths than the reference group. Those with
8 GOLD scores of II-IV were significantly higher on all periodontal measures and had fewer teeth
9 compared to the reference group.
10
11
12
13
14
15
16
17
18
19

20
21 Logistic models for the association between GOLD status and periodontal status using the AAP/CDC
22 definitions appear in Table 5. As in Table 4, small cell size resulted in grouping the more severe GOLD
23 stages together as GOLD II/III/IV. In general, the odds ratios for severe periodontal disease increased
24 with more severe GOLD stage in the crude, minimally adjusted and fully adjusted models. There is also
25 an increase in odds as the severity of periodontal disease increases for the two GOLD categories shown
26 using the periodontally healthy group as the reference group within each GOLD category. Although
27 adjusting for relevant confounders generally decreased the odds, there was still a statistically significant
28 association between higher GOLD stage and severe periodontal disease in the fully adjusted model
29 (OR=1.41; 95% CI 1.07-1.87).
30
31
32
33
34
35
36
37
38
39
40
41

42 The relationship between spirometry based GOLD classification and self-reported physician-based
43 diagnosis of bronchitis or emphysema is shown in Table 6. Participants classified as GOLD III/IV were
44 significantly more likely to have self-reported bronchitis or emphysema. There is a trend for a positive
45 association except for the GOLD zero category. Interestingly, 65.9% of subjects with GOLD III/IV
46 report no diagnosis of COPD suggesting that self-report of bronchitis or emphysema is not a strong
47 predictor of GOLD status. In fact, of all subjects reporting a positive diagnosis of bronchitis or
48 emphysema 53% demonstrated no evidence of chronic respiratory symptoms and no lung function
49 abnormality.
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

We found a statistically significant association between prior COPD and edentulism, with evidence of an increasing prevalence of edentulism with increasing GOLD classification. Although there was some reduction in the magnitude of effect in the adjusted models, the association between dentate status and GOLD classification remained statistically significant even in the fully adjusted model, with a 1.3 fold increased odds of being edentulous in the GOLD II category and a 2.5 fold increased odds of being edentulous in the most severe GOLD III/IV category in the fully adjusted regression model. This represents the first report of an association between edentulism and COPD in a population with GOLD measurements.

In the subgroup of dentate subjects, there was a significant association between the GOLD II - IV group vs. the GOLD reference group and prevalent periodontal disease with a statistically significant increase in the odds of entry level and severe periodontal disease in the crude and minimally adjusted models. In the fully adjusted model, only the association between GOLD stage II/III/IV and severe periodontal disease [OR 1.41 (1.07-1.87)] remained statistically significant. This is the first population report to include both full-mouth periodontal examination data as well as Spirometry assessments, thereby providing direct confirmation of the association between severe periodontal disease and COPD, as well as a dose-response relationship between these conditions.

While there is no previous published information available about the association between edentulism status and COPD, there is evidence in the literature of an association between periodontal disease and COPD, consistent with the findings of the current study^{5,6,7}. The results of this investigation were based upon standardized pulmonary function tests and suggest that studies that are based upon self-reported

1
2
3 diagnoses can potentially result in statistically significant misclassification error. Our data shown in Table
4
5 6 suggest that of the 12,297 subjects reporting no diagnosis, 2,390 subjects or 19.4% were classified as
6
7 GOLD I-IV. Among those 433 subjects with GOLD I-IV the positive predictive rate was only 15.3%.
8
9 These findings suggest that an important strength of the current study was the classification of disease
10
11 based on an objective biological measure (e.g., spirometry) rather than a self-reported measure.
12
13

14
15
16 Additional strengths of the current study include the large sample size, the inclusion of African
17
18 Americans providing some racial diversity of the sample, the use of a full-mouth examination protocol
19
20 and the collection of information about key confounding variables, including but not limited to: tobacco
21
22 use, hypertension, diabetes, and income. Furthermore, the outcomes of interest, edentulism and
23
24 periodontal disease, were evaluated using a comprehensive oral examination at Visit 4.
25
26

27
28
29 The current study must also be considered in light of certain limitations. First, the design of the current
30
31 study does not allow us to determine the directionality or temporality of any association found. It has been
32
33 reported previously that the dental ARIC sub-sample is healthier than the greater ARIC sample and that
34
35 the edentulous subjects within the ARIC dataset have overall poorer health,⁹ as compared to the ARIC
36
37 dental sample. Thus, it is possible that the observed relationship between both edentulism and periodontal
38
39 disease with prior COPD may be due in part to shared risk factors. There is also the potential bias due to
40
41 differential loss to follow-up prior to Visit 4. The current study included only those subjects with dental
42
43 history information at Visit 4. Finally, it is possible that participants with COPD may have been more
44
45 likely to be lost to follow-up (compared to participants without COPD) or that participants with more
46
47 severe COPD may have had a lower probability of surviving to Visit 4 than participants with mild COPD
48
49 or normal lung function. There are no data in this report to mechanistically link edentulism with
50
51 increased risk for COPD. Previous studies of periodontal disease associations with COPD have discussed
52
53 the potential role of periodontal infection serving as a chronic repository of pathogenic oral organisms
54
55 that could be aspirated into the airway to challenge the lungs⁶. Periodontal infections are also associated
56
57
58
59
60

1
2
3 with increases in systemic markers of inflammation including biomarkers such as C-reactive protein
4 (CRP)¹³. By analogy, edentulism is commonly associated with denture infections (denture mucositis)
5
6 which induce mucosal inflammation and unclean dentures can also pose as a chronic reservoir of oral
7
8 pathogens. However, the potential role of denture-based infection as a potential risk factor for COPD
9
10 remains to be elucidated.
11
12

13 14 15 16 17 18 **CONCLUSIONS** 19

20
21
22 Edentulism was associated with prior COPD, with an incremental effect observed with increasing GOLD
23 classification. Additionally, among dentate individuals, severe periodontal disease was associated with
24 GOLD classification. This is one of the first analyses of COPD and dentate status, and expands upon the
25 available literature regarding periodontal disease and COPD. Further research is warranted to explore the
26 relationship of edentulism and periodontal disease with COPD.
27
28
29
30
31
32

33 34 35 **Conflict of Interest** 36

37
38 There are no potential conflicts of interest that exist with any companies/organizations whose products or
39 services may be discussed in this article.
40
41

42 43 **Source of Funding** 44

45 This work was supported by a grant from GlaxoSmithKline and we declare that the sponsor had no role in
46 the study design, collection, analysis and interpretation of data; in the writing of the manuscript; and in
47 the decision to submit the manuscript for publication.
48

49 50 **Responsibility for Data** 51

52 This is to verify that we have full access to all the data in this study and take full responsibility for the
53 integrity of the data and accuracy of the data analysis, including and especially any adverse effects.
54
55
56
57
58
59
60

Abbreviations:

AAP/CDC – American Association of Periodontology/Center for Disease Control

ARIC – Atherosclerosis Risk in Communities

CHF – Congestive Heart Failure

CI – Confidence Intervals

COPD – Chronic Obstructive Pulmonary Disease

CVD – Cardiovascular Disease

D-ARIC – Dental Atherosclerosis Risk in Communities

FEV – Forced Expiratory Volume

FVC – Forced Vital Capacity

GOLD – Global Initiative for Chronic Obstructive Lung Disease

MAL – Mean Attachment Loss

NHANES – National Health and Nutrition Examination Survey

OR – Odds Ratio

SES – Social Economic Status

WHO – World Health Organization

REFERENCES

1. Wouters EFM. Economic analysis of the confronting COPD survey: an overview of results. *Respiratory Medicine* 2003;**97**(C): S3-S14.
2. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006;**3**(11):e442.
3. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med*. 2007;**176**(6):532-55.
4. van Eeden, SF, Sin DD. Chronic Obstructive Pulmonary Disease: A Chronic Systemic Inflammatory Disease. *Respiration* 2008;**75**(2):224-38.
5. Scannapieco FA, Ho AW. Potential associations between chronic respiratory disease and periodontal disease: analysis of National Health and Nutrition Examination Survey III. *J Periodontol* 2001;**72**(1):50-6.
6. Scannapieco FA, Papandonatos GD, Dunford RG. Associations between oral conditions and respiratory disease in a national sample survey population. *Ann Periodontol* 1998;**3**(1):251-6.
7. Garcia RI, Nunn ME, Vokonas PS. Epidemiologic associations between periodontal disease and chronic obstructive pulmonary disease. *Ann Periodontol* 2001;**6**(1):71-7.
8. Glass RT, Conrad RS, Bullard JW, Goodson LB, Mehta N, Lech SJ et al. Evaluation of microbial flora found in previously worn prostheses from the Northeast and Southwest regions of United States. *Journal of Prosthetic Dentistry* 2010;**103**(6):384-9.
9. Elter JR, Offenbacher S, Toole JF, Beck JD. Relationship of periodontal disease and edentulism to stroke/TIA. *J Dent Res* 2003;**82**(12):998-1001.
10. Page RC, Eke PI. Case definitions for use in population-based surveillance of periodontitis. *J Periodontol* 2007;**78**(7 Suppl): 1387-99.
11. Pauwels RA, Buist AS, Calverley PMA, Jenkins CR, Hurd SS. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop Summary. *American Journal of Respiratory and Critical Care Medicine* 2001;**163**(5): 1256-76.
12. Mannino DM, Doherty DE, Buist SA. Global Initiative on Obstructive Lung Disease (GOLD) classification of lung disease and mortality: findings from the Atherosclerosis Risk in Communities (ARIC) study. *Respiratory Med* 2006;**100**(1):115-22.
13. Slade GD, Ghezzi EM, Heiss G, Beck JD, Riche E, Offenbacher S. Relationship between Periodontal Disease and C-Reactive Protein among Adults in the Atherosclerosis Risk in Communities Study. *Arch Intern Med*. 2003;**163**(10):1172-9.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 1. Dentate Status by GOLD Stage

	No Chronic Respiratory Symptoms & No Lung Function Abnormality	Restrictive Disease (FEV ₁ /FVC≥0.7 and FVC<80% predicted)	GOLD 0 (Chronic Respiratory Symptoms & No Lung Function Abnormality)	GOLD I	GOLD II	GOLD III/IV	p-value*
Edentulous	1294 (13.3%)	151 (24.6%)	51 (16.4%)	208 (14.7%)	290 (25.2%)	93 (35.6%)	
Dentate	8422 (86.7%)	464 (75.5%)	260 (83.6%)	1204 (85.3%)	860 (74.8%)	168 (64.4%)	<0.0001

*Chisq

For peer review only

Table 2. Demographics by GOLD Stage

	GOLD Reference*	GOLD I	GOLD II	GOLD III/IV	p-value**
Edentulous	1496 (14.1%)	208 (14.7%)	290 (25.2%)	93 (35.6%)	
Dentate	9146 (85.9%)	1204 (85.3%)	860 (74.8%)	168 (64.4%)	<0.0001
Race/Center					
MS African Americans	2654 (25.1%)	191 (13.6%)	196 (17.1%)	73 (28.4%)	
NC African Americans	285 (2.7%)	35 (2.5%)	27 (2.4%)	11 (4.3%)	
NC White	2211 (20.9%)	401 (28.5%)	344 (30.0%)	84 (32.7%)	
Washington Co, MD	2680 (25.3%)	350 (24.9%)	337 (29.4%)	56 (21.8%)	
Suburban Minneapolis	2747 (26.0%)	430 (30.6%)	243 (21.2%)	33 (12.8%)	<0.0001
Female	6344 (59.6%)	684 (48.4%)	457 (39.7%)	125 (47.9%)	
Male	4298 (40.4%)	728 (51.6%)	693 (60.3%)	136 (52.1%)	<0.0001
Mean Age (StdDev)	62.4 (5.59)	64.2 (5.69)	64.9 (5.57)	64.2 (5.70)	<0.0001
Mean BMI (StdDev)	29.2 (5.68)	27.1 (4.81)	27.6 (5.34)	27.3 (6.16)	<0.0001
Diabetes (Yes)	1573 (17.4%)	132 (10.8%)	179 (19.7%)	36 (20.7%)	
No	7486 (82.6%)	1090 (89.2%)	732 (80.4%)	138 (79.3%)	<0.0001
Hypertension (Yes)	3591 (39.4%)	388 (31.9%)	374 (40.7%)	74 (41.8%)	
No	5535 (60.7%)	829 (68.1%)	545 (59.3%)	103 (58.2%)	<0.0001
Income (Low)	2718 (31.1%)	365 (31.2%)	321 (37.2%)	72 (41.6%)	
Medium	3071 (35.2%)	440 (37.6%)	316 (36.7%)	59 (34.1%)	
High	2945 (33.7%)	366 (31.3%)	225 (26.1%)	42 (24.3%)	<0.0001
Education (Basic)	2226 (21.0%)	291 (20.6%)	332 (28.9%)	93 (35.8%)	
Intermediate	4360 (41.0%)	564 (40.0%)	480 (41.8%)	101 (38.9%)	
Advanced	4038 (38.0%)	556 (39.4%)	336 (29.3%)	66 (25.4%)	<0.0001
Alcohol Use (Yes)	4404 (48.4%)	657 (53.9%)	463 (51.1%)	78 (44.3%)	
No	4695 (51.6%)	561 (46.1%)	444 (49.0%)	98 (55.7%)	0.001
Smoke (Current, Heavy)	748 (8.6%)	191 (16.4%)	256 (29.8%)	33 (20.6%)	
Current, Light	245 (2.8%)	34 (2.9%)	15 (1.7%)	1 (0.6%)	
Former, Heavy	1304 (14.9%)	309 (26.5%)	339 (39.4%)	76 (47.5%)	
Former, Light	1981 (22.7%)	228 (19.6%)	100 (11.6%)	17 (10.6%)	
Never	4448 (51.0%)	403 (34.6%)	150 (17.4%)	33 (20.6%)	<0.0001
Smokeless Tobacco (Yes)	699 (7.7%)	131 (10.8%)	138 (15.2%)	34 (19.3%)	
No	8401 (92.3%)	1088 (89.3%)	769 (84.8%)	142 (80.7%)	<0.0001
Coronary Heart Disease (Yes)	718 (6.9%)	106 (7.6%)	159 (14.2%)	31 (13.0%)	
No	9727 (93.1%)	1281 (92.4%)	963 (85.8%)	208 (87.0%)	<0.0001
Congestive Heart Failure (Yes)	191 (2.1%)	25 (2.1%)	41 (4.6%)	9 (5.1%)	
No	8878 (97.9%)	1194 (98.0%)	861 (95.5%)	166 (94.9%)	<0.0001

*GOLD Reference: No Chronic Respiratory Symptoms & No Lung Function Abnormality + GOLD Zero (Chronic Respiratory Symptoms & No Lung Function Abnormality) + Restrictive (FEV1/FVC \geq 0.70 and FVC < 80% predicted)

**Chisq or t-test

Table 3: Generalized Logits Models: Association between edentulism and GOLD Stages as compared to dentate subjects

	GOLD Reference*	GOLD I	GOLD II	GOLD III/IV
Model 1 Crude				
Edentulous	Reference	1.06 (0.90-1.24)	2.06 (1.79-2.38)	3.38 (2.61-4.39)
Model 2 Minimally Adjusted**				
Edentulous	Reference	1.12 (0.93-1.34)	1.92 (1.61-2.28)	3.17 (2.28-4.41)
Model 3 Fully Adjusted ****				
Edentulous	Reference	1.05 (0.86-1.28)	1.32 (1.08-1.62)	2.47 (1.68-3.63)

*GOLD Reference: No Chronic Respiratory Symptoms & No Lung Function Abnormality + GOLD Zero (Chronic Respiratory Symptoms & No Lung Function Abnormality) + Restrictive (FEV1/FVC \geq 0.70 and FVC < 80% predicted)

**Model 2 is minimally adjusted for race/center (5-level), male, age (years)

***Model 3 is fully adjusted for race/center (5-level), male, age (years), diabetes, hypertension, smoking (5-levels), smokeless tobacco, alcohol, income (3-level), education (3-level), CFH and CHD

Table 4. Periodontal measures among the 5,155 dentate subjects by GOLD stages

	GOLD Reference*	GOLD I	GOLD II/III/IV	Overall p-value
CDC/AAP Definition				
Healthy N (SD)	2378 (43.0%)	281 (38.1%)	151 (28.5%)	
Entry N (SD)	2271 (41.1%)	295 (40.0%)	233 (44.1%)	
Severe N (SD)	878 (15.9%)	161 (21.9%)	145 (27.4%)	<0.0001
Mean (StdDev)				
Extent PD \geq 4mm	7.05 (11.3)	8.12 (12.3) ^a	9.66 (14.8) ^d	<0.0001
Mean PD	1.89 (0.56)	1.94 (0.61)	2.03 (0.74) ^d	<0.0001
Extent AL \geq 3mm	21.7 (21.8)	27.6 (26.0) ^d	34.3 (28.7) ^d	<0.0001
Mean AL	1.72 (0.96)	1.94 (1.15) ^d	2.29 (1.43) ^d	<0.0001
Extent BOP	25.2 (23.4)	25.1 (24.3)	27.7 (26.4) ^a	0.07
Extent PQ \geq 1	42.5 (38.3)	39.3 (37.7) ^a	49.9 (39.3) ^d	<0.0001
Number of Teeth	22.0 (6.99)	21.5 (7.26)	18.8 (8.09) ^d	<0.0001

^a = $p < 0.05$, ^b = $p < 0.01$, ^c = $p < 0.001$, ^d = $p < 0.0001$,

All superscripted p values are compared to reference group by Chi square or GLM

Table 5: Generalized Logits Models: Periodontal status (CDC/AAP Definition) by GOLD stages

	GOLD I	GOLD II/III/IV
Model 1 Crude		
Healthy	Reference	Reference
Entry	1.10 (0.92-1.31)	1.62 (1.31-2.00)
Severe	1.55 (1.26-1.91)	2.60 (2.05-3.31)
Model 2 Minimally Adjusted*		
Healthy	Reference	Reference
Entry	0.99 (0.83-1.19)	1.42 (1.14-1.77)
Severe	1.46 (1.17-1.82)	2.17 (1.67-2.80)
Model 3 Fully Adjusted **		
Healthy	Reference	Reference
Entry	0.95 (0.79-1.14)	1.10 (0.87-1.39)
Severe	1.23 (0.97-1.55)	1.41 (1.07-1.87)

*Model 2 is minimally adjusted for race/center (5-level), male, age (years)

**Model 3 is fully adjusted for race/center (5-level), male, age (years), diabetes, hypertension, smoking (5-levels), smokeless tobacco, alcohol, income (3-level), education (3-level)

Table 6: Self reported physician diagnosis of bronchitis or emphysema according to GOLD stages .


	No Chronic Respiratory Symptoms & No Lung Function Abnormality	GOLD Zero (Chronic Respiratory Symptoms & No Lung Function Abnormality)	Restrictive Disease (FEV₁/FVC\geq0.7 and FVC$<$80% and predicted)	GOLD I	GOLD II	GOLD III/IV	p-value^a
Self Report (Yes)	633 (6.5%)	65 (20.9%)	64 (10.4%)	143 (10.1%)	201 (17.5%)	89 (34.1%)	
No	9083 (93.5%)	246 (79.1%)	551 (89.6%)	1269 (89.7%)	949 (82.5%)	172 (65.9%)	<0.0001

a) Chi square

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Participant Flow Diagram



129x90mm (300 x 300 DPI)

view only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Word Count: 29342763

**OBSTRUCTIVE AIRWAYLUNG DISEASE AND EDENTULISM IN THE ATHEROSCLEROSIS
RISK IN COMMUNITIES (ARIC) STUDY**

Running Title: COPD and Edentulism

Keywords: pulmonary disease, edentulism, chronic obstructive pulmonary disease, bronchitis

Steven Offenbacher, DDS¹, James D Beck, PhD², Silvana P Barros, DDS¹, Robert Suruki ,ScD³ and Zvi
G Loewy, PhD⁴

¹Department of Periodontology, School of Dentistry, University of North Carolina , Chapel Hill, NC
27599

²Department of Dental Ecology, School of Dentistry, University of North Carolina, Chapel Hill, NC
27599

³Worldwide Epidemiology, GlaxoSmithKline R&D, Research Triangle Park, NC 27709-3398

⁴Department of Pharmaceutical and Biomedical Sciences, Touro College of Pharmacy, New York, NY
10027 and Department of Biochemistry & Molecular Biology, New York Medical College, Valhalla, NY,
10595

*Address correspondence to

Steve Offenbacher, DDS, PhD, MMSc

Department of Periodontology, UNC School of Dentistry

North Carolina Oral Health Institute

PO Box 14290

Durham, NC 27709

Phone: (919)425-3597 Fax: (919)425-3530

Email: steve_offenbacher@dentistry.unc.edu

Abstract:

Objectives: We examined the potential association between prior COPD and edentulism, and whether the association varied by COPD severity using data from the Dental ARIC Study.

Design: Cross-sectional ~~Case-control~~

Setting: Community dwelling subjects from four US communities.

Participants and Measurements: Cases were identified as edentulous (without teeth) and subjects with one or more natural teeth were identified as dentate. COPD cases were defined by spirometry measurements that showed the ratio of forced expiratory volume (one second) to vital capacity to be less than 0.7. The severity of COPD cases was also scored using a modified Global Initiative for Chronic Obstructive Lung Disease classification criteria (GOLD scoring I-IV). Multiple-logistic regression was used to examine the association between COPD and edentulism, while adjusting for age, gender, center/race, ethnicity, education level, income, diabetes, hypertension, coronary heart disease and congestive heart failure, body mass index, smoking, smokeless tobacco use and alcohol consumption.

Results: 13,465 participants were included in this analysis (2,087 edentulous; 11,378 dentate). Approximately 28.3% of edentulous participants had prior COPD compared to 19.6% among dentate participants ($p < 0.0001$). After adjustment for potential confounders, we observed a 1.3(1.08-1.62) and 2.5(1.68-3.63) fold increased risk of edentulism among GOLD II and GOLD III/IV COPD, respectively, as compared to the non-COPD/dentate referent. Given the short period of time between the measurements of COPD (Visit2) and dentate status (Visit4) relative to the natural history of both diseases, neither temporality nor insight as to the directionality of the association can be ascertained.

Conclusion: We found a statistically significant association between prior COPD and edentulism, with evidence of a positive incremental effect seen with increasing GOLD classification.

INTRODUCTION

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Chronic obstructive pulmonary disease (COPD) is currently the fifth leading cause of death world-wide, and is expected to rise to fourth place in 2030.¹ Although COPD has a heterogeneous clinical presentation primarily characterized by airflow limitation, generally associated with an abnormal inflammatory response of the lungs², it is also often associated with a significant systemic inflammatory response³ which has been correlated with adverse clinical effects^{3,4}.

Over the past decade there has been an increased interest in the link between respiratory diseases and oral infection. Higher levels of tooth-associated microorganisms (plaque) and periodontal disease have been associated with COPD^{5,6,7}. Recently, in a very comprehensive oral microbiology study, high levels of diverse microbes- including several implicated in COPD and denture stomatitis- were identified on previously worn dentures⁸. Thus both the teeth and the dentures are non-desquamating oral surfaces that allow the oral biofilm to emerge eliciting periodontal disease or denture stomatitis, respectively. This raises the question as to whether denture-associated microbes may contribute to COPD in a manner analogous to that similar to periodontal organisms. Because oral infections and denture-based sources of sepsis are both treatable and preventable, the potential association of high microbial burden of oral origin to COPD may represent modifiable risk factors that may potentially improve respiratory-related morbidity. It has been suggested by Garcia and colleagues⁷ that the observed association between oral health and COPD may be due to aspiration of oral pathogens, or a common underlying host susceptibility trait that places individuals at risk for both conditions such as an exaggerated inflammatory response that is triggered by oral microbes.

Although numerous studies have examined the association between oral health and respiratory diseases in dentate individuals, the published literature are scant with respect to studies that have examined the association between edentulism and COPD status/severity. We undertook the current study to examine the potential association between COPD and edentulism using data from the Atherosclerosis Risk in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Communities (ARIC) Study. We hypothesized that edentulism would be associated with a diagnosis of COPD and display a higher prevalence with more spirometry categories of COPD. These data include not only dentate status and full-mouth periodontal examination measurements, but also includes results from direct pulmonary function tests and symptomology assessments that enabled us to classify the levels of COPD present.

METHODS

We sought to test the hypothesis that the prevalence of edentulism was greater among individuals with a reported history of lung disease, (i.e. COPD) and that the association with edentulism varied by COPD severity as determined by a modified Global Initiative for Chronic Obstructive Lung disease (GOLD) criteria. Our second hypothesis was that among dentate subjects there was a difference in the prevalence of periodontal disease according to the severity of COPD as measured by GOLD criteria.

Study Population and Study Design

The entire cohort from the ARIC study was 15,792 and the available dental cohort Dental Atherosclerosis Risk in Communities (D-ARIC) data were used to study participants with both a periodontal examination and completed spirometry data. The cohort selection process and reasons for exclusion appear in Figure 1. Descriptions of the cohort and the periodontal examination procedure have been described previously⁹. Eligible participants were all aged 45-64 at study entry (i.e., Visit 1) and had spirometry measurements at Visit 2 (3 years later) and periodontal examinations at D-ARIC Visit 4 (6 years from visit 2).

Case Definitions

Edentulism case definitions were defined as study participants without any natural teeth or implants. Participants with one or more natural teeth (comprising 11,378 subjects) were selected as the dentate subjects. The absence, presence or severity of periodontal disease among dentate subjects was determined

using the joint American Association of Periodontology/Center for Disease Control (AAP/CDC) consensus definitions of Health, Early and Advanced disease.¹⁰

This investigation represents the first report that uses actual pulmonary function data to assess lung function as an exposure related to edentulism or periodontal disease. The measured exposure of pulmonary function was classified both as a dichotomous (e.g., COPD vs. no COPD) and ordinal categorical variable (e.g., no lung disease, ~~restrictive~~~~restricted~~ disease, COPD GOLD Stage I, Stage II, Stage III, and Stage IV), using the following criteria.

Airway obstruction

Lung Obstruction

Dichotomous variable (COPD/no COPD)—Prior ~~airway~~~~lung~~ obstruction (or COPD) was defined based on spirometry measurements collected during Visit 2. Participants with a ratio of forced expiratory volume in one second (FEV₁) and forced vital capacity (FVC) less than 0.7 were determined to have ~~airway~~~~lung~~ obstruction consistent with COPD.

Ordinal categorical variable (COPD GOLD Stage)—Severity of COPD was also examined in this study among COPD participants using a modification of the spirometry-based criteria for severity outlined in the GOLD guidelines.^{11,12}

GOLD Stage	Spirometry Criteria
No lung disease	FEV ₁ /FVC ≥ 0.7 and FEV ₁ ≥ 80% predicted
Restrictive Restricted	FEV ₁ /FVC ≥ 0.7 and FVC < 80% predicted
Stage 0	Presence of respiratory symptoms in the absence of any lung function abnormality; and no lung disease
Stage I	FEV ₁ /FVC < 0.7 and FEV ₁ ≥ 80% predicted
Stage II	FEV ₁ /FVC < 0.7 and 50% ≤ FEV ₁ < 80% predicted
Stage III	FEV ₁ /FVC < 0.7 and 30% ≤ FEV ₁ < 50% predicted

Stage IV	FEV ₁ /FVC<0.7 and FEV ₁ <30% predicted
----------	---------------------------------------------------------------

Variables of Interest

Smoking data (both for cigarettes and cigars) collected during Visit 4 were used to adjust for the confounding effect of smoking on the relationship between airwaylung obstruction (COPD) and being edentulous. Smoking was treated as a five-level categorical variable, as described in detail previously⁷.

Cardiovascular disease (CVD) comorbidities that may be associated with both history of airwaylung obstruction and being edentulous were also examined using data from Visit 4. Other potential confounders examined included race and examination center, ethnicity, education level, income, frequency of dental visits, diabetes, body mass index, alcohol use, and medication utilization. These data were extracted from Visit 4.

Data Analysis

Multivariate logistic regression was used to examine the association between COPD and edentulous status. This was followed by a logistic regression analysis of the dentate subjects comparing the COPD prevalence and severity of dentate patients without periodontal disease to that of dentate patients with periodontal disease. The exposure variable, which is airwaylung obstruction, was examined as both a dichotomous variable (e.g., positive prior COPD vs. no COPD) and by logistic models using nominal variables (e.g., no lung disease, restrictiverestricted disease, GOLD 0, Stage I, Stage II, Stage III, and Stage IV) based upon spirometry data collected as part of the ARIC examination. For the no COPD or GOLD reference group we included: subjects with and without chronic respiratory symptoms with no lung function abnormality, and subjects with GOLD 0 and restrictiverestricted disease. Multiple regression models were used to control for confounding. When adjusting for potential confounding variables we included in certain models variables that are known to modify either periodontal disease status or COPD, including Body mass index and smokeless tobacco. We developed minimally and fully-

1
2
3 adjusted logistical models incorporating significant confounders and effect modifiers to compute odds
4 ratios (OR) and 95% confidence intervals (CI). SAS Version 9.2 (Cary, NC). We also included traditional
5 risk factors for edentulism, periodontal disease and COPD- such as age, even if not statistically significant
6 in this cohort. A p-value of <0.05 was considered statistically significant for this study.
7
8
9
10
11

12 13 14 **Sample Size and Power Calculations**

15 We knew *a priori* that the ARIC cohort who had spirometry data from Visit 2 included 2,087 edentulous
16 subjects, as determined by questionnaire data at Visit 4. We also had 11,378 dentate subjects with
17 spirometry data at Visit 2, of which 5,155 subjects had a full-mouth periodontal exam completed. Thus, if
18 approximately 9% of the 13,465 subjects (2,087 edentulous and 11,378 dentate) were determined to have
19 COPD, we estimated that we had >99% power to detect a significant difference of 1% at p=0.05 in the
20 prevalence of COPD when comparing edentulous vs. dentate subjects.
21
22
23
24
25
26
27
28
29
30

31 **RESULTS**

32 ARIC subject participation was based upon the inception cohort of 15,792 subjects. The Visit 4
33 screenings were broadly divided into 11,378 dentate and 2,087 edentulous subjects who also had
34 spirometry data. These subjects are used for analyses in Tables 1-6 on the relationship between dentate
35 status and COPD. Subjects were excluded (n=2,327) from the original inception participants for
36 incomplete data, dental implants only, and missing Visit 4 screening. Table 7 is based upon the 8,274
37 subjects eligible for the oral exam, of which 6967 (84.1%) were actually examined. Of those examined,
38 the periodontal portion of the exam was not completed on 174 subjects, resulting in 6,793 subjects with
39 full periodontal data.
40
41
42
43
44
45
46
47
48
49
50

51 The distribution of study subjects according to dentate status and GOLD classification is shown in Table
52 1. Among the edentulous subjects (2087), there were 591 or 28.3% with COPD (Gold Stages I-IV). With
53 the exception of GOLD I where the prevalence of edentulism was 14.7 percent, the prevalence of
54
55
56
57
58
59
60

1
2
3 edentulous individuals increased from 16.4 percent for GOLD 0 to 35.6% for GOLD III/IV. . The crude
4
5 odds ratio (not shown in Table 1) for edentulism being associated with prior COPD vs no COPD was 1.62
6
7 (95% CI, 1.46-1.80). Approximately 13 percent of individuals with no chronic respiratory symptoms and
8
9 no lung function abnormality (i.e. no lung disease) were edentulous. There were 2232 subjects with a
10
11 prior COPD (19.6%) among 11,378 dentate subjects.

12
13
14 Table 2 presents the prevalence of various subject characteristics according to GOLD classification. In
15
16 this table, the GOLD reference group for statistical testing combines the following three categories of
17
18 subjects: no lung disease, GOLD 0, and subjects with chronic respiratory symptoms but no lung function
19
20 abnormality. The general patterns reflected in Table 2 show that compared to the reference group those
21
22 subjects in the GOLD III-IV category were much more likely to be edentulous, to smoke, to use
23
24 smokeless tobacco, to be African American or North Carolina Caucasians, male, and somewhat more
25
26 likely to have lower levels of income and education, be older and more likely to have chronic heart failure
27
28 and heart disease.
29
30
31
32

33
34 Table 3 presents multivariate logistic models for the relationship between edentulous /dentate status and
35
36 GOLD stages using the GOLD reference group described above. In an unadjusted model individuals with
37
38 GOLD II and GOLD III/IV scores were at significantly higher odds for edentulism that increased with
39
40 increasing GOLD stage severity (OR=2.06; 95% CI 1.79-2.38 and OR=3.38; 95% CI 2.61-4.39,
41
42 respectively). Minimally adjusted models (race/center, sex and age) showed similar significant effects for
43
44 GOLD II and GOLD III/IV with little attenuation. The increased risk of being edentulous for GOLD II
45
46 and GOLD III/IV categories remain significant, but have slightly lower odds ratios. However, the fully
47
48 adjusted model showed more attenuation with the odds for being edentulous being 1.3 (1.08-1.62) and 2.5
49
50 (1.68-3.63) times greater for GOLD II disease and the most severe category GOLD III/IV, respectively.
51
52
53
54

55
56 Table 4 shows the association between periodontal status and GOLD classifications for the dentate
57
58 subjects. Because of small cell sizes in the GOLD III and IV groups, we combined them with GOLD II.
59
60

1
2
3 Using the CDC/AAP classification of periodontal disease, which is based upon clusters of clinical signs,
4 there is a significant positive association between higher GOLD stage and severe periodontal disease
5 (p<0.0001). Subjects with GOLD I or GOLD II-IV were more likely to have severe periodontitis than the
6 reference group. With respect to the individual periodontal variables measured as means or extent scores,
7 individuals with a GOLD classification of I were more likely to have a greater extent of periodontal
8 pocket depths of 4mm or more, greater extent of attachment loss of 3mm. or more, a higher mean
9 attachment loss and a greater extent of plaque in their mouths than the reference group. Those with
10 GOLD scores of II-IV were significantly higher on all periodontal measures and had fewer teeth
11 compared to the reference group.
12
13
14
15
16
17
18
19
20
21
22
23
24

25 Logistic models for the association between GOLD status and periodontal status using the AAP/CDC
26 definitions appear in Table 5. As in Table 4, small cell size resulted in grouping the more severe GOLD
27 stages together as GOLD II/III/IV. In general, the odds ratios for severe periodontal disease increased
28 with more severe GOLD stage in the crude, minimally adjusted and fully adjusted models. There is also
29 an increase in odds as the severity of periodontal disease increases for the two GOLD categories shown
30 using the periodontally healthy group as the reference group within each GOLD category. Although
31 adjusting for relevant confounders generally decreased the odds, there was still a statistically significant
32 association between higher GOLD stage and severe periodontal disease in the fully adjusted model
33 (OR=1.41; 95% CI 1.07-1.87).
34
35
36
37
38
39
40
41
42
43
44
45
46

47 The relationship between spirometry based GOLD classification and self-reported physician-based
48 diagnosis of bronchitis or emphysema is shown in Table 6. Participants classified as GOLD III/IV were
49 significantly more likely to have self-reported bronchitis or emphysema. There is a trend for a positive
50 association except for the GOLD zero category. Interestingly, 65.9% of subjects with GOLD III/IV
51 report no diagnosis of COPD suggesting that self-report of bronchitis or emphysema is not a strong
52 predictor of GOLD status. In fact, of all subjects reporting a positive diagnosis of bronchitis or
53
54
55
56
57
58
59
60

1
2
3 emphysema 53% demonstrated no evidence of chronic respiratory symptoms and no lung function
4
5 abnormality.
6
7
8
9
10

14 DISCUSSION

16 We found a statistically significant association between prior COPD and edentulism, with evidence of an
17
18 increasing prevalence of edentulism with increasing GOLD classification. Although there was some
19
20 reduction in the magnitude of effect in the adjusted models, the association between dentate status and
21
22 GOLD classification remained statistically significant even in the fully adjusted model, with a 1.3 fold
23
24 increased odds of being edentulous in the GOLD II category and a 2.5 fold increased odds of being
25
26 edentulous in the most severe GOLD III/IV category in the fully adjusted regression model. This
27
28 represents the first report of an association between edentulism and COPD in a population with GOLD
29
30 measurements.
31
32
33
34
35

36 In the subgroup of dentate subjects, there was a significant association between the GOLD II - IV group
37
38 vs. the GOLD reference group and prevalent periodontal disease with a statistically significant increase
39
40 in the odds of entry level and severe periodontal disease in the crude and minimally adjusted models. In
41
42 the fully adjusted model, only the association between GOLD stage II/III/IV and severe periodontal
43
44 disease [OR 1.41 (1.07-1.87)] remained statistically significant. This is the first population report to
45
46 include both full-mouth periodontal examination data as well as Spirometry assessments, thereby
47
48 providing direct confirmation of the association between severe periodontal disease and COPD, as well as
49
50 a dose-response relationship between these conditions.
51
52
53
54

55 While there is no previous published information available about the association between edentulism
56
57 status and COPD, there is evidence in the literature of an association between periodontal disease and
58
59
60

1
2
3 COPD, consistent with the findings of the current study^{5,6,7}. The results of this investigation were based
4 upon standardized pulmonary function tests and suggest that studies that are based upon self-reported
5 diagnoses can potentially result in statistically significant misclassification error. Our data shown in Table
6 suggest that of the 12,297 subjects reporting no diagnosis, 2,390 subjects or 19.4% were classified as
7 GOLD I-IV. Among those 433 subjects with GOLD I-IV the positive predictive rate was only 15.3%.
8 These findings suggest that an important strength of the current study was the classification of disease
9 based on an objective biological measure (e.g., spirometry) rather than a self-reported measure.
10
11

12
13
14 Additional strengths of the current study include the large sample size, the inclusion of African
15 Americans providing some racial diversity of the sample, the use of a full-mouth examination protocol
16 and the collection of information about key confounding variables, including but not limited to: tobacco
17 use, hypertension, diabetes, and income. Furthermore, the outcomes of interest, edentulism and
18 periodontal disease, were evaluated using a comprehensive oral examination at Visit 4.
19
20

21
22
23 The current study must also be considered in light of certain limitations. First, the design of the current
24 study does not allow us to determine the directionality or temporality of any association found. It has been
25 reported previously that the dental ARIC sub-sample is healthier than the greater ARIC sample and that
26 the edentulous subjects within the ARIC dataset have overall poorer health,⁹ as compared to the ARIC
27 dental sample. Thus, it is possible that the observed relationship between both edentulism and periodontal
28 disease with prior COPD may be due in part to shared risk factors. There is also the potential bias due to
29 differential loss to follow-up prior to Visit 4. The current study included only those subjects with dental
30 history information at Visit 4. Finally, it is possible that participants with COPD may have been more
31 likely to be lost to follow-up (compared to participants without COPD) or that participants with more
32 severe COPD may have had a lower probability of surviving to Visit 4 than participants with mild COPD
33 or normal lung function. There are no data in this report to mechanistically link edentulism with
34 increased risk for COPD. Previous studies of periodontal disease associations with COPD have discussed
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the potential role of periodontal infection serving as a chronic repository of pathogenic oral organisms
4 that could be aspirated into the airway to challenge the lungs⁶. Periodontal infections are also associated
5 with increases in systemic markers of inflammation including biomarkers such as C-reactive protein
6 (CRP)¹³. By analogy, edentulism is commonly associated with denture infections (denture mucositis)
7 which induce mucosal inflammation and unclean dentures can also pose as a chronic reservoir of oral
8 pathogens. However, the potential role of denture-based infection as a potential risk factor for COPD
9 remains to be elucidated.
10
11
12
13
14
15
16
17
18
19
20
21
22

23 CONCLUSIONS

24
25
26
27 Edentulism was associated with prior COPD, with an incremental effect observed with increasing GOLD
28 classification. Additionally, among dentate individuals, severe periodontal disease was associated with
29 GOLD classification. This is one of the first analyses of COPD and dentate status, and expands upon the
30 available literature regarding periodontal disease and COPD. Further research is warranted to explore the
31 relationship of edentulism and periodontal disease with COPD.
32
33
34
35
36
37
38
39

40 Conflict of Interest

41
42 There are no potential conflicts of interest that exist with any companies/organizations whose products or
43 services may be discussed in this article.
44
45

46 Source of Funding

47
48 This work was supported by a grant from GlaxoSmithKline and we declare that the sponsor had no role in
49 the study design, collection, analysis and interpretation of data; in the writing of the manuscript; and in
50 the decision to submit the manuscript for publication.
51
52

53 Responsibility for Data

54
55 This is to verify that we have full access to all the data in this study and take full responsibility for the
56 integrity of the data and accuracy of the data analysis, including and especially any adverse effects.
57
58
59
60

Abbreviations:

AAP/CDC – American Association of Periodontology/Center for Disease Control
ARIC – Atherosclerosis Risk in Communities
CHF – Congestive Heart Failure
CI – Confidence Intervals
COPD – Chronic Obstructive Pulmonary Disease
CVD – Cardiovascular Disease
D-ARIC – Dental Atherosclerosis Risk in Communities
FEV – Forced Expiratory Volume
FVC – Forced Vital Capacity
GOLD – Global Initiative for Chronic Obstructive Lung Disease
MAL – Mean Attachment Loss
NHANES – National Health and Nutrition Examination Survey
OR – Odds Ratio
SES – Social Economic Status
WHO – World Health Organization

REFERENCES

1. Wouters EFM. Economic analysis of the confronting COPD survey: an overview of results. *Respiratory Medicine* 2003;**97**(C): S3-S14.
2. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006;**3**(11):e442.
3. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med*. 2007;**176**(6):532-55.
4. van Eeden, SF, Sin DD. Chronic Obstructive Pulmonary Disease: A Chronic Systemic Inflammatory Disease. *Respiration* 2008;**75**(2):224-38.
5. Scannapieco FA, Ho AW. Potential associations between chronic respiratory disease and periodontal disease: analysis of National Health and Nutrition Examination Survey III. *J Periodontol* 2001;**72**(1):50-6.
6. Scannapieco FA, Papandonatos GD, Dunford RG. Associations between oral conditions and respiratory disease in a national sample survey population. *Ann Periodontol* 1998;**3**(1):251-6.
7. Garcia RI, Nunn ME, Vokonas PS. Epidemiologic associations between periodontal disease and chronic obstructive pulmonary disease. *Ann Periodontol* 2001;**6**(1):71-7.
8. Glass RT, Conrad RS, Bullard JW, Goodson LB, Mehta N, Lech SJ et al. Evaluation of microbial flora found in previously worn prostheses from the Northeast and Southwest regions of United States. *Journal of Prosthetic Dentistry* 2010;**103**(6):384-9.
9. Elter JR, Offenbacher S, Toole JF, Beck JD. Relationship of periodontal disease and edentulism to stroke/TIA. *J Dent Res* 2003;**82**(12):998-1001.
10. Page RC, Eke PI. Case definitions for use in population-based surveillance of periodontitis. *J Periodontol* 2007;**78**(7 Suppl): 1387-99.
11. Pauwels RA, Buist AS, Calverley PMA, Jenkins CR, Hurd SS. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD) Workshop Summary. *American Journal of Respiratory and Critical Care Medicine* 2001;**163**(5): 1256-76.
12. Mannino DM, Doherty DE, Buist SA. Global Initiative on Obstructive Lung Disease (GOLD) classification of lung disease and mortality: findings from the Atherosclerosis Risk in Communities (ARIC) study. *Respiratory Med* 2006;**100**(1):115-22.
13. Slade GD, Ghezzi EM, Heiss G, Beck JD, Riche E, Offenbacher S. Relationship between Periodontal Disease and C-Reactive Protein among Adults in the Atherosclerosis Risk in Communities Study. *Arch Intern Med*. 2003;**163**(10):1172-9.