

What is a disease? Perspectives of the public, health professionals and legislators

Kari A O Tikkinen,^{1,2} Janne S Leinonen,³ Gordon H Guyatt,^{1,4} Shanil Ebrahim,¹ Teppo L N Järvinen⁵

To cite: Tikkinen KAO, Leinonen JS, Guyatt GH, *et al.* What is a disease? Perspectives of the public, health professionals and legislators. *BMJ Open* 2012;**2**:e001632. doi:10.1136/bmjopen-2012-001632

► Prepublication history and additional material for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2012-001632>).

Received 9 June 2012
Accepted 15 October 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

For numbered affiliations see end of article.

Correspondence to
Dr Kari AO Tikkinen;
kari.tikkinen@fimnet.fi

ABSTRACT

Objective: To assess the perception of diseases and the willingness to use public-tax revenue for their treatment among relevant stakeholders.

Design: A population-based, cross-sectional mailed survey.

Setting: Finland.

Participants: 3000 laypeople, 1500 doctors, 1500 nurses (randomly identified from the databases of the Finnish Population Register, the Finnish Medical Association and the Finnish Nurses Association) and all 200 parliament members.

Main outcome measures: Respondents' perspectives on a five-point Likert scale on two claims on 60 states of being: '(This state of being) is a disease'; and '(This state of being) should be treated with public tax revenue'.

Results: Of the 6200 individuals approached, 3280 (53%) responded. Of the 60 states of being, $\geq 80\%$ of respondents considered 12 to be diseases (Likert scale responses of '4' and '5') and five not to be diseases (Likert scale responses of '1' and '2'). There was considerable variability in most states, and great variability in 10 ($\geq 20\%$ of respondents of all groups considered it a disease and $\geq 20\%$ rejected as a disease). Doctors were more inclined to consider states of being as diseases than laypeople; nurses and members were intermediate ($p < 0.001$), but all groups showed large variability. Responses to the two claims were very strongly correlated ($r = 0.96$ (95% CI 0.94 to 0.98); $p < 0.001$).

Conclusions: There is large disagreement among the public, health professionals and legislators regarding the classification of states of being as diseases and whether their management should be publicly funded. Understanding attitudinal differences can help to enlighten social discourse on a number of contentious public policy issues.

INTRODUCTION

Disease and illness are related concepts: patients suffer from 'illnesses' and doctors diagnose and treat 'diseases'.¹ Illnesses are experiences of discontinuities in states of

ARTICLE SUMMARY

Article focus

- The concept of disease lies at the heart of medicine.
- No study has addressed perceptions of all relevant stakeholders on what, across a wide range of conditions, should be classified as a disease.

Key messages

- Our survey found large differences in the views among Finnish laypeople, doctors, nurses and parliament members regarding whether states of being should be considered diseases and be managed through public revenue.
- Although doctors were more inclined to consider states of being as diseases, disagreement was as evident among health professionals as in other groups.
- Understanding peoples' attitudes about whether states of being should be considered diseases elucidates fundamental underlying attitudes and thus can inform social discourse regarding a number of contentious public policy issues.

Strengths and limitations of this study

- This is the first study to assess whether states of being should be considered diseases and should be managed through public revenue using a broad sample of doctors, nurses, laypeople as well as legislators.
- Our results from the Finnish population may be less generalisable to less affluent countries and countries with different social and cultural values.

being and perceived role performances; when diagnosed as diseases, they are presumed abnormalities in the function or structure of body systems. Disease can refer to a combination of signs and symptoms, phenomena associated with a disorder of function or structure or illness associated with a specific cause(s).² There are, however, no universally accepted criteria for establishing 'disease'.³⁻⁵ Indeed, the complexity of the concept of disease has led to the observation that it can be as difficult to define as beauty, truth or love.⁶

The concept of disease is subject to social, cultural and economic influences that have varied over time: these influences have been particularly evident in the last two decades.^{4 5 7-9} During this time, we witnessed a growing tendency to classify states of being as diseases, a trend with important possible consequences, both positive and negative.^{8 10-13} Possible positive consequences include the facilitation of patient-physician communication^{4 5 11} and increased willingness to use public money and thus enhance equality in the distribution of limited resources.^{4 14} Possible adverse consequences include making relatively healthy individuals perceive themselves as sick, encouraging misguided attempts to treat states that are part of the normal human condition, and individuals being denied employment or insurance.^{4 11 15-17} The extent to which health workers and the public have been influenced by these tendencies and their current perceptions remain uncertain.

Authors have also suggested that the disease label can be used as a social control mechanism.¹⁸⁻²⁰ The 'sick role' theory suggests that illness disrupts normal social functioning, making the individual responsible for adhering to treatment regimes in order to maintain social productivity.²¹⁻²³ However, the relationship between the patient and the medical sphere exists within a socially constructed hierarchy wherein medical institutions ultimately hold the individual accountable for collective social problems.^{19 21 23} When individual behaviour deviates from pre-established social norms, it is not the individual, but the medical community that labels, diagnoses and treats aberrant behaviour as a socially legitimated health condition.¹⁹

No earlier study assessed perceptions' on use of public funding, and only one study² assessed perceptions' on the concept of disease over wider range of conditions.

Campbell *et al*² found that doctors considered more non-infectious conditions to be diseases than laypeople. Because of the importance of the issue, and the paucity of empirical evidence regarding peoples' views, we conducted a survey of the general public, doctors, nurses and parliament members in Finland to determine the extent to which they considered 60 states of being to be diseases and their attitudes towards using public funds for managing these states. On the basis of differences in background, training, and life experience and underlying attitudes, we hypothesised that groups (laypeople, doctors, nurses and parliament members) would vary in their conceptions of disease, and that there would also be large variation in conceptions of disease within groups.

METHODS


The Finnish Disease (FIND) Survey study population

In 2010, we selected a random sample of 3000 laypeople, 1500 doctors, 1500 nurses and all the 200 members of the Parliament of Finland (MPs). We identified laypeople 18-75 years of age from the Finnish Population Register Centre, and doctors and nurses less than 65 years of age from the registers of the Finnish Medical Association and the Finnish Nurses Association. We excluded individuals who had died, emigrated, were deemed seriously disabled or who changed careers and would therefore no longer be members of their respective group (figure 1).

Survey

Referring to the existing literature and the International Classification of Diseases (ICD-10),^{2 11 24 25} we chose, through iterative discussion and consensus-building, 60 states of being that we considered familiar to the

Figure 1 Study flow. We randomised the 60 states of being into three blocks: version A consisted of three blocks (each consisting of 20 states of being) in the order 1-2-3, version B in the order 3-1-2 and version C in the order 2-3-1.



relevant stakeholders. We anticipated that everyone would consider some of these states a disease, none would consider some states a disease, and that some states might elicit disagreement (see online supplementary figures A1 and A2 in the appendix). We asked participants to respond to two claims: (1) '(This state of being) is a disease' (claim A) and (2) '(This state of being) should be treated with public tax revenue' (claim B) on a five-point Likert scale ranging from *strongly disagree* to *strongly agree* (see online supplementary figures A1 and A2 in appendix). We elicited demographic information using questions from earlier surveys (see online supplementary table A1 in the appendix). We pilot tested the questionnaire with 20 laypeople and five doctors, and made minor revisions on the basis of feedback.

We mailed the questionnaires in June 2010 and sent reminders in August and October 2010. We made pre-contacts with MPs by email and telephone. The ethics committee of the Pirkanmaa Hospital District in Finland granted exemption from ethical review (R11110). The reporting of the study conforms to the STROBE statement.²⁶

Randomisation and exclusion criteria

Each participant received a questionnaire eliciting responses to 60 states of being. We randomised the 60 states of being into three blocks (1, 2 and 3; each containing 20 states). We created three versions of the questionnaire: version A consisted of blocks in the order 1-2-3, version B in the order 3-1-2 and version C in the order 2-3-1. Within each sample group (laypeople, doctors, nurses and MPs), we randomised respondents to the three versions (figure 1).

To check comprehension of the questionnaire, we placed three states (myocardial infarction, pneumonia and breast cancer) likely to be considered as disease as the first state of being in each block. Respondents who did not *agree to some extent* or *strongly agree* to the statement '(This state of being) is a disease' (see online supplementary figures A1 and A2 in appendix) for *any* of these three were deemed unlikely to understand the questionnaire and excluded from the analyses (figure 1).

Statistical analysis

For each group (doctors, nurses, laypeople and MPs), we calculated the proportion of states of being where respondents *strongly agreed* or *agreed to some extent* regarding the two claims. Using a Pearson χ^2 test on all possible pair-wise comparisons (altogether six comparisons for each state of being by claim), we evaluated the order of ratings of the perception of disease and expenditure of public tax revenue claims across groups. We calculated the correlation between the proportions of individuals who either *strongly agreed* or *agreed to some extent* across states in the two claims. All other analyses were descriptive.

RESULTS

Of the 6200 people approached, 3280 (53.2%) participated, of whom 36 proved ineligible (figure 1). Of the 3244 eligible individuals who completed and understood the questionnaire, 3246 (99%) responded to at least 55 of the 60 states of being. Among respondents, the mean (SD) age was: laypeople 49.5 (15.5), doctors 46.1 (10.7), nurses 44.9 (11.3) and MPs 54.4 (9.8). There were significantly more women among nurses (97.3%), and fewer among MPs (35.7%) compared with doctors (61.5%) or laypeople (57.3%) ($p < 0.01$ for all comparisons). We found no significant differences in ratings or background characteristics between questionnaire versions and individuals responding at different response rounds. Online supplementary table A1 in the appendix presents the demographic data.

From the 60 states of being, 12 were perceived as diseases by $\geq 80\%$ of respondents from all groups and five were perceived not to be diseases by $\geq 80\%$ (figure 2 and table 1). Doctors were most likely to consider states of being as diseases followed by nurses, MPs and laypeople ($p < 0.001$ for all pairwise comparisons). For a large number of states, there was extreme disagreement regarding classification as a disease among all study groups (figure 2). In 10 states, $\geq 20\%$ of participants considered them diseases and $\geq 20\%$ did not (table 1). There was a very strong correlation between responses to claims ($r = 0.96$ (95% CI 0.94 to 0.98); $p < 0.001$; no differences between groups) (see online supplementary figure A3 in the appendix).

DISCUSSION

Statement of principal findings

Our survey found large discrepancies in the views among laypeople, doctors, nurses and MPs in Finland regarding whether states of being should be considered diseases and should be managed through public revenue. Although physicians were more inclined to consider states of being as diseases, disagreement was as evident among health professionals as in other groups (figure 2 and table 1). In all groups, willingness to pay for treatment from public funds was very strongly correlated with the perception of disease.

Strengths and limitations

The strengths of our study include a large sample of both healthcare professionals and general population, an acceptable response rate, excellent completeness of questionnaires and a large number of states of being that elicited a wide range of responses. Further, the sample proved similar in its characteristics to the target populations in terms of age and gender distribution, education, employment and marital status (for details, see online supplementary table A1 in the appendix and its supplementary references). We found no trend in the perceptions or participants' characteristics by response round, reducing concern regarding selection bias.




Figure 2 Variation of perceptions in the concept of disease among laypeople, doctors, nurses and members of parliament.

The limitations of our study include concern that the strong correlation between the claims may be partly caused by the positioning of questions adjacent to one another in the questionnaire. Second, these results from

the Finnish population may be less generalisable to less affluent countries and those with different social and cultural values. For instance, the high correlation between the disease label and the willingness to fund

Table 1 (A) States of being perceived as a disease by at least 80% of respondents of all groups, (B) states of being not perceived as a disease by at least 80% of respondents of all groups and (C) states of being perceived as a disease by at least 20% and not as a disease by at least another 20% of respondents of all groups (laypeople, doctors, nurses and parliament members)

(A) Perceived as disease by more than 80% (response options '4' and '5')	
Breast cancer	Schizophrenia
Prostate cancer	HIV/AIDS
Pneumonia	Malaria
Lung cancer	Adult-onset diabetes
Juvenile diabetes	Osteoporosis
Myocardial infarction	Autism
(B) Not perceived as disease by more than 80% (response options '1' and '2')	
Wrinkles	Grief
Smoking	Homosexuality
Ageing	
(C) At least 20% perceived as disease (response options '4' and '5') and at least another 20% did not perceive as disease (response options '1' and '2')	
Premenstrual syndrome, PMS	Age-related muscle loss, sarcopenia
Erectile dysfunction	Female menopause
Gambling addiction	Malnutrition
Infertility	Eye refractive error, need for eyeglasses
Drug addiction	Lactose intolerance

socially may be related to Finland's high level of social solidarity. Finland is said to have a strong welfare state, and the high correlation between claims may not be reproduced in other jurisdictions. Third, despite our attempt to address understanding and the potential impact of wording in a pilot study, there is a possibility that a framing effect (ie, individuals reacting differently to a particular response depending on how the question is worded) may have occurred. There is evidence from various populations illustrating the impact of framing on decision-making and preferences.^{27–29} In particular, this may have been an issue for our claim B, whether states of being should be funded by public revenue; an alternative framing of questions may have elicited different results.³⁰

Comparison with other studies

Some investigators have addressed patients' and health-care providers' perceptions regarding the disease concept and use of public funding in specific conditions.^{31–34} However, no earlier study assessed perceptions' on use of public funding over a wider range of conditions, and only one study assessed perceptions' of the disease concept.² In keeping with our finding that physicians were slightly more likely than others to consider states of being to be diseases, Campbell *et al*² found no difference among non-medical faculty, secondary school students, academic internists and general practitioners on how they perceived illnesses due to infections, but found that doctors considered more non-infectious conditions to be diseases.

In another related investigation, the editorial board of the *BMJ* and its readers identified a list of almost 200 *non-diseases* (defined as 'a human process or problem that some have defined as a medical condition but where people may have better outcomes if the problem or process was not defined in that way') including ageing, baldness and boredom.¹¹ As in our survey, there was considerable variation in the states of being deemed 'non-diseases'.

Meaning of the study: possible explanations and implications

The concept of 'disease' lies at the heart of medicine,^{7 14} defining its domain and its role in public policy, including the range of conditions in which sufferers may be entitled to public funding for their treatment.^{35–37} Building on earlier work,^{4 8 11 13–17 38–48} [table 2](#) presents a taxonomy of states of being, exploring the relation between categorisation –or not – as a disease, the implications for action and potential negative consequences. The issues presented in [table 2](#) are subjects of ongoing, often heated, debate. Our results (ie, large differences in views whether states of being should be considered diseases and should be managed through public revenue) provide insight into these debates: why they are so contentious is due at least in part to disparities in views on the fundamental nature of these states of

being. Our study represents only the first steps in understanding the concept of 'disease'. Additional qualitative studies would be useful for obtaining further insight into interpretation of the findings.

As reflected in [table 2](#), people tend to think of diseases as conditions for which individuals do not bear primary responsibility, afflictions of which the sufferer is at least to some extent a victim.³⁴ Thus, if we view addictions as diseases (which substantial proportions of our respondents did, and did not) we are inclined to look for solutions through harm reduction approaches and medical treatment, and to allocate public funding for these interventions.^{42 48} Alternative views include viewing a condition as a moral failing, bad habit or retribution for bad behaviour (all related perspectives) or as a social problem (a quite different perspective).

For instance, a non-disease perspective on addiction includes two alternatives: If we regard addiction as a moral failing, we are likely to demand personal responsibility for dealing with the problem, and institute punitive approaches for those who fail ([table 2](#)).^{40 42} Alternatively, we may see addiction as a social problem and seek social solutions such as poverty reduction.⁴⁴ The general unavailability of safe injection sites for drug users, despite evidence of benefit and eminent advocacy, illustrates how these issues play out in public policy.⁴⁶ Our results suggest that the current contentious debate on social policy towards addiction could benefit not only from evidence regarding the effectiveness of alternative policies, but a more profound understanding of the biology and sociology of addiction.

To take other examples from [table 2](#) with potentially negative consequences of a disease perspective, viewing social anxiety disorder or fibromyalgia as specific biological problems may lead to overdiagnosis and medical overtreatment, and undertreatment with behavioural approaches.^{15 45 49} On the other hand, seeing these conditions as socially mediated adjustment problem risks stigmatisation and underuse of potentially effective medical treatment.^{15 45 49} For other states of being, the ongoing passionate debate has highlighted possible dangers in medicalising conditions that might be considered normal problems of living.^{14 15 17 31}

We found the association between considering a state of being a disease and readiness to fund treatment through public revenue very strong. If we consider obesity a disease, we might devote public funding to weight loss clinics. While this is true of very few jurisdictions,⁵⁰ most high-income countries devote public funding to bariatric surgery for morbid obesity, a policy which—according to a Danish study³⁴—many laypeople may question despite evidence suggesting that it is cost effective.

Advocates argue that placing a disease label on the absence of sexual desire is a step towards helping people,³⁹ while critics deem it a destructive medicalisation of a normal part of living fostering problematic commercialisation.⁴¹ Similarly, creating new diagnostic terms, such as the concept 'overactive bladder' may help to increase

Table 2 Implications of alternative viewpoints regarding accepting or rejecting states of being as diseases

Categories of states of being <i>Examples</i>	Disease?	Conceptualisation	Implications for action	Potential negative consequences/ramifications
Addictions or possible addictions Alcoholism Drug addiction Gambling addiction Obesity Smoking	Yes No	Biological health disorder Lack of self-control Moral failing Social problem	Harm reduction Public funding Medical treatment Abstinence through individual choice and self-discipline Punitive management strategies Preventive social solutions: income redistribution, poverty reduction, education, social marketing Diagnose and treat, possibly with drugs	Focus on individuals and treatments may cause social and moral aspects to be ignored ^{8 43 44 47} Stigma and discrimination, neglect of harm reduction, neglect of social causes, increased suffering for the population ^{40 42–44 46 48} Effective medical treatment underused ^{42 43}
Medical diagnoses with uncertain biological/ psychosocial basis Chronic fatigue syndrome Fibromyalgia Irritable bowel syndrome Panic disorder Personality disorder	Yes No	Specific biological problem Socially mediated adjustment problem	Diagnose and treat, possibly with drugs Behavioural therapy Modify environment	Overdiagnosis and overtreatment with drugs, undertreatment with behavioural approaches ^{11 15 16 31} Patients may feel stigmatised Effective medical treatment may be underused ^{11 16 49}
Diminished function or altered appearance, often age-related Age-related muscle loss Baldness Erectile dysfunction Lack of sexual desire	Yes No	Biological health disorder Normal consequence of living	Diagnose and treat, possibly with drugs Public funding Accept and adjust Responsibility on individual	Overdiagnosis and overtreatment Medicalisation of society, with increased self-perception of illness and poorer coping with suffering that is part of life ^{11 15–17 49} Neglect of treatments that may reduce suffering and improve function ^{11 16 49}
Patterns of behaviour Homosexuality Obesity Smoking Transsexualism	Yes No No	Biological health disorder Lifestyle choice Moral failing	Diagnose and treat, possibly with drugs Negative social stigma Respect person's choice Abstinence/modification of behaviour through individual choice/self-discipline Punitive strategies	Adverse judgment and resulting stigma and discrimination ⁵³ Permissive attitude encourages self-destructive or morally reprehensible behaviour ^{*43} Underuse of effective treatment ^{*34} Stigma and discrimination ⁵³
Syndromes or constellation of patterns of symptoms of unclear basis Attention deficit hyperactivity disorder Fibromyalgia Overactive urinary bladder Panic disorder	Yes No	Essentialist: specific biological disorder Nominalist: collection of symptoms, signs, behaviours, label of convenience	Label all patients with specific category and treat uniformly Acknowledge syndromes as convenient constructions, seek underlying causes, don't attempt to pigeon-hole unusual presentations	Failure to recognise diversity of illness, excessively uniform management, stifle research that could deepen understanding ^{2 5 51} Acknowledgement of complexity may lead to inefficiency, paralysis ^{2 5 51}

*Negative consequences listed here refer particularly to smoking and obesity not to homosexuality and transexualism.

awareness of the symptoms and to simplify management, but it may also cause problematic oversimplification leading to excessive use of ineffective treatment.^{5 51 52}

This discussion can also be seen from a more general perspective: essentialism versus nominalism (table 2). Essentialists regard diseases as causes of illness; the role of a physician, in this view, is to identify the cause and treat it appropriately.⁵¹ Nominalists see diseases as constructs that humans create to bring order to a disorderly world.⁵¹

The concept of disease also helps us understand differing perspectives on patterns of behaviour (table 2), such as homosexuality. The American Psychiatric Association labeled homosexuality as a disease until 1973, when it was removed from its diagnostic and statistical manual of mental disorders (DSM). However, it remained in the ICD until 1992.⁵³ Western societies increasingly view homosexuality as a legitimate lifestyle choice; less than 5% of doctors and nurses and less than 10% of laypeople and MPs in our survey considered homosexuality a disease. Our respondents likewise did not consider trans-sexualism a disease, contrary to the current ICD-10 classification.²⁵ As with addiction, there is another non-disease perspective on sexual orientation: that homosexuality represents a moral failing. Historically, Western societies have deemed homosexual acts criminal behaviour. In many countries in the world, this continues to be the case.

CONCLUSIONS

In summary, the substantial disagreement we found in classifying of states of being as diseases, and the parallel disagreement regarding the legitimacy of public funding for those that warrant treatment provide insight into the attitudes underlying a number of current high profile social debates. The finding suggests that a shared understanding of the biological and social determinants of health conditions and human behaviours could be very useful in helping to facilitate the resolution of these debates.

Author affiliations

¹Department of Clinical Epidemiology & Biostatistics, McMaster University, Hamilton, Ontario, Canada

²Department of Urology, Helsinki University Central Hospital and University of Helsinki, Helsinki, Finland

³Insurance Division, State Treasury, Helsinki, Finland

⁴Department of Medicine, McMaster University, Hamilton, Ontario, Canada

⁵Department of Orthopaedics and Traumatology, Helsinki University Central Hospital and University of Helsinki, Helsinki, Finland

Acknowledgements The authors thank Diane Heels-Ansdell and Brittany B Dennis for constructive comments on data analysis and interpretation of results, Virginia Mattila for linguistic expertise and language revisions, and Paula Hakkarainen, Kustaa Käki, Sanna Käki, Aura Matikainen, Kristiina Mellais, Dr Anita Pienimäki and Markku Viitamäki for support with data acquisition and/or constructive comments on study design and concept.

Contributors KAOT, JSL and TLNJ conceptualised the study. KAOT and TLNJ obtained funding. KAOT collected the data. KAOT and GHG developed the analysis plan with JSL, SE and TLNJ. KAOT analysed the data. All authors

contributed to the interpretation of the results. KAOT and GHG led the writing of the manuscript; all authors contributed. All authors had full access to all the data and take responsibility for the integrity and the accuracy of the data. All authors have approved the final version of the manuscript. KAOT is the guarantor.

Funding This study was supported by the Competitive Research Funding of the Pirkanmaa Hospital District (Tampere, Finland) grant numbers 9L033 and 9K043. The work of KAOT was supported by the Finnish Cultural Foundation and the Finnish Medical Foundation. The work of SE was supported by a Canadian Institutes of Health Research (CIHR) Doctoral Award. The funding sources had no role in design and conduct of the study; collection, management, analysis and interpretation of the data; and preparation, review, or approval of the manuscript. The authors' work was independent of the funders.

Competing interests JSL is a chief medical officer at the Insurance Division of the State Treasury (Helsinki, Finland), which is a government agency that handles statutory employment pension, accident and indemnity insurances and insurance-related employer services of government agencies.

Ethical approval In accordance with the Finnish regulations on questionnaire surveys, the ethics committee of the Pirkanmaa Hospital District in Finland granted exemption from ethical review (R11110).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Extra data can be accessed via the Dryad data repository at <http://datadryad.org/> with the doi:10.5061/dryad.94qh5.

REFERENCES

- Eisenberg L. Disease and illness. Distinctions between professional and popular ideas of sickness. *Cult Med Psychiatry* 1977;1:9–23.
- Campbell EJ, Scadding JG, Roberts RS. The concept of disease. *Br Med J* 1979;2:757–62.
- Wulff HR. The concept of disease: from Newton back to Aristotle. *Lancet* 1999;354(Suppl):SIV50.
- Temple LK, McLeod RS, Gallinger S, *et al.* Essays on science and society. Defining disease in the genomics era. *Science* 2001;293:807–8.
- Pearce JM. Disease, diagnosis or syndrome? *Pract Neurol* 2011;11:91–7.
- McWhinney IR. Health and disease: problems of definition. *CMAJ* 1987;136:815.
- Seguin CA. The concept of disease. *Psychosom Med* 1946;8:252–7.
- Conrad P, Schneider JW. *Deviance and medicalization: from badness to sickness*. Philadelphia: Temple University Press, 1992.
- Hinshaw SP, Cicchetti D. Stigma and mental disorder: conceptions of illness, public attitudes, personal disclosure, and social policy. *Dev Psychopathol* 2000;12:555–98.
- Perry BL. The labeling paradox: stigma, the sick role, and social networks in mental illness. *J Health Soc Behav* 2011;52:460–77.
- Smith R. In search of 'non-disease'. *BMJ* 2002;324:883–5.
- Heath I. Who needs health care—the well or the sick? *BMJ* 2005;330:954–6.
- Moynihan R. Medicalization. A new deal on disease definition. *BMJ* 2011;342:d2548.
- Scully JL. What is a disease? *EMBO Rep* 2004;5:650–3.
- Moynihan R, Heath I, Henry D. Selling sickness: the pharmaceutical industry and disease mongering. *BMJ* 2002;324:886–91.
- Metzl JM, Herzog RM. Medicalisation in the 21st century: introduction. *Lancet* 2007;369:697–8.
- Kleinman A. Culture, bereavement, and psychiatry. *Lancet* 2012;379:608–9.
- Foucault M. *The birth of the clinic: an archaeology of medical perception*. New York: Pantheon Books, 1973.
- Conrad P. Medicalization and social control. *Annu Rev Sociol* 1992;18:209–32.
- Padamsee TJ. The pharmaceutical corporation and the 'good work' of managing women's bodies. *Soc Sci Med* 2011;72:1342–50.
- Parsons T. *The Social System*. New York: The Free Press, 1951.
- Shilling C. Culture, the 'sick role' and the consumption of health. *Br J Sociol* 2002;53:621–38.
- Parsons T. The sick role and the role of the physician reconsidered. *Milbank Mem Fund Q Health Soc* 1975;53:257–78.
- Meador CK. The art and science of nondisease. *N Engl J Med* 1965;272:92–5.

25. International Statistical Classification of Diseases and Health Related Problems, version 10 (ICD-10). Geneva: World Health Organization. <http://apps.who.int/classifications/icd10/browse/2010/en> (accessed 1 Feb 2012).
26. von Elm E, Altman DG, Egger M, *et al*. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007;335:806–8.
27. Tversky A, Kahneman D. The framing of decisions and the psychology of choice. *Science* 1981;211:453–8.
28. Gallagher KM, Updegraff JA. Health message framing effects on attitudes, intentions, and behavior: a meta-analytic review. *Ann Behav Med* 2012;43:101–16.
29. Mishra S, Gregson M, Lalumière ML. Framing effects and risk-sensitive decision making. *Br J Psychol* 2012;103:83–97.
30. Akl EA, Oxman AD, Herrin J, *et al*. Framing of health information messages. *Cochrane Database Syst Rev* 2011;CD006777.
31. Jones MP. What's a disease? *Am J Gastroenterol* 2003;98:2813–14.
32. Tang CH, Liu JT, Chang CW, *et al*. Willingness to pay for drug abuse treatment: results from a contingent valuation study in Taiwan. *Health Policy* 2007;82:251–62.
33. Perry BL, Pescosolido BA, Martin JK, *et al*. Comparison of public attributions, attitudes, and stigma in regard to depression among children and adults. *Psychiatr Serv* 2007;58:632–5.
34. Lund TB, Sandoe P, Lassen J. Attitudes to publicly funded obesity treatment and prevention. *Obesity (Silver Spring)* 2011;19:1580–5.
35. Stronks K, Strijbis AM, Wendte JF, *et al*. Who should decide? Qualitative analysis of panel data from public, patients, healthcare professionals, and insurers on priorities in health care. *BMJ* 1997;315:92–6.
36. Gross CP, Anderson GF, Powe NR. The relation between funding by the National Institutes of Health and the burden of disease. *N Engl J Med* 1999;340:1881–7.
37. Gillum LA, Gouveia C, Dorsey ER, *et al*. NIH disease funding levels and burden of disease. *PLoS One* 2011;6:e16837.
38. Hawkes N. NHS will soon have to specify what care is and what isn't freely available, GPs say. *BMJ* 2012;344:e1493.
39. Basson R, Berman J, Burnett A, *et al*. Report of the international consensus development conference on female sexual dysfunction: definitions and classifications. *J Urol* 2000;163:888–93.
40. Ganley A. US slams Canada over Vancouver's new drug injection site. *CMAJ* 2003;169:1063.
41. Moynihan R. The making of a disease: female sexual dysfunction. *BMJ* 2003;326:45–7.
42. Hyman SE. The neurobiology of addiction: implications for voluntary control of behavior. *Am J Bioeth* 2007;7:8–11.
43. Madueme H. Addiction as an amoral condition? The case remains unproven. *Am J Bioeth* 2007;7:25–7.
44. Levy N. The social: a missing term in the debate over addiction and voluntary control. *Am J Bioeth* 2007;7:35–6.
45. Henningsen P, Zipfel S, Herzog W. Management of functional somatic syndromes. *Lancet* 2007;369:946–55.
46. *Lancet* editorial board. HIV and injecting drug use: a global call for action. *Lancet* 2011;377:1212.
47. Moscrop A. Medicalisation, morality, and addiction: why we should be wary of problem gamblers in primary care. *Br J Gen Pract* 2011;61:e836–8.
48. McNeil DG Jr. An H.I.V. strategy invites addicts in. *The New York Times*, 16 Sept 2011. <http://www.nytimes.com/2011/02/08/health/08vancouver.html?pagewanted=all> (accessed 1 Feb 2012).
49. Broom DH, Woodward RV. Medicalisation reconsidered: a collaborative approach to care. *Sociol Health Illn* 1996;18:357–78.
50. Wharton S, VanderLelie S, Sharma AM, *et al*. Feasibility of an interdisciplinary program for obesity management in Canada. *Can Fam Physician* 2012;58:e32–8.
51. Scadding JG. Essentialism and nominalism in medicine: logic of diagnosis in disease terminology. *Lancet* 1996;348:594–6.
52. Tikkinen KA, Auvinen A. Does the imprecise definition of overactive bladder serve commercial rather than patient interests? *Eur Urol* 2012;61:746–8.
53. King M, Bartlett A. British psychiatry and homosexuality. *Br J Psychiatry* 1999;175:106–13.