

Support and assessment for fall emergency referrals (SAFER 2) research protocol: cluster randomised trial of the clinical and cost effectiveness of new protocols for emergency ambulance paramedics to assess and refer to appropriate community-based care

Helen Snooks,¹ Rebecca Anthony,¹ Robin Chatters,² Wai-Yee Cheung,¹ Jeremy Dale,³ Rachael Donohoe,⁴ Sarah Gaze,¹ Mary Halter,⁵ Marina Koniotou,¹ Philippa Logan,⁶ Ronan Lyons,¹ Suzanne Mason,² Jon Nicholl,² Ceri Phillips,⁷ Judith Phillips,⁸ Ian Russell,⁹ A Niroshan Siriwardena,¹⁰ Mushtaq Wani,¹¹ Alan Watkins,¹² Richard Whitfield,¹³ Lynsey Wilson¹

To cite: Snooks H, Anthony R, Chatters R, *et al*. Support and assessment for fall emergency referrals (SAFER 2) research protocol: cluster randomised trial of the clinical and cost effectiveness of new protocols for emergency ambulance paramedics to assess and refer to appropriate community-based care. *BMJ Open* 2012;**2**:e002169. doi:10.1136/bmjopen-2012-002169

► Prepublication history for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2012-002169>).

Received 28 September 2012
Accepted 3 October 2012

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

For numbered affiliations see end of article

Correspondence to

Professor Helen Snooks;
h.a.snooks@swansea.ac.uk

ABSTRACT

Introduction: Emergency calls to ambulance services are frequent for older people who have fallen, but ambulance crews often leave patients at the scene without ongoing care. Evidence shows that when left at home with no further support older people often experience subsequent falls which result in injury and emergency-department attendances. SAFER 2 is an evaluation of a new clinical protocol which allows paramedics to assess and refer older people who have fallen, and do not need hospital care, to community-based falls services. In this protocol paper, we report methods and progress during trial implementation. SAFER 2 is recruiting patients through three ambulance services. A successful trial will provide robust evidence about the value of this new model of care, and enable ambulance services to use resources efficiently.

Design: Pragmatic cluster randomised trial.

Methods and analysis: We randomly allocated 25 participating ambulance stations (clusters) in three services to intervention or control group. Intervention paramedics received training and clinical protocols for assessing and referring older people who have fallen to community-based falls services when appropriate, while control paramedics deliver care as usual. Patients are eligible for the trial if they are aged 65 or over; resident in a participating falls service catchment area; and attended by a trial paramedic following an emergency call coded as a fall without priority symptoms. The principal outcome is the rate of further emergency contacts (or death), for any cause and for falls. Secondary outcomes include further falls, health-related quality of life, 'fear of falling', patient satisfaction reported by participants through postal questionnaires at 1 and 6 months, and quality and

ARTICLE SUMMARY

Article focus

- Despite recent focus on the benefits of introducing new falls pathways for older adults within the NHS, no studies have evaluated their effectiveness within a pre-hospital setting.
- This article is a protocol of SAFER 2, a cluster randomised trial evaluating the safety, clinical and cost effectiveness of a protocol for paramedics, enabling them to assess and refer older adults to a community based falls service.
- The focus of this article is to detail how the SAFER 2 intervention will be evaluated: what design and methods are being used and what outcomes are included.

Key messages

- The key aim of this article is to detail the design and methods of the SAFER 2 trial, which measures the safety, clinical and cost effectiveness of a new protocol for paramedics enabling them to assess and refer older adults to a community based falls service.
- This article provides an understanding of the groundwork required to set up and implement a trial of a complex intervention within the pre-hospital setting.

Strengths and limitations of this study

- SAFER 2 followed the MRC Framework for developing and evaluating complex interventions.
- Using a cluster randomised trial design will provide robust evidence about the effectiveness of this new model of care, and enable ambulance services to use resources efficiently.

pathways of care at the index incident. We shall compare National Health Service (NHS) and patient/carer costs between intervention and control groups and estimate quality-adjusted life years (QALYs) gained from the intervention and thus incremental cost per QALY. We shall estimate wider system effects on key-performance indicators. We shall interview 60 intervention patients, and conduct focus groups with contributing NHS staff to explore their experiences of the assessment and referral service. We shall analyse quantitative trial data by 'treatment allocated'; and qualitative data using content analysis.

Ethics and dissemination: The Research Ethics Committee for Wales gave ethical approval and each participating centre gave NHS Research and Development approval. We shall disseminate study findings through peer-reviewed publications and conference presentations.

Trial Registration: ISRCTN 60481756

INTRODUCTION

We have written this protocol paper during trial implementation and include information about study progress. We have carried out several minor amendments to the protocol since the original version. We highlight key differences between the original and current protocol, including sample-size calculations and consent processes.

Background

Falls in older people are recognised as an important issue internationally,^{1 2} with high human and organisational costs. Reduction in quality of life and physical activity lead to social isolation and functional deterioration with a high risk of resultant dependency and institutionalisation.³⁻⁵ In the UK, falls account for 3% (about £980 million) of total National Health Service (NHS) expenditure,⁶ and the prevention of falls in older people has been highlighted as a priority.^{7 8}

Although prevention appears effective,⁸ reducing falls and associated morbidity depends on early identification of people at high risk and delivery of interventions across traditional service boundaries⁹, priorities now reflected in national and international guidelines.¹⁰⁻¹² A recent systematic review and meta-analysis found limited evidence of benefit from multifactorial risk assessment and targeted intervention for falls in primary, community or emergency care. However, none of these trials reported quantitative outcome data on health-related quality of life (HRQoL), and although six had been undertaken in emergency care, none were conducted in prehospital care.

Older people taken to emergency departments following a fall are highly likely to fall again in the following year, with a 30% chance of sustaining fracture or dislocation.¹³ Multidisciplinary interventions have increased uptake of preventative advice,^{14 15} and reduced subsequent falls, length of hospital stay and disability.¹³ Older people commonly call an emergency ambulance (999) following a fall.

In London (UK), this group accounts for about 60 000 attendances (8% of emergency ambulance responses) each year. This is very similar to the proportion reported in an urban Emergency Medical Service system in the USA.¹⁶ Non-conveyance to emergency departments (EDs) is high in this group—close to 40% in London,¹⁷ elsewhere in the UK,^{18 19} and the USA.¹⁶ Most (90%) falls not conveyed to ED occur in the home.²⁰ Non-conveyance of patients is recognised internationally as a safety and litigation risk.²¹

More widely, in most UK ambulance services, treatment protocols advise conveying patients to ED unless they refuse to travel to hospital. In practice, however, ambulance services allow their staff to decide who can be safely left at home. Little is known about how, in the absence of specific protocols or training to leave older people who fall at home, paramedics make these decisions. A US study acknowledged the pragmatic nature of negotiation with patients whether to go to hospital or not.²² A UK study identified factors affecting these decisions including: experience and confidence of ambulance staff, time into the shift, presence of carers, quality of the accommodation, waiting times at the local ED and prior knowledge of the patient.²³ There have been few established referral pathways, or even encouragement to inform patients' general practitioners (GPs), or other services of the emergency call. However, recent policy changes in the UK have encouraged the development and implementation of alternative models of care for delivery by the ambulance service, including enhanced training for paramedics, and community-based referral pathways for patients who do not need the ED.^{24 25}

The National Service Framework for Older People⁷ advocates that ambulance crews refer older people who have fallen to community-based care, although this reflects consensus rather than research evidence. A recent study found that referring elderly patients, who had fallen and been left at home by their attending ambulance crew, to a community-based falls-prevention service reduced further falls and improved clinical outcomes.²⁶ Previous studies in this setting have found that change in practice is difficult to achieve and new pathways of care are difficult to exploit.²⁷ Furthermore, there is little evidence about the safety of non-conveyance decisions by paramedics.²⁸

The SAFER 2 trial has followed the MRC framework for developing and evaluating complex interventions.²⁹ Logan²⁶ has since reinforced the case for a multicentre trial of an intervention in which attending ambulance crews assess patients who have fallen and refer them to community-based falls services from the scene.³⁰⁻³⁴ We hypothesise that the intervention works by improving the decision-making of paramedics to use falls services to best effect. If so, we expect better outcomes and reduced costs, both for patients now referred to falls services and for those not now taken to ED unnecessarily. Achieving these improved outcomes for patients requires participating paramedics to change their practice in

relation to assessment, conveyance and referral of patients. Hence, we have designed SAFER 2 to gather data about each of the elements of the pathway and to assess both processes and outcomes.

AIM AND OBJECTIVES

Aim

To assess the benefits and costs to patients and the NHS of a complex intervention comprising education, clinical protocols and pathways enabling paramedics to assess older people who have fallen and refer them to community-based falls services when appropriate.

Objectives

1. To compare outcomes, processes and costs of care between intervention and control groups.
 - ▶ Patient outcomes: rate and pattern of subsequent emergency healthcare contacts or deaths, for any reason and for falls; HRQoL; psychological status, especially fear of falling; and change in place of residence.
 - ▶ Processes of care: pathway of care at index fall; subsequent healthcare contacts; ambulance service operational indicators and protocol compliance including clinical documentation.
 - ▶ Costs of care: provided by NHS and personal social services; incurred by patients or carers in seeking care.
2. To estimate wider system effects of the introduction of the intervention on ambulance service performance and costs.
3. To understand how patients experience the new health technology.
4. To identify factors which facilitate or hinder the use of the intervention.
5. To inform the development of methods for falls research, especially outcome measures recommended for trials of interventions for older people who fall.²

METHODS AND ANALYSIS

Trial design and management

This is a cluster randomised trial (CRT), with economic and qualitative components. We have randomly allocated ambulance stations between trial groups, both to enable us to support change of practice in the intervention group and to minimise contamination between groups in evaluating patient outcomes. The economic component addresses Objectives 1 and 2 by valuing the benefits and costs of the intervention.

The qualitative component addresses Objectives 2 and 3 through two methods: semistructured interviews with participants (or their carers) attended by intervention paramedics; and focus groups with intervention paramedics and NHS service providers.

Following the MRC guidelines for good practice in clinical trials,³⁵ the management structure comprises external Trial Steering Committee (TSC) and Data

Monitoring and Ethics Committee (DMEC); and internal Trial Management Group (TMG), Local Implementation Team (LIT) in each area, and core team. The TSC oversees the trial and provides advice to the Chief Investigator (CI), the Health Technology Assessment (HTA) Programme and the Sponsor on all aspects. The DMEC has access to unblinded comparative data to monitor the data and make recommendations to the TSC whether there are ethical or safety reasons why the trial should not continue. The TMG manages the trial from day to day. The LITs deal with issues emerging at each trial site and provide opportunities to share progress. The core team is smaller, including the CI and research team.

Setting and site selection

We are undertaking the trial in prehospital emergency care, with paramedics delivering the intervention in partnership with community-based falls services. We have selected three ambulance services in England and Wales, covering a mixture of urban and rural areas where a falls service was available, but no process in place for paramedics to make direct referrals from the scene of 999 attendances.

Participants

We invited paramedics based at ambulance stations that normally attend patients within the catchment area of participating falls services, to participate in the trial before allocating those stations randomly between groups.

Patients are eligible for the trial if they: are aged 65 years or over; live in the catchment area of participating falls services; and are attended by a study paramedic following an emergency call to the ambulance service which is coded by a dispatcher as a fall without priority symptoms (Advanced Medical Priority Dispatch System code 17). We exclude patients attended by an Emergency Care Practitioner unless their attendance was at the request of a trial paramedic. We recruit patients to the trial only once, since subsequent falls constitute patient outcomes.

Interventions

The core of the health technology that we are evaluating is a clinical protocol for the care of older people who have fallen, enabling emergency ambulance paramedics to assess and refer them to community-based falls services. Development of the intervention built on previous studies in this field. This complex intervention comprises training, referral pathways to falls services, individual outcome reports to referring paramedics from falls services, and clinical and operational support to change practice including a feedback loop between paramedics and ambulance service managers. Specialist subgroups developed specific components of the intervention for SAFER 2, while modelling and stakeholder feedback facilitated testing of economic viability and expected

affects. Sites agreed common minimum standards for core elements of the intervention at the outset, while permitting local differences in processes like referral and documentation.

In accordance with the National Institute of Clinical Excellence guidelines for the assessment and prevention of falls,¹² the multidisciplinary falls-services team includes nursing, physiotherapy, occupational therapy and rehabilitation provision. They provide risk assessment and treatment including postural stability and balance training, home hazard advice, equipment and adaptations, medical review including osteoporosis risk, advice on fear of falls, social care, benefit advice and referral to other community services.

Control intervention: we have asked paramedics based at control stations to continue their usual practice. Although we know that conveyance rates vary considerably among services, stations and paramedics, we have not sought standardisation of practice as we do not know what is best. Current practice in the control group is therefore care as usual comprising assessment of injury or other conditions requiring immediate care, assistance in moving from where they have fallen, and conveyance to ED unless the patient refuses.

Outcomes

Outcome measures at 1 and 6 months after patients' index calls are consistent with recommendations of Prevention of Falls Network Europe (ProFaNE).³⁶

Principal outcomes

The rate of further contacts with emergency healthcare providers (999 calls, ED attendances, emergency admissions or death)—both for any cause and specifically for falls, as summarised by

- ▶ Proportion of patients who suffer these events
- ▶ Interval to first event
- ▶ Event rate

Secondary outcomes

- ▶ Duration of inpatient episodes
- ▶ Fractures arising from further falls
- ▶ Self-reported further falls
- ▶ Health-related quality of life, as measured by the SF12³⁷
- ▶ 'Fear of falling' as measured by the modified falls efficacy scale³⁸
- ▶ Patient satisfaction as measured by the (quality of care monitor)³⁹
- ▶ Change in place of residence
- ▶ Pathway of care as measured by routine ambulance service data on proportions conveyed to ED, referred to falls service, referred to other providers, or left at scene without further care
- ▶ Durations of: ambulance service job cycle, episode of care, time to falls service response

- ▶ Compliance with guidelines for ambulance service clinical documentation; referrals; and falls services follow-up
- ▶ Costs of care to NHS and personal social services, estimated by routine data from participating services
- ▶ Self-reported costs incurred by patients and carers
- ▶ Views of ambulance service paramedics, managers and partners on implementation of the intervention
- ▶ Experience and satisfaction of patients receiving the intervention

Data collection methods

This CRT does not approach participants at the point of treatment, because they may be in distress and cannot give informed consent. Instead, we seek retrospective consent to follow up through routine medical records and by postal questionnaire. Following experience in several earlier experimental studies, we originally proposed an 'opt-out' procedure, and gained provisional ethical approval from a Multicentre Research Ethics Committee. However, information governance approval for this approach was not forthcoming. We therefore designed an active consent process, in which we contact patients to seek consent to follow up first by post, and then if necessary by telephone or home visit, and gained the necessary approvals. We also include a £5 voucher with each invitation pack to thank participants for their time. We are also undertaking anonymised follow-up through linked records—in Wales, using the Secure Anonymised Information Linkage (SAIL) databank,⁴⁰ and in England, using similar centralised records—again with information governance approval.

Sample size and power

We estimated the sample size for the trial from our principal outcome—the proportion of participants who, within 6 months, die or contact emergency services (999 service or ED). From previous trials of interventions for older people who have sustained a fall and presented for emergency treatment, summarised in a recent systematic review,⁴¹ we make the conservative estimate that trial patients have about 50% chance of making another emergency contact within 6 months. As the intervention appears cheap a priori, we judge that a change of 5% in this proportion may be clinically and economically important. In the absence of clustering, a sample size of 4190 evaluable participants would yield 90% power to detect a change of at least 5% (from 50% to $\leq 45\%$ or $\geq 55\%$) when using two-sided 5% significance level. As participants come from 25 clusters (ambulance stations), we need to adjust this sample size to allow for intracluster correlation (ICC). We estimate this ICC from the findings of the SAFER 1 trial,⁴¹ which evaluated the clinical and cost effectiveness of Computerised Clinical Decision Support software for use by paramedics when attending older adults who had a fall. SAFER 1 estimated the ICC for the same outcome, but over 1 month rather than 6, as zero when clustering

participants by station (as in SAFER 2) but 0.005 when clustering participants by paramedic (as in SAFER 1).⁴² To be conservative, we allow for an ICC of 0.002. Solving a simple algebraic equation then yields: a target of 251.6 evaluable participants per station; a variance inflation factor of $(1+(251.6-1)\times 0.002)$ viz 1.5012; and a total evaluable sample of $4190\times 1.5012=6290$ viz 25×251.6 . This sample will also have more than 90% power to detect a change of 0.18 in the estimated mean of 1.8 emergency contacts per participant over 6 months, given an estimated SD of 1.5. Hence, SAFER 2 can detect a difference of 1 emergency contact in 10 avoided (or induced) by the intervention.

We had originally postulated that patients recruited to the study would have a 40% chance of making an emergency contact within 6 months; and that the ICC could be as high as 0.03. Under those assumptions, our target sample of about 6300 would have yielded 80% power to detect a change of at least 10% (ie, from 40% to $\leq 30\%$ or $\geq 50\%$) when using a two-sided 5% significance level. When SAFER1 showed that the assumed ICC was unduly pessimistic, recruitment was progressing well. Rather than finish the trial early, we decided with the approval of both TSC and DMEC to be less conservative in assuming a worst ICC of 0.02, thus yielding enough power to detect a change of only 5% in the emergency contact rate, still a clinically important difference in the view of our advisers.

Early in SAFER 2, approximately 1 in 10 recruited participants consented to complete questionnaires, confidential but not anonymised. This has now increased to 1 in 4 for the 1-month questionnaire. The trial is on target to achieve a sample size of 6290, and the resulting sample of 800 returned questionnaires at 6 months would yield 90% power of detecting an effect size of 0.25 (equivalent to one-quarter of the population SD) in each of the questionnaire outcomes. Such a low response rate requires that analysis include rigorous non-response analysis to test whether findings extrapolate to the entire population of interest.

Loss to follow-up

We monitor routine outcomes in two forms: anonymised linked data from central NHS databanks for all patients that we can match to NHS administrative records, which needs information governance approvals but no consent; and identifiable data from NHS providers for patients who consent for us to do this. Our experience in the recently completed SAFER 1 trial in a similar population in Wales suggests we can achieve 90% follow-up through anonymised linked data. Although this will reduce statistical power below the 90% postulated in our calculations, it will still exceed the traditional 80%.

Randomisation

An independent statistician randomised the 25 participating ambulance stations between intervention and

control groups after the paramedics had volunteered to participate, thus minimising selection bias; the stratifying variables were the receiving falls service and the number of paramedics participating in each station.

Blinding

Although the trial managers and fieldworkers need to know the allocation of all participating ambulance stations for operational reasons, we keep the trial statistician blind to these allocations.

Statistical analysis

Primary analysis will be by 'treatment allocated'. Analyses will include logistic regression for binary outcomes, cross-tabulations and risk ratios for categorical outcomes, and survival analysis including Cox's proportional hazards models for times to events. We shall use multilevel modelling to estimate (random) station effects and (fixed) group effects and analyse repeated observations as such.

Our principal outcomes comprise a hierarchy, and will undergo analysis incrementally: first deaths; second emergency admissions plus deaths; then ED attendances plus admissions and deaths and finally 999 calls plus attendances, admissions and deaths. Analysis at 1 and 6 months will cover, for all such events, and those coded as a fall: the proportion of patients that call 999, attend ED, get admitted or die; survival analysis of the time to the first subsequent emergency contact; the mean number of further emergency contacts adjusted for time at risk, excluding days in hospital or after death; and recurrent event analysis where feasible. As patients' decision to call 999 after later events could reflect the care they received at the index call, we shall check whether these later calls reflect valid need rather than health-seeking behaviour, by comparing them with self-reported falls and health-related quality of life. We will also examine the effect of the intervention on patient satisfaction, health-related quality of life and costs (as described separately, below).

Potential predictors of triage decisions include the distance between the site of the index event and the ED; patients' age, sex and history of previous falls; type of presentation (eg, out-of-hours); and time since recruitment, as routine data may be less accessible for patients recruited later in the trial. We shall therefore use these as covariates in the analysis. It is possible that patients in the catchment area of one station may receive care for a subsequent event from another station participating in the study but allocated to a different group. Nevertheless, analysis will still be by treatment allocated. Secondary analyses will examine outcomes by treatment received, namely whether participants got referred to falls services.

To identify any wider system effects, we shall compare response times during the trial period across the study catchment area and surrounding areas with pretrial response times and response times elsewhere. We shall also compare the characteristics of those included and

not included through both consented and anonymised routes, to explore whether there are systematic differences between groups and routes that may influence outcomes.

To inform the development of outcome measures for falls research as recommended by ProFaNE, we shall compare SF12 and derived SF6D scores with mFES scores to establish their construct validity. We shall also assess their predictive validity by comparing scores with the number of further events and the time to the first subsequent event.

Economic evaluation

We are undertaking economic evaluation alongside the CRT from the perspective of the UK NHS and personal social services, and patients and their families. Economic analysis will estimate the costs of providing the intervention, the costs to patients and families, and the consequences of the scheme for the NHS and social services in terms of inpatient admissions, ED attendances, GP consultations, out-of-hours GP contacts, NHS direct contacts, and use of social services. We shall collect data on participants' use of health service and social services resources from paramedic records, routine hospital records and patient-completed questionnaires. We shall estimate NHS resource use from routine data including duration of ambulance job cycles and episodes of care, records of resource use and patient records. We shall estimate social services resource costs from discussion with relevant social services departments. We shall calculate the resulting costs using unit costs from published sources. We shall estimate the quality-adjusted life years (QALYs) gained by the intervention from the SF6D. We shall derive an incremental cost-per-QALY and present the resulting cost-effectiveness plane and associated cost-effectiveness acceptability curves. We undertake sensitivity analysis to assess the robustness of the results to changes in the configuration of the scheme and other health-service costs.

We recognise that the follow-up period is not long enough to yield evaluation over the lifetime of participants. We shall therefore develop a decision model to extrapolate costs and effects beyond the data generated by the trial—probably from the hazard rates estimated by the trial. We shall construct alternative scenarios from previous studies and discussions with experts—to judge the extent to which risks will remain constant or change over time, and assess the implications of using a longer time-horizon on the basic analysis of costs and benefits.

Qualitative study

One researcher at each site is undertaking semistructured interviews with older people who have been recruited to the study and focus groups interviews with intervention paramedics and service providers. We are purposively sampling 20 participants in each site attended by intervention paramedics following a fall, to include patients transferred to the ED, patients referred

to the falls service and patients neither transferred or referred. We interview these participants or their carers at a site of their choosing 6–8 weeks after their index fall. We developed the interview schedule to gather information in depth about the experience of patients in the intervention group, and we consulted user groups about the content and acceptability of that schedule. In particular, we are interested in intervention fidelity and the perceptions of those who received the intervention—for instance, whether they feel confident about paramedics' decisions whether they need to go to the ED; and how they feel about the process of referral.

Focus groups are a useful way of understanding organisational change,⁴³ and exploring the success or failure of particular programmes.⁴⁴ At each of the three centres, we are undertaking focus groups with intervention paramedics before and after the trial period. Following the trial period, we shall hold focus groups with a range of ambulance service participants in each centre, including trainers, operational managers, clinical team leaders and dispatch staff; and with participants from other partner services in each centre including falls services, social services and ED. We include six to eight participants in each focus group, to facilitate discussions within manageable groups.⁴⁵ We base the topic guides on previous research in this area and consultation with our LITs. Two researchers lead each focus group, one to facilitate discussion and the other to take notes that link text to speakers and highlight points of consensus or disagreement and issues that draw strong emotional responses such as anger, fear or anxiety.

Systematically comparing and analysing qualitative data in raw form is challenging.⁴⁶ So, we shall record and transcribe all interviews and focus groups with the permission of participants. A protocol will ensure that we use standard format and conventions throughout the transcription process. We shall analyse all these data by using NVivo software to explore commonalities and differences in topics that emerge from the guides. Two researchers will analyse all these data independently and then meet to discuss and agree final coding and interpretation.

Adverse event reporting

SAFER 2 is following the principles of the Standard Operating Procedure for adverse events developed by the West Wales Organisation for Rigorous Trials in Health (WWORTH). As the study population has high mortality and morbidity, we do not routinely record or report adverse events that are neither serious nor adverse reactions (ARs) in the sense of possibly being caused by the new clinical protocol for referring to falls services. The main potential AR is misdiagnosis, which could lead to an inappropriate pathway of care. As misdiagnosis is reliably identifiable only through patient complaints or coroner's inquest, we focus on these, and treat them as serious adverse reactions (SAR). Any patient complaint or coroner's inquest at which the ambulance service is asked to supply information related

to non-conveyance of a trial participant from the index incident will trigger investigation by the local principal investigator and, chief investigator. We also investigate suspected ARs brought to our attention in any other way.

Death or emergency hospital admission is a serious adverse event (SAE). As these form the primary outcome of this trial, and are not unusual or unexpected in the study population, we shall report them at the end of the trial. In particular imbalance between intervention and control groups in the occurrence of SAEs or SARs, will be the subject of statistical analysis at the end of the trial.

ETHICS AND DISSEMINATION

Ethics and R&D governance

Current practice has been shown to carry risks for patients, as many (up to half) are left at scene without further care, and many of these (about half) make further emergency healthcare contacts within 2 weeks.¹⁹ Following a recommendation in the National Service Framework for Older People,⁹ ambulance services around the UK have begun to implement alternative pathways of care for older people who have fallen, either through emergency care practitioner schemes or direct referral from paramedics. Research is urgently needed to understand the safety, costs and clinical effectiveness of this new model of care. Ethical and consent issues in CRTs are acknowledged to present their own unique challenges.⁴⁷ Against this background, we have obtained ethical approval from the Research Ethics Committee for Wales, information governance approval from the National Information Governance Board, and NHS R&D approval from each participating Health Board, NHS Trust and Primary Care Trust.

Service users

We include patients and carers as active participants in the research at all stages. As the relevant service users are often frail, we are using innovative methods to facilitate their contributions. They attend TSC, DMEC, TMG and LIT meetings and additional service user groups, where they contribute to the research process and discuss issues affecting older people with a history of falls. We do not expect them to attend full research team meetings, although they may bring their views to the team meetings, following meetings with service users in other forums. Including service users in emergency care research is a particular challenge,⁴⁸ but is achievable and brings rewards to the trial and the team.

Dissemination

We shall comply with the CONSORT guidelines.⁴⁹ We will present the study results at national and international conferences and publish them in peer-reviewed journals. In accordance with recommendations, we have registered SAFER 2 in a public registry (<http://www.controlled-trials.com/isrctn/>, Identifier: ISRCTN 60481756).

DISCUSSION

Progress so far (September 2012)

The SAFER 2 trial is underway in three ambulance services, in collaboration with eight participating falls services and 12 hospitals with EDs. We have recruited 220 paramedics from 25 ambulance stations (clusters) to the trial. In the first year of the trial, we recruited over 4000 patients. Hence, we are on target to detect clinically important differences in outcomes at 6 months, while monitoring the safety of the intervention at 1 month.

Author affiliations

¹Centre for Health Information Research and Evaluation, Swansea University, Swansea, UK

²School of Health and Related Research (SchARR), University of Sheffield, Sheffield, UK

³Warwick Medical School, University of Warwick, Coventry, UK

⁴Clinical Audit and Research Unit, London Ambulance Service NHS Trust, London, UK

⁵Faculty of Health and Social Services, St Georges University Hospital, London, UK

⁶Community Health Sciences, The University of Nottingham, Nottingham, UK

⁷Swansea Centre for Health Economics, College of Human and Health Sciences, Swansea University, Swansea, UK

⁸Centre for Innovative Ageing, Swansea University, Swansea, UK

⁹West Wales Organisation for Rigorous Trials in Health, College of Medicine, Swansea, UK

¹⁰School of Health and Social Care, University of Lincoln, Lincoln, UK

¹¹Department of Geriatric and Stroke Medicine, Morriston Hospital, Swansea, UK

¹²School of Business and Economics, Swansea University, Swansea, UK

¹³Pre-hospital Emergency Research Unit (PERU), Welsh Ambulance Services NHS Trust, Cardiff, UK

Acknowledgements We thank our collaborators in the community falls services and ambulance services for making this trial a success to date. We are grateful to the service users who also made valuable contributions.

Contributors All authors made a significant contribution to the conception and design of the study protocol. The protocol was written by H Snooks, S Gaze, J Phillips, C Phillips, A Watkins, I Russell and I Cheung and was critically reviewed by J Nicholl, S Mason, R Whitfield, M Wani, N Siriwardena, R Lyons, P Logan, M Halter, R Donohoe, R Anthony, M Konioutou, L Wilson, R Chatters and J Dale. All authors gave approval for the publication.

Funding The National Institute for Health Research (NIHR) Health Technology Assessment (HTA) Programme funds this trial as project 07/01/21; it monitors progress and quality through biannual reports. To inform the British NHS, NIHR will peer-review the full report and publish it in *Health Technology Assessment*. The NIHR also encourages grant-holders to seek external peer-reviewed publication. As sponsor Swansea University monitors legal and financial probity; it delegates monitoring of progress and quality to the WWORTH, the local Registered Clinical Trials Unit.

Competing interests None.

Ethics approval The Research Ethics Committee for Wales gave ethical approval and each participating centre gave NHS Research and Development (R&D) approval.

Provenance and peer review Not commissioned; internally peer-reviewed.

REFERENCES

- Bergeron E, Clement J, Lavoie , *et al*. A simple fall in the elderly: not so simple. *J Trauma Inj Infect Crit Care* 2006;60:268–73.
- Sikron F, Giveon A, Aharonson-Daniel L, *et al*. My home is my castle! Or is it? Hospitalizations following home injury in Israel, 1997–2001. *Isr Med Assoc J* 2004;6:332–5.

3. Keene GS, Parker MJ, Pryor GA. Mortality and morbidity after hip fractures. *Br Med J* 1993;307:1248–50.
4. Kellogg International Work Group: The prevention of falls in later life. KIWG on the prevention of falls by the elderly. *Dan Med Bull* 1987;34:1–24.
5. Tinetti ME. Factors associated with serious injury during falls by ambulatory nursing home residents. *J Am Geriatr* 1987;35:644–8.
6. Scuffham P, Chaplin S, Legood R. Incidence and costs of unintentional falls in older people in the United Kingdom. *J Epidemiol Community Health* 2003;57:740–4.
7. Department of Health: National Service Framework for Older People. 2001. http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/documents/digitalasset/dh_4071283.pdf (accessed Mar 2011).
8. Gillespie LD, Gillespie WJ, Robertson MC, *et al.* *Interventions for preventing falls in elderly people (Cochrane Review)*. The Cochrane Library. Chichester: John Wiley and Sons Ltd, 2003.
9. Close JCT. Interdisciplinary practice in the prevention of falls—a review of working models of care. *Age Ageing* 2001;30:8–12.
10. Aoyagi K, Ross PD, David JW, *et al.* Falls among community-dwelling elderly in Japan. *J Bone Miner Res* 1998;13:1468–74.
11. American Geriatrics Society BGSAAoOS. Guidelines for the prevention of falls in older persons. *J Am Geriatr Soc* 2001;49:664–72.
12. Clinical practice guideline for the assessment and prevention of falls in older people. Guidelines commissioned by the National Institute of Clinical Excellence (NICE, 2004).
13. Close J, Ellis M, Hooper R, *et al.* Prevention of falls in the elderly trial (PROFET): a randomised controlled trial. *Lancet* 1999;353:93–7.
14. Davison J, Bond J, Dawson P, *et al.* Patients with recurrent falls attending accident & emergency benefit from multifactorial intervention—a randomised controlled trial. *Age Ageing* 2005;34:162–8.
15. Whitehead C, Wundke R, Crotty M, *et al.* Evidence-based clinical practice in falls prevention: RCT of a falls prevention service. *Aust Health Rev* 2003;26:88–97.
16. Weiss SJ, Chong R, Ong M, *et al.* Emergency medical services screening of elderly falls in the home. *Prehosp Emerg Care* 2003;7:79–84.
17. Snooks H, Halter M, Close J, *et al.* Emergency care of older people who fall: a missed opportunity. *Qual Saf Healthcare* 2006;15:390–2.
18. Snooks H, Porter A, Gaze S, *et al.* *Non-conveyance of 999 patients and related clinical documentation: Policy, practice and views. Report to the Welsh Ambulance Services NHS Trust*, 2004.
19. Marks PJ, Daniel TD, Afolabi O, *et al.* Emergency (999) calls to the ambulance service that do not result in the patient being transported to hospital: an epidemiological study. *Emerg Med J* 2002;19:449–52.
20. Close JC, Halter M, Elrick A, *et al.* Falls in the older population: a pilot study to assess those attended by London ambulance service but not taken to ED. *Age Ageing* 2002;31:488–9.
21. Goldberg RJ, Zautcke JL, Koenigsberg MD, *et al.* A review of prehospital care litigation in a larger metropolitan EMS system. *Ann Emerg Med* 1990;19:557–61.
22. Cone DC, Kim DT, Davidson SJ. Patient-initiated refusals of prehospital care: ambulance call report documentation, patient outcome and on-line medical command. *Prehosp Disaster Med* 1995;10:22–8.
23. Snooks H, Kearsley N, Dale J, *et al.* Gaps between policy, protocols and practice: a qualitative study of emergency ambulance crews' views and practice concerning the care of patients with non-urgent needs. *Qual Saf Health Care* 2005;14:251–7.
24. Department of Health. *Taking healthcare to the patient. Transforming NHS ambulance services*. Department of Health, London, 2005.
25. Cooper S, Barrett B, Black S, *et al.* The emerging role of the emergency care practitioner. *Emerg Med J* 2004;21:614–18.
26. Logan PA, Coupland CAC, Gladman JRF, *et al.* Community falls prevention for people who call an emergency ambulance after a fall: randomised controlled trial. *BMJ* 2010;340:c2102.
27. Snooks H, Close JCT, Gaze S, *et al.* *Older fallers attended by the ambulance service and left at home. Report to WORD*. February 2003.
28. Snooks H, Kearsley N, Dale J, *et al.* Towards primary care for non-serious callers to the emergency ambulance service: results of a controlled study of 'Treat and Refer' protocols for ambulance crews. *Qual Saf Health Care* 2004;13:435–43.
29. Craig P, Dieppe P, Macintyre S, *et al.* Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008;337:a1655.
30. Halter M, Close JCT, Snooks H, *et al.* Fit to be Left: can ambulance staff use an assessment tool to decide if an older person who has fallen can be safely left at home. *DH Research Programme to Support Implementation of the NSF for Older People*.
31. Mason S, Knowles E, Colwell B, *et al.* Effectiveness of paramedic practitioners in attending 999 calls from elderly people in the community: cluster randomised controlled trial. *BMJ* 2007;335:919.
32. Turner J, Nicholl J, Webber L, *et al.* A randomised controlled trial of prehospital intravenous therapy in serious trauma. *Health Technol Assess* 2000;4:31.
33. Turner J, Snooks H, Youren A, *et al.* *The costs and benefits of managing some low-priority 999 calls by NHS Direct nurse advisers*. Final report to the National Co-ordinating Centre for NHS Service Delivery and Organisation R&D. April 2006.
34. Snooks H, Foster T, Nicholl J. Results of an evaluation of the effectiveness of triage and direct transportation to Minor Injuries Units by ambulance crews. *EMJ* 2004;21:105–11.
35. Medical Research Council. *MRC guidelines for good clinical practice in clinical trials*. Medical Research Council 1998; 20 Park Crescent, London.
36. Lamb SE, Jorstad-Stein EC, Hauer K, *et al.* Development of a common outcome data set for fall injury prevention trials: the Prevention of Falls Network Europe consensus. *J Am Geriatr Soc* 2005;53:1618–22.
37. Ware JE, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey (SF12): construction of scales and preliminary tests of reliability and validity. *Med Care* 1996;32:220–33. <http://www.sf-36.org/tools/sf12.shtml> (accessed March 2011-February 2013)
38. Hill KD, Schwarz JA, Kalogeropoulos AJ, *et al.* Fear of falling revisited. *Arch Phys Med Rehab* 1996;77:1025–9.
39. Carey RC, Seibert JH. A patient survey system to measure quality improvement: questionnaire reliability and validity. *Med Care* 1993;31:834–45.
40. Lyons R, Jones K, John G, *et al.* The SAILdatabank: linking multiple health and social care datasets. *BMC Med Inform Decis Mak* 2009;9:3.
41. Aminzadeh F, Dalziel WB. Older adults in the emergency department: a systematic review of patterns of use, adverse outcomes and effectiveness of interventions. *Ann Emerg Med* 2002;39:238–47.
42. Snooks H, Cheung W, Close J, *et al.* Support and Assessment for Fall Emergency Referrals (SAFER 1) trial protocol. Computerised on-scene decision support for emergency ambulance staff to assess and plan care for older people who have fallen: evaluation of costs and benefits using a pragmatic cluster randomised trial. *BMC Emerg Med* 2010;10:2.
43. Barbour RS. Are focus groups an appropriate tool for studying organizational change? In: Barbour RS, Kitzinger J, eds. *Developing focus group research*. London: Sage, 1999:113–26.
44. Morgan DL, Krueger RA. When to use focus groups and why. In: Morgan DL, ed. *Successful focus groups: advancing the state of the art*. London: Sage, 1993:1–19.
45. Kitzinger J, Barbour RS. Introduction: the challenge and promise of focus groups. In: Barbour RS, Kitzinger J, eds. *Developing focus group research*. London: Sage, 1999:1–20.
46. Miles MB, Huberman MA. *Qualitative data analysis: an expanded source book*. London: Sage, 1994.
47. Hutton JL. Are distinctive ethical principles required for cluster randomised trials? *Statist Med* 2001;20:473–88.
48. Halter M, Jones K, Snooks H, *et al.* Involving consumers in prehospital care research and audit (poster). *Pre-hosp Immediate Care* 1999;3:177.
49. CONSORT. Explanation and Elaboration: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;340:bmj.c869.