



Heat related respiratory hospital admissions in Europe in a changing climate: A health impact assessment

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2012-001842
Article Type:	Research
Date Submitted by the Author:	24-Aug-2012
Complete List of Authors:	Åström, Christofer; Umeå University, Department of Public Health and Clinical Medicine Orru, Hans Rocklöv, Joacim; Umeå University, Department of Public Health and Clinical Medicine Strandberg, Gustav; Swedish Meteorological and Hydrological Institute, Ebi, Kristie; Stanford University, ClimAdapt Forsberg, Bertil; Umeå University, Department of Public Health and Clinical Medicine
Primary Subject Heading:	Epidemiology
Secondary Subject Heading:	Epidemiology, Occupational and environmental medicine
Keywords:	EPIDEMIOLOGY, PUBLIC HEALTH, Epidemiology < THORACIC MEDICINE

SCHOLARONE™
Manuscripts

Heat related respiratory hospital admissions in Europe in a changing climate

Authors:

Christofer Åström^{1*}, Hans Orru^{1,2}, Joacim Rocklöv^{1,3}, Gustav Strandberg⁴, Kristie L. Ebi⁵, Bertil Forsberg¹

Affiliations:

¹ Department of Public Health and Clinical Medicine, Occupational and Environmental Medicine, Umeå University, Umeå, Sweden; ² Department of Public Health, University of Tartu, Tartu, Estonia; ³ Department of Public Health and Clinical Medicine, Epidemiology and Global Health, Umeå University, Umeå, Sweden; ⁴ Swedish Meteorological and Hydrological Institute, Norrköping, Sweden; ⁵ ClimAdapt, Stanford University, Los Altos, California, USA

Corresponding author:

* Author to whom correspondence should be addressed:

Christofer Åström,

Occupational and Environmental Medicine, Department of Public Health and Clinical Medicine, Umeå University, SE 901 87, Sweden

+4690 785 0874

christofer.astrom@envmed.umu.se

Keywords: Climate change, respiratory health, public health, heat

Word count: 3045

ABSTRACT

Objectives: Respiratory diseases are ranked second in Europe in terms of mortality, prevalence and costs. Studies have shown that extreme heat has a large impact on mortality and morbidity, with a large relative increase for respiratory diseases. Expected increases in mean temperature and the number of extreme heat events over the coming decades due to climate change raise questions about the possible health impacts. Here we aim to assess the number of heat related respiratory hospital admissions in a future with a different climate.

Design: A European wide health impact assessment.

Setting: An assessment for each of the EU27 countries.

Methods: Heat-related hospital admissions under a changing climate are projected using multi-city epidemiologic exposure-response relationships applied to gridded population data and country-specific baseline respiratory hospital admission rates. Times-series of temperatures are simulated with a regional climate model based on four global climate models, under two greenhouse gas emission scenarios.

Results: Between a reference period (1981–2010) and a future period (2021–2050), the total number of respiratory hospital admissions attributed to heat is projected to be larger in Southern Europe, with three-times more heat attributed respiratory hospital admissions in the future period. The smallest change was estimated in Eastern Europe with about a two-fold increase. For all of Europe, the number of heat-related respiratory hospital admissions is projected to be 26,000 annually in the future period compared with 11,000 in the reference period.

Conclusions: The results suggest that projected effects of climate change on temperature and the number of extreme heat events could substantially influence respiratory morbidity across Europe.

ARTICLE SUMMARY

Article focus

To assess how heat related respiratory hospital admissions in Europe are expected to change with the global warming in the near future.

With a range of climate change projections explore the range of the different impact estimates.

Key messages

The 30 year annual mean increase in respiratory hospital admissions related to heat can be counted in 10's of thousands.

The increase will be different in different parts of Europe with the largest relative increase in Southern Europe.

Strengths and limitations

The study takes the spatial variation in both climate and population density into account in the impact calculations. This makes the estimates valid for countries with large national differences in population densities and climate.

The study results exemplifies some the variation of the impacts depending on different climate change scenarios as well as the variations within one climate scenario depending on the underlying climate model.

The use of real spatial population data has limited the study to explore the change in all age respiratory hospital admissions without demographic trends. The fact that the elderly is a well-known risk group when studying heat related illness combined with the future rise of the proportion of elderly people in the population may suggest that the estimated impacts are rather conservative.

INTRODUCTION

Respiratory diseases are ranked second in Europe in terms of mortality, prevalence and costs.¹ This burden is expected to increase, partly due to a changing climate.² An environmental factor with a large impact on mortality and morbidity in Europe is extreme heat, with large effects on respiratory diseases.³⁻⁵ Physiological effects of exposure to heat can be directly heat related (heat stroke, heat fatigue and dehydration) or can contribute to worsening of respiratory and cardiovascular diseases, electrolyte disorders, and kidney problems.⁴⁻⁷ A review found that heatwaves have a stronger relative impact on mortality than on emergency room visits and hospital admissions, suggesting that many individuals die before they can get to the hospital.⁸ However, several studies confirm that heat also affects health care utilisation.^{3,9} During the heatwave of 2003 in France, the Assistance Public Hôpitaux de Paris recorded 2,400 additional visits to the emergency care units and 1,900 additional hospital admissions.¹⁰ During the heatwave in California in 2006 there were almost 1,200 hospital admissions together with more than 16,000 visits to the emergency departments.⁶

It is possible that hospital admissions could increase in the future due to temperature changes projected with climate change, assuming no additional acclimatisation. Projected changes include an increase in global mean temperature of 1.8°C to 4.0°C by the end of this century, with larger increases over land areas at high latitudes, changing seasonal temperatures, and increases in the frequency, intensity, duration, and spatial extent of heatwaves.¹¹ Recent projections for 2070–2100 suggest that maximum temperatures experienced once every 20 years during the period 1961–1990 could be expected as often as every year in southern Europe and every 3rd-5th year in northern Europe.¹² This means that Europe should expect an increase in both mean temperatures and the number of extreme heat events. Recent heatwaves are consistent with projections. For example, the heatwave that occurred in Europe 2003 could be expected to return every 46,000 years, based on the temperature distribution for the years 1864–2000.¹³ However, such extreme hot conditions may become normal in the end of this century due to anthropogenic climate changes.¹⁴ The uncertainty in

1
2
3 this estimated return time is quite high, with a lower bound of the 90% confidence interval of 9,000
4
5 years. Even so, another heatwave occurred in 2006 with the most anomalous July temperature ever
6
7 measured in Europe.¹⁵ In 2010, Eastern Europe experienced a heatwave with summer temperatures
8
9 higher than the last 140 years, resulting in roughly 55,000 excess deaths in Russia.¹⁶
10

11
12 The possible impacts of these changes in morbidity, particularly respiratory health, are relatively
13
14 unexplored.
15

16 17 18 **Aim**

19
20 The aim of this study is to assess the extent to which climate change could affect heat related
21
22 respiratory hospital admissions (RHAs) in Europe. Using a range of climate projections, we estimate
23
24 the change in hospital admissions between a reference period and a future period.
25
26
27
28

29 **METHODS**

30 31 32 **Climate change and temperature modelling**

33
34 The Rossby Centre regional atmospheric climate model RCA3¹⁷ was developed by SMHI (Swedish
35
36 meteorological and hydrological institute) to dynamically downscale results from the global climate
37
38 models CCSM3¹⁸, ECHAM5¹⁹, HadleyCM3²⁰ and ECHAM4²¹ to a higher resolution over Europe. RCA3 is
39
40 run on a horizontal grid spacing of 0.44° (corresponding to approximately 50 km) and a time step of
41
42 30 minutes. Projections are based on the global greenhouse gas emission scenarios A1B and A2
43
44 (scenarios are described in detail in the SRES²²). Both have been used in climate change health
45
46 impact assessments (HIA).²³⁻²⁶ A1B is a “middle of the road” scenario and A2 is considered a high
47
48 emission scenario, although recent greenhouse gas emissions have been higher. Data from one
49
50 climate model, under one climate change scenario, is referred to as one climate change projection.
51
52

53
54
55 We use aggregated daily projections of maximum temperature and relative humidity data to
56
57 estimate exposure for the periods 1981–2021 (reference period) and 2021–2050 (future period).
58
59
60

Population and morbidity rates

The annual country specific rate of respiratory hospital admissions between 2005 and 2010 were extracted from the WHO's European Health for All Database (<http://data.euro.who.int/hfad>) for the EU27 countries. The mean value over the six years was used as a baseline morbidity rate. Official population data were from the HYDE theme within the Netherlands Environmental Assessment Agency.²⁷ These data are gridded on a 0.0833° resolution (approximately 9,45km) and matched with the climatic data by summing the population within each climatic grid cell.

Exposure-response assumption

The impact calculations were based on the heat morbidity relationship estimated within the European PHEWE project.³ This relationship was based on analyses of 12 European cities over the 1990s, using the daily maximum of apparent temperature (AT), where AT is a combination of the measured temperature and the dew point temperature:

$$AT = -2.653 + 0.944 * T + 0.0153 * (DT)^2$$

where T is the air temperature and DP the dew point temperature, which can be derived from temperature and relative humidity.²⁸ The AT accounts for heat stress related not only to the absolute temperature but also to the saturation of the surrounding air that makes it harder to regulate body temperature by sweating, the most important mechanism to maintain a healthy body temperature.²⁹

³⁰ The PHEWE study used a 0–3 day lag of the maximum AT and concluded that the 90th percentile of this exposure variable for the summer months, April to September, is the appropriate threshold value for the heat morbidity function.³ We calculated the 90th percentile for the summer months for each grid cell from the climate data for the 1990s yielding individual thresholds for each grid cell for each climate projection.

The PHEWE study calculated the relative risk (RR) coefficients for each of the 12 cities and then combined into two meta-coefficients for North-Continental cities 1.012 (95% CI 1.001–1.022) and

Mediterranean cities 1.021 (95% CI 1.006–1.036), associated with a 1-degree increase above the temperature threshold. Meta-coefficients were more suitable as they allowed calculation of the change in RHAs for countries with and without a city in the PHEWE Study. We assigned the Mediterranean coefficient to Portugal and the Mediterranean countries. The rest of Europe was assigned the RR for the North continental cities. Because the population data were not stratified by age, we used the PHEWE study coefficients for all ages.

Projections

The final data used in the impact calculations were based on the climate data grid with a resolution of 0.44° resulting in 8,075 grid cells over Europe. The population data, RR coefficients and baseline morbidity rates were projected for each grid cell. In addition, the countries within EU27 were grouped into four regions, Northern, Western, Eastern and Southern Europe, according to the United Nations classification scheme (<http://unstats.un.org/unsd/methods/m49/m49regin.htm#europe>).

Thus, each grid cell includes (1) a daily time-series of 0–3 day lag maximum AT for each climate change projection; (2) a grid cell specific temperature morbidity threshold and a location-based RR coefficient; and (3) population. For each grid cell, the AT was compared to the specific threshold for each day in the two time periods and risk estimates were calculated for each day. The expected number of daily RHAs in each cell was calculated using the population and the expected daily number of RHAs per capita, based on the national average of the grid cell.

$$RHA_{i,t} = RR_i^{(AT_{i,t} - Thres)_+} * Pop_i * RHAp_c_i$$

where $RHA_{i,t}$ is the number of RHAs attributable to heat in grid cell i at time t , RR_i the RR coefficient in grid cell i , $(AT_{i,t} - Thres)_+$ the positive difference between the 0–3 lag AT and the threshold for grid cell i at time t , Pop_i and $RHAp_c_i$ the population and expected number RHAs per capita in grid cell i .

For each country, the estimated number of RHAs attributed to heat was calculated for each climate change projection, for both the reference and the future period. The estimated number of RHAs

1
2
3 attributed to heat was then transformed into the proportion of the expected annual number RHAs
4
5 for each country (data available from the authors). Mean estimates were calculated for each region
6
7 for each climate change projection.
8
9

10 11 **Sensitivity analysis**

12 Because the study results could heavily depend on the RR coefficients and thresholds used, we assess
13 whether the estimates appeared sensitive to the region-specific meta-coefficients. We investigated
14 the grids that contained the cities in the PHEWE study using both the location (Mediterranean or
15 North-Continental) and city specific RR coefficients.³ We then compared the results with respect to
16
17 the attributed proportion of RHAs and the proportional change in heat related RHAs.
18
19
20
21
22
23
24
25

26 **RESULTS**

27
28 The periods of warm days will increase in the future. For the cities included in the sensitivity analysis,
29 the temperature threshold was exceeded by 20 days in the baseline period and 40 days in the future
30 period, on average each year.
31
32
33
34

35 In the future period approximately 0.4% of the annual numbers of RHAs in Europe was estimated to
36 be due to heat (Table 1), based on the mean estimates over the climate change projections. In
37 absolute terms, assuming all else equal, this represents about 26,000 cases annually in Europe. This
38 should be compared to the reference period where approximately 0.18% of all RHAs were attributed
39 to heat or about 11,000 cases annually. Thus, the results suggest more than a relative doubling of the
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
RHAs attributed to heat in Europe.

Table 1. The estimated proportion of RHAs attributed to heat for each region. Intervals describe the highest and lowest estimate in each region

Region	1981–2010	2021–2050	Change
Eastern Europe	0.17% (0.16% – 0.19%)	0.31% (0.29% – 0.35%)	0.14% (0.11% – 0.17%)
Northern Europe	0.13% (0.10% – 0.15%)	0.27% (0.19% – 0.32%)	0.14% (0.07% – 0.17%)
Southern Europe	0.23% (0.18% – 0.26%)	0.64% (0.42% – 0.68%)	0.41% (0.23% – 0.44%)
Western Europe	0.18% (0.16% – 0.20%)	0.39% (0.34% – 0.45%)	0.21% (0.17% – 0.26%)
EU27	0.18% (0.10% – 0.26%)	0.40% (0.19% – 0.68%)	0.21% (0.07% – 0.44%)

On the regional level, the five projections estimate increases in the number of heat related RHAs for Europe (Table 2). However, in one climate change projection, The Czech Republic, Hungary, Poland and Slovakia were estimated to have a decrease in heat related RHAs (Fig 1). The countries with the highest estimated increase also show the largest range between the highest and lowest estimate. The Scandinavian and Baltic countries show the smallest range between the highest and lowest estimates along with small increases of the mean estimates.

Table 2. Future increase in heat related RHAs based on the four climate models, under two emission scenarios, as the proportion of the annual expected number of RHAs in each region

Global climate model	CCSM3	ECHAM5	HadCM3	ECHAM4	ECHAM5
Greenhouse gas scenario	A1B		A2		
Eastern Europe	0.32%	0.08%	0.18%	0.12%	0.01%
Northern Europe	0.17%	0.09%	0.14%	0.20%	0.08%
Southern Europe	0.51%	0.29%	0.64%	0.45%	0.14%
Western Europe	0.30%	0.11%	0.26%	0.29%	0.06%
EU27	0.32%	0.13%	0.29%	0.26%	0.07%

There is variation among countries, with the largest increases in the Southern European countries and the smallest increase in Eastern Europe (Table 2). The relative change in the burden of RHA's in relation to the climate change scenarios investigated indicate a larger relative increase in

1
2
3 Mediterranean countries (approximately three times) compared to Northern European countries
4
5 (approximately two times).
6
7

8 9 **Sensitivity analysis**

10 The calculations using the city specific RR coefficients yielded a different proportion and number of
11
12 estimated RHAs than the ones using the two meta-coefficients, as expected. The relative changes in
13
14 the number of hospital admissions attributed to heat between the two time-periods were not
15
16 affected by the change of the RR coefficients.
17
18
19

20 21 **DISCUSSION**

22 The analysis estimates that the number of RHAs will increase in a warmer climate with more hot
23
24 days. Heat related RHAs were projected to increase two- to three-fold in the future due to climate
25
26 change. However, the proportion of respiratory admissions attributed to heat would remain rather
27
28 small. As most heat related health outcomes occur during the warmest period of the year, presenting
29
30 the increase as a change in the proportion of the total number of annual RHAs can underestimate the
31
32 additional burden on the health care system during summer. As the threshold was exceeded for
33
34 approximately 40 days in one year, an annual increase of 0.21% would result in a 1.9% increase on
35
36 average during these 40 days. The annual numbers are used because the available baseline rates of
37
38 RHAs are expressed as an annual average. Applying results to sub-national scales could be
39
40 inappropriate because national averages summarize over considerable heterogeneity.
41
42
43
44
45

46 The results suggest a larger impact from heat in Southern Europe in the future period, centred on
47
48 year 2035, than in the eastern and northern parts. This is in line with many climate change
49
50 projections showing a larger relative increase in the number of extremely hot days in southern
51
52 Europe compared with northern Europe.¹² However, to some extent this might also be explained by
53
54 model bias for northern Europe introduced by the RCA3 model, where the model appears to
55
56 underestimate temperature for the warmest days.¹⁷ This temperature bias is present in the reference
57
58
59
60

1
2
3 period, future period, and threshold values; therefore, the estimated numbers of RHAs from the two
4
5 periods within each scenario are comparable, but comparisons within the same time-period, across
6
7 scenarios may be inaccurate.
8
9

10
11 As the estimates are based on the population size in each grid cell, the added burden will be larger in
12
13 countries with larger increases in temperatures in densely populated areas. Because the population
14
15 of a city is considered a good predictor of the size of the urban heat island,³¹ an increase in the
16
17 population living in urban will increase the numbers exposed and the temperature to which they are
18
19 exposed. Heat islands increase temperatures in urbanized areas compared to surrounding areas, also
20
21 reducing cooling during the night time.³² These factors combined are likely to magnify the health
22
23 burden during a heatwave.³³ The coarse spatial scale of the climate models makes them unable to
24
25 take the urban heat island effect into account. Together with the urbanization of Europe, this could
26
27 potentially result in underestimation of actual consequences/RHAs in the future period because the
28
29 same population size and composition were assumed in both periods. In addition, climate change is
30
31 likely to increase ozone concentrations that would add an additional health burden to people at risk
32
33 of respiratory diseases. The number of deaths and RHAs, due to a change in ozone, is expected to
34
35 change in the future.³⁴
36
37
38

39
40 We used the same thresholds for the reference and future period to isolate the effect of climate
41
42 change. This would tend to overestimate the increased impact because there will undoubtedly be
43
44 biological and/or social adaptations that will reduce future health temperature-related burdens.
45
46 However, given the uncertainty of such adaptation effects, we choose to not incorporate such effects
47
48 in the projections. Studies from the U.S show that the heat related health burdens decreased over
49
50 the last 50 years,³⁵ indicating that some adaptation is taking place. An opposite trend appears to be
51
52 occurring in Stockholm, with an increase of the risk of mortality associated with heat during the
53
54 1990s.³⁶ The magnitude and the extent of future adaptation is, of course, highly uncertain, and will
55
56 vary between and within countries. Nevertheless, cities with higher thresholds seem to have higher
57
58
59
60

1
2
3 risk ratio coefficients.^{3 37} In an effort to estimate future and presumably higher thresholds, one must
4
5 also adjust the risk ratio coefficients. This would result in fewer days of elevated risk, but the risk
6
7 increase on each occasion could be higher due to the higher RRs.
8
9

10
11 Table 1 shows how the results vary by global climate models and greenhouse gas scenarios. This
12
13 study exemplifies the magnitude of the difference between projections made by a model with
14
15 different initial conditions and between different models with the same initial conditions. The results
16
17 indicate the range of these estimates is large. The mean increase over the five projections however,
18
19 provides confidence that the number of RHAs will increase over Europe as the climate continues to
20
21 change.
22
23

24
25 To improve the estimates of the impacts of heat of RHAs, detailed data on emergency room visits
26
27 and admissions during summer are needed, such as the proportion of emergency department cases
28
29 admitted to hospital, and the fatality rate. A better understanding of vulnerable groups is needed,
30
31 including how these groups could change over time. For example, the portion of the population aged
32
33 over 65 will increase from 17 to 29%³⁸ by 2050, significantly increasing the size of this vulnerable
34
35 group.
36
37

38
39 Normal weather patterns in the future will be different than those of today, both for average
40
41 temperature and extreme weather events. As the number of heat related RHAs are expected to
42
43 increase on the sub-continental scale, additional national or regional projections of the future health
44
45 burden from heat are needed, taking into account possible changes in exposure and vulnerability.
46
47
48

49 50 **CONCLUSIONS**

51
52 Projected changes in temperature and the number of extreme heat events with climate change could
53
54 substantially influence respiratory morbidity across Europe. Analyses projected that both the future
55
56 proportion of annual RHAs attributed to heat and the relative change in heat related RHAs will be
57
58 largest in Southern Europe, where they are expected to nearly triple. Eastern Europe can expect the
59
60

1
2
3 smallest increase in heat related RHAs, where they are estimated to approximately double. For all of
4
5 Europe, the number of respiratory heat-related hospital admissions is projected to be 26,000
6
7 annually in the future period (2021–2050) compared with 11,000 in the reference period (1981–
8
9 2010).
10

11
12
13
14
15
16
17
18
19 **Funding Statement** : This work was supported by the EU-funded Climate-Trap project (contract EAHC
20
21 20081108) and by the Swedish Environmental Protection Agency through the research programme
22
23 CLEO – Climate Change and Environmental Objectives. We would also like to acknowledge Estonia's
24
25 Ministry of Education for providing resources to H. Orru with the grant SF0180060s09.
26

27
28 **Contribution statement:** CÅ, BF, HO has planned and designed the study. GS represented SMHI and
29
30 provided climate data. CÅ ran the analyses, drafted the first manuscript and is guarantor. All authors
31
32 have participated in the interpretation of results and revised the manuscript.
33

34
35
36 The specific estimates for each of the EU27 countries, for each of the climate projections is available
37
38 after contact with corresponding author.
39

40
41 **Competing Interests Statement:** There are no competing interests .
42

43
44 **Data Sharing Statement:** The specific estimates for each of the EU27 countries, for each of the
45
46 climate projections is available after contact with corresponding author.
47
48

REFERENCES


1. Loddenkemper R, Gibson G, Sibille Y. European lung white book. *European Respiratory Society/European Lung Foundation, Sheffield, UK, ERSJ* 2003:1-182.
2. Ayres JG, Forsberg B, Annesi-Maesano I, et al. Climate change and respiratory disease: European Respiratory Society position statement. *Eur Respir J* 2009;34(2):295-302.
3. Michelozzi P, Accetta G, De Sario M, et al. High temperature and hospitalizations for cardiovascular and respiratory causes in 12 European cities. *Am J Respir Crit Care Med* 2009;179(5):383-9.
4. Rocklöv J, Forsberg B. Comparing approaches for studying the effects of climate extremes - a case study of hospital admissions in Sweden during an extremely warm summer. *Glob Health Action* 2009;2.
5. Kovats RS, Hajat S. Heat stress and public health: a critical review. *Annu Rev Public Health* 2008;29:41-55.
6. Knowlton K, Rotkin-Ellman M, King G, et al. The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect* 2009;117(1):61-7.
7. Rey G, Jouglu E, Fouillet A, et al. The impact of major heat waves on all-cause and cause-specific mortality in France from 1971 to 2003. *Int Arch Occup Environ Health* 2007;80(7):615-26.
8. Ye X, Wolff R, Yu W, et al. Ambient Temperature and Morbidity: A Review of Epidemiological Evidence. *Environ Health Perspect* 2011.
9. Semenza JC, McCullough JE, Flanders WD, et al. Excess hospital admissions during the July 1995 heat wave in Chicago. *Am J Prev Med* 1999;16(4):269-77.
10. Dhainaut JF, Claessens YE, Ginsburg C, et al. Unprecedented heat-related deaths during the 2003 heat wave in Paris: consequences on emergency departments. *Crit Care* 2004;8(1):1-2.
11. IPCC. *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation - Summary for Policymakers*. Geneva, Switzerland: World Meteorological Organization, 2012.
12. Nikulin G, Kjellström E, Hansson ULF, et al. Evaluation and future projections of temperature, precipitation and wind extremes over Europe in an ensemble of regional climate simulations. *Tellus A* 2011;63(1):41-55.
13. Schar C, Vidale PL, Luthi D, et al. The role of increasing temperature variability in European summer heatwaves. *Nature* 2004;427(6972):332-6.
14. Stott PA, Stone DA, Allen MR. Human contribution to the European heatwave of 2003 (vol 432, pg 610, 2004). *Nature* 2005;436(7054):1200-00.
15. Rebetez M, Dupont O, Giroud M. An analysis of the July 2006 heatwave extent in Europe compared to the record year of 2003. *Theoretical and Applied Climatology* 2009;95(1):1-7.
16. Barriopedro D, Fischer EM, Luterbacher J, et al. The Hot Summer of 2010: Redrawing the Temperature Record Map of Europe. *Science* 2011.
17. Samuelsson P, Jones CG, Willen U, et al. The Rossby Centre Regional Climate model RCA3: model description and performance. *Tellus Series a-Dynamic Meteorology and Oceanography* 2011;63(1):4-23.
18. Collins WD, Bitz CM, Blackmon ML, et al. The Community Climate System Model version 3 (CCSM3). *J Climate* 2006;19(11):2122-43.
19. Roeckner E, Bäuml G, Bonaventura L, et al. The atmospheric general circulation model ECHAM 5. PART I: Model description. Hamburg, Germany: MPI für Meteorologie, 2003.
20. Johns TC, Durman CF, Banks HT, et al. The new Hadley Centre Climate Model (HadGEM1): Evaluation of coupled simulations. *J Climate* 2006;19(7):1327-53.
21. Roeckner E, Arpe K, Bengtsson, et al. The atmospheric general circulation model {ECHAM}-4: {M}odel description and simulation of present-day climate, 1996:103-23.
22. Nakicenovic N, Alcamo J, Davis G, et al. Special report on emissions scenarios: a special report of Working Group III of the Intergovernmental Panel on Climate Change: Pacific Northwest

- 1
2
3 National Laboratory, Richland, WA (US), Environmental Molecular Sciences Laboratory (US),
4 2000.
- 5 23. Ballester J, Robine JM, Herrmann FR, et al. Long-term projections and acclimatization scenarios of
6 temperature-related mortality in Europe. *Nat Commun* 2011;2:358.
- 7 24. Knowlton K, Lynn B, Goldberg RA, et al. Projecting heat-related mortality impacts under a
8 changing climate in the New York City region. *American Journal of Public Health*
9 2007;97(11):2028-34.
- 10 25. Baccini M, Kosatsky T, Analitis A, et al. Impact of heat on mortality in 15 European cities:
11 attributable deaths under different weather scenarios. *J Epidemiol Community Health*
12 2011;65(1):64-70.
- 13 26. Beguin A, Hales S, Rocklov J, et al. The opposing effects of climate change and socio-economic
14 development on the global distribution of malaria. *Global Environmental Change-Human and*
15 *Policy Dimensions* 2011;21(4):1209-14.
- 16 27. Goldewijk KK, Beusen A, Janssen P. Long-term dynamic modeling of global population and built-
17 up area in a spatially explicit way: HYDE 3.1. *Holocene* 2010;20(4):565-73.
- 18 28. Lawrence MG. The Relationship between Relative Humidity and the Dewpoint Temperature in
19 Moist Air: A Simple Conversion and Applications. *Bulletin of the American Meteorological*
20 *Society* 2005;86(2):225-33.
- 21 29. Parsons KC. *Human thermal environments: the effects of hot, moderate, and cold environments*
22 *on human health, comfort, and performance*: Taylor & Francis, 2003.
- 23 30. Steadman RG. A Universal Scale of Apparent Temperature. *J Clim Appl Meteorol*
24 1984;23(12):1674-87.
- 25 31. Oke TR. City size and the urban heat island. *Atmospheric Environment (1967)* 1973;7(8):769-79.
- 26 32. Holmer B, Thorsson S, Eliasson I. COOLING RATES, SKY VIEW FACTORS AND THE DEVELOPMENT
27 OF INTRA - URBAN AIR TEMPERATURE DIFFERENCES. *Geografiska Annaler: Series A, Physical*
28 *Geography* 2007;89(4):237-48.
- 29 33. Nicholls N, Skinner C, Loughnan M, et al. A simple heat alert system for Melbourne, Australia. *Int J*
30 *Biometeorol* 2008;52(5):375-84.
- 31 34. Orru H, Andersson C, Ebi KL, et al. Impact of climate change on ozone related mortality and
32 morbidity in Europe (Accepted). *Eur Respir J* 2012.
- 33 35. Davis RE, Knappenberger PC, Michaels PJ, et al. Changing heat-related mortality in the United
34 States. *Environmental health perspectives* 2003;111(14):1712.
- 35 36. Rocklov J, Forsberg B, Meister K. Winter mortality modifies the heat-mortality association the
36 following summer. *Eur Respir J* 2009;33(2):245-51.
- 37 37. Baccini M, Biggeri A, Accetta G, et al. Heat effects on mortality in 15 European cities.
38 *Epidemiology* 2008;19(5):711-9.
- 39 38. Giannakouris K. Ageing characterises the demographic perspectives of the European societies.
40 *Statistics in focus* 2008;72:2008.
- 41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Figure 1. The range of the absolute increase in RHAs attributed to heat between the two periods**
4 **(1981-2010, 2021-2050) as proportion the annual expected number of RHAs for each of the 27**
5 **countries. The points show the highest and lowest estimate from four climate models under two**
6 **emission scenarios.**
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



406x279mm (200 x 200 DPI)

view only



Heat related respiratory hospital admissions in Europe in a changing climate: A health impact assessment

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2012-001842.R1
Article Type:	Research
Date Submitted by the Author:	16-Nov-2012
Complete List of Authors:	Åström, Christofer; Umeå University, Department of Public Health and Clinical Medicine Orru, Hans Rocklöv, Joacim; Umeå University, Department of Public Health and Clinical Medicine Strandberg, Gustav; Swedish Meteorological and Hydrological Institute, Ebi, Kristie; Stanford University, ClimAdapt Forsberg, Bertil; Umeå University, Department of Public Health and Clinical Medicine
Primary Subject Heading:	Epidemiology
Secondary Subject Heading:	Epidemiology, Occupational and environmental medicine
Keywords:	EPIDEMIOLOGY, PUBLIC HEALTH, Epidemiology < THORACIC MEDICINE

SCHOLARONE™
Manuscripts

Heat related respiratory hospital admissions in Europe in a changing climate

Authors:

Christofer Åström^{1*}, Hans Orru^{1,2}, Joacim Rocklöv^{1,3}, Gustav Strandberg⁴, Kristie L. Ebi⁵, Bertil Forsberg¹

Affiliations:

¹ Department of Public Health and Clinical Medicine, Occupational and Environmental Medicine, Umeå University, Umeå, Sweden; ² Department of Public Health, University of Tartu, Tartu, Estonia; ³ Department of Public Health and Clinical Medicine, Epidemiology and Global Health, Umeå University, Umeå, Sweden; ⁴ Swedish Meteorological and Hydrological Institute, Norrköping, Sweden; ⁵ ClimAdapt, Stanford University, Los Altos, California, USA

Corresponding author:

* Author to whom correspondence should be addressed:

Christofer Åström,

Occupational and Environmental Medicine, Department of Public Health and Clinical Medicine, Umeå University, SE 901 87, Sweden

+4690 785 0874

christofer.astrom@envmed.umu.se

Keywords: Climate change, respiratory health, public health, heat

Word count: 3045

ARTICLE SUMMARY

Article focus

To assess how heat related respiratory hospital admissions in Europe are expected to change with the global warming in the near future. With a range of climate change projections explore the range of the different impact estimates.

Key messages

The 30 year mean annual increase in respiratory hospital admissions related to heat between the periods 1981-2010 and 2021-2050 can be counted in 10's of thousands.

The increase will be different in different parts of Europe with the largest relative increase in Southern Europe.

Strengths and limitations

The study takes the spatial variation in both climate and population density into account in the impact calculations. This makes the estimates valid for countries with large national differences in population densities and climate.

The study results exemplifies some the variation of the impacts depending on different climate change scenarios as well as the variations within one climate scenario depending on the underlying climate model.

The use of real spatial population data has limited the study to explore the change in all age respiratory hospital admissions without demographic trends. The fact that the elderly is a well-known risk group when studying heat related illness combined with the future rise of the proportion of elderly people in the population may suggest that the estimated impacts are rather conservative

ABSTRACT

Respiratory diseases are ranked second in Europe in terms of mortality, prevalence and costs. Studies have shown that extreme heat has a large impact on mortality and morbidity, with a large relative increase for respiratory diseases. Expected increases in mean temperature and the number of extreme heat events over the coming decades due to climate change raise questions about the possible health impacts.

Heat-related respiratory hospital admissions under a changing climate are projected using multi-city epidemiologic exposure-response relationships applied to gridded population data and country-specific baseline respiratory hospital admission rates. Times-series of temperatures are simulated with a regional climate model based on four global climate models, under two greenhouse gas emission scenarios (A1B & A2).

Between a reference period (1981–2010) and a future period (2021–2050), the total number of respiratory hospital admissions attributed to heat is projected to be larger in Southern Europe, with three-times more heat attributed respiratory hospital admissions in the future period. The smallest change was estimated in Eastern Europe with about a two-fold increase. For all of Europe, the number of respiratory heat-related hospital admissions is projected to be 26,000 annually in the future period compared with 11,000 in the reference period.

The results suggest that projected effects of climate change on temperature and the number of extreme heat events could substantially influence respiratory morbidity across Europe.

INTRODUCTION

Respiratory diseases are ranked second in Europe in terms of mortality, prevalence and costs.¹ This burden is expected to increase, partly due to a changing climate.² An environmental factor with a large impact on mortality and morbidity in Europe is extreme heat, with large effects on respiratory diseases.³⁻⁵ Physiological effects of exposure to heat can be directly heat related (heat stroke, heat fatigue and dehydration) or can contribute to worsening of respiratory and cardiovascular diseases, electrolyte disorders, and kidney problems.⁴⁻⁷ The reasons for an increase in respiratory admissions may be several. Elderly with respiratory diseases such as COPD are less fit and suffer often from circulatory problems. Heat influenced admissions due to chronic airway obstruction and asthma increased more than admissions due to chronic bronchitis in a study from New York City,⁸ but the daily number of admissions seldom allow a study of specific diagnoses. In a recent expert elicitation amongst European researchers engaged in environmental medicine or respiratory health, extreme heat stood out as most important climate related pathway to adverse impacts on respiratory health, more important than changes in air pollutants and allergens.⁹ A review found that heatwaves have a stronger relative impact on mortality than on emergency room visits and hospital admissions, suggesting that many individuals die before they can get to the hospital.¹⁰ However, several studies confirm that heat also affects health care utilisation.^{3 11} During the heatwave of 2003 in France, the Assistance Public Hôpitaux de Paris recorded 2,400 additional visits to the emergency care units and 1,900 additional hospital admissions.¹² During the heatwave in California in 2006 there were almost 1,200 hospital admissions together with more than 16,000 visits to the emergency departments.⁶

It is possible that hospital admissions could increase in the future due to temperature changes projected with climate change, assuming no additional acclimatisation. Projected changes include an increase in global mean temperature of 1.8°C to 4.0°C by the end of this century, with larger increases over land areas at high latitudes, changing seasonal temperatures, and increases in the frequency, intensity, duration, and spatial extent of heatwaves.¹³ Recent projections for 2070–2100

1
2
3 suggest that maximum temperatures experienced once every 20 years during the period 1961–1990
4
5 could be expected as often as every year in southern Europe and every 3rd-5th year in northern
6
7 Europe.¹⁴ This means that Europe should expect an increase in both mean temperatures and the
8
9 number of extreme heat events. Recent heatwaves are consistent with projections. For example, the
10
11 heatwave that occurred in Europe 2003 could be expected to return every 46,000 years, based on
12
13 the temperature distribution for the years 1864–2000.¹⁵ However, such extreme hot conditions may
14
15 become much more common in the end of this century due to anthropogenic climate changes.¹⁶ The
16
17 uncertainty in this estimated return time is quite high, with a lower bound of the 90% confidence
18
19 interval of 9,000 years. Even so, another heatwave occurred in 2006 with the most anomalous July
20
21 temperature ever measured in Europe.¹⁷ In 2010, Eastern Europe experienced a heatwave with
22
23 summer temperatures higher than the last 140 years, resulting in roughly 55,000 excess deaths in
24
25 Russia.¹⁸

26
27
28
29
30 The possible impacts of these changes on morbidity, particularly respiratory health, are relatively
31
32 unexplored.

33 34 35 36 **Aim**

37 The aim of this study is to assess the extent to which changes in the frequency of hot days due to
38
39 climate change over the next 40 years could affect heat related respiratory hospital admissions
40
41 (RHAs) in Europe. Using a range of climate projections, we estimate the change in hospital
42
43 admissions between a reference period and a future period.

44 45 46 47 48 **METHODS**

49 50 51 52 **Climate change and temperature modelling**

53 The Rossby Centre regional atmospheric climate model RCA3¹⁹ was developed by SMHI (Swedish
54
55 meteorological and hydrological institute) to dynamically downscale results from the global climate
56
57 models CCSM3²⁰, ECHAM5²¹, HadleyCM3²² and ECHAM4²³ to a higher resolution over Europe. RCA3 is
58
59
60

run on a horizontal grid spacing of 0.44° (corresponding to approximately 50 km) and a time step of 30 minutes. Projections are based on the global greenhouse gas emission scenarios A1B and A2 (scenarios are described in detail in the SRES²⁴). Both have been used in climate change health impact assessments (HIA).²⁵⁻²⁸ A1B is a “middle of the road” scenario and A2 is considered a high emission scenario, although recent greenhouse gas emissions have been higher. Data from one climate model, under one climate change scenario, is referred to as one climate change projection.

We use aggregated daily projections of maximum temperature and relative humidity data to estimate exposure for the periods 1981–2020 (reference period) and 2021–2050 (future period).

Population and morbidity rates

The annual country specific rate of respiratory hospital admissions between 2005 and 2010 were extracted from the WHO’s European Health for All Database (<http://data.euro.who.int/hfadb>) for the EU27 countries. The mean value over the six years was used as a baseline morbidity rate. Official population data were from the HYDE theme within the Netherlands Environmental Assessment Agency.²⁹ These data are gridded on a 0.0833° resolution (approximately 9,45km) and matched with the climatic data by summing the population within each climatic grid cell.

Exposure-response assumption

The impact calculations were based on the relationship between heat and RHAs (ICD-9:460–519) estimated within the European PHEWE (Assessment and Prevention of Acute Health Effects of Weather Conditions in Europe) project.³ This relationship was based on analyses of 12 European cities over the 1990s, using the daily maximum of apparent temperature (AT), where AT is a combination of the measured temperature and the dew point temperature:

$$AT = -2.653 + 0.944 * T + 0.0153 * (DT)^2$$

where T is the air temperature and DP the dew point temperature, which can be derived from temperature and relative humidity.³⁰ The AT accounts for heat stress related not only to the absolute

1
2
3 temperature but also to the saturation of the surrounding air that makes it harder to regulate body
4
5 temperature by sweating, the most important mechanism to maintain a healthy body temperature.³¹
6

7
8 ³² The PHEWE model also included potential confounders such as air pollution, holidays, week days
9
10 etc. The study used a 0–3 day lag of the maximum AT and concluded that the 90th percentile of this
11
12 exposure variable for the summer months, April to September, is the appropriate threshold value for
13
14 the heat-morbidity function.³ We calculated the 90th percentile for the summer months for each grid
15
16 cell from the climate data for the 1990s, yielding individual thresholds for each grid cell for each
17
18 climate projection.
19

20
21 The PHEWE study calculated the relative risk (RR) coefficients for each of the 12 cities and then
22
23 combined these into two meta-coefficients, for North-Continental cities 1.012 (95% CI 1.001–1.022)
24
25 and Mediterranean cities 1.021 (95% CI 1.006–1.036), associated with a 1-degree increase above the
26
27 temperature threshold. Meta-coefficients were more suitable as they allowed calculation of the
28
29 change in RHAs for countries with and without a city in the PHEWE Study. We assigned the
30
31 Mediterranean coefficient to Portugal and the Mediterranean countries. The rest of Europe was
32
33 assigned the RR for the North continental cities. Statistically significant coefficients were found for all
34
35 ages and age 75+ in the PHEWE study. Because the population data were not stratified by age, we
36
37 used the PHEWE study coefficients for all ages.
38
39
40
41

42 Projections

43
44 The final data used in the impact calculations were based on the climate data grid with a resolution
45
46 of 0.44° resulting in 8,075 grid cells over Europe. The population data, RR coefficients and baseline
47
48 morbidity rates were projected for each grid cell. The countries within EU27 were grouped into four
49
50 regions, Northern, Western, Eastern and Southern Europe, according to the United Nations
51
52 classification scheme (<http://unstats.un.org/unsd/methods/m49/m49regin.htm#europe>).
53
54

55
56 Thus, each grid cell includes (1) a daily time-series of 0–3 day lag maximum AT for each climate
57
58 change projection; (2) a grid cell specific temperature morbidity threshold and a location-based RR
59
60

1
2
3 coefficient; and (3) population. For each grid cell, the AT was compared to the specific threshold for
4
5 each day in the two time periods and risk estimates were calculated for each day. The expected
6
7 number of daily RHAs in each cell was calculated using the population and the expected daily number
8
9 of RHAs per capita, based on the national average of the grid cell.
10

$$RHA_{i,t} = RR_i^{(AT_{i,t} - Thres_i)_+} * Pop_i * RHAp_c_i$$

11
12 where $RHA_{i,t}$ is the number of RHAs attributable to heat in grid cell i at time t , RR_i the RR coefficient in
13
14 grid cell i , $(AT_{i,t} - Thres_i)_+$ the difference between the 0–3 lag AT and the threshold for grid cell i at
15
16 time t if $AT_{i,t}$ were greater than the threshold and 0 otherwise, Pop_i and $RHAp_c_i$ the population and
17
18 expected number RHAs per capita in grid cell i .
19
20
21
22
23
24

25 For each country, the estimated number of RHAs attributed to heat was calculated for each climate
26
27 change projection, for both the reference and the future period. The estimated number of RHAs
28
29 attributed to heat was then transformed into the proportion of the expected annual number RHAs
30
31 for each country (data available from the authors). Mean estimates were calculated for each region
32
33 for each climate change projection.
34
35
36

37 Sensitivity analysis

38 Because the study results could heavily depend on the RR coefficients and thresholds used, we assess
39
40 whether the estimates appeared sensitive to the region-specific meta-coefficients. We investigated
41
42 the grids that contained the cities in the PHEWE study using both the location (Mediterranean or
43
44 North-Continental) and city specific RR coefficients.³ We then compared the results with respect to
45
46 the attributed proportion of RHAs and the proportional change in heat related RHAs.
47
48
49
50
51
52

53 RESULTS

54
55
56
57
58
59
60

The periods of warm days will increase in the future. For the cities included in the sensitivity analysis, the temperature threshold was exceeded by 20 days in the baseline period and 40 days in the future period, on average each year.

In the future period approximately 0.4% of the annual numbers of RHAs in Europe was estimated to be due to heat (Table 1), based on the mean estimates over the climate change projections. In absolute terms, assuming all else equal, this represents about 26,000 cases annually in Europe. This should be compared to the reference period where approximately 0.18% of all RHAs were attributed to heat or about 11,000 cases annually. Thus, the results suggest more than a relative doubling of the RHAs attributed to heat in Europe.

Table 1. The estimated proportion of RHAs attributed to heat for each region. Intervals describe the highest and lowest estimate in each region

Region	1981–2010	2021–2050	Change
Eastern Europe	0.17% (0.16% – 0.19%)	0.31% (0.29% – 0.35%)	0.14% (0.11% – 0.17%)
Northern Europe	0.13% (0.10% – 0.15%)	0.27% (0.19% – 0.32%)	0.14% (0.07% – 0.17%)
Southern Europe	0.23% (0.18% – 0.26%)	0.64% (0.42% – 0.68%)	0.41% (0.23% – 0.44%)
Western Europe	0.18% (0.16% – 0.20%)	0.39% (0.34% – 0.45%)	0.21% (0.17% – 0.26%)
EU27	0.18% (0.10% – 0.26%)	0.40% (0.19% – 0.68%)	0.21% (0.07% – 0.44%)

On the regional level, the five projections estimate increases in the number of heat related RHAs for Europe (Table 2). However, in one climate change projection, The Czech Republic, Hungary, Poland and Slovakia were estimated to have a decrease in heat related RHAs (Fig 1). The countries with the highest estimated increase also show the largest range between the highest and lowest estimate. The Scandinavian and Baltic countries show the smallest range between the highest and lowest estimates along with small increases of the mean estimates.

Table 2. Future increase in heat related RHAs based on the four climate models, under two emission scenarios, as the percentage of the annual expected number of RHAs in each region

Climate model	CCSM3	ECHAM5	HadCM3	ECHAM4	ECHAM5
<i>Greenhouse gas scenario</i>	<i>A1B</i>		<i>A2</i>		
Eastern Europe	0.32%	0.08%	0.18%	0.12%	0.01%
Northern Europe	0.17%	0.09%	0.14%	0.20%	0.08%
Southern Europe	0.51%	0.29%	0.64%	0.45%	0.14%
Western Europe	0.30%	0.11%	0.26%	0.29%	0.06%
EU27	0.32%	0.13%	0.29%	0.26%	0.07%

There is variation among countries, with the largest increases in the Southern European countries and the smallest increase in Eastern Europe (Table 2). The relative change in the burden of RHA's in relation to the climate change scenarios investigated indicate a larger relative increase in Mediterranean countries (approximately three times) compared to Northern European countries (approximately two times).

Sensitivity analysis

The calculations using the city specific RR coefficients yielded a different proportion and number of estimated RHAs than the ones using the two meta-coefficients, as expected. However, the relative changes in the number of hospital admissions attributed to heat between the two time-periods were not affected by the change of the RR coefficients.

DISCUSSION

The analysis estimates that the number of RHAs will increase in a warmer climate with more hot days. Heat related RHAs were projected to increase two- to three-fold in the future due to climate change. However, the proportion of respiratory admissions attributed to heat would remain rather small. This projection is in line with the results of a recent study which estimates that the respiratory admissions due to excessive heat in New York State will increase 2-6 times from the period 1991-2004 to the period 2080-2099.³³

1
2
3 As most heat related health outcomes occur during the warmest period of the year, presenting the
4 increase as a change in the proportion of the total number of annual RHAs can underestimate the
5 additional burden on the health care system during summer. As the threshold was exceeded for
6 approximately 40 days in one year, an annual increase of 0.21% would result in a 1.9% increase on
7 average during these 40 days. The annual numbers are used because the available baseline rates of
8 RHAs are expressed as an annual average. Applying results to sub-national scales could be
9 inappropriate because national averages summarize over considerable heterogeneity.
10
11

12
13
14 The results suggest a larger impact from heat in Southern Europe in the future period, centred on
15 year 2035, than in the eastern and northern parts. This is in line with many climate change
16 projections showing a larger relative increase in the number of extremely hot days in southern
17 Europe compared with northern Europe.¹⁴ However, to some extent this might also be explained by
18 model bias for northern Europe introduced by the RCA3 model, where the model appears to
19 underestimate temperature for the warmest days.¹⁹ This temperature bias is present in the reference
20 period, future period, and threshold values; therefore, the estimated numbers of RHAs from the two
21 periods within each scenario are comparable, but comparisons within the same time-period, across
22 scenarios may be inaccurate.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39
40 As the estimates are based on the population size in each grid cell, the added burden will be larger in
41 countries with larger increases in temperatures in densely populated areas. Because the population
42 of a city is considered a good predictor of the size of the urban heat island,³⁴ an increase in the
43 population living in urban areas will increase the numbers exposed and the temperature to which
44 they are exposed. Heat islands increase temperatures in urbanized areas compared to surrounding
45 areas, also reducing cooling during the night time.³⁵ These factors combined are likely to magnify the
46 health burden during a heatwave.³⁶ The coarse spatial scale of the climate models makes them
47 unable to take the urban heat island effect into account. Together with the urbanization of Europe,
48 this could potentially result in underestimation of actual consequences/RHAs in the future period
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 because the same population size and composition were assumed in both periods. In addition,
4
5 climate change is likely to increase ozone concentrations that would add an additional health burden
6
7 for people at risk of respiratory diseases. The number of deaths and RHAs, due to a change in ozone,
8
9 is expected to change in the future.³⁷ A sensitivity analysis in the PHEWE study however showed that
10
11 the exposure-response relationship for heat did not substantially differ between models taking ozone
12
13 into account and models adjusting for NO₂ alone.³
14

15
16 We used the same thresholds for the reference and future period to isolate the effect of climate
17
18 change. This would tend to overestimate the increased impact because there will undoubtedly be
19
20 biological and/or social adaptations that will reduce future health temperature-related burdens.
21
22 However, given the uncertainty of such adaptation effects, we choose to not incorporate such effects
23
24 in the projections. Recent studies in the U.S indicate heat related health burdens have decreased,³⁸
25
26 indicating that some adaptation is taking place. An opposite trend appears to be occurring in
27
28 Stockholm, with an increase of the risk of mortality associated with heat during the 1990s.³⁹ A study
29
30 looking at the impact of heat on mortality, before and after the implementation of a heat warning
31
32 system, in Italy shows that the effects of extreme heat can be reduced while the effects of moderate
33
34 increased average temperatures remains similar.⁴⁰ The magnitude and the extent of future
35
36 adaptation is, of course, highly uncertain, and will vary between and within countries. Nevertheless,
37
38 cities with higher thresholds seem to have higher risk ratio coefficients.^{3 41} In an effort to estimate
39
40 future and presumably higher thresholds, one must also adjust the risk ratio coefficients. This would
41
42 result in fewer days of elevated risk, but the risk increase on each occasion could be higher due to
43
44 the higher RRs.
45
46
47
48

49
50 Table 1 shows how the results vary by global climate models and greenhouse gas scenarios. This
51
52 study exemplifies the magnitude of the difference between projections made by a model with
53
54 different initial conditions and between different models with the same initial conditions. The results
55
56 indicate the range of these estimates is large. The mean increase over the five projections however,
57
58
59
60

1
2
3 provides confidence that the number of RHAs will increase over Europe as the climate continues to
4
5 change.
6

7
8 To improve the estimates of the impacts of heat on RHAs, detailed data on emergency room visits
9
10 and admissions during summer are needed, such as the proportion of emergency department cases
11
12 admitted to hospital, and the fatality rate. A better understanding of vulnerable groups is needed,
13
14 including how these groups could change over time. For example, the portion of the population aged
15
16 over 65 will increase from 17 to 29%⁴² by 2050, significantly increasing the size of this vulnerable
17
18 group. COPD is mainly a disease of the elderly. Persons suffering from COPD appear especially
19
20 vulnerable to heat⁴³ and the incidence of COPD is likely to increase⁴⁴ with the consequence that the
21
22 health impacts of heat may increase in future years. This study was limited to estimating the impacts
23
24 on the total population because few age-specific relationships were reported and because age
25
26 stratified data were only available on a coarser spatial scale than the climate data, which would
27
28 reduce the benefits of having spatial climate data.
29
30

31
32
33 Normal weather patterns in the future will be different than those of today, both for average
34
35 temperature and extreme weather events. As the number of heat related RHAs are expected to
36
37 increase on the sub-continental scale, additional national or regional projections of the future health
38
39 burden from heat are needed, taking into account possible changes in exposure and vulnerability.
40
41

42 43 44 **CONCLUSIONS**

45
46 Projected changes in temperature and the number of extreme heat events with climate change could
47
48 substantially influence respiratory morbidity across Europe. Analyses projected that both the future
49
50 proportion of annual RHAs attributed to heat and the relative change in heat related RHAs will be
51
52 largest in Southern Europe, where they are expected to nearly triple. Eastern Europe can expect the
53
54 smallest increase in heat related RHAs, where they are estimated to approximately double. For all of
55
56 Europe, the number of respiratory heat-related hospital admissions is projected to be 26,000
57
58
59
60

1
2
3 annually in the future period (2021–2050) compared with 11,000 in the reference period (1981–
4
5 2010). The estimates presented rely on the assumption that no additional adaptation occurs. Future
6
7 studies should elaborate and quantify the possible effects of different adaptation assumptions
8
9 applied to regional conditions.
10

11
12
13
14
15
16
17
18
19 This work was supported by the EU-funded Climate-Trap project (contract EAHC 20081108) and by
20
21 the Swedish Environmental Protection Agency through the research programme CLEO – Climate
22
23 Change and Environmental Objectives. We would also like to acknowledge Estonia's Ministry of
24
25 Education for providing resources to H. Orru with the grant SF0180060s09.
26

27
28 Contribution statement: CÅ, BF, HO has planned and designed the study. GS represented SMHI and
29
30 provided climate data. CÅ ran the analyses, drafted the first manuscript and is guarantor. All authors
31
32 have participated in the interpretation of results and revised the manuscript.
33

34
35
36 The specific estimates for each of the EU27 countries, for each of the climate projections is available
37
38 after contact with corresponding author.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Loddenkemper R, Gibson G, Sibille Y. European lung white book. *European Respiratory Society/European Lung Foundation, Sheffield, UK, ERSJ* 2003;1-182.
2. Ayres JG, Forsberg B, Annesi-Maesano I, et al. Climate change and respiratory disease: European Respiratory Society position statement. *Eur Respir J* 2009;34(2):295-302.
3. Michelozzi P, Accetta G, De Sario M, et al. High temperature and hospitalizations for cardiovascular and respiratory causes in 12 European cities. *Am J Respir Crit Care Med* 2009;179(5):383-9.
4. Rocklöv J, Forsberg B. Comparing approaches for studying the effects of climate extremes - a case study of hospital admissions in Sweden during an extremely warm summer. *Glob Health Action* 2009;2.
5. Kovats RS, Hajat S. Heat stress and public health: a critical review. *Annu Rev Public Health* 2008;29:41-55.
6. Knowlton K, Rotkin-Ellman M, King G, et al. The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect* 2009;117(1):61-7.
7. Rey G, Jouglé E, Fouillet A, et al. The impact of major heat waves on all-cause and cause-specific mortality in France from 1971 to 2003. *Int Arch Occup Environ Health* 2007;80(7):615-26.
8. Lin S, Luo M, Walker RJ, et al. Extreme high temperatures and hospital admissions for respiratory and cardiovascular diseases. *Epidemiology* 2009;20(5):738-46.
9. Forsberg B, Braback L, Keune H, et al. An expert assessment on climate change and health - with a European focus on lungs and allergies. *Environ Health* 2012;11 Suppl 1:S4.
10. Ye X, Wolff R, Yu W, et al. Ambient Temperature and Morbidity: A Review of Epidemiological Evidence. *Environ Health Perspect* 2011.
11. Semenza JC, McCullough JE, Flanders WD, et al. Excess hospital admissions during the July 1995 heat wave in Chicago. *Am J Prev Med* 1999;16(4):269-77.
12. Dhainaut JF, Claessens YE, Ginsburg C, et al. Unprecedented heat-related deaths during the 2003 heat wave in Paris: consequences on emergency departments. *Crit Care* 2004;8(1):1-2.
13. IPCC. *Climate Change 2007 - The Physical Science Basis: Working Group I Contribution to the Fourth Assessment Report of the IPCC*. Cambridge, UK and New York, NY, USA: Cambridge University Press, 2007.
14. Nikulin G, Kjellström E, Hansson ULF, et al. Evaluation and future projections of temperature, precipitation and wind extremes over Europe in an ensemble of regional climate simulations. *Tellus A* 2011;63(1):41-55.
15. Schar C, Vidale PL, Luthi D, et al. The role of increasing temperature variability in European summer heatwaves. *Nature* 2004;427(6972):332-6.
16. Stott PA, Stone DA, Allen MR. Human contribution to the European heatwave of 2003. *Nature* 2004;432(7017):610-4.
17. Rebetez M, Dupont O, Giroud M. An analysis of the July 2006 heatwave extent in Europe compared to the record year of 2003. *Theoretical and Applied Climatology* 2009;95(1):1-7.
18. Barriopedro D, Fischer EM, Luterbacher J, et al. The Hot Summer of 2010: Redrawing the Temperature Record Map of Europe. *Science* 2011.
19. Samuelsson P, Jones CG, Willen U, et al. The Rossby Centre Regional Climate model RCA3: model description and performance. *Tellus Series a-Dynamic Meteorology and Oceanography* 2011;63(1):4-23.
20. Collins WD, Bitz CM, Blackmon ML, et al. The Community Climate System Model version 3 (CCSM3). *J Climate* 2006;19(11):2122-43.
21. Roeckner E, Bäuml G, Bonaventura L, et al. The atmospheric general circulation model ECHAM 5. PART I: Model description. Hamburg, Germany: MPI für Meteorologie, 2003.
22. Johns TC, Durman CF, Banks HT, et al. The new Hadley Centre Climate Model (HadGEM1): Evaluation of coupled simulations. *J Climate* 2006;19(7):1327-53.

23. Roeckner E, Arpe K, Bengtsson, et al. The atmospheric general circulation model {ECHAM}-4: {M}odel description and simulation of present-day climate, 1996:103-23.
24. Nakicenovic N, Alcamo J, Davis G, et al. Special report on emissions scenarios: a special report of Working Group III of the Intergovernmental Panel on Climate Change: Pacific Northwest National Laboratory, Richland, WA (US), Environmental Molecular Sciences Laboratory (US), 2000.
25. Ballester J, Robine JM, Herrmann FR, et al. Long-term projections and acclimatization scenarios of temperature-related mortality in Europe. *Nat Commun* 2011;2:358.
26. Knowlton K, Lynn B, Goldberg RA, et al. Projecting heat-related mortality impacts under a changing climate in the New York City region. *American Journal of Public Health* 2007;97(11):2028-34.
27. Baccini M, Kosatsky T, Analitis A, et al. Impact of heat on mortality in 15 European cities: attributable deaths under different weather scenarios. *J Epidemiol Community Health* 2011;65(1):64-70.
28. Beguin A, Hales S, Rocklov J, et al. The opposing effects of climate change and socio-economic development on the global distribution of malaria. *Global Environmental Change-Human and Policy Dimensions* 2011;21(4):1209-14.
29. Goldewijk KK, Beusen A, Janssen P. Long-term dynamic modeling of global population and built-up area in a spatially explicit way: HYDE 3.1. *Holocene* 2010;20(4):565-73.
30. Lawrence MG. The Relationship between Relative Humidity and the Dewpoint Temperature in Moist Air: A Simple Conversion and Applications. *Bulletin of the American Meteorological Society* 2005;86(2):225-33.
31. Parsons KC. *Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort, and performance*: Taylor & Francis, 2003.
32. Steadman RG. A Universal Scale of Apparent Temperature. *J Clim Appl Meteorol* 1984;23(12):1674-87.
33. Lin S, Hsu WH, Van Zutphen AR, et al. Excessive heat and respiratory hospitalizations in new york state: estimating current and future public health burden related to climate change. *Environ Health Perspect* 2012;120(11):1571-7.
34. Oke TR. City size and the urban heat island. *Atmospheric Environment (1967)* 1973;7(8):769-79.
35. Holmer B, Thorsson S, Eliasson I. COOLING RATES, SKY VIEW FACTORS AND THE DEVELOPMENT OF INTRA - URBAN AIR TEMPERATURE DIFFERENCES. *Geografiska Annaler: Series A, Physical Geography* 2007;89(4):237-48.
36. Nicholls N, Skinner C, Loughnan M, et al. A simple heat alert system for Melbourne, Australia. *Int J Biometeorol* 2008;52(5):375-84.
37. Orru H, Andersson C, Ebi KL, et al. Impact of climate change on ozone related mortality and morbidity in Europe. *Eur Respir J* 2012 Jun 27. [Epub ahead of print].
38. Davis RE, Knappenberger PC, Michaels PJ, et al. Changing heat-related mortality in the United States. *Environmental Health Perspectives* 2003;111(14):1712-18.
39. Rocklov J, Forsberg B, Meister K. Winter mortality modifies the heat-mortality association the following summer. *Eur Respir J* 2009;33(2):245-51.
40. Schifano P, Leone M, De Sario M, et al. Changes in the effects of heat on mortality among the elderly from 1998-2010: results from a multicenter time series study in Italy. *Environmental Health* 2012;11(1):58.
41. Baccini M, Biggeri A, Accetta G, et al. Heat effects on mortality in 15 European cities. *Epidemiology* 2008;19(5):711-9.
42. Giannakouris K. Ageing characterises the demographic perspectives of the European societies. *Statistics in focus* 2008;72:2008.
43. Schwartz J. Who is sensitive to extremes of temperature? A case-only analysis. *Epidemiology* 2005;16(1):67-72.
44. Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet* 2007;370(9589):765-73.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. The range of the absolute increase in RHAs attributed to heat between the two periods (1981-2010, 2021-2050) as proportion the annual expected number of RHAs for each of the 27 countries. The points show the highest and lowest estimate from four climate models under two emission scenarios.

For peer review only

BMJ Open: first published as 10.1136/bmjopen-2012-001842 on 24 January 2013. Downloaded from <http://bmjopen.bmj.com/> on April 17, 2024 by guest. Protected by copyright.

Heat related respiratory hospital admissions in Europe in a changing climate

Authors:

Christofer Åström^{1*}, Hans Orru^{1,2}, Joacim Rocklöv^{1,3}, Gustav Strandberg⁴, Kristie L. Ebi⁵, Bertil Forsberg¹

Affiliations:

¹ Department of Public Health and Clinical Medicine, Occupational and Environmental Medicine, Umeå University, Umeå, Sweden; ² Department of Public Health, University of Tartu, Tartu, Estonia; ³ Department of Public Health and Clinical Medicine, Epidemiology and Global Health, Umea University, Umeå, Sweden; ⁴ Swedish Meteorological and Hydrological Institute, Norrköping, Sweden; ⁵ ClimAdapt, Stanford University, Los Altos, California, USA

Corresponding author:

* Author to whom correspondence should be addressed:

Christofer Åström,

Occupational and Environmental Medicine, Department of Public Health and Clinical Medicine, Umeå University, SE 901 87, Sweden

+4690 785 0874

christofer.astrom@envmed.umu.se

Keywords: Climate change, respiratory health, public health, heat

Word count: 3045

ARTICLE SUMMARY

Article focus

To assess how heat related respiratory hospital admissions in Europe are expected to change with the global warming in the near future. With a range of climate change projections explore the range of the different impact estimates.

Key messages

The 30 year mean annual ~~mean~~ increase in respiratory hospital admissions related to heat between the periods 1981-2010 and 2021-2050 can be counted in 10's of thousands.

The increase will be different in different parts of Europe with the largest relative increase in Southern Europe.

Strengths and limitations

The study takes the spatial variation in both climate and population density into account in the impact calculations. This makes the estimates valid for countries with large national differences in population densities and climate.

The study results exemplifies some the variation of the impacts depending on different climate change scenarios as well as the variations within one climate scenario depending on the underlying climate model.

The use of real spatial population data has limited the study to explore the change in all age respiratory hospital admissions without demographic trends. The fact that the elderly is a wellknown well-known risk group when studying heat related illness combined with the future rise of the proportion of elderly people in the population may suggest that the estimated impacts are rather conservative.

ABSTRACT

Respiratory diseases are ranked second in Europe in terms of mortality, prevalence and costs. Studies have shown that extreme heat has a large impact on mortality and morbidity, with a large relative increase for respiratory diseases. Expected increases in mean temperature and the number of extreme heat events over the coming decades due to climate change raise questions about the possible health impacts.

Heat-related respiratory hospital admissions under a changing climate are projected using multi-city epidemiologic exposure-response relationships applied to gridded population data and country-specific baseline respiratory hospital admission rates. Times-series of temperatures are simulated with a regional climate model based on four global climate models, under two greenhouse gas emission scenarios- (A1B & A2).

Between a reference period (1981–2010) and a future period (2021–2050), the total number of respiratory hospital admissions attributed to heat is projected to be larger in Southern Europe, with three-times more heat attributed respiratory hospital admissions in the future period. The smallest change was estimated in Eastern Europe with about a two-fold increase. For all of Europe, the number of respiratory heat-related hospital admissions is projected to be 26,000 annually in the future period compared with 11,000 in the reference period.

The results suggest that projected effects of climate change on temperature and the number of extreme heat events could substantially influence respiratory morbidity across Europe.

INTRODUCTION

Respiratory diseases are ranked second in Europe in terms of mortality, prevalence and costs.¹ This burden is expected to increase, partly due to a changing climate.² An environmental factor with a large impact on mortality and morbidity in Europe is extreme heat, with large effects on respiratory diseases.³⁻⁵ Physiological effects of exposure to heat can be directly heat related (heat stroke, heat fatigue and dehydration) or can contribute to worsening of respiratory and cardiovascular diseases, electrolyte disorders, and kidney problems.⁴⁻⁷ The reasons for an increase in respiratory admissions may be several. Elderly with respiratory diseases such as COPD are less fit and suffer often from circulatory problems. Heat influenced admissions due to chronic airway obstruction and asthma increased more than admissions due to chronic bronchitis in a study from New York City,⁸ but the daily number of admissions seldom allow a study of specific diagnoses. In a recent expert elicitation amongst European researchers engaged in environmental medicine or respiratory health, extreme heat stood out as most important climate related pathway to adverse impacts on respiratory health, more important than changes in air pollutants and allergens.⁹ A review found that heatwaves have a stronger relative impact on mortality than on emergency room visits and hospital admissions, suggesting that many individuals die before they can get to the hospital.^{8,10} However, several studies confirm that heat also affects health care utilisation.^{3, 9,11} During the heatwave of 2003 in France, the Assistance Public Hôpitaux de Paris recorded 2,400 additional visits to the emergency care units and 1,900 additional hospital admissions.^{4,12} During the heatwave in California in 2006 there were almost 1,200 hospital admissions together with more than 16,000 visits to the emergency departments.⁶

It is possible that hospital admissions could increase in the future due to temperature changes projected with climate change, assuming no additional acclimatisation. Projected changes include an increase in global mean temperature of 1.8°C to 4.0°C by the end of this century, with larger increases over land areas at high latitudes, changing seasonal temperatures, and increases in the frequency, intensity, duration, and spatial extent of heatwaves.^{4,13} Recent projections for 2070–2100

1
2
3
4
5
6 suggest that maximum temperatures experienced once every 20 years during the period 1961–1990
7
8 could be expected as often as every year in southern Europe and every 3rd-5th year in northern
9
10 Europe.⁴²¹⁴ This means that Europe should expect an increase in both mean temperatures and the
11
12 number of extreme heat events. Recent heatwaves are consistent with projections. For example, the
13
14 heatwave that occurred in Europe 2003 could be expected to return every 46,000 years, based on
15
16 the temperature distribution for the years 1864–2000.⁴²¹⁵ However, such extreme hot conditions
17
18 may become ~~normal~~ much more common in the end of this century due to anthropogenic climate
19
20 changes.⁴⁴¹⁶ The uncertainty in this estimated return time is quite high, with a lower bound of the
21
22 90% confidence interval of 9,000 years. Even so, another heatwave occurred in 2006 with the most
23
24 anomalous July temperature ever measured in Europe.⁴⁵¹⁷ In 2010, Eastern Europe experienced a
25
26 heatwave with summer temperatures higher than the last 140 years, resulting in roughly 55,000
27
28 excess deaths in Russia.⁴⁶¹⁸

29
30 The possible impacts of these changes ~~in~~ on morbidity, particularly respiratory health, are relatively
31
32 unexplored.

33 34 35 **Aim**

36 The aim of this study is to assess the extent to which changes in the frequency of hot days due to
37
38 climate change over the next 40 years could affect heat related respiratory hospital admissions
39
40 (RHAs) in Europe. Using a range of climate projections, we estimate the change in hospital
41
42 admissions between a reference period and a future period.

43 44 45 **METHODS**

46 47 48 **Climate change and temperature modelling**

49 The Rossby Centre regional atmospheric climate model RCA3⁴⁷RCA3¹⁹ was developed by SMHI
50
51 (Swedish meteorological and hydrological institute) to dynamically downscale results from the global
52
53 climate models CCSM3¹⁸, ECHAM5⁴⁹, HadleyCM3²⁰ and ECHAM4²¹ CCSM3²⁰, ECHAM5²¹, HadleyCM3²²
54
55
56
57
58
59
60

1
2
3
4
5
6
7 | [and ECHAM4²³](#) to a higher resolution over Europe. RCA3 is run on a horizontal grid spacing of 0.44°
8
9 (corresponding to approximately 50 km) and a time step of 30 minutes. Projections are based on the
10 global greenhouse gas emission scenarios A1B and A2 (scenarios are described in detail in the
11
12 | [SRES²²SRES²⁴](#)). Both have been used in climate change health impact assessments (HIA).²³⁻²⁶²⁵⁻²⁸ A1B is
13
14 a “middle of the road” scenario and A2 is considered a high emission scenario, although recent
15
16 greenhouse gas emissions have been higher. Data from one climate model, under one climate
17
18 change scenario, is referred to as one climate change projection.

19
20 We use aggregated daily projections of maximum temperature and relative humidity data to
21
22 | estimate exposure for the periods 1981–~~2021~~2020 (reference period) and 2021–2050 (future
23
24 | period).

25 26 27 **Population and morbidity rates**

28
29 The annual country specific rate of respiratory hospital admissions between 2005 and 2010 were
30
31 extracted from the WHO’s European Health for All Database (<http://data.euro.who.int/hfadb>) for the
32
33 EU27 countries. The mean value over the six years was used as a baseline morbidity rate. Official
34
35 population data were from the HYDE theme within the Netherlands Environmental Assessment
36
37 | Agency.²⁷²⁹ These data are gridded on a 0.0833° resolution (approximately 9,45km) and matched
38
39 with the climatic data by summing the population within each climatic grid cell.

40 41 42 **Exposure-response assumption**

43
44 | The impact calculations were based on the ~~heat morbidity~~ relationship [between heat and RHAs \(ICD-](#)
45
46 [9:460–519\)](#) estimated within the European PHEWE [\(Assessment and Prevention of Acute Health](#)
47
48 [Effects of Weather Conditions in Europe\)](#) project.³ This relationship was based on analyses of 12
49
50 European cities over the 1990s, using the daily maximum of apparent temperature (AT), where AT is
51
52 a combination of the measured temperature and the dew point temperature:

$$53 \quad AT = -2.653 + 0.944 * T + 0.0153 * (DT)^2$$

54
55
56
57
58
59
60

1
2
3
4
5
6
7 where T is the air temperature and DP the dew point temperature, which can be derived from
8 temperature and relative humidity.^{28,30} The AT accounts for heat stress related not only to the
9 absolute temperature but also to the saturation of the surrounding air that makes it harder to
10 regulate body temperature by sweating, the most important mechanism to maintain a healthy body
11 temperature.²⁹⁻³⁰ ~~The PHEWE~~^{31 32} The PHEWE model also included potential confounders such as air
12 pollution, holidays, week days etc. The study used a 0–3 day lag of the maximum AT and concluded
13 that the 90th percentile of this exposure variable for the summer months, April to September, is the
14 appropriate threshold value for the heat-morbidity function.³ We calculated the 90th percentile for
15 the summer months for each grid cell from the climate data for the 1990s, yielding individual
16 thresholds for each grid cell for each climate projection.
17
18
19
20
21
22
23
24

25
26 The PHEWE study calculated the relative risk (RR) coefficients for each of the 12 cities and then
27 combined these into two meta-coefficients, for North-Continental cities 1.012 (95% CI 1.001–1.022)
28 and Mediterranean cities 1.021 (95% CI 1.006–1.036), associated with a 1-degree increase above the
29 temperature threshold. Meta-coefficients were more suitable as they allowed calculation of the
30 change in RHAs for countries with and without a city in the PHEWE Study. We assigned the
31 Mediterranean coefficient to Portugal and the Mediterranean countries. The rest of Europe was
32 assigned the RR for the North continental cities. Statistically significant coefficients were found for all
33 ages and age 75+ in the PHEWE study. Because the population data were not stratified by age, we
34 used the PHEWE study coefficients for all ages.
35
36
37
38
39
40
41
42
43

44 Projections

45 The final data used in the impact calculations were based on the climate data grid with a resolution
46 of 0.44° resulting in 8,075 grid cells over Europe. The population data, RR coefficients and baseline
47 morbidity rates were projected for each grid cell. ~~In addition,~~ The countries within EU27 were
48 grouped into four regions, Northern, Western, Eastern and Southern Europe, according to the United
49 Nations classification scheme (<http://unstats.un.org/unsd/methods/m49/m49regin.htm#europe>).
50
51
52
53
54
55
56
57
58
59
60

Thus, each grid cell includes (1) a daily time-series of 0–3 day lag maximum AT for each climate change projection; (2) a grid cell specific temperature morbidity threshold and a location-based RR coefficient; and (3) population. For each grid cell, the AT was compared to the specific threshold for each day in the two time periods and risk estimates were calculated for each day. The expected number of daily RHAs in each cell was calculated using the population and the expected daily number of RHAs per capita, based on the national average of the grid cell.

$$RHA_{i,t} = RR_i^{(AT_{i,t} - Thres_i)_+} * Pop_i * RHA_{pc_i}$$

where $RHA_{i,t}$ is the number of RHAs attributable to heat in grid cell i at time t , RR_i the RR coefficient in grid cell i , $(AT_{i,t} - Thres_i)_+$ the positive difference between the 0–3 lag AT and the threshold for grid cell i at time t if $AT_{i,t}$ were greater than the threshold and 0 otherwise, Pop_i and RHA_{pc_i} the population and expected number RHAs per capita in grid cell i .

For each country, the estimated number of RHAs attributed to heat was calculated for each climate change projection, for both the reference and the future period. The estimated number of RHAs attributed to heat was then transformed into the proportion of the expected annual number RHAs for each country (data available from the authors). Mean estimates were calculated for each region for each climate change projection.

Sensitivity analysis

Because the study results could heavily depend on the RR coefficients and thresholds used, we assess whether the estimates appeared sensitive to the region-specific meta-coefficients. We investigated the grids that contained the cities in the PHEWE study using both the location (Mediterranean or North-Continental) and city specific RR coefficients.³ We then compared the results with respect to the attributed proportion of RHAs and the proportional change in heat related RHAs.

RESULTS

The periods of warm days will increase in the future. For the cities included in the sensitivity analysis, the temperature threshold was exceeded by 20 days in the baseline period and 40 days in the future period, on average each year.

In the future period approximately 0.4% of the annual numbers of RHAs in Europe was estimated to be due to heat (Table 1), based on the mean estimates over the climate change projections. In absolute terms, assuming all else equal, this represents about 26,000 cases annually in Europe. This should be compared to the reference period where approximately 0.18% of all RHAs were attributed to heat or about 11,000 cases annually. Thus, the results suggest more than a relative doubling of the RHAs attributed to heat in Europe.

Table 1. The estimated proportion of RHAs attributed to heat for each region. Intervals describe the highest and lowest estimate in each region

Region	1981–2010	2021–2050	Change
Eastern Europe	0.17% (0.16% – 0.19%)	0.31% (0.29% – 0.35%)	0.14% (0.11% – 0.17%)
Northern Europe	0.13% (0.10% – 0.15%)	0.27% (0.19% – 0.32%)	0.14% (0.07% – 0.17%)
Southern Europe	0.23% (0.18% – 0.26%)	0.64% (0.42% – 0.68%)	0.41% (0.23% – 0.44%)
Western Europe	0.18% (0.16% – 0.20%)	0.39% (0.34% – 0.45%)	0.21% (0.17% – 0.26%)
EU27	0.18% (0.10% – 0.26%)	0.40% (0.19% – 0.68%)	0.21% (0.07% – 0.44%)

On the regional level, the five projections estimate increases in the number of heat related RHAs for Europe (Table 2). However, in one climate change projection, The Czech Republic, Hungary, Poland and Slovakia were estimated to have a decrease in heat related RHAs (Fig 1). The countries with the highest estimated increase also show the largest range between the highest and lowest estimate. The Scandinavian and Baltic countries show the smallest range between the highest and lowest estimates along with small increases of the mean estimates.

Table 2. Future increase in heat related RHAs based on the four climate models, under two emission scenarios, as the proportionpercentage of the annual expected number of RHAs in each region

Climate model	CCSM3	ECHAM5	HadCM3	ECHAM4	ECHAM5
<i>Greenhouse gas scenario</i>	<i>A1B</i>		<i>A2</i>		
Eastern Europe	0.32%	0.08%	0.18%	0.12%	0.01%
Northern Europe	0.17%	0.09%	0.14%	0.20%	0.08%
Southern Europe	0.51%	0.29%	0.64%	0.45%	0.14%
Western Europe	0.30%	0.11%	0.26%	0.29%	0.06%
EU27	0.32%	0.13%	0.29%	0.26%	0.07%

There is variation among countries, with the largest increases in the Southern European countries and the smallest increase in Eastern Europe (Table 2). The relative change in the burden of RHA's in relation to the climate change scenarios investigated indicate a larger relative increase in Mediterranean countries (approximately three times) compared to Northern European countries (approximately two times).

Sensitivity analysis

The calculations using the city specific RR coefficients yielded a different proportion and number of estimated RHAs than the ones using the two meta-coefficients, as expected. However, the relative changes in the number of hospital admissions attributed to heat between the two time-periods were not affected by the change of the RR coefficients.

DISCUSSION

The analysis estimates that the number of RHAs will increase in a warmer climate with more hot days. Heat related RHAs were projected to increase two- to three-fold in the future due to climate change. However, the proportion of respiratory admissions attributed to heat would remain rather small. This projection is in line with the results of a recent study which estimates that the respiratory

admissions due to excessive heat in New York State will increase 2-6 times from the period 1991-2004 to the period 2080-2099.³³

As most heat related health outcomes occur during the warmest period of the year, presenting the increase as a change in the proportion of the total number of annual RHAs can underestimate the additional burden on the health care system during summer. As the threshold was exceeded for approximately 40 days in one year, an annual increase of 0.21% would result in a 1.9% increase on average during these 40 days. The annual numbers are used because the available baseline rates of RHAs are expressed as an annual average. Applying results to sub-national scales could be inappropriate because national averages summarize over considerable heterogeneity.

The results suggest a larger impact from heat in Southern Europe in the future period, centred on year 2035, than in the eastern and northern parts. This is in line with many climate change projections showing a larger relative increase in the number of extremely hot days in southern Europe compared with northern Europe.¹²¹⁴ However, to some extent this might also be explained by model bias for northern Europe introduced by the RCA3 model, where the model appears to underestimate temperature for the warmest days.¹⁷¹⁹ This temperature bias is present in the reference period, future period, and threshold values; therefore, the estimated numbers of RHAs from the two periods within each scenario are comparable, but comparisons within the same time-period, across scenarios may be inaccurate.

As the estimates are based on the population size in each grid cell, the added burden will be larger in countries with larger increases in temperatures in densely populated areas. Because the population of a city is considered a good predictor of the size of the urban heat island,³⁴³⁴ an increase in the population living in urban areas will increase the numbers exposed and the temperature to which they are exposed. Heat islands increase temperatures in urbanized areas compared to surrounding areas, also reducing cooling during the night time.²²³⁵ These factors combined are likely to magnify the health burden during a heatwave.²²³⁶ The coarse spatial scale of the climate models makes them

1
2
3
4
5
6
7 unable to take the urban heat island effect into account. Together with the urbanization of Europe,
8 this could potentially result in underestimation of actual consequences/RHAs in the future period
9 because the same population size and composition were assumed in both periods. In addition,
10 climate change is likely to increase ozone concentrations that would add an additional health burden
11
12
13
14 ~~to~~for people at risk of respiratory diseases. The number of deaths and RHAs, due to a change in
15 ozone, is expected to change in the future. ^{34,37} A sensitivity analysis in the PHEWE study however
16 showed that the exposure-response relationship for heat did not substantially differ between models
17 taking ozone into account and models adjusting for NO₂ alone.³

22 We used the same thresholds for the reference and future period to isolate the effect of climate
23 change. This would tend to overestimate the increased impact because there will undoubtedly be
24 biological and/or social adaptations that will reduce future health temperature-related burdens.
25
26 However, given the uncertainty of such adaptation effects, we choose to not incorporate such effects
27
28 in the projections. ~~Studies from~~Recent studies in the U.S ~~show that they~~indicate heat related health
29
30 burdens ~~have decreased over the last 50 years,~~^{35, 38} indicating that some adaptation is taking place.
31
32 An opposite trend appears to be occurring in Stockholm, with an increase of the risk of mortality
33 associated with heat during the 1990s. ^{36,39} A study looking at the impact of heat on mortality, before
34 and after the implementation of a heat warning system, in Italy shows that the effects of extreme
35 heat can be reduced while the effects of moderate increased average temperatures remains
36 similar.⁴⁰ The magnitude and the extent of future adaptation is, of course, highly uncertain, and will
37 vary between and within countries. Nevertheless, cities with higher thresholds seem to have higher
38 risk ratio coefficients.^{3, 37,41} In an effort to estimate future and presumably higher thresholds, one
39 must also adjust the risk ratio coefficients. This would result in fewer days of elevated risk, but the
40 risk increase on each occasion could be higher due to the higher RRs.
41
42
43
44
45
46
47
48
49
50

51 Table 1 shows how the results vary by global climate models and greenhouse gas scenarios. This
52 study exemplifies the magnitude of the difference between projections made by a model with
53
54
55
56
57
58
59
60

1
2
3
4
5
6 different initial conditions and between different models with the same initial conditions. The results
7
8 indicate the range of these estimates is large. The mean increase over the five projections however,
9
10 provides confidence that the number of RHAs will increase over Europe as the climate continues to
11
12 change.

13
14
15 To improve the estimates of the impacts of heat ~~of~~ RHAs, detailed data on emergency room visits
16
17 and admissions during summer are needed, such as the proportion of emergency department cases
18
19 admitted to hospital, and the fatality rate. A better understanding of vulnerable groups is needed,
20
21 including how these groups could change over time. For example, the portion of the population aged
22
23 over 65 will increase from 17 to 29%^{38,42} by 2050, significantly increasing the size of this vulnerable
24
25 group. COPD is mainly a disease of the elderly. Persons suffering from COPD appear especially
26
27 vulnerable to heat⁴³ and the incidence of COPD is likely to increase⁴⁴ with the consequence that the
28
29 health impacts of heat may increase in future years. This study was limited to estimating the impacts
30
31 on the total population because few age-specific relationship were reported and because age
32
33 stratified data were only available on a coarser spatial scale than the climate data, which would
34
35 reduce the benefits of having spatial climate data.

36
37 Normal weather patterns in the future will be different than those of today, both for average
38
39 temperature and extreme weather events. As the number of heat related RHAs are expected to
40
41 increase on the sub-continental scale, additional national or regional projections of the future health
42
43 burden from heat are needed, taking into account possible changes in exposure and vulnerability.
44
45

46 CONCLUSIONS

47
48 Projected changes in temperature and the number of extreme heat events with climate change could
49
50 substantially influence respiratory morbidity across Europe. Analyses projected that both the future
51
52 proportion of annual RHAs attributed to heat and the relative change in heat related RHAs will be
53
54 largest in Southern Europe, where they are expected to nearly triple. Eastern Europe can expect the
55
56
57
58
59
60

1
2
3
4
5
6
7 smallest increase in heat related RHAs, where they are estimated to approximately double. For all of
8 Europe, the number of respiratory heat-related hospital admissions is projected to be 26,000
9 annually in the future period (2021–2050) compared with 11,000 in the reference period (1981–
10 2010). The estimates presented rely on the assumption that no additional adaptation occurs. Future
11 studies should elaborate and quantify the possible effects of different adaptation assumptions
12 applied to regional conditions.
13
14
15
16
17
18
19

20
21
22
23
24 This work was supported by the EU-funded Climate-Trap project (contract EAHC 20081108) and by
25 the Swedish Environmental Protection Agency through the research programme CLEO – Climate
26 Change and Environmental Objectives. We would also like to acknowledge Estonia's Ministry of
27 Education for providing resources to H. Orru with the grant SF0180060s09.
28
29
30

31
32 Contribution statement: CÅ, BF, HO has planned and designed the study. GS represented SMHI and
33 provided climate data. CÅ ran the analyses, drafted the first manuscript and is guarantor. All authors
34 have participated in the interpretation of results and revised the manuscript.
35
36
37

38
39 The specific estimates for each of the EU27 countries, for each of the climate projections is available
40 after contact with corresponding author.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Loddenkemper R, Gibson G, Sibille Y. European lung white book. *European Respiratory Society/European Lung Foundation, Sheffield, UK, ERSJ* 2003:1-182.
2. Ayres JG, Forsberg B, Annesi-Maesano I, et al. Climate change and respiratory disease: European Respiratory Society position statement. *Eur Respir J* 2009;34(2):295-302.
3. Michelozzi P, Accetta G, De Sario M, et al. High temperature and hospitalizations for cardiovascular and respiratory causes in 12 European cities. *Am J Respir Crit Care Med* 2009;179(5):383-9.
4. [Rocklöv J, Forsberg B. Comparing approaches for studying the effects of climate extremes - a case study of hospital admissions in Sweden during an extremely warm summer. *Glob Health Action* 2009;2.](#)
5. Kovats RS, Hajat S. Heat stress and public health: a critical review. *Annu Rev Public Health* 2008;29:41-55.
6. Knowlton K, Rotkin-Ellman M, King G, et al. The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect* 2009;117(1):61-7.
7. Rey G, Jouglà E, Fouillet A, et al. The impact of major heat waves on all-cause and cause-specific mortality in France from 1971 to 2003. *Int Arch Occup Environ Health* 2007;80(7):615-26.
8. [Lin S, Luo M, Walker RJ, et al. Extreme high temperatures and hospital admissions for respiratory and cardiovascular diseases. *Epidemiology* 2009;20\(5\):738-46.](#)
9. [Forsberg B, Braback L, Keune H, et al. An expert assessment on climate change and health - with a European focus on lungs and allergies. *Environ Health* 2012;11 Suppl 1:S4.](#)
10. Ye X, Wolff R, Yu W, et al. Ambient Temperature and Morbidity: A Review of Epidemiological Evidence. *Environ Health Perspect* 2011.
11. Semenza JC, McCullough JE, Flanders WD, et al. Excess hospital admissions during the July 1995 heat wave in Chicago. *Am J Prev Med* 1999;16(4):269-77.
12. Dhainaut JF, Claessens YE, Ginsburg C, et al. Unprecedented heat-related deaths during the 2003 heat wave in Paris: consequences on emergency departments. *Crit Care* 2004;8(1):1-2.
13. [IPCC. *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation - Summary for Policymakers*. Geneva, Switzerland: World Meteorological Organization, 2012.](#)
14. [IPCC. *Climate Change 2007 - The Physical Science Basis: Working Group I Contribution to the Fourth Assessment Report of the IPCC*. Cambridge, UK and New York, NY, USA: Cambridge University Press, 2007.](#)
15. Nikulin G, Kjellström E, Hansson ULF, et al. Evaluation and future projections of temperature, precipitation and wind extremes over Europe in an ensemble of regional climate simulations. *Tellus A* 2011;63(1):41-55.
16. Schar C, Vidale PL, Luthi D, et al. The role of increasing temperature variability in European summer heatwaves. *Nature* 2004;427(6972):332-6.
17. Stott PA, Stone DA, Allen MR. Human contribution to the European heatwave of 2003. *Nature* 2004;432(7017):610-4.
18. Rebetez M, Dupont O, Giroud M. An analysis of the July 2006 heatwave extent in Europe compared to the record year of 2003. *Theoretical and Applied Climatology* 2009;95(1):1-7.
19. Barriopedro D, Fischer EM, Luterbacher J, et al. The Hot Summer of 2010: Redrawing the Temperature Record Map of Europe. *Science* 2011.
20. Samuelsson P, Jones CG, Willen U, et al. The Rossby Centre Regional Climate model RCA3: model description and performance. *Tellus Series a-Dynamic Meteorology and Oceanography* 2011;63(1):4-23.
21. Collins WD, Bitz CM, Blackmon ML, et al. The Community Climate System Model version 3 (CCSM3). *J Climate* 2006;19(11):2122-43.
22. [Roeckner E, Bäuml G, Bonaventura L, et al. The atmospheric general circulation model ECHAM 5. PART I: Model description. Hamburg, Germany: MPI für Meteorologie, 2003.](#)

Formatted: Swedish (Sweden)

Formatted: Swedish (Sweden)

- 1
2
3
4
5
6
7 | [2022](#). Johns TC, Durman CF, Banks HT, et al. The new Hadley Centre Climate Model (HadGEM1): Evaluation of coupled simulations. *J Climate* 2006;19(7):1327-53.
- 8 | [2123](#). Roeckner E, Arpe K, Bengtsson, et al. The atmospheric general circulation model {ECHAM-}4: ~~Model~~^{Model} description and simulation of present-day climate, 1996:103-23.
- 9
10
11 | [2224](#). Nakicenovic N, Alcamo J, Davis G, et al. Special report on emissions scenarios: a special report of Working Group III of the Intergovernmental Panel on Climate Change: Pacific Northwest National Laboratory, Richland, WA (US), Environmental Molecular Sciences Laboratory (US), 2000.
- 12
13
14 | [2325](#). Ballester J, Robine JM, Herrmann FR, et al. Long-term projections and acclimatization scenarios of temperature-related mortality in Europe. *Nat Commun* 2011;2:358.
- 15
16 | [2426](#). Knowlton K, Lynn B, Goldberg RA, et al. Projecting heat-related mortality impacts under a changing climate in the New York City region. *American Journal of Public Health* 2007;97(11):2028-34.
- 17
18
19 | [2527](#). Baccini M, Kosatsky T, Analitis A, et al. Impact of heat on mortality in 15 European cities: attributable deaths under different weather scenarios. *J Epidemiol Community Health* 2011;65(1):64-70.
- 20
21
22 | [2628](#). Beguin A, Hales S, Rocklov J, et al. The opposing effects of climate change and socio-economic development on the global distribution of malaria. *Global Environmental Change-Human and Policy Dimensions* 2011;21(4):1209-14.
- 23
24
25 | [2729](#). Goldewijk KK, Beusen A, Janssen P. Long-term dynamic modeling of global population and built-up area in a spatially explicit way: HYDE 3.1. *Holocene* 2010;20(4):565-73.
- 26
27 | [2830](#). Lawrence MG. The Relationship between Relative Humidity and the Dewpoint Temperature in Moist Air: A Simple Conversion and Applications. *Bulletin of the American Meteorological Society* 2005;86(2):225-33.
- 28
29 | [2931](#). Parsons KC. *Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort, and performance*: Taylor & Francis, 2003.
- 30
31 | [3032](#). Steadman RG. A Universal Scale of Apparent Temperature. *J Clim Appl Meteorol* 1984;23(12):1674-87.
- 32
33 | [3133](#). Lin S, Hsu WH, Van Zutphen AR, et al. Excessive heat and respiratory hospitalizations in new york state: estimating current and future public health burden related to climate change. *Environ Health Perspect* 2012;120(11):1571-7.
- 34
35
36 | [34](#). Oke TR. City size and the urban heat island. *Atmospheric Environment (1967)* 1973;7(8):769-79.
- 37
38 | [3235](#). Holmer B, Thorsson S, Eliasson I. COOLING RATES, SKY VIEW FACTORS AND THE DEVELOPMENT OF INTRA - URBAN AIR TEMPERATURE DIFFERENCES. *Geografiska Annaler: Series A, Physical Geography* 2007;89(4):237-48.
- 39
40 | [3336](#). Nicholls N, Skinner C, Loughnan M, et al. A simple heat alert system for Melbourne, Australia. *Int J Biometeorol* 2008;52(5):375-84.
- 41
42 | [3437](#). Orru H, Andersson C, Ebi KL, et al. Impact of climate change on ozone related mortality and morbidity in Europe ~~(Accepted)~~. *Eur Respir J* 2012. Jun 27. [Epub ahead of print].
- 43
44 | [3538](#). Davis RE, Knappenberger PC, Michaels PJ, et al. Changing heat-related mortality in the United States. *Environmental Health Perspectives* 2003;111(14):1712-18.
- 45
46 | [3639](#). Rocklov J, Forsberg B, Meister K. Winter mortality modifies the heat-mortality association the following summer. *Eur Respir J* 2009;33(2):245-51.
- 47
48 | [3740](#). Schifano P, Leone M, De Sario M, et al. Changes in the effects of heat on mortality among the elderly from 1998-2010: results from a multicenter time series study in Italy. *Environmental Health* 2012;11(1):58.
- 49
50 | [41](#). Baccini M, Biggeri A, Accetta G, et al. Heat effects on mortality in 15 European cities. *Epidemiology* 2008;19(5):711-9.
- 51
52 | [3842](#). Giannakouris K. Ageing characterises the demographic perspectives of the European societies. *Statistics in focus* 2008;72:2008.
- 53
54 | [43](#). Schwartz J. Who is sensitive to extremes of temperature? A case-only analysis. *Epidemiology* 2005;16(1):67-72.
- 55
56
57
58
59
60

Formatted: Swedish (Sweden)

Formatted: Swedish (Sweden)


Formatted: Swedish (Sweden)

1
2
3
4
5
6
7 44. Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet*
8 2007;370(9589):765-73.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7 **Figure 1. The range of the absolute increase in RHAs attributed to heat between the two**
8 **periods (1981-2010, 2021-2050) as proportion the annual expected number of RHAs for each of the**
9 **27 countries. The points show the highest and lowest estimate from four climate models under**
10 **two emission scenarios.**
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only



406x279mm (200 x 200 DPI)

view only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Heat related respiratory hospital admissions in Europe in a changing climate: A health impact assessment

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2012-001842.R2
Article Type:	Research
Date Submitted by the Author:	20-Dec-2012
Complete List of Authors:	Åström, Christofer; Umeå University, Department of Public Health and Clinical Medicine Orru, Hans Rocklöv, Joacim; Umeå University, Department of Public Health and Clinical Medicine Strandberg, Gustav; Swedish Meteorological and Hydrological Institute, Ebi, Kristie; Stanford University, ClimAdapt Forsberg, Bertil; Umeå University, Department of Public Health and Clinical Medicine
Primary Subject Heading:	Epidemiology
Secondary Subject Heading:	Epidemiology, Occupational and environmental medicine
Keywords:	EPIDEMIOLOGY, PUBLIC HEALTH, Epidemiology < THORACIC MEDICINE

SCHOLARONE™
Manuscripts

Heat related respiratory hospital admissions in Europe in a changing climate

Authors:

Christofer Åström^{1*}, Hans Orru^{1,2}, Joacim Rocklöv^{1,3}, Gustav Strandberg⁴, Kristie L. Ebi⁵, Bertil Forsberg¹

Affiliations:

¹ Department of Public Health and Clinical Medicine, Occupational and Environmental Medicine, Umeå University, Umeå, Sweden; ² Department of Public Health, University of Tartu, Tartu, Estonia; ³ Department of Public Health and Clinical Medicine, Epidemiology and Global Health, Umeå University, Umeå, Sweden; ⁴ Swedish Meteorological and Hydrological Institute, Norrköping, Sweden; ⁵ ClimAdapt, Stanford University, Los Altos, California, USA

Corresponding author:

* Author to whom correspondence should be addressed:

Christofer Åström,

Occupational and Environmental Medicine, Department of Public Health and Clinical Medicine, Umeå University, SE 901 87, Sweden

+4690 785 0874

christofer.astrom@envmed.umu.se

Keywords: Climate change, respiratory health, public health, heat

Word count: 3246

ARTICLE SUMMARY

Article focus

To assess how heat-related respiratory hospital admissions in Europe could change with increasing temperatures in the near future. Within a range of climate change projections, explore different impact estimates.

Key messages

The 30-year mean annual increase in heat-related respiratory hospital admissions between the periods 1981-2010 and 2021-2050 can be counted in the 10's of thousands because of projected temperature increases. The increase will be different in different parts of Europe with the largest relative increase in Southern Europe.

Strengths and limitations

The study takes the spatial variation in climate and population density into account in impact calculations, making the estimates valid for countries with large national differences in these factors.

The study results exemplify some the variation of impacts under different climate change scenarios, as well as the variations within one scenario depending on the underlying climate model.

The use of spatial population data limited the study to exploring the change in all age respiratory hospital for studying heat-related illness, combined with the future increase in the proportion of elderly people, suggests the estimated impacts are likely to be conservative

ABSTRACT

Objectives: Respiratory diseases are ranked second in Europe in terms of mortality, prevalence and costs. Studies have shown that extreme heat has a large impact on mortality and morbidity, with a large relative increase for respiratory diseases. Expected increases in mean temperature and the number of extreme heat events over the coming decades due to climate change raise questions about the possible health impacts. Here we aim to assess the number of heat related respiratory hospital admissions in a future with a different climate.

Design: A European wide health impact assessment.

Setting: An assessment for each of the EU27 countries.

Methods: Heat-related hospital admissions under a changing climate are projected using multi-city epidemiologic exposure-response relationships applied to gridded population data and country-specific baseline respiratory hospital admission rates. Times-series of temperatures are simulated with a regional climate model based on four global climate models, under two greenhouse gas emission scenarios.

Results: Between a reference period (1981–2010) and a future period (2021–2050), the total number of respiratory hospital admissions attributed to heat is projected to be larger in Southern Europe, with three-times more heat attributed respiratory hospital admissions in the future period. The smallest change was estimated in Eastern Europe with about a two-fold increase. For all of Europe, the number of heat-related respiratory hospital admissions is projected to be 26,000 annually in the future period compared with 11,000 in the reference period.

Conclusions: The results suggest that projected effects of climate change on temperature and the number of extreme heat events could substantially influence respiratory morbidity across Europe.

INTRODUCTION

Respiratory diseases are ranked second in Europe in terms of mortality, prevalence, and costs.¹ This burden is expected to increase, partly due to a changing climate.² An environmental factor with a large impact on mortality and morbidity in Europe is extreme heat, with large effects on respiratory diseases.³⁻⁵ Physiological effects of exposure to heat can be directly heat-related (heat stroke, heat fatigue and dehydration) or can contribute to worsening of respiratory and cardiovascular diseases, electrolyte disorders, and kidney problems.⁴⁻⁷ The reasons for an increase in respiratory admissions may be several. Elderly with respiratory diseases such as COPD are less fit and suffer often from circulatory problems. Heat influenced admissions due to chronic airway obstruction and asthma increased more than admissions due to chronic bronchitis in a study from New York City,⁸ but the daily number of admissions seldom allow a study of specific diagnoses. In a recent expert elicitation amongst European researchers engaged in environmental medicine or respiratory health, extreme heat stood out as most important climate-related pathway to adverse impacts on respiratory health, more important than changes in air pollutants and allergens.⁹ A review found that heatwaves have a stronger relative impact on mortality than on emergency room visits and hospital admissions, suggesting that many individuals die before they can get to the hospital.¹⁰ However, several studies confirm that heat also affects health care utilisation.^{3,11} During the heatwave of 2003 in France, the Assistance Public Hôpitaux de Paris recorded 2,400 additional visits to the emergency care units and 1,900 additional hospital admissions.¹² During the heatwave in California in 2006 there were almost 1,200 hospital admissions together with more than 16,000 visits to the emergency departments.⁶

It is possible that hospital admissions could increase in the future due to temperature changes projected with climate change, assuming no additional acclimatisation. Projected changes include an increase in global mean temperature of 1.8°C to 4.0°C by the end of this century, with larger increases over land areas at high latitudes, changing seasonal temperatures, and increases in the frequency, intensity, duration, and spatial extent of heatwaves.¹³ Recent projections for 2070–2100

1
2
3 suggest that maximum temperatures experienced once every 20 years during the period 1961–1990
4
5 could be expected as often as every year in southern Europe and every 3rd-5th year in northern
6
7 Europe.¹⁴ This means that Europe should expect an increase in both mean temperatures and the
8
9 number of extreme heat events. Recent heatwaves are consistent with projections. For example, the
10
11 heatwave that occurred in Europe 2003 could be expected to return every 46,000 years, based on
12
13 the temperature distribution for the years 1864–2000.¹⁵ However, such extreme hot conditions may
14
15 become much more common in the end of this century due to anthropogenic climate changes.¹⁶ The
16
17 uncertainty in this estimated return time is quite high, with a lower bound of the 90% confidence
18
19 interval of 9,000 years. Even so, another heatwave occurred in 2006 with the most anomalous July
20
21 temperature ever measured in Europe.¹⁷ In 2010, Eastern Europe experienced a heatwave with
22
23 summer temperatures higher than the last 140 years, resulting in roughly 55,000 excess deaths in
24
25 Russia.¹⁸

26
27
28
29
30 The possible impacts of these changes on morbidity, particularly respiratory health, are relatively
31
32 unexplored.

33 34 35 36 **Aim**

37 The aim of this study is to assess the extent to which changes in the frequency of hot days due to
38
39 climate change over the next 40 years could affect heat related respiratory hospital admissions
40
41 (RHAs) in Europe. Using a range of climate projections, we estimate the change in hospital
42
43 admissions between a reference period and a future period.
44
45
46
47

48 49 **METHODS**

50 51 52 **Climate change and temperature modelling**

53 The Rossby Centre regional atmospheric climate model RCA3¹⁹ was developed by SMHI (Swedish
54
55 meteorological and hydrological institute) to dynamically downscale results from the global climate
56
57 models CCSM3²⁰, ECHAM5²¹, HadleyCM3²² and ECHAM4²³ to a higher resolution over Europe. RCA3 is
58
59
60

1
2
3 run on a horizontal grid spacing of 0.44° (corresponding to approximately 50 km) and a time step of
4
5 30 minutes. Projections are based on the global greenhouse gas emission scenarios A1B and A2
6
7 (scenarios are described in detail in the SRES²⁴). Both have been used in climate change health
8
9 impact assessments (HIA).²⁵⁻²⁸ A1B is a “middle of the road” scenario and A2 is considered a high
10
11 emission scenario, although recent greenhouse gas emissions have been higher. Data from one
12
13 climate model, under one climate change scenario, is referred to as one climate change projection.
14
15

16
17 We use aggregated daily projections of maximum temperature and relative humidity data to
18
19 estimate exposure for the periods 1981–2020 (reference period) and 2021–2050 (future period).
20
21

22 **Population and morbidity rates**

23
24 The annual country specific rate of RHAs (ICD-9:460–519) between 2005 and 2010 were extracted
25
26 from the WHO’s European Health for All Database (<http://data.euro.who.int/hfadb>) for the EU27
27
28 countries. In the dataset, national hospital admissions data are provided by national public health
29
30 institutes, health ministries or corresponding functions. The ICD-9 codes were preferred because the
31
32 epidemiological studies in the PHEWE study were based on ICD-9 RHAs. The mean value over the six
33
34 years was used as a baseline morbidity rate. In order to have a fine spatial resolution, official
35
36 population data were from the HYDE theme within the Netherlands Environmental Assessment
37
38 Agency.²⁹ These data are gridded on a 0.0833° resolution (approximately 9,45km) and matched with
39
40 the climatic data by summing the population within each climatic grid cell.
41
42
43
44

45 **Exposure-response assumption**

46
47 The impact calculations were based on the relationship between heat and RHAs (ICD-9:460–519)
48
49 estimated within the European PHEWE (Assessment and Prevention of Acute Health Effects of
50
51 Weather Conditions in Europe) project.³ This relationship was based on analyses of 12 European
52
53 cities over the 1990s, using the daily maximum of apparent temperature (AT), where AT is a
54
55 combination of the measured temperature and the dew point temperature:
56
57
58
59
60

$$AT = -2.653 + 0.944 * T + 0.0153 * (DT)^2$$

where T is the air temperature and DP the dew point temperature, which can be derived from temperature and relative humidity.³⁰ The AT accounts for heat stress related not only to the absolute temperature but also to the saturation of the surrounding air that makes it harder to regulate body temperature by sweating, the most important mechanism to maintain a healthy body temperature.³¹

³² The PHEWE model also included potential confounders such as air pollution, holidays, week days etc. The study used a 0–3 day lag of the maximum AT and concluded that the 90th percentile of this exposure variable for the summer months, April to September, is the appropriate threshold value for the heat-morbidity function.³ We calculated the 90th percentile for the summer months for each grid cell from the climate data for the 1990s, yielding individual thresholds for each grid cell for each climate projection.

The PHEWE study calculated the relative risk (RR) coefficients for each of the 12 cities and then combined these into two meta-coefficients, for North-Continental cities 1.012 (95% CI 1.001–1.022) and Mediterranean cities 1.021 (95% CI 1.006–1.036), associated with a 1-degree increase above the temperature threshold. Meta-coefficients were more suitable as they allowed calculation of the change in RHAs for countries with and without a city in the PHEWE Study. We assigned the Mediterranean coefficient to Portugal and the Mediterranean countries. The rest of Europe was assigned the RR for the North continental cities. Statistically significant coefficients were found for all ages and ages 75+ in the PHEWE study. Because the population data were not stratified by age, we used the PHEWE study coefficients for all ages.

Projections

The final data used in the impact calculations were based on the climate data grid with a resolution of 0.44° resulting in 8,075 grid cells over Europe. The population data, RR coefficients and baseline morbidity rates were projected for each grid cell. The countries within EU27 were grouped into four

regions, Northern, Western, Eastern and Southern Europe, according to the United Nations classification scheme (<http://unstats.un.org/unsd/methods/m49/m49regin.htm#europe>).

Thus, each grid cell includes (1) a daily time-series of 0–3 day lag maximum AT for each climate change projection; (2) a grid cell specific temperature-morbidity threshold and a location-based RR coefficient; and (3) population. For each grid cell, the AT was compared to the specific threshold for each day in the two time periods and risk estimates were calculated for each day. The expected number of daily RHAs in each cell was calculated using the population and the expected daily number of RHAs per capita, based on the national average of the grid cell.

$$RHA_{i,t} = RR_i^{(AT_{i,t} - Thres_i)_+} * Pop_i * RHApc_i$$

where $RHA_{i,t}$ is the number of RHAs attributable to heat in grid cell i at time t , RR_i the RR coefficient in grid cell i , $(AT_{i,t} - Thres_i)_+$ the difference between the 0–3 lag AT and the threshold for grid cell i at time t if $AT_{i,t}$ were greater than the threshold and 0 otherwise, Pop_i and $RHApc_i$ the population and expected number RHAs per capita in grid cell i .

For each country, the estimated number of RHAs attributed to heat was calculated for each climate change projection, for both the reference and the future period. The estimated number of RHAs attributed to heat was then transformed into the proportion of the expected annual number RHAs for each country (data available from the authors). Mean estimates were calculated for each region for each climate change projection.

Sensitivity analysis

Because the study results could heavily depend on the RR coefficients and thresholds used, we assess whether the estimates appeared sensitive to the region-specific meta-coefficients. We investigated the grids that contained the cities in the PHEWE study using both the location (Mediterranean or North-Continental) and city specific RR coefficients.³ We then compared the results with respect to the attributed proportion of RHAs and the proportional change in heat related RHAs.

RESULTS

The periods of warm days will increase in the future. For the cities included in the sensitivity analysis, the temperature threshold was exceeded annually, on average, by 20 days in the baseline period and 40 days in the future period.

In the future period, approximately 0.4% of the annual numbers of RHAs in Europe was estimated to be due to heat (Table 1), based on the mean estimates over the climate change projections. In absolute terms, assuming all else equal, this represents about 26,000 cases annually in Europe. This should be compared to the reference period where approximately 0.18% of all RHAs were attributed to heat or about 11,000 cases annually. Thus, the results suggest more than a relative doubling of the RHAs attributed to heat in Europe.

Table 1. The estimated proportion of RHAs attributed to heat for each region. Intervals describe the highest and lowest national estimate in each region

Region	1981–2010	2021–2050	Change
Eastern Europe	0.17% (0.16% – 0.19%)	0.31% (0.29% – 0.35%)	0.14% (0.11% – 0.17%)
Northern Europe	0.13% (0.10% – 0.15%)	0.27% (0.19% – 0.32%)	0.14% (0.07% – 0.17%)
Southern Europe	0.23% (0.18% – 0.26%)	0.64% (0.42% – 0.68%)	0.41% (0.23% – 0.44%)
Western Europe	0.18% (0.16% – 0.20%)	0.39% (0.34% – 0.45%)	0.21% (0.17% – 0.26%)
EU27	0.18% (0.10% – 0.26%)	0.40% (0.19% – 0.68%)	0.21% (0.07% – 0.44%)

On the regional level, the five projections estimate increases in the number of heat-related RHAs for Europe (Table 2). However, in one climate change projection, The Czech Republic, Hungary, Poland and Slovakia were estimated to have a decrease in heat-related RHAs (Fig 1). The countries with the highest estimated increase also show the largest range between the highest and lowest estimate. The Scandinavian and Baltic countries show the smallest range between the highest and lowest estimates along with small increases of the mean estimates.

Table 2. Future increase in heat-related RHAs based on the four climate models, under two emission scenarios, as the percentage of the annual expected number of RHAs in each region

Climate model	CCSM3	ECHAM5	HadCM3	ECHAM4	ECHAM5
<i>Greenhouse gas scenario</i>	<i>A1B</i>			<i>A2</i>	
Eastern Europe	0.32%	0.08%	0.18%	0.12%	0.01%
Northern Europe	0.17%	0.09%	0.14%	0.20%	0.08%
Southern Europe	0.51%	0.29%	0.64%	0.45%	0.14%
Western Europe	0.30%	0.11%	0.26%	0.29%	0.06%
EU27	0.32%	0.13%	0.29%	0.26%	0.07%

There is variation among countries, with the largest increases in the Southern European countries and the smallest increase in Eastern Europe (Table 2). The relative change in the burden of RHA's in relation to the climate change scenarios investigated indicate a larger relative increase in Mediterranean countries (approximately three times) compared to Northern European countries (approximately two times).

Sensitivity analysis

The calculations using the city specific RR coefficients yielded a different proportion and number of estimated RHAs than the ones using the two meta-coefficients, as expected. However, the relative changes in the number of hospital admissions attributed to heat between the two time-periods were not affected by the change of the RR coefficients.

DISCUSSION

The analysis estimates that the number of RHAs will increase in a warmer climate with more hot days. Heat-related RHAs were projected to increase two- to three-fold in the future due to climate change. However, the proportion of respiratory admissions attributed to heat would remain rather small. This projection is in line with the results of a recent study that estimates respiratory

1
2
3 admissions due to excessive heat in New York State will increase 2-6 times from the period 1991-
4
5 2004 to the period 2080-2099.³³
6
7

8 As most heat-related health outcomes occur during the warmest period of the year, presenting the
9
10 increase as a change in the proportion of the total number of annual RHAs can underestimate the
11
12 additional burden on the health care system during summer. As the threshold was exceeded for
13
14 approximately 40 days in one year, an annual increase of 0.21% would result in a 1.9% increase on
15
16 average during these 40 days. The annual numbers are used because the available baseline rates of
17
18 RHAs are expressed as an annual average. Applying results to sub-national scales could be
19
20 inappropriate because national averages summarize over considerable heterogeneity.
21
22

23
24 The results suggest a larger impact from heat in Southern Europe in the future period, centred on
25
26 year 2035, than in the eastern and northern parts. This is in line with many climate change
27
28 projections showing a larger relative increase in the number of extremely hot days in southern
29
30 Europe compared with northern Europe.¹⁴ However, to some extent this might also be explained by
31
32 model bias for northern Europe introduced by the RCA3 model, where the model appears to
33
34 underestimate temperature for the warmest days.¹⁹ This temperature bias is present in the reference
35
36 period, future period, and threshold values; therefore, the estimated numbers of RHAs from the two
37
38 periods within each scenario are comparable, but comparisons within the same time-period, across
39
40 scenarios may be inaccurate.
41
42

43
44 As the estimates are based on the population size in each grid cell, the added burden will be larger in
45
46 countries with larger increases in temperatures in densely populated areas. Because the population
47
48 of a city is considered a good predictor of the size of the urban heat island,³⁴ an increase in the
49
50 population living in urban areas will increase the numbers exposed and the temperature to which
51
52 they are exposed. Heat islands increase temperatures in urbanized areas compared to surrounding
53
54 areas, also reducing cooling during the night time.³⁵ These factors combined are likely to magnify the
55
56 health burden during a heatwave.³⁶ The spatial scale of the climate models makes them unable to
57
58
59
60

1
2
3 take the urban heat island effect into account. Together with the urbanization of Europe, this could
4
5 potentially result in underestimation of actual consequences/RHAs in the future period because the
6
7 same population size and composition were assumed in both periods. In addition, climate change is
8
9 likely to increase ozone concentrations that would add an additional health burden for people at risk
10
11 of respiratory diseases. The number of deaths and RHAs, due to a change in ozone, is expected to
12
13 change in the future.³⁷ A sensitivity analysis in the PHEWE study however showed that the exposure-
14
15 response relationship for heat did not substantially differ between models taking ozone into account
16
17 and models adjusting for NO₂ alone.³

18
19
20 We used the same thresholds for the reference and future period to isolate the effect of climate
21
22 change. This would tend to overestimate the increased impact because there will undoubtedly be
23
24 biological and/or social adaptations that will reduce future health temperature-related burdens.
25
26 However, given the uncertainty of such adaptation effects, we choose to not incorporate such effects
27
28 in the projections. Recent studies in the U.S indicate heat-related health burdens have decreased,³⁸
29
30 indicating that some adaptation is taking place. An opposite trend appears to be occurring in
31
32 Stockholm, with an increase of the risk of mortality associated with heat during the 1990s.³⁹ A study
33
34 looking at the impact of heat on mortality, before and after the implementation of a heat warning
35
36 system, in Italy shows that the effects of extreme heat can be reduced while the effects of moderate
37
38 increased average temperatures remains similar.⁴⁰ The magnitude and the extent of future
39
40 adaptation is, of course, highly uncertain, and will vary between and within countries. Nevertheless,
41
42 cities with higher thresholds seem to have higher risk ratio coefficients.^{3 41} In an effort to estimate
43
44 future and presumably higher thresholds, one must also adjust the risk ratio coefficients. This would
45
46 result in fewer days of elevated risk, but the risk increase on each occasion could be higher due to
47
48 the higher RRs.

49
50
51 Table 2 shows how the results vary by global climate models and greenhouse gas scenarios. This
52
53
54
55
56
57
58
59
60 study exemplifies the magnitude of the difference between projections made by a model with

1
2
3 different initial conditions and between different models with the same initial conditions. The results
4
5 indicate the range of these estimates is large. The mean increase over the five projections however,
6
7 provides confidence that the number of RHAs will increase over Europe as the climate continues to
8
9 change. The results from the different greenhouse gas emission scenarios are somewhat inconsistent
10
11 with what is expected based on the characteristics of the scenarios. The scenario A2, which is
12
13 considered a high emission scenario, shows lower estimates of the increase in RHAs than the A1B
14
15 scenario, which is considered a middle of the road scenario. This result is from the different regional
16
17 climate models rather than the emission scenarios themselves. Up until 2050, the estimated
18
19 temperature increase is actually estimated to be higher in A1B than A2 according to IPCC.
20
21

22
23 To improve the estimates of the impacts of heat on RHAs, detailed data on emergency room visits
24
25 and admissions during summer are needed, such as the proportion of emergency department cases
26
27 admitted to hospital, and the fatality rate. A better understanding of vulnerable groups is needed,
28
29 including how these groups could change over time. For example, the portion of the population aged
30
31 over 65 will increase from 17 to 29%⁴² by 2050, significantly increasing the size of this vulnerable
32
33 group. COPD is mainly a disease of the elderly. Persons suffering from COPD appear especially
34
35 vulnerable to heat⁴³ and the incidence of COPD is likely to increase⁴⁴ with the consequence that the
36
37 health impacts of heat may increase in future years. This study was limited to estimating the impacts
38
39 on the total population because few age-specific relationships were reported and because age-
40
41 stratified data were only available on a coarser spatial scale than the climate data, which would
42
43 reduce the benefits of having spatial climate data. When spatial population data, such as the
44
45 Eurostat data, have a finer age-stratification, a more detailed assessment will be possible.
46
47

48
49 Normal weather patterns in the future will be different than those of today, both for average
50
51 temperature and extreme weather events. As the number of heat related RHAs are expected to
52
53 increase on the sub-continental scale, additional national or regional projections of the future health
54
55 burden from heat are needed, taking into account possible changes in exposure and vulnerability.
56
57
58
59
60

CONCLUSIONS

Projected changes in temperature and the number of extreme heat events with climate change could substantially influence respiratory morbidity across Europe. Analyses projected that both the future proportion of annual RHAs attributed to heat and the relative change in heat related RHAs will be largest in Southern Europe, where they are expected to nearly triple. Eastern Europe can expect the smallest increase in heat-related RHAs, where they are estimated to approximately double. For all of Europe, the number of respiratory heat-related hospital admissions is projected to be 26,000 annually in the future period (2021–2050) compared with 11,000 in the reference period (1981–2010). The estimates presented rely on the assumption that no additional adaptation occurs. Future studies should elaborate and quantify the possible effects of different adaptation assumptions applied to regional conditions.

This work was supported by the EU-funded Climate-Trap project (contract EAHC 20081108) and by the Swedish Environmental Protection Agency through the research programme CLEO – Climate Change and Environmental Objectives. We would also like to acknowledge Estonia's Ministry of Education for providing resources to H. Orru with the grant SF0180060s09.

Contribution statement: CÅ, BF, HO has planned and designed the study. GS represented SMHI and provided climate data. CÅ ran the analyses, drafted the first manuscript and is guarantor. All authors have participated in the interpretation of results and revised the manuscript.

The specific estimates for each of the EU27 countries, for each of the climate projections is available after contact with corresponding author.

REFERENCES

1. Loddenkemper R, Gibson G, Sibille Y. European lung white book. *European Respiratory Society/European Lung Foundation, Sheffield, UK, ERSJ* 2003;1-182.
2. Ayres JG, Forsberg B, Annesi-Maesano I, et al. Climate change and respiratory disease: European Respiratory Society position statement. *Eur Respir J* 2009;34(2):295-302.
3. Michelozzi P, Accetta G, De Sario M, et al. High temperature and hospitalizations for cardiovascular and respiratory causes in 12 European cities. *Am J Respir Crit Care Med* 2009;179(5):383-9.
4. Rocklöv J, Forsberg B. Comparing approaches for studying the effects of climate extremes - a case study of hospital admissions in Sweden during an extremely warm summer. *Glob Health Action* 2009;2.
5. Kovats RS, Hajat S. Heat stress and public health: a critical review. *Annu Rev Public Health* 2008;29:41-55.
6. Knowlton K, Rotkin-Ellman M, King G, et al. The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect* 2009;117(1):61-7.
7. Rey G, Jouglé E, Fouillet A, et al. The impact of major heat waves on all-cause and cause-specific mortality in France from 1971 to 2003. *Int Arch Occup Environ Health* 2007;80(7):615-26.
8. Lin S, Luo M, Walker RJ, et al. Extreme high temperatures and hospital admissions for respiratory and cardiovascular diseases. *Epidemiology* 2009;20(5):738-46.
9. Forsberg B, Braback L, Keune H, et al. An expert assessment on climate change and health - with a European focus on lungs and allergies. *Environ Health* 2012;11 Suppl 1:S4.
10. Ye X, Wolff R, Yu W, et al. Ambient Temperature and Morbidity: A Review of Epidemiological Evidence. *Environ Health Perspect* 2011.
11. Semenza JC, McCullough JE, Flanders WD, et al. Excess hospital admissions during the July 1995 heat wave in Chicago. *Am J Prev Med* 1999;16(4):269-77.
12. Dhainaut JF, Claessens YE, Ginsburg C, et al. Unprecedented heat-related deaths during the 2003 heat wave in Paris: consequences on emergency departments. *Crit Care* 2004;8(1):1-2.
13. IPCC. *Climate Change 2007 - The Physical Science Basis: Working Group I Contribution to the Fourth Assessment Report of the IPCC*. Cambridge, UK and New York, NY, USA: Cambridge University Press, 2007.
14. Nikulin G, Kjellström E, Hansson ULF, et al. Evaluation and future projections of temperature, precipitation and wind extremes over Europe in an ensemble of regional climate simulations. *Tellus A* 2011;63(1):41-55.
15. Schar C, Vidale PL, Luthi D, et al. The role of increasing temperature variability in European summer heatwaves. *Nature* 2004;427(6972):332-6.
16. Stott PA, Stone DA, Allen MR. Human contribution to the European heatwave of 2003. *Nature* 2004;432(7017):610-4.
17. Rebetez M, Dupont O, Giroud M. An analysis of the July 2006 heatwave extent in Europe compared to the record year of 2003. *Theoretical and Applied Climatology* 2009;95(1):1-7.
18. Barriopedro D, Fischer EM, Luterbacher J, et al. The Hot Summer of 2010: Redrawing the Temperature Record Map of Europe. *Science* 2011.
19. Samuelsson P, Jones CG, Willen U, et al. The Rossby Centre Regional Climate model RCA3: model description and performance. *Tellus Series a-Dynamic Meteorology and Oceanography* 2011;63(1):4-23.
20. Collins WD, Bitz CM, Blackmon ML, et al. The Community Climate System Model version 3 (CCSM3). *J Climate* 2006;19(11):2122-43.
21. Roeckner E, Bäuml G, Bonaventura L, et al. The atmospheric general circulation model ECHAM 5. PART I: Model description. Hamburg, Germany: MPI für Meteorologie, 2003.
22. Johns TC, Durman CF, Banks HT, et al. The new Hadley Centre Climate Model (HadGEM1): Evaluation of coupled simulations. *J Climate* 2006;19(7):1327-53.

23. Roeckner E, Arpe K, Bengtsson, et al. The atmospheric general circulation model {ECHAM}-4: {M}odel description and simulation of present-day climate, 1996:103-23.
24. Nakicenovic N, Alcamo J, Davis G, et al. Special report on emissions scenarios: a special report of Working Group III of the Intergovernmental Panel on Climate Change: Pacific Northwest National Laboratory, Richland, WA (US), Environmental Molecular Sciences Laboratory (US), 2000.
25. Ballester J, Robine JM, Herrmann FR, et al. Long-term projections and acclimatization scenarios of temperature-related mortality in Europe. *Nat Commun* 2011;2:358.
26. Knowlton K, Lynn B, Goldberg RA, et al. Projecting heat-related mortality impacts under a changing climate in the New York City region. *American Journal of Public Health* 2007;97(11):2028-34.
27. Baccini M, Kosatsky T, Analitis A, et al. Impact of heat on mortality in 15 European cities: attributable deaths under different weather scenarios. *J Epidemiol Community Health* 2011;65(1):64-70.
28. Beguin A, Hales S, Rocklov J, et al. The opposing effects of climate change and socio-economic development on the global distribution of malaria. *Global Environmental Change-Human and Policy Dimensions* 2011;21(4):1209-14.
29. Goldewijk KK, Beusen A, Janssen P. Long-term dynamic modeling of global population and built-up area in a spatially explicit way: HYDE 3.1. *Holocene* 2010;20(4):565-73.
30. Lawrence MG. The Relationship between Relative Humidity and the Dewpoint Temperature in Moist Air: A Simple Conversion and Applications. *Bulletin of the American Meteorological Society* 2005;86(2):225-33.
31. Parsons KC. *Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort, and performance*: Taylor & Francis, 2003.
32. Steadman RG. A Universal Scale of Apparent Temperature. *J Clim Appl Meteorol* 1984;23(12):1674-87.
33. Lin S, Hsu WH, Van Zutphen AR, et al. Excessive heat and respiratory hospitalizations in new york state: estimating current and future public health burden related to climate change. *Environ Health Perspect* 2012;120(11):1571-7.
34. Oke TR. City size and the urban heat island. *Atmospheric Environment (1967)* 1973;7(8):769-79.
35. Holmer B, Thorsson S, Eliasson I. COOLING RATES, SKY VIEW FACTORS AND THE DEVELOPMENT OF INTRA - URBAN AIR TEMPERATURE DIFFERENCES. *Geografiska Annaler: Series A, Physical Geography* 2007;89(4):237-48.
36. Nicholls N, Skinner C, Loughnan M, et al. A simple heat alert system for Melbourne, Australia. *Int J Biometeorol* 2008;52(5):375-84.
37. Orru H, Andersson C, Ebi KL, et al. Impact of climate change on ozone related mortality and morbidity in Europe. *Eur Respir J* 2012 Jun 27. [Epub ahead of print].
38. Davis RE, Knappenberger PC, Michaels PJ, et al. Changing heat-related mortality in the United States. *Environmental Health Perspectives* 2003;111(14):1712-18.
39. Rocklov J, Forsberg B, Meister K. Winter mortality modifies the heat-mortality association the following summer. *Eur Respir J* 2009;33(2):245-51.
40. Schifano P, Leone M, De Sario M, et al. Changes in the effects of heat on mortality among the elderly from 1998-2010: results from a multicenter time series study in Italy. *Environmental Health* 2012;11(1):58.
41. Baccini M, Biggeri A, Accetta G, et al. Heat effects on mortality in 15 European cities. *Epidemiology* 2008;19(5):711-9.
42. Giannakouris K. Ageing characterises the demographic perspectives of the European societies. *Statistics in focus* 2008;72:2008.
43. Schwartz J. Who is sensitive to extremes of temperature? A case-only analysis. *Epidemiology* 2005;16(1):67-72.
44. Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet* 2007;370(9589):765-73.

1
2
3 **Figure 1. The range of the absolute increase in RHAs attributed to heat between the two**
4 **periods (1981-2010, 2021-2050) as proportion the annual expected number of RHAs for each of the**
5 **27 countries. The points show the highest and lowest estimate from four climate models under**
6 **two emission scenarios.**
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Heat related respiratory hospital admissions in Europe in a changing climate

Authors:

Christofer Åström^{1*}, Hans Orru^{1,2}, Joacim Rocklöv^{1,3}, Gustav Strandberg⁴, Kristie L. Ebi⁵, Bertil Forsberg¹

Affiliations:

¹ Department of Public Health and Clinical Medicine, Occupational and Environmental Medicine, Umeå University, Umeå, Sweden; ² Department of Public Health, University of Tartu, Tartu, Estonia; ³ Department of Public Health and Clinical Medicine, Epidemiology and Global Health, Umeå University, Umeå, Sweden; ⁴ Swedish Meteorological and Hydrological Institute, Norrköping, Sweden; ⁵ ClimAdapt, Stanford University, Los Altos, California, USA

Corresponding author:

* Author to whom correspondence should be addressed:

Christofer Åström,

Occupational and Environmental Medicine, Department of Public Health and Clinical Medicine, Umeå University, SE 901 87, Sweden

+4690 785 0874

christofer.astrom@envmed.umu.se

Keywords: Climate change, respiratory health, public health, heat

Word count: ~~3045~~3246

ARTICLE SUMMARY

Article focus

To assess how heat-related respiratory hospital admissions in Europe ~~are expected to~~could change with ~~the global warming~~increasing temperatures in the near future. ~~With~~Within a range of climate change projections, explore ~~the range of the~~ different impact estimates.

Key messages

The 30-year mean annual increase in heat-related respiratory hospital admissions ~~related to heat~~ between the periods 1981-2010 and 2021-2050 can be counted in the 10's of thousands. because of projected temperature increases. The increase will be different in different parts of Europe with the largest relative increase in Southern Europe.

Strengths and limitations

The study takes the spatial variation in ~~both~~ climate and population density into account in ~~the~~ impact calculations. ~~This makes,~~ making the estimates valid for countries with large national differences in ~~population densities and climate~~these factors.

The study results ~~exemplifies~~exemplify some the variation of ~~the~~ impacts ~~depending on~~under different climate change scenarios, as well as the variations within one ~~climate~~ scenario depending on the underlying climate model.

The use of ~~real~~ spatial population data ~~has~~ limited the study to ~~explore~~exploring the change in all age respiratory hospital ~~admissions without demographic trends.~~ ~~The fact that the elderly is a well-known risk group when~~for studying heat-related illness, combined with the future ~~rise of~~increase in the proportion of elderly people ~~in the population may suggest that,~~ suggests the estimated impacts are ~~rather~~likely to be conservative

ABSTRACT

Respiratory diseases are ranked second in Europe in terms of mortality, prevalence, and costs. Studies have shown that extreme heat has a large impact on mortality and morbidity, with a large relative increase for respiratory diseases. Expected increases in mean temperature and the number of extreme heat events over the coming decades due to climate change raise questions about the possible health impacts.

Heat-related respiratory hospital admissions under a changing climate are projected using multi-city epidemiologic exposure-response relationships applied to gridded population data and country-specific baseline respiratory hospital admission rates. Times-series of temperatures are simulated with a regional climate model based on four global climate models, under two greenhouse gas emission scenarios (A1B & A2).

Between a reference period (1981–2010) and a future period (2021–2050), the total number of respiratory hospital admissions attributed to heat is projected to be larger in Southern Europe, with three-times more heat attributed respiratory hospital admissions in the future period. The smallest change was estimated in Eastern Europe with about a two-fold increase. For all of Europe, the number of respiratory heat-related hospital admissions is projected to be 26,000 annually in the future period compared with 11,000 in the reference period.

The results suggest that projected effects of climate change on temperature and the number of extreme heat events could substantially influence respiratory morbidity across Europe.

INTRODUCTION

Respiratory diseases are ranked second in Europe in terms of mortality, prevalence, and costs.¹ This burden is expected to increase, partly due to a changing climate.² An environmental factor with a large impact on mortality and morbidity in Europe is extreme heat, with large effects on respiratory diseases.³⁻⁵ Physiological effects of exposure to heat can be directly heat-related (heat stroke, heat fatigue and dehydration) or can contribute to worsening of respiratory and cardiovascular diseases, electrolyte disorders, and kidney problems.⁴⁻⁷ The reasons for an increase in respiratory admissions may be several. Elderly with respiratory diseases such as COPD are less fit and suffer often from circulatory problems. Heat influenced admissions due to chronic airway obstruction and asthma increased more than admissions due to chronic bronchitis in a study from New York City,⁸ but the daily number of admissions seldom allow a study of specific diagnoses. In a recent expert elicitation amongst European researchers engaged in environmental medicine or respiratory health, extreme heat stood out as most important climate-related pathway to adverse impacts on respiratory health, more important than changes in air pollutants and allergens.⁹ A review found that heatwaves have a stronger relative impact on mortality than on emergency room visits and hospital admissions, suggesting that many individuals die before they can get to the hospital.¹⁰ However, several studies confirm that heat also affects health care utilisation.^{3 11} During the heatwave of 2003 in France, the Assistance Public Hôpitaux de Paris recorded 2,400 additional visits to the emergency care units and 1,900 additional hospital admissions.¹² During the heatwave in California in 2006 there were almost 1,200 hospital admissions together with more than 16,000 visits to the emergency departments.⁶

It is possible that hospital admissions could increase in the future due to temperature changes projected with climate change, assuming no additional acclimatisation. Projected changes include an increase in global mean temperature of 1.8°C to 4.0°C by the end of this century, with larger increases over land areas at high latitudes, changing seasonal temperatures, and increases in the frequency, intensity, duration, and spatial extent of heatwaves.¹³ Recent projections for 2070–2100

1
2
3 suggest that maximum temperatures experienced once every 20 years during the period 1961–1990
4 could be expected as often as every year in southern Europe and every 3rd-5th year in northern
5 Europe.¹⁴ This means that Europe should expect an increase in both mean temperatures and the
6 number of extreme heat events. Recent heatwaves are consistent with projections. For example, the
7 heatwave that occurred in Europe 2003 could be expected to return every 46,000 years, based on
8 the temperature distribution for the years 1864–2000.¹⁵ However, such extreme hot conditions may
9 become much more common in the end of this century due to anthropogenic climate changes.¹⁶ The
10 uncertainty in this estimated return time is quite high, with a lower bound of the 90% confidence
11 interval of 9,000 years. Even so, another heatwave occurred in 2006 with the most anomalous July
12 temperature ever measured in Europe.¹⁷ In 2010, Eastern Europe experienced a heatwave with
13 summer temperatures higher than the last 140 years, resulting in roughly 55,000 excess deaths in
14 Russia.¹⁸

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30 The possible impacts of these changes on morbidity, particularly respiratory health, are relatively
31 unexplored.
32
33

34 35 36 **Aim**

37 The aim of this study is to assess the extent to which changes in the frequency of hot days due to
38 climate change over the next 40 years could affect heat related respiratory hospital admissions
39 (RHAs) in Europe. Using a range of climate projections, we estimate the change in hospital
40 admissions between a reference period and a future period.
41
42
43
44
45
46
47

48 49 **METHODS**

50 51 52 **Climate change and temperature modelling**

53 The Rossby Centre regional atmospheric climate model RCA3¹⁹ was developed by SMHI (Swedish
54 meteorological and hydrological institute) to dynamically downscale results from the global climate
55 models CCSM3²⁰, ECHAM5²¹, HadleyCM3²² and ECHAM4²³ to a higher resolution over Europe. RCA3 is
56
57
58
59
60

1
2
3 run on a horizontal grid spacing of 0.44° (corresponding to approximately 50 km) and a time step of
4
5 30 minutes. Projections are based on the global greenhouse gas emission scenarios A1B and A2
6
7 (scenarios are described in detail in the SRES²⁴). Both have been used in climate change health
8
9 impact assessments (HIA).²⁵⁻²⁸ A1B is a “middle of the road” scenario and A2 is considered a high
10
11 emission scenario, although recent greenhouse gas emissions have been higher. Data from one
12
13 climate model, under one climate change scenario, is referred to as one climate change projection.
14
15

16
17 We use aggregated daily projections of maximum temperature and relative humidity data to
18
19 estimate exposure for the periods 1981–2020 (reference period) and 2021–2050 (future period).
20
21

22 **Population and morbidity rates**

23
24 The annual country specific rate of ~~respiratory hospital admissions~~ RHAs (ICD-9:460–519) between
25
26 2005 and 2010 were extracted from the WHO’s European Health for All Database
27
28 (<http://data.euro.who.int/hfadb>) for the EU27 countries. In the dataset, national hospital admissions
29
30 data are provided by national public health institutes, health ministries or corresponding functions.
31
32 The ICD-9 codes were preferred because the epidemiological studies in the PHEWE study were based
33
34 on ICD-9 RHAs. The mean value over the six years was used as a baseline morbidity rate. In order to
35
36 have a fine spatial resolution, official population data were from the HYDE theme within the
37
38 Netherlands Environmental Assessment Agency.²⁹ These data are gridded on a 0.0833° resolution
39
40 (approximately 9,45km) and matched with the climatic data by summing the population within each
41
42 climatic grid cell.
43
44
45

46 **Exposure-response assumption**

47
48 The impact calculations were based on the relationship between heat and RHAs (ICD-9:460–519)
49
50 estimated within the European PHEWE (Assessment and Prevention of Acute Health Effects of
51
52 Weather Conditions in Europe) project.³ This relationship was based on analyses of 12 European
53
54 cities over the 1990s, using the daily maximum of apparent temperature (AT), where AT is a
55
56 combination of the measured temperature and the dew point temperature:
57
58
59
60

$$AT = -2.653 + 0.944 * T + 0.0153 * (DT)^2$$

where T is the air temperature and DP the dew point temperature, which can be derived from temperature and relative humidity.³⁰ The AT accounts for heat stress related not only to the absolute temperature but also to the saturation of the surrounding air that makes it harder to regulate body temperature by sweating, the most important mechanism to maintain a healthy body temperature.³¹

³² The PHEWE model also included potential confounders such as air pollution, holidays, week days etc. The study used a 0–3 day lag of the maximum AT and concluded that the 90th percentile of this exposure variable for the summer months, April to September, is the appropriate threshold value for the heat-morbidity function.³ We calculated the 90th percentile for the summer months for each grid cell from the climate data for the 1990s, yielding individual thresholds for each grid cell for each climate projection.

The PHEWE study calculated the relative risk (RR) coefficients for each of the 12 cities and then combined these into two meta-coefficients, for North-Continental cities 1.012 (95% CI 1.001–1.022) and Mediterranean cities 1.021 (95% CI 1.006–1.036), associated with a 1-degree increase above the temperature threshold. Meta-coefficients were more suitable as they allowed calculation of the change in RHAs for countries with and without a city in the PHEWE Study. We assigned the Mediterranean coefficient to Portugal and the Mediterranean countries. The rest of Europe was assigned the RR for the North continental cities. Statistically significant coefficients were found for all ages and ageages 75+ in the PHEWE study. Because the population data were not stratified by age, we used the PHEWE study coefficients for all ages.

Projections

The final data used in the impact calculations were based on the climate data grid with a resolution of 0.44° resulting in 8,075 grid cells over Europe. The population data, RR coefficients and baseline morbidity rates were projected for each grid cell. The countries within EU27 were grouped into four

regions, Northern, Western, Eastern and Southern Europe, according to the United Nations classification scheme (<http://unstats.un.org/unsd/methods/m49/m49regin.htm#europe>).

Thus, each grid cell includes (1) a daily time-series of 0–3 day lag maximum AT for each climate change projection; (2) a grid cell specific temperature-morbidity threshold and a location-based RR coefficient; and (3) population. For each grid cell, the AT was compared to the specific threshold for each day in the two time periods and risk estimates were calculated for each day. The expected number of daily RHAs in each cell was calculated using the population and the expected daily number of RHAs per capita, based on the national average of the grid cell.

$$RHA_{i,t} = RR_i^{(AT_{i,t} - Thres_i)_+} * Pop_i * RHApc_i$$

where $RHA_{i,t}$ is the number of RHAs attributable to heat in grid cell i at time t , RR_i the RR coefficient in grid cell i , $(AT_{i,t} - Thres_i)_+$ the difference between the 0–3 lag AT and the threshold for grid cell i at time t if $AT_{i,t}$ were greater than the threshold and 0 otherwise, Pop_i and $RHApc_i$ the population and expected number RHAs per capita in grid cell i .

For each country, the estimated number of RHAs attributed to heat was calculated for each climate change projection, for both the reference and the future period. The estimated number of RHAs attributed to heat was then transformed into the proportion of the expected annual number RHAs for each country (data available from the authors). Mean estimates were calculated for each region for each climate change projection.

Sensitivity analysis

Because the study results could heavily depend on the RR coefficients and thresholds used, we assess whether the estimates appeared sensitive to the region-specific meta-coefficients. We investigated the grids that contained the cities in the PHEWE study using both the location (Mediterranean or North-Continental) and city specific RR coefficients.³ We then compared the results with respect to the attributed proportion of RHAs and the proportional change in heat related RHAs.

RESULTS

The periods of warm days will increase in the future. For the cities included in the sensitivity analysis, the temperature threshold was exceeded annually, on average, by 20 days in the baseline period and 40 days in the future period, ~~on average each year~~.

In the future period, approximately 0.4% of the annual numbers of RHAs in Europe was estimated to be due to heat (Table 1), based on the mean estimates over the climate change projections. In absolute terms, assuming all else equal, this represents about 26,000 cases annually in Europe. This should be compared to the reference period where approximately 0.18% of all RHAs were attributed to heat or about 11,000 cases annually. Thus, the results suggest more than a relative doubling of the RHAs attributed to heat in Europe.

Table 1. The estimated proportion of RHAs attributed to heat for each region. Intervals describe the highest and lowest national estimate in each region

Region	1981–2010	2021–2050	Change
Eastern Europe	0.17% (0.16% – 0.19%)	0.31% (0.29% – 0.35%)	0.14% (0.11% – 0.17%)
Northern Europe	0.13% (0.10% – 0.15%)	0.27% (0.19% – 0.32%)	0.14% (0.07% – 0.17%)
Southern Europe	0.23% (0.18% – 0.26%)	0.64% (0.42% – 0.68%)	0.41% (0.23% – 0.44%)
Western Europe	0.18% (0.16% – 0.20%)	0.39% (0.34% – 0.45%)	0.21% (0.17% – 0.26%)
EU27	0.18% (0.10% – 0.26%)	0.40% (0.19% – 0.68%)	0.21% (0.07% – 0.44%)

On the regional level, the five projections estimate increases in the number of heat-related RHAs for Europe (Table 2). However, in one climate change projection, The Czech Republic, Hungary, Poland and Slovakia were estimated to have a decrease in heat-related RHAs (Fig 1). The countries with the highest estimated increase also show the largest range between the highest and lowest estimate. The Scandinavian and Baltic countries show the smallest range between the highest and lowest estimates along with small increases of the mean estimates.

Table 2. Future increase in heat-related RHAs based on the four climate models, under two emission scenarios, as the percentage of the annual expected number of RHAs in each region

Climate model	CCSM3	ECHAM5	HadCM3	ECHAM4	ECHAM5
<i>Greenhouse gas scenario</i>	<i>A1B</i>			<i>A2</i>	
Eastern Europe	0.32%	0.08%	0.18%	0.12%	0.01%
Northern Europe	0.17%	0.09%	0.14%	0.20%	0.08%
Southern Europe	0.51%	0.29%	0.64%	0.45%	0.14%
Western Europe	0.30%	0.11%	0.26%	0.29%	0.06%
EU27	0.32%	0.13%	0.29%	0.26%	0.07%

There is variation among countries, with the largest increases in the Southern European countries and the smallest increase in Eastern Europe (Table 2). The relative change in the burden of RHA's in relation to the climate change scenarios investigated indicate a larger relative increase in Mediterranean countries (approximately three times) compared to Northern European countries (approximately two times).

Sensitivity analysis

The calculations using the city specific RR coefficients yielded a different proportion and number of estimated RHAs than the ones using the two meta-coefficients, as expected. However, the relative changes in the number of hospital admissions attributed to heat between the two time-periods were not affected by the change of the RR coefficients.

DISCUSSION

The analysis estimates that the number of RHAs will increase in a warmer climate with more hot days. Heat-related RHAs were projected to increase two- to three-fold in the future due to climate change. However, the proportion of respiratory admissions attributed to heat would remain rather small. This projection is in line with the results of a recent study ~~which that~~ estimates ~~that the~~

1
2
3 respiratory admissions due to excessive heat in New York State will increase 2-6 times from the
4
5 period 1991-2004 to the period 2080-2099.³³
6
7

8 | As most heat-related health outcomes occur during the warmest period of the year, presenting the
9
10 increase as a change in the proportion of the total number of annual RHAs can underestimate the
11
12 additional burden on the health care system during summer. As the threshold was exceeded for
13
14 approximately 40 days in one year, an annual increase of 0.21% would result in a 1.9% increase on
15
16 average during these 40 days. The annual numbers are used because the available baseline rates of
17
18 RHAs are expressed as an annual average. Applying results to sub-national scales could be
19
20 inappropriate because national averages summarize over considerable heterogeneity.
21
22
23

24 The results suggest a larger impact from heat in Southern Europe in the future period, centred on
25
26 year 2035, than in the eastern and northern parts. This is in line with many climate change
27
28 projections showing a larger relative increase in the number of extremely hot days in southern
29
30 Europe compared with northern Europe.¹⁴ However, to some extent this might also be explained by
31
32 model bias for northern Europe introduced by the RCA3 model, where the model appears to
33
34 underestimate temperature for the warmest days.¹⁹ This temperature bias is present in the reference
35
36 period, future period, and threshold values; therefore, the estimated numbers of RHAs from the two
37
38 periods within each scenario are comparable, but comparisons within the same time-period, across
39
40 scenarios may be inaccurate.
41
42
43

44 As the estimates are based on the population size in each grid cell, the added burden will be larger in
45
46 countries with larger increases in temperatures in densely populated areas. Because the population
47
48 of a city is considered a good predictor of the size of the urban heat island,³⁴ an increase in the
49
50 population living in urban areas will increase the numbers exposed and the temperature to which
51
52 they are exposed. Heat islands increase temperatures in urbanized areas compared to surrounding
53
54 areas, also reducing cooling during the night time.³⁵ These factors combined are likely to magnify the
55
56 health burden during a heatwave.³⁶ The coarse-spatial scale of the climate models makes them
57
58
59
60

1
2
3 unable to take the urban heat island effect into account. Together with the urbanization of Europe,
4
5 this could potentially result in underestimation of actual consequences/RHAs in the future period
6
7 because the same population size and composition were assumed in both periods. In addition,
8
9 climate change is likely to increase ozone concentrations that would add an additional health burden
10
11 for people at risk of respiratory diseases. The number of deaths and RHAs, due to a change in ozone,
12
13 is expected to change in the future.³⁷ A sensitivity analysis in the PHEWE study however showed that
14
15 the exposure-response relationship for heat did not substantially differ between models taking ozone
16
17 into account and models adjusting for NO₂ alone.³

18
19
20 We used the same thresholds for the reference and future period to isolate the effect of climate
21
22 change. This would tend to overestimate the increased impact because there will undoubtedly be
23
24 biological and/or social adaptations that will reduce future health temperature-related burdens.
25
26 However, given the uncertainty of such adaptation effects, we choose to not incorporate such effects
27
28 in the projections. Recent studies in the U.S indicate heat-related health burdens have decreased,³⁸
29
30 indicating that some adaptation is taking place. An opposite trend appears to be occurring in
31
32 Stockholm, with an increase of the risk of mortality associated with heat during the 1990s.³⁹ A study
33
34 looking at the impact of heat on mortality, before and after the implementation of a heat warning
35
36 system, in Italy shows that the effects of extreme heat can be reduced while the effects of moderate
37
38 increased average temperatures remains similar.⁴⁰ The magnitude and the extent of future
39
40 adaptation is, of course, highly uncertain, and will vary between and within countries. Nevertheless,
41
42 cities with higher thresholds seem to have higher risk ratio coefficients.^{3 41} In an effort to estimate
43
44 future and presumably higher thresholds, one must also adjust the risk ratio coefficients. This would
45
46 result in fewer days of elevated risk, but the risk increase on each occasion could be higher due to
47
48 the higher RRs.
49
50
51
52

53
54 Table [12](#) shows how the results vary by global climate models and greenhouse gas scenarios. This
55
56 study exemplifies the magnitude of the difference between projections made by a model with
57
58
59
60

different initial conditions and between different models with the same initial conditions. The results indicate the range of these estimates is large. The mean increase over the five projections however, provides confidence that the number of RHAs will increase over Europe as the climate continues to change. The results from the different greenhouse gas emission scenarios are somewhat inconsistent with what is expected based on the characteristics of the scenarios. The scenario A2, which is considered a high emission scenario, shows lower estimates of the increase in RHAs than the A1B scenario, which is considered a middle of the road scenario. This result is from the different regional climate models rather than the emission scenarios themselves. Up until 2050, the estimated temperature increase is actually estimated to be higher in A1B than A2 according to IPCC.

To improve the estimates of the impacts of heat on RHAs, detailed data on emergency room visits and admissions during summer are needed, such as the proportion of emergency department cases admitted to hospital, and the fatality rate. A better understanding of vulnerable groups is needed, including how these groups could change over time. For example, the portion of the population aged over 65 will increase from 17 to 29%⁴² by 2050, significantly increasing the size of this vulnerable group. COPD is mainly a disease of the elderly. Persons suffering from COPD appear especially vulnerable to heat⁴³ and the incidence of COPD is likely to increase⁴⁴ with the consequence that the health impacts of heat may increase in future years. This study was limited to estimating the impacts on the total population because few age-specific relationships were reported and because age-stratified data were only available on a coarser spatial scale than the climate data, which would reduce the benefits of having spatial climate data. When spatial population data, such as the Eurostat data, have a finer age-stratification, a more detailed assessment will be possible.

Normal weather patterns in the future will be different than those of today, both for average temperature and extreme weather events. As the number of heat related RHAs are expected to increase on the sub-continental scale, additional national or regional projections of the future health burden from heat are needed, taking into account possible changes in exposure and vulnerability.

CONCLUSIONS

Projected changes in temperature and the number of extreme heat events with climate change could substantially influence respiratory morbidity across Europe. Analyses projected that both the future proportion of annual RHAs attributed to heat and the relative change in heat related RHAs will be largest in Southern Europe, where they are expected to nearly triple. Eastern Europe can expect the smallest increase in heat-related RHAs, where they are estimated to approximately double. For all of Europe, the number of respiratory heat-related hospital admissions is projected to be 26,000 annually in the future period (2021–2050) compared with 11,000 in the reference period (1981–2010). The estimates presented rely on the assumption that no additional adaptation occurs. Future studies should elaborate and quantify the possible effects of different adaptation assumptions applied to regional conditions.

This work was supported by the EU-funded Climate-Trap project (contract EAHC 20081108) and by the Swedish Environmental Protection Agency through the research programme CLEO – Climate Change and Environmental Objectives. We would also like to acknowledge Estonia's Ministry of Education for providing resources to H. Orru with the grant SF0180060s09.

Contribution statement: CÅ, BF, HO has planned and designed the study. GS represented SMHI and provided climate data. CÅ ran the analyses, drafted the first manuscript and is guarantor. All authors have participated in the interpretation of results and revised the manuscript.

The specific estimates for each of the EU27 countries, for each of the climate projections is available after contact with corresponding author.

REFERENCES


1. Loddenkemper R, Gibson G, Sibille Y. European lung white book. *European Respiratory Society/European Lung Foundation, Sheffield, UK, ERSJ* 2003;1-182.
2. Ayres JG, Forsberg B, Annesi-Maesano I, et al. Climate change and respiratory disease: European Respiratory Society position statement. *Eur Respir J* 2009;34(2):295-302.
3. Michelozzi P, Accetta G, De Sario M, et al. High temperature and hospitalizations for cardiovascular and respiratory causes in 12 European cities. *Am J Respir Crit Care Med* 2009;179(5):383-9.
4. Rocklöv J, Forsberg B. Comparing approaches for studying the effects of climate extremes - a case study of hospital admissions in Sweden during an extremely warm summer. *Glob Health Action* 2009;2.
5. Kovats RS, Hajat S. Heat stress and public health: a critical review. *Annu Rev Public Health* 2008;29:41-55.
6. Knowlton K, Rotkin-Ellman M, King G, et al. The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect* 2009;117(1):61-7.
7. Rey G, Jouglé E, Fouillet A, et al. The impact of major heat waves on all-cause and cause-specific mortality in France from 1971 to 2003. *Int Arch Occup Environ Health* 2007;80(7):615-26.
8. Lin S, Luo M, Walker RJ, et al. Extreme high temperatures and hospital admissions for respiratory and cardiovascular diseases. *Epidemiology* 2009;20(5):738-46.
9. Forsberg B, Braback L, Keune H, et al. An expert assessment on climate change and health - with a European focus on lungs and allergies. *Environ Health* 2012;11 Suppl 1:S4.
10. Ye X, Wolff R, Yu W, et al. Ambient Temperature and Morbidity: A Review of Epidemiological Evidence. *Environ Health Perspect* 2011.
11. Semenza JC, McCullough JE, Flanders WD, et al. Excess hospital admissions during the July 1995 heat wave in Chicago. *Am J Prev Med* 1999;16(4):269-77.
12. Dhainaut JF, Claessens YE, Ginsburg C, et al. Unprecedented heat-related deaths during the 2003 heat wave in Paris: consequences on emergency departments. *Crit Care* 2004;8(1):1-2.
13. IPCC. *Climate Change 2007 - The Physical Science Basis: Working Group I Contribution to the Fourth Assessment Report of the IPCC*. Cambridge, UK and New York, NY, USA: Cambridge University Press, 2007.
14. Nikulin G, Kjellström E, Hansson ULF, et al. Evaluation and future projections of temperature, precipitation and wind extremes over Europe in an ensemble of regional climate simulations. *Tellus A* 2011;63(1):41-55.
15. Schar C, Vidale PL, Luthi D, et al. The role of increasing temperature variability in European summer heatwaves. *Nature* 2004;427(6972):332-6.
16. Stott PA, Stone DA, Allen MR. Human contribution to the European heatwave of 2003. *Nature* 2004;432(7017):610-4.
17. Rebetez M, Dupont O, Giroud M. An analysis of the July 2006 heatwave extent in Europe compared to the record year of 2003. *Theoretical and Applied Climatology* 2009;95(1):1-7.
18. Barriopedro D, Fischer EM, Luterbacher J, et al. The Hot Summer of 2010: Redrawing the Temperature Record Map of Europe. *Science* 2011.
19. Samuelsson P, Jones CG, Willen U, et al. The Rossby Centre Regional Climate model RCA3: model description and performance. *Tellus Series a-Dynamic Meteorology and Oceanography* 2011;63(1):4-23.
20. Collins WD, Bitz CM, Blackmon ML, et al. The Community Climate System Model version 3 (CCSM3). *J Climate* 2006;19(11):2122-43.
21. Roeckner E, Bäuml G, Bonaventura L, et al. The atmospheric general circulation model ECHAM 5. PART I: Model description. Hamburg, Germany: MPI für Meteorologie, 2003.
22. Johns TC, Durman CF, Banks HT, et al. The new Hadley Centre Climate Model (HadGEM1): Evaluation of coupled simulations. *J Climate* 2006;19(7):1327-53.

23. Roeckner E, Arpe K, Bengtsson, et al. The atmospheric general circulation model {ECHAM}-4: {M}odel description and simulation of present-day climate, 1996:103-23.
24. Nakicenovic N, Alcamo J, Davis G, et al. Special report on emissions scenarios: a special report of Working Group III of the Intergovernmental Panel on Climate Change: Pacific Northwest National Laboratory, Richland, WA (US), Environmental Molecular Sciences Laboratory (US), 2000.
25. Ballester J, Robine JM, Herrmann FR, et al. Long-term projections and acclimatization scenarios of temperature-related mortality in Europe. *Nat Commun* 2011;2:358.
26. Knowlton K, Lynn B, Goldberg RA, et al. Projecting heat-related mortality impacts under a changing climate in the New York City region. *American Journal of Public Health* 2007;97(11):2028-34.
27. Baccini M, Kosatsky T, Analitis A, et al. Impact of heat on mortality in 15 European cities: attributable deaths under different weather scenarios. *J Epidemiol Community Health* 2011;65(1):64-70.
28. Beguin A, Hales S, Rocklov J, et al. The opposing effects of climate change and socio-economic development on the global distribution of malaria. *Global Environmental Change-Human and Policy Dimensions* 2011;21(4):1209-14.
29. Goldewijk KK, Beusen A, Janssen P. Long-term dynamic modeling of global population and built-up area in a spatially explicit way: HYDE 3.1. *Holocene* 2010;20(4):565-73.
30. Lawrence MG. The Relationship between Relative Humidity and the Dewpoint Temperature in Moist Air: A Simple Conversion and Applications. *Bulletin of the American Meteorological Society* 2005;86(2):225-33.
31. Parsons KC. *Human thermal environments: the effects of hot, moderate, and cold environments on human health, comfort, and performance*: Taylor & Francis, 2003.
32. Steadman RG. A Universal Scale of Apparent Temperature. *J Clim Appl Meteorol* 1984;23(12):1674-87.
33. Lin S, Hsu WH, Van Zutphen AR, et al. Excessive heat and respiratory hospitalizations in new york state: estimating current and future public health burden related to climate change. *Environ Health Perspect* 2012;120(11):1571-7.
34. Oke TR. City size and the urban heat island. *Atmospheric Environment (1967)* 1973;7(8):769-79.
35. Holmer B, Thorsson S, Eliasson I. COOLING RATES, SKY VIEW FACTORS AND THE DEVELOPMENT OF INTRA - URBAN AIR TEMPERATURE DIFFERENCES. *Geografiska Annaler: Series A, Physical Geography* 2007;89(4):237-48.
36. Nicholls N, Skinner C, Loughnan M, et al. A simple heat alert system for Melbourne, Australia. *Int J Biometeorol* 2008;52(5):375-84.
37. Orru H, Andersson C, Ebi KL, et al. Impact of climate change on ozone related mortality and morbidity in Europe. *Eur Respir J* 2012 Jun 27. [Epub ahead of print].
38. Davis RE, Knappenberger PC, Michaels PJ, et al. Changing heat-related mortality in the United States. *Environmental Health Perspectives* 2003;111(14):1712-18.
39. Rocklov J, Forsberg B, Meister K. Winter mortality modifies the heat-mortality association the following summer. *Eur Respir J* 2009;33(2):245-51.
40. Schifano P, Leone M, De Sario M, et al. Changes in the effects of heat on mortality among the elderly from 1998-2010: results from a multicenter time series study in Italy. *Environmental Health* 2012;11(1):58.
41. Baccini M, Biggeri A, Accetta G, et al. Heat effects on mortality in 15 European cities. *Epidemiology* 2008;19(5):711-9.
42. Giannakouris K. Ageing characterises the demographic perspectives of the European societies. *Statistics in focus* 2008;72:2008.
43. Schwartz J. Who is sensitive to extremes of temperature? A case-only analysis. *Epidemiology* 2005;16(1):67-72.
44. Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet* 2007;370(9589):765-73.

1
2
3 **Figure 1. The range of the absolute increase in RHAs attributed to heat between the two**
4 **periods (1981-2010, 2021-2050) as proportion the annual expected number of RHAs for each of the**
5 **27 countries. The points show the highest and lowest estimate from four climate models under**
6 **two emission scenarios.**
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



130x90mm (300 x 300 DPI)

view only