



**A prospective cohort study to refine and validate the *Panic Screening Score* for identifying panic attacks associated with unexplained chest pain in the Emergency Department**

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-003877
Article Type:	Protocol
Date Submitted by the Author:	22-Aug-2013
Complete List of Authors:	Foldes-Busque, Guillaume; École de psychologie, Université Laval, ; Centre de santé et de services sociaux Alphonse-Desjardins, Research Center of the University-Affiliated Hospital of Lévis, Emergency medicine Denis, Isabelle; Centre de santé et de services sociaux Alphonse-Desjardins, Research Center of the University-Affiliated Hospital of Lévis, Poitras, Julien; Centre de santé et de services sociaux Alphonse-Desjardins, Research Center of the University-Affiliated Hospital of Lévis, Emergency medicine; Université Laval, Family and Emergency Medicine Fleet, Richard; Centre de santé et de services sociaux Alphonse-Desjardins, Research Center of the University-Affiliated Hospital of Lévis, Emergency medicine; Université Laval, Family and Emergency Medicine Archambault, Patrick; Centre de santé et de services sociaux Alphonse-Desjardins, Research Center of the University-Affiliated Hospital of Lévis, Emergency medicine; Université Laval, Family and Emergency medicine Dionne, Clermont; Centre de recherche FRQS du Centre hospitalier universitaire (CHU) de Québec, Hôpital du St-Sacrement,
<b>Primary Subject Heading</b>:	Emergency medicine
Secondary Subject Heading:	Epidemiology, Mental health
Keywords:	Unexplained chest pain, Non-cardiac chest pain, Panic attacks, Panic disorder, Emergency medicine
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
Authorization for funding.pdf Ethic approval.pdf	

SCHOLARONE™  
Manuscripts

**Title:** A prospective cohort study to refine and validate the *Panic Screening Score* for identifying panic attacks associated with unexplained chest pain in the Emergency

Department

**Guillaume Foldes-Busque, Ph.D., Psy.D.<sup>a,b\*</sup>, Isabelle Denis, Psy.D., Ph.D.<sup>b</sup>, Julien Poitras, M.D.<sup>b,c</sup>, Richard P. Fleet, M.D., Ph.D.<sup>b,c</sup>, Patrick Archambault M.D., M.Sc.<sup>b,c</sup> & Clermont E. Dionne, Ph.D.<sup>d</sup>**

<sup>a</sup> École de psychologie, Faculté des sciences sociales, Université Laval, Québec (Québec), Canada

<sup>b</sup> Centre de santé et de services sociaux Alphonse-Desjardins, Research Center of the University-Affiliated Hospital of Lévis, Québec (Québec), Canada

<sup>c</sup> Department of Family and Emergency Medicine, Faculté de médecine, Université Laval, Québec (Québec), Canada

<sup>d</sup> Centre de recherche FRQS du Centre hospitalier universitaire (CHU) de Québec, Hôpital du St-Sacrement, Québec (Québec), Canada

**\*Corresponding author:**

Guillaume Foldes-Busque; Centre hospitalier affilié universitaire de Lévis, Centre de recherche, 143, rue Wolfe, Lévis, Québec, Canada, G6V 3Z1, E-mail: [guillaume.foldes-busque@psy.ulaval.ca](mailto:guillaume.foldes-busque@psy.ulaval.ca)

**Keywords:** Unexplained chest pain, Non-cardiac chest pain, Panic attacks, Panic disorder, Emergency medicine.

**Word count:** 4280

## ABSTRACT

### Introduction

Panic-like anxiety (panic attacks with or without panic disorder), a highly treatable condition, is the most prevalent condition associated with unexplained chest pain in the Emergency Department. Panic-like anxiety may be responsible for a significant portion of the negative consequences of unexplained chest pain, such as functional limitations and chronicity. However, more than 92% of panic-like anxiety cases remain undiagnosed at the time of discharge from the Emergency Department.

The 4-item *Panic Screening Score* questionnaire was derived in order to increase the identification of panic-like anxiety in Emergency Department patients with unexplained chest pain.

### Methods and analysis

The goals of this prospective cohort study are to 1) refine the *Panic Screening Score*; 2) validate the revised version of the *Panic Screening Score*; 3) measure the reliability of the revised version of the *Panic Screening Score*; and 4) assess the acceptability of the instrument among emergency physicians. Eligible and consenting patients will be administered the *Panic Screening Score* in a large Emergency Department. Patients will be contacted by phone for administration of the criterion standard for panic attacks as well as by a standardized interview to collect information for other predictors of panic attacks. Multivariate analysis will be used to refine the *Panic Screening Score*. The new version will be prospectively validated in an independent sample and inter-rater agreement will be assessed in 10% of cases. The screening instrument acceptability will be assessed with the *Ottawa Acceptability of Decision Rules Instrument* (OADRI).

## Ethics and dissemination

This study protocol has been reviewed and approved by the Alphonse–Desjardins research ethics committee. The results of the study will be presented in scientific conferences and published in peer-reviewed scientific journals. Further dissemination via workshops and a dedicated website is planned.

## ARTICLE SUMMARY

### Article focus

- This research protocol aims to refine and validate a brief screening instrument for panic attacks in patients with unexplained chest pain.

### Key messages

- The resulting clinical instrument will help emergency physicians determine the likelihood of panic attacks, in turn facilitating appropriate referrals to mental health professionals or family physicians for confirmation of the diagnosis and treatment.
- This project will likely contribute to improve care for patients with unexplained chest pain in the emergency department.

### Strengths and limitations of this study

- This study protocol is based on well-established methodological standards.
- While great efforts were made to insure inclusion of the most relevant predictors of panic-like anxiety, it is possible that some unidentified but relevant variables were missed.

## INTRODUCTION

### Chest pain in Emergency Medicine

Chest pain accounts for approximately 5% of all Emergency Department (ED) consultations,[1] and over 50% of cases remain unexplained at discharge.[2-6] In Canada, approximately 400,000 patients per year present in the ED with unexplained chest pain (UCP).[2, 4]

### The burden of UCP

Despite a generally favorable prognosis, 80% of cases of UCP persist for up to 12 years after initial medical evaluation.[7-13] Many patients with UCP (41%-60%) report limitations in daily functioning (e.g., housework, walking, and exercising) and work absenteeism or disability (17-35 %).[7, 8, 10-12, 14-22] Moreover, the occupational impairments associated with UCP are comparable or more severe than those associated with cardiac chest pain.[7, 22] The negative impact of UCP on quality of life and day-to-day functioning is considerable and may be observed for up to 10 years after symptom onset.[7-9, 11, 12, 14-22]

Despite the benign origin of their pain, patients with UCP report persistent fear of serious health conditions.[11, 20, 22, 23] They are frequent users of health care services, including emergency care, and often undergo multiple invasive tests (e.g., coronary angiograms).[9, 14, 22, 24-27] In Canada, the average duration of an ED consultation for a UCP patient is 11 hours and one in four patients arrives in the ED by ambulance.[3] The direct annual cost associated with UCP in the United States is estimated to amount to eight billion US dollars.[28, 29]

UCP is also associated with significant psychological distress that can become

1  
2  
3 chronic in the absence of targeted interventions.[30-32] In fact, 20% to 40% of patients  
4 present a psychiatric disorder at the time of ED consultation [3, 6, 33-35] and 15% report  
5 suicidal ideation.[3, 33, 34, 36] Unfortunately, fewer than 5% of patients are referred to a  
6 mental health professional for psychiatric or psychological treatment.[3]  
7  
8  
9  
10  
11

12 While the cause of UCP may be unclear, the literature clearly demonstrates that  
13 UCP is highly prevalent and often chronic, and that it constitutes a significant burden for  
14 patients and society alike.  
15  
16  
17  
18

### 19 **Etiology of UCP**

20 Although pathologies such as micro-vascular angina and gastroesophageal reflux  
21 may be at the origin of some cases of UCP,[29] panic attacks are the most prevalent  
22 condition associated with UCP in the ED.[3, 34, 35, 37-39] As many as 44% of patients  
23 with UCP experience panic attacks in the month prior to ED consultation.[3, 34, 35, 37-  
24 39]  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 A panic attack is defined as a discrete period of intense fear or discomfort that  
35 peaks in a few minutes.[40, 41] Fear or discomfort is accompanied by at least four of the  
36 following symptoms: chest pain, palpitations, dyspnea, a feeling of suffocation, hot or  
37 cold flashes, sweating, nausea, feeling faint, paresthesia, trembling, fear of death,  
38 depersonalization, and fear of losing control or going crazy.[40, 41] Panic attacks may be  
39 an isolated phenomenon or may occur in the context of a psychiatric disorder; the most  
40 common psychiatric disorder in which panic attacks occur is panic disorder.[42] The one-  
41 year prevalence of panic attacks in the general adult population is 8%-11%[42, 43] the  
42 prevalence is four to six times higher among patients presenting UCP.[3, 42, 43]  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The literature clearly demonstrates that panic attacks with and without panic disorder constitute a significant mental health problem with serious consequences.[3, 36, 42-47] For simplicity, the term *panic-like anxiety* (PLA) will be used to refer to panic attacks with or without panic disorder.

### **Consequences of PLA in patients with UCP**

PLA may be responsible for a significant portion of the negative consequences of UCP.[3, 6, 15, 30-32, 34, 35, 37] PLA is associated with a greater frequency of UCP episodes and increased risk of chronicity.[15, 30-32] Quality of life is lower and functional limitations levels are higher in patients with UCP and PLA.[15, 30, 31, 37] Moreover, in patients with UCP, PLA is associated with at least a threefold increase in psychiatric morbidity and suicidal ideation.[3, 34] Similarly, use of medical resources nearly doubles when PLA is present.[31, 32]

In patients with UCP, PLA is associated with elevated morbidity, excessive health services use, and a negative prognosis. Unfortunately, more than 92% of cases of PLA remain undiagnosed at the time of discharge from the ED.[3, 34, 35]

### **Identifying PLA in patients with UCP**

Several factors may contribute to the current low rate of PLA identification in patients in the ED. First, PLA patients and physicians alike tend to focus on physical symptoms and on potential organic causes.[48] Second, the identification of PLA is complicated by the similarity between PLA symptoms and symptoms of medical conditions such as coronary artery disease. Third, the limited time available for clinical evaluation in ED settings may be insufficient to identify psychological causes of symptoms.[49] Finally, some ED physicians are unfamiliar with PLA or believe that it is

1  
2  
3 not their role to identify psychiatric problems.[50] However, other physicians recognize  
4 the importance of improving identification and treatment of PLA in the ED settings.[51-  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

not their role to identify psychiatric problems.[50] However, other physicians recognize the importance of improving identification and treatment of PLA in the ED settings.[51-53]

Researchers and clinicians seeking methods for increasing PLA identification rates must take into consideration certain constraints related to the clinical practice of Emergency Medicine, notably the brief period of time available to assess patients.

### **The importance of screening for PLA in ED patients with UCP**

Increasing the rate of identification of a problem is not in itself sufficient to improve clinical outcomes for patients.[54, 55] Gates [54] and Stiell and Wells [55] propose five criteria for determining the importance of a detection procedure and its potential impact on patients' clinical outcomes: 1) The problem has an impact on public health; 2) the problem is sufficiently prevalent; 3) effective treatments are available to reduce morbidity; 4) early diagnosis improves patient prognosis; and 5) additional investigations or treatments are acceptable to patients.

The current data demonstrate that more accurate identification of PLA in ED patients with UCP could improve clinical outcomes. First, PLA in patients with UCP is a prevalent health problem with serious consequences for patients and society. Second, research demonstrates that morbidity associated with PLA in patients suffering from UCP can be greatly reduced via evidence-based treatments.[56-59] For example, 80%-95% of patients with PLA show significant improvement and attain an adequate level of functioning following cognitive-behavioral therapy.[58, 60, 61] Several evidence-based treatment methods for PLA have proven to be effective in patients with UCP.[38, 62-64] Third, given that PLA tends to worsen over time,[32, 42, 65-68] negatively influencing



1  
2  
3 treatment response, early diagnosis improves prognosis.[25, 42, 65, 67, 69-72] Finally,  
4 the criterion of acceptability to patients appears to be met. Participation rates for patients  
5 with PLA and UCP approached for inclusion in a study are generally over 70%.[38, 62-  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

care.[73]

The development of interventions designed to improve identification of PLA associated with UCP in the ED appears to be indicated. A central factor for such an intervention is the availability of a suitable screening instrument, that is, an instrument that is efficient and acceptable to emergency physicians.[54, 55] The use of decision aids such as screening instruments is recognized as an effective method for improving clinical decision-making.[55]

### **The *Panic Screening Score***

To our knowledge, our team has developed the only two screening instruments for PLA in ED patients with UCP.[74, 75] One instrument, the *Panic Screening Score* (PSS) (Figure 1) was designed for use with patients with UCP and identifies panic attacks with and without panic disorder.[74] In addition, the PSS has the advantage of being brief (four items) and easy to use. We have shown that the PSS is eight times more sensitive in detecting PLA associated with UCP than is clinical evaluation by an emergency physician.[3, 74] In addition, the PSS offers a good combination of sensitivity and specificity (Table 1) and these properties have been shown to be stable in a retrospective validation and preliminary prospective validation.[74, 76] To date, the PSS is the briefest and most effective screening instrument for PLA associated with UCP in emergency

settings. Although the PSS has good specificity, its sensitivity needs to be improved (Table 1).

**Table 1 Predictive validity of the PSS [74]**

	<b>Derivation</b> ( <i>n</i> = 201)	<b>Retrospective validation</b> ( <i>n</i> = 305)
Sensitivity	63% (95% CI, 52-73%)	53% (95% CI, 44-62%)
Specificity	84% (95% CI, 76-90%)	85% (95% CI, 78-89%)
Positive predictive value	74% (95% CI, 62-83%)	72% (95% CI, 62-80%)
Negative predictive value	76% (95% CI, 68-89%)	71% (95% CI, 65-77%)
Positive likelihood ratio	3.89 (95% CI, 2.5-6.05)	3.45 (95% CI, 2.35-5.04)
Negative likelihood ratio	0.44 (95% CI, 0.33-0.59)	0.55 (95% CI, 0.46-0.67)

Note. CI = Confidence Interval

### Summary

Early identification of PLA in ED patients with UCP appears to be the strategy of choice for reducing morbidity, chronicity, and overuse of health care services. The PSS is a concise and effective instrument that represents the most promising method for achieving this objective. The present study will represent a major step toward the clinical application of the PSS and the early diagnosis and treatment of PLA in ED patients with UCP.

## METHODS AND ANALYSIS

### Objectives

The objectives of this prospective cohort study are to: 1) refine the PSS; 2) validate the revised version of the PSS in an independent sample; 3) estimate the reliability of the revised PSS; and 4) assess the acceptability of the instrument among ED physicians.

## Methodological framework

The research methods and statistical analyses used in this study are based on clinical decision rule standards:[55]

- 1) The outcome must be clearly defined and assessed blindly;
- 2) The predictors must be clearly defined, standardized, and evaluated without knowledge of patient status;
- 3) The reliability of the variables studied must be demonstrated;
- 4) Participants must be selected without bias and must represent a large range of clinical and demographic characteristics. Ideally, the study should be conducted in several centers in order to increase external validity;
- 5) Appropriate statistical analyses must be used;
- 6) Sample size must be sufficient to permit valid statistical analyses;
- 7) The sensibility of the instrument must be adequate. It must have a clear objective, good content validity, and clinical relevance, and must be easy to use in the context of targeted practice;
- 8) The capacity of the instrument to identify patients with (sensitivity) and without (specificity) the condition must be demonstrated;
- 9) Measures must be taken to guarantee the appropriate application of the instrument.

The procedures for refining the PSS (objective 1) are based on Stiell and Wells' recommendations.[55] First, the PLA predictors that were valuable but not essential in previous studies will be reassessed.[74] Next, potential predictors of PLA that were not assessed at the time of the initial study will be evaluated.

## Participants

This study will consecutively recruit 3000 patients at the two emergency services of the Centre de santé et de services sociaux Alphonse-Desjardins (University-Affiliated Hospital of Lévis and Paul-Gilbert Hospital). To be eligible, patients will have to present UCP as defined by: 1) absence of an identifiable cause (e.g., pneumothorax, pneumonia); 2) absence of chest trauma; 3) absence of new malignant cardiac arrhythmia; and 4) a score of 2 or less on the modified version of the *Thrombolysis in myocardial infarction score* (TIMI score).[77, 78] This simple instrument stratifies patients presenting chest pain according to the probability of mortality or myocardial infarction in the 30 days following the ED visit.[77-79] A result  $\leq 2$  is associated with low incidence (3%) of mortality or cardiac events; scores  $> 2$  are associated with a higher incidence (28%). Scores are obtained by summing values that correspond to the following characteristics: 1) age  $\geq 65$  years (1 point); 2) known coronary stenosis  $\geq 50\%$  or history of revascularization (1 point); 3) deviation of the ST segment  $\geq 0.5$  mm (5 points); and 4) elevated rate of cardiac enzymes defined as troponins I  $\geq 99^{\text{th}}$  percentile.[78] A value of zero is assigned for criteria 3 and 4 if the physician does not order the tests necessary to obtain the information.

Patients will be excluded if they: 1) present a terminal illness; 2) present a severe communication problem that could interfere with the administration of the questionnaire; 3) present a psychotic state, major cognitive deficit, or other condition that could invalidate the interview; and 4) are legally incompetent or younger than 18 years old.

## Procedure

With the assistance of a research nurse, the emergency physicians will assess the eligibility of all patients presenting UCP. Physicians will complete the PSS for every eligible and consenting patient. To assess the reliability of PSS items, a second physician will independently

complete the PSS for at least 10% of patients. As in other similar studies and due to constraints such as the availability of a second physician, the reliability assessment will be conducted using a convenience sample.[80-82]

To assess primary outcome and potential predictors of PLA, all participants will complete a telephone interview within 72 hours following recruitment. Interviewers will be blind to patient PSS scores. Since the evaluation of the criterion standard will occur after patient discharge, physicians will be blind to the results of the PLA evaluations.

The research nurse will review the ED computerized database every day to ensure that all potentially eligible patients were assessed. A registered nurse will contact patients who were missed and request consent for the telephone interview. For non-consenting patients, only age, gender, and time of ED visit will be recorded. This step will enable us to identify potential selection biases and ensure quality control.

The acceptability of the PSS to emergency physicians will be assessed at the end of the study. The time taken to administer the PSS will also be evaluated in 10% of cases randomly selected. The initiation of the questionnaire will be defined as the moment the physician asks the first question; the end of the questionnaire will be defined as the moment the patient finishes answering the final question. This measure will be used to assess whether or not the PSS is sufficiently brief for the ED.

### Measures

- Eligibility evaluation form: This form contains the inclusion and exclusion criteria. It also records patients contact information, age, gender and time of ED visit.
- The Panic Screening Score and the Panic Screening Score – revised version: PSS score is calculated by summing the points assigned to the answer for each of the four

1  
2  
3 questions. A score  $\geq 6$  is considered to be a positive result: the patient presents an  
4 elevated probability of PLA. A revised version of the PSS will be administered to  
5 patients during the validation phase (objective 2). The revised version will include  
6 only the items selected during the optimization phase.  
7  
8  
9

- 10  
11  
12  
13 • Anxiety Disorders Interview Schedule for Diagnostic and Statistical Manual of  
14 Mental Disorders, Fourth Edition (ADIS-IV):[83, 84] The Panic disorder module of  
15 the French (or English) version of the interview will be administered by telephone.  
16 As recommended by experts in the field, the ADIS-IV will serve as the criterion  
17 standard for the identification of PLA.[85] The Panic disorder module has shown  
18 excellent reliability for the identification of PLA ( $k \geq 0.80$ ).[34] PLA is defined as  
19 either the presence of a panic attack during the previous month or the presence of  
20 panic disorder.[74]  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32 Additional predictors for the refinement phase (objective 1): Additional predictors  
33 include the four potential predictors of PLA identified but not included in the final version of the  
34 PSS,[74] as well as the items selected following a pilot study [76] and a review of the literature:  
35  
36  
37

- 38 - Fear of dying associated with chest pain;[86, 87]
- 39 - Item number 4 (fear of fainting) of the Anxiety Sensitivity Index;[88, 89]
- 40 - Item number 28 (feeling overwhelmed by one's problems) on the State-Trait  
41 Anxiety Inventory;[90, 91]
- 42 - The *Autonomic Nervous System Questionnaire*:[92]
- 43 - The four panic disorder items from the Patient Health Questionnaire-15;[93]

44 these items were selected because they demonstrated sensitivity and  
45  
46  
47  
48  
49  
50  
51  
52

specificity between 75% and 96% for PLA in patients in primary care and psychosomatic settings;[94]

- The modified version of the *Life Events Stress Scale*. [95, 96] This scale has excellent internal consistency (0.87). The questionnaire includes 10 items, each of which corresponds to a category of stressful events. Patients are asked to specify whether or not each event has occurred and, if so, whether or not it occurred within the previous six months. For the purpose of this study, patients will be asked to specify whether or not the event occurred in the last month. The intensity of the stress associated with each event will be evaluated on a five-point Likert scale. This questionnaire was selected because the occurrence or aggravation of PLA is preceded by stressful life events in 80% of cases. [97]

- Modified version of the *Ottawa Acceptability of Decision Rules Instrument* (OADRI): [98] This 12-item questionnaire assesses the acceptability of clinical decision rules by physicians. The questionnaire has good internal consistency (0.80 to 0.86) and good construct validity. [98] Ten items will be used in the present study as some information will not be available to ED physicians at the time of assessment. The first excluded item concerns the instrument's validation data; the second excluded item concerns the impact of the instrument on the use of clinical resources.
- PSS administration time assessment record sheet: This document includes instructions for assessing the PSS administration time and recording the result.

## Quality control

Emergency physicians will receive a 30-minute training session on how to use the PSS. The session will include an overview of the study, inclusion and exclusion criteria, and guidelines for the administration of the PSS and for scoring patients' answers. The training session will be developed jointly with the Centre de liaison sur l'intervention et la prévention psychosociale (CLIPP). This non-profit organization is specialized in knowledge transfer and dissemination of research results.

Over a one-month period, a research nurse will periodically observe each physician as he or she administers the PSS, in order to obtain feedback and to identify problems. This step will be repeated during the implementation of the refined version of the PSS.

Telephone interviews will be conducted by graduate students in psychology. Each student will receive one day of training on the administration of the ADIS-IV, followed by weekly supervision with a clinical psychologist. Telephone interviews will be recorded to facilitate supervision. A random sample of 25% of recorded interviews will be used to evaluate inter-rater agreement on the diagnosis of PLA. During the optimization phase (objective 1), the recordings will also be used to evaluate inter-rater agreement on each of the additional interview items. These supervision and inter-rater agreement procedures have been proven effective in our previous studies and generated excellent diagnostic reliability for PLA.[3, 34]

## Data analysis

Participants' sociodemographic data will be presented in descriptive form. To evaluate the representativeness of the sample, participant data will be compared with data of the eligible patients who declined to participate. Continuous variables that meet the assumptions of normality will be analysed with Student's *t* test. Otherwise, Wilcoxon Mann-Whitney's



1  
2  
3 nonparametric test will be used. The Chi-square test will be used for categorical variables.  
4  
5 Inter-rater agreement for PLA diagnosis on the ADIS-IV [83, 84] will be assessed with  
6  
7  
8 Cohen's kappa coefficient.  
9

10  
11 Because the predictive performance of a clinical rule is usually overestimated in the  
12  
13 sample used in its development, it is important to evaluate the rule in an independent  
14  
15 sample.[55, 99] In this study, a temporal validation procedure will be used. The refinement  
16  
17 analyses (objective 1) will be conducted using data from the first 1500 patients; the validation  
18  
19 of the refined version of the PSS (objective 2) will be performed using data from the  
20  
21 subsequent 1500 patients.  
22  
23

#### 24 Refinement of the Panic Screening Score (objective 1)

25  
26 The reliability of the four PSS items and the 15 potential predictors will be evaluated  
27  
28 using Cohen's kappa tests or weighted kappa. Only the items with a good Kappa's coefficient ( $k$   
29  
30  $\geq 0.6$ ) will be included in further analysis.[55] We will use the two types of analysis  
31  
32 recommended by experts in the field, recursive partitioning and regression,[55] to refine the PSS  
33  
34 data collected from the first 1500 patients. Recursive partitioning generally results in a more  
35  
36 sensitive instrument, whereas regression yields models with a higher global predictive value.[55,  
37  
38  
39  
40  
41 100]  
42

43  
44 This study will use the recursive partitioning technique known as Classification and  
45  
46 Regression Trees.[101] The construction of Classification and Regression Trees will be  
47  
48 automated, but manual intervention will be used if some of the concurrent predictors are more  
49  
50 useful than others (e.g., more reliable, more representative or easier to use). We will  
51  
52 dichotomize continuous variables by selecting the most effective cut-off point for identifying  
53  
54 patients with PLA.  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

Log-binomial regression analysis will also be performed. This type of analysis is preferred to logistic regression because it provides exact relative risks rather than odds ratios. Furthermore, when the prevalence of the dependent variable is greater than 10%, the estimate of the relative risk by logistic regression is imprecise.[102] In this study, the expected prevalence of PLA is 44%.[3] Variables associated with PLA in univariate log-binomial regression ( $p \leq 0.15$ ) will be considered in the multivariate analysis. The multivariate log-binomial regression will be performed using the ascending stepwise method. Multi-collinearity between variables will be verified. If correlations  $\geq 0.80$  are obtained, the analysis will be repeated using only one of the inter-correlated items, in order to obtain the most effective model. The regression equation will be converted into a score by assigning points to each answer; point assignment will be based on the magnitude of corresponding regression coefficients according to the *Framingham study risk score function*. [103] The result is a simple score that provides probability estimates that correspond to the score generated by the multivariate regression model. The cut-off score that yields the best predictive validity will be selected based on the area underneath the receiver operating characteristic (ROC) curve and the measures of predictive validity.

41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

The calibration (Hosmer-Lemeshow test) and discriminating validity (area under the ROC curve, sensitivity, specificity, likelihood ratios, predictive values) of the two optimized versions of the PSS will be evaluated. Confidence intervals of 95% will be calculated for each of the discrimination measures. The version of the PSS that is simplest (i.e., the version with the fewest items) and offers the best discrimination will be evaluated (objective 2).

53  
54  
55  
56  
57  
58  
59  
60

Validation of the refined PSS (objective 2)

1  
2  
3 The optimized version of the PSS will be prospectively validated in a validation sample  
4  
5 ( $n = 1500$ ). Sensitivity, specificity, predictive values, and likelihood ratios will be calculated  
6  
7 with 95% confidence intervals.  
8  
9

#### 10 Evaluation of PSS reliability

11  
12 Cohen's Kappa coefficient will be used to assess the level of inter-rater agreement  
13  
14 (reliability) for the refined PSS (presence or absence of PLA) and for each of its components. In  
15  
16 the case of variables including three or more categories, a weighted Kappa coefficient will be  
17  
18 calculated.[74, 104]  
19  
20

#### 21 Acceptability of the PSS

22  
23 Descriptive data on PSS administration time will be reported, including the mean,  
24  
25 median and range. The total score on the OADRI and the level of endorsement for each item will  
26  
27 also be reported in descriptive form.  
28  
29  
30

#### 31 **Justification of the sample size and feasibility**

32  
33 Based on our previous study, we estimate the minimum prevalence of PLA in these  
34  
35 settings to be 40%.[3] The sample composed of the first 1500 patients will be used for analyses  
36  
37 related to the refinement of the PSS. It will include approximately 600 patients with PLA. This  
38  
39 number exceeds the minimum ratio of 10 cases for each variable in the regression analysis.[105]  
40  
41 Each subsample of 1500 patients will allow us to obtain a confidence interval of  $95\% \pm 3.9\%$ ,  
42  
43 for a level of sensitivity equivalent to that reported in the original PSS study (63%).  
44  
45  
46  
47

48 The EDs at the Centre de santé et de services sociaux Alphonse-Desjardins receive  
49  
50 approximately 110,000 patients each year, and approximately 2% of patients visiting the two  
51  
52 EDs present UCP. Of 2200 eligible patients per year, we expect that 10% will be missed and  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 20% will decline to participate. Our final recruitment estimate is therefore 3000 participants in  
4  
5  
6 24 months (1500 per year).  
7  
8  
9

## 10 **ETHICS AND DISSEMINATION**

11  
12  
13 The research ethics committee at the Centre de santé et de services sociaux  
14  
15 Alphonse-Desjardins approved this protocol. The study will not affect usual care and the  
16  
17 ethical considerations are minimal. Patients' verbal consent to complete the PSS and to be  
18  
19 contacted by telephone will be solicited. Verbal consent will be obtained again at the time  
20  
21 of the telephone interview. All data will be treated according to standard guidelines for  
22  
23 ensuring patient confidentiality.  
24  
25  
26

27  
28 The results of the study will be presented in scientific conferences and published in  
29  
30 peer-reviewed scientific journals. Further dissemination via workshops aimed at emergency  
31  
32 physicians in clinical settings and a dedicated website is planned.  
33  
34  
35

## 36 **DISCUSSION AND CONCLUSIONS**

37  
38  
39 This study is designed to validate an effective and efficient screening instrument for  
40  
41 PLA in ED patients with UCP. The PSS will help emergency physicians determine the  
42  
43 likelihood of PLA, in turn facilitating appropriate referrals to mental health professionals or  
44  
45 family physicians for confirmation of the diagnosis and treatment. Treatment for PLA  
46  
47 significantly reduces associated morbidity and excessive use of health care services, and has  
48  
49 an overall favorable cost/benefit ratio.[26, 38, 62-64, 106-108] This study will result in a  
50  
51 screening tool with the potential to have a tangible clinical impact for ED patients with UCP  
52  
53 and PLA. Further research will focus on assessing the impact of use of the PSS and on  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 validating the instrument in other settings, such as cardiology and primary care clinics.[25,  
4  
5 31]  
6  
7

### 8 **Acknowledgements**

9  
10 The authors wish to thank Stéphanie Hamel for her help in preparing this manuscript  
11 and Sarah Roberts for her help in editing this manuscript.  
12  
13

### 14 **Competing interests**

15  
16 The authors declare that they have no competing interests.  
17  
18

### 19 **Funding**

20  
21 This work was supported by the Canadian Institutes of Health Research (CIHR),  
22 grant number: 126125.  
23  
24

### 25 **Contributorship**

26  
27 GFB was responsible for the original idea, literature review and study design. ID helped draft  
28 the manuscript, assisted with the methodology and revised the manuscript for important  
29 intellectual content. PA and JP revised the manuscript for important intellectual content. RPF  
30 assisted with the methodology and revised the manuscript for important intellectual content.  
31  
32 CED assisted with the statistical design and methodology and revised the manuscript  
33 critically for important intellectual content. All authors read and approved the final  
34 manuscript.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

### 45 **Figure legend**

46  
47  
48 Figure 1 – The Panic Screening Score (PSS)  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

- 1 McCaig LF, Burt CW. National Hospital Ambulatory Medical Care Survey: 2002 Emergency Department Summary. *Adv Data* 2004;**340**:1-34.
- 2 Christenson J, Innes G, Mcknight D, *et al.* Safety and efficiency of emergency department assessment of chest discomfort. *CMAJ* 2004;**170**:1803-7.
- 3 Foldes-Busque G, Marchand A, Chauny JM, *et al.* Unexplained chest pain in the ED: could it be panic? *Am J Emerg Med* 2011;**29**:743-51.
- 4 Karlson BW, Herlitz J, Pettersin P. Patients admitted to the emergency room with symptoms indicative of acute myocardial infarction. *J Intern Med* 1991;**230**:251-8.
- 5 Kontos MC, Jesse RL. Evaluation of the emergency department chest pain patient. *Am J Cardiol* 2000;**85**:32B-9B.

- 1  
2  
3  
4 6 Srinivasan K, Joseph W. A study of lifetime prevalence of anxiety and depressive  
5 disorders in patients presenting with chest pain to emergency medicine. *Gen Hosp*  
6 *Psychiatry* 2004;**26**:470-4.  
7  
8  
9  
10 7 Karlson BW, Wiklund I, Bengtson A, *et al.* Prognosis, severity of symptoms, and  
11 aspects of well-being among patients in whom myocardial infarction was ruled  
12 out. *Clin Cardiol* 1994;**17**:427-31.  
13  
14  
15 8 Papanicolaou MN, Califf RM, Hlatky MA, *et al.* Prognostic implications of  
16 angiographically normal and insignificantly narrowed coronary arteries. *Am J*  
17 *Cardiol* 1986;**56**:1181-7.  
18  
19  
20 9 Eslick GD, Talley NJ. Natural history and predictors of outcome for non-cardiac  
21 chest pain: a prospective 4-year cohort study. *Neurogastroenterol Motil*  
22 2008;**20**:989-97.  
23  
24  
25 10 Rolls M, Kollind M, Theorel T. Five-year follow-up of young adults visiting an  
26 emergency unit because of atypical chest pain. *J Intern Med* 1992;**231**:59-65.  
27  
28  
29 11 Dumville JC, MacPherson H, Griffith K, *et al.* Non-cardiac chest pain: a  
30 retrospective cohort study of patients who attended a Rapid Access Chest Pain  
31 Clinic. *Fam Pract* 2007;**24**:152-7.  
32  
33  
34 12 Herlitz J, Karlson BW, Wiklund I, *et al.* Prognosis and gender differences in chest  
35 pain patients discharged from an ED. *Am J Emerg Med* 1995;**13**:127-32.  
36  
37  
38 13 Lichtlen PR, Bargheer F, Wenzlaff P. Long-term prognosis of patients with  
39 anginalike chest pain and normal coronary angiographic findings. *J Am Coll*  
40 *Cardiol* 1995;**25**:1013-8.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 14 Eslick GD, Talley NJ. Non-cardiac chest pain: predictors of health care seeking,  
5 the types of health care professional consulted, work absenteeism and interruption  
6 of daily activities. *Aliment Pharmacol Ther* 2004;**20**:909-15.  
7  
8  
9  
10  
11 15 Dammen T, Ekeberg O, Arnesen H, *et al.* Health-related quality of life in non-  
12 cardiac chest pain patients with and without panic disorder. *Int J Psychiatry Med*  
13 2008;**38**:271-86.  
14  
15  
16  
17 16 Jerlock M, Kjellgren KI, Gaston-Johanson F, *et al.* Psychosocial profile in men  
18 and women with unexplained chest pain. *J Intern Med* 2008;**264**:265-74.  
19  
20  
21  
22 17 Jerlock M, Gaston-Johanson F, Danielson E. Living with unexplained chest pain.  
23 *J Clin Nurs* 2005;**14**:956-64.  
24  
25  
26  
27 18 Fagring A, Kjellgren KI, Rosengren A, *et al.* Depression, anxiety, stress, social  
28 interaction and health-related quality of life in men and women with unexplained  
29 chest pain. *BMC Public Health* 2008;**8**:165.  
30  
31  
32  
33  
34 19 Wong WM, Lam KF, Cheng C, *et al.* Population based study of noncardiac chest  
35 pain in southern Chinese: prevalence, psychosocial factors and health care  
36 utilization. *World J Gastroenterol* 2004;**10**:707-12.  
37  
38  
39  
40  
41 20 Fagring A, Gaston-Johanson F, Kjellgren KI, *et al.* Unexplained chest pain in  
42 relation to psychosocial factors and health-related quality of life in men and  
43 women. *Eur J Cardiovasc Nurs* 2007;**6**:329-36.  
44  
45  
46  
47  
48 21 Rolls M, Kollind M, Theorel T. Five years follow-up of young adults visiting an  
49 emergency unit because of atypical chest pain. *J Intern Med* 1992;**231**:59-65.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 22 Potts SG, Bass CM. Psychosocial outcome and use of medical resources in patients with chest pain and normal or near-normal coronary arteries: a long-term follow-up study. *Q J Med* 1993;**86**:583-93.
- 23 Fagring AJ, Gaston-Johansson F, Danielson E. Description of unexplained chest pain and its influence on daily life in men and women. *Eur J Cardiovasc Nurs* 2005;**4**:337-44.
- 24 Coley K, Saul MI, Seybert AL. Economic burden of not recognizing panic disorder in the emergency department. *J Emerg Med* 2007;**36**:3-7.
- 25 Carter C, Maddock R, Zoglio M, *et al.* Panic disorder and chest pain: a study of cardiac stress scintigraphy patients. *Am J Cardiol* 1994;**74**:296-8.
- 26 Abbass A, Campbell S, Magee K, *et al.* Intensive short-term dynamic psychotherapy to reduce rates of emergency department return visits for patients with medically unexplained symptoms: preliminary evidence from a pre-post intervention study. *CJEM* 2009;**11**:529-34.
- 27 Katerndahl DA. Factors associated with persons with panic attacks seeking medical care. *Fam Med* 1990;**22**:462-6.
- 28 Kahn S. The challenge of evaluating the patient with chest pain. *Arch Pathol Lab Med* 2000;**124**:1418-9.
- 29 Eslick GD, Coulshed DS, Talley NJ. Review article: the burden of illness of non-cardiac chest pain. *Aliment Pharmacol Ther* 2002;**16**:1217-23.
- 30 Bringager CB, Arnesen H, Friis S, *et al.* A long term follow-up study of chest pain patients: effect of panic disorder on mortality, morbidity and quality of life. *Cardiology* 2008;**110**:8-14.

- 1  
2  
3  
4 31 Dammen T, Bringager CB, Arnesen H, *et al.* A 1-year follow-up study of chest-  
5 pain patients with and without panic disorder. *Gen Hosp Psychiatry* 2006;**28**:516-  
6 24.  
7  
8  
9  
10 32 Fleet RP, Lavoie KL, Martel JP, *et al.* Two-year follow-up status of emergency  
11 department patients with chest pain: was it panic disorder? *CJEM* 2003;**5**:247-54.  
12  
13 33 Fleet RP, Dupuis G, Kaczorowski J, *et al.* Suicidal ideation in emergency  
14 department chest pain patients: panic disorder a risk factor. *Am J Emerg Med*  
15 1997;**15**:345-9.  
16  
17  
18 34 Fleet RP, Dupuis G, Marchand A, *et al.* Panic disorder in emergency department  
19 chest pain patients: prevalence, comorbidity, suicidal ideation, and physician  
20 recognition. *Am J Med* 1996;**101**:371-80.  
21  
22  
23 35 Yingling KW, Wulsin LR, Arnold LM, *et al.* Estimated prevalences of panic  
24 disorder and depression among consecutive patients seen in an emergency  
25 department with acute chest pain. *J Gen Intern Med* 1993;**8**:231-5.  
26  
27  
28 36 Foldes-Busque G, Fleet RP, Poitras J, *et al.* Suicidality and panic in emergency  
29 department patients with unexplained chest pain. *Gen Hosp Psychiatry*  
30 2012;**34**:178-84.  
31  
32  
33 37 Worthington JJ, Pollack MH, Otto MW, *et al.* Panic disorder in emergency ward  
34 patients with chest pain. *J Nerv Ment Dis* 1997;**185**:274-6.  
35  
36  
37 38 Wulsin L, Liu T, Storrow A, *et al.* A randomized controlled trial of panic disorder  
38 treatment initiation in an emergency department chest pain center. *Ann Emerg*  
39 *Med* 2002;**39**:139-43.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 39 Wulsin LR, Hillard JR, Geier P, *et al.* Screening emergency room patients with  
5 atypical chest pain for depression and panic disorder. *Int J Psychiatry Med*  
6  
7 1988;**18**:315-23.  
8  
9  
10 40 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental*  
11  
12 *Disorders, Fourth Edition, Text Revised.* Washington, DC: American Psychiatric  
13 Association 2000.  
14  
15  
16 41 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental*  
17  
18 *Disorders, Fifth Edition: DSM-5.* Washington, DC: American Psychiatric  
19 Association 2013.  
20  
21  
22 42 Kessler RC, Chiu WT, Jin R, *et al.* The epidemiology of panic attacks, panic  
23  
24 disorder, and agoraphobia in the National Comorbidity Survey Replication. *Arch*  
25  
26 *Gen Psychiatry* 2006;**63**:415-24.  
27  
28  
29 43 Ramage-Morin PL. Trouble panique et comportements d'adaptation. *Suppléments*  
30  
31 *aux Rapports sur la Santé* 2004;**15**:37-72.  
32  
33  
34 44 Baillie AJ, Rapee RM. Panic attacks as risk markers for mental disorders. *Soc*  
35  
36 *Psychiatry Psychiatr Epidemiol* 2005;**40**:240-4.  
37  
38  
39 45 Batelaan N, De Graaf R, Van Balkom A, *et al.* Thresholds for health and  
40  
41 thresholds for illness: panic disorder versus subthreshold panic disorder. *Psychol*  
42  
43 *Med* 2007;**37**:247-56.  
44  
45  
46 46 Batelaan N, Smit F, de Graaf R, *et al.* Economic costs of full-blown and  
47  
48 subthreshold panic disorder. *J Affect Disord* 2007;**104**:127-36.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 47 Kinley DJ, Cox BJ, Clara I, *et al.* Panic attacks and their relation to psychological  
4 and physical functioning in Canadians: results from a nationally representative  
5  
6  
7  
8 sample. *Can J Psychiatry* 2009;**54**:113-22.  
9
- 10 48 Rubens M. Chronic anxiety: aspects of diagnosis and treatment. *Canadian*  
11  
12 *Journal of CME* 1997;**February**:147-67.  
13
- 14 49 Stephenson DT, Price JR. Medically unexplained physical symptoms in  
15  
16 emergency medicine. *Emerg Med J* 2006;**23**:595-600.  
17
- 18 50 Epstein NL. Chest pain and panic disorder in the ED? *CJEM* 2003;**5**:308.  
19
- 20 51 Pollard CA, Lewis LM. Managing panic attacks in emergency patients. *J Emerg*  
21  
22 *Med* 1989;**7**:547-52.  
23
- 24 52 Lee J, Dade LA. The buck stops where? What is the role of the emergency  
25  
26 physician in managing panic disorder in chest pain patients? *CJEM* 2003;**5**:237-8.  
27
- 28 53 Larkin GL, Beautrais AL, Spirito A, *et al.* Mental health and emergency  
29  
30 medicine: a research agenda. *Acad Emerg Med* 2009;**16**:1110-9.  
31
- 32 54 Gates TJ. Screening for cancer: evaluating the evidence. *Am Fam Physician*  
33  
34 2001;**63**:513-22.  
35
- 36 55 Stiehl IG, Wells GA. Methodologic standards for the development of clinical  
37  
38 decision rules in emergency medicine. *Ann Emerg Med* 1999;**33**:437-47.  
39
- 40 56 American Psychiatric Association. *Practice guideline for the treatment of patients*  
41  
42 *with panic disorder*. In: *Practice Guidelines for the Treatment of Psychiatric*  
43  
44 *Disorders: Compendium 2006*. Arlington, VA: American Psychiatric Association  
45  
46 2006:945-1002.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63
- 57 American Psychiatric Association. *Practice Guideline for the Treatment of Patients with Panic Disorder, Second Edition*. Washington, DC: American Psychiatric Association 2009.
- 58 Foldes-Busque G, Marchand A, Landry P. [Early detection and treatment of panic disorder with or without agoraphobia: update]. *Can Fam Physician* 2007;**53**:1686-93.
- 59 National Collaborating Centre for Mental Health and National Collaborating Centre for Primary Care. *Generalised Anxiety Disorder and Panic Disorder (With or Without Agoraphobia) in Adults. Management in Primary, Secondary and Community Care*. London, UK: National Institute for Health and Clinical Expertise 2011.
- 60 Starcevic V. *Anxiety Disorder in Adults: a Clinical Guide*. New York, NY: Oxford University Press 2005.
- 61 Marchand A, Boivin I. *Le trouble panique*. In: Ladouceur R, Marchand A, Boisvert JM, eds. *Les Troubles Anxieux: Approche Cognitive et Comportementale*. Montréal, QC: Gaétan Morin 1999:59-94.
- 62 Swinson RP, Soulios C, Cox BJ, *et al*. Brief treatment of emergency room patients with panic attacks. *Am J Psychiatry* 1992;**149**:944-6.
- 63 Pelland ME, Marchand A, Lessard MJ, *et al*. Efficacy of 2 interventions for panic disorder in patients presenting to the ED with chest pain. *Am J Emerg Med* 2011;**29**:1051-61.

- 1  
2  
3  
4 64 Lessard MJ, Marchand A, Pelland MÈ, *et al.* Comparing two brief psychological  
5 interventions to usual care in panic disorder patients presenting to the emergency  
6 department with chest pain. *Behav Cogn Psychother* 2012;**40**:129-47.  
7  
8  
9  
10  
11 65 Roy-Byrne PP, Stein MB, Russo J, *et al.* Panic disorder in the primary care  
12 setting: comorbidity, disability, service utilization, and treatment. *J Clin*  
13 *Psychiatry* 1999;**60**:492-9.  
14  
15  
16  
17 66 Swinson RP, Kuch K. *Clinical-features of panic and related disorders.* In:  
18 Ballenger JC.ed. *Clinical Aspects of Panic Disorder.* New York, NY: Wiley-Liss  
19 1990:13-30.  
20  
21  
22  
23  
24 67 Reich J, Warshaw M, Peterson LG, *et al.* Comorbidity of panic and major  
25 depressive disorder. *J Psychiatr Res* 1993;**27**:23-33.  
26  
27  
28  
29 68 Kessler RC, Ustün TB. The World Mental Health (WMH) Survey Initiative  
30 Version of the World Health Organization (WHO) Composite International  
31 Diagnostic Interview (CIDI). *Int J Methods Psychiatr Res* 2004;**13**:93-121.  
32  
33  
34  
35 69 Beitman BD, Kushner MG, Basha I, *et al.* Follow-up status of patients with  
36 angiographically normal coronary arteries and panic disorder. *JAMA*  
37 1991;**265**:1545-9.  
38  
39  
40  
41  
42 70 Brown TA, Barlow DH. Comorbidity among anxiety disorders: implications for  
43 treatment and DSM-IV. *J Consult Clin Psychol* 1992;**60**:835-44.  
44  
45  
46  
47 71 Kessler RC, McGonagle KA, Zhao S, *et al.* Lifetime and 12-month prevalence of  
48 DSM-III-R psychiatric disorders in the United States: results from the National  
49 Comorbidity Survey. *Arch Gen Psychiatry* 1994;**51**:8-19.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 72 Schmidt NB. Panic disorder: cognitive behavioral and pharmacological treatment  
5 strategies. *J Clin Psychol Med Settings* 1999;**6**:89-111.  
6  
7  
8 73 Johnson MR, Gold PB, Siemon L, *et al*. Panic disorder in primary care: patients'  
9 attributions of illness causes and willingness to accept psychiatric treatment. *Int J*  
10 *Psychiatry Med* 2000;**30**:367-84.  
11  
12  
13 74 Foldes-Busque G, Fleet RP, Poitras J, *et al*. Preliminary investigation of the *Panic*  
14 *Screening Score* for emergency department patients with unexplained chest pain.  
15 *Acad Emerg Med* 2011;**18**:322-5.  
16  
17  
18 75 Fleet RP, Dupuis G, Marchand A, *et al*. Detecting panic disorder in emergency  
19 department chest pain patients: a validated model to improve recognition. *Ann*  
20 *Behav Med* 1997;**19**:124-31.  
21  
22  
23 76 Foldes-Busque G, Hamel S, Castonguay J, *et al*. Prospective validation of the  
24 *Panic Screening Score* for emergency department patients with unexplained chest  
25 pain: a preliminary investigation. *Psychosom Med* 2013;**75**:A142.  
26  
27  
28 77 Jaffery Z, Hudson MP, Jacobsen G, *et al*. Modified thrombolysis in myocardial  
29 infarction (TIMI) risk score to risk stratify patients in the emergency department  
30 with possible acute coronary syndrome. *J Thromb Thrombolysis* 2007;**24**:137-44.  
31  
32  
33 78 Hess EP, Perry JJ, Calder LA, *et al*. Prospective validation of a modified  
34 thrombolysis in myocardial infarction risk score in emergency department  
35 patients with chest pain and possible acute coronary syndrome. *Acad Emerg Med*  
36 2010;**17**:368-75.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 79 Chase M, Robey JL, Zogby KE, *et al.* Prospective validation of the thrombolysis  
5  
6 in myocardial infarction risk score in the emergency department chest pain  
7  
8 population. *Ann Emerg Med* 2006;**48**:252-9.  
9
- 10  
11 80 Stiell IG, Clement CM, McKnight RD, *et al.* The Canadian C-spine rule versus  
12  
13 the NEXUS low-risk criteria in patients with trauma. *N Engl J Med*  
14  
15 2003;**349**:2510-8.  
16
- 17  
18 81 Stiell IG. Ottawa ankle rules. *Can Fam Physician* 1996;**42**:478-80.  
19
- 20  
21 82 Stiell IG, Clement CM, Rowe BH, *et al.* Comparison of the Canadian CT Head  
22  
23 Rule and the New Orleans Criteria in patients with minor head injury. *JAMA*  
24  
25 2005;**294**:1511-8.  
26
- 27  
28 83 Brown TA, Di Nardo PA, Barlow DH. *Anxiety Disorders Interview for DSM-IV*  
29  
30 (*ADIS-IV*).USA: Oxford University Press 1994.  
31
- 32  
33 84 Boivin I, Marchand A. *Évaluation structurée pour les troubles anxieux du DSM-*  
34  
35 *IV*. Montréal, QC: Université du Québec à Montréal 1996.  
36
- 37  
38 85 Shear MK, Maser JD. Standardized assessment of panic disorder research. A  
39  
40 conference report. *Arch Gen Psychiatry* 1994;**51**:346-54.  
41
- 42  
43 86 Fraenkel YM, Kindler S, Melmed RN. Differences in cognitions during chest pain  
44  
45 of patients with panic disorder and ischemic heart disease. *Depress Anxiety* 1996-  
46  
47 97;**4**:217-22.  
48
- 49  
50 87 First MB, Spitzer RL, Gibbon M, *et al.* *Structured Clinical Interview for DSM-IV*  
51  
52 *Axis I Disorders-Patient Edition (SCID-I/P, Version 2.0)*. New York, NY: New  
53  
54 York State Psychiatric Institute 1995.  
55  
56  
57  
58  
59  
60



- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 88 Reiss S, Peterson RA, Gursky DM, *et al.* Anxiety sensitivity, anxiety frequency and the prediction of fearfulness. *Behav Res Ther* 1986;**24**:1-8.
- 89 Verreault N, Marchand A, Labrecque J, *et al.* Validation de l'Index de sensibilité à l'anxiété auprès de la population québécoise francophone. *Revue Québécoise de Psychologie* 2007;**28**:253-68.
- 90 Gauthier J, Bouchard S. Adaptation canadienne-française de la forme révisée de State-Trait Anxiety Inventory de Spielberger. *Can J Behav Sci* 1993;**25**:559-78.
- 91 Spielberger CD. *Manual for the State-Trait Anxiety Inventory STAI-form Y*. Palo Alto, CA: Consulting Psychologists Press Inc 1983.
- 92 Stein MB, Roy-Byrne PP, McQuaid JR, *et al.* Development of a brief diagnostic screen for panic disorder in primary care. *Psychosom Med* 1999;**61**:359-64.
- 93 Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *JAMA* 1999;**282**:1737-44.
- 94 Löwe B, Gräfe K, Zipfel S, *et al.* Detecting panic disorder in medical and psychosomatic outpatients: comparative validation of the Hospital Anxiety and Depression Scale, the Patient Health Questionnaire, a screening question, and physicians' diagnosis. *J Psychosom Res* 2003;**55**:515-9.
- 95 Hurst MW, Jenkins CD, Rose RM. The assessment of life change stress: a comparative and methodological inquiry. *Psychosom Med* 1978;**40**:126-41.
- 96 Czajkowski SM, Terrin M, Lindquist R, *et al.* Comparison of preoperative characteristics of men and women undergoing coronary artery bypass grafting (the Post Coronary Artery Bypass Graft [CABG] Biobehavioral Study). *Am J Cardiol* 1997;**79**:1017-24.

- 1  
2  
3 97 Klauke B, Deckert J, Reif A, *et al.* Life events in panic disorder—an update on  
4 "candidate stressors". *Depress Anxiety* 2010;**27**:716-30.  
5  
6  
7  
8 98 Berhaut JC, Graham ID, Wood TJ, *et al.* Measuring acceptability of clinical  
9 decision rules: validation of the Ottawa acceptability of decision rules instrument  
10 (OADRI) in four countries. *Med Decis Making* 2010;**30**:398-408.  
11  
12  
13 99 Altman DG, Royston P. What do we mean by validating a prognostic model? *Stat*  
14 *Med* 2000;**19**:453-73.  
15  
16  
17 100 Zhang H, Singer B. *Recursive Partitioning in the Health Sciences*. New York,  
18 NY: Springer-Verlag 1999.  
19  
20  
21 101 Breiman L, Friedman J, Stone CJ, *et al.* *Classification and Regression Trees*.  
22 Belmont, CA: Wadsworth International Group 1984.  
23  
24  
25 102 Skov T, Deddens J, Petersen MR, *et al.* Prevalence proportion ratios: estimation  
26 and hypothesis testing. *Int J Epidemiol* 1998;**27**:91-5.  
27  
28  
29 103 Sullivan LM, Massaro JM, D'Agostino RB Sr. Presentation of multivariate data  
30 for clinical use: the Framingham Study risk score functions. *Stat Med*  
31 2004;**23**:1631-60.  
32  
33  
34 104 Fleiss JL. *Statistical Methods for Rates and Proportions. Second Edition*. New  
35 York, NY: John Wiley & Sons 1981.  
36  
37  
38 105 Hosmer DW, Lemeshow S. *Applied Logistic Regression. Second Edition*. New  
39 York, NY: John Wiley & Sons 2000.  
40  
41  
42 106 Roberge P, Marchand A, Reinharz D, *et al.* Health care utilization following  
43 cognitive-behavioral treatment for panic disorder with agoraphobia. *Cogn Behav*  
44 *Ther* 2005;**34**:79-88.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- 107 Katon W, Russo J, Sherbourne C, *et al.* Incremental cost-effectiveness of a collaborative care intervention for panic disorder. *Psychol Med* 2006;**36**:353-63.
- 108 Marchand A, Belleville G, Fleet R, *et al.* Treatment of panic in chest pain patients from emergency departments: efficacy of different interventions focusing on panic management. *Gen Hosp Psychiatry* 2012;**34**:671-80.

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11 **Figure 1 The Panic Screening Score (PSS)**  
12

13 1) Does the patient have a history of anxiety disorders?  
14 Yes = 7  
15 No = 0, ask other questions.

16 2) Please indicate how often this thought occurs when you are nervous:  
17 "I will choke to death."  
18 0: Never occurs, 2: Rarely occurs, 4: Occurs during half the times,  
19 6: Usually occurs, 8: Always occurs.

20 3) Did the patient arrive in the ED by ambulance?  
21 Yes = 3  
22 No = 0

23 4) Please answer the statement by circling the number that best applies to you.  
24 "When I notice my heart beating rapidly, I worry that I might be having a heart attack."  
25 0: Very little, 1: A little, 2: Some, 3: Much, 4: Very much.

26 Scoring:  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- Total score = sum of all items value
- A total score  $\geq 6$  indicates probable panic.

90x116mm (300 x 300 DPI)