



Respiratory infections in preterm infants and subsequent asthma: a cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-004034
Article Type:	Research
Date Submitted by the Author:	15-Sep-2013
Complete List of Authors:	Montgomery, Scott; Örebro University Hospital and Örebro University, Clinical Epidemiology and Biostatistics; Karolinska Institutet, Clinical Epidemiology Unit, Department of Medicine Bahmanyar, Shahram; The Clinical Epidemiology Unit, Karolinska Institutet; The Centre for Pharmacoepidemiology, Karolinska Institutet Brus, Ole; Clinical Epidemiology and Biostatistics, Örebro University Hospital & Örebro University Hussein, Oula; Örebro University Hospital, Clinical Epidemiology and Biostatistics Kosma, Paraskevi; Department of Woman and Child health, Karolinska Institutet; Division of Neonatology, Karolinska hospital Palme-Kilander, Charlotte; Karolinska Institutet, Department of Woman and Child health
Primary Subject Heading:	Respiratory medicine
Secondary Subject Heading:	Epidemiology
Keywords:	Asthma < THORACIC MEDICINE, gestational age, bronchiolitis, cohort study

SCHOLARONE™
Manuscripts

Respiratory infections in preterm infants and subsequent asthma: a cohort study

Scott Montgomery¹²³ (scott.montgomery@orebroll.se),
Shahram Bahmanyar²⁴ (shahram.bahmanyar@ki.se),
Ole Brus¹ (ole.brus@orebroll.se),
Oula Hussein¹ (oula.hussein@orebroll.se),
Paraskevi Kosma⁵⁶ (evaki782002@yahoo.com),
Charlotte Palme-Kilander⁵ (charlotte.palme-kilander@karolinska.se)

1. Clinical Epidemiology and Biostatistics, Örebro University Hospital & Örebro University, Örebro, Sweden.
2. The Clinical Epidemiology Unit, Karolinska Institutet, Stockholm, Sweden.
3. Department of Epidemiology and Public Health, University College London, UK.
4. The Centre for Pharmacoepidemiology, Karolinska Institutet, Stockholm, Sweden.
5. Department of Woman and Child health, Karolinska Institutet, Stockholm, Sweden
6. Division of Neonatology, Karolinska Hospital, Stockholm, Sweden

Correspondence to:

Prof. Scott Montgomery
Clinical Epidemiology and Biostatistics
S-huset
Örebro University Hospital
701 85 Örebro
Sweden

Scott.montgomery@orebroll.se
Tel: +46 19 602 6210

Word count 2356 (main text); 238 (abstract)

Keywords gestational age; bronchiolitis; asthma; cohort study

ABSTRACT

Objectives To investigate if gestational age modifies the association of airway infections that result in hospital admission during the first year after birth, with subsequent asthma risk after age 5 years.

Setting Hospital inpatients and a general population comparison group in Sweden followed for subsequent diagnoses in primary and secondary care.

Participants National registers identified 42,334 children admitted to hospital for respiratory infection in their first year after birth during 1981-1995, individually matched with 211,594 children not admitted to hospital for infection during their first year.

Primary outcome Asthma diagnoses and prescribed asthma treatments after age 5 years identified through registers.

Results Cox regression was used to identify a hazard ratio (and 95% confidence interval) of 1.51 (1.47 to 1.51) for the association of respiratory infection before one year of age with asthma after age 5 years, after adjustment for sex, gestational age, chronic lung disease, maternal asthma and maternal smoking. When stratified by gestational age (and with additional adjustment for birth weight), there is statistically significant effect modification by gestational age, with the highest magnitude asthma risk among those born with a gestational age of less than 28 weeks, producing an adjusted hazard ratio of 2.22 (1.59 to 3.09). This higher magnitude asthma risk persisted until after age 10 years, but differences in risk by gestational age were less pronounced for asthma after age 16 years.

Conclusions Extremely preterm infants are most likely to have chronic respiratory sequelae following respiratory infections in early life.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

BMJ Open: first published as 10.1136/bmjopen-2013-004034 on 29 October 2013. Downloaded from <http://bmjopen.bmj.com/> on April 23, 2021 by guest. Protected by copyright.

STRENGTHS AND LIMITATIONS OF THIS STUDY

Strengths

- The vast majority of infants admitted to hospital with airway infections in their first year after birth in Sweden during the study period will have been identified.
- The use of prospectively recorded national register data allow for longitudinal analysis of asthma risk, with follow-up into early adulthood.

Limitations

- The infections are incompletely characterised, particularly during earlier years.
- Asthma was identified based on inpatient diagnoses and prescribed medication, but it was not possible to confirm the accuracy of diagnoses.

INTRODUCTION

Respiratory infection in the first year after birth^[1-3], and preterm birth^[4-7] are both markers of subsequent asthma risk. How the interrelationship of these factors relates to persistent respiratory disease is incompletely understood.

The lungs of preterm infants are underdeveloped with immature airways and the association of prematurity with later respiratory disease indicates asthma's developmental origin^[8-10]. Respiratory infections requiring hospital admission during the first year after birth may signal susceptibility to asthma or may be causally implicated in asthma aetiology; and these explanations are not necessarily mutually exclusive. Respiratory infection *epidemics* that produce infections among infants are associated with a raised risk of persistent respiratory disease^[11], consistent with a causal explanation for at least a proportion of the association.

Earlier research has demonstrated that degree of prematurity is relevant to risk of persistent asthma and bronchiolitis in the first year^[6]. This paper investigates whether the *combination*

1
2
3 (interaction) of early respiratory infection with prematurity, when the immature lungs may
4
5 be particularly susceptible to long-term damage, represents a disproportionately raised risk
6
7 for persistent respiratory disease. Swedish general population registers were used to identify
8
9 a cohort of children who were admitted to hospital due to respiratory infection in their first
10
11 year after birth and they were matched with a cohort of children not admitted to hospital
12
13 for infection in their first year.
14
15
16
17
18
19
20
21
22
23

24 **METHODS**

25
26 All children in Sweden admitted to hospital for severe infections that affect the airways in
27
28 the 12 months following birth between 1981 and 1995 were identified (N=4233). They were
29
30 individually matched with children who were not admitted to hospital for these infections in
31
32 their first year (N=211594), with matching for gestational age, region of birth, and the month
33
34 and year of their birth. Each exposed child was matched with five unexposed children,
35
36 selected at random from among those with relevant characteristics. If fewer than five
37
38 unexposed infants were available for matching with an exposed subject, all suitable children
39
40 were selected. Follow-up to identify chronic airway disease (asthma) was up to 2010.
41
42
43
44
45
46
47

48 **Registers**

49
50 This study utilised national Swedish register data that can be linked at the individual level
51
52 using the unique personal identity number issued to all residents. Information on pregnancy
53
54 and delivery, and other characteristics of mother and baby were obtained through the
55
56 Medical Birth Register, which has recorded almost every delivery in Sweden since 1973. The
57
58
59
60

1
2
3 Patient Register has had complete national coverage of inpatient diagnoses since 1987 and
4
5 included outpatient diagnoses since 2000. The Prescription Register has provided
6
7 information on all prescribed medication since 2005.
8
9

10 11 12 **Measures**

13
14 The Patient Register was used to identify respiratory infections in the first year after birth
15
16 that resulted in hospital admission; and these diagnoses are listed in Table 1. This register
17
18 also identified diagnoses of lung disease through ICD codes consistent with asthma in
19
20 children and young adults. To avoid surveillance bias (incidental reporting of asthma in
21
22 children admitted or investigated for other diseases) asthma was only defined as the
23
24 outcome here when it was the sole diagnosis made at the hospital visit. The Prescription
25
26 Register – which records all treatments dispensed through pharmacies (including all
27
28 prescriptions from primary care) was used to identify pharmaceutical treatments for asthma
29
30 in children and young adults and the selected treatments are presented in table 1. The
31
32 Medical Birth Register provided information on gestational age, birth weight, maternal age
33
34 at delivery, maternal smoking during pregnancy, maternal asthma, sex of the child, parity
35
36 and a diagnosis of neonatal chronic lung disease (CLD: ICD-8 776.6 and ICD-9 770H): these
37
38 measures were categorised as presented in table 2.
39
40
41
42
43
44
45
46
47

48 **Statistical analysis**

49
50 Cox regression was used to estimate hazard ratios associated with asthma after 5 years of
51
52 age (hospital diagnosis or treatment), with follow-up to first asthma event, exit from the
53
54 study population or end of the study period, whichever came first. The following
55
56 independent measures were also included in models individually, then mutually adjusted for
57
58
59
60

1
2
3 each other: respiratory infection in the first year after birth, gestational age, sex, maternal
4
5 age at delivery, parity, maternal smoking, a maternal diagnosis of asthma, CLD and year of
6
7 delivery (all modelled as series of binary dummy variables in the categories shown in tables 2
8
9 and 3). The non-stratified models (table 3) were not adjusted for birth weight, as it was
10
11 collinear with gestational age while the stratified models (tables 4 and 5) and interaction
12
13 tests were adjusted for a continuous measure of birth weight. Log-minus-log plots were
14
15 used to assess whether the proportional hazards assumption was violated: the curves
16
17 comparing the exposed and unexposed cohorts did not converge and proportionality was
18
19 maintained.
20
21
22
23
24
25

26
27 The main analysis concerns the *combination* of early infection with gestational age and this
28
29 was investigated in models for the association of first-year infection with asthma after age 5
30
31 years, stratified by gestational age, with adjustment for all of the potential confounding
32
33 factors described above, with the addition of birth weight modelled as a continuous
34
35 measure. To assess effect modification by gestational age for the association of early
36
37 infection with asthma, interaction testing using the entire (non-stratified) study population
38
39 was undertaken. The interaction terms for gestational age (categorical) by infection were
40
41 adjusted for the main effect (gestational age and infection), as well as for all of the
42
43 previously described potential confounding factors.
44
45
46
47
48
49

50
51 To assess whether the combination of prematurity with first year infection is a risk for
52
53 asthma that persists beyond early childhood, the above analysis stratified for gestational age
54
55 was repeated, but with truncated follow-up to identify asthma after age 10 years and after
56
57 age 16 years.
58
59
60

1
2
3
4
5 SPSS software was used and statistical significance was defined as confidence intervals that
6
7 do not include 1.00 and p values below 0.05.
8
9

10 11 12 **RESULTS**

13
14 As the two cohorts were matched, the distributions for gestational age and year of birth are
15 almost identical (table 2). This matching is also reflected in the very similar distributions for
16 birth weight. Children admitted to hospital with infections were more often male and their
17 mothers were more likely to be younger, smokers, with a diagnosis of asthma and to have
18 had a greater number of previous pregnancies. CLD was also more common in the infection
19 cohort, particularly among the extremely preterm infants (12.3% compared with 7.9%
20 among the extremely preterm).
21
22
23
24
25
26
27
28
29
30
31
32

33
34 Respiratory infection during the first year is statistically significantly associated with a raised
35 risk of asthma after age 5 years, both before and after adjustment for potential confounding
36 factors (table 3). Female sex, neonatal lung disease, maternal asthma, fewer previous
37 pregnancies are associated with a statistically significant raised asthma risk in the adjusted
38 model. Gestational age was a matching characteristic, so associations with this measure are
39 not presented.
40
41
42
43
44
45
46
47
48
49

50 Table 4 shows the association of hospital admission for infection in the first year with
51 asthma after age 5 years, *stratified by gestational age*. Infection is associated with asthma
52 across all of the gestational age groups after adjustment for potential confounding factors.
53
54
55
56
57
58
59
60 The highest magnitude association is among the most preterm infants. A notably raised risk

1
2
3 was also observed among those with uncertain gestational age. Interaction testing among
4
5 the entire study population confirms statistically significant effect modification by the most
6
7 premature gestational age group, at less than 28 weeks compared with the infants who were
8
9 born at term. The interaction results were not statistically significant for the other
10
11 gestational age groups, including where it was unspecified. After adjustment for the main
12
13 effects the hazard ratio (and 95% confidence interval) is 1.41 (1.02 to 1.95; p for
14
15 interaction=0.034) and with additional adjustment for the potential confounding factors it is
16
17 1.40 (1.01 to 1.93; p for interaction=0.042).
18
19
20
21
22
23

24 Early Infection is associated with asthma after age 10 and after age 16 years for all
25
26 gestational age groups, but although still statistically significant, the notably increased
27
28 magnitude for the most premature is somewhat reduced after age 16 years (table 5).
29
30
31
32

33 The analysis was repeated excluding those with missing maternal smoking information and
34
35 the results were not notably altered (data not shown).
36
37
38
39

40 DISCUSSION

41
42 In this register-based cohort study, hospital admission for respiratory infection in the 12
43
44 months following birth was associated with an increased risk of asthma after age 5 years,
45
46 particularly among those born extremely prematurely.
47
48
49
50

51
52 Another study using Swedish register data found that early infection and prematurity were
53
54 independently associated with later increased corticosteroid use^[6]. While the results
55
56 between the earlier study and ours are similar for associations with infection, the cohorts in
57
58
59
60

our study were matched for gestational age, which was considered as a potential modifying factor for the effect of early infections. Here, asthma after age 5 years was identified through prescribed corticosteroids and other asthma medication, as well as inpatient or outpatient hospital diagnoses. We used several strategies to ensure that the associations are with asthma rather than other comorbidity associated with prematurity and early infection. Only diagnoses of asthma made after age 5 years were included, as these are more likely to represent persistent asthma. We limited diagnoses to instances where asthma was the sole diagnosis recorded by the outpatient or inpatient discharge summary to minimise surveillance bias, where admission for another condition results in an incidental diagnosis of asthma: this can produce a conservative estimate of association. While it can be difficult to differentiate between the effects of infection and susceptibility to infection, we attempted to tackle this through adjustment for a variety of potential confounding factors, including birth weight within gestational age categories to provide fine-grain control for foetal development. The extremely preterm infants more often had CLD, possibly signalling to both susceptibility to early infection and also asthma: we therefore adjusted for neonatal CLD.

This study has some potential limitations. Identification of asthma through registers does not provide information about disease phenotype, including whether allergic sensitisation is implicated and confirmation of individual diagnoses is not possible. The infections defining the exposed cohort are not always well characterised (particularly from the earlier periods), but all will affect the lungs and were sufficiently serious to result in hospital admission. Prior to 1990, respiratory syncytial virus (RSV) was often diagnosed as an unspecified viral infection, so we included some non-specific infections. Earlier studies found that RSV was responsible for the majority of bronchiolitis diagnoses during the first year after birth^[12 13],

1
2
3 while rhinovirus was associated with 18% of bronchiolitis diagnoses, although hospital
4 admission was less common^[14]. It is possible that use of more specific diagnoses, not
5 available through the Patient Register, would result in higher magnitude associations. We
6 were unable to examine delivery mode as a potential confounding factor as this was not
7 available in our data set: caesarean section has been linked with an increased asthma risk in
8 term infants^[15]. Number of older siblings is an important indicator of asthma risk. While this
9 was not measured directly, we adjusted for parity as a reliable indicator of number of older
10 siblings. Also, it has been hypothesised that previous pregnancies – directly measured by
11 parity - may influence immune-mediated disease risk through changes in the mother's
12 immune profile^[16]. The measure of maternal smoking was incomplete as it was not always
13 included in the Medical Birth Register. The proportions without smoking data were almost
14 identical in both the exposed and unexposed cohorts. When the analysis was restricted to
15 mothers with complete smoking data, the results were not materially altered. Our study will
16 have underestimated the total number of asthma diagnoses, as we did not have data from
17 primary care, and because we excluded asthma with comorbid diagnoses to avoid
18 confounding by comorbidity. The use of prescribed medication will have improved our ability
19 to identify chronic asthma as this will identify those diagnosed and treated in primary care.
20 The two strategies to identify asthma were combined to maximise reliable asthma
21 identification and statistical power. Unavoidably, gestational age was unavailable for some
22 infants: the higher magnitude asthma risk associated with infections in this group may be
23 due to a proportion who were also very premature, and possibly other risks. As interaction
24 testing by gestational age was undertaken, adjustment rather than internal stratification for
25 this factor was performed. Despite this, use of a matching characteristic may still have
26 resulted in underestimation of interaction effects between infection and gestational age.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5 Both RSV and rhinovirus infections in infancy have been linked with subsequent childhood
6 wheezing^[1 3], but the association of early life infections with allergy and possible asthma risk
7 has a complicated pattern. While some markers of infection, such as positive serology for
8 Epstein-Barr virus (EBV) and cytomegalovirus (CMV) are associated with a reduced risk of
9 allergic sensitisation^[17], there is evidence that early infection with RSV may increase the risk
10 of allergic asthma through its influence on regulatory T-cell function^[18]. However, pre-
11 existing atopy – or associated characteristics - may also influence subsequent infection risk,
12 as rhinovirus bronchiolitis has been found to occur more frequently in infants who had prior
13 allergic sensitization^[19]. Whether or not allergy is implicated, respiratory infections in
14 infancy, both RSV and non-RSV infections, are thought to contribute to airway impairment^{[3}
15 20-23]. The possible non-mutually exclusive mechanisms include through structural
16 damage^[24], modification of the mucosal immunology of the lungs^[25] or through epigenetic
17 changes^[26]. Treatment of infections by antibiotics has been suggested as a risk for asthma,
18 but the evidence is weak; and reviews suggest that the association is due to confounding by
19 indication.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Lung development continues throughout foetal life and babies who are born prematurely
44 have developmental deficits of airway^[27-30] and may result in chronic asthma^[8], which can
45 diminish with increasing age as lung function improves^[31 32]. Atopic sensitization – and thus
46 allergic asthma that may be more persistent in adulthood - is less common in those born
47 preterm compared with term deliveries^[33]. Thus, the combination of early infection and
48 prematurity may conspire to increase both allergic and non-allergic risks for asthma, helping
49 to explain the multiplicative effect of this combination of exposures for the risk of
50
51
52
53
54
55
56
57
58
59
60

1
2
3 subsequent asthma. The notably higher risk of asthma associated with this exposure
4
5 combination continues throughout childhood, but appears to be less profound after age 16
6
7 years, possibly reflecting age-associated changes.
8
9

10
11
12 This study provides evidence that respiratory infections resulting in hospital admission
13
14 during the first year after birth are a risk for subsequent asthma after age 5 years. The
15
16 increase in childhood asthma risk is particularly high among extremely preterm infants.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

What is already known on this subject

- Infants born preterm have an increased risk of airway disease and subsequent childhood asthma.
- Respiratory infection in infancy is also associated with an increased risk of childhood asthma, independent of gestational age.

What this study adds

- Extremely premature infants who experience respiratory infections before one year of age have an even higher increase in risk of subsequent asthma than term infants.

Funding The study was supported by the Abbvie Research Fund and Örebro University strategic funding.

Contributors SM and CPK conceived the study and prepared the first draft of the manuscript; and with SB they developed the methods. SB and PK were responsible for obtaining and preparing the data. OB and OH conducted the statistical analysis and produced the tables. All authors were involved in critical editing of the manuscript.

Competing interests None.

Ethics approval Karolinska Regional Ethics Committee, Stockholm

Data sharing: We do not have ethical permission to share the data, however they are register data and available to other academic researchers.

REFERENCES

1. Sigurs N, Aljassim F, Kjellman B, et al. Asthma and allergy patterns over 18 years after severe RSV bronchiolitis in the first year of life. *Thorax* 2010;**65**(12):1045-52.
2. Hyvarinen MK, Kotaniemi-Syrjanen A, Reijonen TM, et al. Lung function and bronchial hyper-responsiveness 11 years after hospitalization for bronchiolitis. *Acta Paediatr* 2007;**96**(10):1464-9.
3. Jackson DJ, Gangnon RE, Evans MD, et al. Wheezing rhinovirus illnesses in early life predict asthma development in high-risk children. *Am J Respir Crit Care Med*. 2008;**178**(7):667-72.
4. Kallen B, Finnstrom O, Nygren KG, et al. Association between preterm birth and intrauterine growth retardation and child asthma. *Eur Respir J* 2013;**41**(3):671-6.
5. Vrijlandt EJ, Kerstjens JM, Duiverman EJ, et al. Moderately Preterm Children Have More Respiratory Problems during Their First 5 Years of Life Than Children Born Full Term. *Am J Respir Crit Care Med* 2013;**187**(11):1234-40.
6. Vogt H, Lindstrom K, Braback L, et al. Preterm birth and inhaled corticosteroid use in 6- to 19-year-olds: a Swedish national cohort study. *Pediatrics* 2011;**127**(6):1052-9.
7. Doyle LW. Respiratory function at age 8-9 years in extremely low birthweight/very preterm children born in Victoria in 1991-1992. *Pediatr Pulmonol* 2006;**41**(6):570-6.
8. Fawke J, Lum S, Kirkby J, et al. Lung function and respiratory symptoms at 11 years in children born extremely preterm: the EPICure study. *Am J Respir Crit Care Med* 2010;**182**(2):237-45.
9. Jaakkola JJ, Ahmed P, Ieromnimon A, et al. Preterm delivery and asthma: a systematic review and meta-analysis. *J Allergy Clin Immunol* 2006;**118**(4):823-30.
10. Greenough A. Long-term pulmonary outcome in the preterm infant. *Neonatology* 2008;**93**(4):324-7.
11. Wu P, Hartert TV. Evidence for a causal relationship between respiratory syncytial virus infection and asthma. *Expert Rev Anti Infect Ther* 2011;**9**(9):731-45.
12. Broughton S, Roberts A, Fox G, et al. Prospective study of healthcare utilisation and respiratory morbidity due to RSV infection in prematurely born infants. *Thorax* 2005;**60**(12):1039-44.
13. Wickman M, Farahmand BY, Persson PG, et al. Hospitalization for lower respiratory disease during 20 yrs among under 5 yr old children in Stockholm County: a population based survey. *Eur Respir J* 1998;**11**(2):366-70.
14. Miller EK, Williams JV, Gebretsadik T, et al. Host and viral factors associated with severity of human rhinovirus-associated infant respiratory tract illness. *J Allergy Clin Immunol* 2011;**127**(4):883-91.
15. Almqvist C, Cnattingius S, Lichtenstein P, et al. The impact of birth mode of delivery on childhood asthma and allergic diseases--a sibling study. *Clin Exp Allergy* 2012;**42**(9):1369-76.
16. Amoudruz P, Holmlund U, Schollin J, et al. Maternal country of birth and previous pregnancies are associated with breast milk characteristics. *Pediatr Allergy Immunol* 2009;**20**(1):19-29.

17. Nilsson C, Linde A, Montgomery SM, et al. Does early EBV infection protect against IgE sensitization? *J Allergy Clin Immunol* 2005;**116**(2):438-44.
18. Krishnamoorthy N, Khare A, Oriss TB, et al. Early infection with respiratory syncytial virus impairs regulatory T cell function and increases susceptibility to allergic asthma. *Nat Med* 2012;**18**(10):1525-30.
19. Jackson DJ, Evans MD, Gangnon RE, et al. Evidence for a causal relationship between allergic sensitization and rhinovirus wheezing in early life. *Am J Respir Crit Care Med* 2012;**185**(3):281-5.
20. Fjaerli HO, Farstad T, Rod G, et al. Acute bronchiolitis in infancy as risk factor for wheezing and reduced pulmonary function by seven years in Akershus County, Norway. *BMC Pediatr* 2005;**5**:31.
21. Romero JR, Stewart DL, Buysman EK, et al. Serious early childhood wheezing after respiratory syncytial virus lower respiratory tract illness in preterm infants. *Clin Ther* 2010;**32**(14):2422-32.
22. Koponen P, Helminen M, Paassilta M, et al. Preschool asthma after bronchiolitis in infancy. *Eur Respir J* 2012;**39**(1):76-80.
23. Mikalsen IB, Halvorsen T, Oymar K. The outcome after severe bronchiolitis is related to gender and virus. *Pediatr Allergy Immunol* 2012;**23**(4):391-8.
24. Benjamin JT, Carver BJ, Plosa EJ, et al. NF-kappaB activation limits airway branching through inhibition of Sp1-mediated fibroblast growth factor-10 expression. *J Immunol* 2010;**185**(8):4896-903.
25. Han J, Takeda K, Gelfand EW. The Role of RSV Infection in Asthma Initiation and Progression: Findings in a Mouse Model. *Pulm Med* 2011;**2011**:748038.
26. Joss-Moore LA, Lane RH. Epigenetics and the developmental origins of disease: the key to unlocking the door of personalized medicine. *Epigenomics* 2012;**4**(5):471-3.
27. Hjalmarson O, Sandberg K. Abnormal lung function in healthy preterm infants. *Am J Respir Crit Care Med* 2002;**165**(1):83-7.
28. Friedrich L, Pitrez PM, Stein RT, et al. Growth rate of lung function in healthy preterm infants. *Am J Respir Crit Care Med* 2007;**176**(12):1269-73.
29. Boyle EM, Poulsen G, Field DJ, et al. Effects of gestational age at birth on health outcomes at 3 and 5 years of age: population based cohort study. *BMJ* 2012;**344**:e896.
30. Kotecha SJ, Watkins WJ, Paranjothy S, et al. Effect of late preterm birth on longitudinal lung spirometry in school age children and adolescents. *Thorax* 2012;**67**(1):54-61.
31. Narang I, Rosenthal M, Cremonesini D, et al. Longitudinal evaluation of airway function 21 years after preterm birth. *Am J Respir Crit Care Med* 2008;**178**(1):74-80.
32. Crump C, Winkleby MA, Sundquist J, et al. Risk of asthma in young adults who were born preterm: a Swedish national cohort study. *Pediatrics* 2011;**127**(4):e913-20.
33. Siltanen M, Wehkalampi K, Hovi P, et al. Preterm birth reduces the incidence of atopy in adulthood. *J Allergy Clin Immunol* 2011;**127**(4):935-42.

Table 1. Respiratory infections and prescribed asthma treatments

Respiratory infections during the first year of life	
Adenovirus infection	Pneumococcal pneumonia
Other specified Adenovirus infection	Pneumonia
Bordetella Pertussis	Pneumonia due to streptococcus
Bronchitis	Pneumonia due to Haemophilus influenzae
Bronchiolitis	Pneumonia due to respiratory syncytial virus
Bronchopneumonia organism unspecified	Other unspecified pneumonia
Coxsackie virus infection	Bacterial pneumonia unspecified
Cytomegalovirus	Other specified viral pneumonia
Echo virus infection	Pneumonia in other infectious diseases classified elsewhere
Influenza	Rhinovirus infection
Influenza with other respiratory manifestations	Other specified viral infections
Influenza unspecified	Suspected viral infections
Influenza with pleuropneumonia	Unspecified viral infections
Prescribed treatments	
Beclomethasone	Omalizumab
Budesonide	Salbutamol
Flutikasone	Salmeterol
Formoterol	Sodiumchromoglicate
Indakaterol	Terbutaline
Ipratropium	Theophylline
Mometasone	Tiotropium bromide
Montelukast	

Table 2. Characteristics of the cohorts defined by respiratory infection during the first year of life

	Respiratory infection in first year	No respiratory infection in first year	Total
	n= 42334 (16.7%)	n= 211594 (83.3%)	n= 253928 (100%)
Prescribed asthma medication*	7485 (17.7)	26591 (12.6)	34076 (13.4)
Hospital asthma diagnosis*	420 (1.0)	761 (0.4)	1181 (0.5)
No asthma*	34749 (82.1)	184775 (87.3)	219524 (86.5)
Gestational age (weeks)			
≤27	252 (0.6)	1221 (0.6)	1473 (0.6)
28-31	723 (1.7)	3578 (1.7)	4301 (1.7)
32-36	3906 (9.2)	19528 (9.2)	23434 (9.2)
37- 41	34839 (82.3)	174194 (82.3)	209033 (82.3)
> 41	2507 (5.9)	12535 (5.9)	15042 (5.9)
Uncertain	107 (0.3)	538 (0.3)	645 (0.3)
Birth weight			
<1000g	241(0.6)	1292 (0.6)	1533 (0.6)
> 1000-1500g	580 (1.4)	2714 (1.3)	3294 (1.3)
>1500-2000g	953 (2.3)	4300 (2.0)	5253 (2.1)
>2000-2750g	4099 (9.7)	18815 (8.9)	22914 (9.0)
>2750-3250g	9727 (23.0)	47617 (22.5)	57344 (22.6)
>3250-4000g	20178 (47.7)	104693 (49.5)	124871 (49.2)
> 4000g	6426 (15.2)	31524 (14.9)	37950 (14.9)
Not recorded	130 (0.3)	639 (0.3)	769 (0.3)
Maternal asthma			
Current	721 (1.7)	2754 (1.3)	3475 (1.4)
Previous	210 (0.5)	749 (0.4)	959 (0.4)
None	41403 (97.8)	208091 (98.3)	249494 (98.3)
Maternal age at delivery			
13-18	755 (1.8)	2917 (1.4)	3672 (1.5)
19-25	14230 (33.6)	66604 (31.5)	80834 (31.8)
26-30	15312 (36.2)	76752 (36.3)	92064 (36.3)
31-35	8911 (21.0)	46336 (21.9)	55247 (21.8)
36-40	2757 (6.5)	16673 (7.9)	19430 (7.7)
41-54	369 (0.9)	2312 (1.1)	2681 (1.1)
Maternal smoking			
Non-smoker	23871 (56.4)	134802 (63.7)	158673 (62.5)
Smoker	12429 (29.4)	46817 (22.1)	59246 (23.3)
Not recorded	6034 (14.3)	29975 (14.2)	36009 (14.2)
Child's sex			
Female	17619 (41.6)	103007 (48.7)	120626 (47.5)
Male	24715 (58.4)	108587 (51.3)	133302 (52.5)
Parity			
1	12714 (30.0)	87275 (41.2)	99989 (39.4)
2	17247 (40.7)	74534 (35.2)	91781 (36.1)
3	8270 (19.5)	34880 (16.5)	43150 (17.0)
4 or more	4103 (9.7)	14905 (7.0)	19008 (7.5)
Chronic lung disease			
No	42272 (99.9)	211416 (99.9)	253688 (99.9)
Yes	62 (0.1)	178 (0.1)	240 (0.1)

* These categories are not mutually exclusive.

Table 3. Early life characteristics and subsequent asthma risk after age 5 years

Characteristic	Asthma n (%)	No asthma n (%)	Unadjusted HR (95% CI)	P value	Adjusted ¹ HR (95% CI)	P value
Respiratory infection in first year						
Yes	7585 (17.9)	34749 (82.1)	1.45 (1.41 to 1.49)	<0.001	1.51 (1.47 to 1.54)	<0.001
No	26819 (12.7)	184775 (87.3)	Ref.		Reference	
Maternal asthma						
Current	921 (26.5)	2554 (73.5)	3.86 (3.61 to 4.12)	<0.001	1.94 (1.82 to 2.08)	<0.001
Previous	222 (23.1)	737 (76.9)	3.05 (2.67 to 3.48)	<0.001	1.38 (1.21 to 1.58)	<0.001
None	33261 (13.3)	216233 (86.7)	Ref.		Ref.	
Maternal age at delivery (years)						
13-18	517 (14.1)	3155 (85.9)	0.96 (0.88 to 1.05)	0.372	0.98 (0.90 to 1.07)	0.625
19-25	11451 (14.2)	69383 (85.8)	1.02 (1.00 to 1.05)	0.069	1.03 (1.00 to 1.06)	0.026
26-30	12249 (13.3)	79815 (86.7)	Ref.		Ref.	
31-35	7276 (13.2)	47971 (86.8)	0.99 (0.96 to 1.02)	0.490	1.02 (0.99 to 1.05)	0.140
36-40	2562 (13.2)	16868 (86.8)	1.00 (0.96 to 1.05)	0.830	1.05 (1.01 to 1.10)	0.029
41-54	349 (13.0)	2332 (87.0)	1.00 (0.90 to 1.12)	0.934	1.05 (0.94 to 1.17)	0.391
Maternal smoking						
Non-smoker	21701 (13.7)	136972 (86.3)	Ref.		Ref.	
Smoker	8278 (14.0)	50968 (86.0)	0.93 (0.91 to 0.96)	<0.001	1.01 (0.99 to 1.04)	0.380
Not recorded	4425 (12.3)	31584 (87.7)	0.59 (0.57 to 0.61)	<0.001	0.90 (0.87 to 0.93)	<0.001
Child's sex						
Female	18150 (15.0)	102476 (85.0)	1.25 (1.22 to 1.27)	<0.001	1.27 (1.24 to 1.29)	<0.001
Male	16254 (12.2)	117048 (87.8)	Ref.		Ref.	
Parity						
1	14371 (14.4)	85618 (85.6)	Ref.		Ref.	
2	12230 (13.3)	79551 (86.7)	0.93 (0.90 to 0.95)	<0.001	0.90 (0.88 to 0.93)	<0.001
3	5437 (12.6)	37713 (87.4)	0.87 (0.84 to 0.89)	<0.001	0.84 (0.82 to 0.87)	<0.001
4 or more	2366 (12.4)	16642 (87.6)	0.88 (0.84 to 0.92)	<0.001	0.81 (0.78 to 0.85)	<0.001
Chronic lung disease						
No	34360	219328	Ref.		Ref.	
Yes	44	196	1.59 (1.18 to 2.14)	0.002	1.44 (1.06 to 1.96)	0.019

¹ Adjusted for child's sex, maternal asthma, maternal age at delivery, year of birth, maternal smoking, gestational age, infection during first year of life, parity, and chronic lung disease

Table 4. Respiratory infection in the first year of life subsequent asthma risk after age 5 years, stratified by gestational age

Infection stratified by gestational age	Asthma n (%)	No asthma n (%)	HR (95% CI)	P Value	HR ¹ (95% CI)	P Value
Week 0-27						
Infection	52 (20.6)	200 (79.4)	2.03 (1.47 to 2.80)	< 0.001	2.22 (1.59 to 3.09)	< 0.001
No infection	130 (10.6)	1091 (89.4)	Ref.		Ref.	
Week 28-31						
Infection	132 (18.3)	591 (81.7)	1.47 (1.21 to 1.78)	< 0.001	1.57 (1.29 to 1.92)	< 0.001
No infection	458 (12.8)	3130 (87.2)	Ref.		Ref.	
Week 32-36						
Infection	738 (18.9)	3168 (81.1)	1.47 (1.35 to 1.59)	< 0.001	1.54 (1.42 to 1.68)	< 0.001
No infection	2588 (13.3)	16940 (86.7)	Ref.		Ref.	
Week 37-41						
Infection	6194 (17.8)	28645 (82.2)	1.44 (1.40 to 1.48)	< 0.001	1.49 (1.45 to 1.53)	< 0.001
No infection	22075 (12.7)	152119 (87.3)	Ref.		Ref.	
Week > 41						
Infection	446 (17.8)	2061 (82.2)	1.51 (1.36 to 1.68)	< 0.001	1.57 (1.41 to 1.75)	< 0.001
No infection	1512(12.1)	11023 (87.9)	Ref.		Ref.	
Week uncertain						
Infection	23 (21.5)	84 (78.5)	2.10 (1.29 to 3.42)	0.003	2.21 (1.34 to 3.65)	0.002
No infection	56 (10.4)	482 (89.6)	Ref.		Ref.	

¹ Adjusted for child's sex, maternal asthma, maternal age at delivery, year of birth, birth weight, maternal smoking, infection during first year of life, parity, and chronic lung disease

Table 5. Respiratory infection in the first year of life subsequent asthma risk after age 10 years and after age 16 years, stratified by gestational age

	Asthma n (%)	No asthma n (%)	HR (95% CI)	P Value	HR ¹ (95% CI)	P Value
≥ Age 10						
Gestational age (weeks)						
≤27						
Infection	45 (17.9)	207 (82.1)	1.83 (1.30 to 2.57)	0.001	1.99 (1.40 to 2.82)	< 0.001
No infection	123 (10.1)	1098 (89.9)	Ref.		Ref.	
28-31						
Infection	116 (16.0)	607 (84.0)	1.43 (1.17 to 1.76)	0.001	1.52 (1.24 to 1.87)	< 0.001
No infection	411 (11.5)	3167 (88.5)	Ref.		Ref.	
32-36						
Infection	635 (16.3)	3271 (83.7)	1.39 (1.27 to 1.52)	< 0.001	1.46 (1.33 to 1.59)	< 0.001
No infection	2328 (11.)	17200 (88.1)	Ref.		Ref.	
37- 41						
Infection	5541 (15.9)	29298 (84.1)	1.44 (1.40 to 1.48)	< 0.001	1.49 (1.45 to 1.54)	< 0.001
No infection	19637 (11.3)	154557 (88.7)	Ref.		Ref.	
> 41						
Infection	397 (15.8)	2119 (84.2)	1.51 (1.35 to 1.69)	< 0.001	1.57 (1.41 to 1.76)	< 0.001
No infection	1340 (10.7)	11195 (89.3)	Ref.		Ref.	
Uncertain						
Infection	19 (17.8)	88 (82.2)	1.93 (1.14 to 3.28)	0.015	2.05 (1.19 to 3.55)	0.010
No infection	50 (9.3)	488 (90.7)	Ref.		Ref.	
≥ Age 16						
Gestational age (weeks)						
≤27						
Infection	22 (8.8)	227 (91.2)	1.56 (0.96 to 2.52)	0.071	1.65 (1.01 to 2.71)	0.047
No infection	69 (5.7)	1132 (94.3)	Ref.		Ref.	
28-31						
Infection	65 (9.2)	638 (90.8)	1.41 (1.07 to 1.86)	0.014	1.45 (1.10 to 1.91)	0.009
No infection	233 (6.7)	3249 (93.3)	Ref.		Ref.	
32-36						
Infection	396 (10.5)	3362 (89.5)	1.43 (1.28 to 1.60)	< 0.001	1.49 (1.34 to 1.68)	< 0.001
No infection	1410 (7.5)	17379 (92.5)	Ref.		Ref.	
37- 41						
Infection	3227 (9.7)	30040 (90.3)	1.37 (1.32 to 1.42)	< 0.001	1.42 (1.37 to 1.48)	< 0.001
No infection	11987 (7.2)	154347 (92.8)	Ref.		Ref.	
> 41						
Infection	248 (10.3)	2161 (89.7)	1.58 (1.37 to 1.82)	< 0.001	1.64 (1.42 to 1.90)	< 0.001
No infection	801 (6.7)	11244 (93.3)	Ref.		Ref.	
Uncertain						
Infection	16 (15.2)	89 (84.8)	2.48 (1.37 to 4.51)	0.003	2.72 (1.47 to 5.04)	0.002
No infection	33 (6.2)	496 (93.8)	Ref.		Ref.	

¹ Adjusted for child's sex, maternal asthma, maternal age at delivery, year of birth, maternal smoking, birth weight, infection during first year of life, parity, and chronic lung disease.

Respiratory infections in preterm infants and subsequent asthma

Scott Montgomery^{1 2 3} (scott.montgomery@orebroll.se),
Shahram Bahmanyar^{2 4} (shahram.bahmanyar@ki.se),
Ole Brus¹ (ole.brus@orebroll.se),
Oula Hussein¹ (oula.hussein@orebroll.se),
Paraskevi Kosma^{5 6} (evaki782002@yahoo.com),
Charlotte Palme-Kilander⁵ (charlotte.palme-kilander@karolinska.se)

1. Clinical Epidemiology and Biostatistics, Örebro University Hospital & Örebro University, Örebro, Sweden.
2. The Clinical Epidemiology Unit, Karolinska Institutet, Stockholm, Sweden.
3. Department of Epidemiology and Public Health, University College London, UK.
4. The Centre for Pharmacoepidemiology, Karolinska Institutet, Stockholm, Sweden.
5. Department of Woman and Child health, Karolinska Institutet, Stockholm, Sweden
6. Division of Neonatology, Karolinska Hospital, Stockholm, Sweden

Correspondence to:

Prof. Scott Montgomery
Clinical Epidemiology and Biostatistics
S-huset
Örebro University Hospital
701 85 Örebro
Sweden

Scott.montgomery@orebroll.se
Tel: +46 19 602 6210

Word count 2356 (main text); 238 (abstract)

Keywords gestational age; bronchiolitis; asthma; cohort study

Funding The study was supported by the Abbvie Research Fund and Örebro University strategic funding.

Contributors SM and CPK conceived the study and prepared the first draft of the manuscript; and with SB they developed the methods. SB and PK were responsible for obtaining and preparing the data. OB and OH conducted the statistical analysis and produced the tables. All authors were involved in critical editing of the manuscript.

Competing interests None.

Ethics approval Karolinska Regional Ethics Committee, Stockholm

ABSTRACT

Objective To investigate if gestational age modifies the association of airway infections, resulting in hospital admission during the first year after birth, with asthma risk after age 5 years.

Patients Swedish general population registers identified 42,334 children admitted to hospital for respiratory infection in their first year after birth during 1981-1995. They were individually matched with 211,594 children not admitted to hospital for infection.

Methods Cox regression analysis was used to estimate asthma risk after age 5 years (identified through diagnoses and prescribed treatments in registers), with adjustment for sex, gestational age, chronic lung disease, maternal asthma and smoking; and additionally for birth weight in models stratified for gestational age.

Results Hospital admission for airway infection in the first year is associated with an increased risk for subsequent asthma, producing an adjusted hazard ratio (and 95% confidence interval) of 1.51 (1.47 to 1.54). When stratified, the highest magnitude risk (statistically significant effect modification) was among those born with a gestational age of less than 28 weeks, producing an adjusted hazard ratio of 2.22 (1.59 to 3.09). This higher magnitude risk asthma persisted until after age 10 years. Early infection was statistically significantly associated with asthma after age 16 years for all gestational age groups, but the difference in risk magnitude between the extremely prematurely born and other children was reduced.

Conclusions Extremely preterm infants are most likely to have chronic respiratory sequelae following respiratory infections in early life.

STRENGTHS AND LIMITATIONS OF THIS STUDY

Strengths

- The vast majority of infants admitted to hospital with airway infections in their first year after birth in Sweden during the study period will have been identified.
- The use of prospectively recorded national register data allow for longitudinal analysis of asthma risk, with follow-up into early adulthood.

Limitations

- The infections are incompletely characterised, particularly during earlier years.
- Asthma was identified based on inpatient diagnoses and prescribed medication, but it was not possible to confirm the accuracy of diagnoses.

INTRODUCTION

Respiratory infection in the first year after birth^[1-3], and preterm birth^[4-7] are both markers of subsequent asthma risk. How the interrelationship of these factors relates to persistent respiratory disease is incompletely understood.

The lungs of preterm infants are underdeveloped with immature airways and the association of prematurity with later respiratory disease indicates asthma's developmental origin^[8-10]. Respiratory infections requiring hospital admission during the first year after birth may signal susceptibility to asthma or may be causally implicated in asthma aetiology; and these explanations are not necessarily mutually exclusive. Respiratory infection *epidemics* that produce infections among infants are associated with a raised risk of persistent respiratory disease^[11], consistent with a causal explanation for at least a proportion of the association.

Earlier research has demonstrated that degree of prematurity is relevant to risk of persistent asthma and bronchiolitis in the first year^[6]. This paper investigates whether the *combination* (interaction) of early respiratory infection with prematurity, when the immature lungs may

1
2
3 be particularly susceptible to long-term damage, represents a disproportionately raised risk
4
5 for persistent respiratory disease. Swedish general population registers were used to identify
6
7 a cohort of children who were admitted to hospital due to respiratory infection in their first
8
9 year after birth and they were matched with a cohort of children not admitted to hospital
10
11 for infection in their first year.
12
13

14 15 16 17 18 19 20 21 **METHODS**

22
23 All children in Sweden admitted to hospital for severe infections that affect the airways in
24
25 the 12 months following birth between 1981 and 1995 were identified (N=4233). They were
26
27 individually matched with children who were not admitted to hospital for these infections in
28
29 their first year (N=211594), with matching for gestational age, region of birth, and the month
30
31 and year of their birth. Each exposed child was matched with five unexposed children,
32
33 selected at random from among those with relevant characteristics. If fewer than five
34
35 unexposed infants were available for matching with an exposed subject, all suitable children
36
37 were selected. Follow-up to identify chronic airway disease (asthma) was up to 2010.
38
39
40
41
42
43
44

45 **Registers**

46
47 This study utilised national Swedish register data that can be linked at the individual level
48
49 using the unique personal identity number issued to all residents. Information on pregnancy
50
51 and delivery, and other characteristics of mother and baby were obtained through the
52
53 Medical Birth Register, which has recorded almost every delivery in Sweden since 1973. The
54
55 Patient Register has had complete national coverage of inpatient diagnoses since 1987 and
56
57
58
59
60

1
2
3 included outpatient diagnoses since 2000. The Prescription Register has provided
4
5 information on all prescribed medication since 2005.
6
7
8
9

10 **Measures**

11
12 The Patient Register was used to identify respiratory infections in the first year after birth
13
14 that resulted in hospital admission; and these diagnoses are listed in Table 1. This register
15
16 also identified diagnoses of lung disease through ICD codes consistent with asthma in
17
18 children and young adults. To avoid surveillance bias (incidental reporting of asthma in
19
20 children admitted or investigated for other diseases) asthma was only defined as the
21
22 outcome here when it was the sole diagnosis made at the hospital visit. **The Prescription**
23
24 **Register – which records all treatments dispensed through pharmacies (including all**
25
26 **prescriptions from primary care)** was used to identify pharmaceutical treatments for asthma
27
28 in children and young adults and the selected treatments are presented in table 1. The
29
30 Medical Birth Register provided information on gestational age, birth weight, maternal age
31
32 at delivery, maternal smoking during pregnancy, maternal asthma, sex of the child, parity
33
34 and a diagnosis of neonatal chronic lung disease (CLD: ICD-8 776.6 and ICD-9 770H): these
35
36 measures were categorised as presented in table 2.
37
38
39
40
41
42
43
44
45

46 **Statistical analysis**

47
48 Cox regression was used to estimate hazard ratios associated with asthma after 5 years of
49
50 age (hospital diagnosis or treatment), with follow-up to first asthma event, exit from the
51
52 study population or end of the study period, whichever came first. The following
53
54 independent measures were also included in models individually, then mutually adjusted for
55
56 each other: respiratory infection in the first year after birth, gestational age, sex, maternal
57
58
59
60

1
2
3 age at delivery, parity, maternal smoking, a maternal diagnosis of asthma, CLD and year of
4
5 delivery (all modelled as series of binary dummy variables in the categories shown in tables 2
6
7 and 3). The non-stratified models (table 3) were not adjusted for birth weight, as it was
8
9 collinear with gestational age while the stratified models (tables 4 and 5) and interaction
10
11 tests were adjusted for a continuous measure of birth weight. Log-minus-log plots were
12
13 used to assess whether the proportional hazards assumption was violated: the curves
14
15 comparing the exposed and unexposed cohorts did not converge and proportionality was
16
17 maintained.
18
19
20
21
22
23

24 The main analysis concerns the *combination* of early infection with gestational age and this
25
26 was investigated in models for the association of first-year infection with asthma after age 5
27
28 years, stratified by gestational age, with adjustment for all of the potential confounding
29
30 factors described above, with the addition of birth weight modelled as a continuous
31
32 measure. To assess effect modification by gestational age for the association of early
33
34 infection with asthma, interaction testing using the entire (non-stratified) study population
35
36 was undertaken. The interaction terms for gestational age (categorical) by infection were
37
38 adjusted for the main effect (gestational age and infection), as well as for all of the
39
40 previously described potential confounding factors.
41
42
43
44
45
46
47

48 To assess whether the combination of prematurity with first year infection is a risk for
49
50 asthma that persists beyond early childhood, the above analysis stratified for gestational age
51
52 was repeated, but with truncated follow-up to identify asthma after age 10 years and after
53
54 age 16 years.
55
56
57
58
59
60

1
2
3 SPSS software was used and statistical significance was defined as confidence intervals that
4
5 do not include 1.00 and p values below 0.05.
6
7
8

9 10 **RESULTS**

11
12 As the two cohorts were matched, the distributions for gestational age and year of birth are
13
14 almost identical (table 2). This matching is also reflected in the very similar distributions for
15
16 birth weight. Children admitted to hospital with infections were more often male and their
17
18 mothers were more likely to be younger, smokers, with a diagnosis of asthma and to have
19
20 had a greater number of previous pregnancies. CLD was also more common in the infection
21
22 cohort, particularly among the extremely preterm infants (12.3% compared with 7.9%
23
24 among the extremely preterm).
25
26
27

28
29
30
31 Respiratory infection during the first year is statistically significantly associated with a raised
32
33 risk of asthma after age 5 years, both before and after adjustment for potential confounding
34
35 factors (table 3). Female sex, neonatal lung disease, maternal asthma, fewer previous
36
37 pregnancies are associated with a statistically significant raised asthma risk in the adjusted
38
39 model. Gestational age was a matching characteristic, so associations with this measure are
40
41 not presented.
42
43
44

45
46
47 Table 4 shows the association of hospital admission for infection in the first year with
48
49 asthma after age 5 years, *stratified by gestational age*. Infection is associated with asthma
50
51 across all of the gestational age groups after adjustment for potential confounding factors.
52
53

54
55 The highest magnitude association is among the most preterm infants. A notably raised risk
56
57 was also observed among those with uncertain gestational age. **Interaction testing among**
58
59
60

1
2
3 the entire study population confirms statistically significant effect modification by the most
4
5 premature gestational age group, at less than 28 weeks compared with the infants who were
6
7 born at term. The interaction results were not statistically significant for the other
8
9 gestational age groups, including where it was unspecified. After adjustment for the main
10
11 effects the hazard ratio (and 95% confidence interval) is 1.41 (1.02 to 1.95; p for
12
13 interaction=0.034) and with additional adjustment for the potential confounding factors it is
14
15 1.40 (1.01 to 1.93; p for interaction=0.042).
16
17
18
19
20
21

22 Early Infection is associated with asthma after age 10 and after age 16 years for all
23
24 gestational age groups, but although still statistically significant, the notably increased
25
26 magnitude for the most premature is somewhat reduced after age 16 years (table 5).
27
28
29
30

31 The analysis was repeated excluding those with missing maternal smoking information and
32
33 the results were not notably altered (data not shown).
34
35
36
37

38 DISCUSSION

39
40 In this register-based cohort study, hospital admission for respiratory infection in the 12
41
42 months following birth was associated with an increased risk of asthma after age 5 years,
43
44 particularly among those born extremely prematurely.
45
46
47
48
49

50 Another study using Swedish register data found that early infection and prematurity were
51
52 independently associated with later increased corticosteroid use^[6]. While the results
53
54 between the earlier study and ours are similar for associations with infection, the cohorts in
55
56 our study were matched for gestational age, which was considered as a potential modifying
57
58
59
60

1
2
3 factor for the effect of early infections. Here, asthma after age 5 years was identified
4
5 through prescribed corticosteroids and other **asthma medication**, as well as inpatient or
6
7 outpatient hospital diagnoses. We used several strategies to ensure that the associations
8
9 are with asthma rather than other comorbidity associated with prematurity and early
10
11 infection. Only diagnoses of asthma made after age 5 years were included, as these are more
12
13 likely to represent persistent asthma. We limited diagnoses to instances where asthma was
14
15 the sole diagnosis recorded by the outpatient or inpatient discharge summary to minimise
16
17 surveillance bias, where admission for another condition results in an incidental diagnosis of
18
19 asthma: this can produce a conservative estimate of association. While it can be difficult to
20
21 differentiate between the effects of infection and susceptibility to infection, we attempted
22
23 to tackle this through adjustment for a variety of potential confounding factors, including
24
25 birth weight within gestational age categories to provide fine-grain control for foetal
26
27 development. The extremely preterm infants more often had CLD, possibly signalling to both
28
29 susceptibility to early infection and also asthma: we therefore adjusted for neonatal CLD.
30
31
32
33
34
35
36
37

38 **This study has some potential limitations. Identification of asthma through registers does not**
39 **provide information about disease phenotype, including whether allergic sensitisation is**
40 **implicated and confirmation of individual diagnoses is not possible.** The infections defining
41
42 the exposed cohort are not always well characterised (particularly from the earlier periods),
43
44 but all will affect the lungs and were sufficiently serious to result in hospital admission. Prior
45
46 to 1990, respiratory syncytial virus (RSV) was often diagnosed as an unspecified viral
47
48 infection, so we included some non-specific infections. Earlier studies found that RSV was
49
50 responsible for the majority of bronchiolitis diagnoses during the first year after birth^[12 13],
51
52 while rhinovirus was associated with 18% of bronchiolitis diagnoses, although hospital
53
54
55
56
57
58
59
60

1
2
3 admission was less common^[14]. It is possible that use of more specific diagnoses, not
4
5 available through the Patient Register, would result in higher magnitude associations. We
6
7 were unable to examine delivery mode as a potential confounding factor as this was not
8
9 available in our data set: caesarean section has been linked with an increased asthma risk in
10
11 term infants^[15]. Number of older siblings is an important indicator of asthma risk. While this
12
13 was not measured directly, we adjusted for parity as a reliable indicator of number of older
14
15 siblings. Also, it has been hypothesised that previous pregnancies – directly measured by
16
17 parity - may influence immune-mediated disease risk through changes in the mother's
18
19 immune profile^[16]. The measure of maternal smoking was incomplete as it was not always
20
21 included in the Medical Birth Register. The proportions without smoking data were almost
22
23 identical in both the exposed and unexposed cohorts. When the analysis was restricted to
24
25 mothers with complete smoking data, the results were not materially altered. Our study will
26
27 have underestimated the total number of asthma diagnoses, as we did not have data from
28
29 primary care, and because we excluded asthma with comorbid diagnoses to avoid
30
31 confounding by comorbidity. The use of prescribed medication will have improved our ability
32
33 to identify chronic asthma as this will identify those diagnosed and treated in primary care.
34
35 The two strategies to identify asthma were combined to maximise reliable asthma
36
37 identification and statistical power. Unavoidably, gestational age was unavailable for some
38
39 infants: the higher magnitude asthma risk associated with infections in this group may be
40
41 due to a proportion who were also very premature, and possibly other risks. As interaction
42
43 testing by gestational age was undertaken, adjustment rather than internal stratification for
44
45 this factor was performed. Despite this, use of a matching characteristic may still have
46
47 resulted in underestimation of interaction effects between infection and gestational age.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Both RSV and rhinovirus infections in infancy have been linked with subsequent childhood
4 wheezing^[1 3], but the association of early life infections with allergy and possible asthma risk
5 has a complicated pattern. While some markers of infection, such as positive serology for
6 Epstein-Barr virus (EBV) and cytomegalovirus (CMV) are associated with a reduced risk of
7 allergic sensitisation^[17], there is evidence that early infection with RSV may increase the risk
8 of allergic asthma through its influence on regulatory T-cell function^[18]. However, pre-
9 existing atopy – or associated characteristics - may also influence subsequent infection risk,
10 as rhinovirus bronchiolitis has been found to occur more frequently in infants who had prior
11 allergic sensitization^[19]. Whether or not allergy is implicated, respiratory infections in
12 infancy, both RSV and non-RSV infections, are thought to contribute to airway impairment^{[3}
13 20-23]. The possible non-mutually exclusive mechanisms include through structural
14 damage^[24], modification of the mucosal immunology of the lungs^[25] or through epigenetic
15 changes^[26]. Treatment of infections by antibiotics has been suggested as a risk for asthma,
16 but the evidence is weak; and reviews suggest that the association is due to confounding by
17 indication.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40 Lung development continues throughout foetal life and babies who are born prematurely
41 have developmental deficits of airway^[27-30] and may result in chronic asthma^[8], which can
42 diminish with increasing age as lung function improves^[31 32]. Atopic sensitization – and thus
43 allergic asthma that may be more persistent in adulthood - is less common in those born
44 preterm compared with term deliveries^[33]. Thus, the combination of early infection and
45 prematurity may conspire to increase both allergic and non-allergic risks for asthma, helping
46 to explain the multiplicative effect of this combination of exposures for the risk of
47 subsequent asthma. The notably higher risk of asthma associated with this exposure
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 combination continues throughout childhood, but appears to be less profound after age 16
4
5 years, possibly reflecting age-associated changes.
6
7
8
9

10 This study provides evidence that respiratory infections resulting in hospital admission
11 during the first year after birth are a risk for subsequent asthma after age 5 years. The
12
13 increase in childhood asthma risk is particularly high among extremely preterm infants.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

What is already known on this subject

- Infants born preterm have an increased risk of airway disease and subsequent childhood asthma.
- Respiratory infection in infancy is also associated with an increased risk of childhood asthma, independent of gestational age.

What this study adds

- Extremely premature infants who experience respiratory infections before one year of age have an even higher increase in risk of subsequent asthma than term infants.

REFERENCES

1. Sigurs N, Aljassim F, Kjellman B, et al. Asthma and allergy patterns over 18 years after severe RSV bronchiolitis in the first year of life. *Thorax* 2010;**65**(12):1045-52.
2. Hyvarinen MK, Kotaniemi-Syrjanen A, Reijonen TM, Korhonen K, Korppi MO. Lung function and bronchial hyper-responsiveness 11 years after hospitalization for bronchiolitis. *Acta Paediatr* 2007;**96**(10):1464-9.
3. Jackson DJ, Gangnon RE, Evans MD, et al. Wheezing rhinovirus illnesses in early life predict asthma development in high-risk children. *Am J Respir Crit Care Med* 2008;**178**(7):667-72.
4. Kallen B, Finnstrom O, Nygren KG, Otterblad Olausson P. Association between preterm birth and intrauterine growth retardation and child asthma. *Eur Respir J* 2013;**41**(3):671-6.
5. Vrijlandt EJ, Kerstjens JM, Duiverman EJ, Bos AF, Reijneveld SA. Moderately Preterm Children Have More Respiratory Problems during Their First 5 Years of Life Than Children Born Full Term. *Am J Respir Crit Care Med* 2013;**187**(11):1234-40.
6. Vogt H, Lindstrom K, Braback L, Hjern A. Preterm birth and inhaled corticosteroid use in 6- to 19-year-olds: a Swedish national cohort study. *Pediatrics* 2011;**127**(6):1052-9.
7. Doyle LW. Respiratory function at age 8-9 years in extremely low birthweight/very preterm children born in Victoria in 1991-1992. *Pediatr Pulmonol* 2006;**41**(6):570-6.
8. Fawke J, Lum S, Kirkby J, et al. Lung function and respiratory symptoms at 11 years in children born extremely preterm: the EPICure study. *Am J Respir Crit Care Med* 2010;**182**(2):237-45.
9. Jaakkola JJ, Ahmed P, Ieromnimon A, et al. Preterm delivery and asthma: a systematic review and meta-analysis. *J Allergy Clin Immunol* 2006;**118**(4):823-30.
10. Greenough A. Long-term pulmonary outcome in the preterm infant. *Neonatology* 2008;**93**(4):324-7.
11. Wu P, Hartert TV. Evidence for a causal relationship between respiratory syncytial virus infection and asthma. *Expert Rev Anti Infect Ther* 2011;**9**(9):731-45.
12. Broughton S, Roberts A, Fox G, et al. Prospective study of healthcare utilisation and respiratory morbidity due to RSV infection in prematurely born infants. *Thorax* 2005;**60**(12):1039-44.
13. Wickman M, Farahmand BY, Persson PG, Pershagen G. Hospitalization for lower respiratory disease during 20 yrs among under 5 yr old children in Stockholm County: a population based survey. *Eur Respir J* 1998;**11**(2):366-70.
14. Miller EK, Williams JV, Gebretsadik T, et al. Host and viral factors associated with severity of human rhinovirus-associated infant respiratory tract illness. *J Allergy Clin Immunol* 2011;**127**(4):883-91.
15. Almqvist C, Cnattingius S, Lichtenstein P, Lundholm C. The impact of birth mode of delivery on childhood asthma and allergic diseases--a sibling study. *Clin Exp Allergy* 2012;**42**(9):1369-76.
16. Amoudruz P, Holmlund U, Schollin J, Sverremark-Ekstrom E, Montgomery SM. Maternal country of birth and previous pregnancies are associated with breast milk characteristics. *Pediatr Allergy Immunol* 2009;**20**(1):19-29.
17. Nilsson C, Linde A, Montgomery SM, et al. Does early EBV infection protect against IgE sensitization? *J Allergy Clin Immunol* 2005;**116**(2):438-44.

18. Krishnamoorthy N, Khare A, Oriss TB, et al. Early infection with respiratory syncytial virus impairs regulatory T cell function and increases susceptibility to allergic asthma. *Nat Med* 2012;**18**(10):1525-30.
19. Jackson DJ, Evans MD, Gangnon RE, et al. Evidence for a causal relationship between allergic sensitization and rhinovirus wheezing in early life. *Am J Respir Crit Care Med* 2012;**185**(3):281-5.
20. Fjaerli HO, Farstad T, Rod G, Ufert GK, Gulbrandsen P, Nakstad B. Acute bronchiolitis in infancy as risk factor for wheezing and reduced pulmonary function by seven years in Akershus County, Norway. *BMC Pediatr* 2005;**5**:31.
21. Romero JR, Stewart DL, Buysman EK, Fernandes AW, Jafri HS, Mahadevia PJ. Serious early childhood wheezing after respiratory syncytial virus lower respiratory tract illness in preterm infants. *Clin Ther* 2010;**32**(14):2422-32.
22. Koponen P, Helminen M, Paassilta M, Luukkaala T, Korppi M. Preschool asthma after bronchiolitis in infancy. *Eur Respir J* 2012;**39**(1):76-80.
23. Mikalsen IB, Halvorsen T, Oymar K. The outcome after severe bronchiolitis is related to gender and virus. *Pediatr Allergy Immunol* 2012;**23**(4):391-8.
24. Benjamin JT, Carver BJ, Plosa EJ, et al. NF-kappaB activation limits airway branching through inhibition of Sp1-mediated fibroblast growth factor-10 expression. *J Immunol* 2010;**185**(8):4896-903.
25. Han J, Takeda K, Gelfand EW. The Role of RSV Infection in Asthma Initiation and Progression: Findings in a Mouse Model. *Pulm Med* 2011;**2011**:748038.
26. Joss-Moore LA, Lane RH. Epigenetics and the developmental origins of disease: the key to unlocking the door of personalized medicine. *Epigenomics* 2012;**4**(5):471-3.
27. Hjalmarson O, Sandberg K. Abnormal lung function in healthy preterm infants. *Am J Respir Crit Care Med* 2002;**165**(1):83-7.
28. Friedrich L, Pitrez PM, Stein RT, Goldani M, Tepper R, Jones MH. Growth rate of lung function in healthy preterm infants. *Am J Respir Crit Care Med* 2007;**176**(12):1269-73.
29. Boyle EM, Poulsen G, Field DJ, et al. Effects of gestational age at birth on health outcomes at 3 and 5 years of age: population based cohort study. *BMJ* 2012;**344**:e896.
30. Kotecha SJ, Watkins WJ, Paranjothy S, Dunstan FD, Henderson AJ, Kotecha S. Effect of late preterm birth on longitudinal lung spirometry in school age children and adolescents. *Thorax* 2012;**67**(1):54-61.
31. Narang I, Rosenthal M, Cremonesini D, Silverman M, Bush A. Longitudinal evaluation of airway function 21 years after preterm birth. *Am J Respir Crit Care Med* 2008;**178**(1):74-80.
32. Crump C, Winkleby MA, Sundquist J, Sundquist K. Risk of asthma in young adults who were born preterm: a Swedish national cohort study. *Pediatrics* 2011;**127**(4):e913-20.
33. Siltanen M, Wehkalaampi K, Hovi P, et al. Preterm birth reduces the incidence of atopy in adulthood. *J Allergy Clin Immunol* 2011;**127**(4):935-42.

Table 1. Respiratory infections and prescribed asthma treatments

Respiratory infections during the first year of life	
Adenovirus infection	Pneumococcal pneumonia
Other specified Adenovirus infection	Pneumonia
Bordetella Pertussis	Pneumonia due to streptococcus
Bronchitis	Pneumonia due to Haemophilus influenzae
Bronchiolitis	Pneumonia due to respiratory syncytial virus
Bronchopneumonia organism unspecified	Other unspecified pneumonia
Coxsackie virus infection	Bacterial pneumonia unspecified
Cytomegalovirus	Other specified viral pneumonia
Echo virus infection	Pneumonia in other infectious diseases classified elsewhere
Influenza	Rhinovirus infection
Influenza with other respiratory manifestations	Other specified viral infections
Influenza unspecified	Suspected viral infections
Influenza with pleuropneumonia	Unspecified viral infections
Prescribed treatments	
Beclomethasone	Omalizumab
Budesonide	Salbutamol
Flutikasone	Salmeterol
Formoterol	Sodiumchromoglicate
Indakaterol	Terbutaline
Ipratropium	Theophylline
Mometasone	Tiotropium bromide
Montelukast	

Table 2. Characteristics of the cohorts defined by respiratory infection during the first year of life

	Respiratory infection in first year	No respiratory infection in first year	Total
	n= 42334 (16.7%)	n= 211594 (83.3%)	n= 253928 (100%)
Prescribed asthma medication*	7485 (17.7)	26591 (12.6)	34076 (13.4)
Hospital asthma diagnosis*	420 (1.0)	761 (0.4)	1181 (0.5)
No asthma*	34749 (82.1)	184775 (87.3)	219524 (86.5)
Gestational age (weeks)			
≤27	252 (0.6)	1221 (0.6)	1473 (0.6)
28-31	723 (1.7)	3578 (1.7)	4301 (1.7)
32-36	3906 (9.2)	19528 (9.2)	23434 (9.2)
37-41	34839 (82.3)	174194 (82.3)	209033 (82.3)
>41	2507 (5.9)	12535 (5.9)	15042 (5.9)
Uncertain	107 (0.3)	538 (0.3)	645 (0.3)
Birth weight			
<1000g	241(0.6)	1292 (0.6)	1533 (0.6)
> 1000-1500g	580 (1.4)	2714 (1.3)	3294 (1.3)
>1500-2000g	953 (2.3)	4300 (2.0)	5253 (2.1)
>2000-2750g	4099 (9.7)	18815 (8.9)	22914 (9.0)
>2750-3250g	9727 (23.0)	47617 (22.5)	57344 (22.6)
>3250-4000g	20178 (47.7)	104693 (49.5)	124871 (49.2)
> 4000g	6426 (15.2)	31524 (14.9)	37950 (14.9)
Not recorded	130 (0.3)	639 (0.3)	769 (0.3)
Maternal asthma			
Current	721 (1.7)	2754 (1.3)	3475 (1.4)
Previous	210 (0.5)	749 (0.4)	959 (0.4)
None	41403 (97.8)	208091 (98.3)	249494 (98.3)
Maternal age at delivery			
13-18	755 (1.8)	2917 (1.4)	3672 (1.5)
19-25	14230 (33.6)	66604 (31.5)	80834 (31.8)
26-30	15312 (36.2)	76752 (36.3)	92064 (36.3)
31-35	8911 (21.0)	46336 (21.9)	55247 (21.8)
36-40	2757 (6.5)	16673 (7.9)	19430 (7.7)
41-54	369 (0.9)	2312 (1.1)	2681 (1.1)
Maternal smoking			
Non-smoker	23871 (56.4)	134802 (63.7)	158673 (62.5)
Smoker	12429 (29.4)	46817 (22.1)	59246 (23.3)
Not recorded	6034 (14.3)	29975 (14.2)	36009 (14.2)
Child's sex			
Female	17619 (41.6)	103007 (48.7)	120626 (47.5)
Male	24715 (58.4)	108587 (51.3)	133302 (52.5)
Parity			
1	12714 (30.0)	87275 (41.2)	99989 (39.4)
2	17247 (40.7)	74534 (35.2)	91781 (36.1)
3	8270 (19.5)	34880 (16.5)	43150 (17.0)
4 or more	4103 (9.7)	14905 (7.0)	19008 (7.5)
Chronic lung disease			
No	42272 (99.9)	211416 (99.9)	253688 (99.9)
Yes	62 (0.1)	178 (0.1)	240 (0.1)

* These categories are not mutually exclusive.

Table 3. Early life characteristics and subsequent asthma risk after age 5 years

Characteristic	Asthma n (%)	No asthma n (%)	Unadjusted HR (95% CI)	P value	Adjusted ¹ HR (95% CI)	P value
Respiratory infection in first year						
Yes	7585 (17.9)	34749 (82.1)	1.45 (1.41 to 1.49)	<0.001	1.51 (1.47 to 1.54)	<0.001
No	26819 (12.7)	184775 (87.3)	Ref.		Reference	
Maternal asthma						
Current	921 (26.5)	2554 (73.5)	3.86 (3.61 to 4.12)	<0.001	1.94 (1.82 to 2.08)	<0.001
Previous	222 (23.1)	737 (76.9)	3.05 (2.67 to 3.48)	<0.001	1.38 (1.21 to 1.58)	<0.001
None	33261 (13.3)	216233 (86.7)	Ref.		Ref.	
Maternal age at delivery (years)						
13-18	517 (14.1)	3155 (85.9)	0.96 (0.88 to 1.05)	0.372	0.98 (0.90 to 1.07)	0.625
19-25	11451 (14.2)	69383 (85.8)	1.02 (1.00 to 1.05)	0.069	1.03 (1.00 to 1.06)	0.026
26-30	12249 (13.3)	79815 (86.7)	Ref.		Ref.	
31-35	7276 (13.2)	47971 (86.8)	0.99 (0.96 to 1.02)	0.490	1.02 (0.99 to 1.05)	0.140
36-40	2562 (13.2)	16868 (86.8)	1.00 (0.96 to 1.05)	0.830	1.05 (1.01 to 1.10)	0.029
41-54	349 (13.0)	2332 (87.0)	1.00 (0.90 to 1.12)	0.934	1.05 (0.94 to 1.17)	0.391
Maternal smoking						
Non-smoker	21701 (13.7)	136972 (86.3)	Ref.		Ref.	
Smoker	8278 (14.0)	50968 (86.0)	0.93 (0.91 to 0.96)	<0.001	1.01 (0.99 to 1.04)	0.380
Not recorded	4425 (12.3)	31584 (87.7)	0.59 (0.57 to 0.61)	<0.001	0.90 (0.87 to 0.93)	<0.001
Child's sex						
Female	18150 (15.0)	102476 (85.0)	1.25 (1.22 to 1.27)	<0.001	1.27 (1.24 to 1.29)	<0.001
Male	16254 (12.2)	117048 (87.8)	Ref.		Ref.	
Parity						
1	14371 (14.4)	85618 (85.6)	Ref.		Ref.	
2	12230 (13.3)	79551 (86.7)	0.93 (0.90 to 0.95)	<0.001	0.90 (0.88 to 0.93)	<0.001
3	5437 (12.6)	37713 (87.4)	0.87 (0.84 to 0.89)	<0.001	0.84 (0.82 to 0.87)	<0.001
4 or more	2366 (12.4)	16642 (87.6)	0.88 (0.84 to 0.92)	<0.001	0.81 (0.78 to 0.85)	<0.001
Chronic lung disease						
No	34360	219328	Ref.		Ref.	
Yes	44	196	1.59 (1.18 to 2.14)	0.002	1.44 (1.06 to 1.96)	0.019

¹ Adjusted for child's sex, maternal asthma, maternal age at delivery, year of birth, maternal smoking, gestational age, infection during first year of life, parity, and chronic lung disease

Table 4. Respiratory infection in the first year of life subsequent asthma risk after age 5 years, stratified by gestational age

Infection stratified by gestational age	Asthma n (%)	No asthma n (%)	HR (95% CI)	P Value	HR ¹ (95% CI)	P Value
Week 0-27						
Infection	52 (20.6)	200 (79.4)	2.03 (1.47 to 2.80)	< 0.001	2.22 (1.59 to 3.09)	< 0.001
No infection	130 (10.6)	1091 (89.4)	Ref.		Ref.	
Week 28-31						
Infection	132 (18.3)	591 (81.7)	1.47 (1.21 to 1.78)	< 0.001	1.57 (1.29 to 1.92)	< 0.001
No infection	458 (12.8)	3130 (87.2)	Ref.		Ref.	
Week 32-36						
Infection	738 (18.9)	3168 (81.1)	1.47 (1.35 to 1.59)	< 0.001	1.54 (1.42 to 1.68)	< 0.001
No infection	2588 (13.3)	16940 (86.7)	Ref.		Ref.	
Week 37-41						
Infection	6194 (17.8)	28645 (82.2)	1.44 (1.40 to 1.48)	< 0.001	1.49 (1.45 to 1.53)	< 0.001
No infection	22075 (12.7)	152119 (87.3)	Ref.		Ref.	
Week > 41						
Infection	446 (17.8)	2061 (82.2)	1.51 (1.36 to 1.68)	< 0.001	1.57 (1.41 to 1.75)	< 0.001
No infection	1512(12.1)	11023 (87.9)	Ref.		Ref.	
Week uncertain						
Infection	23 (21.5)	84 (78.5)	2.10 (1.29 to 3.42)	0.003	2.21 (1.34 to 3.65)	0.002
No infection	56 (10.4)	482 (89.6)	Ref.		Ref.	

¹ Adjusted for child's sex, maternal asthma, maternal age at delivery, year of birth, birth weight, maternal smoking, infection during first year of life, parity, and chronic lung disease

Table 5. Respiratory infection in the first year of life subsequent asthma risk after age 10 years and after age 16 years, stratified by gestational age

	Asthma n (%)	No asthma n (%)	HR (95% CI)	P Value	HR ¹ (95% CI)	P Value
≥ Age 10						
Gestational age (weeks)						
≤27						
Infection	45 (17.9)	207 (82.1)	1.83 (1.30 to 2.57)	0.001	1.99 (1.40 to 2.82)	< 0.001
No infection	123 (10.1)	1098 (89.9)	Ref.		Ref.	
28-31						
Infection	116 (16.0)	607 (84.0)	1.43 (1.17 to 1.76)	0.001	1.52 (1.24 to 1.87)	< 0.001
No infection	411 (11.5)	3167 (88.5)	Ref.		Ref.	
32-36						
Infection	635 (16.3)	3271 (83.7)	1.39 (1.27 to 1.52)	< 0.001	1.46 (1.33 to 1.59)	< 0.001
No infection	2328 (11.)	17200 (88.1)	Ref.		Ref.	
37- 41						
Infection	5541 (15.9)	29298 (84.1)	1.44 (1.40 to 1.48)	< 0.001	1.49 (1.45 to 1.54)	< 0.001
No infection	19637 (11.3)	154557 (88.7)	Ref.		Ref.	
> 41						
Infection	397 (15.8)	2119 (84.2)	1.51 (1.35 to 1.69)	< 0.001	1.57 (1.41 to 1.76)	< 0.001
No infection	1340 (10.7)	11195 (89.3)	Ref.		Ref.	
Uncertain						
Infection	19 (17.8)	88 (82.2)	1.93 (1.14 to 3.28)	0.015	2.05 (1.19 to 3.55)	0.010
No infection	50 (9.3)	488 (90.7)	Ref.		Ref.	
≥ Age 16						
Gestational age (weeks)						
≤27						
Infection	22 (8.8)	227 (91.2)	1.56 (0.96 to 2.52)	0.071	1.65 (1.01 to 2.71)	0.047
No infection	69 (5.7)	1132 (94.3)	Ref.		Ref.	
28-31						
Infection	65 (9.2)	638 (90.8)	1.41 (1.07 to 1.86)	0.014	1.45 (1.10 to 1.91)	0.009
No infection	233 (6.7)	3249 (93.3)	Ref.		Ref.	
32-36						
Infection	396 (10.5)	3362 (89.5)	1.43 (1.28 to 1.60)	< 0.001	1.49 (1.34 to 1.68)	< 0.001
No infection	1410 (7.5)	17379 (92.5)	Ref.		Ref.	
37- 41						
Infection	3227 (9.7)	30040 (90.3)	1.37 (1.32 to 1.42)	< 0.001	1.42 (1.37 to 1.48)	< 0.001
No infection	11987 (7.2)	154347 (92.8)	Ref.		Ref.	
> 41						
Infection	248 (10.3)	2161 (89.7)	1.58 (1.37 to 1.82)	< 0.001	1.64 (1.42 to 1.90)	< 0.001
No infection	801 (6.7)	11244 (93.3)	Ref.		Ref.	
Uncertain						
Infection	16 (15.2)	89 (84.8)	2.48 (1.37 to 4.51)	0.003	2.72 (1.47 to 5.04)	0.002
No infection	33 (6.2)	496 (93.8)	Ref.		Ref.	

¹ Adjusted for child's sex, maternal asthma, maternal age at delivery, year of birth, maternal smoking, birth weight, infection during first year of life, parity, and chronic lung disease.

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cohort studies*

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	3
Objectives	3	State specific objectives, including any prespecified hypotheses	3
Methods			
Study design	4	Present key elements of study design early in the paper	4
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	4-5
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	4-5
		(b) For matched studies, give matching criteria and number of exposed and unexposed	4
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	4-6
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	4-5
Bias	9	Describe any efforts to address potential sources of bias	4-6
Study size	10	Explain how the study size was arrived at	4
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	5-6, 16

Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	5-6
		(b) Describe any methods used to examine subgroups and interactions	5-6
		(c) Explain how missing data were addressed	7
		(d) If applicable, explain how loss to follow-up was addressed	5-6
		(e) Describe any sensitivity analyses	6-7
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	4
		(b) Give reasons for non-participation at each stage	4
		(c) Consider use of a flow diagram	
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	6-7, 16
		(b) Indicate number of participants with missing data for each variable of interest	5-7, 16
		(c) Summarise follow-up time (eg, average and total amount)	4
Outcome data	15*	Report numbers of outcome events or summary measures over time	
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	6-7, 16-19
		(b) Report category boundaries when continuous variables were categorized	16-17
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	5-7, 17-19

Discussion			
Key results	18	Summarise key results with reference to study objectives	7-8
Limitations			
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	8-11
Generalisability	21	Discuss the generalisability (external validity) of the study results	11
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	1

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.