



**Muscle Strength in Adolescent Men and Future
Musculoskeletal Pain:
A Cohort Study with 17 Years of Follow-up**

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-002656
Article Type:	Research
Date Submitted by the Author:	30-Jan-2013
Complete List of Authors:	Timpka, Simon; Lund University, Department of Orthopedics Pettersson, Ingemar; Lund University, Department of Orthopedics Zhou, Caddie; Lund University, Department of Orthopedics Englund, Martin; Lund University, Department of Orthopedics; Boston University School of Medicine, Clinical Epidemiology Research & Training Unit
Primary Subject Heading:	Epidemiology
Secondary Subject Heading:	Sports and exercise medicine, Rheumatology, Public health
Keywords:	Musculoskeletal disorders < ORTHOPAEDIC & TRAUMA SURGERY, SPORTS MEDICINE, PREVENTIVE MEDICINE

SCHOLARONE™
Manuscripts

Muscle Strength in Adolescent Men and Future Musculoskeletal Pain:

A Cohort Study with 17 Years of Follow-up

Simon Timpka¹, Ingemar F Petersson¹, Caddie Zhou¹, Martin Englund^{1, 2}

1 Department of Orthopedics, Clinical Sciences Lund, Lund University, Lund, Sweden

2 Clinical Epidemiology Research & Training Unit, Boston, University School of Medicine, Boston, Massachusetts, United States of America

Corresponding author:

Simon Timpka
Skånes universitetssjukhus
Epi-centrum Skåne
Medicon Village
223 81 Lund
Sweden

Phone: +46 738 395 276
Mail: simon.timpka@med.lu.se

Words in main text: 2800
Number of figures: 1
Number of tables: 2

Abstract

Objectives

Musculoskeletal pain is highly prevalent throughout adulthood with a major impact on health, function and participation in the society. Still, the association between muscle strength and development of musculoskeletal pain is unclear. We aimed to study whether overall muscle strength in adolescent men is associated with self-reported musculoskeletal pain in adulthood.

Design

Cohort study with baseline data from the Swedish Conscription Register and outcome information from the random population-based Swedish Living Conditions Surveys.

Setting

Sweden, 1970-2005.

Participants

We studied 5492 men who at age 17-19 years tested their isometric muscle strength (hand grip, arm flexion, knee extension) during the compulsory conscription.

Outcome measures

The men were surveyed regarding self-reported musculoskeletal pain; mean follow-up time of 17 (range 1-35) years. Our primary outcome was a self-report of musculoskeletal pain, and secondary outcomes were a report of "severe pain", "pain in back/hips", "pain in neck/shoulders", or "pain in arms/legs", respectively. We categorized muscle strength into three groups; low, average, and high using the 25th to 75th percentile to define the reference category (average). We calculated incidence rate ratios using log binomial regression with adjustment for smoking, body mass index, education, and physical activity.

Results

1
2
3 In the adjusted model, men with low overall muscle strength had decreased risk of self-reported
4 musculoskeletal pain (incidence rate ratio 0.93, 95% confidence interval 0.87-0.99). We observed no
5 such association in men with high strength (0.99, 0.93-1.05). Furthermore, no statistically significant
6 increase or decrease in risk was observed for any of the secondary outcomes.
7
8
9
10

11 **Conclusion**

12
13
14 In men, low overall isometric muscle strength in youth was not associated with an increased risk of
15 future musculoskeletal pain. Contrarily, we observed a slightly decreased risk of self-reported
16 musculoskeletal pain in adulthood. Residual confounding by adult occupational exposures, leisure
17 time physical activity level and psychosocial factors may have impacted on results.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article summary

Article focus

- We aimed to study whether overall muscle strength in youth is inversely associated with the development of musculoskeletal pain in adulthood.

Key messages

- In contradiction to our expectations, we observed a decreased risk of self-reported musculoskeletal pain in adult men with low overall isometric muscle strength in youth.
- Low muscle strength may potentially serve as a deselection criterion for activities with high risk of acute injuries or chronic physical overload, factors with negative impact on musculoskeletal health.

Strengths and limitations of study

- The main strengths of the study are a large sample, the use of three different measures of muscle strength, and comparably long time to follow-up.
- The main limitations of the study are the following; the cohort does not include women, musculoskeletal pain only identified with one question per site, motivation for military service might influence measurement of muscle strength and the potential for unmeasured or residual confounding.

Introduction

Musculoskeletal disorders (MSDs), such as low back pain, osteoarthritis, and widespread pain, are highly prevalent in the adult population.¹⁻³ MSDs also contribute to a substantial burden of disease at middle and older ages.⁴ Although pain emanating from the musculoskeletal system might be attributed to a wide range of diseases with diverse causal chains, many MSDs have common risk factors such as heavy occupational work load,^{5,6} a high body mass index (BMI),⁷⁻⁹ and a low socio-economy.¹⁰⁻¹² Although smoking in some studies have been identified as a risk factor for certain MSDs,^{13,14} its main effect on musculoskeletal pain might be as an effect modifier of the pain sensation.¹⁵ As physical work load is a risk factor for many MSDs,¹⁶ a model in which muscle strength in the loaded parts of the body are protective for future disorders is appealing. Furthermore, physical exercise with focus on muscle strength is an important secondary and tertiary prevention of MSDs.¹⁷

¹⁸ A handful of studies have hitherto longitudinally investigated the strength of isolated muscle groups in adulthood as a determinant of later MSDs.¹⁹⁻²² However, in adult subjects, there is for the time being conflicting evidence of the value of muscle strength as a protective factor of musculoskeletal pain, such as neck/shoulder pain and low back pain.¹⁹

In the longer perspective, relatively little is known about the association of muscle strength in youth and later disease, including musculoskeletal pain.²³ Two studies have investigated the result of single muscle strength tests as determinants of musculoskeletal complaints decades later.^{24,25} The first, using number of sit-ups during 30 seconds as a strength measure, found no association with later low back pain or tension neck in men. In women, the high strength group had a decreased odds ratio (OR) of tension neck.²⁴ The second study found a decreased OR for MSDs in men who either had a strong performance in isotonic bench press or in a isometric two hand lift test.²⁵ Neither of the two studies includes a measure of overall muscular capacity. In summary, although there is some evidence of an association between low muscle strength in youth and later risk of MSDs, the association between overall muscle strength in adolescence and later musculoskeletal pain has not

1
2
3 been studied. Furthermore, with a larger sample size, data on common risk factors, testing of three
4
5 different muscle groups, and data on physical work capacity, we address some of the limitations of
6
7 earlier studies.
8
9

10 Our aim in this study was to investigate the general muscle strength in adolescent males as a
11
12 determinant of later self-reported musculoskeletal pain. We hypothesized that low general muscle
13
14 strength in youth is associated with an increased risk of having musculoskeletal pain in adulthood.
15
16

17 **Methods**

18
19 We used two main two main criteria to identify the study cohort. First, the subjects should have
20
21 performed mandatory conscription testing in Sweden between 1970-1994, with the exception of the
22
23 years 1978 and 1985. Secondly, they should have been included in the Swedish Living Conditions
24
25 Surveys any year between 1980 and 2005 when questions regarding musculoskeletal problems,
26
27 smoking status, and physical activity were simultaneously included.
28
29
30

31
32 Furthermore, we excluded all men who were surveyed prior to the baseline testing or were younger
33
34 than 17 years or older than 19 years at baseline (Table 1). We also excluded men with an existing
35
36 musculoskeletal disorder (Diseases of the musculoskeletal system and connective tissue according to
37
38 the International classification of disease version 8 or 9) and those who had missing data on variables
39
40 included in the primary model (muscle strength, smoking, BMI, physical activity, level of education).
41
42 In the final study sample we included 5492 men (figure 1). Data from the Swedish conscription
43
44 testing has been previously used for research purposes.²⁶⁻²⁸ During the period of the study sample
45
46 testing, conscription was mandatory by law for all Swedish men. Specially trained employees at six
47
48 regional conscription centers administrated the conscription tests during a two-day session that also
49
50 included separate evaluations by a medical doctor and a psychologist. Only men with serious health
51
52 complaints, were excused from conscription. The procedure included measurements of each
53
54 subject's weight in underwear to the kilogram and height without shoes to the centimeter. Using
55
56
57
58
59
60

1
2
3 height and weight, we calculated BMI as height/kg². Probably due to rare errors of data entry, there
4
5 are unlikely extreme values in the dataset. Therefore, we excluded all subjects with registered
6
7 extreme values on height (<150, >210 cm), weight (<40, >150 kg), or an extreme calculated value for
8
9 BMI (<15, >60) (figure 1). The study was approved by the Ethical Review Board at Lund University
10
11 and the manuscript was prepared according to the STROBE-statement.²⁹

13 *Muscle strength*

14
15 The men performed three tests of isometric muscle strength during conscription; hand grip, elbow
16
17 flexion, and knee extension. At the start of test period in 1970, the tests were performed as
18
19 previously described³⁰ and remained unchanged in general throughout the test period. In summary,
20
21 hand grip was measured with a 90° flexion at the elbow with the humerus in parallel to the torso.
22
23 Knee extension was measured in a sitting position with 90° knee flexion and arms crossed over chest.
24
25 The pelvis was fixed to the seat and a strap fastened above the lateral malleolus. Also, elbow flexion
26
27 was measured in a sitting position with 90° flexion at the elbow and the humerus in parallel to the
28
29 torso. A strap was fastened at the level of the radial styloid process.

30
31 We calculated a measure of general muscle strength by standardizing and combining the three tests
32
33 of muscle strength. To avoid bias due to change of testing procedure over time, we categorized the
34
35 cohort into five subgroups based on period of conscription (1970-1973, 1974-1977, 1979-1984, 1986-
36
37 1990, 1990-1994) and for each subgroup calculated the relative muscle strength. We standardized
38
39 the three tests of muscle strength [standardized value= (value–mean)/standard deviation] within
40
41 each subgroup and used the mean of the three test scores as a proxy for general muscle strength.
42
43 Using percentiles, we then categorized the cohort into three groups of muscle strength, where the
44
45 25th to 75th percentile defined the average category, the bottom 25th percentile configured the low
46
47 category, and the top 25th percentile defined the high muscle strength category.

52 **Survey of musculoskeletal pain**

53
54 The Swedish Living Conditions Surveys (ULF) is a random population based survey conducted by
55
56 Statistics Sweden, previously used for research purposes.³¹⁻³³ For the present study, we used data
57
58
59
60

1
2
3 collected during a total of 10 years (1980, 1988, 1989, 1997-1999, 2002-2005). The surveys were
4
5 generally performed as interviews in person by trained interviewers, with a minority of the
6
7 interviews performed by phone. For men included in more than one survey, we used the last survey
8
9 without relevant study data missing.

10
11 At follow-up, the men were asked three questions regarding any current musculoskeletal pain: 1) Do
12
13 you have pain in neck or shoulders? 2) Do you have back-pain, hip-pain or sciatica? 3) Do you have
14
15 ache, pain in hands, elbows, legs or knees? For each type of complaint one of three answers was
16
17 possible: 1) Yes, severe 2) Yes, mild and 3) No. Our primary outcome was having reported either
18
19 severe or mild musculoskeletal pain, whereas our secondary outcomes we defined as follows: 1)
20
21 Having reported severe musculoskeletal pain 2) Having reported pain in back/hips 3) Having reported
22
23 pain in shoulders/neck 4) Having reported pain in arms/legs. From the surveys, we also included data
24
25 on self-reported current smoking status (yes/no), physical activity (practically none, now and then,
26
27 regularly, regularly strenuous), and level of education (compulsory school or less, secondary
28
29 education, higher education). Drop-outs from the survey, i.e. those who have declined participation,
30
31 cannot be individually identified. However, of the men who were asked to be included in the relevant
32
33 age-groups, 11.3-31.0% declined participation in the years of study. The mean rate of non-
34
35 participation over the relevant years in the same group was 22.9%.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Statistical analyses

All analyses were performed with SAS 9.3 (SAS Institute Inc). We used logistic binomial regression to estimate incidence rate ratios (IRR) and control potential confounders. In the multivariate model (primary model), we included muscle strength, BMI, smoking status, physical activity and level of education.

To test whether cardiovascular aspects of physical capacity confounded our results, we used physical work capacity measured as $W_{\max 6 \text{ min}}$ in a sensitivity analysis. For $W_{\max 6 \text{ min}}$, the test result is an estimate of maximum work sustainable for 6 minutes³⁴ and is in young men correlated with maximum oxygen uptake ($r=0.9$).^{35 36} Acceptable data quality on work capacity was available in the subsample of men performing the baseline testing in 1976-1982. Out of all men in the cohort conscripted during the time period, 1154 men (74.6 %) completed an acceptable physical work capacity test on a bicycle ergometer (i.e. heart rate >174 at the end of testing). We added the work capacity in relation to body weight as a continuous variable to the univariate model. Furthermore, we also performed two sensitivity analyses on the multivariate model with musculoskeletal pain as the dependent variable. In the first, we added test center to the model. For the second, to test whether our categorization of muscle strength influenced the results, we 1) treated the standardized muscle strength as a continuous variable 2) treated the standardized muscle strength as a categorical variable based on quintiles.

Results

The mean time to follow-up was 17 years (table 1). Men with low muscle strength did not have an increased risk, but rather a statistically significant decreased risk, for the primary outcome "Musculoskeletal pain" (table 2). To summarize the observations of the secondary outcomes, we did not observe any statistically significant risk increases for neither men with a low nor high muscle strength. Compared to the crude model, the multivariate model produced similar risk estimates.

Sensitivity analyses

Work capacity had a significant effect in the subsample analysis ($p=0.04$) whereas it had only minor effect on the risk estimates for a musculoskeletal problem, being 0.94 (0.82-1.08) and 1.02 (0.90-1.16) for low and high strength, respectively. The pattern of association in the secondary outcomes were in general somewhat strengthened when we adjusted for work capacity (data not shown).

Using muscle strength as a continuous variable (hence assuming a linear relationship) did weaken the association with later musculoskeletal pain ($p=0.22$). When we instead used quintiles to categorize muscle strength, we observed no increased risk for the group with lowest strength compared to average strength (IRR=0.93, 95% confidence interval 0.85-1.01).

Discussion

Investigating the overall isometric muscle strength in adolescent men as a determinant of future musculoskeletal pain, we observed a decreased risk of self-reported musculoskeletal pain in men with low muscle strength. No such association was observed for men with high strength. We also found a similar, however not statistically significant, pattern for "pain in back/hips" and "pain in neck/shoulders", whereas no association was found for future problems in arms/legs. Noticeably the observed associations were in contradiction to our expectations.

1
2
3 Using a historical cohort design with prospective registration of exposure and the outcome, our study
4 includes a large sample and thus allows better control for known confounders compared to previous
5 studies. All covariates (muscle strength, BMI, smoking, education, physical activity) included in the
6 multivariate model for musculoskeletal pain also had a significant association with the outcome.
7

8
9
10
11 Furthermore, by controlling for work capacity, we aimed to isolate the direct effect of muscle
12 strength from other aspects of physical capacity. However, the study also has important limitations.

13
14 First and foremost, we have used strength data from military conscription testing. Although it
15 provides a rich dataset from a structured environment, we do not know how the subject's motivation
16 for military service may have biased the performance during the testing procedure. However,
17 assuming there is no association between motivation at conscription testing and later risk of
18 musculoskeletal pain (or the loss to follow-up) any bias would at most dilute our result. Nevertheless,
19 stronger recruits are more likely to be assigned to positions with heavy load duty. Secondly, the
20 conscription was mandatory for men only. As the pattern of physical activity and occupational
21 exposure differ between men and women, any generalization of the results to women must be made
22 with great caution. Third, the physical activity measurement consisted of a single question and did
23 neither allow calculation of metabolic equivalent of task (MET) nor included occupational exposure.
24 Also musculoskeletal pain is measured by questions that combine more than one site, decreasing the
25 precision. However, the categories are fairly well demarcated anatomically save for the question
26 regarding pain in arms/legs. Fourth, the covariates collected with musculoskeletal pain at follow-up
27 (smoking, physical activity, level of education) are cross-sectional and might thus be mediators of
28 reverse causation. However, the adjusted estimates are much in line with the crude estimates.
29

30
31 It has been suggested that there is a U-shaped association between physical activity and later back
32 pain.^{37 38} Furthermore, as former occupational exposure and certain sport participation are
33 established risk factors for future MSDs,^{16 39} it lends some evidence for a more general model, in
34 which certain forms of physical activity is negative for the musculoskeletal health. Primarily, our
35 observations do not support low muscle strength in youth as a risk factor for later musculoskeletal
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 pain. Instead, we suggest that our results can be explained by muscle strength in youth being one
4
5 selection criterion for future high risk activities with a negative influence on the musculoskeletal
6
7 health, e.g. higher risk of joint injury due to sports participation or manual repetitive work load. This
8
9 would also include more immediate exposure such as more physically demanding military service.⁴⁰
10
11 Although we have controlled for level of education, which might serve as a proxy for occupational
12
13 exposure, there is potential for residual confounding as we did not have more appropriate data. In
14
15 other words, individuals with low general muscle strength might to a certain degree be deselected
16
17 for high risk activities compared to men with an average or high strength. However, the strength of
18
19 an individual is associated with the muscle fiber type distribution, which have a large genetic
20
21 component.⁴¹ Type I fibers are more common in endurance athletes⁴² whereas high type II
22
23 percentage have been reported to be associated with isometric muscle strength⁴³ as well as low back
24
25 pain.⁴⁴ Thus, we cannot exclude that the decreased risk observed in men with low strength is
26
27 mediated by factors related to muscle fiber type.
28
29

30
31 Partly in contrast with a previous study,²⁵ we did not observe a negative effect of low muscle
32
33 strength on the risk future of musculoskeletal problems in men. Although we in the present study
34
35 only include measurements of isometric strength, the previous study observed associations with
36
37 both an isometric strength measure (static two hand lift) and an isotonic strength measure (bench
38
39 press). In a study on the same cohort, it is reported that the result in bench press, but not two hand
40
41 lift, was associated with both future cardiovascular fitness and future physical activity,⁴⁵ potentially
42
43 explaining part of the difference. As another study reported flexibility as a sit and reach test,²⁴ but
44
45 not strength measured as sit-ups, to be negatively associated with future risk of back pain, it is
46
47 possible that other aspects of muscular and musculoskeletal function is of greater importance of
48
49 future risk of MSDs than isometric muscle strength.
50
51

52
53 The association between muscle strength and later musculoskeletal pain diminished when we used
54
55 muscle strength as a continuous variable. This was not surprising, as the observed association was
56
57
58
59
60

1
2
3 non linear in the primary model. It is to be expected that the test centers differ somewhat in their
4 reported test results, as they were assigned adolescents based on geography. However, including
5 test centers in the model did not have any effect on the overall association. When we included work
6 capacity in the model, most risk estimates decreased in absolute values, furthering strengthening the
7 observations in the primary model. By using the relative muscle strength during testing periods of
8 five years, we partially address the potential systematic change in testing procedure over the years.
9 Although the methods of measurements have not changed at large, minor adjustments cannot be
10 excluded.
11
12

13
14
15
16
17
18
19
20
21 Although we found no increased risk of future musculoskeletal problems in men with low muscle
22 strength in adolescence, future studies need to better quantify the occupational exposure and leisure
23 time physical activity. Since the physical activity pattern and physical fitness profiles differ between
24 men and women, further investigations are also needed to investigate if similar associations can be
25 found in women.
26
27
28
29
30
31

32
33 We observed a decreased risk of self-reported musculoskeletal pain in adult men with low overall
34 isometric muscle strength in youth. We hypothesize that low muscle strength *per se* is not protective
35 of future musculoskeletal pain. Instead, low muscle strength might serve as a deselection criterion
36 for professions or types of leisure time physical activity with higher risk of acute injuries or chronic
37 physical overload, factors with negative impact on musculoskeletal health.
38
39
40
41
42
43

44 **Acknowledgements**

45
46
47 No author has a conflict of interest to report. The authors would like to thank Charlotte Bergknut for
48 the initial data management.
49
50
51
52
53
54
55
56
57
58
59
60

Funding

The authors are supported by grants from The Swedish Research Council, Skåne Regional Council, and the Faculty of Medicine at Lund University. The funders neither influenced the study process nor manuscript preparation.

Author contributions

ST proposed the study idea, collected data, designed the study, and drafted the manuscript. IP designed the study and provided input during manuscript preparation. CZ designed the study and performed the statistical analyses. ME designed the study, collected data, and provided input during manuscript preparation. All authors approved the final version of the manuscript.

Data sharing statement

No additional data to share.

Competing Interests

None

References

1. Bergman S, Herrstrom P, Hogstrom K, Petersson IF, Svensson B, Jacobsson LT. Chronic musculoskeletal pain, prevalence rates, and sociodemographic associations in a Swedish population study. *J Rheumatol* 2001;28(6):1369-77.
2. Picavet HS, Schouten JS. Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study. *Pain* 2003;102(1-2):167-78.
3. Urwin M, Symmons D, Allison T, Brammah T, Busby H, Roxby M, et al. Estimating the burden of musculoskeletal disorders in the community: the comparative prevalence of symptoms at

- 1
2
3 different anatomical sites, and the relation to social deprivation. *Ann Rheum Dis*
4
5 1998;57(11):649-55.
6
7 4. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years
8
9 (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the
10
11 Global Burden of Disease Study 2010. *Lancet* 2013;380(9859):2197-223.
12
13
14 5. van der Windt DA, Thomas E, Pope DP, de Winter AF, Macfarlane GJ, Bouter LM, et al.
15
16 Occupational risk factors for shoulder pain: a systematic review. *Occup Environ Med*
17
18 2000;57(7):433-42.
19
20
21 6. Hoogendoorn WE, van Poppel MN, Bongers PM, Koes BW, Bouter LM. Physical load during work
22
23 and leisure time as risk factors for back pain. *Scand J Work Environ Health* 1999;25(5):387-
24
25 403.
26
27 7. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, et al. Risk factors for incident
28
29 radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum*
30
31 1997;40(4):728-33.
32
33
34 8. Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between obesity
35
36 and low back pain: a meta-analysis. *Am J Epidemiol* 2010;171(2):135-54.
37
38
39 9. Grotle M, Hagen KB, Natvig B, Dahl FA, Kvien TK. Obesity and osteoarthritis in knee, hip and/or
40
41 hand: an epidemiological study in the general population with 10 years follow-up. *BMC*
42
43 *Musculoskelet Disord* 2008;9:132.
44
45
46 10. Hagen K, Zwart JA, Svebak S, Bovim G, Jacob Stovner L. Low socioeconomic status is associated
47
48 with chronic musculoskeletal complaints among 46,901 adults in Norway. *Scand J Public*
49
50 *Health* 2005;33(4):268-75.
51
52
53 11. Dalstra JA, Kunst AE, Borrell C, Breeze E, Cambois E, Costa G, et al. Socioeconomic differences in
54
55 the prevalence of common chronic diseases: an overview of eight European countries. *Int J*
56
57 *Epidemiol* 2005;34(2):316-26.
58
59
60

- 1
2
3 12. Bengtsson C, Nordmark B, Klareskog L, Lundberg I, Alfredsson L. Socioeconomic status and the
4 risk of developing rheumatoid arthritis: results from the Swedish EIRA study. *Ann Rheum Dis*
5 2005;64(11):1588-94.
6
7
8
9 13. Palmer KT, Syddall H, Cooper C, Coggon D. Smoking and musculoskeletal disorders: findings from
10 a British national survey. *Ann Rheum Dis* 2003;62(1):33-6.
11
12
13 14. Holth HS, Werpen HK, Zwart JA, Hagen K. Physical inactivity is associated with chronic
14 musculoskeletal complaints 11 years later: results from the Nord-Trondelag Health Study.
15 *BMC Musculoskelet Disord* 2008;9:159.
16
17
18 15. McBeth J, Jones K. Epidemiology of chronic musculoskeletal pain. *Best Pract Res Clin Rheumatol*
19 2007;21(3):403-25.
20
21
22 16. Roquelaure Y, Ha C, Rouillon C, Fouquet N, Leclerc A, Descatha A, et al. Risk factors for upper-
23 extremity musculoskeletal disorders in the working population. *Arthritis Rheum*
24 2009;61(10):1425-34.
25
26
27 17. Hakkinen A, Sokka T, Kotaniemi A, Hannonen P. A randomized two-year study of the effects of
28 dynamic strength training on muscle strength, disease activity, functional capacity, and bone
29 mineral density in early rheumatoid arthritis. *Arthritis Rheum* 2001;44(3):515-22.
30
31
32 18. Baker KR, Nelson ME, Felson DT, Layne JE, Sarno R, Roubenoff R. The efficacy of home based
33 progressive strength training in older adults with knee osteoarthritis: a randomized
34 controlled trial. *J Rheumatol* 2001;28(7):1655-65.
35
36
37 19. Hamberg-van Reenen HH, Ariens GA, Blatter BM, van Mechelen W, Bongers PM. A systematic
38 review of the relation between physical capacity and future low back and neck/shoulder
39 pain. *Pain* 2007;130(1-2):93-107.
40
41
42 20. Leino P, Aro S, Hasan J. Trunk muscle function and low back disorders: a ten-year follow-up study.
43 *J Chronic Dis* 1987;40(4):289-96.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 21. Kujala UM, Taimela S, Viljanen T, Jutila H, Viitasalo JT, Videman T, et al. Physical loading and
4 performance as predictors of back pain in healthy adults. A 5-year prospective study. *Eur J*
5 *Appl Physiol Occup Physiol* 1996;73(5):452-8.
6
7
8
9
10 22. Faber A, Sell L, Hansen JV, Burr H, Lund T, Holtermann A, et al. Does muscle strength predict
11 future musculoskeletal disorders and sickness absence? *Occupational medicine (Oxford,*
12 *England)* 2012;62(1):41-6.
13
14
15
16 23. Ruiz JR, Castro-Pinero J, Artero EG, Ortega FB, Sjostrom M, Suni J, et al. Predictive validity of
17 health-related fitness in youth: a systematic review. *Br J Sports Med* 2009;43(12):909-23.
18
19
20 24. Mikkelsen LO, Nupponen H, Kaprio J, Kautiainen H, Mikkelsen M, Kujala UM. Adolescent
21 flexibility, endurance strength, and physical activity as predictors of adult tension neck, low
22 back pain, and knee injury: a 25 year follow up study. *Br J Sports Med* 2006;40(2):107-13.
23
24
25
26 25. Barnekow-Bergkvist M, Hedberg GE, Janlert U, Jansson E. Determinants of self-reported neck-
27 shoulder and low back symptoms in a general population. *Spine (Phila Pa 1976)*
28 1998;23(2):235-43.
29
30
31
32 26. Neovius M, Sundstrom J, Rasmussen F. Combined effects of overweight and smoking in late
33 adolescence on subsequent mortality: nationwide cohort study. *BMJ* 2009;338:b496.
34
35
36 27. Nilsson PM, Nyberg P, Ostergren PO. Increased susceptibility to stress at a psychological
37 assessment of stress tolerance is associated with impaired fetal growth. *Int J Epidemiol*
38 2001;30(1):75-80.
39
40
41
42 28. Hemmingsson T, Melin B, Allebeck P, Lundberg I. The association between cognitive ability
43 measured at ages 18-20 and mortality during 30 years of follow-up--a prospective
44 observational study among Swedish males born 1949-51. *Int J Epidemiol* 2006;35(3):665-70.
45
46
47
48 29. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. Strengthening the
49 Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for
50 reporting observational studies. *BMJ* 2007;335(7624):806-8.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 30. Nordesjo LO, Schéle R. Validity of Ergometer Cycle Test and Measures of Isometric Muscle
4 Strength when Predicting some Aspects of Military Performance. *Swedish J Defence Med*
5 1974(10):11-23.
6
7
8
9
10 31. Wolk A, Rossner S. Effects of smoking and physical activity on body weight: developments in
11 Sweden between 1980 and 1989. *J Intern Med* 1995;237(3):287-91.
12
13 32. Sundquist J, Johansson SE. The influence of socioeconomic status, ethnicity and lifestyle on body
14 mass index in a longitudinal study. *Int J Epidemiol* 1998;27(1):57-63.
15
16 33. Bygren LO, Konlaan BB, Johansson SE. Attendance at cultural events, reading books or
17 periodicals, and making music or singing in a choir as determinants for survival: Swedish
18 interview survey of living conditions. *BMJ* 1996;313(7072):1577-80.
19
20 34. Tornvall G. Assessment of physical capabilities. *Acta physiologica Scandinavica* 1963;58(Suppl
21 201):1-102.
22
23 35. Nordesjo LO. A comparison between the Thornvall maximal ergometer test, submaximal
24 ergometer tests and maximal oxygen uptake. *Swedish J Defence Med* 1974(10):3-10.
25
26 36. Nordesjo LO. Estimation of the maximal work rate sustainable for 6 minutes using a single-level
27 load or stepwise increasing loads. *Ups J Med Sci* 1974;79(1):45-50.
28
29 37. Heneweer H, Vanhees L, Picavet HS. Physical activity and low back pain: a U-shaped relation?
30 *Pain* 2009;143(1-2):21-5.
31
32 38. Heneweer H, Picavet HS, Staes F, Kiers H, Vanhees L. Physical fitness, rather than self-reported
33 physical activities, is more strongly associated with low back pain: evidence from a working
34 population. *Eur Spine J* 2012;21(7):1265-72.
35
36 39. Kujala UM, Kettunen J, Paananen H, Aalto T, Battie MC, Impivaara O, et al. Knee osteoarthritis in
37 former runners, soccer players, weight lifters, and shooters. *Arthritis Rheum* 1995;38(4):539-
38 46.
39
40 40. Almeida SA, Williams KM, Shaffer RA, Brodine SK. Epidemiological patterns of musculoskeletal
41 injuries and physical training. *Med Sci Sports Exerc* 1999;31(8):1176-82.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 41. Simoneau JA, Bouchard C. Genetic determinism of fiber type proportion in human skeletal
4 muscle. *FASEB J* 1995;9(11):1091-5.
5
6
7 42. Costill DL, Daniels J, Evans W, Fink W, Krahenbuhl G, Saltin B. Skeletal muscle enzymes and fiber
8 composition in male and female track athletes. *J Appl Physiol* 1976;40(2):149-54.
9
10
11 43. Tesch P, Karlsson J. Isometric strength performance and muscle fibre type distribution in man.
12 *Acta physiologica Scandinavica* 1978;103(1):47-51.
13
14
15 44. Mannion AF, Weber BR, Dvorak J, Grob D, Muntener M. Fibre type characteristics of the lumbar
16 paraspinal muscles in normal healthy subjects and in patients with low back pain. *J Orthop*
17 *Res* 1997;15(6):881-7.
18
19
20
21 45. Barnekow-Bergkvist M, Hedberg G, Janlert U, Jansson E. Prediction of physical fitness and
22 physical activity level in adulthood by physical performance and physical activity in
23 adolescence--an 18-year follow-up study. *Scand J Med Sci Sports* 1998;8(5 Pt 1):299-308.
24
25
26
27
28
29
30
31
32

33 Tables and Figures

34
35 **Table 1:** Description of study sample

36
37 **Table 2:** Risk estimates for the muscle strength in youth as a determinant of self-reported
38 musculoskeletal pain in adulthood
39

40
41 **Figure 1:** The identification of the study sample and the loss to follow-up
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Description of study sample

Mean age at baseline (SD)	18.2 (0.5)
Mean time to follow-up (SD, range)	17.2 years (8.4, 1-35)
BMI (%)	
<18.5	477 (8.7)
18.5-24.9	4500 (81.9)
25-29.9	448 (8.2)
>30	67 (1.2)
Muscle Strength* (%)	
Low	1371 (25.0)
Average	2745 (50.0)
High	1371 (25.0)
Type of interview (%)	
In person	4351 (79.2)
By telephone	1141 (20.8)
Pain in back/hips (%)	
Yes	1647 (30.0)
Yes, severe	321 (5.8)
No	3843 (70.0)
Missing	2 (0.0)
Pain in neck/shoulders (%)	
Yes	1564 (28.5)
Yes, severe	247 (4.5)
No	3926 (71.5)
Missing	2 (0.0)
Pain in arms/legs (%)	
Yes	1243 (22.6)
Yes, severe	196 (3.6)
No	4246 (77.4)
Missing	3 (0.0)
Pain, independent of location (%)	
Yes	2850 (51.9)
Yes, severe	576 (10.0)
No	2639 (48.1)
Missing	3 (0.0)
Smoking status (%)	
Yes	827 (15.1)
No	4665 (84.9)
Level of education (%)	
Compulsory	582 (10.6)
Secondary	2891 (52.7)
Higher	2011 (36.7)
Physical Activity (%)	
Practically non	584 (10.6)
Now and then	1615 (29.4)
Regularly	2077 (37.8)
Regularly strenuous	1216 (22.2)

*= categorization based on quartiles

Table 2 Risk estimates for the muscle strength in youth as a determinant of self-reported musculoskeletal pain in adulthood

Outcomes	Models				
	Univariate Muscle strength			Multivariate ^a Muscle strength	
	Low (N=1371) IRR (N)	Average (N=2745) Reference (N)	High (N=1371) IRR (N)	Low IRR (95% CI)	High IRR (95% CI)
Musculoskeletal pain	0.92 (668)	1 (1457)	0.99 (722)	0.93 (0.87-0.99)	0.99 (0.93-1.05)
Severe musculoskeletal pain	0.96 (135)	1 (283)	1.12 (158)	0.96 (0.79-1.17)	1.08 (0.90-1.30)
Pain in back/hips	0.92 (384)	1 (832)	1.03 (429)	0.93 (0.84-1.04)	1.03 (0.94-1.14)
Pain in neck/shoulders	0.91 (366)	1 (799)	0.99 (397)	0.92 (0.83-1.03)	1.00 (0.90-1.10)
Pain in arms/legs	0.97 (297)	1 (616)	1.07 (330)	0.97 (0.85-1.10)	1.06 (0.94-1.19)

N= Number of cases

IRR= Incidence rate ratio

CI= Confidence interval

a= adjusted for smoking status, physical activity, education, body mass index




Figure 1: The identification of the study sample and the loss to follow-up
254x190mm (96 x 96 DPI)

Review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of cohort studies

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5-6
Objectives	3	State specific objectives, including any prespecified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	6+
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6-9
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	6-7
		(b) For matched studies, give matching criteria and number of exposed and unexposed	-
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6-9
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	6-9
Bias	9	Describe any efforts to address potential sources of bias	9
Study size	10	Explain how the study size was arrived at	Figure 1
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	6-9
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	9
		(b) Describe any methods used to examine subgroups and interactions	9
		(c) Explain how missing data were addressed	-
		(d) If applicable, explain how loss to follow-up was addressed	-
		(e) Describe any sensitivity analyses	9
Results			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Figure 1
		(b) Give reasons for non-participation at each stage	Figure 1
		(c) Consider use of a flow diagram	Figure 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Table 1
		(b) Indicate number of participants with missing data for each variable of interest	Table 1
		(c) Summarise follow-up time (eg, average and total amount)	Table 1
Outcome data	15*	Report numbers of outcome events or summary measures over time	Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 1
		(b) Report category boundaries when continuous variables were categorized	6-9
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	-
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	9
Discussion			
Key results	18	Summarise key results with reference to study objectives	10-11
Limitations			
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	13-14
Generalisability	21	Discuss the generalisability (external validity) of the study results	12-13
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>



**Muscle Strength in Adolescent Men and Future
Musculoskeletal Pain:
A Cohort Study with 17 Years of Follow-up**

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-002656.R1
Article Type:	Research
Date Submitted by the Author:	01-Mar-2013
Complete List of Authors:	Timpka, Simon; Lund University, Department of Orthopedics Pettersson, Ingemar; Lund University, Department of Orthopedics Zhou, Caddie; Lund University, Department of Orthopedics Englund, Martin; Lund University, Department of Orthopedics; Boston University School of Medicine, Clinical Epidemiology Research & Training Unit
Primary Subject Heading:	Epidemiology
Secondary Subject Heading:	Sports and exercise medicine, Rheumatology, Public health
Keywords:	Musculoskeletal disorders < ORTHOPAEDIC & TRAUMA SURGERY, SPORTS MEDICINE, PREVENTIVE MEDICINE

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Muscle Strength in Adolescent Men and Future Musculoskeletal Pain: A Cohort Study with 17 Years of Follow-up

Simon Timpka¹, Ingemar F Petersson¹, Caddie Zhou¹, Martin Englund^{1, 2}

1 Department of Orthopedics, Clinical Sciences Lund, Lund University, Lund, Sweden

2 Clinical Epidemiology Research & Training Unit, Boston University School of Medicine, Boston, Massachusetts, United States of America

Corresponding author:

Simon Timpka
Skånes universitetssjukhus
Epi-centrum Skåne
Medicon Village
223 81 Lund
Sweden

Phone: +46 738 395 276
Mail: simon.timpka@med.lu.se

Words in main text: 2800
Number of figures: 1
Number of tables: 2

Abstract

Objectives

Musculoskeletal pain is highly prevalent throughout adulthood with a major impact on health, function and participation in the society. Still, the association between muscle strength and development of musculoskeletal pain is unclear. We aimed to study whether overall muscle strength in adolescent men is associated with self-reported musculoskeletal pain in adulthood.

Design

Cohort study with baseline data from the Swedish Conscription Register and outcome information from the random population-based Swedish Living Conditions Surveys.

Setting

Sweden, 1970-2005.

Participants

We studied 5489 men who at age 17-19 years tested their isometric muscle strength (hand grip, arm flexion, knee extension) during the compulsory conscription.

Outcome measures

The men were surveyed regarding self-reported musculoskeletal pain; mean follow-up time of 17 (range 1-35) years. Our primary outcome was a self-report of musculoskeletal pain, and secondary outcomes were a report of "severe pain", "pain in back/hips", "pain in neck/shoulders", or "pain in arms/legs", respectively. We categorized muscle strength into three groups; low, average, and high using the 25th to 75th percentile to define the reference category (average). We estimated relative risks using log binomial regression with adjustment for smoking, body mass index, education, and physical activity.

Results

In the adjusted model, men with low overall muscle strength had decreased risk of self-reported musculoskeletal pain (0.93, 95% confidence interval 0.87-0.99). We observed no such association in men with high strength (0.99, 0.93-1.05). Furthermore, no statistically significant increase or decrease in risk was observed for any of the secondary outcomes.

Conclusion

In men, low overall isometric muscle strength in youth was not associated with an increased risk of future musculoskeletal pain. Contrarily, we observed a slightly decreased risk of self-reported musculoskeletal pain in adulthood. Residual confounding by adult occupational exposures, leisure time physical activity level and psychosocial factors may have impacted on results.

Article summary

Article focus

- We aimed to study whether overall muscle strength in youth is inversely associated with the development of musculoskeletal pain in adulthood.

Key messages

- In contradiction to our expectations, we observed a decreased risk of self-reported musculoskeletal pain in adult men with low overall isometric muscle strength in youth.
- We speculate that low muscle strength may serve as a deselection criterion for activities with high risk of acute injuries or chronic physical overload, factors with negative impact on musculoskeletal health.

Strengths and limitations of study

- The main strengths of the study are a large sample, the use of three different measures of muscle strength, and comparably long time to follow-up.
- The main limitations of the study are the following; the cohort does not include women, musculoskeletal pain only identified with one question per site, motivation for military service might influence measurement of muscle strength, and the potential for unmeasured or residual confounding.

Introduction

Musculoskeletal disorders (MSDs), such as low back pain, osteoarthritis, and widespread pain, are highly prevalent in the adult population.¹⁻³ MSDs also contribute to a substantial burden of disease at middle and older ages.⁴ Although pain emanating from the musculoskeletal system might be attributed to a wide range of diseases with diverse causal chains, many MSDs have common risk factors such as heavy occupational work load,^{5,6} a high body mass index (BMI),⁷⁻⁹ and a low socio-economy.¹⁰⁻¹² Although smoking in some studies have been identified as a risk factor for certain MSDs,^{13,14} its main effect on musculoskeletal pain might be as an effect modifier of the pain sensation.¹⁵ As physical work load is a risk factor for many MSDs,¹⁶ a model in which the muscle strength in the loaded parts of the body are protective for future disorders is appealing.

Furthermore, physical exercise with focus on muscle strength is an important secondary and tertiary prevention of MSDs.^{17,18} A handful of studies have hitherto longitudinally investigated the strength of isolated muscle groups in adulthood as a determinant of later MSDs.¹⁹⁻²² However, in adult subjects, there is for the time being conflicting evidence of the value of muscle strength as a protective factor of musculoskeletal pain, such as neck/shoulder pain and low back pain.¹⁹

In the longer perspective, relatively little is known about the association of muscle strength in youth and later disease, including musculoskeletal pain.²³ Two studies have investigated the result of single muscle strength tests as determinants of musculoskeletal complaints decades later.^{24,25} The first, using number of sit-ups during 30 seconds as a strength measure, found no association with later low back pain or tension neck in men. In women, the high strength group had a decreased odds ratio (OR) of tension neck.²⁴ The second study found a decreased OR for MSDs in men who either had a strong performance in isotonic bench press or in an isometric two hand lift test.²⁵ Neither of the two studies includes a measure of overall muscular capacity. In summary, although there is some evidence of an association between low muscle strength in youth and later risk of MSDs, the association between overall muscle strength in adolescence and later musculoskeletal pain has not

1
2
3 been studied. Furthermore, with a larger sample size, data on common risk factors, testing of three
4
5 different muscle groups, and data on physical work capacity, we address some of the limitations of
6
7 earlier studies.
8
9

10 Our aim in this study was to investigate the general muscle strength in adolescent males as a
11
12 determinant of later self-reported musculoskeletal pain. We hypothesized that low general muscle
13
14 strength in youth is associated with an increased risk of having musculoskeletal pain in adulthood.
15
16

17 **Methods**

20 For this prospective register-based cohort study, we used two main criteria to identify the study
21
22 sample. First, when typically aged 18, the subjects should have performed mandatory conscription
23
24 testing in Sweden between 1970-1994, with the exception of the years 1978 and 1985. Secondly,
25
26 they should have been included in the Swedish Living Conditions Surveys any year between 1980 and
27
28 2005 when questions regarding musculoskeletal problems, smoking status, and physical activity were
29
30 simultaneously included.
31
32

33
34 Furthermore, we excluded all men who were surveyed prior to the baseline testing or were younger
35
36 than 17 years or older than 19 years at baseline (Table 1). We also excluded men with an existing
37
38 musculoskeletal disorder (Diseases of the musculoskeletal system and connective tissue according to
39
40 the International classification of disease version 8 or 9) and those who had missing data on variables
41
42 included in the primary model (muscle strength, smoking, BMI, physical activity, level of education).
43
44 In the final study sample we included 5489 men (figure 1). Data from the Swedish conscription
45
46 testing has been previously used for research purposes.²⁶⁻²⁸ During the period of the study sample
47
48 testing, conscription was mandatory by law for all Swedish men. Specially trained employees at six
49
50 regional conscription offices administrated the conscription tests during a two-day session that also
51
52 included separate evaluations by a medical doctor and a psychologist. Only men with serious health
53
54 complaints, were excused from conscription. The procedure included measurements of each
55
56
57
58
59
60

1
2
3 subject's weight in underwear to the kilogram and height without shoes to the centimeter. Using
4 height and weight, we calculated BMI as height/kg². Probably due to rare errors of data entry, there
5 are unlikely extreme values in the dataset. Therefore, we excluded all subjects with registered
6 extreme values on height (<150, >210 cm), weight (<40, >150 kg), or an extreme calculated value for
7 BMI (<15, >60) (figure 1). The study was approved by the Ethical Review Board at Lund University
8 and the manuscript was prepared according to the STROBE-statement.²⁹

15 *Muscle strength*

16
17 The men performed three tests of isometric muscle strength during conscription; hand grip, elbow
18 flexion, and knee extension. At the start of test period in 1970, the tests were performed as
19 previously described³⁰ and remained unchanged in general throughout the test period. In summary,
20 hand grip strength on the preferred side was measured with 90° flexion at the elbow and the
21 humerus in parallel to the torso. Knee extension was measured in a sitting position with 90° knee
22 flexion and arms crossed over chest. The pelvis was fixed to the seat and a strap fastened above the
23 lateral malleolus. Also, elbow flexion was measured in a sitting position with 90° flexion at the elbow
24 and the humerus in parallel to the torso. A strap was fastened at the level of the radial styloid
25 process.

26
27 We calculated a measure of general muscle strength by standardizing and combining the three tests
28 of muscle strength. To avoid bias due to change of testing procedure over time, we categorized the
29 cohort into five subgroups based on period of conscription (1970-1973, 1974-1977, 1979-1984, 1986-
30 1990, 1990-1994) and for each subgroup calculated the relative muscle strength. We standardized
31 the three tests of muscle strength [standardized value= (value–mean)/standard deviation] within
32 each subgroup and used the mean of the three test scores as a proxy for general muscle strength.
33 Using percentiles, we then categorized the cohort into three groups of muscle strength, where the
34 25th to 75th percentile defined the average category, the bottom 25th percentile configured the low
35 category, and the top 25th percentile defined the high muscle strength category.

Survey of musculoskeletal pain

The Swedish Living Conditions Surveys (ULF) is a random population based survey conducted by Statistics Sweden, previously used for research purposes.³¹⁻³³ For the present study, we used data collected during a total of 10 years (1980, 1988, 1989, 1997-1999, 2002-2005). The surveys were generally performed as interviews in person by trained interviewers, with a minority of the interviews performed by phone. For men included in more than one survey, we used the last survey without relevant study data missing.

At follow-up, the men were asked three questions regarding any current musculoskeletal pain: 1) Do you have pain in neck or shoulders? 2) Do you have back-pain, hip-pain or sciatica? 3) Do you have ache, pain in hands, elbows, legs or knees? For each type of complaint one of three answers was possible: 1) Yes, severe 2) Yes, mild and 3) No. Our primary outcome was having reported either severe or mild musculoskeletal pain, whereas our secondary outcomes we defined as follows: 1) Having reported severe musculoskeletal pain 2) Having reported pain in back/hips 3) Having reported pain in shoulders/neck 4) Having reported pain in arms/legs. From the surveys, we also included data on self-reported current smoking status (yes/no), physical activity (practically none, now and then, regularly, regularly strenuous), and level of education (compulsory school or less, secondary education, higher education). Drop-outs from the survey, i.e. those who have declined participation, cannot be individually identified. However, during the years of survey used in this study, the participation rate in the survey among men in relevant age groups were 70.0-88.7%.

Statistical analyses

All analyses were performed with SAS 9.3 (SAS Institute Inc). We used logistic binomial regression to estimate relative risks (RR) and control potential confounders. In the multivariate model (primary model), we included muscle strength, BMI, smoking status, physical activity and level of education.

Sensitivity analyses

To test whether cardiovascular aspects of physical capacity confounded our results, we used physical work capacity measured as $W_{\max 6 \text{ min}}$ in a sensitivity analysis. For $W_{\max 6 \text{ min}}$, the test result is an estimate of maximum work sustainable for 6 minutes³⁴ and is in young men correlated with maximum oxygen uptake ($r=0.9$).^{35 36} Acceptable data quality on work capacity was available in the subsample of men performing the baseline testing in 1976-1982. Out of all men in the cohort conscripted during the time period, 1154 men (74.6 %) completed an acceptable physical work capacity test on a bicycle ergometer (i.e. heart rate >174 at the end of testing). We added the work capacity in relation to body weight as a continuous variable to the univariate model. Furthermore, we also performed two sensitivity analyses on the multivariate model with musculoskeletal pain as the dependent variable. In the first, we added test center to the model. For the second, to test whether our categorization of muscle strength influenced the results, we 1) treated the standardized muscle strength as a continuous variable 2) treated the standardized muscle strength as a categorical variable based on quintiles.

Results

The mean time to follow-up was 17 years (table 1). Men with low muscle strength did not have an increased risk, but rather a statistically significant decreased risk, for the primary outcome "Musculoskeletal pain" (table 2). To summarize the observations of the secondary outcomes, we did not observe any statistically significant risk increases for neither men with a low nor high muscle strength. Compared to the crude model, the multivariate model produced similar risk estimates.

Sensitivity analyses

Work capacity had a significant effect in the subsample analysis ($p=0.03$) whereas it had only minor effect on the risk estimates for musculoskeletal pain, being 0.94 (95% confidence interval 0.82-1.08) and 1.02 (0.90-1.16) for low and high strength, respectively. The pattern of association in the secondary outcomes were in general somewhat strengthened when we adjusted for work capacity (data not shown). Using muscle strength as a continuous variable (hence assuming a linear relationship) did weaken the association with later musculoskeletal pain ($p=0.23$). When we instead used quintiles to categorize muscle strength, we observed no increased risk for the group with lowest strength compared to average strength (RR=0.93, 0.85-1.01).

Discussion

Investigating the overall isometric muscle strength in adolescent men as a determinant of future musculoskeletal pain, we observed a decreased risk of self-reported musculoskeletal pain in men with low muscle strength. No such association was observed for men with high strength. We also found a similar, however not statistically significant, pattern for "pain in back/hips" and "pain in neck/shoulders", whereas no association was found for future problems in arms/legs. Noticeably the observed associations were in contradiction to our expectations.

1
2
3 Using a historical cohort design with prospective registration of exposure and the outcome, our study
4 includes a large sample and thus allows better control for known confounders compared to previous
5 studies. All covariates (muscle strength, BMI, smoking, education, physical activity) included in the
6 multivariate model for musculoskeletal pain also had a significant association with the outcome.
7

8
9
10
11 Furthermore, by controlling for work capacity, we aimed to isolate the direct effect of muscle
12 strength from other aspects of physical capacity. However, the study also has important limitations.
13
14 First and foremost, we have used strength data from military conscription testing. Although it
15 provides a rich dataset from a structured environment, we do not know how the subject's motivation
16 for military service may have biased the performance during the testing procedure. However,
17 assuming there is no association between motivation at conscription testing and later risk of
18 musculoskeletal pain (or the loss to follow-up) any bias would at most dilute our result. Nevertheless,
19 stronger recruits are more likely to be assigned to positions with heavy load duty. Secondly, the
20 conscription was mandatory for men only. As the pattern of physical activity and occupational
21 exposure differ between men and women, any generalization of the results to women must be made
22 with great caution. Third, the physical activity measurement consisted of a single question and did
23 neither allow calculation of metabolic equivalent of task (MET) nor included occupational exposure.
24 Also musculoskeletal pain is measured by questions that combine more than one site, decreasing the
25 precision. However, the categories are fairly well demarcated anatomically save for the question
26 regarding pain in arms/legs. Fourth, the covariates collected with musculoskeletal pain at follow-up
27 (smoking, physical activity, level of education) are cross-sectional and might thus be mediators of
28 reverse causation. However, the adjusted estimates are much in line with the crude estimates.
29

30
31
32
33 It has previously been suggested that there is a U-shaped association between physical activity and
34 later back pain.^{37 38} Furthermore, as former occupational exposure and certain sport participation
35 are established risk factors for future MSDs,^{16 39} it lends some evidence for a more general model, in
36 which certain forms of physical activity is negative for the musculoskeletal health. Primarily, our
37 observations do not support low muscle strength in youth as a risk factor for later musculoskeletal
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 pain. Instead, we speculate that our results can be explained by muscle strength in youth being one
4
5 selection criterion for future high risk activities with a negative influence on the musculoskeletal
6
7 health, e.g. higher risk of joint injury due to sports participation or manual repetitive work load. This
8
9 would also include more immediate exposure such as more physically demanding military service.⁴⁰
10
11 Although we have controlled for level of education, which might serve as a proxy for occupational
12
13 exposure, there is potential for residual confounding as we did not have more appropriate data. In
14
15 other words, individuals with low general muscle strength might to a certain degree be deselected
16
17 for high risk activities compared to men with an average or high strength. However, this is only one
18
19 of many possible explanations. For example, the strength of an individual is associated with the
20
21 muscle fiber type distribution, which have a large genetic component.⁴¹ Type I fibers are more
22
23 common in endurance athletes⁴² whereas high type II percentage have been reported to be
24
25 associated with isometric muscle strength⁴³ as well as low back pain.⁴⁴ Thus, our observations could
26
27 potentially be explained by both social and biological factors.
28
29

30
31 Partly in contrast with a previous study,²⁵ we did not observe a negative effect of low muscle
32
33 strength on the risk future of musculoskeletal problems in men. Although we in the present study
34
35 only include measurements of isometric strength, the previous study observed associations with
36
37 both an isometric strength measure (static two hand lift) and an isotonic strength measure (bench
38
39 press). In a study on the same cohort, it is reported that the result in bench press, but not two hand
40
41 lift, was associated with both future cardiovascular fitness and future physical activity,⁴⁵ potentially
42
43 explaining part of the difference. Another study reported flexibility as a sit and reach test,²⁴ but not
44
45 strength measured as sit-ups, to be negatively associated with future risk of back pain. Hence, it is
46
47 possible that other aspects of muscular and musculoskeletal function is of greater importance of
48
49 future risk of MSDs than isometric muscle strength.
50
51

52
53 The association between muscle strength and later musculoskeletal pain diminished when we used
54
55 muscle strength as a continuous variable. This was not surprising, as the observed association was
56
57
58
59
60

1
2
3 non linear in the primary model. It is to be expected that the test offices differ somewhat in their
4 reported test results, as they were assigned adolescents based on geography. However, including
5 test offices in the model did not have any effect on the overall association. When we included work
6 capacity in the model, most risk estimates decreased in absolute values, furthering strengthening the
7 observations in the primary model. By using the relative muscle strength during testing periods of
8 five years, we partially address the potential systematic change in testing procedure over the years.
9 Although the methods of measurements have not changed at large, minor adjustments cannot be
10 excluded.
11
12

13
14
15
16
17
18
19
20
21 Although we found no increased risk of future musculoskeletal problems in men with low muscle
22 strength in adolescence, future studies need to better quantify the occupational exposure and leisure
23 time physical activity. Since the physical activity pattern and physical fitness profiles differ between
24 men and women, further investigations are also needed to investigate if similar associations can be
25 found in women.
26
27
28
29
30
31

32
33 In contrast to our hypothesis, we observed a decreased risk of self-reported musculoskeletal pain in
34 adult men with low overall isometric muscle strength in youth. We speculate that low muscle
35 strength *per se* is not protective of future musculoskeletal pain. Instead, low muscle strength might
36 serve as a deselection criterion for professions or types of leisure time physical activity with higher
37 risk of acute injuries or chronic physical overload, factors with negative impact on musculoskeletal
38 health.
39
40
41
42
43
44
45

46 **Acknowledgements**

47
48
49 No author has a conflict of interest to report. The authors would like to thank Charlotte Bergknot for
50 the initial data management.
51
52
53
54

55 **Funding**

1
2
3 The authors are supported by grants from The Swedish Research Council, Skåne Regional Council,
4 and the Faculty of Medicine at Lund University. The funders neither influenced the study process nor
5
6 manuscript preparation.
7
8
9

10 **Author contributions**

11
12
13
14 ST proposed the study idea, collected data, designed the study, and drafted the manuscript. IP
15
16 designed the study and provided input during manuscript preparation. CZ designed the study and
17
18 performed the statistical analyses. ME designed the study, collected data, and provided input during
19
20 manuscript preparation. All authors approved the final version of the manuscript.
21
22

23 **Data sharing statement**

24
25
26
27 There is no additional data to share.
28
29

30 **References**

- 31
32
33
34 1. Bergman S, Herrstrom P, Hogstrom K, Petersson IF, Svensson B, Jacobsson LT. Chronic
35
36 musculoskeletal pain, prevalence rates, and sociodemographic associations in a Swedish
37
38 population study. *J Rheumatol* 2001;28(6):1369-77.
39
40 2. Picavet HS, Schouten JS. Musculoskeletal pain in the Netherlands: prevalences, consequences and
41
42 risk groups, the DMC(3)-study. *Pain* 2003;102(1-2):167-78.
43
44 3. Urwin M, Symmons D, Allison T, Brammah T, Busby H, Roxby M, et al. Estimating the burden of
45
46 musculoskeletal disorders in the community: the comparative prevalence of symptoms at
47
48 different anatomical sites, and the relation to social deprivation. *Ann Rheum Dis*
49
50 1998;57(11):649-55.
51
52 4. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years
53
54 (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the
55
56 Global Burden of Disease Study 2010. *Lancet* 2013;380(9859):2197-223.
57
58
59
60

- 1
2
3 5. van der Windt DA, Thomas E, Pope DP, de Winter AF, Macfarlane GJ, Bouter LM, et al.
4
5 Occupational risk factors for shoulder pain: a systematic review. *Occup Environ Med*
6
7 2000;57(7):433-42.
8
- 9
10 6. Hoogendoorn WE, van Poppel MN, Bongers PM, Koes BW, Bouter LM. Physical load during work
11
12 and leisure time as risk factors for back pain. *Scand J Work Environ Health* 1999;25(5):387-
13
14 403.
- 15
16 7. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, et al. Risk factors for incident
17
18 radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum*
19
20 1997;40(4):728-33.
21
- 22
23 8. Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between obesity
24
25 and low back pain: a meta-analysis. *Am J Epidemiol* 2010;171(2):135-54.
26
- 27
28 9. Grotle M, Hagen KB, Natvig B, Dahl FA, Kvien TK. Obesity and osteoarthritis in knee, hip and/or
29
30 hand: an epidemiological study in the general population with 10 years follow-up. *BMC*
31
32 *Musculoskelet Disord* 2008;9:132.
- 33
34 10. Hagen K, Zwart JA, Svebak S, Bovim G, Jacob Stovner L. Low socioeconomic status is associated
35
36 with chronic musculoskeletal complaints among 46,901 adults in Norway. *Scand J Public*
37
38 *Health* 2005;33(4):268-75.
39
- 40
41 11. Dalstra JA, Kunst AE, Borrell C, Breeze E, Cambois E, Costa G, et al. Socioeconomic differences in
42
43 the prevalence of common chronic diseases: an overview of eight European countries. *Int J*
44
45 *Epidemiol* 2005;34(2):316-26.
- 46
47 12. Bengtsson C, Nordmark B, Klareskog L, Lundberg I, Alfredsson L. Socioeconomic status and the
48
49 risk of developing rheumatoid arthritis: results from the Swedish EIRA study. *Ann Rheum Dis*
50
51 2005;64(11):1588-94.
- 52
53 13. Palmer KT, Syddall H, Cooper C, Coggon D. Smoking and musculoskeletal disorders: findings from
54
55 a British national survey. *Ann Rheum Dis* 2003;62(1):33-6.
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
14. Holth HS, Werpen HK, Zwart JA, Hagen K. Physical inactivity is associated with chronic musculoskeletal complaints 11 years later: results from the Nord-Trondelag Health Study. *BMC Musculoskelet Disord* 2008;9:159.
15. McBeth J, Jones K. Epidemiology of chronic musculoskeletal pain. *Best Pract Res Clin Rheumatol* 2007;21(3):403-25.
16. Roquelaure Y, Ha C, Rouillon C, Fouquet N, Leclerc A, Descatha A, et al. Risk factors for upper-extremity musculoskeletal disorders in the working population. *Arthritis Rheum* 2009;61(10):1425-34.
17. Hakkinen A, Sokka T, Kotaniemi A, Hannonen P. A randomized two-year study of the effects of dynamic strength training on muscle strength, disease activity, functional capacity, and bone mineral density in early rheumatoid arthritis. *Arthritis Rheum* 2001;44(3):515-22.
18. Baker KR, Nelson ME, Felson DT, Layne JE, Sarno R, Roubenoff R. The efficacy of home based progressive strength training in older adults with knee osteoarthritis: a randomized controlled trial. *J Rheumatol* 2001;28(7):1655-65.
19. Hamberg-van Reenen HH, Ariens GA, Blatter BM, van Mechelen W, Bongers PM. A systematic review of the relation between physical capacity and future low back and neck/shoulder pain. *Pain* 2007;130(1-2):93-107.
20. Leino P, Aro S, Hasan J. Trunk muscle function and low back disorders: a ten-year follow-up study. *J Chronic Dis* 1987;40(4):289-96.
21. Kujala UM, Taimela S, Viljanen T, Jutila H, Viitasalo JT, Videman T, et al. Physical loading and performance as predictors of back pain in healthy adults. A 5-year prospective study. *Eur J Appl Physiol Occup Physiol* 1996;73(5):452-8.
22. Faber A, Sell L, Hansen JV, Burr H, Lund T, Holtermann A, et al. Does muscle strength predict future musculoskeletal disorders and sickness absence? *Occupational medicine (Oxford, England)* 2012;62(1):41-6.

- 1
2
3 23. Ruiz JR, Castro-Pinero J, Artero EG, Ortega FB, Sjostrom M, Suni J, et al. Predictive validity of
4 health-related fitness in youth: a systematic review. *Br J Sports Med* 2009;43(12):909-23.
5
6
7 24. Mikkelsen LO, Nupponen H, Kaprio J, Kautiainen H, Mikkelsen M, Kujala UM. Adolescent
8 flexibility, endurance strength, and physical activity as predictors of adult tension neck, low
9 back pain, and knee injury: a 25 year follow up study. *Br J Sports Med* 2006;40(2):107-13.
10
11
12 25. Barnekow-Bergkvist M, Hedberg GE, Janlert U, Jansson E. Determinants of self-reported neck-
13 shoulder and low back symptoms in a general population. *Spine (Phila Pa 1976)*
14 1998;23(2):235-43.
15
16
17 26. Neovius M, Sundstrom J, Rasmussen F. Combined effects of overweight and smoking in late
18 adolescence on subsequent mortality: nationwide cohort study. *BMJ* 2009;338:b496.
19
20
21 27. Nilsson PM, Nyberg P, Ostergren PO. Increased susceptibility to stress at a psychological
22 assessment of stress tolerance is associated with impaired fetal growth. *Int J Epidemiol*
23 2001;30(1):75-80.
24
25
26 28. Hemmingsson T, Melin B, Allebeck P, Lundberg I. The association between cognitive ability
27 measured at ages 18-20 and mortality during 30 years of follow-up--a prospective
28 observational study among Swedish males born 1949-51. *Int J Epidemiol* 2006;35(3):665-70.
29
30
31 29. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. Strengthening the
32 Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for
33 reporting observational studies. *BMJ* 2007;335(7624):806-8.
34
35
36 30. Nordesjo LO, Schéle R. Validity of Ergometer Cycle Test and Measures of Isometric Muscle
37 Strength when Predicting some Aspects of Military Performance. *Swedish J Defence Med*
38 1974(10):11-23.
39
40
41 31. Wolk A, Rossner S. Effects of smoking and physical activity on body weight: developments in
42 Sweden between 1980 and 1989. *J Intern Med* 1995;237(3):287-91.
43
44
45 32. Sundquist J, Johansson SE. The influence of socioeconomic status, ethnicity and lifestyle on body
46 mass index in a longitudinal study. *Int J Epidemiol* 1998;27(1):57-63.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 33. Bygren LO, Konlaan BB, Johansson SE. Attendance at cultural events, reading books or
4
5 periodicals, and making music or singing in a choir as determinants for survival: Swedish
6
7 interview survey of living conditions. *BMJ* 1996;313(7072):1577-80.
8
9
10 34. Tornvall G. Assessment of physical capabilities. *Acta physiologica Scandinavica* 1963;58(Suppl
11
12 201):1-102.
13
14 35. Nordesjo LO. A comparison between the Thornvall maximal ergometer test, submaximal
15
16 ergometer tests and maximal oxygen uptake. *Swedish J Defence Med* 1974(10):3-10.
17
18 36. Nordesjo LO. Estimation of the maximal work rate sustainable for 6 minutes using a single-level
19
20 load or stepwise increasing loads. *Ups J Med Sci* 1974;79(1):45-50.
21
22 37. Heneweer H, Vanhees L, Picavet HS. Physical activity and low back pain: a U-shaped relation?
23
24
25 *Pain* 2009;143(1-2):21-5.
26
27 38. Heneweer H, Picavet HS, Staes F, Kiers H, Vanhees L. Physical fitness, rather than self-reported
28
29 physical activities, is more strongly associated with low back pain: evidence from a working
30
31 population. *Eur Spine J* 2012;21(7):1265-72.
32
33 39. Kujala UM, Kettunen J, Paananen H, Aalto T, Battie MC, Impivaara O, et al. Knee osteoarthritis in
34
35 former runners, soccer players, weight lifters, and shooters. *Arthritis Rheum* 1995;38(4):539-
36
37 46.
38
39 40. Almeida SA, Williams KM, Shaffer RA, Brodine SK. Epidemiological patterns of musculoskeletal
40
41 injuries and physical training. *Med Sci Sports Exerc* 1999;31(8):1176-82.
42
43 41. Simoneau JA, Bouchard C. Genetic determinism of fiber type proportion in human skeletal
44
45 muscle. *FASEB J* 1995;9(11):1091-5.
46
47 42. Costill DL, Daniels J, Evans W, Fink W, Krahenbuhl G, Saltin B. Skeletal muscle enzymes and fiber
48
49 composition in male and female track athletes. *J Appl Physiol* 1976;40(2):149-54.
50
51 43. Tesch P, Karlsson J. Isometric strength performance and muscle fibre type distribution in man.
52
53
54
55 *Acta physiologica Scandinavica* 1978;103(1):47-51.
56
57
58
59
60

- 1
2
3 44. Mannion AF, Weber BR, Dvorak J, Grob D, Muntener M. Fibre type characteristics of the lumbar
4 paraspinal muscles in normal healthy subjects and in patients with low back pain. *J Orthop*
5 *Res* 1997;15(6):881-7.
6
7
8
9 45. Barnekow-Bergkvist M, Hedberg G, Janlert U, Jansson E. Prediction of physical fitness and
10 physical activity level in adulthood by physical performance and physical activity in
11 adolescence--an 18-year follow-up study. *Scand J Med Sci Sports* 1998;8(5 Pt 1):299-308.
12
13
14
15
16
17
18
19

20 Tables and Figures

21 **Table 1:** Description of study sample

22
23 **Table 2:** Risk estimates for the muscle strength in youth as a determinant of self-reported
24 musculoskeletal pain in adulthood
25
26
27

28
29 **Figure 1:** The identification of the study sample and the loss to follow-up
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Description of study sample

Number of men	5489
Mean age at baseline (SD)	18.2 (0.5)
Mean time to follow-up (SD, range)	17.2 years (8.4, 1-35)
Mean hand grip* (SD)	617 (98)
Mean elbow flexion* (SD)	385 (83)
Mean knee extension* (SD)	567 (116)
Muscle Strength (%)	
Low	1371 (25.0)
Average	2747 (50.0)
High	1371 (25.0)
BMI (%)	
<18.5	477 (8.7)
18.5-24.9	4498 (81.9)
25-29.9	448 (8.2)
>30	66 (1.2)
Type of interview (%)	
In person	4349 (79.2)
By telephone	1140 (20.8)
Pain in back/hips (%)	
Yes	1645 (30.0)
of which severe	321 (5.8)
No	3842 (70.0)
Missing	2 (0.0)
Pain in neck/shoulders (%)	
Yes	1562 (28.5)
of which severe	246 (4.5)
No	3925 (71.5)
Missing	2 (0.0)
Pain in arms/legs (%)	
Yes	1243 (22.6)
of which severe	196 (3.6)
No	4243 (77.3)
Missing	3 (0.0)
Pain, independent of location (%)	
Yes	2847 (51.9)
of which severe	576 (10.5)
No	2639 (48.1)
Missing	3 (0.0)
Smoking status (%)	
Yes	827 (15.1)
No	4662 (84.9)
Level of education (%)	
Compulsory	589 (10.7)
Secondary	2889 (52.6)
Higher	2011 (36.6)
Physical Activity (%)	
Practically non	583 (10.6)
Now and then	1613 (29.4)
Regularly	2077 (37.8)
Regularly strenuous	1216 (22.2)

* In Newton.

Table 2 Risk estimates for the muscle strength in youth as a determinant of self-reported musculoskeletal pain in adulthood

Outcomes	Models				
	Univariate Muscle strength			Multivariate ^a Muscle strength	
	Low (N=1371) RR (N)	Average (N=2747) Reference (N)	High (N=1371) RR (N)	Low RR (95% CI)	High RR (95% CI)
Musculoskeletal pain	0.92 (668)	1 (1457)	0.99 (722)	0.93 (0.87-0.99)	0.99 (0.93-1.05)
Severe musculoskeletal pain	0.96 (135)	1 (283)	1.12 (158)	0.96 (0.79-1.18)	1.07 (0.89-1.29)
Pain in back/hips	0.92 (384)	1 (832)	1.03 (429)	0.93 (0.84-1.03)	1.03 (0.94-1.13)
Pain in neck/shoulders	0.92 (366)	1 (799)	0.99 (397)	0.93 (0.83-1.03)	1.00 (0.90-1.10)
Pain in arms/legs	0.97 (297)	1 (616)	1.07 (330)	0.97 (0.86-1.10)	1.06 (0.94-1.19)

N= Number of cases

RR= Relative risk estimates

CI= Confidence interval

a= adjusted for smoking status, physical activity, education, body mass index

1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Muscle Strength in Adolescent Men and Future Musculoskeletal Pain: A Cohort Study with 17 Years of Follow-up

Simon Timpka¹, Ingemar F Petersson¹, Caddie Zhou¹, Martin Englund^{1, 2}

1 Department of Orthopedics, Clinical Sciences Lund, Lund University, Lund, Sweden

2 Clinical Epidemiology Research & Training Unit, Boston, University School of Medicine, Boston, Massachusetts, United States of America

Corresponding author:

Simon Timpka
Skånes universitetssjukhus
Epi-centrum Skåne
Medicon Village
223 81 Lund
Sweden

Phone: +46 738 395 276
Mail: simon.timpka@med.lu.se

Words in main text: 2800
Number of figures: 1
Number of tables: 2

Abstract

Objectives

Musculoskeletal pain is highly prevalent throughout adulthood with a major impact on health, function and participation in the society. Still, the association between muscle strength and development of musculoskeletal pain is unclear. We aimed to study whether overall muscle strength in adolescent men is associated with self-reported musculoskeletal pain in adulthood.

Design

Cohort study with baseline data from the Swedish Conscription Register and outcome information from the random population-based Swedish Living Conditions Surveys.

Setting

Sweden, 1970-2005.

Participants

We studied 54 8992 men who at age 17-19 years tested their isometric muscle strength (hand grip, arm flexion, knee extension) during the compulsory conscription.

Outcome measures

The men were surveyed regarding self-reported musculoskeletal pain; mean follow-up time of 17 (range 1-35) years. Our primary outcome was a self-report of musculoskeletal pain, and secondary outcomes were a report of "severe pain", "pain in back/hips", "pain in neck/shoulders", or "pain in arms/legs", respectively. We categorized muscle strength into three groups; low, average, and high using the 25th to 75th percentile to define the reference category (average). We ~~calculated incidence rate ratios~~ estimated relative risks using log binomial regression with adjustment for smoking, body mass index, education, and physical activity.

3

Results

In the adjusted model, men with low overall muscle strength had decreased risk of self-reported musculoskeletal pain (~~incidence rate ratio~~ 0.93, 95% confidence interval 0.87-0.99). We observed no such association in men with high strength (0.99, 0.93-1.05). Furthermore, no statistically significant increase or decrease in risk was observed for any of the secondary outcomes.

Conclusion

In men, low overall isometric muscle strength in youth was not associated with an increased risk of future musculoskeletal pain. Contrarily, we observed a slightly decreased risk of self-reported musculoskeletal pain in adulthood. Residual confounding by adult occupational exposures, leisure time physical activity level and psychosocial factors may have impacted on results.

Formatted: Keep with next

Article summary

Article focus

- We aimed to study whether overall muscle strength in youth is inversely associated with the development of musculoskeletal pain in adulthood.

Key messages

- In contradiction to our expectations, we observed a decreased risk of self-reported musculoskeletal pain in adult men with low overall isometric muscle strength in youth.
- We speculate that low muscle strength may ~~potentially~~ serve as a deselection criterion for activities with high risk of acute injuries or chronic physical overload, factors with negative impact on musculoskeletal health.

Strengths and limitations of study

- The main strengths of the study are a large sample, the use of three different measures of muscle strength, and comparably long time to follow-up.
- The main limitations of the study are the following; the cohort does not include women, musculoskeletal pain only identified with one question per site, motivation for military service might influence measurement of muscle strength, and the potential for unmeasured or residual confounding.

Introduction

Musculoskeletal disorders (MSDs), such as low back pain, osteoarthritis, and widespread pain, are highly prevalent in the adult population.¹⁻³ MSDs also contribute to a substantial burden of disease at middle and older ages.⁴ Although pain emanating from the musculoskeletal system might be attributed to a wide range of diseases with diverse causal chains, many MSDs have common risk factors such as heavy occupational work load,^{5,6} a high body mass index (BMI),⁷⁻⁹ and a low socio-economy.¹⁰⁻¹² Although smoking in some studies have been identified as a risk factor for certain MSDs,^{13,14} its main effect on musculoskeletal pain might be as an effect modifier of the pain sensation.¹⁵ As physical work load is a risk factor for many MSDs,¹⁶ a model in which the muscle strength in the loaded parts of the body are protective for future disorders is appealing.

Furthermore, physical exercise with focus on muscle strength is an important secondary and tertiary prevention of MSDs.^{17,18} A handful of studies have hitherto longitudinally investigated the strength of isolated muscle groups in adulthood as a determinant of later MSDs.¹⁹⁻²² However, in adult subjects, there is for the time being conflicting evidence of the value of muscle strength as a protective factor of musculoskeletal pain, such as neck/shoulder pain and low back pain.¹⁹

In the longer perspective, relatively little is known about the association of muscle strength in youth and later disease, including musculoskeletal pain.²³ Two studies have investigated the result of single muscle strength tests as determinants of musculoskeletal complaints decades later.^{24,25} The first, using number of sit-ups during 30 seconds as a strength measure, found no association with later low back pain or tension neck in men. In women, the high strength group had a decreased odds ratio (OR) of tension neck.²⁴ The second study found a decreased OR for MSDs in men who either had a strong performance in isotonic bench press or in a isometric two hand lift test.²⁵ Neither of the two studies includes a measure of overall muscular capacity. In summary, although there is some evidence of an association between low muscle strength in youth and later risk of MSDs, the association between overall muscle strength in adolescence and later musculoskeletal pain has not

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

been studied. Furthermore, with a larger sample size, data on common risk factors, testing of three different muscle groups, and data on physical work capacity, we address some of the limitations of earlier studies.

Our aim in this study was to investigate the general muscle strength in adolescent males as a determinant of later self-reported musculoskeletal pain. We hypothesized that low general muscle strength in youth is associated with an increased risk of having musculoskeletal pain in adulthood.

Methods

For this prospective register-based cohort study, we used two main criteria to identify the study sample cohort. First, when typically aged 18, the subjects should have performed mandatory conscription testing in Sweden between 1970-1994, with the exception of the years 1978 and 1985. Secondly, they should have been included in the Swedish Living Conditions Surveys any year between 1980 and 2005 when questions regarding musculoskeletal problems, smoking status, and physical activity were simultaneously included.

Furthermore, we excluded all men who were surveyed prior to the baseline testing or were younger than 17 years or older than 19 years at baseline (Table 1). We also excluded men with an existing musculoskeletal disorder (Diseases of the musculoskeletal system and connective tissue according to the International classification of disease version 8 or 9) and those who had missing data on variables included in the primary model (muscle strength, smoking, BMI, physical activity, level of education).

In the final study sample we included 54 ~~8992~~ men (figure 1). Data from the Swedish conscription testing has been previously used for research purposes.²⁶⁻²⁸ During the period of the study sample testing, conscription was mandatory by law for all Swedish men. Specially trained employees at six regional conscription ~~center~~ offices administrated the conscription tests during a two-day session that also included separate evaluations by a medical doctor and a psychologist. Only men with serious health complaints, were excused from conscription. The procedure included measurements of each

7

subject's weight in underwear to the kilogram and height without shoes to the centimeter. Using height and weight, we calculated BMI as height/kg². Probably due to rare errors of data entry, there are unlikely extreme values in the dataset. Therefore, we excluded all subjects with registered extreme values on height (<150, >210 cm), weight (<40, >150 kg), or an extreme calculated value for BMI (<15, >60) (figure 1). The study was approved by the Ethical Review Board at Lund University and the manuscript was prepared according to the STROBE-statement.²⁹

Muscle strength

The men performed three tests of isometric muscle strength during conscription; hand grip, elbow flexion, and knee extension. At the start of test period in 1970, the tests were performed as previously described³⁰ and remained unchanged in general throughout the test period. In summary, hand grip strength on the preferred side was measured with a 90° flexion at the elbow and with the humerus in parallel to the torso. Knee extension was measured in a sitting position with 90° knee flexion and arms crossed over chest. The pelvis was fixed to the seat and a strap fastened above the lateral malleolus. Also, elbow flexion was measured in a sitting position with 90° flexion at the elbow and the humerus in parallel to the torso. A strap was fastened at the level of the radial styloid process.

We calculated a measure of general muscle strength by standardizing and combining the three tests of muscle strength. To avoid bias due to change of testing procedure over time, we categorized the cohort into five subgroups based on period of conscription (1970-1973, 1974-1977, 1979-1984, 1986-1990, 1990-1994) and for each subcohortgroup calculated the relative muscle strength. We standardized the three tests of muscle strength [standardized value= (value–mean)/standard deviation] within each subgroup and used the mean of the three test scores as a proxy for general muscle strength. Using percentiles, we then categorized the cohort into three groups of muscle strength, where the 25th to 75th percentile defined the average category, the bottom 25th percentile configured the low category, and the top 25th percentile defined the high muscle strength category.

8

Survey of musculoskeletal pain

The Swedish Living Conditions Surveys (ULF) is a random population based survey conducted by Statistics Sweden, previously used for research purposes.³¹⁻³³ For the present study, we used data collected during a total of 10 years (1980, 1988, 1989, 1997-1999, 2002-2005). The surveys were generally performed as interviews in person by trained interviewers, with a minority of the interviews performed by phone. For men included in more than one survey, we used the last survey without relevant study data missing.

At follow-up, the men were asked three questions regarding any current musculoskeletal pain: 1) Do you have pain in neck or shoulders? 2) Do you have back-pain, hip-pain or sciatica? 3) Do you have ache, pain in hands, elbows, legs or knees? For each type of complaint one of three answers was possible: 1) Yes, severe 2) Yes, mild and 3) No. Our primary outcome was having reported either severe or mild musculoskeletal pain, whereas our secondary outcomes we defined as follows: 1) Having reported severe musculoskeletal pain 2) Having reported pain in back/hips 3) Having reported pain in shoulders/neck 4) Having reported pain in arms/legs. From the surveys, we also included data on self-reported current smoking status (yes/no), physical activity (practically none, now and then, regularly, regularly strenuous), and level of education (compulsory school or less, secondary education, higher education). Drop-outs from the survey, i.e. those who have declined participation, cannot be individually identified. However, during the years of survey used in this study, the participation rate in the survey among men in relevant age groups were 70.0-88.7%. of the men who were asked to be included in the relevant age groups, 11.3-31.0% declined participation in the years of study. The mean rate of non-participation over the relevant years in the same group was 22.9%.

Formatted: Keep with next

Statistical analyses

All analyses were performed with SAS 9.3 (SAS Institute Inc). We used logistic binomial regression to estimate ~~incidence rate ratios~~ *relative risks* (RR) and control potential confounders. In the multivariate model (primary model), we included muscle strength, BMI, smoking status, physical activity and level of education.

Sensitivity analyses

Formatted: Font: Italic

To test whether cardiovascular aspects of physical capacity confounded our results, we used physical work capacity measured as $W_{\max 6 \text{ min}}$ in a sensitivity analysis. For $W_{\max 6 \text{ min}}$, the test result is an estimate of maximum work sustainable for 6 minutes³⁴ and is in young men correlated with maximum oxygen uptake ($r=0.9$).^{35 36} Acceptable data quality on work capacity was available in the subsample of men performing the baseline testing in 1976-1982. Out of all men in the cohort conscripted during the time period, 1154 men (74.6 %) completed an acceptable physical work capacity test on a bicycle ergometer (i.e. heart rate >174 at the end of testing). We added the work capacity in relation to body weight as a continuous variable to the univariate model. Furthermore, we also performed two sensitivity analyses on the multivariate model with musculoskeletal pain as the dependent variable. In the first, we added test center to the model. For the second, to test whether our categorization of muscle strength influenced the results, we 1) treated the standardized muscle strength as a continuous variable 2) treated the standardized muscle strength as a categorical variable based on quintiles.

Results

The mean time to follow-up was 17 years (table 1). Men with low muscle strength did not have an increased risk, but rather a statistically significant decreased risk, for the primary outcome "Musculoskeletal pain" (table 2). To summarize the observations of the secondary outcomes, we did not observe any statistically significant risk increases for neither men with a low nor high muscle strength. Compared to the crude model, the multivariate model produced similar risk estimates.

Sensitivity analyses

Work capacity had a significant effect in the subsample analysis ($p=0.034$) whereas it had only minor effect on the risk estimates for musculoskeletal ~~problem~~, being 0.94 (95% confidence interval 0.82-1.08) and 1.02 (0.90-1.16) for low and high strength, respectively. The pattern of association in the secondary outcomes were in general somewhat strengthened when we adjusted for work capacity (data not shown). Using muscle strength as a continuous variable (hence assuming a linear relationship) did weaken the association with later musculoskeletal pain ($p=0.232$). When we instead used quintiles to categorize muscle strength, we observed no increased risk for the group with lowest strength compared to average strength (HR=0.93, 95% confidence interval 0.85-1.01).

Discussion

Investigating the overall isometric muscle strength in adolescent men as a determinant of future musculoskeletal pain, we observed a decreased risk of self-reported musculoskeletal pain in men with low muscle strength. No such association was observed for men with high strength. We also found a similar, however not statistically significant, pattern for "pain in back/hips" and "pain in neck/shoulders", whereas no association was found for future problems in arms/legs. Noticeably the observed associations were in contradiction to our expectations.

Using a historical cohort design with prospective registration of exposure and the outcome, our study includes a large sample and thus allows better control for known confounders compared to previous studies. All covariates (muscle strength, BMI, smoking, education, physical activity) included in the multivariate model for musculoskeletal pain also had a significant association with the outcome. Furthermore, by controlling for work capacity, we aimed to isolate the direct effect of muscle strength from other aspects of physical capacity. However, the study also has important limitations. First and foremost, we have used strength data from military conscription testing. Although it provides a rich dataset from a structured environment, we do not know how the subject's motivation for military service may have biased the performance during the testing procedure. However, assuming there is no association between motivation at conscription testing and later risk of musculoskeletal pain (or the loss to follow-up) any bias would at most dilute our result. Nevertheless, stronger recruits are more likely to be assigned to positions with heavy load duty. Secondly, the conscription was mandatory for men only. As the pattern of physical activity and occupational exposure differ between men and women, any generalization of the results to women must be made with great caution. Third, the physical activity measurement consisted of a single question and did neither allow calculation of metabolic equivalent of task (MET) nor included occupational exposure. Also musculoskeletal pain is measured by questions that combine more than one site, decreasing the precision. However, the categories are fairly well demarcated anatomically save for the question regarding pain in arms/legs. Fourth, the covariates collected with musculoskeletal pain at follow-up (smoking, physical activity, level of education) are cross-sectional and might thus be mediators of reverse causation. However, the adjusted estimates are much in line with the crude estimates.

It has [previously](#) been suggested that there is a U-shaped association between physical activity and later back pain.^{37 38} Furthermore, as former occupational exposure and certain sport participation are established risk factors for future MSDs,^{16 39} it lends some evidence for a more general model, in which certain forms of physical activity is negative for the musculoskeletal health. Primarily, our observations do not support low muscle strength in youth as a risk factor for later musculoskeletal

1
2
3
4
5
6
7 pain. Instead, we ~~speculatesuggest~~ that our results can be explained by muscle strength in youth
8 being one selection criterion for future high risk activities with a negative influence on the
9 musculoskeletal health, e.g. higher risk of joint injury due to sports participation or manual repetitive
10 work load. This would also include more immediate exposure such as more physically demanding
11 military service.⁴⁰ Although we have controlled for level of education, which might serve as a proxy
12 for occupational exposure, there is potential for residual confounding as we did not have more
13 appropriate data. In other words, individuals with low general muscle strength might to a certain
14 degree be deselected for high risk activities compared to men with an average or high strength.

15
16 However, this is only one of many possible explanations. For example, the strength of an individual is
17 associated with the muscle fiber type distribution, which have a large genetic component.⁴¹ Type I
18 fibers are more common in endurance athletes⁴² whereas high type II percentage have been
19 reported to be associated with isometric muscle strength⁴³ as well as low back pain.⁴⁴ Thus, our
20 observations could potentially be explained by both social and biological factors, we cannot exclude
21 that the decreased risk observed in men with low strength is mediated by factors related to muscle
22 fiber type.

23
24 Partly in contrast with a previous study,²⁵ we did not observe a negative effect of low muscle
25 strength on the risk future of musculoskeletal problems in men. Although we in the present study
26 only include measurements of isometric strength, the previous study observed associations with
27 both an isometric strength measure (static two hand lift) and an isotonic strength measure (bench
28 press). In a study on the same cohort, it is reported that the result in bench press, but not two hand
29 lift, was associated with both future cardiovascular fitness and future physical activity,⁴⁵ potentially
30 explaining part of the difference. ~~As a~~ another study reported flexibility as a sit and reach test,²⁴ but
31 not strength measured as sit-ups, to be negatively associated with future risk of back pain. Hence, it
32 is possible that other aspects of muscular and musculoskeletal function is of greater importance of
33 future risk of MSDs than isometric muscle strength.

The association between muscle strength and later musculoskeletal pain diminished when we used muscle strength as a continuous variable. This was not surprising, as the observed association was non linear in the primary model. It is to be expected that the test ~~offices~~~~centers~~ differ somewhat in their reported test results, as they were assigned adolescents based on geography. However, including test ~~office~~~~center~~s in the model did not have any effect on the overall association. When we included work capacity in the model, most risk estimates decreased in absolute values, furthering strengthening the observations in the primary model. By using the relative muscle strength during testing periods of five years, we partially address the potential systematic change in testing procedure over the years. Although the methods of measurements have not changed at large, minor adjustments cannot be excluded.

Although we found no increased risk of future musculoskeletal problems in men with low muscle strength in adolescence, future studies need to better quantify the occupational exposure and leisure time physical activity. Since the physical activity pattern and physical fitness profiles differ between men and women, further investigations are also needed to investigate if similar associations can be found in women.

In contrast to our hypothesis, wWe observed a decreased risk of self-reported musculoskeletal pain in adult men with low overall isometric muscle strength in youth. We ~~speculate~~~~hypothesize~~ that low muscle strength *per se* is not protective of future musculoskeletal pain. Instead, low muscle strength might serve as a deselection criterion for professions or types of leisure time physical activity with higher risk of acute injuries or chronic physical overload, factors with negative impact on musculoskeletal health.

Acknowledgements

No author has a conflict of interest to report. The authors would like to thank Charlotte Bergknut for the initial data management.

Funding

The authors are supported by grants from The Swedish Research Council, Skåne Regional Council, and the Faculty of Medicine at Lund University. The funders neither influenced the study process nor manuscript preparation.

Author contributions

ST proposed the study idea, collected data, designed the study, and drafted the manuscript. IP designed the study and provided input during manuscript preparation. CZ designed the study and performed the statistical analyses. ME designed the study, collected data, and provided input during manuscript preparation. All authors approved the final version of the manuscript.

Data sharing statement

There is no additional data to share.

References

1. Bergman S, Herrstrom P, Hogstrom K, Petersson IF, Svensson B, Jacobsson LT. Chronic musculoskeletal pain, prevalence rates, and sociodemographic associations in a Swedish population study. *J Rheumatol* 2001;28(6):1369-77.
2. Picavet HS, Schouten JS. Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study. *Pain* 2003;102(1-2):167-78.
3. Urwin M, Symmons D, Allison T, Brammah T, Busby H, Roxby M, et al. Estimating the burden of musculoskeletal disorders in the community: the comparative prevalence of symptoms at different anatomical sites, and the relation to social deprivation. *Ann Rheum Dis* 1998;57(11):649-55.

4. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2013;380(9859):2197-223.
5. van der Windt DA, Thomas E, Pope DP, de Winter AF, Macfarlane GJ, Bouter LM, et al. Occupational risk factors for shoulder pain: a systematic review. *Occup Environ Med* 2000;57(7):433-42.
6. Hoogendoorn WE, van Poppel MN, Bongers PM, Koes BW, Bouter LM. Physical load during work and leisure time as risk factors for back pain. *Scand J Work Environ Health* 1999;25(5):387-403.
7. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, et al. Risk factors for incident radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum* 1997;40(4):728-33.
8. Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between obesity and low back pain: a meta-analysis. *Am J Epidemiol* 2010;171(2):135-54.
9. Grotle M, Hagen KB, Natvig B, Dahl FA, Kvien TK. Obesity and osteoarthritis in knee, hip and/or hand: an epidemiological study in the general population with 10 years follow-up. *BMC Musculoskelet Disord* 2008;9:132.
10. Hagen K, Zwart JA, Svebak S, Bovim G, Jacob Stovner L. Low socioeconomic status is associated with chronic musculoskeletal complaints among 46,901 adults in Norway. *Scand J Public Health* 2005;33(4):268-75.
11. Dalstra JA, Kunst AE, Borrell C, Breeze E, Cambois E, Costa G, et al. Socioeconomic differences in the prevalence of common chronic diseases: an overview of eight European countries. *Int J Epidemiol* 2005;34(2):316-26.
12. Bengtsson C, Nordmark B, Klareskog L, Lundberg I, Alfredsson L. Socioeconomic status and the risk of developing rheumatoid arthritis: results from the Swedish EIRA study. *Ann Rheum Dis* 2005;64(11):1588-94.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
13. Palmer KT, Syddall H, Cooper C, Coggon D. Smoking and musculoskeletal disorders: findings from a British national survey. *Ann Rheum Dis* 2003;62(1):33-6.
14. Holth HS, Werpen HK, Zwart JA, Hagen K. Physical inactivity is associated with chronic musculoskeletal complaints 11 years later: results from the Nord-Trøndelag Health Study. *BMC Musculoskelet Disord* 2008;9:159.
15. McBeth J, Jones K. Epidemiology of chronic musculoskeletal pain. *Best Pract Res Clin Rheumatol* 2007;21(3):403-25.
16. Roquelaure Y, Ha C, Rouillon C, Fouquet N, Leclerc A, Descatha A, et al. Risk factors for upper-extremity musculoskeletal disorders in the working population. *Arthritis Rheum* 2009;61(10):1425-34.
17. Hakkinen A, Sokka T, Kotaniemi A, Hannonen P. A randomized two-year study of the effects of dynamic strength training on muscle strength, disease activity, functional capacity, and bone mineral density in early rheumatoid arthritis. *Arthritis Rheum* 2001;44(3):515-22.
18. Baker KR, Nelson ME, Felson DT, Layne JE, Sarno R, Roubenoff R. The efficacy of home based progressive strength training in older adults with knee osteoarthritis: a randomized controlled trial. *J Rheumatol* 2001;28(7):1655-65.
19. Hamberg-van Reenen HH, Ariens GA, Blatter BM, van Mechelen W, Bongers PM. A systematic review of the relation between physical capacity and future low back and neck/shoulder pain. *Pain* 2007;130(1-2):93-107.
20. Leino P, Aro S, Hasan J. Trunk muscle function and low back disorders: a ten-year follow-up study. *J Chronic Dis* 1987;40(4):289-96.
21. Kujala UM, Taimela S, Viljanen T, Jutila H, Viitasalo JT, Videman T, et al. Physical loading and performance as predictors of back pain in healthy adults. A 5-year prospective study. *Eur J Appl Physiol Occup Physiol* 1996;73(5):452-8.

- 1
2
3
4
5
6
7 22. Faber A, Sell L, Hansen JV, Burr H, Lund T, Holtermann A, et al. Does muscle strength predict
8 future musculoskeletal disorders and sickness absence? *Occupational medicine (Oxford,*
9 *England)* 2012;62(1):41-6.
10
11 23. Ruiz JR, Castro-Pinero J, Artero EG, Ortega FB, Sjostrom M, Suni J, et al. Predictive validity of
12 health-related fitness in youth: a systematic review. *Br J Sports Med* 2009;43(12):909-23.
13
14 24. Mikkelsen LO, Nupponen H, Kaprio J, Kautiainen H, Mikkelsen M, Kujala UM. Adolescent
15 flexibility, endurance strength, and physical activity as predictors of adult tension neck, low
16 back pain, and knee injury: a 25 year follow up study. *Br J Sports Med* 2006;40(2):107-13.
17
18 25. Barnekow-Bergkvist M, Hedberg GE, Janlert U, Jansson E. Determinants of self-reported neck-
19 shoulder and low back symptoms in a general population. *Spine (Phila Pa 1976)*
20 1998;23(2):235-43.
21
22 26. Neovius M, Sundstrom J, Rasmussen F. Combined effects of overweight and smoking in late
23 adolescence on subsequent mortality: nationwide cohort study. *BMJ* 2009;338:b496.
24
25 27. Nilsson PM, Nyberg P, Ostergren PO. Increased susceptibility to stress at a psychological
26 assessment of stress tolerance is associated with impaired fetal growth. *Int J Epidemiol*
27 2001;30(1):75-80.
28
29 28. Hemmingsson T, Melin B, Allebeck P, Lundberg I. The association between cognitive ability
30 measured at ages 18-20 and mortality during 30 years of follow-up--a prospective
31 observational study among Swedish males born 1949-51. *Int J Epidemiol* 2006;35(3):665-70.
32
33 29. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. Strengthening the
34 Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for
35 reporting observational studies. *BMJ* 2007;335(7624):806-8.
36
37 30. Nordesjo LO, Schéle R. Validity of Ergometer Cycle Test and Measures of Isometric Muscle
38 Strength when Predicting some Aspects of Military Performance. *Swedish J Defence Med*
39 1974(10):11-23.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
31. Wolk A, Rossner S. Effects of smoking and physical activity on body weight: developments in Sweden between 1980 and 1989. *J Intern Med* 1995;237(3):287-91.
32. Sundquist J, Johansson SE. The influence of socioeconomic status, ethnicity and lifestyle on body mass index in a longitudinal study. *Int J Epidemiol* 1998;27(1):57-63.
33. Bygren LO, Konlaan BB, Johansson SE. Attendance at cultural events, reading books or periodicals, and making music or singing in a choir as determinants for survival: Swedish interview survey of living conditions. *BMJ* 1996;313(7072):1577-80.
34. Tornvall G. Assessment of physical capabilities. *Acta physiologica Scandinavica* 1963;58(Suppl 201):1-102.
35. Nordesjo LO. A comparison between the Thornvall maximal ergometer test, submaximal ergometer tests and maximal oxygen uptake. *Swedish J Defence Med* 1974(10):3-10.
36. Nordesjo LO. Estimation of the maximal work rate sustainable for 6 minutes using a single-level load or stepwise increasing loads. *Ups J Med Sci* 1974;79(1):45-50.
37. Heneweer H, Vanhees L, Picavet HS. Physical activity and low back pain: a U-shaped relation? *Pain* 2009;143(1-2):21-5.
38. Heneweer H, Picavet HS, Staes F, Kiers H, Vanhees L. Physical fitness, rather than self-reported physical activities, is more strongly associated with low back pain: evidence from a working population. *Eur Spine J* 2012;21(7):1265-72.
39. Kujala UM, Kettunen J, Paananen H, Aalto T, Battie MC, Impivaara O, et al. Knee osteoarthritis in former runners, soccer players, weight lifters, and shooters. *Arthritis Rheum* 1995;38(4):539-46.
40. Almeida SA, Williams KM, Shaffer RA, Brodine SK. Epidemiological patterns of musculoskeletal injuries and physical training. *Med Sci Sports Exerc* 1999;31(8):1176-82.
41. Simoneau JA, Bouchard C. Genetic determinism of fiber type proportion in human skeletal muscle. *FASEB J* 1995;9(11):1091-5.

- 1
2
3
4
5
6
7 42. Costill DL, Daniels J, Evans W, Fink W, Krahenbuhl G, Saltin B. Skeletal muscle enzymes and fiber
8 composition in male and female track athletes. *J Appl Physiol* 1976;40(2):149-54.
9
10 43. Tesch P, Karlsson J. Isometric strength performance and muscle fibre type distribution in man.
11
12 *Acta physiologica Scandinavica* 1978;103(1):47-51.
13
14 44. Mannion AF, Weber BR, Dvorak J, Grob D, Muntener M. Fibre type characteristics of the lumbar
15 paraspinal muscles in normal healthy subjects and in patients with low back pain. *J Orthop*
16
17 *Res* 1997;15(6):881-7.
18
19 45. Barnekow-Bergkvist M, Hedberg G, Janlert U, Jansson E. Prediction of physical fitness and
20 physical activity level in adulthood by physical performance and physical activity in
21
22 adolescence--an 18-year follow-up study. *Scand J Med Sci Sports* 1998;8(5 Pt 1):299-308.
23
24
25
26
27

28 Tables and Figures

29
30
31 **Table 1:** Description of study sample

32
33 **Table 2:** Risk estimates for the muscle strength in youth as a determinant of self-reported
34 musculoskeletal pain in adulthood

35
36 **Figure 1:** The identification of the study sample and the loss to follow-up
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Description of study sample

Number of men	5489
Mean age at baseline (SD)	18.2 (0.5)
Mean time to follow-up (SD, range)	17.2 years (8.4, 1-35)
Mean hand grip* (SD)	617 (98)
Mean elbow flexion* (SD)	385 (83)
Mean knee extension* (SD)	567 (116)
Muscle Strength (%)	
Low	1371 (25.0)
Average	2747 (50.0)
High	1371 (25.0)
BMI (%)	
<18.5	477 (8.7)
18.5-24.9	4498 (81.9)
25-29.9	448 (8.2)
>30	66 (1.2)
Type of interview (%)	
In person	4349 (79.2)
By telephone	1140 (20.8)
Pain in back/hips (%)	
Yes	1645 (30.0)
of which severe	321 (5.8)
No	3842 (70.0)
Missing	2 (0.0)
Pain in neck/shoulders (%)	
Yes	1562 (28.5)
of which severe	246 (4.5)
No	3925 (71.5)
Missing	2 (0.0)
Pain in arms/legs (%)	
Yes	1243 (22.6)
of which severe	196 (3.6)
No	4243 (77.3)
Missing	3 (0.0)
Pain, independent of location (%)	
Yes	2847 (51.9)
of which severe	576 (10.5)
No	2639 (48.1)
Missing	3 (0.0)
Smoking status (%)	
Yes	827 (15.1)
No	4662 (84.9)
Level of education (%)	
Compulsory	589 (10.7)
Secondary	2889 (52.6)
Higher	2011 (36.6)
Physical Activity (%)	
Practically non	583 (10.6)
Now and then	1613 (29.4)
Regularly	2077 (37.8)
Regularly strenuous	1216 (22.2)

* In Newton.

Table 2 Risk estimates for the muscle strength in youth as a determinant of self-reported musculoskeletal pain in adulthood


Outcomes	Models				
	Univariate Muscle strength			Multivariate ^a Muscle strength	
	Low (N=1371) IRR (N)	Average (N=27475) Reference (N)	High (N=1371) IRR (N)	Low IRR (95% CI)	High IRR (95% CI)
Musculoskeletal pain	0.92 (668)	1 (1457)	0.99 (722)	0.93 (0.87-0.99)	0.99 (0.93-1.05)
Severe musculoskeletal pain	0.96 (135)	1 (283)	1.12 (158)	0.96 (0.79- 1.187)	1.078 (0.8990- 1.2930)
Pain in back/hips	0.92 (384)	1 (832)	1.03 (429)	0.93 (0.84- 1.034)	1.03 (0.94- 1.1344)
Pain in neck/shoulders	0.924 (366)	1 (799)	0.99 (397)	0.932 (0.83- 1.03)	1.00 (0.90-1.10)
Pain in arms/legs	0.97 (297)	1 (616)	1.07 (330)	0.97 (0.865- 1.10)	1.06 (0.94-1.19)

N= Number of cases

IRR= ~~incidence rate ratio~~Relative risk estimates

CI= Confidence interval

a= adjusted for smoking status, physical activity, education, body mass index



343x452mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cohort studies*

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5-6
Objectives	3	State specific objectives, including any prespecified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	6+
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6-9
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	6-7
		(b) For matched studies, give matching criteria and number of exposed and unexposed	-
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6-9
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	6-9
Bias	9	Describe any efforts to address potential sources of bias	9
Study size	10	Explain how the study size was arrived at	Figure 1
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	6-9
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	9
		(b) Describe any methods used to examine subgroups and interactions	9
		(c) Explain how missing data were addressed	-
		(d) If applicable, explain how loss to follow-up was addressed	-
		(e) Describe any sensitivity analyses	9
Results			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Figure 1
		(b) Give reasons for non-participation at each stage	Figure 1
		(c) Consider use of a flow diagram	Figure 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Table 1
		(b) Indicate number of participants with missing data for each variable of interest	Table 1
		(c) Summarise follow-up time (eg, average and total amount)	Table 1
Outcome data	15*	Report numbers of outcome events or summary measures over time	Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 1
		(b) Report category boundaries when continuous variables were categorized	6-9
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	-
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	9
Discussion			
Key results	18	Summarise key results with reference to study objectives	10-11
Limitations			
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	13-14
Generalisability	21	Discuss the generalisability (external validity) of the study results	12-13
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.



**Muscle Strength in Adolescent Men and Future
Musculoskeletal Pain:
A Cohort Study with 17 Years of Follow-up**

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-002656.R2
Article Type:	Research
Date Submitted by the Author:	21-Apr-2013
Complete List of Authors:	Timpka, Simon; Lund University, Department of Orthopedics Pettersson, Ingemar; Lund University, Department of Orthopedics Zhou, Caddie; Lund University, Department of Orthopedics Englund, Martin; Lund University, Department of Orthopedics; Boston University School of Medicine, Clinical Epidemiology Research & Training Unit
Primary Subject Heading:	Epidemiology
Secondary Subject Heading:	Sports and exercise medicine, Rheumatology, Public health
Keywords:	Musculoskeletal disorders < ORTHOPAEDIC & TRAUMA SURGERY, SPORTS MEDICINE, PREVENTIVE MEDICINE

SCHOLARONE™
Manuscripts

Muscle Strength in Adolescent Men and Future Musculoskeletal Pain:

A Cohort Study with 17 Years of Follow-up

Simon Timpka¹, Ingemar F Petersson¹, Caddie Zhou¹, Martin Englund^{1, 2}

1 Department of Orthopedics, Clinical Sciences Lund, Lund University, Lund, Sweden

2 Clinical Epidemiology Research & Training Unit, Boston University School of Medicine, Boston, Massachusetts, United States of America

Corresponding author:

Simon Timpka
Skånes universitetssjukhus
Epi-centrum Skåne
Medicon Village
223 81 Lund
Sweden

Phone: +46 738 395 276
Mail: simon.timpka@med.lu.se

Words in main text: 2868
Number of figures: 1
Number of tables: 2

Abstract

Objectives

Musculoskeletal pain is highly prevalent throughout adulthood with a major impact on health, function and participation in the society. Still, the association between muscle strength and development of musculoskeletal pain is unclear. We aimed to study whether overall muscle strength in adolescent men is associated with self-reported musculoskeletal pain in adulthood.

Design

Cohort study with baseline data from the Swedish Conscription Register and outcome information from the random population-based Swedish Living Conditions Surveys.

Setting

Sweden, 1970-2005.

Participants

We studied 5489 men who at age 17-19 years tested their isometric muscle strength (hand grip, arm flexion, knee extension) during the compulsory conscription.

Outcome measures

The men were surveyed regarding self-reported musculoskeletal pain; mean follow-up time of 17 (range 1-35) years. Our primary outcome was a self-report of musculoskeletal pain, and secondary outcomes were a report of "severe pain", "pain in back/hips", "pain in neck/shoulders", or "pain in arms/legs", respectively. We categorized muscle strength into three groups; low, average, and high using the 25th to 75th percentile to define the reference category (average). We estimated relative risks using log binomial regression with adjustment for smoking, body mass index, education, and physical activity.

Results

In the adjusted model, men with low overall muscle strength had decreased risk of self-reported musculoskeletal pain (0.93, 95% confidence interval 0.87-0.99). We observed no such association in men with high strength (0.99, 0.93-1.05). Furthermore, no statistically significant increase or decrease in risk was observed for any of the secondary outcomes.

Conclusion

In men, low overall isometric muscle strength in youth was not associated with an increased risk of future musculoskeletal pain. Contrarily, we observed a slightly decreased risk of self-reported musculoskeletal pain in adulthood. Our results does not support a model in which low muscle strength is a risk factor for future musculoskeletal pain.

Article summary

Article focus

- We aimed to study whether overall muscle strength in youth is inversely associated with the development of musculoskeletal pain in adulthood.

Key messages

- In contradiction to our expectations, we observed a decreased risk of self-reported musculoskeletal pain in adult men with low overall isometric muscle strength in youth. The study does not provide evidence in support of a theoretical model in which low muscle strength in young men is associated with an increased risk of musculoskeletal pain later in life.

Strengths and limitations of study

- The main strengths of the study are a large sample, the independent evaluation of exposure and outcome, the combined use of three different measures of muscle strength, and the comparably long time to follow-up.
- The main limitations of the study are the following; the cohort does not include women, musculoskeletal pain only identified with one question per site, the motivation for military service might influence performance in muscle strength testing, and the potential for unmeasured or residual confounding.

Introduction

Musculoskeletal disorders (MSDs), such as low back pain, osteoarthritis, and widespread pain, are highly prevalent in the adult population.¹⁻³ MSDs also contribute to a substantial burden of disease at middle and older ages.⁴ Although pain emanating from the musculoskeletal system might be attributed to a wide range of diseases with diverse causal chains, many MSDs have common risk factors such as heavy occupational work load,^{5,6} a high body mass index (BMI),⁷⁻⁹ and a low socio-economy.¹⁰⁻¹² Although smoking in some studies have been identified as a risk factor for certain MSDs,^{13,14} its main effect on musculoskeletal pain might be as an effect modifier of the pain sensation.¹⁵ As physical work load is a risk factor for many MSDs,¹⁶ a model in which the muscle strength in the loaded parts of the body are protective for future disorders is appealing. This is also the main rationale of studies in the area; does general or demarcated muscle strength have a protective effect on future complaints in the adjacent structures? It is also known that physical exercise with focus on muscle strength is an important secondary and tertiary prevention of MSDs.¹⁷ ¹⁸ A handful of studies have hitherto investigated the strength of isolated muscle groups as a determinant of later MSDs in adulthood.¹⁹⁻²² However, there is conflicting evidence of the value of muscle strength as a protective factor of musculoskeletal pain, such as neck/shoulder pain and low back pain in adult subjects.¹⁹

When considering muscle strength in youth as a potential risk factor in the longer perspective, its association with later disease of any kind is relatively unknown, including musculoskeletal pain.²³ At least two studies have investigated the result of single muscle strength tests as determinants of musculoskeletal complaints decades later.^{24,25} The first study, using number of sit-ups during 30 seconds as a strength measure, found no association with later low back pain or tension neck in men. In women however, the high strength group had a decreased odds ratio (OR) of tension neck.²⁴ The second study found a decreased OR for MSDs in men who either had a strong performance in isotonic bench press or in a isometric two hand lift test.²⁵ Neither of the two studies includes a

1
2
3 measure of overall muscular capacity. Hence, although there is some evidence of an association
4
5 between low muscle strength in youth and later risk of MSDs, the association between overall muscle
6
7 strength in adolescence and later musculoskeletal pain has never been studied. Furthermore, with a
8
9 larger sample size, data on common risk factors, testing of three different muscle groups, and data
10
11 on physical work capacity, we also address some of the limitations of earlier studies.
12

13
14 Our aim in this study was to investigate the general muscle strength in adolescent males as a
15
16 determinant of later self-reported musculoskeletal pain. We hypothesized that low general muscle
17
18 strength in youth is associated with an increased risk of having musculoskeletal pain in adulthood.
19

20 21 22 **Methods**

23
24 For this prospective register-based cohort study, we used two main inclusion criteria to identify the
25
26 study sample. First, when typically aged 18, the subjects should have performed mandatory
27
28 conscription testing in Sweden between 1970-1994, with the exception of the years 1978 and 1985.
29
30 Second, they should have been included in the Swedish Living Conditions Surveys any year between
31
32 1980 and 2005 when questions regarding musculoskeletal problems, smoking status, and physical
33
34 activity were simultaneously included.
35
36

37
38 We excluded all men who were surveyed prior to the baseline testing or were younger than 17 years
39
40 or older than 19 years at baseline (Table 1). We also excluded men with an existing musculoskeletal
41
42 disorder (Diseases of the musculoskeletal system and connective tissue according to the
43
44 International classification of disease version 8 or 9) and those who had missing data on variables
45
46 included in the primary model (muscle strength, smoking, BMI, physical activity, level of education).
47
48 In the final study sample we included 5489 men (figure 1). The Swedish conscription register is well
49
50 characterized and has been used for research purposes previously.²⁶⁻²⁸ At the time of study sample
51
52 testing, conscription was mandatory by law for all Swedish men. Specially trained employees at six
53
54 regional conscription offices administrated the conscription tests during a two-day session. The
55
56
57
58
59
60

1
2
3 procedure also included separate evaluations by a medical doctor and a psychologist. Only men with
4
5 serious health complaints were excused from conscription. The procedure included measurements of
6
7 each man's weight in underwear to the kilogram and height without shoes to the centimeter. Using
8
9 height and weight, we calculated BMI as height/kg². Probably due to rare errors of data entry, there
10
11 are unlikely extreme values in the dataset. Therefore, we excluded all subjects with registered
12
13 extreme values on height (<150, >210 cm), weight (<40, >150 kg), or an extreme calculated value for
14
15 BMI (<15, >60) (figure 1). The study was approved by the Ethical Review Board at Lund University
16
17 and the manuscript was prepared according to the STROBE-statement.²⁹

18 19 20 *Muscle strength*

21
22 The men performed three tests of isometric muscle strength during conscription; hand grip, elbow
23
24 flexion, and knee extension. At the start of test period in 1970, the tests were performed as
25
26 previously described³⁰ and remained unchanged in general throughout the test period. In summary,
27
28 hand grip strength on the preferred side was measured with 90° flexion at the elbow and the
29
30 humerus in parallel to the torso. Knee extension was measured in a sitting position with 90° knee
31
32 flexion and arms crossed over chest. The pelvis was fixed to the seat and a strap fastened above the
33
34 lateral malleolus. Also, elbow flexion was measured in a sitting position with 90° flexion at the elbow
35
36 and the humerus in parallel to the torso. A strap was fastened at the level of the radial styloid
37
38 process.

39
40
41 We calculated a measure of general muscle strength by standardizing and combining the three tests
42
43 of muscle strength. To avoid bias due to change of testing procedure over time, we categorized the
44
45 cohort into five subgroups based on period of conscription (1970-1973, 1974-1977, 1979-1984, 1986-
46
47 1990, 1990-1994) and for each subcohort calculated the relative muscle strength. First, we
48
49 standardized the three tests of muscle strength [standardized value= (value-mean)/standard
50
51 deviation] within each subgroup and used the mean of the three test scores as a proxy for general
52
53 muscle strength. Using percentiles, we then categorized the cohort into three groups of muscle
54
55
56
57
58
59
60

1
2
3 strength, where the 25th to 75th percentile defined the average category, the bottom 25th percentile
4
5 configured the low category, and the top 25th percentile defined the high muscle strength category.
6
7

8 **Survey of musculoskeletal pain**

9
10 The Swedish Living Conditions Surveys (ULF) is a random population based survey conducted by
11
12 Statistics Sweden, previously used for research purposes.³¹⁻³³ For the present study, we used data
13
14 collected during a total of 10 years (1980, 1988, 1989, 1997-1999, 2002-2005). The surveys were
15
16 generally performed as interviews in person by trained interviewers, with a minority of the
17
18 interviews performed by phone. For men included in more than one survey, we used the last survey
19
20 without relevant study data missing.
21
22

23 At follow-up, the men were asked three questions regarding any current musculoskeletal pain: 1) Do
24
25 you have pain in neck or shoulders? 2) Do you have back-pain, hip-pain or sciatica? 3) Do you have
26
27 ache, pain in hands, elbows, legs or knees? For each type of complaint one of three answers was
28
29 possible: 1) Yes, severe 2) Yes, mild and 3) No. Our primary outcome was having reported either
30
31 severe or mild musculoskeletal pain, whereas our secondary outcomes we defined as follows: 1)
32
33 Having reported severe musculoskeletal pain 2) Having reported pain in back/hips 3) Having reported
34
35 pain in shoulders/neck 4) Having reported pain in arms/legs. From the surveys, we also included data
36
37 on self-reported current smoking status (yes/no), physical activity (practically none, now and then,
38
39 regularly, regularly strenuous), and level of education (compulsory school or less, secondary
40
41 education, higher education). Drop-outs from the survey, i.e. those who have declined participation,
42
43 cannot be individually identified. However, during the years of survey used in this study, the
44
45 participation rate in the survey among men in relevant age groups were 70.0-88.7%.
46
47
48

49 **Statistical analyses**

50
51 All analyses were performed with SAS 9.3 (SAS Institute Inc). We used logistic binomial regression to
52
53 estimate relative risks (RR) and control potential confounders. In the multivariate model (primary
54
55 model), we included muscle strength, BMI, smoking status, physical activity and level of education.
56
57
58
59
60

Sensitivity analyses

To test whether cardiovascular aspects of physical capacity confounded our results, we used physical work capacity measured as $W_{\max 6}$ in a sensitivity analysis. For $W_{\max 6}$, the test result is an estimate of maximum work sustainable for 6 minutes³⁴ and is in young men correlated with maximum oxygen uptake ($r=0.9$).^{35 36} Acceptable data quality on work capacity was available in the subsample of men performing the baseline testing in 1976-1982. Out of all men in the cohort conscripted during the time period, 1154 men (74.6 %) completed an acceptable physical work capacity test on a bicycle ergometer (i.e. heart rate >174 at the end of testing). We added the work capacity in relation to body weight as a continuous variable to the univariate model. Furthermore, we also performed two sensitivity analyses on the multivariate model with musculoskeletal pain as the dependent variable. In the first, we added test center to the model. For the second, to test whether our categorization of muscle strength influenced the results, we used the standardized muscle strength both as a continuous variable and as a categorical variable based on quintiles.

Results

The mean time to follow-up was 17 years (table 1). Men with low muscle strength did not have an increased risk for the primary outcome "Musculoskeletal pain", but rather a statistically significant decreased risk (table 2). To summarize the observations of the secondary outcomes, we did not observe any statistically significant risk increases for neither men with a low nor high muscle strength. Compared to the crude model, the multivariate model produced similar risk estimates.

Sensitivity analyses

Work capacity had a significant effect in the subsample analysis ($p=0.03$) whereas it had only minor effect on the risk estimates for musculoskeletal pain, being 0.94 (95% confidence interval 0.82-1.08) and 1.02 (0.90-1.16) for low and high strength, respectively. The pattern of association in the secondary outcomes were in general somewhat strengthened when we adjusted for work capacity (data not shown). Using muscle strength as a continuous variable (hence assuming a linear relationship) did weaken the association with later musculoskeletal pain ($p=0.23$). When we instead used quintiles to categorize muscle strength, we observed no increased risk for the group with lowest strength compared to average strength (RR=0.93, 0.85-1.01).

Discussion

Investigating the general isometric muscle strength in adolescent men as a determinant of future musculoskeletal pain, we observed a decreased risk of self-reported musculoskeletal pain in men with low muscle strength. We also found a similar pattern, however not statistically significant, for "pain in back/hips" and "pain in neck/shoulders", whereas no association was found for future problems in arms/legs. Noticeably, the current study adds no support to a model in which low muscle strength in men is a risk factor for later musculoskeletal pain.

1
2
3 Using a historical cohort design with prospective registration of exposure and the outcome, the study
4
5 includes a large sample and thus allows better control for known confounders compared to previous
6
7 studies. All covariates (muscle strength, BMI, smoking, education, physical activity) included in the
8
9 multivariate model for musculoskeletal pain also had a significant association with the outcome.

10
11 Furthermore, by investigating the effects of physical work capacity in a sensitivity analysis, we aimed
12
13 to isolate the direct effect of muscle strength from other aspects of physical capacity. However, the
14
15 study also has important limitations. First and foremost, we have used strength data from military
16
17 conscription testing. Although it provides a rich dataset from a structured environment, we do not
18
19 know how the subjects' motivation for military service may have biased the performance during the
20
21 testing procedure. However, assuming there is no association between motivation at conscription
22
23 testing and later risk of musculoskeletal pain (or the loss to follow-up) any bias would at most dilute
24
25 our result. Nevertheless, stronger recruits are more likely to be assigned to positions with heavy load
26
27 duty. Secondly, the conscription was mandatory for men only. As the pattern of physical activity and
28
29 occupational exposure differ between men and women, any generalization of the results to women
30
31 must be made with great caution. Third, the physical activity measurement consisted of a single
32
33 question and did neither allow calculation of metabolic equivalent of task (MET) nor included
34
35 occupational exposure. Also musculoskeletal pain is measured by questions that combine more than
36
37 one site, decreasing the precision. However, the categories are fairly well demarcated anatomically
38
39 save for the question regarding pain in arms/legs. Fourth, the covariates collected with
40
41 musculoskeletal pain at follow-up (smoking, physical activity, level of education) are cross-sectional
42
43 and might thus be mediators of reverse causation. However, the adjusted estimates are much in line
44
45 with the crude estimates.
46
47
48
49

50
51 Primarily, we do not suggest that low muscle strength in youth is a protective factor for later
52
53 musculoskeletal pain. However, our observations could potentially be explained by both social and
54
55 biological factors. First, as former occupational exposure¹⁶ and certain sport participation³⁷ are
56
57 established risk factors for future MSDs, it lends some evidence for a general model in which certain
58
59
60

forms of physical activity is negative for the musculoskeletal health. It has also previously been suggested that there is a U-shaped association between physical activity and later back pain,^{38 39} i.e. that subjects with low and high levels of physical activity has an increased risk compared to the group with average activity. First, we speculate that our results may be explained by muscle strength in youth being one selection criterion for future high risk activities with a negative influence on the musculoskeletal health, e.g. higher risk of joint injury due to sports participation or manual repetitive work load. This would also include more immediate exposure in youth such as more physically demanding military service.⁴⁰ Although we have controlled for level of education, which we regard as a proxy for occupational exposure, there is potential for residual confounding. In other words, individuals with low general muscle strength might to a certain degree be deselected for high risk activities compared to stronger men.. A second potential explanation for our observation can be based on that, the strength of an individual is associated with the muscle fiber type distribution, which have a large genetic component.⁴¹ Whereas type I fibers are more common in endurance athletes⁴², a high type II percentage have been reported to be associated with both isometric muscle strength⁴³ as well as low back pain.⁴⁴

It is important to note that our main result are not the significantly decreased risk of later musculoskeletal pain observed in men with low strength but the nonexistent risk increase in the same group. This is partly in contrast with one of the few previous studies in the area. Although we in the present study only include measurements of isometric strength, the previous study²⁵ reported associations with both an isometric strength measure (static two hand lift) and an isotonic strength measure (bench press). In another study on the same cohort, it is reported that the result in bench press, but not two hand lift, was associated with both future cardiovascular fitness and future physical activity,⁴⁵ potentially explaining part of the difference. Another study reported flexibility as a sit and reach test,²⁴ but not strength measured as sit-ups, to be negatively associated with future risk of back pain. Hence, it is possible that other aspects of muscular and musculoskeletal function is of greater importance of future risk of MSDs than the isometric muscle strength as such.

1
2
3 The association between muscle strength and later musculoskeletal pain diminished when we used
4 muscle strength as a continuous variable. This was not surprising, as the observed association was
5 non linear in the primary model. It is to be expected that the test offices differ somewhat in their
6 reported test results, as they were assigned adolescents based on geography. However, including
7 test offices in the model did not have any effect on the overall association. When we included work
8 capacity in the model, most risk estimates decreased in absolute values, furthering strengthening the
9 observations in the primary model. By using the relative muscle strength during testing periods of
10 five years, we partially address the potential systematic change in testing procedure over the years.
11 Although the methods of measurements have not changed at large, minor adjustments cannot be
12 excluded.
13

14
15
16 In future studies, it would be of interest to investigate if low muscle strength serves as a deselection
17 criterion for professions or types of leisure time physical activity with higher risk of acute injuries or
18 chronic physical overload, factors with negative impact on musculoskeletal health.. Since the physical
19 activity pattern and physical fitness profiles differ between men and women, further investigations
20 are also needed to investigate if similar associations can be found in women.
21

22
23
24 In conclusion, we observed no increased risk of self-reported musculoskeletal pain in adult men with
25 low overall isometric muscle strength in youth. Thus, this study add no support to a model in which
26 muscle strength is a risk factor for future musculoskeletal pain in men.
27

28 **Acknowledgements**

29
30 No author has a conflict of interest to report. The authors would like to thank Charlotte Bergknut for
31 the initial data management.
32

33 **Funding**

1
2
3 The authors are supported by grants from The Swedish Research Council, Skåne Regional Council,
4
5 and the Faculty of Medicine at Lund University. The funders neither influenced the study process nor
6
7 manuscript preparation.
8
9

10 **Author contributions**

11
12
13
14 ST proposed the study idea, collected data, designed the study, and drafted the manuscript. IP
15
16 designed the study and provided input during manuscript preparation. CZ designed the study and
17
18 performed the statistical analyses. ME designed the study, collected data, and provided input during
19
20 manuscript preparation. All authors approved the final version of the manuscript.
21
22

23 **Data sharing statement**

24
25
26
27 No additional data available.
28
29

30 **Funding**

31
32
33
34 None
35
36

37 **Competing Interests**

38
39
40
41 None
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Bergman S, Herrstrom P, Hogstrom K, et al. Chronic musculoskeletal pain, prevalence rates, and sociodemographic associations in a Swedish population study. *J Rheumatol* 2001;28(6):1369-77.
2. Picavet HS, Schouten JS. Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study. *Pain* 2003;102(1-2):167-78.
3. Urwin M, Symmons D, Allison T, et al. Estimating the burden of musculoskeletal disorders in the community: the comparative prevalence of symptoms at different anatomical sites, and the relation to social deprivation. *Ann Rheum Dis* 1998;57(11):649-55.
4. Murray CJ, Vos T, Lozano R, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2013;380(9859):2197-223.
5. van der Windt DA, Thomas E, Pope DP, et al. Occupational risk factors for shoulder pain: a systematic review. *Occup Environ Med* 2000;57(7):433-42.
6. Hoogendoorn WE, van Poppel MN, Bongers PM, et al. Physical load during work and leisure time as risk factors for back pain. *Scand J Work Environ Health* 1999;25(5):387-403.
7. Felson DT, Zhang Y, Hannan MT, et al. Risk factors for incident radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum* 1997;40(4):728-33.
8. Shiri R, Karppinen J, Leino-Arjas P, et al. The association between obesity and low back pain: a meta-analysis. *Am J Epidemiol* 2010;171(2):135-54.
9. Grotle M, Hagen KB, Natvig B, et al. Obesity and osteoarthritis in knee, hip and/or hand: an epidemiological study in the general population with 10 years follow-up. *BMC Musculoskeletal Disord* 2008;9:132.

10. Hagen K, Zwart JA, Svebak S, et al. Low socioeconomic status is associated with chronic musculoskeletal complaints among 46,901 adults in Norway. *Scand J Public Health* 2005;33(4):268-75.
11. Dalstra JA, Kunst AE, Borrell C, et al. Socioeconomic differences in the prevalence of common chronic diseases: an overview of eight European countries. *Int J Epidemiol* 2005;34(2):316-26.
12. Bengtsson C, Nordmark B, Klareskog L, et al. Socioeconomic status and the risk of developing rheumatoid arthritis: results from the Swedish EIRA study. *Ann Rheum Dis* 2005;64(11):1588-94.
13. Palmer KT, Syddall H, Cooper C, et al. Smoking and musculoskeletal disorders: findings from a British national survey. *Ann Rheum Dis* 2003;62(1):33-6.
14. Holth HS, Werpen HK, Zwart JA, et al. Physical inactivity is associated with chronic musculoskeletal complaints 11 years later: results from the Nord-Trøndelag Health Study. *BMC Musculoskelet Disord* 2008;9:159.
15. McBeth J, Jones K. Epidemiology of chronic musculoskeletal pain. *Best Pract Res Clin Rheumatol* 2007;21(3):403-25.
16. Roquelaure Y, Ha C, Rouillon C, et al. Risk factors for upper-extremity musculoskeletal disorders in the working population. *Arthritis Rheum* 2009;61(10):1425-34.
17. Hakkinen A, Sokka T, Kotaniemi A, et al. A randomized two-year study of the effects of dynamic strength training on muscle strength, disease activity, functional capacity, and bone mineral density in early rheumatoid arthritis. *Arthritis Rheum* 2001;44(3):515-22.
18. Baker KR, Nelson ME, Felson DT, et al. The efficacy of home based progressive strength training in older adults with knee osteoarthritis: a randomized controlled trial. *J Rheumatol* 2001;28(7):1655-65.
19. Hamberg-van Reenen HH, Ariens GA, Blatter BM, et al. A systematic review of the relation between physical capacity and future low back and neck/shoulder pain. *Pain* 2007;130(1-2):93-107.

- 1
2
3 20. Leino P, Aro S, Hasan J. Trunk muscle function and low back disorders: a ten-year follow-up study.
4
5 *J Chronic Dis* 1987;40(4):289-96.
6
7 21. Kujala UM, Taimela S, Viljanen T, et al. Physical loading and performance as predictors of back
8
9 pain in healthy adults. A 5-year prospective study. *Eur J Appl Physiol Occup Physiol*
10
11 1996;73(5):452-8.
12
13 22. Faber A, Sell L, Hansen JV, et al. Does muscle strength predict future musculoskeletal disorders
14
15 and sickness absence? *Occupational medicine (Oxford, England)* 2012;62(1):41-6.
16
17 23. Ruiz JR, Castro-Pinero J, Artero EG, et al. Predictive validity of health-related fitness in youth: a
18
19 systematic review. *Br J Sports Med* 2009;43(12):909-23.
20
21 24. Mikkelsen LO, Nupponen H, Kaprio J, et al. Adolescent flexibility, endurance strength, and
22
23 physical activity as predictors of adult tension neck, low back pain, and knee injury: a 25 year
24
25 follow up study. *Br J Sports Med* 2006;40(2):107-13.
26
27 25. Barnekow-Bergkvist M, Hedberg GE, Janlert U, et al. Determinants of self-reported neck-shoulder
28
29 and low back symptoms in a general population. *Spine (Phila Pa 1976)* 1998;23(2):235-43.
30
31 26. Neovius M, Sundstrom J, Rasmussen F. Combined effects of overweight and smoking in late
32
33 adolescence on subsequent mortality: nationwide cohort study. *BMJ* 2009;338:b496.
34
35 27. Nilsson PM, Nyberg P, Ostergren PO. Increased susceptibility to stress at a psychological
36
37 assessment of stress tolerance is associated with impaired fetal growth. *Int J Epidemiol*
38
39 2001;30(1):75-80.
40
41 28. Hemmingsson T, Melin B, Allebeck P, et al. The association between cognitive ability measured at
42
43 ages 18-20 and mortality during 30 years of follow-up--a prospective observational study
44
45 among Swedish males born 1949-51. *Int J Epidemiol* 2006;35(3):665-70.
46
47 29. von Elm E, Altman DG, Egger M, et al. Strengthening the Reporting of Observational Studies in
48
49 Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*
50
51 2007;335(7624):806-8.
52
53
54
55
56
57
58
59
60

- 1
2
3 30. Nordesjo LO, Schéle R. Validity of Ergometer Cycle Test and Measures of Isometric Muscle
4 Strength when Predicting some Aspects of Military Performance. *Swedish J Defence Med*
5 1974(10):11-23.
6
7
8
9
10 31. Wolk A, Rossner S. Effects of smoking and physical activity on body weight: developments in
11 Sweden between 1980 and 1989. *J Intern Med* 1995;237(3):287-91.
12
13 32. Sundquist J, Johansson SE. The influence of socioeconomic status, ethnicity and lifestyle on body
14 mass index in a longitudinal study. *Int J Epidemiol* 1998;27(1):57-63.
15
16 33. Bygren LO, Konlaan BB, Johansson SE. Attendance at cultural events, reading books or
17 periodicals, and making music or singing in a choir as determinants for survival: Swedish
18 interview survey of living conditions. *BMJ* 1996;313(7072):1577-80.
19
20 34. Tornvall G. Assessment of physical capabilities. *Acta physiologica Scandinavica* 1963;58(Suppl
21 201):1-102.
22
23 35. Nordesjo LO. A comparison between the Thornvall maximal ergometer test, submaximal
24 ergometer tests and maximal oxygen uptake. *Swedish J Defence Med* 1974(10):3-10.
25
26 36. Nordesjo LO. Estimation of the maximal work rate sustainable for 6 minutes using a single-level
27 load or stepwise increasing loads. *Ups J Med Sci* 1974;79(1):45-50.
28
29 37. Heneweer H, Picavet HS, Staes F, et al. Physical fitness, rather than self-reported physical
30 activities, is more strongly associated with low back pain: evidence from a working
31 population. *Eur Spine J* 2012;21(7):1265-72.
32
33 38. Kujala UM, Kettunen J, Paananen H, et al. Knee osteoarthritis in former runners, soccer players,
34 weight lifters, and shooters. *Arthritis Rheum* 1995;38(4):539-46.
35
36 39. Heneweer H, Vanhees L, Picavet HS. Physical activity and low back pain: a U-shaped relation?
37 *Pain* 2009;143(1-2):21-5.
38
39 40. Almeida SA, Williams KM, Shaffer RA, et al. Epidemiological patterns of musculoskeletal injuries
40 and physical training. *Med Sci Sports Exerc* 1999;31(8):1176-82.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 41. Simoneau JA, Bouchard C. Genetic determinism of fiber type proportion in human skeletal
4 muscle. *FASEB J* 1995;9(11):1091-5.
5
6
7 42. Costill DL, Daniels J, Evans W, et al. Skeletal muscle enzymes and fiber composition in male and
8 female track athletes. *J Appl Physiol* 1976;40(2):149-54.
9
10
11 43. Tesch P, Karlsson J. Isometric strength performance and muscle fibre type distribution in man.
12 *Acta physiologica Scandinavica* 1978;103(1):47-51.
13
14
15 44. Mannion AF, Weber BR, Dvorak J, et al. Fibre type characteristics of the lumbar paraspinal
16 muscles in normal healthy subjects and in patients with low back pain. *J Orthop Res*
17 1997;15(6):881-7.
18
19
20 45. Barnekow-Bergkvist M, Hedberg G, Janlert U, et al. Prediction of physical fitness and physical
21 activity level in adulthood by physical performance and physical activity in adolescence--an
22 18-year follow-up study. *Scand J Med Sci Sports* 1998;8(5 Pt 1):299-308.
23
24
25
26
27
28
29
30
31
32
33
34
35

36 Tables and Figures

37
38 **Table 1:** Description of study sample
39

40 **Table 2:** Risk estimates for the muscle strength in youth as a determinant of self-reported
41 musculoskeletal pain in adulthood
42
43

44 **Figure 1:** The identification of the study sample and the loss to follow-up
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Description of study sample

Number of men	5489
Mean age at baseline (SD)	18.2 (0.5)
Mean time to follow-up (SD, range)	17.2 years (8.4, 1-35)
Mean hand grip* (SD)	617 (98)
Mean elbow flexion* (SD)	385 (83)
Mean knee extension* (SD)	567 (116)
Muscle Strength (%)	
Low	1371 (25.0)
Average	2747 (50.0)
High	1371 (25.0)
BMI (%)	
<18.5	477 (8.7)
18.5-24.9	4498 (81.9)
25-29.9	448 (8.2)
>30	66 (1.2)
Type of interview (%)	
In person	4349 (79.2)
By telephone	1140 (20.8)
Pain in back/hips (%)	
Yes	1645 (30.0)
of which severe	321 (5.8)
No	3842 (70.0)
Missing	2 (0.0)
Pain in neck/shoulders (%)	
Yes	1562 (28.5)
of which severe	246 (4.5)
No	3925 (71.5)
Missing	2 (0.0)
Pain in arms/legs (%)	
Yes	1243 (22.6)
of which severe	196 (3.6)
No	4243 (77.3)
Missing	3 (0.0)
Pain, independent of location (%)	
Yes	2847 (51.9)
of which severe	576 (10.5)
No	2639 (48.1)
Missing	3 (0.0)
Smoking status (%)	
Yes	827 (15.1)
No	4662 (84.9)
Level of education (%)	
Compulsory	589 (10.7)
Secondary	2889 (52.6)
Higher	2011 (36.6)
Physical Activity (%)	
Practically non	583 (10.6)
Now and then	1613 (29.4)
Regularly	2077 (37.8)
Regularly strenuous	1216 (22.2)

* In Newton.

Table 2 Risk estimates for the muscle strength in youth as a determinant of self-reported musculoskeletal pain in adulthood

Outcomes	Models				
	Univariate Muscle strength			Multivariate ^a Muscle strength	
	Low (N=1371) RR (N)	Average (N=2747) Reference (N)	High (N=1371) RR (N)	Low RR (95% CI)	High RR (95% CI)
Musculoskeletal pain	0.92 (668)	1 (1457)	0.99 (722)	0.93 (0.87-0.99)	0.99 (0.93-1.05)
Severe musculoskeletal pain	0.96 (135)	1 (283)	1.12 (158)	0.96 (0.79-1.18)	1.07 (0.89-1.29)
Pain in back/hips	0.92 (384)	1 (832)	1.03 (429)	0.93 (0.84-1.03)	1.03 (0.94-1.13)
Pain in neck/shoulders	0.92 (366)	1 (799)	0.99 (397)	0.93 (0.83-1.03)	1.00 (0.90-1.10)
Pain in arms/legs	0.97 (297)	1 (616)	1.07 (330)	0.97 (0.86-1.10)	1.06 (0.94-1.19)

N= Number of cases

RR= Relative risk estimates

CI= Confidence interval

a= adjusted for smoking status, physical activity, education, body mass index

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Muscle Strength in Adolescent Men and Future Musculoskeletal Pain: A Cohort Study with 17 Years of Follow-up

Simon Timpka¹, Ingemar F Petersson¹, Caddie Zhou¹, Martin Englund^{1, 2}

1 Department of Orthopedics, Clinical Sciences Lund, Lund University, Lund, Sweden

2 Clinical Epidemiology Research & Training Unit, Boston University School of Medicine, Boston, Massachusetts, United States of America

Corresponding author:

Simon Timpka
Skånes universitetssjukhus
Epi-centrum Skåne
Medicon Village
223 81 Lund
Sweden

Phone: +46 738 395 276
Mail: simon.timpka@med.lu.se

Words in main text: 2868
Number of figures: 1
Number of tables: 2

Abstract

Objectives

Musculoskeletal pain is highly prevalent throughout adulthood with a major impact on health, function and participation in the society. Still, the association between muscle strength and development of musculoskeletal pain is unclear. We aimed to study whether overall muscle strength in adolescent men is associated with self-reported musculoskeletal pain in adulthood.

Design

Cohort study with baseline data from the Swedish Conscription Register and outcome information from the random population-based Swedish Living Conditions Surveys.

Setting

Sweden, 1970-2005.

Participants

We studied 5489 men who at age 17-19 years tested their isometric muscle strength (hand grip, arm flexion, knee extension) during the compulsory conscription.

Outcome measures

The men were surveyed regarding self-reported musculoskeletal pain; mean follow-up time of 17 (range 1-35) years. Our primary outcome was a self-report of musculoskeletal pain, and secondary outcomes were a report of "severe pain", "pain in back/hips", "pain in neck/shoulders", or "pain in arms/legs", respectively. We categorized muscle strength into three groups; low, average, and high using the 25th to 75th percentile to define the reference category (average). We estimated relative risks using log binomial regression with adjustment for smoking, body mass index, education, and physical activity.

Results

In the adjusted model, men with low overall muscle strength had decreased risk of self-reported musculoskeletal pain (0.93, 95% confidence interval 0.87-0.99). We observed no such association in men with high strength (0.99, 0.93-1.05). Furthermore, no statistically significant increase or decrease in risk was observed for any of the secondary outcomes.

Conclusion

In men, low overall isometric muscle strength in youth was not associated with an increased risk of future musculoskeletal pain. Contrarily, we observed a slightly decreased risk of self-reported

musculoskeletal pain in adulthood. Our results does not support a model in which low muscle strength is a risk factor for future musculoskeletal pain. Residual confounding by adult occupational exposures, leisure time physical activity level and psychosocial factors may have impacted on results.

Article summary

Article focus

- We aimed to study whether overall muscle strength in youth is inversely associated with the development of musculoskeletal pain in adulthood.

Key messages

- In contradiction to our expectations, we observed a decreased risk of self-reported musculoskeletal pain in adult men with low overall isometric muscle strength in youth.
- The study does not provide evidence in support of a theoretical model in which low muscle strength in young men is associated with an increased risk of musculoskeletal pain later in life.
- ~~We speculate that low muscle strength may serve as a deselection criterion for activities with high risk of acute injuries or chronic physical overload, factors with negative impact on musculoskeletal health.~~

Strengths and limitations of study

- The main strengths of the study are a large sample, the independent evaluation of exposure and outcome, the combined use of three different measures of muscle strength, and the comparably long time to follow-up.
- The main limitations of the study are the following; the cohort does not include women, musculoskeletal pain only identified with one question per site, the motivation for military service might influence ~~measurement~~ performance in-of muscle strength testing, and the potential for unmeasured or residual confounding.

Introduction

Musculoskeletal disorders (MSDs), such as low back pain, osteoarthritis, and widespread pain, are highly prevalent in the adult population.¹⁻³ MSDs also contribute to a substantial burden of disease at middle and older ages.⁴ Although pain emanating from the musculoskeletal system might be attributed to a wide range of diseases with diverse causal chains, many MSDs have common risk factors such as heavy occupational work load,^{5,6} a high body mass index (BMI),⁷⁻⁹ and a low socio-economy.¹⁰⁻¹² Although smoking in some studies have been identified as a risk factor for certain MSDs,^{13,14} its main effect on musculoskeletal pain might be as an effect modifier of the pain sensation.¹⁵ As physical work load is a risk factor for many MSDs,¹⁶ a model in which the muscle strength in the loaded parts of the body are protective for future disorders is appealing. This is also the main rationale of studies in the area; does general or demarcated muscle strength have a protective effect on future complaints in the adjacent structures? Furthermore, it is also known that physical exercise with focus on muscle strength is an important secondary and tertiary prevention of MSDs.^{17,18} A handful of studies have hitherto ~~longitudinally~~ investigated the strength of isolated muscle groups as a determinant of later MSDs in adulthood ~~as a determinant of later MSDs.~~¹⁹⁻²² However, ~~in adult subjects,~~ there is ~~for the time being~~ conflicting evidence of the value of muscle strength as a protective factor of musculoskeletal pain, such as neck/shoulder pain and low back pain in adult subjects.¹⁹ When considering muscle strength in youth as a potential risk factor in the longer perspective, relatively little is known about its the association with of muscle strength in youth and later disease of any kind is relatively unknown, including musculoskeletal pain.²³ ~~At least Ft~~ two studies have investigated the result of single muscle strength tests as determinants of musculoskeletal complaints decades later.^{24,25} ~~The first study,~~ using number of sit-ups during 30 seconds as a strength measure, found no association with later low back pain or tension neck in men. In women however, the high strength group had a decreased odds ratio (OR) of tension neck.²⁴ The second study found a

1
2
3 decreased OR for MSDs in men who either had a strong performance in isotonic bench press or in a
4
5 isometric two hand lift test.²⁵ Neither of the two studies includes a measure of overall muscular
6
7 capacity. Hence, although there is some evidence of an association between low muscle strength in
8
9 youth and later risk of MSDs, the association between overall muscle strength in adolescence and
10
11 later musculoskeletal pain has ~~never~~ been studied. Furthermore, with a larger sample size, data
12
13 on common risk factors, testing of three different muscle groups, and data on physical work capacity,
14
15 we also address some of the limitations of earlier studies.

16
17
18 Our aim in this study was to investigate the general muscle strength in adolescent males as a
19
20 determinant of later self-reported musculoskeletal pain. We hypothesized that low general muscle
21
22 strength in youth is associated with an increased risk of having musculoskeletal pain in adulthood.
23
24

25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Methods

For this prospective register-based cohort study, we used two main inclusion criteria to identify the study sample. First, when typically aged 18, the subjects should have performed mandatory conscription testing in Sweden between 1970-1994, with the exception of the years 1978 and 1985. Second, they should have been included in the Swedish Living Conditions Surveys any year between 1980 and 2005 when questions regarding musculoskeletal problems, smoking status, and physical activity were simultaneously included.

We excluded all men who were surveyed prior to the baseline testing or were younger than 17 years or older than 19 years at baseline (Table 1). We also excluded men with an existing musculoskeletal disorder (Diseases of the musculoskeletal system and connective tissue according to the International classification of disease version 8 or 9) and those who had missing data on variables included in the primary model (muscle strength, smoking, BMI, physical activity, level of education).

In the final study sample we included 5489 men (figure 1). The Swedish conscription register is well characterized and has been used for research purposes previously. Data from the Swedish

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

scription testing has been previously used for research purposes.²⁶⁻²⁸ During the period At the
time of the study sample testing, conscription was mandatory by law for all Swedish men. Specially
trained employees at six regional conscription offices administrated the conscription tests during a
two-day session. The procedure that also included separate evaluations by a medical doctor and a
psychologist. Only men with serious health complaints, were excused from conscription. The
procedure included measurements of each mansubject's weight in underwear to the kilogram and
height without shoes to the centimeter. Using height and weight, we calculated BMI as height/kg².
Probably due to rare errors of data entry, there are unlikely extreme values in the dataset. Therefore,
we excluded all subjects with registered extreme values on height (<150, >210 cm), weight (<40,
>150 kg), or an extreme calculated value for BMI (<15, >60) (figure 1). The study was approved by
the Ethical Review Board at Lund University and the manuscript was prepared according to the
STROBE-statement.²⁹

Muscle strength

The men performed three tests of isometric muscle strength during conscription; hand grip, elbow
flexion, and knee extension. At the start of test period in 1970, the tests were performed as
previously described³⁰ and remained unchanged in general throughout the test period. In summary,
hand grip strength on the preferred side was measured with 90° flexion at the elbow and the
humerus in parallel to the torso. Knee extension was measured in a sitting position with 90° knee
flexion and arms crossed over chest. The pelvis was fixed to the seat and a strap fastened above the
lateral malleolus. Also, elbow flexion was measured in a sitting position with 90° flexion at the elbow
and the humerus in parallel to the torso. A strap was fastened at the level of the radial styloid
process.

We calculated a measure of general muscle strength by standardizing and combining the three tests
of muscle strength. To avoid bias due to change of testing procedure over time, we categorized the
cohort into five subgroups based on period of conscription (1970-1973, 1974-1977, 1979-1984, 1986-
1990, 1990-1994) and for each subcohort calculated the relative muscle strength. First, ~~We~~

1
2
3 standardized the three tests of muscle strength [standardized value= (value–mean)/standard
4 deviation] within each subgroup and used the mean of the three test scores as a proxy for general
5 muscle strength. Using percentiles, we then categorized the cohort into three groups of muscle
6 strength. Using percentiles, we then categorized the cohort into three groups of muscle
7 strength, where the 25th to 75th percentile defined the average category, the bottom 25th percentile
8 configured the low category, and the top 25th percentile defined the high muscle strength category.
9
10
11
12

13 **Survey of musculoskeletal pain**

14
15 The Swedish Living Conditions Surveys (ULF) is a random population based survey conducted by
16 Statistics Sweden, previously used for research purposes.³¹⁻³³ For the present study, we used data
17 collected during a total of 10 years (1980, 1988, 1989, 1997-1999, 2002-2005). The surveys were
18 generally performed as interviews in person by trained interviewers, with a minority of the
19 interviews performed by phone. For men included in more than one survey, we used the last survey
20 without relevant study data missing.
21
22
23
24
25
26
27
28

29 At follow-up, the men were asked three questions regarding any current musculoskeletal pain: 1) Do
30 you have pain in neck or shoulders? 2) Do you have back-pain, hip-pain or sciatica? 3) Do you have
31 ache, pain in hands, elbows, legs or knees? For each type of complaint one of three answers was
32 possible: 1) Yes, severe 2) Yes, mild and 3) No. Our primary outcome was having reported either
33 severe or mild musculoskeletal pain, whereas our secondary outcomes we defined as follows: 1)
34 Having reported severe musculoskeletal pain 2) Having reported pain in back/hips 3) Having reported
35 pain in shoulders/neck 4) Having reported pain in arms/legs. From the surveys, we also included data
36 on self-reported current smoking status (yes/no), physical activity (practically none, now and then,
37 regularly, regularly strenuous), and level of education (compulsory school or less, secondary
38 education, higher education). Drop-outs from the survey, i.e. those who have declined participation,
39 cannot be individually identified. However, during the years of survey used in this study, the
40 participation rate in the survey among men in relevant age groups were 70.0-88.7%.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Statistical analyses

All analyses were performed with SAS 9.3 (SAS Institute Inc). We used logistic binomial regression to estimate relative risks (RR) and control potential confounders. In the multivariate model (primary model), we included muscle strength, BMI, smoking status, physical activity and level of education.

Sensitivity analyses

To test whether cardiovascular aspects of physical capacity confounded our results, we used physical work capacity measured as $W_{\max-6\text{min}}$ in a sensitivity analysis. For $W_{\max-6\text{min}}$, the test result is an estimate of maximum work sustainable for 6 minutes³⁴ and is in young men correlated with maximum oxygen uptake ($r=0.9$).^{35 36} Acceptable data quality on work capacity was available in the subsample of men performing the baseline testing in 1976-1982. Out of all men in the cohort conscripted during the time period, 1154 men (74.6 %) completed an acceptable physical work capacity test on a bicycle ergometer (i.e. heart rate >174 at the end of testing). We added the work capacity in relation to body weight as a continuous variable to the univariate model. Furthermore, we also performed two sensitivity analyses on the multivariate model with musculoskeletal pain as the dependent variable. In the first, we added test center to the model. For the second, to test whether our categorization of muscle strength influenced the results, we ~~1) treated~~ used the standardized muscle strength both as a continuous variable and 2) treated the standardized muscle strength as a categorical variable based on quintiles.

Results

The mean time to follow-up was 17 years (table 1). Men with low muscle strength did not have an increased risk for the primary outcome "Musculoskeletal pain", but rather a statistically significant decreased risk, ~~for the primary outcome "Musculoskeletal pain"~~ (table 2). To summarize the observations of the secondary outcomes, we did not observe any statistically significant risk increases for neither men with a low nor high muscle strength. Compared to the crude model, the multivariate model produced similar risk estimates.

Sensitivity analyses

Work capacity had a significant effect in the subsample analysis ($p=0.03$) whereas it had only minor effect on the risk estimates for musculoskeletal pain, being 0.94 (95% confidence interval 0.82-1.08) and 1.02 (0.90-1.16) for low and high strength, respectively. The pattern of association in the secondary outcomes were in general somewhat strengthened when we adjusted for work capacity (data not shown). Using muscle strength as a continuous variable (hence assuming a linear relationship) did weaken the association with later musculoskeletal pain ($p=0.23$). When we instead used quintiles to categorize muscle strength, we observed no increased risk for the group with lowest strength compared to average strength (RR=0.93, 0.85-1.01).

Discussion

Investigating the general overall isometric muscle strength in adolescent men as a determinant of future musculoskeletal pain, we observed a decreased risk of self-reported musculoskeletal pain in men with low muscle strength. ~~No such association was observed for men with high strength.~~ We also found a similar pattern, however not statistically significant, ~~pattern~~ for "pain in back/hips" and "pain in neck/shoulders", whereas no association was found for future problems in arms/legs.

Noticeably, the current study adds no support to a model in which low muscle strength in men is a

~~risk factor for later musculoskeletal pain. The observed associations were in contradiction to our expectations.~~

Using a historical cohort design with prospective registration of exposure and the outcome, ~~the our~~ study includes a large sample and thus allows better control for known confounders compared to previous studies. All covariates (muscle strength, BMI, smoking, education, physical activity) included in the multivariate model for musculoskeletal pain also had a significant association with the outcome. Furthermore, by ~~investigating the effects of controlling for physical~~ work capacity ~~in a~~ sensitivity analysis, we aimed to isolate the direct effect of muscle strength from other aspects of physical capacity. However, the study also has important limitations. First and foremost, we have used strength data from military conscription testing. Although it provides a rich dataset from a structured environment, we do not know how the subject's¹ motivation for military service may have biased the performance during the testing procedure. However, assuming there is no association between motivation at conscription testing and later risk of musculoskeletal pain (or the loss to follow-up) any bias would at most dilute our result. Nevertheless, stronger recruits are more likely to be assigned to positions with heavy load duty. Secondly, the conscription was mandatory for men only. As the pattern of physical activity and occupational exposure differ between men and women, any generalization of the results to women must be made with great caution. Third, the physical activity measurement consisted of a single question and did neither allow calculation of metabolic equivalent of task (MET) nor included occupational exposure. Also musculoskeletal pain is measured by questions that combine more than one site, decreasing the precision. However, the categories are fairly well demarcated anatomically save for the question regarding pain in arms/legs. Fourth, the covariates collected with musculoskeletal pain at follow-up (smoking, physical activity, level of education) are cross-sectional and might thus be mediators of reverse causation. However, the adjusted estimates are much in line with the crude estimates.

1
2
3 Primarily, we do not suggest that low muscle strength in youth is a protective factor for later
4 musculoskeletal pain. However, our observations could potentially be explained by both social and
5 biological factors. It has previously been suggested that there is a U-shaped association between
6 physical activity and later back pain.³⁷⁻³⁸ ~~Furthermore, a~~ First, as former occupational exposure¹⁶ and
7 certain sport participation³⁷ are established risk factors for future MSDs,^{16 39} it lends some evidence
8 for a ~~more~~ general model, in which certain forms of physical activity is negative for the
9 musculoskeletal health. It has also previously been suggested that there is a U-shaped association
10 between physical activity and later back pain,^{38 39} i.e. that subjects with low and high levels of physical
11 activity has an increased risk compared to the group with average activity. Primarily, our
12 observations do not support low muscle strength in youth as a risk factor for later musculoskeletal
13 pain. ~~First~~ Instead, we speculate that our results ~~can~~ may be explained by muscle strength in youth
14 being one selection criterion for future high risk activities with a negative influence on the
15 musculoskeletal health, e.g. higher risk of joint injury due to sports participation or manual repetitive
16 work load. This would also include more immediate exposure in youth such as more physically
17 demanding military service.⁴⁰ Although we have controlled for level of education, which we regard as
18 might serve as a proxy for occupational exposure, there is potential for residual confounding. ~~as we~~
19 ~~did not have more appropriate data.~~ In other words, individuals with low general muscle strength
20 might to a certain degree be deselected for high risk activities compared to stronger men. ~~with an~~
21 ~~average or high strength.~~ A second potential explanation for our observation can be based on
22 that ~~However, this is only one of many possible explanations. For example,~~ the strength of an
23 individual is associated with the muscle fiber type distribution, which have a large genetic
24 component.⁴¹ Whereas Type I fibers are more common in endurance athletes⁴², ~~whereas a~~ high type
25 II percentage have been reported to be associated with both isometric muscle strength⁴³ as well as
26 low back pain.⁴⁴ ~~Thus, our observations could potentially be explained by both social and biological~~
27 ~~factors.~~

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

It is important to note that our main result are not the significantly decreased risk of later musculoskeletal pain observed in men with low strength but the nonexistent risk increase in the same group. This is Ppartly in contrast with one of the fewa previous studiesy in the area.²⁵ ~~we did not observe a negative effect of low muscle strength on the risk future of musculoskeletal problems in men.~~ Although we in the present study only include measurements of isometric strength, the previous study²⁵ ~~reported~~observed associations with both an isometric strength measure (static two hand lift) and an isotonic strength measure (bench press). In another study on the same cohort, it is reported that the result in bench press, but not two hand lift, was associated with both future cardiovascular fitness and future physical activity,⁴⁵ potentially explaining part of the difference. Another study reported flexibility as a sit and reach test,²⁴ but not strength measured as sit-ups, to be negatively associated with future risk of back pain. Hence, it is possible that other aspects of muscular and musculoskeletal function is of greater importance of future risk of MSDs than the isometric muscle strength as such.

The association between muscle strength and later musculoskeletal pain diminished when we used muscle strength as a continuous variable. This was not surprising, as the observed association was non linear in the primary model. It is to be expected that the test offices differ somewhat in their reported test results, as they were assigned adolescents based on geography. However, including test offices in the model did not have any effect on the overall association. When we included work capacity in the model, most risk estimates decreased in absolute values, furthering strengthening the observations in the primary model. By using the relative muscle strength during testing periods of five years, we partially address the potential systematic change in testing procedure over the years. Although the methods of measurements have not changed at large, minor adjustments cannot be excluded.

In future studies, it would be of interest to investigate if low muscle strength serves as a deselection criterion for professions or types of leisure time physical activity with higher risk of acute injuries or

~~chronic physical overload, factors with negative impact on musculoskeletal health. Although we found no increased risk of future musculoskeletal problems in men with low muscle strength in adolescence, future studies need to better quantify the occupational exposure and leisure time physical activity.~~ Since the physical activity pattern and physical fitness profiles differ between men and women, further investigations are also needed to investigate if similar associations can be found in women.

~~In contrast to our hypothesis~~ In conclusion, we observed ~~no~~ an ~~inde~~creased risk of self-reported musculoskeletal pain in adult men with low overall isometric muscle strength in youth. Thus, this study add no support to a model in which muscle strength is a risk factor for future musculoskeletal pain in men. ~~We speculate that low muscle strength per se is not protective of future musculoskeletal pain. Instead, low muscle strength might serve as a deselection criterion for professions or types of leisure time physical activity with higher risk of acute injuries or chronic physical overload, factors with negative impact on musculoskeletal health.~~

Acknowledgements

No author has a conflict of interest to report. The authors would like to thank Charlotte Bergknut for the initial data management.

Funding

The authors are supported by grants from The Swedish Research Council, Skåne Regional Council, and the Faculty of Medicine at Lund University. The funders neither influenced the study process nor manuscript preparation.

Author contributions

ST proposed the study idea, collected data, designed the study, and drafted the manuscript. IP designed the study and provided input during manuscript preparation. CZ designed the study and

performed the statistical analyses. ME designed the study, collected data, and provided input during manuscript preparation. All authors approved the final version of the manuscript.

Data sharing statement

There is no additional data to share.

References

1. Bergman S, Herrstrom P, Hogstrom K, Petersson IF, Svensson B, Jacobsson LT. Chronic musculoskeletal pain, prevalence rates, and sociodemographic associations in a Swedish population study. *J Rheumatol* 2001;28(6):1369-77.
2. Picavet HS, Schouten JS. Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC(3)-study. *Pain* 2003;102(1-2):167-78.
3. Urwin M, Symmons D, Allison T, Brammah T, Busby H, Roxby M, et al. Estimating the burden of musculoskeletal disorders in the community: the comparative prevalence of symptoms at different anatomical sites, and the relation to social deprivation. *Ann Rheum Dis* 1998;57(11):649-55.
4. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2013;380(9859):2197-223.
5. van der Windt DA, Thomas E, Pope DP, de Winter AF, Macfarlane GJ, Bouter LM, et al. Occupational risk factors for shoulder pain: a systematic review. *Occup Environ Med* 2000;57(7):433-42.
6. Hoogendoorn WE, van Poppel MN, Bongers PM, Koes BW, Bouter LM. Physical load during work and leisure time as risk factors for back pain. *Scand J Work Environ Health* 1999;25(5):387-403.

- 1
2
3 7. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, et al. Risk factors for incident
4
5 radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum*
6
7 1997;40(4):728-33.
- 8
9 8. Shiri R, Karppinen J, Leino-Arjas P, Solovieva S, Viikari-Juntura E. The association between obesity
10
11 and low back pain: a meta-analysis. *Am J Epidemiol* 2010;171(2):135-54.
- 12
13 9. Grotle M, Hagen KB, Natvig B, Dahl FA, Kvien TK. Obesity and osteoarthritis in knee, hip and/or
14
15 hand: an epidemiological study in the general population with 10 years follow-up. *BMC*
16
17 *Musculoskelet Disord* 2008;9:132.
- 18
19 10. Hagen K, Zwart JA, Svebak S, Bovim G, Jacob Stovner L. Low socioeconomic status is associated
20
21 with chronic musculoskeletal complaints among 46,901 adults in Norway. *Scand J Public*
22
23 *Health* 2005;33(4):268-75.
- 24
25 11. Dalstra JA, Kunst AE, Borrell C, Breeze E, Cambois E, Costa G, et al. Socioeconomic differences in
26
27 the prevalence of common chronic diseases: an overview of eight European countries. *Int J*
28
29 *Epidemiol* 2005;34(2):316-26.
- 30
31 12. Bengtsson C, Nordmark B, Klareskog L, Lundberg I, Alfredsson L. Socioeconomic status and the
32
33 risk of developing rheumatoid arthritis: results from the Swedish EIRA study. *Ann Rheum Dis*
34
35 2005;64(11):1588-94.
- 36
37 13. Palmer KT, Syddall H, Cooper C, Coggon D. Smoking and musculoskeletal disorders: findings from
38
39 a British national survey. *Ann Rheum Dis* 2003;62(1):33-6.
- 40
41 14. Holth HS, Werpen HK, Zwart JA, Hagen K. Physical inactivity is associated with chronic
42
43 musculoskeletal complaints 11 years later: results from the Nord-Trøndelag Health Study.
44
45 *BMC Musculoskelet Disord* 2008;9:159.
- 46
47 15. McBeth J, Jones K. Epidemiology of chronic musculoskeletal pain. *Best Pract Res Clin Rheumatol*
48
49 2007;21(3):403-25.
- 50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 16. Roquelaure Y, Ha C, Rouillon C, Fouquet N, Leclerc A, Descatha A, et al. Risk factors for upper-
4 extremity musculoskeletal disorders in the working population. *Arthritis Rheum*
5 2009;61(10):1425-34.
6
7
8
9 17. Hakkinen A, Sokka T, Kotaniemi A, Hannonen P. A randomized two-year study of the effects of
10 dynamic strength training on muscle strength, disease activity, functional capacity, and bone
11 mineral density in early rheumatoid arthritis. *Arthritis Rheum* 2001;44(3):515-22.
12
13
14 18. Baker KR, Nelson ME, Felson DT, Layne JE, Sarno R, Roubenoff R. The efficacy of home based
15 progressive strength training in older adults with knee osteoarthritis: a randomized
16 controlled trial. *J Rheumatol* 2001;28(7):1655-65.
17
18
19 19. Hamberg-van Reenen HH, Ariens GA, Blatter BM, van Mechelen W, Bongers PM. A systematic
20 review of the relation between physical capacity and future low back and neck/shoulder
21 pain. *Pain* 2007;130(1-2):93-107.
22
23
24 20. Leino P, Aro S, Hasan J. Trunk muscle function and low back disorders: a ten-year follow-up study.
25
26
27
28
29
30
31
32
33 21. Kujala UM, Taimela S, Viljanen T, Jutila H, Viitasalo JT, Videman T, et al. Physical loading and
34 performance as predictors of back pain in healthy adults. A 5-year prospective study. *Eur J*
35
36
37
38
39
40 22. Faber A, Sell L, Hansen JV, Burr H, Lund T, Holtermann A, et al. Does muscle strength predict
41 future musculoskeletal disorders and sickness absence? *Occupational medicine (Oxford,*
42
43
44
45
46 23. Ruiz JR, Castro-Pinero J, Artero EG, Ortega FB, Sjostrom M, Suni J, et al. Predictive validity of
47 health-related fitness in youth: a systematic review. *Br J Sports Med* 2009;43(12):909-23.
48
49
50 24. Mikkelsen LO, Nupponen H, Kaprio J, Kautiainen H, Mikkelsen M, Kujala UM. Adolescent
51 flexibility, endurance strength, and physical activity as predictors of adult tension neck, low
52
53
54
55
56
57
58
59
60

- 1
2
3 25. Barnekow-Bergkvist M, Hedberg GE, Janlert U, Jansson E. Determinants of self-reported neck-
4
5 shoulder and low back symptoms in a general population. *Spine (Phila Pa 1976)*
6
7 1998;23(2):235-43.
8
9
10 26. Neovius M, Sundstrom J, Rasmussen F. Combined effects of overweight and smoking in late
11
12 adolescence on subsequent mortality: nationwide cohort study. *BMJ* 2009;338:b496.
13
14 27. Nilsson PM, Nyberg P, Ostergren PO. Increased susceptibility to stress at a psychological
15
16 assessment of stress tolerance is associated with impaired fetal growth. *Int J Epidemiol*
17
18 2001;30(1):75-80.
19
20 28. Hemmingsson T, Melin B, Allebeck P, Lundberg I. The association between cognitive ability
21
22 measured at ages 18-20 and mortality during 30 years of follow-up--a prospective
23
24 observational study among Swedish males born 1949-51. *Int J Epidemiol* 2006;35(3):665-70.
25
26 29. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. Strengthening the
27
28 Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for
29
30 reporting observational studies. *BMJ* 2007;335(7624):806-8.
31
32
33 30. Nordesjo LO, Schéle R. Validity of Ergometer Cycle Test and Measures of Isometric Muscle
34
35 Strength when Predicting some Aspects of Military Performance. *Swedish J Defence Med*
36
37 1974(10):11-23.
38
39
40 31. Wolk A, Rossner S. Effects of smoking and physical activity on body weight: developments in
41
42 Sweden between 1980 and 1989. *J Intern Med* 1995;237(3):287-91.
43
44 32. Sundquist J, Johansson SE. The influence of socioeconomic status, ethnicity and lifestyle on body
45
46 mass index in a longitudinal study. *Int J Epidemiol* 1998;27(1):57-63.
47
48
49 33. Bygren LO, Konlaan BB, Johansson SE. Attendance at cultural events, reading books or
50
51 periodicals, and making music or singing in a choir as determinants for survival: Swedish
52
53 interview survey of living conditions. *BMJ* 1996;313(7072):1577-80.
54
55 34. Tornvall G. Assessment of physical capabilities. *Acta physiologica Scandinavica* 1963;58(Suppl
56
57 201):1-102.
58
59
60

- 1
2
3 35. Nordesjo LO. A comparison between the Thornvall maximal ergometer test, submaximal
4 ergometer tests and maximal oxygen uptake. *Swedish J Defence Med* 1974(10):3-10.
5
6
7 36. Nordesjo LO. Estimation of the maximal work rate sustainable for 6 minutes using a single-level
8 load or stepwise increasing loads. *Ups J Med Sci* 1974;79(1):45-50.
9
10
11 ~~37. Heneweer H, Vanhees L, Picavet HS. Physical activity and low back pain: a U-shaped relation?~~
12 ~~*Pain* 2009;143(1-2):21-5.~~
13
14 378. Heneweer H, Picavet HS, Staes F, Kiers H, Vanhees L. Physical fitness, rather than self-reported
15 physical activities, is more strongly associated with low back pain: evidence from a working
16 population. *Eur Spine J* 2012;21(7):1265-72.
17
18
19
20
21 389. Kujala UM, Kettunen J, Paananen H, Aalto T, Battie MC, Impivaara O, et al. Knee osteoarthritis in
22 former runners, soccer players, weight lifters, and shooters. *Arthritis Rheum* 1995;38(4):539-
23 46.
24
25
26
27
28 397. Heneweer H, Vanhees L, Picavet HS. Physical activity and low back pain: a U-shaped relation?
29 *Pain* 2009;143(1-2):21-5.
30
31
32
33
34
35 40. Almeida SA, Williams KM, Shaffer RA, Brodine SK. Epidemiological patterns of musculoskeletal
36 injuries and physical training. *Med Sci Sports Exerc* 1999;31(8):1176-82.
37
38
39 41. Simoneau JA, Bouchard C. Genetic determinism of fiber type proportion in human skeletal
40 muscle. *FASEB J* 1995;9(11):1091-5.
41
42
43 42. Costill DL, Daniels J, Evans W, Fink W, Krahenbuhl G, Saltin B. Skeletal muscle enzymes and fiber
44 composition in male and female track athletes. *J Appl Physiol* 1976;40(2):149-54.
45
46
47 43. Tesch P, Karlsson J. Isometric strength performance and muscle fibre type distribution in man.
48 *Acta physiologica Scandinavica* 1978;103(1):47-51.
49
50
51 44. Mannion AF, Weber BR, Dvorak J, Grob D, Muntener M. Fibre type characteristics of the lumbar
52 paraspinal muscles in normal healthy subjects and in patients with low back pain. *J Orthop*
53 *Res* 1997;15(6):881-7.
54
55
56
57
58
59
60

- 1
2
3 45. Barnekow-Bergkvist M, Hedberg G, Janlert U, Jansson E. Prediction of physical fitness and
4 physical activity level in adulthood by physical performance and physical activity in
5 adolescence--an 18-year follow-up study. *Scand J Med Sci Sports* 1998;8(5 Pt 1):299-308.
6
7
8
9
10
11
12

13 **Tables and Figures**

14
15
16 **Table 1:** Description of study sample

17
18 **Table 2:** Risk estimates for the muscle strength in youth as a determinant of self-reported
19 musculoskeletal pain in adulthood
20
21

22 **Figure 1:** The identification of the study sample and the loss to follow-up
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Description of study sample

Number of men	5489
Mean age at baseline (SD)	18.2 (0.5)
Mean time to follow-up (SD, range)	17.2 years (8.4, 1-35)
Mean hand grip* (SD)	617 (98)
Mean elbow flexion* (SD)	385 (83)
Mean knee extension* (SD)	567 (116)
Muscle Strength (%)	
Low	1371 (25.0)
Average	2747 (50.0)
High	1371 (25.0)
BMI (%)	
<18.5	477 (8.7)
18.5-24.9	4498 (81.9)
25-29.9	448 (8.2)
>30	66 (1.2)
Type of interview (%)	
In person	4349 (79.2)
By telephone	1140 (20.8)
Pain in back/hips (%)	
Yes	1645 (30.0)
of which severe	321 (5.8)
No	3842 (70.0)
Missing	2 (0.0)
Pain in neck/shoulders (%)	
Yes	1562 (28.5)
of which severe	246 (4.5)
No	3925 (71.5)
Missing	2 (0.0)
Pain in arms/legs (%)	
Yes	1243 (22.6)
of which severe	196 (3.6)
No	4243 (77.3)
Missing	3 (0.0)
Pain, independent of location (%)	
Yes	2847 (51.9)
of which severe	576 (10.5)
No	2639 (48.1)
Missing	3 (0.0)
Smoking status (%)	
Yes	827 (15.1)
No	4662 (84.9)
Level of education (%)	
Compulsory	589 (10.7)
Secondary	2889 (52.6)
Higher	2011 (36.6)
Physical Activity (%)	
Practically non	583 (10.6)
Now and then	1613 (29.4)
Regularly	2077 (37.8)
Regularly strenuous	1216 (22.2)

* In Newton.

Table 2 Risk estimates for the muscle strength in youth as a determinant of self-reported musculoskeletal pain in adulthood


Outcomes	Models				
	Univariate Muscle strength			Multivariate ^a Muscle strength	
	Low (N=1371) RR (N)	Average (N=2747) Reference (N)	High (N=1371) RR (N)	Low RR (95% CI)	High RR (95% CI)
Musculoskeletal pain	0.92 (668)	1 (1457)	0.99 (722)	0.93 (0.87-0.99)	0.99 (0.93-1.05)
Severe musculoskeletal pain	0.96 (135)	1 (283)	1.12 (158)	0.96 (0.79-1.18)	1.07 (0.89-1.29)
Pain in back/hips	0.92 (384)	1 (832)	1.03 (429)	0.93 (0.84-1.03)	1.03 (0.94-1.13)
Pain in neck/shoulders	0.92 (366)	1 (799)	0.99 (397)	0.93 (0.83-1.03)	1.00 (0.90-1.10)
Pain in arms/legs	0.97 (297)	1 (616)	1.07 (330)	0.97 (0.86-1.10)	1.06 (0.94-1.19)

N= Number of cases

RR= Relative risk estimates

CI= Confidence interval

a= adjusted for smoking status, physical activity, education, body mass index



90x118mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of cohort studies

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5-6
Objectives	3	State specific objectives, including any prespecified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	6+
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6-9
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	6-7
		(b) For matched studies, give matching criteria and number of exposed and unexposed	-
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6-9
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	6-9
Bias	9	Describe any efforts to address potential sources of bias	9
Study size	10	Explain how the study size was arrived at	Figure 1
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	6-9
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	9
		(b) Describe any methods used to examine subgroups and interactions	9
		(c) Explain how missing data were addressed	-
		(d) If applicable, explain how loss to follow-up was addressed	-
		(e) Describe any sensitivity analyses	9
Results			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Figure 1
		(b) Give reasons for non-participation at each stage	Figure 1
		(c) Consider use of a flow diagram	Figure 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Table 1
		(b) Indicate number of participants with missing data for each variable of interest	Table 1
		(c) Summarise follow-up time (eg, average and total amount)	Table 1
Outcome data	15*	Report numbers of outcome events or summary measures over time	Table 1
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 1
		(b) Report category boundaries when continuous variables were categorized	6-9
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	-
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	9
Discussion			
Key results	18	Summarise key results with reference to study objectives	10-11
Limitations			
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	13-14
Generalisability	21	Discuss the generalisability (external validity) of the study results	12-13
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>