

BMJ Open Measuring the relationship between interruptions, multitasking and prescribing errors in an emergency department: a study protocol

Magdalena Z Raban,¹ Scott R Walter,¹ Heather E Douglas,¹ Dana Strumpman,² John Mackenzie,³ Johanna I Westbrook¹

To cite: Raban MZ, Walter SR, Douglas HE, *et al*. Measuring the relationship between interruptions, multitasking and prescribing errors in an emergency department: a study protocol. *BMJ Open* 2015;**5**:e009076. doi:10.1136/bmjopen-2015-009076

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2015-009076>).

Received 13 June 2015
Revised 18 August 2015
Accepted 21 September 2015



CrossMark

¹Centre for Health Systems and Safety Research, Australian Institute of Health Innovation, Macquarie University, Sydney, New South Wales, Australia

²Pharmacy Department, Prince of Wales Hospital, Sydney, New South Wales, Australia

³Emergency Department, Prince of Wales Hospital, Sydney, New South Wales, Australia

Correspondence to

Dr Magdalena Z Raban;
magda.raban@mq.edu.au

ABSTRACT

Introduction: Interruptions and multitasking are frequent in clinical settings, and have been shown in the cognitive psychology literature to affect performance, increasing the risk of error. However, comparatively less is known about their impact on errors in clinical work. This study will assess the relationship between prescribing errors, interruptions and multitasking in an emergency department (ED) using direct observations and chart review.

Methods and analysis: The study will be conducted in an ED of a 440-bed teaching hospital in Sydney, Australia. Doctors will be shadowed at proximity by observers for 2 h time intervals while they are working on day shift (between 0800 and 1800). Time stamped data on tasks, interruptions and multitasking will be recorded on a handheld computer using the validated Work Observation Method by Activity Timing (WOMBAT) tool. The prompts leading to interruptions and multitasking will also be recorded. When doctors prescribe medication, type of chart and chart sections written on, along with the patient's medical record number (MRN) will be recorded. A clinical pharmacist will access patient records and assess the medication orders for prescribing errors. The prescribing error rate will be calculated per prescribing task and is defined as the number of errors divided by the number of medication orders written during the prescribing task. The association between prescribing error rates, and rates of prompts, interruptions and multitasking will be assessed using statistical modelling.

Ethics and dissemination: Ethics approval has been obtained from the hospital research ethics committee. Eligible doctors will be provided with written information sheets and written consent will be obtained if they agree to participate. Doctor details and MRNs will be kept separate from the data on prescribing errors, and will not appear in the final data set for analysis. Study results will be disseminated in publications and feedback to the ED.

INTRODUCTION

Since the publication of the Institute of Medicine's report *To Err Is Human* in 1999,

Strengths and limitations of this study

- This will be the first study to provide empirical evidence on the relationship between interruptions, multitasking and prescribing errors in situ.
- The study will use validated direct observation work measurement methods and prescribing error measurement to generate evidence.
- Although cognitive failures are similar across settings, the study will be carried out in one hospital potentially limiting the generalisability of the rates of errors to other settings.

there has been growing concern about the potential for medical errors due to the disruptive nature of clinical work environments.¹ This has led to the implementation of a variety of interventions to reduce interruptions during clinical work.²⁻⁴ The negative consequences of interruptions have been demonstrated in experimental studies in the psychology literature.⁵⁻⁸ Similarly, multitasking has been shown to have a negative effect on task performance in experimental studies,⁹⁻¹² notably during driving.¹³⁻¹⁵ However, while there are numerous descriptive studies detailing the frequency of interruptions and multitasking by health professionals in healthcare settings,¹⁶⁻²⁴ few studies have assessed the impact of interruptions or multitasking on clinical outcomes or errors.

One study, using direct observation of nurses during medication administration across six wards in two Australian hospitals, found that each interruption was associated with 12.1% and 12.7% increased odds of procedural and clinical medication administration errors.²⁵ Other studies have demonstrated interruptions that can result in changes in workflow and non-resumption of tasks.^{19 26-28} Frequent interruptions and

multitasking have also been found to be associated with increased self-reported strain in hospital doctors.^{21 29}

Prescribing errors are one indicator of clinical performance. A systematic review of prescribing errors in hospitals reported that they affect 7% of medication orders, 2% of patient days and 50% of hospital admissions.³⁰ Prescribing errors also occur in emergency departments (EDs), and have been reported at a rate of 21.5% and 12.5% per medication order for adult and paediatric patients, respectively.^{31 32} Prescribing represents a discrete task during which interruptions or multitasking behaviours can be directly observed. Since EDs are high-pressure environments, where doctors have been shown to be interrupted more frequently than ward doctors,^{20 26} they present an ideal setting for observing interruptions and multitasking during prescribing. In the context of a lack of evidence about the association of interruption and multitasking with errors in clinical practice, this study aims to assess the relationship between interruptions, multitasking and prescribing errors in an ED.

METHODS AND ANALYSIS

Study setting and participants

The study will be carried out in the acute section of an ED in a 440-bed teaching hospital in Sydney, Australia.³³ The study ED operates 24 h a day and in 2014 treated approximately 50 000 adult patients. The ED is organised in separate sections or 'models of care' in which patients of varying severity are treated. The acute section of the ED sees the most severe presentations, and includes three resuscitation bays and 12 beds. The resuscitation bays and beds are placed at the perimeter of the acute section, with a raised platform with computer terminals, termed 'the bridge', at the centre of the department. The bridge is where the majority of documentation and department management takes place.

All fully qualified ED doctors will be invited to participate in the study. A qualified doctor in Australia has completed a medical degree and 1 year of on the job training. This includes resident medical officers (RMO; 1–2 postgraduate years' experience), senior resident medical officers (SRMO; 2–3 postgraduate years' experience), registrars (at least 3 postgraduate years' experience and completion of relevant training) and staff specialists (at least 5 postgraduate years' experience and completion of relevant training). Written consent will be obtained from doctors who agree to participate. Interns, who have completed their university education but who have not yet obtained full medical registration, will not be invited as they may be less comfortable with the presence of an observer.

All qualified doctors listed above can prescribe medications. In Australian EDs, doctors prescribe medications that need to be administered in the ED, as well as on wards when a patient is admitted to the hospital.

In the acute section of the study ED, approximately 60–70% of patients are admitted to the hospital.

Direct observations

Doctors who consent to participate in the study will be observed for 2 h time intervals while they are working on day shift (beginning at 08:00 and ending at 18:00). Observers will shadow doctors at proximity in all areas of the ED, excluding the resuscitation bays, as required by the ethics study approval. Approximately equal numbers of doctors by seniority will be followed, and observation sessions will be spaced so that any time between 08:00 and 18:00 is covered by roughly the same number of sessions so as to be representative of ED work during the full time span of the day shift. An observational sampling matrix has been prepared to support this process.

Sophisticated time-and-motion methods will be applied to document doctors' work using the Work Observation Method by Activity Timing (WOMBAT) system.³⁴ WOMBAT provides data collection software which allows observers to record tasks undertaken during doctors' work using a handheld tablet computer. All tasks entered are automatically time-stamped. The task categories and their definitions to be used in this study are shown in [table 1](#), and have been adapted to this study setting based on other similar studies.^{23 26 34 35} For each task, the person with whom it is performed (staff specialist, registrar, resident or intern, nurse, patient, relative, ambulance officer, pharmacist, allied health, administrative personnel, security personnel or police, other, no one), the location, communication tools used (mobile phone, landline phone, computer, paper) and any interruptions or multitasking will be recorded.

When prescribing is observed, this will be recorded in the WOMBAT system as a prescribing task ([table 1](#)). In the ED, medication orders are handwritten on the following types of paper-based charts: (1) the Australian national inpatient medication chart (NIMC),³⁶ (2) fluid form, (3) heparin infusion chart, (4) insulin chart and (5) personally controlled analgesia form. For the NIMC, the chart sections include once only, when required, variable dose, venous thromboembolism (VTE) prophylaxis, warfarin, regular page one and regular page two. Medication orders written during observed prescribing tasks will be assessed for errors after the observational session. To facilitate this, details of which medication chart, sections and lines were completed during the observed prescribing task will be recorded in WOMBAT, and the patient medical record number (MRN) will be noted on a separate sheet. The MRNs for each subsequent prescribing task will be noted in consecutive order on the sheet during an observation session to enable linking to the observed, time-stamped prescribing tasks recorded in WOMBAT.

Observers will undergo extensive training in using the data collection tool. Prior to starting data collection, inter-rater reliability (IRR) testing will be undertaken

Table 1 Dimensions, categories and subcategories to be used for data collection in the WOMBAT system

| Task | Definition |
|--------------------------|--|
| Direct care | An activity directly related to the care of one patient for example, examining a patient, performing a procedure, communication with a patient or relative |
| Indirect care | An activity indirectly related to patient care for example, ordering tests, reading documents, washing hands, gathering and returning equipment |
| Documentation | Recording of patient-related information on paper or computer for example, writing patient notes |
| Clinical communication | Any discussion on clinical issues, excluding communication with patients/relatives for example, planning care with another health professional, requesting medical or nursing consultation, handover |
| Management communication | Discussion related to running of the department and other administrative issues for example, rosters, bed allocations, employment issues |
| Social communication | Conversations unrelated to work for example, personal phone calls |
| Prescribing | Writing one or more medication orders on a medication chart, form or prescription pad |
| In transit | Moving between areas of the department |
| On break | Breaks from work for example, eating lunch |

WOMBAT, Work Observation Method by Activity Timing.

and data collection will only begin once observers have reached high agreement. IRR will be reassessed at several time points during the data collection period to ensure consistency over time between observers.

Prescribing errors

A senior clinical pharmacist working at the study hospital and providing clinical services to the ED will review all medication orders observed during the observation sessions in order to identify prescribing errors. The pharmacist will be provided with the date and time an order was written, as well as the prescribing doctor's name, which will aid in identifying the correct order for assessment.

Following the protocol used for previous prescribing error studies, errors will be classified into three broad categories: clinical, procedural and legal errors.^{37 38} Online supplementary appendix 1 shows the types of errors within each category and their definitions. Each medication order can have more than one error. For example, an order may use an unapproved abbreviation for the dose, as well as have a wrong dose error. To assess a medication order for clinical errors, the pharmacist will access the patient's clinical notes, including reason for their ED visit, clinical history and pathology results. Once prescribing orders are assessed by the clinical pharmacist for errors, a researcher with pharmacy training will enter the details into a Microsoft Access database, double checking the consistency of coding. Each patient's year of birth, drug name, route, dose and frequency will also be entered into the database. Details of clinical errors will be noted for verification by a professor of clinical pharmacology.

The severity of the potential harm of prescribing errors identified, except legal errors, will be independently assessed by the clinical pharmacist and pharmacy researcher and rated as follows: (1) insignificant, that is, incident is likely to have little or no effect on the patient; (2) minor, that is, incident is likely to lead to

an increase in level of care (eg, review, investigations, or referral to another clinician); (3) moderate, that is, incident is likely to lead to permanent reduction in bodily functioning, increased length of stay, or surgical intervention; (4) major, that is, incident is likely to lead to a major permanent loss of function; (5) serious, that is, incident is likely to lead to death.³⁹ In cases where there is disagreement between the severity codes assigned and one of the assessors has assigned a code of moderate or higher severity, a final severity code will be assigned through discussion and consultation with an experienced clinical pharmacist independent of the study and if required a professor of clinical pharmacology.

Definitions of multitasking and interruptions

For this study multitasking is defined as performing two tasks simultaneously (in parallel). An interruption is defined as a response to an external stimulus (prompt) which causes the participant to stop the task they are engaged in (the primary task), and switch to the secondary (interrupting) task. This may also be described as task-switching, a term used in the cognitive psychology literature.⁴⁰ In this study, we will classify the nature of the observable external stimulus, which we refer to as a *prompt*. The type of prompt will be recorded as communication of a clinical, management or social nature (following the definitions in table 1), a phone call, public address (PA) announcement relevant to the observed doctor, equipment alarm or emergency alarm. If the prompt is communication, the person who initiated it will also be recorded.

Both multitasking and task-switching may be triggered by a prompt. For example, a nurse may approach a doctor while typing notes on a computer and ask about a patient's clinical management. This prompt may lead the doctor to stop typing and engage in a conversation with the nurse, which indicates he/she has chosen to task-switch. Alternatively the doctor may decide to

respond to the prompt by speaking with the nurse while continuing to type, thus multitasking. Additionally, possible responses to external prompts which will be captured in the study include *deferring* the prompt, for example, by replying they will speak to the nurse in 5 min, *deflecting* the prompt by telling the nurse to ask another doctor, or *acknowledging* the nurse's statement or question with a short reply of 'ok'. Both the prompt and response observed will be recorded in the WOMBAT system during the direct observations of doctors in the study. Recording the prompt and the responses in this way, allows for a more nuanced understanding of how doctors handle competing work demands, while maintaining comparability with previous study definitions.

Sample size and statistical analysis plan

During pilot testing, prescribing tasks were observed to occur about three times per hour with an average of two orders per prescribing task. Detectable effect sizes for differences in error rates were assessed for a sample size of 360 tasks (120 h of observation) across a range of plausible assumed values. Calculations were performed for multivariate Poisson regression with a main binary independent variable of interest (eg, was the prescribing task interrupted or not), assumed to occur in 50% of prescribing tasks, with the R^2 of the other covariates assumed to be in the range 0.1–0.5. This is a slightly simplified scenario, compared to the proposed analysis, in order to facilitate the sample size calculation. Error rates were allowed to vary from 0.2 to 0.5 per order consistent with clinical ED errors previously reported at around 20%³¹ and assuming legal errors will be more common. With significance set at 0.05 and power at 0.8, the detectable increase in error rate due to either task-switching or multitasking ranged from 23% to 53%.

Analyses will be conducted at the prescribing task level, where each task may include multiple orders. The prescribing error rate for a given prescribing task is the number of errors divided by the number of medication orders. Prescribing error rates will be calculated for clinical, procedural and legal prescribing errors separately, as well as for all types of errors combined. Multivariate Poisson regression will be applied to the number of errors per task with the number of orders per task as the offset (ie, denominator), hence modelling the error rate at task level. If there is a preponderance of prescribing tasks with no errors then a multivariate zero-inflated Poisson model will be applied. The main covariates of interest will be rates of task-switching and multitasking per unit time. Since the number of errors, task-switches and multitasks are proportional to task length it is necessary to include all of these in the model as rates. We will also adjust for other factors including patient age, doctor seniority and patient load at the time of the prescription. Patient load data will be extracted from the ED information management system.

A generalised estimating equations approach will be used to adjust SE estimates for clustering of outcomes within individual doctors. Sensitivity analyses will be performed in which task-switching and multitasking are included as the proportion of time spent performing a secondary task. Where sample size allows, a sub-analysis will apply the same Poisson model stratified by error severity.

Limitations

The 'Hawthorne' effect is when observed participants change their behaviour due to being under observation. Studies of this effect in clinicians suggest it is minimal,⁴¹ as does our prior research using direct observation of clinicians.³⁵ However, observers will be trained and conduct extensive IRR testing in the ED prior to starting the study, which will ensure staff in the ED are accustomed to being observed and seeing the observers in the department. If doctors were to change their behaviour, either by minimising their responses to external stimuli or being more diligent in prescribing, this will lead to an underestimation of studied effects.

Factors associated with the study setting need to be acknowledged when considering the generalisability of the study findings. First, the study ED has paper based prescribing, but electronic patient records. Settings with electronic prescribing systems are likely to show lower prescribing error rates than those seen in our study.³⁸ However, the evidence suggests that electronic prescribing is more likely to reduce the rates of procedural and legal errors, rather than clinical errors.³⁸ Our analysis plans to examine all errors, and procedural, legal and clinical error rates separately, and thus will provide evidence for comparison to settings with electronic prescribing. Additionally, our study will be based in one ED in Australia and only observe doctors on day shift. However, errors due to interruptions and multitasking, occur due to an overload on cognitive processing, irrespective of settings. The focus on day shift will limit variability in errors due to fatigue, which is linked to decreased cognitive function. Thus, though the prescribing errors described in our study may not be the same across all ED settings, their relationship with interruptions or multi-tasking should be relevant to other settings. Naturally, the frequency of prompts, interruptions and multi-tasking may also vary across settings, but several studies of clinicians in the ED report these at consistently high levels.^{16 20 42 43}

ETHICS AND DISSEMINATION

All participants meeting the inclusion criteria will be invited to participate and will be provided with a participant information sheet. Those agreeing to participate will provide written informed consent. Doctors participating in the study will be assigned a unique participant identifier. The list of doctors and identification numbers will be kept separate from the data collection. This

data-key list will be destroyed at the end of the study. Any prescribing error identified that could result in serious patient harm will be followed up by the clinical pharmacist as per hospital incident procedures and reporting policies. No patient names will be recorded, all MRNs collected as part of the study will be destroyed on study completion, and a de-identified code will replace MRNs in the final dataset. Ethics clearance for the study protocol has been obtained from the hospital human research ethics committee (reference number 13/310).

The study results will be disseminated through scientific publications and presentations. Aggregate feedback will be given to the staff within the ED in relation to prescribing errors and patterns of work identified.

Contributors JIW, SRW and JM conceptualised the direct observation of ED doctors work and SRW designed the initial protocol. JIW, SRW, MZR, HD, DS and JM then expanded the scope of the design to encompass prescribing errors. MZR and DS designed the prescribing record review process. MZR, HD and SRW devised the integration between data from record review and direct observation. MZR led the drafting of the manuscript. SRW designed and drafted the statistical analysis plan. All authors contributed to critical revisions of the manuscript and approved the final version for publication.

Funding SRW was supported by an Australian National Health and Medical Research Council Postgraduate Scholarship (1055498).

Competing interests None declared.

Ethics approval South Eastern Sydney Local Health District Human Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Institute of Medicine. *To err is human: building a safer health system*. Washington DC: National Academy Press, 1999.
- Raban MZ, Westbrook JI. Are interventions to reduce interruptions and errors during medication administration effective? A systematic review. *BMJ Qual Saf* 2014;23:414–21.
- Weigl M, Hoffmann F, Müller A, et al. Hospital paediatricians' workflow interruptions, performance, and care quality: a unit-based controlled intervention. *Eur J Pediatr* 2014;173:637–45.
- Weigl M, Hornung S, Glaser J, et al. Reduction of hospital physicians' workflow interruptions: a controlled unit-based intervention study. *J Healthc Eng* 2012;3:605–20.
- Gillie T, Broadbent D. What makes interruptions disruptive? A study of length, similarity, and complexity. *Psychol Res* 1989;50:243–50.
- Hodgetts HM, Jones DM. Interruption of the tower of London task: support for a goal-activation approach. *J Exp Psychol Gen* 2006;135:103–15.
- Foroughi CK, Werner NE, Nelson ET, et al. Do interruptions affect quality of work? *Hum Factors* 2014;56:1262–71.
- Li SYW, Blandford A, Cairns P, et al. The effect of interruptions on postcompletion and other procedural errors: an account based on the activation-based goal memory model. *J Exp Psychol Appl* 2008;14:314–28.
- Monsell S. Task switching. *Trends Cogn Sci* 2003;7:134–40.
- Pashler H. Task switching and multitask performance. In: Monsell S, Driver J, eds. *Control of cognitive processes: attention and performance XVIII*. USA: International Association for the Study of Attention and Performance, 2000:277–307.
- Rogers RD, Monsell S. Costs of a predictable switch between simple cognitive tasks. *J Exp Psychol Gen* 1995;124:207–31.
- Rubinstein JS, Meyer DE, Evans JE. Executive control of cognitive processes in task switching. *J Exp Psychol Hum Percept Perform* 2001;27:763–97.
- Klauer SG, Guo F, Simons-Morton BG, et al. Distracted driving and risk of road crashes among novice and experienced drivers. *N Engl J Med* 2014;370:54–9.
- Strayer DL, Drews FA. Multi-tasking in the automobile. In: Kramer AF, Weigmann D, Kirlik A, eds. *Attention: from theory to practice*. Oxford, UK: Oxford University Press, 2006:121–33.
- Olson R, Hanowski R, Hickman J, et al. *Driver distraction in commercial vehicle operations: federal motor carrier safety administration*. US Department of Transportation, 2009.
- Chisholm CD, Dornfeld AM, Nelson DR, et al. Work interrupted: a comparison of workplace interruptions in emergency departments and primary care offices. *Ann Emerg Med* 2001;38:146–51.
- Coiera EW, Jayasuriya RA, Hardy J, et al. Communication loads on clinical staff in the emergency department. *Med J Aust* 2002;176:415–18.
- Kuhn K, ed. *Multitasking by clinicians in the context of CPOE and CIS use*. Brisbane, Australia: Medinfo, 2007.
- Grundgeiger T, Sanderson P, MacDougall HG, et al. Interruption management in the intensive care unit: predicting resumption times and assessing distributed support. *J Exp Psychol Appl* 2010;16:317–44.
- Walter SR, Li L, Dunsmuir WTM, et al. Managing competing demands through task-switching and multitasking: a multi-setting observational study of 200 clinicians over 1000 hours. *BMJ Qual Saf* 2014;23:231–41.
- Weigl M, Müller A, Sevdalis N, et al. Relationships of multitasking, physicians' strain, and performance: an observational study in ward physicians. *J Patient Saf* 2013;9:18–23.
- Weigl M, Müller A, Zupanc A, et al. Participant observation of time allocation, direct patient contact and simultaneous activities in hospital physicians. *BMC Health Serv Res* 2009;9:110.
- Westbrook JI, Ampt A, Kearney L, et al. All in a day's work: an observational study to quantify how and with whom doctors on hospital wards spend their time. *Med J Aust* 2008;188:506.
- Westbrook JI, Duffield C, Li L, et al. How much time do nurses have for patients? A longitudinal study quantifying hospital nurses' patterns of task time distribution and interactions with health professionals. *BMC Health Serv Res* 2011;11:319.
- Westbrook JI, Woods A, Rob MI, et al. Association of interruptions with an increased risk and severity of medication administration errors. *Arch Intern Med* 2010;170:683.
- Westbrook JI, Coiera E, Dunsmuir WTM, et al. The impact of interruptions on clinical task completion. *Qual Saf Health Care* 2010;19:284–9.
- Drews FA. The frequency and impact of task interruptions in the ICU. *Proc Hum Factors Ergon Soc Ann Meeting* 2007;51:683–6.
- Laxmisan A, Hakimzada F, Sayan OR, et al. The multitasking clinician: decision-making and cognitive demand during and after team handoffs in emergency care. *Int J Med Inform* 2007;76:801–11.
- Weigl M, Müller A, Angerer P, et al. Workflow interruptions and mental workload in hospital paediatricians: an observational study. *BMC Health Serv Res* 2014;14:433.
- Lewis PJ, Dorman T, Taylor D, et al. Prevalence, incidence and nature of prescribing errors in hospital inpatients: a systematic review. *Drug Saf* 2009;32:379–89.
- Lifshitz AE, Goldstein LH, Sharist M, et al. Medication prescribing errors in the prehospital setting and in the ED. *Am J Emerg Med* 2012;30:726–31.
- Rinke ML, Moon M, Clark JS, et al. Prescribing errors in a pediatric emergency department. *Pediatr Emerg Care* 2008;24:1–8.
- South Eastern Sydney Local Health District—Government of New South Wales. About the Prince of Wales Hospital and Community Health Services. <http://www.seslhd.health.nsw.gov.au/POWH/about.asp>
- Ballermann MA, Shaw NT, Mayes DC, et al. Validation of the Work Observation Method By Activity Timing (WOMBAT) method of conducting time-motion observations in critical care settings: an observational study. *BMC Med Inform Decis Mak* 2011;11:32.
- Westbrook JI, Li L, Georgiou A, et al. Impact of an electronic medication management system on hospital doctors' and nurses' work: a controlled pre-post, time and motion study. *J Am Med Inform Assoc* 2013;20:1150–8.



36. Australian Commission on Safety and Quality in Health Care. NIMC for adults. <http://www.safetyandquality.gov.au/our-work/medication-safety/medication-chart/nimc/>
37. Reckmann M, Li L, Westbrook JI. *Protocol for measuring prescribing errors pre and post electronic medication management system introduction*. Unpublished Protocol, Kensington: Centre for Health Systems and Safety Research, University of New South Wales, 2011.
38. Westbrook JI, Reckmann M, Li L, *et al*. Effects of two commercial electronic prescribing systems on prescribing error rates in hospital in-patients: a before and after study. *PLoS Med* 2012;9:e1001164.
39. New South Wales Health Department. *Severity Assessment Code (SAC) Matrix*. Sydney: NSW Health, 2005.
40. Vandierendonck A, Liefoghe B, Verbruggen F. Task switching: interplay of reconfiguration and interference control. *Psychol Bull* 2010;136:601–26.
41. Dean B, Barber N. Validity and reliability of observational methods for studying medication administration errors. *Am J Health Syst Pharm* 2001;58:54–9.
42. Chisholm CD, Weaver CS, Whenmouth L, *et al*. A task analysis of emergency physician activities in academic and community settings. *Ann Emerg Med* 2011;58:117–22.
43. Chisholm CD, Collison EK, Nelson DR, *et al*. Emergency department workplace interruptions: are emergency physicians “interrupt-driven” and “multitasking”? *Acad Emerg Med* 2000;7:1239–43.

APPENDIX

Table A1: Classification of prescribing errors (from^{1,2})

| CLINICAL ERRORS | | |
|------------------------|--|---|
| Error category | Definition | Examples |
| Wrong drug | <p>Occurs when an inappropriate medication or IV fluid is prescribed</p> <p>e.g. the drug prescribed is not indicated for the patient's condition; the drug or IV fluid is contraindicated for a coexisting condition (drug-disease interaction); or an IV drug is prescribed with an incompatible diluent</p> <p>Note: Excludes generic substitution</p> | <p>e.g. hydrocortisone 25mg oral mane was prescribed instead of cortisone;</p> <p>chamomile lotion was ordered instead of calamine lotion</p> |
| Wrong dose / volume | <p>Occurs when the prescribed medication dose or IV fluid volume is higher or lower than that recommended for the condition, taking into account the patient's age, weight, renal and liver function</p> <p>May also occur when a dose is not altered in response to abnormal drug serum levels or laboratory tests</p> <p>Note: A dose may differ from normal recommended reference ranges and not be classed as an error where it is accepted practice to do so, i.e. the dose may have been queried by a pharmacist, but the specialist physician insisted on the prescribed dose, e.g. high dose flucloxacillin despite severe renal impairment in patients with severe infection when recommended by the infectious diseases team; low doses of tricyclic antidepressants initiated by the pain team.</p> | |
| Wrong rate / frequency | Occurs when the prescribed frequency of administration of a drug or an IV rate falls outside the recommended range | |
| Wrong route | Occurs when a medication is prescribed via an incorrect route of administration | <p>e.g. IV medication was prescribed orally;</p> <p>left eye was written instead of right eye</p> |
| Wrong formulation | Occurs when the wrong dosage form of a medication is ordered | e.g. an immediate release tablet was prescribed when an extended release form was |

| | | |
|-------------------------|---|---|
| | | required |
| Wrong timing | Occurs when a drug is prescribed at the wrong time of day | e.g. simvastatin prescribed in the morning instead of the evening (it is more efficacious when taken at night) |
| Wrong strength | Occurs when the prescribed drug strength is incorrect; the concentration of an IV infusion is prescribed incorrectly; or a dose is prescribed that does not exist or would not be able to be obtained easily from the current dose forms | e.g. mg was prescribed instead of micrograms (or vice versa) e.g. alendronate 75mg tab oral, take one tab once weekly (weekly dose only available as 70mg tablets) |
| Wrong patient | Occurs when a medication is prescribed for the wrong patient e.g. the prescriber writes a drug order intended for patient A on the medication chart belonging to patient B | |
| Not indicated | Occurs when a drug which is not indicated is prescribed for the patient; a drug is continued following a clinically significant adverse drug reaction; a drug which is no longer indicated is reordered; or a drug which should have been discontinued has not been ceased May also occur when a prescriber fails to cease/withhold a drug in response to abnormal drug serum levels or laboratory tests | e.g. fluticasone/ salmeterol inhaler prescribed for a patient without chronic obstructive airways disease e.g. an antibiotic which was not discontinued after completion of the course |
| Duplicated drug therapy | Occurs when two orders have been prescribed for one medication and both orders are active; there are two active orders for the same medication on two different charts; or the same drug is prescribed twice, as a single agent and as a combination product May also occur when two drugs are prescribed for the same indication when only one is necessary | e.g. one order was prescribed by generic and one by brand name e.g. ranitidine and omeprazole for gastro-oesophageal reflux disease |
| Drug-drug interaction | Occurs if two of the drugs prescribed for a patient are known to have a clinically significant interaction and this interaction is not acknowledged and monitored | |
| Allergy | Occurs when a drug is prescribed for a patient with a documented clinically significant allergy to that drug/class of drugs | |
| Inadequate monitoring | Occurs when the prescriber fails to order appropriate and timely clinical or laboratory tests to assess the patient's response to prescribed therapy Note: if adequate lab tests are ordered, but the results are not acted upon accordingly, resulting in potential or actual compromised patient care, | |

| | | |
|--------------------------|--|---|
| | this may be classed as wrong dose/volume error | |
| PROCEDURAL ERRORS | | |
| Unclear order | Occurs when the prescription is unclear or ambiguous e.g. the writing is illegible; or the order contains additional comments which apparently contradict the medication order | e.g. clotrimoxazole topical interdental BD (the prescriber was confused between cotrimoxazole and clotrimazole) |
| Incomplete order | Occurs when the order does not include all the necessary information i.e. drug name; strength (if appropriate); formulation (if appropriate); dose; route of administration; frequency ; the diluent for injectables; duration of time and/or rate of infusion (IV infusions); duration of time (IV fluids) | |
| LEGAL ERRORS | | |
| Legal/Procedural | Occurs when an aspect related to the prescription does not comply with the law, the NSW Department of Health or the hospital policy (and has not been assigned as an unclear order); the allergy field of the medication chart has not been completed; or the strength, dose, route or frequency of an existing handwritten medication order has been altered (such a change legally requires the entire order to be recharted) Use of unapproved abbreviations ³ and brand names instead of molecule names falls within this category | e.g. 'q4h' to convey 'every four hours' or 'µg' to convey 'micrograms' are considered unapproved abbreviations |

References

1. **Reckmann M, Li L, et al.** Protocol for Measuring Prescribing Errors Pre and Post Electronic Medication Management System Introduction. Unpublished Protocol, Kensington: Centre for Health Systems and Safety Research, University of New South Wales, 2011.
2. **Westbrook JI, Reckmann M, et al.** Effects of two commercial electronic prescribing systems on prescribing error rates in hospital in-patients: a before and after study. *PLoS Med.* 2012;9:e1001164.
3. **Australian Commission on Safety and Quality in Health Care.** Recommendations for Terminology, Abbreviations and Symbols used in the Prescribing and Administration of Medicines. Sydney: Australian Commission on Safety and Quality in Health Care - Government of Australia, 2011.