

BMJ Open

Cohort profile: Bushehr Elderly Health (BEH) Programme, Phase I (Cardiovascular System)

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2015-009597
Article Type:	Cohort profile
Date Submitted by the Author:	07-Aug-2015
Complete List of Authors:	<p>Ostovar, Afshin; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Nabipour, Iraj; Bushehr University of Medical Sciences, The Persian Gulf Biotechnology Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Iarijani, Bagher; Endocrinology & Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran.,</p> <p>Heshmat, Ramin; Tehran University of Medical Sciences, Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute</p> <p>Darabi, Hossein; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Vahdat, Katayoun; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Ravanipour, Maryam; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Mehrdad, Neda; Tehran University of Medical Sciences, Elderly Health Research Center, Endocrinology and Metabolism Population Sciences Institute</p> <p>Raeisi, Alireza; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Heidari, Gholamreza; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Shafiee, Gita; Tehran University of Medical Sciences, Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute</p> <p>Haeri, Mohammadjavad; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Pourbehi, Mohammadreza; Bushehr University of Medical Sciences, The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Sharifi, Farshad; Tehran University of Medical Sciences, Elderly Health Research Center, Endocrinology and Metabolism Population Sciences</p>

	<p>Institute Noroozi, Azita; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute Tahmasebi, Rahim; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute Aghaei Meybodi, Hamid Reza; Tehran University of Medical Sciences, Osteoporosis Research Center, Endocrinology and Metabolism Clinical Sciences Institute Assadi, Majid; Bushehr University of Medical Sciences, The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute Farrokhi, Shokrollah; Bushehr University of Medical Sciences, The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute Nemati, Reza; Bushehr University of Medical Sciences, The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute Amini, Mohammad Reza; Tehran University of Medical Sciences, Endocrinology & Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Barekat, Maryam; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute Amini, Abdullatif; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute Salimipour, Hooman; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute Dobaradaran, Sina; Bushehr University of Medical Sciences, The Persian Gulf Marine Biotechnology Research Center, The Persian Gulf Biomedical Sciences Research Institute Moshtaghi, Darab; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p>
Primary Subject Heading:	Cardiovascular medicine
Secondary Subject Heading:	Geriatric medicine
Keywords:	Cardiac Epidemiology < CARDIOLOGY, Coronary heart disease < CARDIOLOGY, Hypertension < CARDIOLOGY, Ischaemic heart disease < CARDIOLOGY, General diabetes < DIABETES & ENDOCRINOLOGY, Lipid disorders < DIABETES & ENDOCRINOLOGY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **TITLE:** Cohort profile: Bushehr Elderly Health (BEH) Programme, Phase I (Cardiovascular
4 System)
5
6
7
8
9

10
11 Afshin Ostovar¹; Iraj Nabipour^{2*}; Bagher Larijani³; Ramin Heshmat⁴; Hossein Darabi¹; Katayoun
12 Vahdat¹; Maryam Ravanipour¹; Neda Mehrdad⁵; Alireza Raeisi¹; Gholamreza Heidari¹; Gita
13 Shafiee⁴; Mohammadjavad Haeri¹; Mohammadreza Pourbehi²; Farshad Sharifi⁵; Azita Noroozi¹;
14 Rahim Tahmasebi¹; Hamidreza Aghaei Meybodi⁷; Majid Assadi⁶; Shokrollah Farrokhi⁶; Reza
15 Nemati⁶; Mohammad Reza Amini³; Maryam Barekat¹; Abdullatif Amini¹; Hooman Salimipour¹;
16 Sina Dobaradaran²; Darab Moshtaghi¹
17
18
19
20
21
22
23
24
25
26
27
28
29

30
31 ¹The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences
32 Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran
33
34

35
36 ²The Persian Gulf Biotechnology Research Center, The Persian Gulf Biomedical Sciences
37 Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran
38
39

40
41 ³Endocrinology & Metabolism Research Center, Endocrinology and Metabolism Clinical
42 Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran.
43
44

45
46 ⁴Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences
47 Institute, Tehran University of Medical Sciences, Tehran, Iran.
48
49

50
51 ⁵Elderly Health Research Center, Endocrinology and Metabolism Population Sciences Institute,
52 Tehran University of Medical Sciences, Tehran, Iran
53
54
55
56
57
58
59
60

1
2
3 ⁶The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences
4
5
6 Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran
7

8
9 ⁷Osteoporosis Research Center, Endocrinology and Metabolism Clinical Sciences Institute,
10
11 Tehran University of Medical Sciences, Tehran, Iran
12

13
14
15
16
17
18 **Correspondence:** Iraj Nabipour
19

20
21
22
23
24 **Email:** inabipour@gmail.com
25
26
27

28
29
30 **Postal Address:** The Persian Gulf Tropical Medicine Research Centre, No. 19, Boostan Alley,
31
32 Imam St., Bushehr, Iran
33
34

35 **Tel No:** +98 77 33341828
36
37
38
39

40
41 **Keywords:** Aged; Cardiovascular System; Risk factors; Cohort Studies; Iran
42
43
44

45
46 **Word Count:** 1738
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Purpose: The main objective of the Bushehr Elderly Health Programme, in its first phase, is to investigate the prevalence of cardiovascular risk factors and their association to major adverse cardiovascular events.

Participants: Between March 2013 and October 2014, a total of 3000 men and women aged \geq 60 years old residing in Bushehr, Iran participated in this prospective cohort study (participation rate=90.2%).

Findings to date: Baseline data on risk factors, including demographic and socio-economic status, smoking and medical history were collected through a modified WHO MONICA questionnaire. Vital signs and anthropometric measures, including systolic and diastolic blood pressure, weight, height, waist and hip circumference were also measured. A 12-lead electrocardiography and echocardiography were conducted on all participants, and total of 10cc venous blood was taken and sera was separated and stored at -80 degrees Centigrade for possible future use. Preliminary data analyses showed a noticeably higher prevalence of risk factors among older women compared to men.

Future plans: Risk factor assessments will be repeated every five years, and the participants will be followed during the study to measure the occurrence of major adverse cardiac events. Moreover, the second phase which includes investigation of the bone health and cognition in the elderly will be started in September 2015. Data are available at the Persian Gulf Biomedical Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran for any collaboration.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This is the first large sample prospective cohort study in Iran focusing on cardiovascular risk factors in the elderly as a growing population.
- The long-term follow-up in this study will allow for the assessment of many relevant outcomes.
- The participation rate in this study was high; however, outcome ascertainment may be incomplete because of incomplete death and major adverse cardiovascular events registries.

INTRODUCTION

The world's population is ageing rapidly. The proportion of those over the age of 60 will double, from about 11% to 22%, between 2000 and 2050. The absolute number of people aged 60 and over is expected to increase from 605 million to 2 billion over the same period.[1] According to the latest Iranian Census Data, about 8.2% of the population was over 60 years of age in 2011.[2] By the year 2020, the elderly population (over 60 years of age) is estimated to reach 10%.

Coronary heart disease (CHD) is one of the leading causes of disease burden in developing countries. In other words, three-fourths of global deaths due to CHD occurred in low- and middle-income countries.[3] In the year 2003, there were 21 572 Disability Adjusted Lost Years (DALY) due to all diseases and injuries per 100 000 Iranian people of all ages and both sexes. From this total number of DALYs, 58% were due to non-communicable diseases; ischemic heart disease was the fourth cause of DALYs in males and the first cause of DALYs in females.[4]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

There is a need for effective public health action on cardiovascular disease (CVD) prevention, especially in low- and middle-income countries, and for assessments of the cost-effectiveness of feasible interventions.[5] In addition, there is a need to try to balance the fight against the existing burden of infectious diseases with the growing epidemic of chronic diseases such as heart disease and diabetes.[6] The high costs of direct medical care and the indirect costs of cardiovascular disease, according to the American Heart Association, were approaching \$450 billion annually in 2010 and were projected to rise to over \$1 trillion annually by 2030.[7] Prevention of premature deaths due to non-communicable diseases and the reduction of related health care costs should be the main goals of health policy. Improving the detection and treatment of non-communicable diseases and preventing complications and catastrophic events from occurring should be the major goals of clinical medicine.[8] Trends in CVD risk factors and blood glucose will result in substantial CVD burden in developing countries and economies in transition in the near future. Periodic and consistent monitoring of trends and the effects of these risk factors on disease burden is needed for prioritising prevention programmes.[9] In recent decades, cohort studies have played a major role in investigating the incidence and causes of common health outcomes. As the major adverse cardiovascular events are quite common in old age, there are many cohort studies investigating these outcomes. A vast majority of these cohort studies are being conducted in developed countries.[10-20] However, the number of such prospective studies in the developing world is also rising.[21-25] In Iran, there are a few large prospective studies being conducted in various fields. Golestan's cohort study, a prospective study of oesophageal cancer in northern Iran,[26] the Tehran Lipid and Glucose Study (TLGS)[27] and the Shahrood Eye Cohort Study (ShECS)[28] are among these studies. The Amirkola Health and Ageing Project (AHAP) is the only large sample prospective cohort

1
2
3 study in Iran specifically aimed at investigating falling, bone fragility and fractures, cognitive
4
5 impairment and dementia, poor mobility and functional dependence in the elderly.[29]
6
7

8
9 This paper describes the rationale for, design and preliminary results of the Bushehr Elderly
10
11 Health programme (BEH Programme), a population-based prospective cohort study conducted in
12
13 Bushehr, a southern province of Iran. The main objective of this study, in its first phase, is to
14
15 investigate the prevalence of cardiovascular risk factors and their association to major adverse
16
17 cardiovascular events (MACE).
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

COHORT DESCRIPTION

The target population of this prospective cohort study was all men and women aged 60 years and over residing in the city of Bushehr. Based on the information available from the Bushehr District Health Centre, this population was estimated to be around 10 000. Table 1 shows eligibility criteria for participation in the study.

Table 1 List of inclusion and exclusion criteria of participants

Inclusion criteria

- Age more than or equal to 60
- Both sexes
- Residency in Bushehr port since at least 1 year prior to the recruitment
- No plan to leave Bushehr for the following 2 years after the recruitment
- Adequate physical and mental ability to participate in evaluation program
- Signing written informed consent

Exclusion criteria

- No residence in Bushehr
- No willingness to participate in the study

The participants in BEH Programme were selected through a multi-stage, stratified cluster random sampling method. Based on the classifications made by the municipality, as shown in Figure 1, we stratified Bushehr to 75 strata. Numbers were assigned to the blocks (as clusters) of each stratum and then randomly sorted. We invited all eligible older people residing in each

block selected to participate and then moved to the next block, repeating the invitation process, until gaining the sample required for that stratum. Sample sizes for strata were determined proportional to the number of households residing in each stratum. As shown in Figure 2, between March 2013 and October 2014, from among 3297 older people aged 60 years and over who were invited, a total of 3000 participated in this study (participation rate: 90.2%). Non-respondents were found among all strata, and there was no obvious pattern indicating selection bias. Baseline characteristics of participants are shown in Table 2.

Table 2 Baseline socio-demographic characteristics of participants in Bushehr Elderly Health Program

Characteristics [N (%)]	Men [1455 (48.5)]	Women [1545 (51.5)]	
Age Group	≤64	616 (42.3)	674 (43.6)
	65-69	317 (21.8)	378 (24.5)
	70-74	230 (15.8)	200 (12.9)
	75-79	166 (11.4)	181 (11.7)
	≥80	126 (8.7)	112 (7.2)
Marital status	Single	5 (0.3)	20 (1.3)
	Married	1378 (94.7)	884 (57.2)
	Widowed	68 (4.7)	619 (40.1)
	Divorced	4 (0.3)	22 (1.4)
Current occupation	Employed	133 (9.1)	23 (1.5)
	Retired	1195 (82.1)	126 (8.2)
	Unemployed	127 (8.7)	1396 (90.4)
Education	No education	315 (21.6)	777 (50.3)

Primary school	400 (27.5)	459 (29.7)
Secondary School	276 (19.0)	151 (9.8)
High school	287 (19.7)	125 (8.1)
University	177 (12.2)	33 (2.1)

*homemaker for female

The main follow-up assessments of risk factors will be done every five years for three consecutive periods (a total of 15 years of follow-up), in which all assessments will be repeated with comparable methods and modalities.

In addition, interim assessments will be made to ascertain outcomes and risk factor changes.

All participants will be contacted by a trained nurse once annually, and a checklist will be completed to check if any outcome of interest (major adverse cardiac events) has occurred during the 12 months prior to the assessment. A form has also been distributed to the participants to self-report as soon as possible after the occurrence of any of the targeted outcomes. Focal points in the two main hospitals in Bushehr (Shohadaye-Khalij-e-Fars and Salman-e-Farsi hospitals) are responsible for monthly checking of the hospital information system (HIS) and reporting admissions. If any outcome is reported, a general physician will review the in/outpatient medical records and documents, and detailed information will be entered into the special outcome forms. Our database will be cross-linked with the death registry system database, available via the public health system. The death registry system receives information from hospitals, forensic medicine departments, cemeteries and vital events records offices, and duplicate records are deleted. The International Classification of Diseases-10 (ICD-10), WHO online version, is used to classify causes of death. A person's national identification number is used as a unique identifier to cross-link databases. In the case of a death report but no reliable death certificate, a

1
2
3 verbal autopsy will be performed to determine cause of death. Table 3 presents list of outcomes
4
5
6 ascertained in the present study.
7

8
9 **Table 3 List of Outcomes, BEHP Study**

- 10
11 - Death
12 - Myocardial Infarction
13 - Stroke
14 - Cardiac Interventions
15 - Hospital admission
16
17
18
19

20
21
22
23 **Baseline examination**

24 A modified WHO MONICA questionnaire[30] translated into Persian was used as the core
25
26 questionnaire to gather baseline information on demographic and socio-economic variables and
27
28 risk factors. Table 4 presents the components of the core questionnaire.
29
30
31

32
33 **Table 4 Section and Topics of the Core Questionnaire in the BEHP Study**

34
35
36

Type of data	Components
Demographic	<ul style="list-style-type: none"> - Personal information (Name, Nickname, Surname) - National Identification Number - Age and Sex - Marital status - Contact Information
Socio-economic	<ul style="list-style-type: none"> - Employment status - Education level - Insurance - Family income - Family assets - Residence status

37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Cardiovascular risk factors	- Smoking status and history
	- Physical activity
	- Menstrual and menopause history
Medical history	- Blood pressure history
	- Diabetes history
	- Lipid Profile history
	- Ischemia and myocardial infarction history
	- Weakness, impaired sensation and stroke history
	- Rose and claudication history
	- Heart failure history
Drug history	- Administered by physician
	- Over the counter drugs
	- Supplements

Phlebotomy and laboratory analyses

Participants were asked to provide a venous blood sample for laboratory tests. A total of 10cc of whole blood were taken by a trained nurse after 8-12 hours of fasting. Table 5 presents laboratory tests done at the baseline and their methods of measurement. Sera were also separated and stored at -80 degrees Centigrade for possible future use.

Clinical assessment

A comprehensive physical examination, including vital signs, weight and height measurements and waist and hip circumference were taken at the baseline. A 12-lead electrocardiography, performed by a trained nurse, and echocardiography, carried out by a cardiologist, were conducted for all participants (see Table 5).

Table 5 Baseline Physical Examinations and Laboratory Tests, BEHP Study

	Item	Method of measurement
Physical Examination	Arterial Blood Pressure	Manually by Standard Mercury sphygmomanometer in sitting position
	Weight in cm	Stadiometer; Heavy outer garments will be removed.
	Height in cm	Stadiometer; Shoes will be removed.
	Waist circumference in cm	At the midway level between the costal margins and the iliac crests.
	Hip circumference in cm	At the level of the greater trochanters.
Laboratory tests	Complete Blood Count (CBC)	Automated Hematology Analyzer
	Fasting Blood Sugar	Enzymatic (glucose oxidase) colorimetric method using a commercial kit (Pars Azmun Inc., Tehran, Iran)
	Lipid Profile (Total cholesterol, LDL, HDL, triglyceride)	Serum total cholesterol and HDL cholesterol will be measured using a Cholesterol oxidase phenol aminoantipyrine and triglycerides using a glycerol-3 phosphate oxidase phenol aminoantipyrine enzymatic method. Serum LDL cholesterol will be calculated using the Friedwald formula; LDL cholesterol will not be calculated when triglycerides concentration is >400 mg/dl.
Procedures	Electrocardiography	12-lead, by a trained nurse
	Echocardiography	Using M-Turbo™ Ultrasound System, Manufactured by SonoSite, Inc. by cardiologist

FINDING TO DATE

Baseline data are being analysed, and preliminary findings are presented in Table 6. The prevalence of cardiovascular risk factors was remarkably higher among older women compared to men. Smoking behaviour was different among older men and women. Older women did not smoke cigarettes as often as men; however, the prevalence of hookah smoking was higher in women. One third of older women were obese, which was two times more than the prevalence of obesity among men. Metabolic syndrome was also about two times more prevalent among older women.

Table 6 Frequency of Risk Factors in Participants of Bushehr Elderly Health Program

Risk Factor [N(%)]	Men [1455(48.5)]	Women [1545(51.5)]
Smoking		
Hookah		
Non-smoker	1143 (78.6)	951 (61.6)
Current Smoker	110 (7.6)	267 (17.3)
Intermittent Smoker	4 (0.3)	7 (0.5)
Former Smoker	198 (13.6)	320 (20.7)
Cigarette		
Non- Smoker	1023 (70.3)	1522 (98.5)
Current Smoker	198 (13.6)	14 (0.9)
Intermittent Smoker	1 (0.1)	0 (0.0)
Former Smoker	233 (16.0)	9 (0.6)
Hypertension	820 (56.4)	1054 (68.2)
Diabetes Mellitus	390 (26.8)	506 (32.8)

Hypercholesterolemia		886 (60.9)	1034 (66.9)
BMI*	<25	615 (42.8)	414 (27.3)
	25-29.9	612 (42.6)	591 (39.0)
	≥30	211 (14.7)	511 (33.7)
Metabolic Syndrome**		403 (28.4)	826 (53.5)

*46 missing

**Based on the ATP III Criteria

STRENGTHS AND LIMITATIONS

This is the first large sample prospective cohort study in Iran focusing on cardiovascular risk factors in the elderly as a growing population. The long-term follow-up in this study will allow for the assessment of many relevant outcomes. The participation rate in this study was high; however, outcome ascertainment may be incomplete because of incomplete death and major adverse cardiovascular events registries.

COLLABORATIONS

This study is carrying out at The Persian Gulf Tropical Medicine Research Centre, which is affiliated with Bushehr University of Medical Sciences, in Bushehr, Iran. We welcome any collaboration, and researchers should contact AO via email, a.ostovar@bpums.ac.ir or IN at inabipour@gmail.com.

ETHICS AND DISSEMINATION

This study is being conducted in agreement with Declaration of Helsinki and in accordance with Iranian national guidelines for ethics in research. The protocol of the study was approved by the

1
2
3 regional research ethics committee of Bushehr University of Medical Sciences on 23 September
4
5 2013, Reference Number: B-91-14-2. All participants were asked to sign a written informed
6
7 consent, which was approved by the research ethics committee. The participants are able to
8
9 withdraw from the study at any time without any explanation. Data collected are stored in a
10
11 reidentifiable form by national ID code. The results will be presented at national and
12
13 international meetings and published in a peer-reviewed journal. We aim to translate the key
14
15 findings to an easily understandable format for local residents and to present them through local
16
17 media. Relevant findings will also be presented as policy briefs to national and local health
18
19 policy makers.
20
21
22
23

24 25 **Acknowledgments** 26

27
28 We are grateful to the staff of both research centres at BPUMS and TUMS for their commitment
29
30 to the study's protocol and objectives. We are also indebted to all participants who accepted our
31
32 invitation and patiently underwent exhausting measurements and examinations.
33
34
35

36 37 **Contributors** 38

39 AO drafted the manuscript, participated in study design and conduction, performed data analysis
40
41 and interpretation. IN conceived the study, helped draft the manuscript, and participated in the
42
43 study design and conduction, and data analysis and interpretation. BL participated in the study
44
45 design and conduction and reviewed the manuscript. RH helped draft the manuscript and
46
47 participated in the study design and conduction and data analysis and interpretation. HD, KV,
48
49 NM, GH, AR, and MH participated in the study design and reviewed the manuscript. MP, AA,
50
51 RN, HS, and MB participated in the study design and data collection and reviewed the
52
53 manuscript. MR, GS, FS, AN, and RT participated in questionnaire development, study design,
54
55
56
57
58
59
60

1
2
3 and staff training and reviewed the manuscript. HAM, MRA, SF, SD, and DM participated in
4
5 data collection and reviewed the manuscript.
6
7

8 9 **Funding**

10
11 The Persian Gulf Biomedical Sciences Research Institute affiliated with Bushehr (Port)
12
13 University of Medical Sciences (BPUMS) and the Endocrinology and Metabolism Research
14
15 Institute, affiliated with Tehran University of Medical Sciences (TUMS), have jointly provided
16
17 funding for this research project.
18
19

20 21 22 **Conflict of interest**

23
24
25 None declared.
26
27

28 29 **Data sharing statement**

30
31 Huge data have been collected. Access to the data is available for interested researchers from
32
33 corresponding author IN (inabipour@gmail.com) or AO (a.ostovar@bpums.ac.ir).
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Interesting facts about ageing. Secondary Interesting facts about ageing 2012. <http://www.who.int/ageing/about/facts/en/index.html>.
2. Selected Findings of National Population and Housing Census, 2011. In: Cooperations. PRAI, ed. Tehran: Statistical Center of Iran, 2012:27.
3. Gaziano TA, Bitton A, Anand S, et al. Growing epidemic of coronary heart disease in low- and middle-income countries. *Curr Probl Cardiol* 2010;**35**(2):72-115 doi: 10.1016/j.cpcardiol.2009.10.002[published Online First: Epub Date]].
4. Naghavi M, Abolhassani F, Pourmalek F, et al. The burden of disease and injury in Iran 2003. *Popul Health Metr* 2009;**7**:9 doi: 10.1186/1478-7954-7-9[published Online First: Epub Date]].
5. Labarthe DR, Dunbar SB. Global cardiovascular health promotion and disease prevention: 2011 and beyond. *Circulation* 2012;**125**(21):2667-76 doi: 10.1161/circulationaha.111.087726[published Online First: Epub Date]].
6. Shetty P. Grey matter: ageing in developing countries. *The Lancet* 2012;**379**(9823):1285-87
7. Weintraub WS, Daniels SR, Burke LE, et al. Value of primordial and primary prevention for cardiovascular disease: a policy statement from the American Heart Association. *Circulation* 2011;**124**(8):967-90 doi: 10.1161/CIR.0b013e3182285a81[published Online First: Epub Date]].
8. Hunter DJ, Reddy KS. Noncommunicable diseases. *N Engl J Med* 2013;**369**(14):1336-43 doi: 10.1056/NEJMra1109345[published Online First: Epub Date]].
9. Singh GM, Danaei G, Farzadfar F, et al. The Age-Specific Quantitative Effects of Metabolic Risk Factors on Cardiovascular Diseases and Diabetes: A Pooled Analysis. *PLoS one* 2013;**8**(7):e65174
10. Schoeni-Affolter F, Ledergerber B, Rickenbach M, et al. Cohort profile: the Swiss HIV Cohort study. *Int J Epidemiol* 2010;**39**(5):1179-89 doi: 10.1093/ije/dyp321[published Online First: Epub Date]].
11. Hofman A, van Duijn CM, Franco OH, et al. The Rotterdam Study: 2012 objectives and design update. *Eur J Epidemiol* 2011;**26**(8):657-86 doi: 10.1007/s10654-011-9610-5[published Online First: Epub Date]].
12. Mitchell GF, Hwang S-J, Vasan RS, et al. Arterial stiffness and cardiovascular events the Framingham Heart Study. *Circulation* 2010;**121**(4):505-11
13. Christiansen CB, Olesen JB, Gislason G, et al. Cardiovascular and non-cardiovascular hospital admissions associated with atrial fibrillation: a Danish nationwide, retrospective cohort study. *BMJ open* 2013;**3**(1)
14. Welmer AK, Angleman S, Rydwick E, et al. Association of cardiovascular burden with mobility limitation among elderly people: a population-based study. *PLoS One* 2013;**8**(5):e65815 doi: 10.1371/journal.pone.0065815[published Online First: Epub Date]].
15. Vu T-HT, Stamler J, Liu K, et al. Prospective Relationship of Low Cardiovascular Risk Factor Profile at Younger Ages to Ankle-Brachial Index: 39-Year Follow-Up—The Chicago Healthy Aging Study. *Journal of the American Heart Association* 2012;**1**(6)
16. Marmot M, Brunner E. Cohort Profile: the Whitehall II study. *Int J Epidemiol* 2005;**34**(2):251-6 doi: 10.1093/ije/dyh372[published Online First: Epub Date]].

17. Lahelma E, Aittomaki A, Laaksonen M, et al. Cohort profile: the Helsinki Health Study. *Int J Epidemiol* 2013;**42**(3):722-30 doi: 10.1093/ije/dys039[published Online First: Epub Date]].
18. Cumming RG, Handelsman D, Seibel MJ, et al. Cohort Profile: the Concord Health and Ageing in Men Project (CHAMP). *Int J Epidemiol* 2009;**38**(2):374-8 doi: 10.1093/ije/dyn071[published Online First: Epub Date]].
19. Magnus P, Irgens LM, Haug K, et al. Cohort profile: the Norwegian Mother and Child Cohort Study (MoBa). *Int J Epidemiol* 2006;**35**(5):1146-50 doi: 10.1093/ije/dyl170[published Online First: Epub Date]].
20. Osler M, Lund R, Kriebbaum M, et al. Cohort profile: the Metropolit 1953 Danish male birth cohort. *Int J Epidemiol* 2006;**35**(3):541-5 doi: 10.1093/ije/dyi300[published Online First: Epub Date]].
21. Dugee O, Oum S, Buckley BS, et al. Assessment of total cardiovascular risk using WHO/ISH risk prediction charts in three low and middle income countries in Asia. *BMC public health* 2013;**13**(1):539
22. Kok VC, Horng J-T, Lin H-L, et al. Gout and subsequent increased risk of cardiovascular mortality in non-diabetics aged 50 and above: a population-based cohort study in Taiwan. *BMC cardiovascular disorders* 2012;**12**(1):108
23. Victora CG, Barros FC. Cohort profile: the 1982 Pelotas (Brazil) birth cohort study. *Int J Epidemiol* 2006;**35**(2):237-42 doi: 10.1093/ije/dyi290[published Online First: Epub Date]].
24. Woodward M, Barzi F, Martiniuk A, et al. Cohort profile: the Asia Pacific Cohort Studies Collaboration. *Int J Epidemiol* 2006;**35**(6):1412-6 doi: 10.1093/ije/dyl222[published Online First: Epub Date]].
25. Jiang C, Thomas GN, Lam TH, et al. Cohort profile: The Guangzhou Biobank Cohort Study, a Guangzhou-Hong Kong-Birmingham collaboration. *Int J Epidemiol* 2006;**35**(4):844-52 doi: 10.1093/ije/dyl131[published Online First: Epub Date]].
26. Pourshams A, Khademi H, Malekshah AF, et al. Cohort Profile: The Golestan Cohort Study- a prospective study of oesophageal cancer in northern Iran. *Int J Epidemiol* 2010;**39**(1):52-9 doi: 10.1093/ije/dyp161[published Online First: Epub Date]].
27. Azizi F, Rahmani M, Emami H, et al. Cardiovascular risk factors in an Iranian urban population: Tehran lipid and glucose study (phase 1). *Soz Praventivmed* 2002;**47**(6):408-26
28. Fotouhi A, Hashemi H, Shariati M, et al. Cohort profile: shahroud eye cohort study. *Int J Epidemiol* 2013;**42**(5):1300-8 doi: 10.1093/ije/dys161[published Online First: Epub Date]].
29. Hosseini SR, Cumming RG, Kheirkhah F, et al. Cohort profile: the Amirkola Health and Ageing Project (AHAP). *International journal of epidemiology* 2013:dyt089
30. Bothig S. WHO MONICA Project: objectives and design. *Int J Epidemiol* 1989;**18**(3 Suppl 1):S29-37

1
2
3 **Figure 1 Map of Bushehr and Distribution of Participants in the Strata**
4
5

6 **Figure 2 Flowchart of enrolments**
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1 Map of Bushehr and Distribution of Participants in the Strata
286x432mm (96 x 96 DPI)




Figure 2 Flowchart of enrolments
140x150mm (120 x 120 DPI)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

BMJ Open

Cohort profile: Bushehr Elderly Health (BEH) Programme, Phase I (Cardiovascular System)

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2015-009597.R1
Article Type:	Cohort profile
Date Submitted by the Author:	22-Oct-2015
Complete List of Authors:	<p>Ostovar, Afshin; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Nabipour, Iraj; Bushehr University of Medical Sciences, The Persian Gulf Biotechnology Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Larjani, Bagher; Endocrinology & Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran.,</p> <p>Heshmat, Ramin; Tehran University of Medical Sciences, Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute</p> <p>Darabi, Hossein; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Vahdat, Katayoun; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Ravanipour, Maryam; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Mehrdad, Neda; Tehran University of Medical Sciences, Elderly Health Research Center, Endocrinology and Metabolism Population Sciences Institute</p> <p>Raeisi, Alireza; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Heidari, Gholamreza; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Shafiee, Gita; Tehran University of Medical Sciences, Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute</p> <p>Haeri, Mohammadjavad; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Pourbehi, Mohammadreza; Bushehr University of Medical Sciences, The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Sharifi, Farshad; Tehran University of Medical Sciences, Elderly Health Research Center, Endocrinology and Metabolism Population Sciences</p>

	<p>Institute Noroozi, Azita; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Tahmasebi, Rahim; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Aghaei Meybodi, Hamid Reza; Tehran University of Medical Sciences, Osteoporosis Research Center, Endocrinology and Metabolism Clinical Sciences Institute</p> <p>Assadi, Majid; Bushehr University of Medical Sciences, The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Farrokhi, Shokrollah; Bushehr University of Medical Sciences, The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Nemati, Reza; Bushehr University of Medical Sciences, The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Amini, Mohammad Reza; Tehran University of Medical Sciences, Endocrinology & Metabolism Research Center, Endocrinology and Metabolism Clinical Sciences Institute,</p> <p>Barekat, Maryam; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Amini, Abdullatif; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Salimipour, Hooman; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Dobaradaran, Sina; Bushehr University of Medical Sciences, The Persian Gulf Marine Biotechnology Research Center, The Persian Gulf Biomedical Sciences Research Institute</p> <p>Moshtaghi, Darab; Bushehr University of Medical Sciences, The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences Research Institute</p>
Primary Subject Heading:	Cardiovascular medicine
Secondary Subject Heading:	Geriatric medicine
Keywords:	Cardiac Epidemiology < CARDIOLOGY, Coronary heart disease < CARDIOLOGY, Hypertension < CARDIOLOGY, Ischaemic heart disease < CARDIOLOGY, General diabetes < DIABETES & ENDOCRINOLOGY, Lipid disorders < DIABETES & ENDOCRINOLOGY

SCHOLARONE™
Manuscripts

1
2
3 **TITLE:** Cohort profile: Bushehr Elderly Health (BEH) Programme, Phase I (Cardiovascular
4 System)
5
6
7
8
9

10
11 Afshin Ostovar¹; Iraj Nabipour^{2*}; Bagher Larijani³; Ramin Heshmat⁴; Hossein Darabi¹; Katayoun
12 Vahdat¹; Maryam Ravanipour¹; Neda Mehrdad⁵; Alireza Raeisi¹; Gholamreza Heidari¹; Gita
13 Shafiee⁴; Mohammadjavad Haeri¹; Mohammadreza Pourbehi²; Farshad Sharifi⁵; Azita Noroozi¹;
14 Rahim Tahmasebi¹; Hamidreza Aghaei Meybodi⁷; Majid Assadi⁶; Shokrollah Farrokhi⁶; Reza
15 Nemati⁶; Mohammad Reza Amini³; Maryam Barekat¹; Abdullatif Amini¹; Houman Salimipour¹;
16 Sina Dobaradaran²; Darab Moshtaghi¹
17
18
19
20
21
22
23
24
25
26
27
28
29

30
31 ¹The Persian Gulf Tropical Medicine Research Center, The Persian Gulf Biomedical Sciences
32 Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran
33
34

35
36 ²The Persian Gulf Biotechnology Research Center, The Persian Gulf Biomedical Sciences
37 Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran
38
39

40
41 ³Endocrinology & Metabolism Research Center, Endocrinology and Metabolism Clinical
42 Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran.
43
44

45
46 ⁴Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences
47 Institute, Tehran University of Medical Sciences, Tehran, Iran.
48
49

50
51 ⁵Elderly Health Research Center, Endocrinology and Metabolism Population Sciences Institute,
52 Tehran University of Medical Sciences, Tehran, Iran
53
54
55
56
57
58
59
60

1
2
3 ⁶The Persian Gulf Nuclear Medicine Research Center, The Persian Gulf Biomedical Sciences
4
5
6 Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran
7

8
9 ⁷Osteoporosis Research Center, Endocrinology and Metabolism Clinical Sciences Institute,
10
11 Tehran University of Medical Sciences, Tehran, Iran
12

13
14
15
16
17
18 **Correspondence:** Iraj Nabipour
19

20
21
22
23
24 **Email:** inabipour@gmail.com
25
26
27

28
29
30 **Postal Address:** The Persian Gulf Tropical Medicine Research Centre, No. 19, Boostan Alley,
31
32 Imam St., Bushehr, Iran
33
34

35 **Tel No:** +98 77 33341828
36
37
38

39
40
41 **Keywords:** Aged; Cardiovascular System; Risk factors; Cohort Studies; Iran
42
43
44

45
46 **Word Count:** 1738
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Purpose: The main objective of the Bushehr Elderly Health Programme, in its first phase, is to investigate the prevalence of cardiovascular risk factors and their association to major adverse cardiovascular events.

Participants: Between March 2013 and October 2014, a total of 3000 men and women aged \geq 60 years old residing in Bushehr, Iran participated in this prospective cohort study (participation rate=90.2%).

Findings to date: Baseline data on risk factors, including demographic and socio-economic status, smoking and medical history were collected through a modified WHO MONICA questionnaire. Vital signs and anthropometric measures, including systolic and diastolic blood pressure, weight, height, waist and hip circumference were also measured. A 12-lead electrocardiography and echocardiography were conducted on all participants, and total of 10cc venous blood was taken and sera was separated and stored at -80 degrees Centigrade for possible future use. Preliminary data analyses showed a noticeably higher prevalence of risk factors among older women compared to men.

Future plans: Risk factor assessments will be repeated every five years, and the participants will be followed during the study to measure the occurrence of major adverse cardiac events. Moreover, the second phase which includes investigation of the bone health and cognition in the elderly will be started in September 2015. Data are available at the Persian Gulf Biomedical Research Institute, Bushehr University of Medical Sciences, Bushehr, Iran for any collaboration.

STRENGTHS AND LIMITATIONS

- This is the first large sample prospective cohort study in Iran focusing on cardiovascular risk factors in the elderly as a growing population.
- The long-term follow-up in this study will allow for the assessment of many relevant outcomes.
- The participation rate in this study was high; however, outcome ascertainment may be incomplete because of incomplete death and MACE registries.

INTRODUCTION

The world's population is ageing rapidly. The proportion of those over the age of 60 will double, from about 11% to 22%, between 2000 and 2050. The absolute number of people aged 60 and over is expected to increase from 605 million to 2 billion over the same period.[1] According to the latest Iranian Census Data, about 8.2% of the population was over 60 years of age in 2011.[2] By the year 2020, the elderly population (over 60 years of age) is estimated to reach 10%.

Coronary heart disease (CHD) is one of the leading causes of disease burden in developing countries. In other words, three-fourths of global deaths due to CHD occurred in low- and middle-income countries.[3] In the year 2003, there were 21 572 Disability Adjusted Lost Years (DALY) due to all diseases and injuries per 100 000 Iranian people of all ages and both sexes. From this total number of DALYs, 58% were due to non-communicable diseases; ischemic heart disease was the fourth cause of DALYs in males and the first cause of DALYs in females.[4]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

There is a need for effective public health action on cardiovascular disease (CVD) prevention, especially in low- and middle-income countries, and for assessments of the cost-effectiveness of feasible interventions.[5] In addition, there is a need to try to balance the fight against the existing burden of infectious diseases with the growing epidemic of chronic diseases such as heart disease and diabetes.[6] The high costs of direct medical care and the indirect costs of cardiovascular disease, according to the American Heart Association, were approaching \$450 billion annually in 2010 and were projected to rise to over \$1 trillion annually by 2030.[7] Prevention of premature deaths due to non-communicable diseases and the reduction of related health care costs should be the main goals of health policy. Improving the detection and treatment of non-communicable diseases and preventing complications and catastrophic events from occurring should be the major goals of clinical medicine.[8] Trends in CVD risk factors and blood glucose will result in substantial CVD burden in developing countries and economies in transition in the near future. Periodic and consistent monitoring of trends and the effects of these risk factors on disease burden is needed for prioritising prevention programmes.[9] In recent decades, cohort studies have played a major role in investigating the incidence and causes of common health outcomes. As the major adverse cardiovascular events are quite common in old age, there are many cohort studies investigating these outcomes. A vast majority of these cohort studies are being conducted in developed countries.[10-20] However, the number of such prospective studies in the developing world is also rising.[21-25] In Iran, there are a few large prospective studies being conducted in various fields. Golestan's cohort study, a prospective study of oesophageal cancer in northern Iran,[26] the Tehran Lipid and Glucose Study (TLGS)[27] and the Shahrood Eye Cohort Study (ShECS)[28] are among these studies. The Amirkola Health and Ageing Project (AHAP) is the only large sample prospective cohort

1
2
3 study in Iran specifically aimed at investigating falling, bone fragility and fractures, cognitive
4
5 impairment and dementia, poor mobility and functional dependence in the elderly.[29]
6
7

8
9 This paper describes the rationale for, design and preliminary results of the Bushehr Elderly
10
11 Health programme (BEH Programme), a population-based prospective cohort study conducted in
12
13 Bushehr, a southern province of Iran. The main objective of this study, in its first phase, is to
14
15 investigate the prevalence of cardiovascular risk factors and their association to major adverse
16
17 cardiovascular events (MACE).
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

COHORT DESCRIPTION

The target population of this prospective cohort study was all men and women aged 60 years and over residing in the city of Bushehr. Based on the information available from the Bushehr District Health Centre, this population was estimated to be around 10 000. Table 1 shows eligibility criteria for participation in the study.

Table 1 List of inclusion and exclusion criteria of participants

Inclusion criteria

- Age more than or equal to 60
- Both sexes
- Residency in Bushehr port since at least 1 year prior to the recruitment
- No plan to leave Bushehr for the following 2 years after the recruitment
- Adequate physical and mental ability to participate in evaluation program
- Signing written informed consent

Exclusion criteria

- No residence in Bushehr
- No willingness to participate in the study

The participants in BEH Programme were selected through a multi-stage, stratified cluster random sampling method. Based on the classifications made by the municipality, as shown in Figure 1, we stratified Bushehr to 75 strata. Numbers were assigned to the blocks (as clusters) of each stratum and then randomly sorted. We invited all eligible older people residing in each

block selected to participate and then moved to the next block, repeating the invitation process, until gaining the sample required for that stratum. Sample sizes for strata were determined proportional to the number of households residing in each stratum. As shown in Figure 2, between March 2013 and October 2014, from among 3297 older people aged 60 years and over who were invited, a total of 3000 participated in this study (participation rate: 91.0%). Non-respondents were found among all strata, and there was no obvious pattern indicating selection bias. Baseline characteristics of participants are shown in Table 2.

Table 2 Baseline socio-demographic characteristics of participants in Bushehr Elderly Health Program

Characteristics [N (%)]	Men [1455 (48.5)]	Women [1545 (51.5)]	
Age Group	≤64	616 (42.3)	674 (43.6)
	65-69	317 (21.8)	378 (24.5)
	70-74	230 (15.8)	200 (12.9)
	75-79	166 (11.4)	181 (11.7)
	≥80	126 (8.7)	112 (7.2)
Marital status	Single	5 (0.3)	20 (1.3)
	Married	1378 (94.7)	884 (57.2)
	Widowed	68 (4.7)	619 (40.1)
	Divorced	4 (0.3)	22 (1.4)
Current occupation	Employed	133 (9.1)	23 (1.5)
	Retired	1195 (82.1)	126 (8.2)
	Unemployed*	127 (8.7)	1396 (90.4)
Education	No education	315 (21.6)	777 (50.3)

Primary school	400 (27.5)	459 (29.7)
Secondary School	276 (19.0)	151 (9.8)
High school	287 (19.7)	125 (8.1)
University	177 (12.2)	33 (2.1)

*homemaker for female

The main follow-up assessments of risk factors will be done every five years for three consecutive periods (a total of 15 years of follow-up), in which all assessments will be repeated with comparable methods and modalities.

In addition, interim assessments will be made to ascertain outcomes and risk factor changes.

All participants will be contacted by a trained nurse once annually, and a checklist will be completed to check if any outcome of interest (major adverse cardiac events) has occurred during the 12 months prior to the assessment. A form has also been distributed to the participants to self-report as soon as possible after the occurrence of any of the targeted outcomes. Focal points in the two main hospitals in Bushehr (Shohadaye-Khalij-e-Fars and Salman-e-Farsi hospitals) are responsible for monthly checking of the hospital information system (HIS) and reporting admissions. If any outcome is reported, a general physician will review the in/outpatient medical records and documents, and detailed information will be entered into the special outcome forms. Our database will be cross-linked with the death registry system database, available via the public health system. The death registry system receives information from hospitals, forensic medicine departments, cemeteries and vital events records offices, and duplicate records are deleted. The International Classification of Diseases-10 (ICD-10), WHO online version, is used to classify causes of death. A person's national identification number is used as a unique identifier to cross-link databases. In the case of a death report but no reliable death certificate, a

1
2
3 verbal autopsy will be performed to determine cause of death. Table 3 presents list of outcomes
4
5
6 ascertained in the present study.
7
8

9 **Table 3 List of Outcomes, BEHP Study**

- 10
11 - Death
12 - Myocardial Infarction
13 - Stroke
14 - Cardiac Interventions
15 - Hospital admission
16
17
18
19

20
21
22
23 **Baseline examination**

24 A modified WHO MONICA questionnaire[30] translated into Persian was used as the core
25
26 questionnaire to gather baseline information on demographic and socio-economic variables and
27
28 risk factors. Table 4 presents the components of the core questionnaire.
29
30
31

32
33 **Table 4 Section and Topics of the Core Questionnaire in the BEHP Study**

34
35

Type of data	Components
Demographic	<ul style="list-style-type: none"> - Personal information (Name, Nickname, Surname) - National Identification Number - Age and Sex - Marital status - Contact Information
Socio-economic	<ul style="list-style-type: none"> - Employment status - Education level - Insurance - Family income - Family assets - Residence status

36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Cardiovascular risk factors	- Smoking status and history
	- Physical activity
	- Menstrual and menopause history
Medical history	- Blood pressure history
	- Diabetes history
	- Lipid Profile history
	- Ischemia and myocardial infarction history
	- Weakness, impaired sensation and stroke history
	- Rose and claudication history
	- Heart failure history
Drug history	- Administered by physician
	- Over the counter drugs
	- Supplements

Phlebotomy and laboratory analyses

Participants were asked to provide a venous blood sample for laboratory tests. A total of 10cc of whole blood were taken by a trained nurse after 8-12 hours of fasting. Table 5 presents laboratory tests done at the baseline and their methods of measurement. Sera were also separated and stored at -80 degrees Centigrade for possible future use.

Clinical assessment

A comprehensive physical examination, including vital signs, weight and height measurements and waist and hip circumference were taken at the baseline. A 12-lead electrocardiography, performed by a trained nurse, and echocardiography, carried out by a cardiologist, were conducted for all participants (see Table 5).

Table 5 Baseline Physical Examinations and Laboratory Tests, BEHP Study

	Item	Method of measurement
Physical Examination	Arterial Blood Pressure	Manually by Standard Mercury sphygmomanometer in sitting position
	Weight in kg	Stadiometer; Heavy outer garments will be removed.
	Height in cm	Stadiometer; Shoes will be removed.
	Waist circumference in cm	At the midway level between the costal margins and the iliac crests.
	Hip circumference in cm	At the level of the greater trochanters.
Laboratory tests	Complete Blood Count (CBC)	Automated Hematology Analyzer
	Fasting Blood Sugar	Enzymatic (glucose oxidase) colorimetric method using a commercial kit (Pars Azmun Inc., Tehran, Iran)
	Lipid Profile (Total cholesterol, LDL, HDL, triglyceride)	Serum total cholesterol and HDL cholesterol will be measured using a Cholesterol oxidase phenol aminoantipyrine and triglycerides using a glycerol-3 phosphate oxidase phenol aminoantipyrine enzymatic method. Serum LDL cholesterol will be calculated using the Friedwald formula; LDL cholesterol will not be calculated when triglycerides concentration is >400 mg/dl.
Procedures	Electrocardiography	12-lead, by a trained nurse
	Echocardiography	Using M-Turbo™ Ultrasound System, Manufactured by SonoSite, Inc. by cardiologist

FINDING TO DATE

Baseline data are being analysed, and preliminary findings are presented in Table 6. The prevalence of cardiovascular risk factors was remarkably higher among older women compared to men. Smoking behaviour was different among older men and women. Older women did not smoke cigarettes as often as men; however, the prevalence of hookah smoking was higher in women. One third of older women were obese, which was two times more than the prevalence of obesity among men. Metabolic syndrome was also about two times more prevalent among older women.

Table 6 Frequency and Age-Adjusted Proportions of Risk Factors in Participants of Bushehr Elderly Health Program

Risk Factor [N(%)]	Men [1455(48.5)]	Women [1545(51.5)]
Smoking		
Hookah		
Non-smoker	1143 (78.8)	951 (61.4)
Current Smoker	110 (7.5)	267 (17.2)
Intermittent Smoker	4 (0.3)	7 (0.5)
Former Smoker	198 (13.4)	320 (20.9)
Cigarette		
Non- Smoker	1023 (70.3)	1522 (98.5)
Current Smoker	198 (13.7)	14 (0.9)
Intermittent Smoker	1 (0.1)	0 (0.0)
Former Smoker	233 (16.0)	9 (0.6)
Hypertension	820 (56.1)	1054 (68.4)

Diabetes Mellitus		390 (27.0)	506 (32.6)
Hypercholesterolemia		886 (61.0)	1034 (66.9)
BMI*	<25	615 (42.6)	414 (27.8)
	25-29.9	612 (42.7)	591 (39.0)
	≥30	211 (14.7)	511 (33.2)
Metabolic Syndrome**		403 (28.5)	826 (53.3)

*46 missing

**Based on the ATP III Criteria

***Percentages are age-adjusted

STRENGTHS AND LIMITATIONS

This is the first large sample prospective cohort study in Iran focusing on cardiovascular risk factors in the elderly as a growing population. The long-term follow-up in this study will allow for the assessment of many relevant outcomes. The participation rate in this study was high; however, outcome ascertainment may be incomplete because of incomplete death and MACE registries.

COLLABORATIONS

This study is carrying out at The Persian Gulf Tropical Medicine Research Centre, which is affiliated with Bushehr University of Medical Sciences, in Bushehr, Iran. We welcome any collaboration, and researchers should contact AO via email, a.ostovar@bpums.ac.ir or IN at inabipour@gmail.com.

FUNDING

1
2
3 The Persian Gulf Biomedical Sciences Research Institute affiliated with Bushehr (Port)
4 University of Medical Sciences (BPUMS) and the Endocrinology and Metabolism Research
5 Institute, affiliated with Tehran University of Medical Sciences (TUMS), have jointly provided
6 funding for this research project.
7
8
9
10
11
12

13 **ETHICS AND DISSEMINATION**

14
15 This study is being conducted in agreement with Declaration of Helsinki and in accordance with
16 Iranian national guidelines for ethics in research. The protocol of the study was approved by the
17 regional research ethics committee of Bushehr University of Medical Sciences on 23 September
18 2013, Reference Number: B-91-14-2. All participants were asked to sign a written informed
19 consent, which was approved by the research ethics committee. The participants are able to
20 withdraw from the study at any time without any explanation. Data collected are stored in a
21 reidentifiable form by national ID code. The results will be presented at national and
22 international meetings and published in a peer-reviewed journal. We aim to translate the key
23 findings to an easily understandable format for local residents and to present them through local
24 media. Relevant findings will also be presented as policy briefs to national and local health
25 policy makers.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **ACKNOWLEDGMENTS**

44
45 We are grateful to the staff of both research centres at BPUMS and TUMS for their commitment
46 to the study's protocol and objectives. We are also indebted to all participants who accepted our
47 invitation and patiently underwent exhausting measurements and examinations.
48
49
50
51
52
53
54

55 **CONTRIBUTORSHIP STATEMENT**

1
2
3 AO drafted the manuscript, participated in study design and conduction, performed data analysis
4 and interpretation. IN conceived the study, helped draft the manuscript, and participated in the
5 study design and conduction, and data analysis and interpretation. BL participated in the study
6 design and conduction and reviewed the manuscript. RH helped draft the manuscript and
7 participated in the study design and conduction and data analysis and interpretation. HD, KV,
8 NM, GH, AR, and MH participated in the study design and reviewed the manuscript. MP, AA,
9 RN, HS, and MB participated in the study design and data collection and reviewed the
10 manuscript. MR, GS, FS, AN, and RT participated in questionnaire development, study design,
11 and staff training and reviewed the manuscript. HAM, MRA, SF, SD, and DM participated in
12 data collection and reviewed the manuscript.

13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 **DATA SHARING STATEMENT**

29
30
31 Huge data have been collected. Access to data is available for interested researchers from
32 corresponding author IN (inabipour@gmail.com) or AO (a.ostovar@bpums.ac.ir).
33
34

35 36 37 **CONFLICT OF INTEREST**

38
39
40 No, there are no competing interests.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Interesting facts about ageing. Secondary Interesting facts about ageing 2012. <http://www.who.int/ageing/about/facts/en/index.html>.
2. Selected Findings of National Population and Housing Census, 2011. In: Cooperations. PRAI, ed. Tehran: Statistical Center of Iran, 2012:27.
3. Gaziano TA, Bitton A, Anand S, et al. Growing epidemic of coronary heart disease in low- and middle-income countries. *Curr Probl Cardiol* 2010;**35**(2):72-115 doi: 10.1016/j.cpcardiol.2009.10.002[published Online First: Epub Date]].
4. Naghavi M, Abolhassani F, Pourmalek F, et al. The burden of disease and injury in Iran 2003. *Popul Health Metr* 2009;**7**:9 doi: 10.1186/1478-7954-7-9[published Online First: Epub Date]].
5. Labarthe DR, Dunbar SB. Global cardiovascular health promotion and disease prevention: 2011 and beyond. *Circulation* 2012;**125**(21):2667-76 doi: 10.1161/circulationaha.111.087726[published Online First: Epub Date]].
6. Shetty P. Grey matter: ageing in developing countries. *The Lancet* 2012;**379**(9823):1285-87
7. Weintraub WS, Daniels SR, Burke LE, et al. Value of primordial and primary prevention for cardiovascular disease: a policy statement from the American Heart Association. *Circulation* 2011;**124**(8):967-90 doi: 10.1161/CIR.0b013e3182285a81[published Online First: Epub Date]].
8. Hunter DJ, Reddy KS. Noncommunicable diseases. *N Engl J Med* 2013;**369**(14):1336-43 doi: 10.1056/NEJMra1109345[published Online First: Epub Date]].
9. Singh GM, Danaei G, Farzadfar F, et al. The Age-Specific Quantitative Effects of Metabolic Risk Factors on Cardiovascular Diseases and Diabetes: A Pooled Analysis. *PLoS one* 2013;**8**(7):e65174
10. Schoeni-Affolter F, Ledergerber B, Rickenbach M, et al. Cohort profile: the Swiss HIV Cohort study. *Int J Epidemiol* 2010;**39**(5):1179-89 doi: 10.1093/ije/dyp321[published Online First: Epub Date]].
11. Hofman A, van Duijn CM, Franco OH, et al. The Rotterdam Study: 2012 objectives and design update. *Eur J Epidemiol* 2011;**26**(8):657-86 doi: 10.1007/s10654-011-9610-5[published Online First: Epub Date]].
12. Mitchell GF, Hwang S-J, Vasan RS, et al. Arterial stiffness and cardiovascular events the Framingham Heart Study. *Circulation* 2010;**121**(4):505-11
13. Christiansen CB, Olesen JB, Gislason G, et al. Cardiovascular and non-cardiovascular hospital admissions associated with atrial fibrillation: a Danish nationwide, retrospective cohort study. *BMJ open* 2013;**3**(1)
14. Welmer AK, Angleman S, Rydwick E, et al. Association of cardiovascular burden with mobility limitation among elderly people: a population-based study. *PLoS One* 2013;**8**(5):e65815 doi: 10.1371/journal.pone.0065815[published Online First: Epub Date]].
15. Vu T-HT, Stamler J, Liu K, et al. Prospective Relationship of Low Cardiovascular Risk Factor Profile at Younger Ages to Ankle-Brachial Index: 39-Year Follow-Up—The Chicago Healthy Aging Study. *Journal of the American Heart Association* 2012;**1**(6)
16. Marmot M, Brunner E. Cohort Profile: the Whitehall II study. *Int J Epidemiol* 2005;**34**(2):251-6 doi: 10.1093/ije/dyh372[published Online First: Epub Date]].

17. Lahelma E, Aittomaki A, Laaksonen M, et al. Cohort profile: the Helsinki Health Study. *Int J Epidemiol* 2013;**42**(3):722-30 doi: 10.1093/ije/dys039[published Online First: Epub Date]].
18. Cumming RG, Handelsman D, Seibel MJ, et al. Cohort Profile: the Concord Health and Ageing in Men Project (CHAMP). *Int J Epidemiol* 2009;**38**(2):374-8 doi: 10.1093/ije/dyn071[published Online First: Epub Date]].
19. Magnus P, Irgens LM, Haug K, et al. Cohort profile: the Norwegian Mother and Child Cohort Study (MoBa). *Int J Epidemiol* 2006;**35**(5):1146-50 doi: 10.1093/ije/dyl170[published Online First: Epub Date]].
20. Osler M, Lund R, Kriegaum M, et al. Cohort profile: the Metropolit 1953 Danish male birth cohort. *Int J Epidemiol* 2006;**35**(3):541-5 doi: 10.1093/ije/dyi300[published Online First: Epub Date]].
21. Dugee O, Oum S, Buckley BS, et al. Assessment of total cardiovascular risk using WHO/ISH risk prediction charts in three low and middle income countries in Asia. *BMC public health* 2013;**13**(1):539
22. Kok VC, Horng J-T, Lin H-L, et al. Gout and subsequent increased risk of cardiovascular mortality in non-diabetics aged 50 and above: a population-based cohort study in Taiwan. *BMC cardiovascular disorders* 2012;**12**(1):108
23. Victora CG, Barros FC. Cohort profile: the 1982 Pelotas (Brazil) birth cohort study. *Int J Epidemiol* 2006;**35**(2):237-42 doi: 10.1093/ije/dyi290[published Online First: Epub Date]].
24. Woodward M, Barzi F, Martiniuk A, et al. Cohort profile: the Asia Pacific Cohort Studies Collaboration. *Int J Epidemiol* 2006;**35**(6):1412-6 doi: 10.1093/ije/dyl222[published Online First: Epub Date]].
25. Jiang C, Thomas GN, Lam TH, et al. Cohort profile: The Guangzhou Biobank Cohort Study, a Guangzhou-Hong Kong-Birmingham collaboration. *Int J Epidemiol* 2006;**35**(4):844-52 doi: 10.1093/ije/dyl131[published Online First: Epub Date]].
26. Pourshams A, Khademi H, Malekshah AF, et al. Cohort Profile: The Golestan Cohort Study- a prospective study of oesophageal cancer in northern Iran. *Int J Epidemiol* 2010;**39**(1):52-9 doi: 10.1093/ije/dyp161[published Online First: Epub Date]].
27. Azizi F, Rahmani M, Emami H, et al. Cardiovascular risk factors in an Iranian urban population: Tehran lipid and glucose study (phase 1). *Soz Praventivmed* 2002;**47**(6):408-26
28. Fotouhi A, Hashemi H, Shariati M, et al. Cohort profile: shahroud eye cohort study. *Int J Epidemiol* 2013;**42**(5):1300-8 doi: 10.1093/ije/dys161[published Online First: Epub Date]].
29. Hosseini SR, Cumming RG, Kheirkhah F, et al. Cohort profile: the Amirkola Health and Ageing Project (AHAP). *International journal of epidemiology* 2013:dyt089
30. Bothig S. WHO MONICA Project: objectives and design. *Int J Epidemiol* 1989;**18**(3 Suppl 1):S29-37

1
2
3 **Figure 1 Map of Bushehr and Distribution of Participants in the Strata**
4
5

6 **Figure 2 Flowchart of enrolments**
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1 Map of Bushehr and Distribution of Participants in the Strata
286x432mm (300 x 300 DPI)




Figure 2 Flowchart of enrolments
144x147mm (300 x 300 DPI)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60