

# BMJ Open Exposure to air pollution and meteorological factors associated with children's primary care visits at night due to asthma attack: case-crossover design for 3-year pooled patients

Shin Yamazaki,<sup>1,2</sup> Masayuki Shima,<sup>3</sup> Yoshiko Yoda,<sup>3</sup> Katsumi Oka,<sup>4</sup> Fumitake Kurosaka,<sup>4</sup> Shigeta Shimizu,<sup>4</sup> Hironobu Takahashi,<sup>4</sup> Yuji Nakatani,<sup>4</sup> Jittoku Nishikawa,<sup>4</sup> Katsuhiko Fujiwara,<sup>4</sup> Yasuyuki Mizumori,<sup>4</sup> Akira Mogami,<sup>4</sup> Taku Yamada,<sup>4</sup> Nobuharu Yamamoto<sup>4</sup>

**To cite:** Yamazaki S, Shima M, Yoda Y, *et al.* Exposure to air pollution and meteorological factors associated with children's primary care visits at night due to asthma attack: case-crossover design for 3-year pooled patients. *BMJ Open* 2015;**5**:e005736. doi:10.1136/bmjopen-2014-005736

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2014-005736>).

Received 20 May 2014  
Revised 26 March 2015  
Accepted 8 April 2015



CrossMark

For numbered affiliations see end of article.

**Correspondence to**  
Dr Masayuki Shima;  
shima-m@hyo-med.ac.jp

## ABSTRACT

**Objective:** We examined the association of outdoor air pollution and meteorological parameters with primary care visits (PCVs) at night due to asthma attack.

**Setting:** A case-crossover study was conducted in a primary care clinic in Himeji City, Japan.

**Participants:** Participants were 1447 children aged 0–14 years who visited the clinic with an asthma attack from April 2010 until March 2013.

**Exposure:** Daily concentrations of air pollutants and meteorological parameters were measured.

**Primary outcome:** PCVs at night due to asthma attack. A conditional logistic regression model was used to estimate ORs of PCVs per unit increment of air pollutants or meteorological parameters (the per-unit increments of particulate matter with an aerodynamic diameter  $\leq 2.5 \mu\text{m}$  (PM<sub>2.5</sub>) and ozone were 10  $\mu\text{g}/\text{m}^3$  and 10 ppb, respectively). Analyses took into consideration the effects of seasonality.

**Results:** We noted an association between PCVs and daily ozone levels on the day before a PCV (OR=1.17; 95% CI 1.01 to 1.35;  $p=0.04$ ), as well as between PCVs and 3-day mean ozone levels before a PCV (OR=1.29; 95% CI 1.00 to 1.46;  $p=0.04$ ), from April until June. We also observed an association between PCVs and daily PM<sub>2.5</sub> levels on the day before a PCV from December until March (OR=1.16; 95% CI 1.01 to 1.33;  $p=0.05$ ). Meteorological parameters, such as hours of sunshine from September until November, atmospheric pressure from April until June, and temperature from April until August, were also found to be associated with PCVs.

**Conclusions:** The findings in the present study supported an association between ozone and PCVs and suggest that certain meteorological items may be associated with PCVs.

## Strengths and limitations of this study

- The findings in this study support an association between ozone and primary care visits in spring.
- Moreover, certain meteorological items such as atmospheric pressure, temperature and hours of daylight may be associated with primary care visits.
- One of the limitations of this study is that our use of a number of statistical test produces led to issues with multiple comparisons.

## INTRODUCTION

Exposure to ambient air pollutants such as particulate matter (PM), ozone and nitrogen dioxide (NO<sub>2</sub>) is associated with a number of adverse health outcomes ranging from increased symptoms of allergic airway disease to increased mortality.<sup>1–6</sup> Children are considered to be particularly susceptible to air pollution and meteorological factors because their lungs are still growing and their immune and pulmonary systems are developing, making them vulnerable to infection with respiratory pathogens. Children are considered to be more sensitive to air pollution than adults,<sup>7</sup> and children with asthma are particularly vulnerable to the adverse health effects of air pollution. Studies in children with asthma have concluded that exposure to high concentrations of ozone or PM significantly enhances the risk of respiratory symptoms, asthma medication use and reduced lung function, characterised by a decline in peak expiratory flow or forced expiratory flow in a short time.<sup>8–14</sup> However, while a

number of studies have been conducted in the field of air pollution epidemiology, the adverse health effects of low-level exposure of air pollutants and seasonal effects of air pollutants on respiratory symptoms remain unclear.

Further, while many studies have examined the association between ambient air pollutants and asthma attack,<sup>1–6</sup> few have assessed that between meteorological parameters and asthma attack. Previous studies have shown that the meteorological risk factors of care visits due to asthma attack were high temperature,<sup>15–17</sup> large changes in temperature,<sup>18</sup> high atmospheric pressure,<sup>18</sup> low relative humidity,<sup>18</sup> large changes in humidity<sup>19</sup> and hours of daylight.<sup>17</sup> However, these data may suffer from publication bias.

We previously collected and analysed daily mean air pollutant data and meteorological parameters as well as records of primary care visits (PCVs) at night due to asthma attack in Himeji City, western Japan from April 2010 to March 2012. The results of that 2-year analysis suggested no stable association between air pollutants and PCVs, although some meteorological parameters—such as temperature in summer and hours of daylight in fall—were found to be associated with PCVs.<sup>17</sup> In addition, at the beginning of 2013, extremely high concentrations of PM with an aerodynamic diameter  $\leq 2.5$   $\mu\text{m}$  (PM<sub>2.5</sub>) were observed around Beijing, China.<sup>21</sup> In Japan as well, the health effects of transboundary air pollution have been a matter of concern. We previously examined the association between PM<sub>2.5</sub> and PCVs at night from January until March 2013 (3-month analysis). This analysis was conducted in the same setting as the previous 2-year analysis,<sup>17</sup> and the results did not support any association between the daily mean concentration of PM<sub>2.5</sub> and PCVs,<sup>22</sup> although we did note an association between ozone and PCVs. However, both of these previous studies suffered from statistical power that was too low to detect the association between air pollutants and PCVs.

We conducted a 3-year (April 2010–March 2013) analysis to clarify any potential association of air pollutants and meteorological parameters with PCVs due to asthma attacks at night.

## METHODS

### Participants

The setting of this study was the same as that of the previous 2-year analysis<sup>17</sup>—namely, Himeji City Emergency Clinic (Himeji, Japan), which was established for the purpose of treating emergency cases between 21:00 and 6:00 on weekdays. Himeji City is located in western Japan, within 100 km of central Osaka, and facing the Seto Inland Sea. The city is 534 km<sup>2</sup> in area, and its population was about 540 000, some 80 000 of whom were aged less than 15 years. Participants were city residents aged <15 years with a history of asthma attack who visited the clinic for asthma attack between 21:00 and

6:00 from April 2010 to March 2013 and who had received a prescription for bronchodilators from their general practitioner. In Himeji City, primary care at night is generally only available at this emergency clinic, and almost all patients who suffer asthma attacks at night visit the clinic. Patients who visited the clinic on national holidays were excluded (see *Statistical methods* section). The medical records of all patients were provided retrospectively, and patient age, sex, diagnosis and date of visit were recorded.

### Air pollutants and meteorological parameters

From April 2010 until March 2013, concentrations of PM<sub>2.5</sub>, PM<sub>10</sub> and optical black carbon (OBC) were measured hourly using an SPM-613D dichotomous  $\beta$  gauge monitor (Kimoto Electric Co. Inc, Osaka, Japan) at the monitoring station, which was located in the central site of the residential districts in the city. In this study, OBC in PM<sub>2.5</sub> was measured. Data on daily concentrations of NO<sub>2</sub> and ozone from during the same period were obtained from the Himeji local government. All participants resided within 10 km of the monitoring station. Data on meteorological parameters, such as daily mean values for atmospheric pressure, relative humidity, temperature, wind speed, as well as total hours of daylight, were obtained from the Japan Meteorological Agency.

### Statistical methods

The study was conducted under a time-stratified case-crossover design, which is used to assess brief changes in risk associated with transient exposures.<sup>23</sup> Case-crossover analyses require exposure data for cases only. They can be regarded as a special type of case-control study in which each case serves as its own control providing inherent control of potential confounding by fixed individual characteristics, such as sex, race, diet and age. ‘Time-stratified’ indicates the method by which the control periods were chosen. Specifically, we stratified time into months to select days for control periods that fell on the same day of the week within the same month in the same year as the date of a PCV (day of the index period). For example, if a PCV at night due to asthma attack occurred on 18 March 2012, the three control days were 4, 11 and 25 March 2012. This approach therefore also controls for long-term trends, seasonality and day of the week.

We excluded patients who visited the clinic on national holidays because of bias in control selection. That is, if patients whose visits occurred on holidays were included as participants, the estimated relative risks were lower than expected because the concentration of air pollutants on holidays (days for the index periods) was usually/systematically lower than that on non-holidays (days for control periods).<sup>25</sup>

We examined associations of daily mean concentrations (day-of and day before PCVs) of each air pollutant before PCVs with the risk of PCVs at night due to asthma attack. We also examined associations of 3-day

mean concentrations of each air pollutant before PCVs with the risk of PCVs at night due to asthma attack. We estimated ORs of PCVs at night due to asthma attack per  $10 \mu\text{g}/\text{m}^3$  increment in  $\text{PM}_{2.5}$  in a single-pollutant model adjusted for 1-day/3-day mean atmospheric pressure (hPa), relative humidity (%), temperature ( $^{\circ}\text{C}$ ), wind speed (m/s) and hours of daylight (h). Similarly, we also estimated ORs of PCVs per  $10 \mu\text{g}/\text{m}^3$  increment in  $\text{PM}_{10}$ , per  $0.1 \mu\text{g}/\text{m}^3$  increment for OBC, and per 10 ppb increment in  $\text{NO}_2$  and ozone. In addition, we simultaneously assessed the associations between PCVs at night due to asthma attack and increment of  $\text{PM}_{2.5}$ ,  $\text{NO}_2$  and ozone using a multipollutant model adjusted for the same variables as the single-pollutant model. When 3-day mean concentrations of air pollutants were explanatory variables, adjusted variables were also included as 3-day mean values. In addition, we considered interactions between ozone and temperature in the single-pollutant model as a sensitivity analysis.

Modified effects of season were examined using a four-level indicator variable for spring (April until June), summer (July and August), fall (September until November), and winter months (December until March). These classifications were the same as those used in our previous study.<sup>17</sup>

Conditional logistic regression was carried out using the PHREG procedures of SAS release V.9.2 (SAS Institute, Inc, Cary, North Carolina, USA). All tests were two-tailed, and  $\alpha$  was set at 0.05. We computed ORs and their 95% CIs.

## RESULTS

Participant characteristics are shown in table 1. Of the 1447 participants evaluated, 1005 were boys, and 115 were aged less than 2 years. The number of cases in spring, summer, fall and winter were 390, 291, 481 and 285, respectively. Daily mean concentrations of air pollutants and other meteorological data by season are shown in table 2. Ozone levels were highest in spring among all seasons (table 2). Figure 1 shows daily concentrations of  $\text{PM}_{2.5}$  and ozone from April 2010 until March 2013. The correlation coefficients among daily concentrations of

air pollutants and daily means of meteorological parameters are shown in table 3.

Associations between air pollutants and PCVs at night are described using the single-pollutant model in table 4. Findings with the multipollutant model are shown in table 5.

## Ozone

While we noted no association between PCVs and concentrations of air pollutants on the same day as a PCV, we did observe an association between PCVs and ozone levels on the day before a PCV in spring and winter, with respective ORs per 10 ppb increment of ozone of 1.16 (95% CI 1.01 to 1.34;  $p=0.04$ ) and 1.29 (95% CI 1.03 to 1.61;  $p=0.03$ ) (table 4). With respect to the association between PCVs and 3-day mean concentrations of air pollutants before PCVs, ozone was associated with PCVs in spring and winter, with respective ORs per 10 ppb increment of ozone of 1.21 (95% CI 1.00 to 1.46;  $p=0.05$ ) and 1.56 (95% CI 1.12 to 2.17;  $p=0.01$ ). We also detected a statistically significant association of daily (day before a PCV) and 3-day mean ozone levels before a PCV with PCVs in spring in the multipollutant model (table 5), with respective ORs per 10 ppb increment of daily (day before a PCV) and 3-day mean ozone levels of 1.17 (95% CI 1.01 to 1.35;  $p=0.04$ ) and 1.21 (95% CI 1.00 to 1.46;  $p=0.04$ ). With respect to annual effects of ozone, we also observed a significant association between daily (day before a PCV) and 3-day mean ozone levels before a PCV and PCVs (tables 4 and 5). In addition, when we considered interactions between ozone and temperature in the single-pollutant model, there was no significant interaction between ozone and temperature ( $p$  values of the interactions on the same day of a PCV, on the day before a PCV, and for 3-days mean were 0.17, 0.21 and 0.11, respectively); the association between ozone and PCV remained significant (data not shown).

## PM and $\text{NO}_2$

$\text{PM}_{10}$  levels on the day before a PCV were also associated with PCVs in summer. OR per  $10 \mu\text{g}/\text{m}^3$  increment of  $\text{PM}_{10}$  was 1.19 (95% CI 1.04 to 1.38;  $p=0.01$ ) (table 4). In addition,  $\text{PM}_{2.5}$  levels on the day before a PCV were

**Table 1** Age and gender in participants


	April until June (spring)	July and August (summer)	September until November (fall)	December until March (winter)	Total
Sex					
Male (n)	275	213	340	177	1005
Female (n)	115	78	141	108	442
Age					
0–1 year (n)	30	16	39	30	115
2–5 years (n)	177	120	250	126	673
6–14 years (n)	183	155	192	129	659
Total (n)	390	291	481	285	1447

**Table 2** Summary statistics of daily concentration of air pollutants and meteorological parameters

	April until June (spring)		July and August (summer)		September until November (fall)		December until March (winter)		Over 1 year	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
PM <sub>2.5</sub> (µg/m <sup>3</sup> )	23.4	12.5	21.0	9.6	20.4	11.1	22.7	11.8	22.0	11.5
PM <sub>10</sub> (µg/m <sup>3</sup> )	40.3	26.5	33.9	11.8	32.2	18.9	31.7	17.4	34.4	20.0
OBC (µg/m <sup>3</sup> )	0.5	0.3	0.4	0.3	0.5	0.3	0.5	0.4	0.5	0.4
NO <sub>2</sub> (ppb)	12.0	4.7	8.6	2.7	10.2	3.9	12.9	5.8	11.3	4.9
Ozone (ppb)	36.0	10.0	22.3	10.6	22.2	8.0	23.7	9.1	26.1	11.0
Atmospheric pressure (hPa)	1007.8	5.8	1005.7	4.9	1011.4	5.9	1014.2	5.5	1010.5	6.4
Relative humidity (%)	69.2	11.5	74.1	7.4	72.8	8.3	67.8	9.4	70.5	9.8
Temperature (°C)	17.9	4.8	27.7	1.9	18.3	6.0	5.5	3.2	15.5	9.0
Wind speed (m/s)	2.7	0.8	2.7	0.8	2.5	0.9	2.6	0.9	2.6	0.9
Hours of daylight (h)	5.6	4.4	6.6	3.9	5.5	3.6	5.2	3.2	5.6	3.7

NO<sub>2</sub>, nitrogen dioxide; OBC, optical black carbon; PM, particulate matter.

**Figure 1** Daily concentrations of particulate matter with an aerodynamic diameter  $\leq 2.5$  µm (PM<sub>2.5</sub>) (A) and ozone (B) from April 2010 until March 2013.





**Table 3** Correlation coefficients among daily concentrations of air pollutants and daily means of meteorological parameters

	PM <sub>2.5</sub>	PM <sub>10</sub>	OBC	NO <sub>2</sub>	Ozone	Atmospheric pressure	Relative humidity	Temperature	Wind speed	Hours of daylight
PM <sub>2.5</sub>	1	0.87	0.62	0.31	0.06	-0.01	0.13	0.02	-0.32	-0.04
PM <sub>10</sub>		1	0.44	0.24	0.13	-0.05	0.03	0.11	-0.20	0.02
OBC			1	0.50	-0.16	0.06	0.27	-0.04	-0.51	-0.09
NO <sub>2</sub>				1	-0.23	0.26	0.01	-0.30	-0.34	-0.00
Ozone					1	-0.12	-0.33	0.05	0.14	0.16
Atmospheric pressure						1	-0.38	-0.55	-0.27	0.25
Temperature							1	0.28	-0.24	-0.61
Relative humidity								1	0.02	0.10
Wind speed									1	0.03
Hours of daylight										1

NO<sub>2</sub>, nitrogen dioxide; OBC, optical black carbon; PM, particulate matter.

associated with PCVs in winter, with an OR per 10 µg/m<sup>3</sup> increment of PM<sub>2.5</sub> of 1.16 (95% CI 1.01 to 1.33; p=0.05) (table 5). In contrast, we noted statistical significance in the inverse association between 3-day mean NO<sub>2</sub> levels and PCVs in winter. Neither any association was noted between OBC and PCVs using the single-pollutant model (table 4), nor any association was noted between OBC and PCVs using the multipollutant model, which was adjusted for PM<sub>2.5</sub>, NO<sub>2</sub>, ozone and meteorological factors (see online supplementary table S1).

### Meteorological parameters

In spring, we found that temperature on the day-of and the day before PCV and 3-day mean temperature before PCVs were associated with PCVs, and further noted that atmospheric pressure on the day before PCVs were associated with PCVs (table 5). In summer, we also observed that temperature on the day-of a PCV was associated with PCVs (table 5). In fall, we found that the day-of and 3-day mean decline in the number of hours of daylight before PCVs as well as a decline in 3-day mean relative humidity were associated with PCVs (table 5). Over the course of a year, we found that increases in daily atmospheric pressure and temperature and decreases in wind speed were associated with PCVs (table 5).

### DISCUSSION

The findings from this study support an association between ozone levels and PCVs in spring, when ozone levels are the highest. We also noted an association between single-day PM<sub>2.5</sub> levels on the day before a PCV and PCVs in winter. Further, our findings also suggest that meteorological parameters such as temperature, hours of sunshine, relative humidity and atmospheric pressure may be associated with PCVs to some extent. Of note, the effect of 3-day mean ozone levels in spring in the single-pollutant model did not differ markedly from that in the multipollutant model. Our findings support the notion that the association between air pollution

and meteorological parameters with PCVs due to asthma attack has seasonality.

In our previous 2-year analysis,<sup>17</sup> we noted statistical significance in the inverse association between PCVs and ozone levels on the day-of (but not before) a PCV in winter. We also noted in that study that the OR per degree increment in daily mean temperature was 1.31 (95% CI 1.09 to 1.56) in summer, and OR per hourly increment in the number of daily hours of sunshine was 0.94 (95% CI 0.90 to 0.99) in fall. The results of the present 3-year analysis supported these previous findings with respect to the association between PCVs and meteorological parameters, such as temperature in summer and daily hours of sunshine in fall. In addition, the results of another previous study conducted when extremely high concentrations of PM<sub>2.5</sub> were observed around Beijing, China (3-month analysis) suggested that ozone on the day before PCVs was positively associated with PCVs in winter.<sup>21</sup> Our findings in the present 3-year analysis supported these previous 3-month results. Further, we also detected the association between PCVs and ozone levels on the day before in spring, when the concentrations of ozone were the highest. We consider the results of the present 3-year analysis to have a smaller random error than those of the 2-year or 3-month analyses, due to the sample size of this study being 1.5 times larger than that of the 2-year analysis and approximately 2.5 times that of the 3-month study.

Our findings regarding a potential association between ozone and physician visits due to asthma attack were consistent with those of previous studies. For example, a recent US EPA<sup>5</sup> analysis of ambient ozone health effects concluded that children with asthma suffer acute adverse health consequences even at ambient ozone levels, and Babin *et al*<sup>26</sup> also observed an association between paediatric emergency room visits for asthma exacerbation and outdoor ozone levels. With respect to the seasonality effects of ozone on asthma attack, Moore *et al*<sup>27</sup> indicated that ozone contributes to an increased risk of hospitalisation for children with asthma in warmer months. In a previous Japanese study,

**Table 4** Associations between air pollutants and PCVs at night due to asthma attack (single-pollutant model)

Unit increment	April until June (spring)		July and August (summer)		September until November (fall)		December until March (winter)		Annual effects					
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI				
Concentration on the same day as a primary care visit														
PM <sub>2.5</sub> 10 µg/m <sup>3</sup>	0.935	0.849	1.031	0.921	1.096	0.985	0.891	1.089	1.060	0.923	1.217	0.984	0.931	1.041
PM <sub>10</sub> 10 µg/m <sup>3</sup>	0.965	0.920	1.012	0.947	0.822	1.091	0.901	1.030	1.018	0.924	1.121	0.975	0.944	1.007
OBC 0.1 µg/m <sup>3</sup>	0.990	0.949	1.032	0.956	0.886	1.031	0.984	1.076	0.998	0.957	1.041	0.999	0.976	1.022
NO <sub>2</sub> 10 ppb	0.913	0.671	1.242	1.128	0.619	2.058	1.184	1.643	0.909	0.670	1.232	0.992	0.841	1.170
Ozone 10 ppb	1.019	0.884	1.176	1.124	0.971	1.302	0.979	1.177	1.021	0.820	1.270	1.057	0.978	1.142
Concentration on the day before a primary care visit														
PM <sub>2.5</sub> 10 µg/m <sup>3</sup>	0.964	0.869	1.070	1.185	0.996	1.409	0.970	1.070	1.130	0.986	1.294	1.021	0.965	1.081
PM <sub>10</sub> 10 µg/m <sup>3</sup>	0.961	0.912	1.013	1.194*	1.036	1.377	0.980	1.048	1.096	0.995	1.207	0.998	0.965	1.032
OBC 0.1 µg/m <sup>3</sup>	1.002	0.959	1.047	1.036	0.957	1.121	0.972	1.018	1.001	0.960	1.045	0.997	0.974	1.021
NO <sub>2</sub> 10 ppb	1.060	0.774	1.451	0.864	0.455	1.639	1.132	1.587	0.746	0.550	1.011	0.929	0.783	1.103
Ozone 10 ppb	1.160*	1.005	1.340	1.077	0.932	1.246	0.943	1.138	1.286*	1.028	1.609	1.114*	1.029	1.206
3-day mean concentration before a primary care visit														
PM <sub>2.5</sub> 10 µg/m <sup>3</sup>	0.959	0.834	1.104	1.119	0.882	1.420	0.928	1.048	1.103	0.921	1.320	0.990	0.921	1.065
PM <sub>10</sub> 10 µg/m <sup>3</sup>	0.951	0.892	1.014	1.180	0.973	1.431	0.938	1.016	1.079	0.946	1.231	0.973	0.934	1.014
OBC 0.1 µg/m <sup>3</sup>	1.004	0.943	1.070	0.977	0.882	1.083	0.959	1.018	0.985	0.930	1.042	0.981	0.951	1.012
NO <sub>2</sub> 10 ppb	1.002	0.667	1.506	0.880	0.363	2.134	1.057	1.628	0.581*	0.381	0.887	0.849	0.678	1.063
Ozone 10 ppb	1.207*	1.000	1.457	1.086	0.895	1.318	0.940	1.216	1.562*	1.124	2.170	1.163*	1.046	1.293

Associations are shown as ORs and their 95% CIs per unit increment of each pollutant. Adjusted for daily/3-day mean atmospheric pressure, relative humidity, temperature, wind speed and hours of daylight.

\*p<0.05.

NO<sub>2</sub>, nitrogen dioxide; OBC, optical black carbon; PCV, primary care visit; PM, particulate matter.

**Table 5** Associations between air pollutants and PCVs at night due to asthma attack (multipollutant model)

	April until June (spring)		July and August (summer)		September until November (fall)		December until March (winter)		Annual effects						
	Unit increment	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI				
Concentration on the same day as a primary care visit															
PM <sub>2.5</sub>	10 µg/m <sup>3</sup>	0.936	0.843	1.039	0.891	0.743	1.067	0.863	1.074	1.081	0.939	1.245	0.977	0.920	1.036
NO <sub>2</sub>	10 ppb	0.990	0.709	1.383	0.980	0.528	1.820	0.861	1.821	0.722	0.414	1.258	1.028	0.858	1.230
Ozone	10 ppb	1.025	0.888	1.184	1.130	0.967	1.321	0.841	1.263	0.864	0.585	1.277	1.065	0.981	1.157
Atmospheric pressure	1 hPa	1.027*	1.002	1.053	0.995	0.959	1.032	0.984	1.041	1.002	0.968	1.038	1.016*	1.002	1.031
Temperature	1°C	1.095*	1.043	1.150	1.229*	1.057	1.430	0.996	1.034	0.997	0.935	1.062	1.030*	1.004	1.056
Relative humidity	10%	0.979	0.823	1.165	1.226	0.816	1.843	0.834	0.678	0.956	0.748	1.221	0.960	0.864	1.066
Wind speed	1 m/s	0.892	0.738	1.077	0.786	0.609	1.014	0.923	0.777	0.863	0.677	1.099	0.904*	0.821	0.996
Hours of daylight	1 h	0.971	0.935	1.009	0.994	0.928	1.066	0.953*	0.916	1.002	0.948	1.059	0.979	0.957	1.000
Concentration on the day before a primary care visit															
PM <sub>2.5</sub>	10 µg/m <sup>3</sup>	0.949	0.850	1.059	1.172	0.982	1.399	0.957	1.064	1.157*	1.006	1.331	1.017	0.958	1.079
NO <sub>2</sub>	10 ppb	1.161	0.832	1.619	0.760	0.378	1.530	1.173	1.722	0.857	0.487	1.509	0.979	0.813	1.179
Ozone	10 ppb	1.168*	1.011	1.349	1.087	0.929	1.272	0.977	1.197	1.216	0.809	1.827	1.115*	1.025	1.212
Atmospheric pressure	1 hPa	1.026*	1.000	1.054	1.021	0.985	1.058	0.998	1.025	1.034	0.999	1.070	1.018*	1.003	1.032
Temperature	1°C	1.102*	1.050	1.157	1.081	0.922	1.268	0.980	1.019	0.971	0.913	1.033	1.017	0.992	1.043
Relative humidity	10%	1.009	0.843	1.207	1.035	0.691	1.551	0.854	0.702	1.139	0.891	1.458	0.987	0.889	1.097
Wind speed	1 m/s	0.965	0.805	1.157	1.000	0.795	1.257	0.925	0.781	1.123	0.897	1.407	0.986	0.899	1.081
Hours of daylight	1 h	0.993	0.956	1.032	0.987	0.924	1.056	0.990	1.032	0.983	0.930	1.040	0.993	0.971	1.016
3-day mean concentration before a primary care visit															
PM <sub>2.5</sub>	10 µg/m <sup>3</sup>	0.952	0.824	1.101	1.128	0.877	1.451	0.916	1.046	1.146	0.953	1.377	0.988	0.916	1.065
NO <sub>2</sub>	10 ppb	1.036	0.680	1.578	0.584	0.200	1.708	1.166	1.902	0.665	0.329	1.345	0.897	0.708	1.138
Ozone	10 ppb	1.209*	1.001	1.460	1.123	0.896	1.407	0.989	1.300	1.249	0.733	2.130	1.158*	1.039	1.290
Atmospheric pressure	1 hPa	1.024	0.993	1.057	1.020	0.974	1.069	0.998	1.033	1.026	0.978	1.076	1.017	1.000	1.035
Temperature	1°C	1.121*	1.061	1.185	1.065	0.882	1.287	0.988	1.030	1.010	0.937	1.090	1.036*	1.007	1.065
Relative humidity	10%	0.953	0.752	1.209	1.167	0.688	1.979	0.743*	0.562	1.155	0.782	1.704	0.955	0.826	1.105
Wind speed	1 m/s	0.936	0.722	1.214	0.903	0.609	1.337	0.804	0.622	1.001	0.665	1.508	0.907	0.789	1.042
Hours of daylight	1 h	0.986	0.928	1.049	1.034	0.932	1.147	0.942*	0.888	0.972	0.889	1.063	0.979	0.947	1.013

A multipollutant model is a model which simultaneously assessed the associations between PCVs at night due to asthma attack and increment of PM<sub>2.5</sub>, NO<sub>2</sub> and ozone.

Associations are shown as ORs and their 95% CIs per unit increment of each parameter.

\**p*<0.05.

NO<sub>2</sub>, nitrogen dioxide; PCV, primary care visit; PM, particulate matter.

Yamazaki *et al*<sup>13</sup> also noted an association between ozone and PCVs due to asthma attack in warmer months. Samoli *et al*<sup>14</sup> also noted an association between ozone levels and risk of asthma admission among older children in summer. This study further added that, even in winter, ozone levels might be associated with PCVs.

While we did note an association between single-day PM<sub>2.5</sub> levels on the day before a PCV and PCVs in winter, it was a weak one, observed only in the multipollutant model. The findings in our previous study,<sup>21</sup> which was restricted to cases from January until March 2013, did not support any association between the daily mean concentration of PM<sub>2.5</sub> and PCVs at night due to asthma attack.

As with ozone, the findings regarding the seasonal effects of PM<sub>2.5</sub> on emergency department visits or hospital admission for asthma have proven controversial. Several studies in elderly participants have shown stronger seasonal effects in winter/cooler seasons than in summer/warmer seasons.<sup>28–29</sup> Similarly, Ostro *et al*<sup>30</sup> reported that the association of PM<sub>2.5</sub> with total respiratory admissions was stronger during cool seasons than warm ones for those aged less than 19 years. In contrast, however, Stieb *et al*<sup>31</sup> reported that the effect was stronger in warm seasons than cool seasons in all ages, and Strickland *et al*<sup>32</sup> also observed associations with several traffic-related primary pollutants during the warm season in a large study analysing over 90 000 emergency department visits for paediatric asthma in relation to ambient air pollutant concentrations. Muddying the waters still further, Yamazaki *et al*<sup>13–17</sup> reported an unstable association between PM<sub>2.5</sub> and PCVs in warmer months among patients aged less than 15 years, and Sheppard *et al*<sup>33</sup> reported that the increase in risk of hospital admission for asthma was greatest in spring and fall among patients aged less than 65 years. Additional studies are needed to clarify the seasonal effects of PM<sub>2.5</sub>/air pollutants on respiratory symptoms.

With respect to an inverse association between 3-day mean concentration of NO<sub>2</sub> and PCVs in winter in a single-pollutant model, we speculate that photochemical radiation decreased the concentration of NO<sub>2</sub> and increased the concentration of ozone. In winter, Pearson's correlation coefficient between NO<sub>2</sub> and ozone was  $-0.69$ . Since ozone is secondarily formed by the action of short-wavelength radiation from the sun on NO<sub>2</sub>, when the concentration of ozone was increased, the concentration of NO<sub>2</sub> subsequently tended to be decreased. Therefore, when the 3-day mean concentration of ozone was positively associated with the risk of PCV in winter, the 3-day mean concentration of NO<sub>2</sub> was inversely associated with the risk of PCV. In the multipollutant model, we noted no significant association between PCVs and NO<sub>2</sub>.

While we did note that some meteorological parameters—such as hours of sunshine in fall, atmospheric pressure in spring, and temperature in spring and

summer—were associated with PCVs, additional studies will still be needed to confirm these findings.

### Limitations

Several limitations to this study warrant mention. First, our study design is limited with respect to control of within-person confounding, which is still possible for multiple correlated transient factors that change over time within a participant. For example, if a patient in this study was exposed to certain air pollution or meteorological factors and concurrently was using an inhaler of corticosteroids or albuterol, this confounding would be uncontrolled and would be a threat to the internal validity of the findings. Second, the selection of participants for this study may have been subject to issues with external validity, as we restricted our population to nighttime patients. Third, the estimated ORs in this study may suffer from non-differential misclassification, causing our results to be biased towards null, as single values for air pollution concentrations or meteorological data were assigned to all individuals living in a certain area. Fourth, the association between concentrations of air pollutants and PCV might not be adequately adjusted for meteorological factors, because associations between meteorological factors and PCV are not clear.

### CONCLUSION

The findings in this study supported an association between ozone levels and PCVs in spring and winter and between PM<sub>2.5</sub> levels and PCVs in winter. Further, certain meteorological parameters—such as atmospheric pressure in spring, temperature in spring or summer, and hours of daylight in fall—may be associated with PCVs due to asthma attack. Informing local residents of air pollution levels and meteorological parameters may help prevent asthmatic events in the area.

### Author affiliations

<sup>1</sup>Department of Environmental Epidemiology, National Institute for Environmental Studies, Tsukuba, Japan

<sup>2</sup>Department of Healthcare Epidemiology, Kyoto University School of Public Health, Kyoto, Japan

<sup>3</sup>Department of Public Health, Hyogo College of Medicine, Nishinomiya, Japan

<sup>4</sup>Himeji Medical Association, Himeji, Japan

**Contributors** SY analysed the data and drafted the manuscript. MS developed the survey and study and supervised all aspects of its implementation. YY, KO, FK, SS, HT, YN, JN, KF, YM, AM, TY and NY organised the field survey. All the authors contributed to the study, interpreted the results and reviewed the final draft.

**Funding** This research was supported by the Environmental Research and Technology Development Fund (C-1005 and 5-1456) of the Ministry of the Environment, Japan.

**Competing interests** None declared.

**Ethics approval** The study protocol was approved by the Ethics Committee of Hyogo College of Medicine.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional data are available.



**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

## REFERENCES

- U.S. EPA. Integrated Science Assessment for Particulate Matter. U.S. Environmental Protection Agency, Research Triangle Park, NC. 2009. [http://www.epa.gov/ncea/pdfs/partmatt/Dec2009/PM\\_ISA\\_full.pdf](http://www.epa.gov/ncea/pdfs/partmatt/Dec2009/PM_ISA_full.pdf) (accessed 2 Dec 2014).
- U.S. EPA. *Provisional assessment of recent studies on health effects of particulate matter exposure*. Washington, DC: U.S. Environmental Protection Agency, 2012. <http://cfpub.epa.gov/ncea/isa/recordisplay.cfm?deid=247132> (accessed 2 Dec 2014).
- WHO. *Air quality guidelines, Global update 2005, particulate matter, ozone, nitrogen dioxide and sulfur dioxide*. Geneva, Switzerland: WHO Press, 2005. [http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0005/78638/E90038.pdf](http://www.euro.who.int/__data/assets/pdf_file/0005/78638/E90038.pdf) (accessed 2 Dec 2014).
- The WHO European Centre for Environment and Health, WHO Regional Office for Europe. *Review of evidence on health aspects of air pollution—REVIHAAP project: technical report*. Copenhagen, Denmark: WHO Regional Office for Europe, 2013. [http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0004/193108/REVIHAAP-Final-technical-report-final-version.pdf?ua=1](http://www.euro.who.int/__data/assets/pdf_file/0004/193108/REVIHAAP-Final-technical-report-final-version.pdf?ua=1) (accessed 2 Dec 2014).
- U.S. EPA. Integrated Science Assessment for Ozone and Related Photochemical Oxidants. US Environmental Protection Agency, Research Triangle Park, NC. 2013. <http://epa.gov/ncea/isa/index.htm> (accessed 2 Dec 2014).
- Atkinson RW, Kang S, Anderson HR, *et al*. Epidemiological time series studies of PM<sub>2.5</sub> and daily mortality and hospital admissions: a systematic review and meta-analysis. *Thorax* 2014;69:660–5.
- Pope CA III, Dockery DW. Acute health effects of PM<sub>10</sub> pollution on symptomatic and asymptomatic children. *Am Rev Respir Dis* 1992;145:1123–8.
- Romieu I, Meneses F, Ruiz S, *et al*. Effects of air pollution on the respiratory health of asthmatic children living in Mexico City. *Am J Respir Crit Care Med* 1996;154:300–7.
- Peters A, Dockery DW, Heinrich J, *et al*. Medication use modifies the health effects of particulate sulfate air pollution in children with asthma. *Environ Health Perspect* 1997;105:430–5.
- Gielen MH, van der Zee SC, van Wijnen JH, *et al*. Acute effects of summer air pollution on respiratory health of asthmatic children. *Am J Respir Crit Care Med* 1997;155:2105–8.
- Delfino RJ, Zeiger RS, Seltzer JM, *et al*. Symptoms in pediatric asthmatics and air pollution: differences in effects by symptom severity, anti-inflammatory medication use and particulate averaging time. *Environ Health Perspect* 1998;106:751–61.
- Mortimer KM, Tager IB, Dockery DW, *et al*. The effect of ozone on inner-city children with asthma: identification of susceptible subgroups. *Am J Respir Crit Care Med* 2000;162:1838–45.
- Yamazaki S, Shima M, Ando M, *et al*. Modifying effect of age on the association between ambient ozone and nighttime primary care visits due to asthma attack. *J Epidemiol* 2009;19:143–51.
- Samoli E, Nastos PT, Paliatatos AG, *et al*. Acute effects of air pollution on pediatric asthma exacerbation: evidence of association and effect modification. *Environ Res* 2011;111:418–24.
- Lin S, Luo M, Walker RJ, *et al*. Extreme high temperatures and hospital admissions for respiratory and cardiovascular diseases. *Epidemiology* 2009;20:738–46.
- Abe T, Tokuda Y, Ohde S, *et al*. The relationship of short-term air pollution and weather to ED visits for asthma in Japan. *Am J Emerg Med* 2009;27:153–9.
- Yamazaki S, Shima M, Yoda Y, *et al*. Association of ambient air pollution and meteorological factors with primary care visits at night due to asthma attack. *Environ Health Perspect* 2013;18:401–6.
- Ehara A, Takasaki H, Takeda Y, *et al*. Are high barometric pressure, low humidity and diurnal change of temperature related to the onset of asthmatic symptoms? *Pediatr Int* 2000;42:272–4.
- Mireku N, Wang Y, Ager J, *et al*. Changes in weather and the effects on pediatric asthma exacerbations. *Ann Allergy Asthma Immunol* 2009;103:220–4.
- Chen CH, Xirasagar S, Lin HC. Seasonality in adult asthma admissions, air pollutant levels, and climate: a population-based study. *J Asthma* 2006;43:287–92.
- Yamazaki S, Shima M, Yoda Y, *et al*. Association between PM<sub>2.5</sub> and primary care visits due to asthma attack in Japan: relation to Beijing's air pollution episode in January 2013. *Environ Health Perspect* 2014;19:172–6.
- Wang Z, Li J, Wang Z, *et al*. Modeling study of regional severe hazes over mid-eastern China in January 2013 and its implications on pollution prevention and control. *Sci China Earth Sci* 2014;57:3–13.
- Maclure M. The case-crossover design: a method for studying transient effects on the risk of acute events. *Am J Epidemiol* 1991;133:144–53.
- Levy D, Lumley T, Sheppard L, *et al*. Referent selection in case-crossover analyses of acute health effects of air pollution. *Epidemiology* 2001;12:186–92.
- Kinki Regional Development Bureau, Ministry of Land, Infrastructure, Transport and Tourism. The FY2005 road traffic census results of the general traffic volume surveys in Kinki, Japan (in Japanese). 2006. [http://www.kkr.mlit.go.jp/road/ir/kisyu\\_pdf/18\\_n\\_x/0630\\_02.pdf](http://www.kkr.mlit.go.jp/road/ir/kisyu_pdf/18_n_x/0630_02.pdf) (accessed 2 Dec 2014).
- Babin SM, Burkorn HS, Holtry RS, *et al*. Pediatric patient asthma-related emergency department visits and admissions in Washington, DC, from 2001–2004, and associations with air quality, socio-economic status and age group. *Environ Health* 2007;6:9.
- Moore K, Neugebauer R, Lurmann F, *et al*. Ambient ozone concentrations cause increased hospitalizations for asthma in children: an 18-year study in Southern California. *Environ Health Perspect* 2008;116:1063–70.
- Bell ML. *Assessment of the Health Impacts of Particulate Matter Characteristics*. Health Effect Institute Research Report 161. Boston, MA, 2012:5–38. <http://pubs.healththeeffects.org/getfile.php?u=685> (accessed 2 Dec 2014).
- Linares C, Diaz J. Short-term effect of concentrations of fine particulate matter on hospital admissions due to cardiovascular and respiratory causes among the over-75 age group in Madrid, Spain. *Public Health* 2010;124:28–36.
- Ostro B, Roth L, Malig B, *et al*. The effects of fine particle components on respiratory hospital admissions in children. *Environ Health Perspect* 2009;117:475–80.
- Stieb DM, Szyszkowicz M, Rowe BH, *et al*. Air pollution and emergency department visits for cardiac and respiratory conditions: a multi-city time-series analysis. *Environ Health* 2009;8:25.
- Strickland MJ, Darrow LA, Klein M, *et al*. Short-term associations between ambient air pollutants and pediatric asthma emergency department visits. *Am J Respir Crit Care Med* 2010;182:307–16.
- Sheppard L, Levy D, Norris G, *et al*. Effects of ambient air pollution on nonelderly asthma hospital admissions in Seattle, Washington, 1987–1994. *Epidemiology* 1999;10:23–30.