

BMJ Open Managing work–family conflict in the medical profession: working conditions and individual resources as related factors

Stefanie Mache,¹ Monika Bernburg,² Karin Vitzthum,² David A Groneberg,³ Burghard F Klapp,⁴ Gerhard Danzer⁴

To cite: Mache S, Bernburg M, Vitzthum K, *et al.* Managing work–family conflict in the medical profession: working conditions and individual resources as related factors. *BMJ Open* 2015;5:e006871. doi:10.1136/bmjopen-2014-006871

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2014-006871>).

Received 9 October 2014
Revised 23 January 2015
Accepted 1 February 2015



CrossMark

For numbered affiliations see end of article.

Correspondence to
Dr Stefanie Mache;
s.mache@uke.de

ABSTRACT

Objectives: This study developed and tested a research model that examined the effects of working conditions and individual resources on work–family conflict (WFC) using data collected from physicians working at German clinics.

Material and methods: This is a cross-sectional study of 727 physicians working in German hospitals. The work environment, WFC and individual resources were measured by the Copenhagen Psychosocial Questionnaire, the WFC Scale, the Brief Resilient Coping Scale and the Questionnaire for Self-efficacy, Optimism and Pessimism. Descriptive, correlation and linear regression analyses were applied.

Results: Clinical doctors working in German hospitals perceived high levels of WFC (mean=76).

Sociodemographic differences were found for age, marital status and presence of children with regard to WFC. No significant gender differences were found. WFCs were positively related to high workloads and quantitative job demands. Job resources (eg, influence at work, social support) and personal resources (eg, resilient coping behaviour and self-efficacy) were negatively associated with physicians' WFCs. Interaction terms suggest that job and personal resources buffer the effects of job demands on WFC.

Conclusions: In this study, WFC was prevalent among German clinicians. Factors of work organisation as well as factors of interpersonal relations at work were identified as significant predictors for WFC. Our results give a strong indication that both individual and organisational factors are related to WFC. Results may play an important role in optimising clinical care. Practical implications for physicians' career planning and recommendations for future research are discussed.

INTRODUCTION

The total number of physicians working in the German clinical sector diminished drastically during the past decade.¹ The introduction of the diagnosis-related-groups

Strengths and limitations of this study

- This is one of the first large studies to analyse and discuss work–family conflicts in the medical profession.
- The main predictors for work family conflicts within the work domain of physicians are illustrated.
- The relatively large sample of physicians, together with few missing data, strengthens the study and makes it possible to generalise the findings.
- The cross-sectional design limits the value in supporting causal effects.
- Data were assessed by self-report measures that limit the objectivity of the study results.

(DRG)-based financing system in 2004 has led to major changes in hospital work organisation and workload for clinical doctors. These days health insurance reimbursements are granted diagnosis-centered, which has led to a number of problems such as early dismissal, underfunding of mandatory treatment with complications etc.² Probably, in consequence of this, an increasing number of physicians decided to work in non-medical fields (such as pharmaceutical industry, etc.) or abroad.³ Studies investigated the motives why physicians decided not to work in the clinical sector,⁴ for example, overwork, insufficient income, etc.^{5 6} Since incompatibility between work and family life is one of the main factors for physicians' dissatisfaction and a central reason for stopping work in the clinical sector, research studies should investigate more thoroughly the topic 'work–family conflict'.

Work–family conflict

Work–family conflict (WFC) has been defined as “a type of inter-role conflict in which the role pressures from work and family are not compatible in some

respect".^{7 8} The definition of WFC implies a bidirectional relation between work and family life in such a way that work can interfere with family life (WIF) and family life can interfere with work demands (ie, family-to-work conflict: FIW).

According to the role theory,⁷ WFC may occur when people are involved in multiple roles (eg, private and work related); these roles tend to drain them and cause stress or inter-role conflict.^{9 10} Role theory argues that inter-role conflicts experienced by individuals will result in an undesirable state, if it becomes difficult to fulfil each role successfully owing to conflicting demands on time and behaviour among roles.^{7 11}

Working conditions in the healthcare sector and WFC

The medical profession is characterised by an intense and high work commitment. Physicians working in clinics have often reported facing a high workload, low autonomy and job control.^{2 12} In addition, physicians often report that the time of residency coincides with the family-founding life stage leading to high levels of WFC.^{3 13} Young physicians are more likely to have young children and consequently experience high family or parenting demands, resulting in high levels of WFCs. This has been found to be significantly associated with physicians' job demands (ie, workload).^{14–16} As the study by Fuss *et al*³ showed, German physicians perceive a high extent of WFC compared to the general population. Moreover, they demonstrated that a high workload, the number of working hours per day, the amount and frequency of overtime work, an inflexible work schedule and rare support from colleagues and supervisors can increase the likelihood of employees experiencing a conflict between their work and family roles.

Although this research topic is of high interest, data examining WFCs within the medical profession are rare in general.¹⁷ We expect that both physicians' work-related demands (eg, high workload) and perceived resources (eg, social support) relate directly to WFC.

The Job Demands–Resources (JD–R) Model¹⁸ and the Demand–Control–Support Model¹⁹ build the rationale for this hypothesis. According to the JD–R Model, effects of job demands (ie, organisational job factors that require sustained physical and/or psychological efforts and are associated with certain physiological and/or psychological costs)²⁰ are related to WFC.²¹ A key proposition of the JD–R model is that interactions between job demands and resources are important by the way that certain job resources can buffer negative effects of job stress.²² Given that WFC is a chronic stressful condition, one could expect that the availability of resources (ie, working part-time, childcare, maternity leave) would help the individual to successfully manage this conflict.^{20 23} Research supports this assumption; a number of cross-sectional studies have shown negative associations of job resources with WFCs. These job resources include job control,^{24 25} social support at work,^{26 27} reward, feedback and supervisory coaching.²¹

The Demand–Control–Support model¹⁹ presumes that working situations have negative psychological or physical consequences in particular when high demands coincide with limited decision latitude and low social support at the workplace.¹⁹ Several studies focused on the question how the model interacts when WFC is considered. They showed that increased job demands are associated with more WFC, while control has a reducing effect.^{28–31}

Gender differences, personal resources and WFC

Recently, the number of female physicians working in the German clinical sector has increased.³² At the same time, expectations within family roles have neither been modified nor reduced.³³ Findings of previous research in various samples concerning gender difference in WFCs are inconsistent. Some studies report higher WFC scores among women compared to men.^{34 35} A study by Adám *et al*³⁶ showed that female physicians reported significantly higher levels of WFC compared to male physicians and more often.

In other studies, WFCs were found to be similar for men and women.^{37 38} Some studies demonstrated that female doctors with children spend more time with their family than working men do^{7 34 39} by trying to combine a medical career with child caring and working part-time.⁴⁰

Presently, there are few studies that have focused on associations between personal resources and WFC as the outcome variable. The construct of personal resources refers to those personal characteristics that promote an individual's sense of control and self-evaluation, and are linked to resilience, positive coping with environmental demands and well-being.⁴¹ Some studies have examined the role of personal characteristics in WFC.^{42–44} According to Hobfoll,⁴⁵ the resource category of personal characteristics enhances general resistance to stress. A study by Bernas and Major⁴⁶ examined the role of self-efficacy and demonstrated a negative correlation to WFC. Bruck and Allen⁴⁷ reported the following results for the relationship between personality variables (Big Five Personality) and WFC: agreeableness and conscientiousness have significant and negative prediction effects on WFC. In contrast, neuroticism was positively associated with the WFC dimensions.

Research questions and hypotheses

The aims of this study are: (1) to analyse the prevalence of WFCs among German hospital physicians, (2) to expose antecedents of WFC within the work domain of physicians and (3) to investigate differences in sociodemographic variables (gender, age) with regard to WFC. Another focus lies in the research question whether personal resources are related to WFC. Expecting personal and work-related resources to moderate relations between work-related demands and WFCs, we also concentrate on interactions between WFCs and different types of personal and work-related resources and demands.

In summary of the outlined theoretical frame and the cited empirical results, we work on the following questions by testing hypotheses:

- ▶ Question I: Do German hospital physicians perceive levels of WFC and, if so, do they differ across various sociodemographic variables?
- ▶ Hypothesis I: Levels of WFC differ significantly among hospital physicians depending on sociodemographic variables (eg, age, gender, medical specialty).
- ▶ Question II: What are the main predictors for WFC within the work domain of physicians?
- ▶ Hypothesis IIa: Perceptions of workplace resources (eg, social support) will be negatively related to WFC.
- ▶ Hypothesis IIb: Perceptions of workplace demands (eg, quantitative demands) will be positively related to WFC.
- ▶ Hypothesis IIc: Perceptions of resources will moderate the relationships between job demands and WFC.
- ▶ Question III: Do personal resources predict WFC in the medical profession?
- ▶ Hypothesis: Personal resources (eg, self-efficacy) will be negatively related to WFC.

MATERIAL AND METHODS

Study design, participants and setting

The study was designed as a cross-sectional questionnaire evaluation. Data collection took place between 2010 and 2013 in 15 different hospitals in the northern and eastern parts of Germany. Hospital departments were Internal Medicine, Neurology, Surgery, Paediatrics, Anaesthesiology and Gynaecology and Obstetrics. On the basis of information of the German Federal Office of Statistics in 2012, the chosen hospitals are comparable to other German hospitals.⁴⁸ The hospitals included in the study had similarities in the following variables: size, number of patients/beds as well as employed medical staff (eg, residents, junior doctors, nurses).

Data collection

In the beginning, we presented our study design to clinic management, physicians' supervisors and physicians during clinical conferences/meetings. Afterwards, the questionnaire was distributed together with an informative and invitational letter to participate. The cover letter explained that participation in this study was voluntary and anonymous.

We did not ask for written consent of the participants as such, but were given the opportunity to review their voluntary participation after having read the informative letter (an informed consent letter). We asked physicians (N=1154) to fill out the questionnaire within 3 weeks and to return it via locked boxes at the hospital wards. Reminders were sent by email after 3 weeks to increase the response rate. At the end, we received 727 questionnaires from the N=1154 contacted physicians (response rate of 63%).

Variables

WFC was the dependent outcome variable. Organisational factors were included as independent variables:

- ▶ Quantitative and emotional job demands;
- ▶ Degree of freedom at work, influence at work, social relations, social support, sense of community, quality of leadership.

In addition, specific psychological trait variables were chosen in the set of questionnaires (eg, self-efficacy, etc.) presumed by us as predictors for WFC:

- ▶ Resilience, self-efficacy, optimism.

We expected physicians' sociodemographic characteristics to have an effect on their ratings; therefore, we analysed differences in age, gender, marital status, presence of children, medical specialty and work experience in years at the time of the study.

Instruments

Work-family conflict

We analysed the WFCs using the German version of the original instrument by Netemeyer.⁴⁹ This instrument consists of five items including questions on the influence of work on personal or family life, to be answered with a five-point Likert scale (from 'strongly agree' to 'strongly disagree'):

1. The demands of my work interfere with my home, personal and family life.
2. The amount of time my work takes up makes it difficult to fulfil family responsibilities or personal obligations.
3. Things I want to do at home do not get done because of the demands my job puts on me.
4. My job produces strain that makes it difficult to fulfil family duties or personal duties.
5. Due to work-related duties, I have to make changes to my plans for family or personal activities.

Items described above were transformed to a scale ranging from 0 (minimum value, eg, 'strongly disagree') to 100 points (maximum value, eg, 'strongly agree').

The internal consistency of the scale with our sample was good ($\alpha=0.81$).

Organisational resources

The German version of the Copenhagen Psychosocial Questionnaire (COPSOQ) was used to evaluate job-related and psychosocial factors at work.^{50 51}

The questionnaire includes 12 scales; we analysed the following eight scales of the COPSOQ: emotional and quantitative job demands (workload, working under pressure), job resources (quality of leadership, feedback, opportunities for development, social support, sense of community, social relationships, degrees of freedom at work and influence at work). Items are scored on a five-point Likert scale and transformed to a scale ranging from 0 (minimum value, eg, 'do not agree at all') to 100 points (maximum value, eg, 'fully agree'). Previous investigations proved quality criteria of the COPSOQ.⁵¹ We also verified them: Cronbach's α coefficients ranged

between $\alpha=0.71$ to $\alpha=0.85$, while intercorrelations ranged between $r=0.30$ and $r=0.68$.

Personal resources

The questionnaire 'Self-Efficacy, Optimism and Pessimism' (SWOP-K9) was used to analyse physicians' personal resources.⁵² This instrument assesses individuals' perception of self-efficacy, optimism and pessimism on three scales (with nine items in total). The good test quality criteria of the SWOP-K9 questionnaire has been discussed in a previous publication.⁵²

Resilience was evaluated by using the German version of the 'Brief Resilient Coping Scale' (BRCS),⁵³ which consists of four self-assessing items for resilient coping behaviour in difficult or unpleasant situations in the past.

Statistics

Data analysis included descriptive analyses, correlation and reliability analyses. Analyses of variance, and linear bivariate and multiple regression analysis were also performed. Data collected on age, gender, number of years in training and marital status were used as covariates and predictive factors. *p* Values of less than 0.05 were considered significant, and all *p* values given were two tailed.⁵⁴ Data were calculated using the SPSS software package for social sciences; V.21.0.

RESULTS

Descriptive statistics

Physicians' sociodemographic characteristics are given in table 1. In all, 54% of the respondents were female physicians; 46% were male, 62% were married or lived in a partnership and 41% had children; the mean age was 35 years (SD=7.9 years), and work experience was rated with a mean of 8 years (SD=7.2 years).

WFC and working conditions

On the WFC scale, the study sample of 727 hospital physicians reached a mean of 76.1 (SD=20.4) (range 0–100) (table 2). With regard to the single items of the WFC scale, more than 40% ($n=291$) of the respondents agreed with the statements "Things I want to do at home do not get done because of the demands my job puts on me" (45%) and "Due to work-related duties, I have to make changes to my plans for family or private activities" (41%). 94% ($n=683$) agreed with the statement "My job produces strain that makes it difficult to fulfill family duties or personal duties". The other two items were also important for the majority of doctors (>50%).

Descriptive values of physicians' working conditions are listed in table 3. Quantitative job demands were rated with high levels ($M=74.56$, $SD=12.17$), whereas job resources were rated with significantly lower scores (range $M=44.13$ – 62.66).

Sociodemographic differences

In addition, we examined differences in sociodemographic variables. Men and women differed significantly in age and presence of children ($p<0.05$). Female physicians were younger than their male colleagues. Men reported more often having a child or children than their female colleagues ($p<0.05$). For none of the other sociodemographic predictors, a gender difference could be found. Additionally, we analysed the predictors of WFC for gender differences via multivariate analysis of variance. Sense of community (sc) and social support (ss) were the predicting scales which were significantly different for female physicians (sc: mean=79.26, ss: mean=76.61) and male physicians (sc: mean=68.34; ss: mean=70.15; $p<0.05$). The analysis showed a non-significant result for gender differences in WFC ($F(1,726)=0.411$; $p=0.52$) (table 2). We also found that the factor age stayed significant ($F(3,724)=3.17$; $p=0.024$): younger physicians reported lower levels of WFC. In addition, for WFC we identified significant differences for presence of children ($F(1,726)=4.619$; $p=0.032$): physicians with children reported higher levels of WFC.

Table 1 Demographic characteristics of the participants ($n=727$)

	N	Per cent
Gender		
Female	393	54
Male	334	46
Age		
<25	0	0
26–30	211	29
31–35	182	25
36–40	138	19
41–45	87	12
46–50	73	10
>50	36	5
Professional status		
Intern/resident	575	79
Attending physician	116	16
Senior physician	36	5
Work experience (years)		
Less than 1	73	10
1–2	102	14
3–5	247	34
More than 5	305	42
Area of specialisation		
Internal medicine	181	25
Surgery	124	17
Paediatrics	102	14
Anaesthesiology	95	13
Neurology	110	15
Gynaecology	115	16
Marital status		
Single	276	38
Married/partnership	451	62
Children		
Child/children	298	41
Without children	429	59

Table 2 Sociodemographic differences in WFC

WFC	M	SD
Gender		
Female	74.1	19.5
Male	73.5	19.4
Age		
<25	61.2	20.1
26–30	68.9	19.6
31–35	75.8	20.4
36–40	76.1	19.7
41–45	75.5	18.3
46–50	74.3	18.5
>50	72.4	16.1
Work experience		
Less than 1 year	61.6	19.8
1–2 years	68.2	20.1
3–5 years	74.5	20.4
More than 5 years	75.3	19.7
Professional status/presence of children		
Intern, no children	58.9	18.9
Intern, children	68.1	19.3
Attending, no children	70.9	20.5
Attending, children	75.4	21.8
Senior, no children	74.9	20.1
Senior, children	72.1	19.5
Medical specialty		
Internal medicine	75.4	21.3
Surgery	76.9	20.7
Paediatrics	73.5	19.6
Anaesthesiology	75.8	21.1
Neurology	72.5	19.2
Gynaecology	76.7	22.6
Sum score	76.1	20.4

WFC, work–family conflict.

We found no significant differences in WFC depending on medical specialty ($p>0.05$). Mean values of WFC per medical specialty are illustrated in [table 2](#).

Table 3 Descriptive values of independent and dependent variables

Dimensions	M	SD
Working conditions		
Quantitative demands	74.56	12.17
Emotional demands	67.54	12.23
Possibilities for development	43.58	13.51
Influence at work	44.13	11.37
Degree of freedom at work	47.29	12.59
Social support	61.16	13.14
Social relationships	59.52	12.62
Sense of community	62.64	11.45
Quality of leadership	50.73	15.27
Feedback	49.65	12.49
Psychological variables		
Self-efficacy	3.23	0.65
Optimism	3.46	0.69
Resilience	3.82	0.75

Table 4 Correlation coefficients between work family conflicts and psychological, organisational resources

Dimensions	WFC
Organisational demands	
Quantitative demands	0.32**
Emotional demands	0.25*
Organisational resources	
Possibilities for development	–0.31**
Influence at work	–0.36**
Degree of freedom at work	–0.33**
Sense of community	–0.35**
Social support	–0.38**
Social relationships	–0.34**
Quality of leadership	–0.29*
Feedback	–0.21*
Psychological variables	
Self-efficacy	–0.32**
Optimism	–0.27*
Resilience	–0.35**

* $p<0.05$; ** $p<0.01$.

WFC, work–family conflict.

Associations with WFC

Bivariate analyses revealed several significant negative correlations between WFC and psychosocial (trait) dimensions “resilient coping, self-efficacy and optimism” (see [table 4](#)). Similar correlations were found between WFC and influence at work, sense of community, degree of freedom at work, social support and quality of leadership (see [table 4](#)).

Finally, we performed the regression analysis in which WFC was set as the dependent variable. At first, the variables age, work experience, cohabitation with a partner and medical specialty were statistically controlled ([table 5](#)). These control variables accounted for 5% of the variance in WFCs (model 1). The variables age and years of experience were found to predict WFC (age: $\beta=-0.12$, $p=0.03$; years of experience: $\beta=-0.11$, $p=0.04$). In the second step, the included personal resources accounted for an additional 10% of the variance (R^2 increased to 0.15) (model 2). Personal resources like resilient coping were found to predict WFC ($\beta=-0.18$, $p=0.01$), similar to self-efficacy ($\beta=-0.12$; $p=0.01$) and optimism ($\beta=-0.10$, $p=0.03$), while pessimism did not.

The third step included ‘quantitative demands’ and ‘emotional demands’, both positive predictors for WFC. Quantitative job demands revealed a significant β weight ($\beta=0.26$, $p=0.01$). Emotional demands showed a non-significant β weight of $\beta=-0.04$. In the fourth step, physicians’ job resources were included in the regression analysis (see [table 5](#)). The regression model explained 41% of the variance in the final model. The included organisational resources accounted for an additional 17% of the variance. Job resources revealed significant β weights for the following variables: influence at work ($\beta=-0.12$, $p=0.03$), possibilities for development ($\beta=-0.23$, $p=0.01$), degree of freedom at work ($\beta=-0.16$, $p=0.01$), sense of community ($\beta=-0.18$, $p=0.01$), feedback

Table 5 Multiple hierarchical regressions (ratios of variance and standardised β weights)

Explanatory variables	WFC		
	β	R ²	R ² change
Step 1: Sociodemographic variables		0.05	0.05
Age	-0.12*		
Gender	0.07		
Marital status	0.05		
Children	0.06		
Years of experience	-0.11*		
Step 2: Personal resources		0.15	0.10
Resilience	-0.18**		
Optimism	-0.10*		
Pessimism	0.07		
Self-efficacy	-0.12*		
Step 3: Job demands		0.24	0.09
Quantitative demands	0.26**		
Emotional demands	-0.04		
Step 4: Job resources		0.41	0.17
Influence at work	-0.12*		
Possibilities for development	-0.23**		
Degree of freedom at work	-0.16*		
Sense of community	-0.18**		
Feedback	-0.10*		
Quality of leadership	-0.16**		
Social support	-0.14*		
Social relationships	-0.13*		
Total R ²	0.41		

β , standardised β -coefficients from the *final step* of the model. R², explanation rate. Δ R², change in the explanation rate *at each step*.

*p<0.05; **p<0.01.

WFC, work-family conflict.

($\beta=-0.10$, $p=0.04$), quality of leadership ($\beta=-0.16$, $p=0.01$), social support ($\beta=-0.14$, $p=0.02$) and social relationships ($\beta=-0.13$, $p=0.04$). In addition, we used hierarchical multiple regression to conduct the moderation effect analysis. All variables entered the regression equation following the first regression analysis in table 6. In the first step, age and years of experience have been entered explaining 4% of the variance in WFC ($R^2=0.04$, $p<0.05$). In the second step, personal resources, job demands (quantitative and emotional demands) and job resources (degree of freedom at work, influence at work, social relations, social support, sense of community, quality of leadership) were entered together, explaining an additional 35% of the variance in WFC ($R^2=0.35$, $p<0.01$). In the third step, the interaction terms were significant (range between $\beta=0.20$ and 0.23 , $p<0.01$), explaining an additional variance of 6% (R^2 final=0.45, $p<0.01$). As table 6 shows, there was a significant increase in R^2 as an indicator of the moderation effect.

DISCUSSION

By sampling German hospital physicians ($n=727$) working in different hospitals and medical disciplines, we conducted a cross-sectional study focusing on physicians' WFC, working conditions and personal resources. The present study aimed to discover person related and organisational circumstances in relation to WFC. In addition, we focused on gender differences in these issues.

In comparison to the study of Fuss *et al.*,³ we found similar levels of WFC among German hospital physicians. In accordance with other studies comparing male and female physicians,⁵⁵ this study did not find a gender difference for WFC. In contrast, a study by Dumelow^{55a} found higher WFC levels for female physicians.⁵⁶ Adam⁵⁷ also reported that female physicians reported a significantly higher mean level and prevalence of WFC compared to men. Warde *et al.*⁵⁸ demonstrated that more female than male physicians and more younger than older female physicians experienced levels of role conflict.

On the basis of gender role theory, some researchers confirm that women are more likely to report higher levels of WFC. Nevertheless, there are several studies that revealed the contrary.^{35 59} A possible explanation for our findings might be the fact that men in our sample reported more often than women that they lived with children in their household.

Several predictive factors for WFC were identified in our study. In line with other research findings,⁶⁰ our study shows that quantitative job demands (defined as high workload and working under pressure) are a significant predictor for WFC. Previous research also showed that job demands such as high workloads and working under pressure are related to.^{61 62} In addition, this result supports past findings by researchers such as Boles *et al.*^{63 64} and Noor⁶⁵ who also found significant positive

Table 6 Moderated multiple regression analyses testing the moderating effect of personal resources and job resources in the relationship between work–family conflict and job demands

	Step 1	Step 2	Step3
Criterion variable: work–family conflict			
Step 1: Control variables			
Age	−0.15*	−0.13*	−0.12*
Years of experience	−0.13*	−0.12*	−0.11*
Step 2: Main effects			
Resilience		−0.17**	−0.18**
Self-efficacy		−0.12**	−0.13**
Optimism		−0.10*	−0.10*
Job demands		0.25**	0.21**
Job resources		−0.18**	−0.22**
Step 3: Interaction			
Job demands×resilience			0.20**
Job demands×self-efficacy			0.21**
Job demands×optimism			0.23**
Job demands×job resources			0.21**
R ²	0.04	0.39	0.45
ΔR ²		0.35	0.06

Standardised regression coefficients are provided for each of the three steps; *p<0.05; **p<0.01.

relationships between WFC and working demands. In a meta-analysis of Allen *et al*,²³ all studies reviewed suggested a positive relationship between WFC and high job demands as well as for job-strain.

In line, our study also demonstrated relatively high scores for quantitative job demands (workload, working under pressure) compared with data of other studies performed in different work settings in Germany.^{66 67} With regard to the medical setting, this finding is not surprising since intensification of workload in the course of restructuring of the German health system has been identified as one of the most direct effects on German doctors.⁶⁸ Previous studies analysing job demands in the medical setting with the same questionnaire illustrated comparable scores of job demands (eg, quantitative job demands).^{3 69} Study results describe that one of the sources of WFC lies in work schedule irregularities.⁶² Previous studies also identified the frequent necessity of delaying planned holidays as a predictor for WFC.⁷⁰

Similar to the results of studies on other professionals, ‘influence at work’, ‘social support’ and ‘sense of community’ were identified as protective factors against WFC: the high scores of these job resources predict low values of WFC and low scores predict WFC. These results indicate that a good working atmosphere and support among colleagues and teamwork should be encouraged in hospitals, and that satisfactory working conditions will be facilitated by supportive leadership behaviour.⁷¹

In addition, personal resources seem to help physicians balance WFC, that is, doctors who show high levels of self-efficacy, resilient coping behaviour and optimism may be more capable of preventing WFC. These physicians seem more likely to be able to recognise what role

work and family play for them, and have a better insight into strategies of how job and family demands can be managed. In the case of managing conflicts between private and work responsibilities, self-efficacy and resilient coping behaviour can provide a perspective on what might ultimately help to reduce negative outcomes (such as lower life and/or job satisfaction).^{72 73}

In line with previous findings,³ we found younger (ie, less experienced) physicians scoring higher on the WFC scale than more mature ones. Advanced age could be a protective factor for WFC due to better coping strategies with job stress based on longer job experience.⁷⁴ Moreover, older physicians are not in the family-founding phase; thus, fewer conflicts in work-related and family-related fields may exist.

Limitations

Some limitations of the study shall be addressed: its cross-sectional design limits the value in supporting causal effects and generalisability of the findings. Since data on WFC, working conditions and personal resources were assessed by self-report measures and no observers’ data are available to cross-validate the data of participants, we can only claim limited objectivity of our results. Furthermore, our study sample due to logistic and pragmatic reasons includes only physicians working in clinic departments of the northern and eastern parts of Germany, which may also limit the generalisability of the study findings.

Finally, additional variables should be included such as structure and composition of families as well as measures of conflict orientation.

CONCLUSIONS

The findings of this study provide new information on relations between German physicians’ WFCs, working conditions and personal resources.

The study results can be summarised as follows:

1. Physicians working at German hospitals perceive levels of WFCs.
2. Physicians’ WFCs are *negatively* associated with:
 - A. Job resources, particularly influence at work, sense of community and social support;
 - B. Personal resources (eg, self-efficacy and resilience).
3. Physicians’ WFCs are *positively* associated with job demands such as workload.
4. Personal resources and job resources moderate the relationship between job demands and WFCs.
5. Significant relations between WFCs and sociodemographic variables could be identified (eg, age).

Our findings, in turn, have practical implications:

Hospital management/administration should provide more resources just enough to enable health professionals to balance work and family demands by offering support, influence at work and opportunities for personal development. In addition, to address the needs especially of younger physicians, the start of their

working life might be improved by, for example, structured vocational training,²¹ mentoring programmes and supervision.⁷⁵ The compatibility between career and family should be improved (eg, by offering part-time jobs). Services such as in-house or supported childcare are suggested by different authors as another opportunity of support.^{76–77} By improving these working conditions, not only would WFC be reduced but also the overall job satisfaction and job performance may be increased.

Author affiliations

¹Institute for Occupational Medicine and Maritime Medicine (ZfAM), University Medical Center Hamburg-Eppendorf, Hamburg, Germany

²Institute of Occupational Medicine, Charité—Universitätsmedizin Berlin, Free University and Humboldt University, Berlin, Germany

³Institute of Occupational Medicine, Social Medicine and Environmental Medicine, Goethe-University, Frankfurt am Main, Germany

⁴Charité Center for Internal Medicine and Dermatology, Division of General Internal and Psychosomatic Medicine, Charité—Universitätsmedizin Berlin, Berlin, Germany

Contributors SM designed the study and wrote the manuscript. SM and MB performed the investigation and analysed the data. SM, MB, KV, DAG, BFK, GD interpreted the data and contributed substantially to the writing and revising of the manuscript.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Kopetsch T. Ärzttestatistik. Moderater Zugang, Überalterung setzt sich fort. *Dtsch Arztebl* 2004;101:A-1396–98.
- Janus K, Amelung VE, Gaitanides M, et al. German physicians "on strike"—shedding light on the roots of physician dissatisfaction. *Health Policy* 2007;82:357–65.
- Fuss I, Nubling M, Hasselhorn HM, et al. Working conditions and Work-Family Conflict in German hospital physicians: psychosocial and organisational predictors and consequences. *BMC Public Health* 2008;8:353.
- Bohrer T, Koller M, Schlitt HJ, et al. Workload and quality of life of surgeons. Results and implications of a large-scale survey by the German Society of Surgery. *Langenbecks Arch Surg* 2011;396:669–76.
- Bornschein S, Erbas B, Borelli S, et al. [Working hours and job satisfaction among physicians in hospitals and general practice in Munich. Results of an anonymous questionnaire]. *Gesundheitswesen* 2006;68:535–44.
- Jurkat H, Reimer C, Schröder K. Erwartungen und Einstellungen von Medizinstudentinnen und -studenten zu den Belastungen und Folgen ihrer späteren ärztlichen Tätigkeit. *Psychother Psych Med* 2000;50:215–21.
- Greenhaus J, Beutell N. Sources of conflict between work and family roles. *Acad Manage Rev* 1985;10:76–88.
- Frone M. Work-family balance. In: Quick J, Tetrick LE, eds. *Handbook of occupational health psychology*. Washington, DC: American Psychological Association, 2003:143–62.
- Grandey A, Cropanzano R. The conservation of resources model applied to work-family conflict and strain. *J Vocational Behav* 1999;54:350–70.
- Marks S. Multiple roles and role strain: some notes and human energy, time, and commitment. *Am Sociolog Rev* 1977;42:921–36.

- Kahn RL, Wolfe DM, Quinn RP, et al. *Organizational stress: studies in role conflict and ambiguity*. Oxford, UK: John Wiley, 1964.
- Mache S, Vitzthum K, Nienhaus A, et al. Physicians' working conditions and job satisfaction: does hospital ownership in Germany make a difference? *BMC Health Serv Res* 2009;9:148.
- Estryn-Behar M, Fry C, Guetarni K, et al. Work week duration, work-family balance and difficulties encountered by female and male physicians: results from the French SESMAT study. *Work* 2011;40 (Suppl 1):S83–100.
- Veldhoven M, Beijer S. Workload, work-to-family conflict, and health: gender differences and the influence of private life context. *J Soc Issue* 2012;68:665–83.
- Ilies R, Schwind KM, Wagner DT, et al. When can employees have a family life? The effects of daily workload and affect on work-family conflict and social behaviors at home. *J Appl Psychol* 2007;92:1368–79.
- Ohta H, Wada K, Kawashima M, et al. Work-family conflict and prolonged fatigue among Japanese married male physicians. *Int Arch Occup Environ Health* 2011;84:937–42.
- Tayfur O, Arslan M. The role of lack of reciprocity, supervisory support, workload and work-family conflict on exhaustion: evidence from physicians. *Psychol Health Med* 2013;18:564–75.
- Bakker AB, Demerouti E. The job demands-resources model: state of the art. *J Manag Psychol* 2007;22:309–28.
- Johnson JV, Hall EM. Job strain, work place social support and cardio-vascular disease: a cross-sectional study of a random sample of the Swedish population. *Am J Public Health* 1988;78:1336–42.
- Demerouti E, Bakker AB, Nachreiner F, et al. The Job Demands-Resources model of burnout. *J Appl Psychol* 2001;86:499–512.
- Schaufeli W, Bakker AB. Job demands, job resources, and their relationship with burnout and engagement. *J Organ Behav* 2004;25:293–315.
- Karasek RA. Job demands, job decision latitude and mental strain: implications for job redesign. *Adm Sci Q* 1979;24:285–308.
- Allen TD, Herst DEL, Bruck CS, et al. Consequences associated with work-to-family conflict: a review and agenda for future research. *J Occup Health Psychol* 2000;5:278–308.
- Demerouti E, Shimazu A, Bakker AB, et al. Work-self balance: a longitudinal study on the effects of job demands and resources on personal functioning in Japanese working parents. *Work and Stress* 2013;27:223–43.
- Demerouti E, Peeters MC, van der Heijden BI. Work-family interface from a life and career stage perspective: the role of demands and resources. *Int J Psychol* 2012;47:241–58.
- Ford MT, Heinen BA, Langkamer KL. Work and family satisfaction and conflict: a metaanalysis of cross-domain relations. *J Appl Psychol* 2007;92:57–80.
- Hammer LB, Kossek EE, Anger WK, et al. Clarifying work-family intervention processes: the roles of work-family conflict and family-supportive supervisor behaviors. *J Appl Psychol* 2011;96:134–50.
- Thomas LT, Ganster DC. Impact of family-supportive work variables on work-family conflict and strain: a control perspective. *J Appl Psychol* 1995;80:6–15.
- Duxbury L, Higgins C, Lee C. Work-family conflict: a comparison by gender, family type, and perceived control. *J Fam Issue* 1994;15:449–66.
- Wallace JE. Job Stress, depression and work-to-family conflict: a test of the strain and buffer hypotheses. *Relations Industrielles/Industrial Relations* 2005;60:510–15.
- Grönlund A. More control, less conflict? Job demand-control, gender and work-family conflict. *Scand Work Organ* 2007;14:476–97.
- Bundesärztekammer: Ergebnisse der Ärzttestatistik zum 31. Dezember 2011. <http://www.bundesaerztekammer.de/page.asp?his=0.3.10275.10280>
- Ergeneli A, Ilsev A, Karapinar PB. Work-family conflict and job satisfaction relationship: the roles of gender and interpretive habits. *Scand Work Organ* 2010;17:679–95.
- Hammer L, Allen E, Grigsby T. Work-family conflict in dual-earner couples: within-individual and crossover effects of work and family. *J Vocational Behav* 1997;50:185–203.
- Hill E. Work-family facilitation and conflict, working fathers and mothers, work-family stressors and support. *J Fam Issue* 2005;26:793–819.
- Adám S, Györfy Z, László K. High prevalence of job dissatisfaction among female physicians: work-family conflict as a potential stressor. *Orv Hetil* 2009;150:1451–6.
- Grandey AA, Cordeiro BL, Crouter AC. A longitudinal and multisource test of the work-family conflict and job satisfaction relationship. *J Occup Organ Psychol* 2005;78:1–20.
- Milkie M, Peltola P. Playing all the roles: gender and the work-family balancing act. *J Marriage Fam* 1999;61:476–90.

39. Avison WR, Ali J, Walters D. Family structure, stress, and psychological distress: a demonstration of the impact of differential exposure. *J Health Soc Behav* 2007;48:301–17.
40. Gibson H. Are part time doctors better doctors? *BMJ* 1997;315:7113.
41. Hobfoll SE, Johnson RJ, Ennis N, et al. Resource loss, resource gain, and emotional outcomes among inner city women. *J Pers Soc Psychol* 2003;84:632–43.
42. Boyar SL, Mosley DC. The relationship between core self evaluations and work and family satisfaction: the mediating role of work–family conflict and facilitation. *J Vocat Behav* 2007;71:265–81.
43. Carlson DS. Personality and role variables as predictors of three forms of work–family conflict. *J Vocat Behav* 1999;55:236–53.
44. Blanch A, Aluja A. Work, family and personality: a study of work–family conflict. *Pers Individ Diff* 2009;46:520–4.
45. Hobfoll SE. Conservation of resources: a new attempt at conceptualizing stress. *Am Psychol* 1989;44:513–24.
46. Bernas KH, Major AD. Contributors to stress resistance: testing a model of female's work–family conflict. *Psychol Women Q* 2000;24:170–8.
47. Bruck CS, Allen TD. The relationship between big five personality traits, negative affectivity, type A behavior, and work–family conflict. *J Vocat Behav* 2003;63:457–72.
48. Bundesamt S. *Gesundheit. Grunddaten der Krankenhäuser*. Fachserie 2012. Wiesbaden, 2012.
49. Netemeyer RG, Boles JS, McMurrian R. Development and validation of work–family conflict and family–work conflict scales. *J Appl Psychol* 1996;81:400–10.
50. Kristensen T, Hannerz H, Høgh A, et al. The Copenhagen Psychosocial Questionnaire (COPSOQ). A tool for the assessment and improvement of the psychosocial work environment. *Scand J Work Environ Health* 2005;31:438–49.
51. Nuebling M, Hasselhorn HM. The Copenhagen Psychosocial Questionnaire in Germany: from the validation of the instrument to the formation of a job-specific database of psychosocial factors at work. *Scand J Public Health* 2010;38(3 Suppl):120–4.
52. Scholler G, Fliege H, Klapp BF. [Questionnaire of self-efficacy, optimism and pessimism: reconstruction, selection of items and validation of an instrument by means of examinations of clinical samples]. *Psychother Psychosom Med Psychol* 1999;49:275–83.
53. Sinclair VG, Wallston KA. The development and psychometric evaluation of the Brief Resilient Coping Scale. *Assessment* 2004;11:94–101.
54. Bortz J, Döring N. *Forschungsmethoden und Evaluation für Human- und Sozialwissenschaftler*. Berlin, Heidelberg, New York: Springer, 2002.
55. Innstrand ST, Langballe EM, Falkum E. Exploring occupational differences in work–family interaction: who is at risk? *Int J Stress Manag* 2010;17:38–55.
- 55a. Dumelow C, Littlejohns P, Griffiths S. Relation between a career and family life for English hospital consultants: qualitative, semistructured interview study. *BMJ* 2000;320:1437–40.
56. Pololi LH, Civian JT, Brennan RT, et al. Experiencing the culture of academic medicine: gender matters, a national study. *J Gen Intern Med* 2013;28:201–7.
57. Adam S. High prevalence of work–family conflict among female physicians: lack of social support as a potential antecedent. *Orv Hetil* 2009;150:2274–81.
58. Warde C, Allen W, Gelberg L. Physician role conflict and resulting career changes. Gender and generational differences. *J Gen Intern Med* 1996;11:729–35.
59. Yavas O, Babakus E, Karatepe OM. Attitudinal and behavioral consequences of work–family conflict and family–work conflict. Does gender matter? *Int J Serv Indust Manag* 2008;19:7–31.
60. Frone MR, Yardley JK, Markel KS. Developing and testing an integrative model of the work–family interface. *J Vocat Behav* 1997;50:145–67.
61. Ramirez AJ, Graham J, Richards MA, et al. Mental health of hospital consultants: the effects of stress and satisfaction at work. *Lancet* 1996;347:724–8.
62. Yildirim D, Aycan Z. Nurses' work demands and work–family conflict: a questionnaire survey. *Int J Nurs Stud* 2008;45:1366–78.
63. Boles JS, Johnston MW, Hair JF Jr. Role stress, work–family conflict and emotional exhaustion: inter-relationships and the effects on some work-related consequences. *J Pers Sel Sal Manag* 1997;17:17–29.
64. Boles JS, Howard WG, Donofrio HH. An investigation into the interrelationships of work–family conflict, family–work conflict and work–satisfaction. *J Manag Issue* 2001;13:376–87.
65. Noor NM. Work–family conflict, locus of control, and women's well-being: tests of alternative pathways. *J Soc Psychol* 2002;142:645–53.
66. Kossek EE, Ozeki C. Work–family conflict, policies, and job-life satisfaction relationship: a review and directions of organizational behavior human resources research. *J Appl Psychol* 1998;83:139–49.
67. Rivera-Torres P, Araque-Padilla RA, Montero-Simo MJ. Job stress across gender: the importance of emotional and intellectual demands and social support in women. *Int J Environ Res Public Health* 2013;10:375–89.
68. Almeida DM, Wethington E, Chandler AL. Daily transmission of tensions between marital dyads and parent–child dyads. *J Marriage Fam* 1999;61:49–61.
69. Heinke W, Dunkel P, Brähler E, et al. Burnout in anesthesiology and intensive care: is there a problem in Germany. *Anaesthesist* 2011;60:1109–18.
70. Matthews S, Hertzman C, Ostro A, et al. Gender, work roles and psychosocial work characteristics as determinants of health. *Soc Sci Med* 1998;46:1417–24.
71. Jeffrey Hill E, Jacob JI, Shannon LL, et al. Exploring the relationship of workplace flexibility, gender, and life stage to family-to-work conflict, and stress and burnout. *Community Work Fam* 2008;11:165–81.
72. Halbesleben JRB, Harvey J, Bolino MC. Too engaged? A conservation of resources view of the relationship between work engagement and work interference with family. *J Appl Psychol* 2009;94:1452–65.
73. Bakker AB, Demerouti E. Towards a model of work engagement. *Career Dev Int* 2008;13:209–23.
74. Thomas KW. *Intrinsic motivation at work: what really drives employee engagement*. San Francisco: Berrett-Koehler, 2009.
75. Cudeck R, Browne MW. Alternative ways of assessing model fit. In: Bollen KA, Scott Long J, eds. *Testing structural equation models*. Newbury Park, CA: Sage, 1993:230–58.
76. Baron RM, Kenny DA. The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986;51:1173–82.
77. Langhammer F. *Forscher, Facharzt und Vater*. DIE ZEIT, 2013.